An evolution-based method of epistasis measurement:

theory and application to influenza

Gabriele Pedruzzi and Igor M. Rouzine*

Sorbonne Université, Institute de Biologie Paris-Seine
Laboratoire de Biologie Computationelle et Quantitative LCQB
F-75004, Paris, France

Influenza   protein   sequence   data   are   from   public   database
https://www.fludb.org.

Short Title: Epistasis measurement

Key words: mathematical model, quasi-equilibrium, false-positive, linkage

*Corresponding author:

Igor Rouzine

igor.rouzine@sorbonne-universite.fr

Equipe "Evolution and immunology of pathogens"

Laboratory of Computational and Quantitative Biology

7238 CNRS-UPMC

Bâtiment C, C311C

Sorbonne Université Campus Pierre et Marie Curie

4 rue Jussieu, Paris 75005

+33 7 69 54 91 28

http://www.ibps.upmc.fr/fr/Recherche/umr-7238/evolution-immunologie-

pathogenes

Abstract

Linkage effects in a multi-locus population strongly influence its evolution. Recent models based on the traveling wave approach enable us to predict the speed of evolution and the statistics of phylogeny. However, predicting the evolution of specific sites and pairs of sites in the multi-locus context remains a mathematical challenge. In particular, the effect of epistasis, the interaction of gene regions contributing to phenotype, is difficult both to predict theoretically and detect experimentally in sequence data. A large number of false interactions arising from linkage and indirect interactions which mask true interactions. Here we develop a method to clean false-positive interactions. We start by demonstrating that averaging of the two-way haplotype frequencies over a hundred of independent populations is not enough to clear false interactions. Then, to address this problem, we develop analytically and use a triple-way haplotype test, which isolates true interactions. Next, the fidelity of the test is confirmed on simulated genetic sequences, where the epistatic network known in advance. Finally, we apply the test to a large database on influenza A H1N1 virus sequences of neurominidase from various geographic locations to predict the epistatic network responsible for the transition from the pre-pandemic virus to the pandemic strain. We predict a primary mutation and 15-22 secondary compensatory mutations of variable strength, as many as typically observed for drug resistance and immune escape mutations in HIV. These results present a simple and reliable method to measure epistatic interaction from sequence data.

Author's summary

Interaction of genomic sites creating "fitness landscape" is very important for

predicting the escape of viruses from drugs and immune response and for pssinf

through fitness valleys. Many efforts have been invested into measuring these

interactions from DNA sequence sets. Unfortunately, reproducibility of the

results remains low, due partly to a very small fraction of interaction pairs, and

partly to stochastic noise intrinsic for evolution masking true interactions. Here

we propose a method based on analysis of genetic sequences at three genomic

sites to clean stochastic linkage and apply it to influenza virus sequence data.

Introduction

It was realized long ago that properties of an evolving populations are strongly

affected by the fact that alleles at different genetic sites, loci, are co-inherited unless

separated by recombination. These *linkage* effects include clonal interference

(FISHER 1930; MULLER 1932), genetic background and genetic hitchhiking effect (RICE

2004), enhanced accumulation of deleterious mutations (Muller ratchet)

(FELSENSTEIN 1974), and the increase of random genetic drift on one locus due to

selection at another locus (HILL AND ROBERTSON 1966a). Linkage slows down

substantially adaptation of population, and creates random associations between

pairs of mutations that occurred on the same branch of the ancestral tree.

These effects has been taken into account by early mathematical models

considering two loci (KIMURA 1994) and, more recently, by the traveling wave

approach, which can describe an arbitrarily large number of connected sites

(ROUZINE *et al.* 2003; DESAI AND FISHER 2007; HALLATSCHEK 2011; GOOD *et al.* 2012).

These models usually consider fitness classes of sequences subjected to selection,

mutation, random genetic drift. They show that population represent a narrow

fitness distribution traveling in the fitness space in a direction depending on initial

conditions and parameters. It consists from the deterministic bulk subject to

selection alone, and the stochastic leading edge. At the edge, generation and

establishment of new beneficial mutations, limits the speed of adaptation.

Alternatively, the distribution may move backwards accumulating deleterious

alleles (Muller ratchet). These models were able to predict accurately several

integral parameters of evolution in terms of model parameters, including population

size, mutation rate, and the distribution of selection coefficients over loci. Predicted

parameters include the adaptation rate, Muller ratchet rate, equilibrium state

(ROUZINE *et al.* 2003; ROUZINE *et al.* 2008), the probability of fixation and the

probable selection coefficient (GOOD *et al.* 2012), and the statistics of the ancestral

tree (WALCZAK *et al.* 2012; NEHER AND HALLATSCHEK 2013).

With all this progress, the evolution of separate sites in the multi-site still

remains an open question. How do allelic frequencies at each site change in time?

Although they are known to follow random trajectories, what can be said about the

average allelic frequency of a given site with a given fitness effect of mutation?

What can we say about the evolution of site pairs, especially, in the presence of the

epistatic interaction?

Epistasis, the interaction of genes and gene regions contributing to

phenotype, is an omnipresent phenomenon (CARLBORG *et al.* 2006). Gene

interactions are responsible for 70% of the organism's genetic inheritance (ZUK *et al.*

2012) and create fitness valleys in the evolutionary path (WEISSMAN *et al.* 2009). In

pathogens, epistasis accelerates the development of drug resistance and immune

escape and impedes its reversion (NIJHUIS *et al.* 1999; LEVIN *et al.* 2000; PIANA *et al.*

2002; HANDEL *et al.* 2006; GONZALEZ-ORTEGA *et al.* 2011; WU *et al.* 2018). Most of HIV

variation in untreated patients has been argued to arise from mutations

compensating virus escape from the immune response (ROUZINE AND COFFIN 1999).

A number of approaches have been proposed to measure epistasis from

genomic data (CORDELL 2009; CHEN *et al.* 2011; UEKI AND CORDELL 2012). The simplest

methods are based on pairwise allelic correlations (HILL AND ROBERTSON 1966b;

BARTON 1995). However, other forces, such as linkage and indirect interactions,

create strong correlations even between non-interacting pairs of loci, much more

numerous than epistatic pairs. Stochastic effects are a serious obstacle to the
detection of epistatic effects (WEI *et al.* 2014). Mathematical modeling shows that,
in a single asexual population with a relative short genome, stochastic linkage
overshadows the epistatic footprint, except in a narrow range of times and
parameters (PEDRUZZI AND ROUZINE 2019).

A tree-based real-time technique was used to detect epistatic interactions in
influenza A including those that are known to confer resistance to drug oseltamavir
(KRYAZHIMSKIY *et al.* 2011; NEVEROV *et al.* 2015). They identified compensatory
mutations as closely following primary mutations and appearing in the same
lineage. A method to discern directly interacting pairs from false (indirect)
interactions in a highly-diverged single-time protein sequence set from different
species has also been proposed (WEIGT *et al.* 2009; COCCO *et al.* 2018). The idea was
to combine a covariance method based on mutual information with the estimates of
the direct mutual information. The direct interactions are estimated from the
condition that entropy is maximal given observed frequencies of single sites and
pairs of sites. A similar technique has been used to measure the fitness landscape of
Ab binding regions of HIV surface protein gp120 (LOUIE *et al.* 2018).

While the last data-based technique allows to eliminate indirect interactions
for highly-divergent sequences from different species, it cannot eliminate linkage
effects present in a single or several recently-diverged populations. The divergence
of independent populations from a common ancestor in a random direction creates
high covariance (linkage disequilibrium) of random sign for most of site pairs
masking the covariance from epistasis (PEDRUZZI AND ROUZINE 2019).

142   Recently, we proposed another, evolution-based approach to the problem.
The challenge we faced was to predict analytically the co-evolution of a specific pair
of loci linked to a multi-site genome and then use this information to estimate
epistatic interaction (PEDRUZZI *et al.* 2018). Using a simulated population, we verified
that, in a broad parameter range, the distribution of alleles across genomes and
sites, within each fitness class, is controlled by the condition of maximum entropy
at given fitness. We demonstrated the existence of a broad parameter and time
range where each fitness class of an adapting population has sufficient time to

assume the most likely and disordered state, given by the current average fitness,

which changes slowly in time. This situation is referred to as "quasi-equilibrium"

(not to be confused with "linkage quasi-equilibrium" or "mutation-selection

balance") . In quasi-equilibrium, the entropy S is at its current maximum and is a

function of fitness W, which slowly changes in time during the process of

adaptation: $S = F[W(t)]$. The following simple derivation employs the fact, known

from all the traveling-wave models cited above, that the distribution of fitness W is

relatively narrow, $\text{std}[W] \ll W$. Quasi-equilibrium is possible due to a very slow

motion of the fitness distribution, limited by the slow establishment of new

beneficial alleles at the front edge of the distribution (ROUZINE *et al.* 2003; DESAI AND

FISHER 2007; NEHER *et al.* 2010; HALLATSCHEK 2011).   This situation exists very far

mutation-selection balance, i.e., full equilibrium.

Consider a pair of loci (sites). Assume that each site can have two variants,

either a better-fit allele or a less-fit allele, denoted 0 or 1, and s_1, s_2 are the

respective fitness effects of mutation at the sites. We can have 4 possible haplotypes,

00, 01, 10, 11. Let us also assume that difference in fitness between two alleles is the

two sites interact epistatically, so that two deleterious alleles emerging together,

denoted 11, partly compensate each other to the degree E, where $0 < E < 1$. The

log fitness values of the four haplotypes are

$$W_{00} = 0, \ W_{01} = -s_1, W_{10} = -s_2, \ W_{11} = -(s_1 + s_2)(1 - E),$$

respectively (Fig. 1). If E is allowed to change below 0 or above 1, we can obtain any

type and sign of epistasis: positive, negative, overcompensation.

The frequency of haplotype in population f_{ij} is proportional to the number of

possible sequence configurations of the rest of the genome (grey box in Fig. 1a),

exp(S_{rest}). Entropy S_{rest}, as we explained above, is restricted by the fitness of the rest

of genome equal to $W - W_{ij}$. Hence, the entropy of each haplotype subset is $S_{rest} = S(W$

$- W_{ij}$). Since the genome is long, we can safely assume that W_{pair} is much smaller

than *W,* so that the corresponding change in entropy is small and proportional to
W_{ij}. Hence, we can approximate

$$S(W - W_{ij}) \approx S(W) - \beta W_{ij} \tag{1}$$

where $\beta = |dS/dW|$. The frequency of each haplotype is proportional to the
corresponding configuration number, $\exp[S(W - W_{ij})]$. Combining Eqs. 1 and 2, we
can express the haplotype frequencies as

$$
\begin{aligned}
f_{10} &= f_{00}e^{-\beta s_1} \\
f_{01} &= f_{00}e^{-\beta s_2} \\
188 \quad f_{11} &= f_{00}e^{-\beta(s_1+s_2)(1-E)} \tag{2}
\end{aligned}
$$

where β slowly depends on W and, hence, on time, but is the same for all genomic
sites (PEDRUZZI *et al.* 2018). This simple derivation assumes that the site pair does
not interact with other sites of the genome.

This assumption can be lifted and generalized for any interaction network,
and the function $S(W)$ can be calculated explicitly from a more microsopic model
(PEDRUZZI *et al.* 2018). We also do not need to restrict ourselves to the bi-allelic
model, but we will keep this assumption, since it is fairly good for a relatively short
term evolution.   (A physicist will recognize in Eq. 2 the Boltzman distribution
derived originally for a small subsystem connecting to a large thermal reservoir.)

Eqs. 1, 2 predicts how the average frequencies of single site and pairs of sites
evolve in time. It is also tempting to use it for the practical goal of estimating
parameters E, s_1, s_1 from the observed haplotype frequencies, and thus find the
fitness landscape for the entire genome, at least, if epistasis is pairwise, as we
assumed. In reality, these formulae are not readily applicable to real-life sequences.
because of the strong random variation of f_{ij} among pairs of loci caused by
stochastic linkage effects.

206   Of course, the problem of strong linkage noise causing poor reproducibility of
various methods [see (Wei *et al.* 2014) for review] is not specific for this method of
estimating epistasis. Any attempt to measure epistasis in a single population,
whether by measuring covariance [D', r^2, mutual entropy, (Pedruzzi and Rouzine
2019)] or by tree-based methods used, for example, by (Neverov *et al.* 2015), faces
the same problem. The epistatic pairs in a single population are not observable in
principle, due to the the phylogenetic relationship of sequences in population. As
shown previously by simulation, all sequences in a population closely resemble the
common ancestor, which diverges from the origin in a random direction (Pedruzzi
and Rouzine 2019). As a result, any measure of co-variance, or even the use of the
entire tree, produces only strong noise of random sign. Co-variation due to random
linkage completely masks the epistasis signature in a population. The only way to
resolve this issue is by averaging over many independent populations. But how
many populations do we need to have a small fraction of false positives?

220   As we show in the present work, the number of required populations is
unrealistically large even for a short genome of 50 loci, not to mention more realistic
genomes. The averaging over populations helps only to enrich the observed
network in true bonds, but does not solve the problem: false-positive interactions
dominate. We also not that the existing technique developed for measuring epistasis
by comparing sequences from remote species (Weigt *et al.* 2009; Cocco *et al.* 2018),
does not help with this problem either, because it is designed for indirect
interactions, not for filtering out linkage effects for closely-related populations.

228   Here we offer a new technique, based on the application of the quasi-
229   equilibrium argument (above) to triple-way haplotypes. We show analytically and
230   by simulation that a triple-way haplotype test represents an effective way to
231   eliminate residual false links, including linkage and indirect interactions. We will
232   demonstrate its high fidelity in a broad parameter range on the mock sequences
233   obtained by computer simulation.

234   After training the method on simulated sequences, we apply this technique to
235   real virus sequences from an adapting population. Influenza virus evolving in a
236   population was demonstrated to obey the traveling wave theory with an effective

selection pressure caused by the accumulating memory cells (Rouzine and Rozhnova 2018; Yan *et al.* 2019). Therefore, it satisfies the same quasi-equilibrium criteria (the evolution is slow and the distribution of fitness is narrow) and is amenable to our method. Using 8000 influenza sequences obtained from various geographic locations sequences, both before and after the pandemic of 2009, we detect and estimate quantitatively the epistatic network responsible for the transition of two influenza proteins (NA and HA) between the old and new strains, which share 80% of homology. To avoid a possible misunderstanding, the origin and the cause of pandemics, such as antigenic shifts, are not addressed in this paper; we focus only on the epistatic network, which allows to obtain the new variant from the old variant.

RESULTS

Simulation model to generate sequences for the test. We start by simulating *in silico* stochastic asexual evolution of a virus using a Wright-Fisher model in the presence of the factors of mutation, random genetic drift, and selection with known epistatic interactions (Fig. 2b). Simulation generates genome sequences demonstrating LD originating from epistatic interactions, indirect interactions, and stochastic linkage effects. We assume only two alleles per amino acid position (site): better-fit (denoted 0) or less-fit (denoted 1). The binary simplification provides a major reduction in the computational cost and is especially accurate for relatively conserved sequences (Rouzine and Coffin 2005). We set some pairs of sites to interact epistatically, with a mutual degree of compensation between deleterious alleles set at $E = 0.75$. We consider the same example of double arches (Fig. 1 or 2B).

First step: averaging over populations. Out task is to recover true epistatic interactions from simulated sequences using pair-wise association analysis (Fig. 2A). Previous work has shown that their detection is masked by stochastic linkage, which is known for its strong effects on evolution, including clonal interference,

genetic background effect, and alterations in the adaptation speed and genealogy
(FISHER 1930; BARTON 1995; ROUZINE *et al.* 2003; BRUNET *et al.* 2008; ROUZINE AND
COFFIN 2010; GOOD *et al.* 2012; NEHER AND HALLATSCHEK 2013). To decrease the
linkage effects on detection, we calculate pairwise haplotypes frequencies f_{ij} for each
pair of sites (*Methods*) and average them over multiple evolutionary-independent
populations of the same size (PEDRUZZI AND ROUZINE 2019). Then, we perform a
pairwise association analysis using a correlation measure shown to be a direct
estimator of the degree of mutual compensation of two deleterious alleles, *E*
(PEDRUZZI *et al.* 2018). More traditional LD measures, including D' and Pearson
coefficient r^2, have similar performance but do not estimate *E* directly (PEDRUZZI AND
ROUZINE 2019). We select pairs with high correlation, UFE > 0.6. For a single
population, the raw cluster of predicted pairs is extremely complex and completely
hides true epistatic interactions, in agreement with previous results (PEDRUZZI *et al.*
2018) (Fig 2C). A significant complexity reduction is obtained by averaging f_{ij} over
200 independent populations (Fig. 2D).

Second step: Triple-site haplotype method. Although much simpler, the resulting
network in Fig. 1D is still densely populated by residual stochastic LD and indirect
links, which are as strong as direct interactions. To filter the residual false links, we
use a procedure based on the quasi-equilibrium approximation *(Introduction* and
Methods, Analytic Derivation), and then test its validity by comparison of the
predicted network with the network set in simulation, as follows.

We re-calculate UFE for every potential link *i,j* by considering only the
sequences that contain 0 at one of the neighbor nodes of the link (3-locus
haplotypes 110, 100, 010, and 000), and denote it UFE_{ij0}. A 0-node cuts a possible
indirect correlation for the link even it is very strong (*Methods*). Considering
different 0 nodes, we find the minimum value of UFE_{ij0} over all possible detours. The
scatter plot in Fig. 2E demonstrates that, for the false pairs, $\min(UFE_{ij0})$ is several-
fold smaller than UFE_{ij}. The reason for this effect is that a 0-node removes a detour
path around the link that causes indirect correlations. Therefore, we identify and
remove false links as those with low UFE_{ij0}/UFE. The exact cutoff is not crucial, as

long as we average f_{ij} over, at least, 20 populations, so that the two groups, false and

true interactions, remain distinct. As a result, we obtain 100% perfect detection. We

even obtain correct estimates for the compensation strength, within 15% accuracy

UFE = E (Fig. 2F, green links).

303   We can conclude that, as predicted previously [2008, 2009], the quasi-

equilibrium approximation works in this parameter range, which corresponds to

the traveling wave regime, and that the triple-haplotype method has a high fidelity

with simulated sequences with a known epistatic network.

Application to influenza A virus data. After training the method on simulated

sequences, we calculate interaction network for virus sequences from an evolving

population. Influenza evolution in human population was shown to be described by

the traveling wave framework [rr, neher], which justifies the use of quasi-

equilibrium approximation during the process of adaptation (not to be confused

with mutation-selection equilibrium) [PBR]. Therefore, we have the right to apply

the technique to influenza virus sequences.

315   We focus on the two surface proteins of Influenza A virus strain H1N1,

316   Neuraminidase (NA) and Hemagglutinin (HA), both important targets of immune

317   response and drugs. Our aim is to identify subnetwork responsible for the transition

318   of the pre-2009 strain of H1N1 to the post-2009 of the same strain. For this aim, we

319   need to compare sequences of the first strain sampled worldwide to sequences of

320   the second strain sample worldwide.

321   We have downloaded 8440 and 9811 sequences for NA and HA, respectively,

from a public database (https://www.fludb.org). They were collected worldwide,

from different geographic locations, between years 2005 and 2010, and included

both pre-pandemic and post-pandemic strain.   After aligning and binarizing

sequences by setting consensus residues to 0 and non-consensus to 1, we observed

a bimodal distribution of sequences in allele frequency, with two separate maxima.

The bimodal distribution is related to the emergence of a new strain with 80%

homology with the old strain in 2009.

In order to compensate for unequal sampling from the early strain and the
late strain, the sequences with mutation frequency per genome less than a preset
value d_v were randomly sampled and down-weighted by a coefficient D_w ranging
from 5% to 50%. To obtain the average pairwise haplotype frequencies f_{ij} , we
repeated the sampling multiple times. Then, we followed the procedure described
above to predict the intra-protein network of interactions.

Sensitivity to sampling. The resulting networks are fairly robust to the variation of
(d_v, D_w) (Fig. 3A-F). There exists a plateau region of (d_v, D_w) where the total number
of links varies weakly (Fig. 3E). The dependence of results on (d_v, D_w), which predict
between 15 and 22 compensatory sites, probably originates from the unequal
presentation of local subpopulations in the database. As a result, stochastic linkage
effects average out less than optimally. Below, we choose network variants shown in
Fig 3D and 3J as representative for NA and HA, respectively. We observe that site
248 in NA serves the primary site for multiple compensatory links (Fig. 3D).

Structural interpretation. The predicted epistatic sites are shown on the three-
dimensional protein structure of NA. The active pocket of NA (purple) serves to bind
sialic acid on target cell surface (Fig 4A). These results suggest that a mutation at
residue 248 near the pocket was the primary mutation responsible for the
replacement of the old variant of NA with the new variant of NA, and the other
mutations shown on the wheel are compensatory mutations helping NA to restore
and improve its fitness. Mutation 248 was shown to enhance the low-pH stability of
NA (TAKAHASHI *et al.* 2013), and it is found in all influenza A H1N1 strains after 2009
pandemic, regardless of a geographic location (SEIBERT *et al.* 2012; BYARUGABA *et al.*
2016; OTTE *et al.* 2016).

DISCUSSION

In the present work, we propose an efficient evolution-based method to identify the co-variance caused by epistasis from the co-variance caused by stochastic linkage effects and indirect interaction. The idea (*Methods*) is to pre-average the observed haplotype frequencies over independent (or quasi-independent) populations, select the links with high co-variance, and then apply a tri-way haplotype test for each candidate bond to eliminate the effects of linked sites due to common inheritance.

The test is designed analytically ,based on the quasi-equilibrium state within the traveling fitness wave (not to be confused with quasi-linkage equilibrium or mutation-selection balance) tested by simulation in a broad parameter range (PEDRUZZI *et al.* 2018; BARLUKOVA *et al.* 2020). Intuitively, because population evolves slowly, the distribution of alleles between sites has sufficient time to obtain the most likely and disordered state given a current average fitness of population.

We use a mock sequence set evolved in a Wright-Fisher population with a known epistatic network to demonstrate the high fidelity of the method in the controlled environment. The method eliminates false bonds caused by both indirect interactions and linkage, at least, in the case of relatively simple epistatic network and moderately diverse populations (< 40% diversity).

We apply this technique to identify primary and secondary mutations in the two surface proteins of influenza A H1N1 responsible for the transition between the pre-2009 variant and the post-2009 variants of these proteins. We do not address the complex origins of 2009 pandemic in this work, which involves antigenic shifts. Influenza virus has been shown to map to the traveling wave theory (ROUZINE AND ROZHNOVA 2018; YAN *et al.* 2019), which justifies the use of the quasi-equlibrium assumption. For NA protein, we observe a single primary site and 15-20 strong multiple compensatory mutations, whose is in the same range as the number of compensatory mutations observed for HIV for drug-resistant strains.

As compared to the existing technique of removal of the indirect bonds developed for highly diverged species (WEIGT *et al.* 2009; COCCO *et al.* 2018), the method is designed to be used for recently diverged populations of the same species with small to moderate diversity. Furthermore, our method primarily targets stochastic linkage, which is less important when comparing sequences from

differeent species (WEIGT *et al.* 2009; COCCO *et al.* 2018). Where both methods can be

potentially applied, such as calculating the fitness landscape of HIV Ab-binding

regions (LOUIE *et al.* 2018), our method is computationally faster, because it is local.

Indeed, we can consider one pair of loci at a time and not worry about simultaneous

optimization of $L^2/2$ parameters of the full interaction matrix. Also, it helps to avoid

the situation when the number of fitting parameter is too large, and the system is

over-defined.

To summarize, we proposed a technique to predict evolution of pairs of sites

and tease it out from stochastic linkage effects. We hope that our quasi-equilibrium

approach and further development of this technique will prove useful for all

researchers interested in finding fitness landscapes of various organisms from

genetic samples.

METHODS

Wright-Fisher model for simulation. We simulate the evolution of a haploid

population of N binary sequences (ACEVEDO *et al.*), where each genome site

(nucleotide position) numbered by i =1, 2, …, L has either K_i =0 or K_i =1, in discrete

generations (Wright-Fisher model). The processes included are random mutation at

rate μL, per genome, natural selection, random genetic drift, and an epistatic

network with a set strength and topology. Recombination is absent. Previous work

in silico shows that sufficient levels of recombination would enhance epistatic

detection (PEDRUZZI AND ROUZINE 2019). In the presence of epistasis, the average

progeny number (Darwinian fitness) of sequence [K_i] is set to e^W where

$$W = -\sum_{i=1}^{L} s_i K_i - \sum_{i<j}^{L} s_{ij} K_i K_j \tag{3}$$

$$s_{ij} = E_{ij}(s_i + s_j)T_{ij} \tag{4}$$

where the selection coefficients s_i and s_j represent the fitness costs of two

deleterious mutations that are partially compensated by each other. By definition, E_{ij}

is the degree of compensation of deleterious alleles at sites i and j. Values $E = 0$ and

1 represents no epistasis and full compensation, respectively.

Quasi-equilibrium approximation. In a broad parameter range, an adapting

asexual or partly sexual population represents a slowly-moving, narrow peak in

fitness space (ROUZINE *et al.* 2003; ROUZINE AND COFFIN 2005; ROUZINE *et al.* 2008;

ROUZINE AND COFFIN 2010; GOOD *et al.* 2012). Its speed is restricted by the formation

and fixation of rare beneficial mutations in the few best-fit genomes (the leading

edge). Because the fitness distribution moves slowly, the alleles entropy of the

mutation distribution over genomes has enough time to reach its current maximum,

restricted by the current average fitness of the population. This situation is called

"quasi-equilibrium". We verified the validity of Eq. 3 in a broad range of parameters

and initial conditions after time $\sim 1/\langle s \rangle$ (PEDRUZZI *et al.* 2018). When the population

finally arrives in mutation-selection equilibrium, the presence of deleterious

mutation events compensates the natural selection and creates linkage equilibrium

and the mutation-selection balance. At this point, the method stops working.

Linkage measure. In our recent work (PEDRUZZI *et al.* 2018), we introduced a binary

measure of LD

$$\text{UFE} = 1 - \frac{\log\left(f_{11}/f_{00}\right)}{\log\left(f_{01}f_{10}/f_{00}{}^2\right)} \tag{5}$$

where f_{00} , f_{01}, f_{10} f_{11} are the average-over-ensemble haplotype frequencies, where 0

corresponds to consensus sequence. As follows from Eqs. (1) and (2), UFE = E for an

isolated epistatic pair (PEDRUZZI *et al.* 2018).

UFE performs similarly to more widely used measures, such as Lewontin's D'

and Pearson correlation coefficient, r^2 (PEDRUZZI AND ROUZINE 2019) or mutual

information (WEIGT *et al.* 2009; COCCO *et al.* 2018). As compared to these measures,

UFE has the unique advantage of directly measuring the degree of mutual

compensation of two alleles E (Eq. 2), provided they are epistatically isolated. If they
are a part of a network, this measure overestimates E. We calculate UFE for every
potential epistatic link in our simulated sequence set (*Results*).

Tri-site test of false bonds. To test whether the interaction is direct, and not due to
linkage or an indirect detour, we calculate for each suspected pair i, j the value

$$\text{UFE}_{ij0} = 1 - \frac{\log\left(f_{110}/f_{000}\right)}{\log\left(f_{010}f_{100}/f_{000}{}^2\right)} \tag{6}$$

where the third 0 selects only for the sequences with the consensus allele 0 at a
chosen neighbor site of the pair. We try all possible neighbor sites and calculate the
minimum value of UFE_{ij0} over all neighbors. This method cuts the most important
indirect correlation by "detour" for a site pair. It works for both linkage and a
modest amount of indirect correlation.

To illustrate how this method works on indirect bonds, let us consider a
simple analytic example in Fig. 1. We assume that some sites carry deleterious
alleles with equal selection coefficient $s_i = s_0$ in Eq. 3, with a fixed epistatic strength
$E_{ij} = E$. In this simple case, we can fully characterize a genome by the numbers of
interacting allelic clusters of different size. To do so, let k_i define the number of
clusters with i nodes and b_i bonds. Then, from Eq. 3, we can express fitness as a sum
over clusters of different size (PEDRUZZI *et al.* 2018)

$$W \equiv -s_0 f_0 L = -s_0 \sum_{i=1}^{i_{max}} k_i(i - 2Eb_i) \tag{7}$$

New notation f_0 represents the frequency of uncompensated mutations with total
fitness W. The number of bonds b_i depends on the topology, and $b_1 = 0$, $b_2 = 1$. In our
example with double arches, $b_3 = 2$ (Fig. 1).

As it has been shown previously in the literature on traveling wave (see
Introduction), the population of genomes represents a slowly moving, narrow
fitness peak. Because it in the state of quasi-equilibrium determined by its current

average fitness, at each moment of time, numbers k_i are determined by the condition

that the entropy of the system is maximum given the value of fitness, Eq. 7. Entropy

S is defined as the log number of sequence configurations

484
$$S = \log\left[\prod_{i=1}^{i_{max}} C_{L_i}^{k_i} (n_i)^{k_i}\right] \tag{8}$$

where L_i is the number of all possible locations for a cluster of size i, and n_i is the

number of each cluster's configurations (shapes). The values of L_i and n_i depend on

the network topology.

Previously, we applied this argument for several topologies to predict cluster

numbers (PEDRUZZI *et al.* 2018). We showed, for the one in Fig. 1, that the numbers of

clusters of size i = 1, 2 and 3 in, normalized as $f_i = k_i/L$, are related as

493
$$f_2 = \tfrac{1}{3} f_1^{2-2E}, \ \ f_3 = \tfrac{2}{3} f_1^{3-4E} \qquad E < \tfrac{3}{4} \tag{9}$$

495 The 1st and 3d site in each triplet in Fig. 1 do not interact directly, but only

496 indirectly through site 2. For these two sites, the haplotype frequencies are

498
$$f_{11} = 3f_3 + f_1^2, \quad f_{10} = \tfrac{3}{2} f_2 + f_1 \tag{10}$$

500 When epistatic interaction is sufficiently strong, as given by the condition $E > 1/2$,

501 triplets dominate over single alleles and their doubles, as given by $f_1 \ll f_2 \ll f_3$

502 (Eq. 9). In this case, we can approximate haplotype frequencies as

504
$$f_{11} \approx 2f_1^{3-4E}, \ f_{10} \approx \tfrac{1}{2} f_1^{2-2E}, \qquad \tfrac{1}{2} < E < \tfrac{3}{4} \tag{11}$$

Using covariance measure UFE_{ij} defined in Eq. 3, we obtain $\text{UFE}_{ij} \approx \frac{2E-1}{2-2E}$. We

observe that the indirect covariance between sites 1 and 3 is very strong, ~ 1. [In

fact, covariance is approximately the same for directly interacting sites 1 and 2 (Fig.

1), for which we previously obtained $f_{11}^{dir} \approx f_1^{3-4E}, f_{10}^{dir} \approx \frac{1}{3} f_1^{2-2E}$, see (PEDRUZZI *et*

al. 2018), *Supplement*, Eqs. (3.29) and (3.30)].

However, if we calculate UFE_{ij0} instead of UFE_{ij} by including only the

sequences with majority allele "0" found at position 2, we obtain

$$f_{101} \approx f_1^2, \; f_{100} \approx f_1 \qquad\qquad (10)$$

instead of Eq. 9, and Eq. 6 yields $\text{UFE}_{ij0} = 0$. Thus, phantom covariance disappears,

when we select only sequences with a majority allele between the sites of interest.

This result is intuitively clear: by the definition of fitness (Eq. 1), only minority

alleles "1" interact with each other, while majority alleles are neutral background.

The same method turns out to be extremely effective for eliminating false bonds

created by linkage (see *Results*, Fig. 3).

Sequence preparation.   We have applied the three-way test to influenza virus.

Evolution of the virus in population has been mapped recently to traveling-wave

models (ROUZINE AND ROZHNOVA 2018; YAN *et al.* 2019), which justifies the use of

quasi-equilibrium approximation dependent on current average fitness (see Intro

for the explanation). We performed a multiple progressive alignment for amino acid

sequences obtained for two proteins of Influenza A virus strain H1N1,

Neuraminidase (NA) and Hemagglutinin (HA) from public database

https://www.fludb.org.

We have used all the sequences discovered in the database for the period

2000-2010, which all come from various geographic locations. Our aim was to

understand the transition between the protein variants before and after pandemic

of 2009, which have 80% of mutual homology. We aimed at discovering only the

epistatic sub-network related to that transition, and are not interested in any other

epistatic interactions in these proteins. For this end, we compared worldwide

samples of sequences from the two strains. We randomly sampled similar amounts

of sequences from the first and second strains, and re-sampled them several

hundred times (*Results*). We also checked the robustness of the results to exact

sampling size (see *Results*).

Pairwise distances between sequences were computed using pairwise

alignment with the Gonnet scoring matrix implemented in MATLAB. To calculate the

guide tree we used the neighbor-joining method assuming equal variance and

independence of evolutionary distance estimates. The obtained consensus served as

a universal reference to binarize data sequences. Before applying the detection

algorithm, the protein sequences were binarized, by direct comparison of each

sequence to the consensus. Each amino-acid residue was set to 0 or 1 for consensus

or non–consensus. Pooling all mutations per site ignores the chemistry of

substitutions. In exchange, this approach greatly reduces the number of haplotype

combinations and increases the sensitivity by effectively increasing the haplotype

frequencies. Next, we measure the mutational frequency for each sequence, for each

site. The subset of low-diversity sequences with allelic frequency below a cut-off d_v

is randomly sampled and down-weighted according to the coefficient D_w. Then, we

determined the average pairwise and triple haplotype frequencies for pairs and

triplets of sites, to be used in the method described above.

3D structures. To characterize the three-dimensional network of epistatic

interaction, we used software package ChimeraX from internet site

https://www.rbvi.ucsf.edu/chimerax/.

Data availability. Influenza sequence data are from public database

https://www.fludb.org.

Acknowledgements. We thank Martin Weigt and Alessandra Carbone for useful

comments. This research has been funded by Agence Nationale de la Recherche

grant J16R389 to IMR, http://www.agence-nationale-recherche.fr/.

References

Acevedo, A., L. Brodsky and R. Andino, 2014 Mutational and fitness landscapes of an
RNA virus revealed through population sequencing. Nature 505**:** 686-690.
Barlukova, A., G. Pedruzzi and I. M. Rouzine, 2020 The evolutionary origin of the
universal   distribution   of   mutation   fitness   effect
https://biorxiv.org/cgi/content/short/867390v1.
Barton, N. H., 1995 Linkage and the limits to natural selection. Genetics 140**:** 821-
841.
Brunet, E., I. M. Rouzine and C. O. Wilke, 2008 The stochastic edge in adaptive
evolution. Genetics 179**:** 603-620.
Byarugaba, D. K., B. Erima, M. Millard, H. Kibuuka, L. Lkwago *et al.*, 2016 Whole-
genome analysis of influenza A(H1N1)pdm09 viruses isolated in Uganda
from 2009 to 2011. Influenza Other Respir Viruses 10**:** 486-492.
Carlborg, O., L. Jacobsson, P. Ahgren, P. Siegel and L. Andersson, 2006 Epistasis and
the release of genetic variation during long-term selection. Nat Genet 38**:**
418-420.
Chen, C. C., H. Schwender, J. Keith, R. Nunkesser, K. Mengersen *et al.*, 2011 Methods
for identifying SNP interactions: a review on variations of Logic Regression,
Random Forest and Bayesian logistic regression. IEEE/ACM Trans Comput
Biol Bioinform 8**:** 1580-1591.
Cocco, S., C. Feinauer, M. Figliuzzi, R. Monasson and M. Weigt, 2018 Inverse
statistical physics of protein sequences: a key issues review. Rep Prog Phys
81**:** 032601.
Cordell, H. J., 2009 Detecting gene-gene interactions that underlie human diseases.
Nature reviews. Genetics 10**:** 392-404.
Desai, M. M., and D. S. Fisher, 2007 Beneficial mutation selection balance and the
effect of linkage on positive selection. Genetics 176**:** 1759-1798.
Felsenstein, J., 1974 The evolutionary advantage of recombination. Genetics 78**:**
737-756.
Fisher, R. A., 1930 *The genetical theory of natural selection*. Clarendon Press, 1958,
Oxford, United Kingdom.
Gonzalez-Ortega, E., E. Ballana, R. Badia, B. Clotet and J. A. Este, 2011 Compensatory
mutations rescue the virus replicative capacity of VIRIP-resistant HIV-1.
Antiviral Res 92**:** 479-483.
Good, B. H., I. M. Rouzine, D. J. Balick, O. Hallatschek and M. M. Desai, 2012
Distribution of fixed beneficial mutations and the rate of adaptation in
asexual populations. Proc Natl Acad Sci U S A 109**:** 4950-4955.
Hallatschek, O., 2011 The noisy edge of traveling waves. Proc Natl Acad Sci U S A
108**:** 1783-1787.
Handel, A., R. R. Regoes and R. Antia, 2006 The role of compensatory mutations in
the emergence of drug resistance. PLoS Comput Biol 2**:** e137.
Hill, W. G., and A. Robertson, 1966a The effect of linkage on limits to artificial
selection. Genet. Res. 8**:** 269-294.
Hill, W. G., and A. Robertson, 1966b The effect of linkage on limits to artificial
selection. Genet Res 8**:** 269-294.

Kimura, M., 1994 *Population genetics, molecular evolution, and the neutral theory.*
Selected papers. The University of Chicago Press, Chicago.
Kryazhimskiy, S., J. Dushoff, G. A. Bazykin and J. B. Plotkin, 2011 Prevalence of
epistasis in the evolution of influenza A surface proteins. PLoS Genet 7**:**
e1001301.
Levin, B. R., V. Perrot and N. Walker, 2000 Compensatory mutations, antibiotic
resistance and the population genetics of adaptive evolution in bacteria.
Genetics 154**:** 985-997.
Louie, R. H. Y., K. J. Kaczorowski, J. P. Barton, A. K. Chakraborty and M. R. McKay,
2018 Fitness landscape of the human immunodeficiency virus envelope
protein that is targeted by antibodies. Proc Natl Acad Sci U S A 115**:** E564-
E573.
Muller, H. J., 1932 Some genetic aspects of sex. Am. Nat. 66**:** 118-128.
Neher, R. A., and O. Hallatschek, 2013 Genealogies of rapidly adapting populations.
Proc Natl Acad Sci U S A 110**:** 437-442.
Neher, R. A., B. I. Shraiman and D. S. Fisher, 2010 Rate of adaptation in large sexual
populations. Genetics 184**:** 467-481.
Neverov, A. D., S. Kryazhimskiy, J. B. Plotkin and G. A. Bazykin, 2015 Coordinated
Evolution of Influenza A Surface Proteins. PLoS Genet 11**:** e1005404.
Nijhuis, M., R. Schuurman, D. de Jong, J. Erickson, E. Gustchina *et al.*, 1999 Increased
fitness of drug resistant HIV-1 protease as a result of acquisition of
compensatory mutations during suboptimal therapy. AIDS 13**:** 2349-2359.
Otte, A., A. C. Marriott, C. Dreier, B. Dove, K. Mooren *et al.*, 2016 Evolution of 2009
H1N1 influenza viruses during the pandemic correlates with increased viral
pathogenicity and transmissibility in the ferret model. Sci Rep 6**:** 28583.
Pedruzzi, G., A. Barlukova and I. M. Rouzine, 2018 Evolutionary footprint of
epistasis. PLoS Comput Biol 14**:** e1006426.
Pedruzzi, G., and I. M. Rouzine, 2019 Epistasis detectably alters correlations
between genomic sites in a narrow parameter window. PLoS ONE in press.
Piana, S., P. Carloni and U. Rothlisberger, 2002 Drug resistance in HIV-1 protease:
Flexibility-assisted mechanism of compensatory mutations. Protein Sci 11**:**
2393-2402.
Rice, S. H., 2004 *Evolutionary Theory: Mathematical And Conceptual Foundations.*
Sinauer Associated.
Rouzine, I. M., E. Brunet and C. O. Wilke, 2008 The traveling-wave approach to
asexual evolution: Muller's ratchet and speed of adaptation. Theor Popul Biol
73**:** 24-46.
Rouzine, I. M., and J. M. Coffin, 1999 Search for the mechanism of genetic variation in
the pro gene of human immunodeficiency virus. J Virol 73**:** 8167-8178.
Rouzine, I. M., and J. M. Coffin, 2005 Evolution of human immunodeficiency virus
under selection and weak recombination. Genetics 170**:** 7-18.
Rouzine, I. M., and J. M. Coffin, 2010 Multi-site adaptation in the presence of
infrequent recombination. Theor Popul Biol 77**:** 189-204.
Rouzine, I. M., and G. Rozhnova, 2018 Antigenic evolution of viruses in host
populations. PLoS Pathog 14**:** e1007291.

Rouzine, I. M., J. Wakeley and J. M. Coffin, 2003 The solitary wave of asexual
evolution. Proc Natl Acad Sci U S A 100**:** 587-592.
Seibert, C. W., S. Rahmat, F. Krammer, P. Palese and N. M. Bouvier, 2012 Efficient
transmission of pandemic H1N1 influenza viruses with high-level oseltamivir
resistance. J Virol 86**:** 5386-5389.
Takahashi, T., J. Song, T. Suzuki and Y. Kawaoka, 2013 Mutations in NA that induced
low pH-stability and enhanced the replication of pandemic (H1N1) 2009
influenza A virus at an early stage of the pandemic. PLoS One 8**:** e64439.
Ueki, M., and H. J. Cordell, 2012 Improved statistics for genome-wide interaction
analysis. PLoS Genet 8: e1002625.
Walczak, A. M., L. E. Nicolaisen, J. B. Plotkin and M. M. Desai, 2012 The structure of
genealogies in the presence of purifying selection: a fitness-class coalescent.
Genetics 190: 753-779.
Wei, W.-H., G. Hemani and C. S. Haley, 2014 Detecting epistasis in human complex
traits. Nature Reviews Genetics 15**:** 722.
Weigt, M., R. A. White, H. Szurmant, J. A. Hoch and T. Hwa, 2009 Identification of
direct residue contacts in protein-protein interaction by message passing.
Proc Natl Acad Sci U S A 106**:** 67-72.
Weissman, D. B., M. M. Desai, D. S. Fisher and M. W. Feldman, 2009 The rate at which
asexual populations cross fitness valleys. Theor Popul Biol 75**:** 286-300.
Wu, N. C., A. J. Thompson, J. Xie, C. W. Lin, C. M. Nycholat *et al.*, 2018 A complex
epistatic network limits the mutational reversibility in the influenza
hemagglutinin receptor-binding site. Nat Commun 9**:** 1264.
Yan, L., R. A. Neher and B. I. Shraiman, 2019 Phylodynamic theory of persistence,
extinction and speciation of rapidly adapting pathogens. Elife 8.
Zuk, O., E. Hechter, S. R. Sunyaev and E. S. Lander, 2012 The mystery of missing
heritability: Genetic interactions create phantom heritability. Proc Natl Acad
Sci U S A 109**:** 1193-1198.
687
688

FIGURES AND CAPTIONS

Fig. 1. A simplest epistatic network illustrating the effect of indirect interactions. Black and white circles are deleterious and favorable alleles, respectively, arches show direct (potential) epistatic interactions, red arches are actual interactions between deleterious alleles. Numbers k_i denote the numbers of i-clusters in a genome.

Fig. 2 Schematic diagram of the method and its training on simulated
sequence data. A. The computer model of asexual evolution includes the factors of
random mutation, selection, epistasis, and random genetic drift. Pairwise haplotype
frequencies f_{ij} are averaged over an ensemble of independent simulation runs
(populations). The strength of interactions UFE_{ij} is calculated. The indirect links and
the residual stochastic LD are excluded by using triple-site haplotype frequencies,
UFE_{ij0}. **B.** Epistatic links (green), indirect links (red), and stochastic LD (blue). **C-D.**
The network of strong candidate epistatic interactions predicted (**C**) from a single
population and (**D**) after averaging over 200 populations. **E.** Scatter plot of 3-locus
haplotype min(UFE_{ij0}) shown against UFE_{ij} for the pairs identified in (**D**). Dashed
sector (green): Predicted strong, direct interactions. **F.** Predicted network
accurately recapitulates the pre-set epistatic network. Parameters are shown (**E**).

Fig. 3 Epistatic network predicted from sequence data on surface protein sequences of Influenza A H1N1 obtained between year 2005 and year 2010.

The circular diagrams show the network of interaction for (**A-E**) neuraminidase and (**G-K**) hemagglutinin. (**F, L**) Sequences with homology to the consensus less than d_v are randomly re-sampled many times, with their number downweighted by coefficient D_w. (**F, L**) 2D heatmaps showing the total number of links as a function of d_v (X-axis) and D_w (Y-axis). Different versions of wheels in **A-E** (or **G-K**) correspond to different choices of D_w and d_v shown by crosses in **F, L.**

Fig. 4 Structural location of the predicted epistasis network for the two influenza virus envelope proteins. The figure shows the three-dimensional structure of Influenza A H1N1 proteins (**A**) neuraminidase (PDB ID code 4QVZ) and (**B**) hemagglutinin (PDB ID code 1RUZ). Colored spheres represent predicted epistatic residues from Fig. 3D for (**A**) and Fig. 3J for (**B**). Red sphere: Predicted primary mutation (residue 248 in Fig 3D). Orange spheres: Compensatory residues from Fig. 3D and 3J.