Expansion and re-classification of the extracytoplasmic function (ECF) σ factor family

Delia Casas-Pastor[1], Raphael Rene Müller[2], Anke Becker[1], Mark Buttner[3], Carol Gross[4], Thorsten Mascher[5], Alexander Goesmann[2], Georg Fritz[1,6*]

[1]Center for Synthetic Microbiology (SYNMIKRO), Philipps-Universität Marburg, Germany

[2]Bioinformatics and Systems Biology, Justus-Liebig-Universität, Giessen, Germany

[3]Department of Molecular Microbiology, John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

[4]Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, CA 94158, USA; Department of Cell and Tissue Biology, University of California, San Francisco, San Francisco, CA 94158, USA; California Institute of Quantitative Biology, University of California, San Francisco, San Francisco, CA 94158, USA

[5]Institute of Microbiology, Technische Universität Dresden, Germany

[6]Present address: School of Molecular Sciences, The University of Western Australia, Perth, Western Australia 6009, Australia

* Corresponding author: georg.fritz@uwa.edu.au

Abstract

Extracytoplasmic function σ factors (ECFs) represent one of the major bacterial signal transduction mechanisms in terms of abundance, diversity and importance, particularly in mediating stress responses. Here, we performed a comprehensive phylogenetic analysis of this protein family by scrutinizing all proteins in the NCBI database. As result, we identified ~10 ECFs per bacterial genome on average and classified them into 157 phylogenetic ECF groups that feature a conserved genetic neighborhood and a similar regulation mechanism. Our analysis expands the number of unique ECF sequences ~50-fold relative to previous classification efforts, enriches many original ECF groups with previously unclassified proteins and identifies 22 entirely new ECF groups. The ECF groups are hierarchically related to each other and are further composed of subgroups with closely related sequences. This two-tiered classification allows for the accurate prediction of common promoter motifs and the inference of putative regulatory mechanisms across subgroups composing an ECF group. This comprehensive, high-resolution description of the phylogenetic distribution of the ECF family, together with the massive expansion of classified ECF sequences, enables the application of *in silico* tools for the prediction of important functional residues, and serves as a powerful hypothesis-generator to guide future research in the field.

Introduction

Bacterial homeostasis is achieved through signal transduction mechanisms that connect the extracellular environment with the cytoplasm. Extracytoplasmic function σ factors (ECFs) are amongst the major signal transduction components in bacteria in terms of abundance and importance of the stress responses they mediate (1). As members of the σ[70] family, ECFs guide the RNA polymerase (RNAP) to specific promoter sequences, and thereby enable bacteria to redirect gene expression in response to deteriorating environmental conditions (2, 3). Although ECFs are generally less prevalent than one-component systems (1CS) and two-component systems (2CS), previous studies revealed a

large ECF abundance, with an average of six ECFs per bacterial genome (1), a large diversity, with

more than 90 phylogenetic groups (1, 4–6), and a diverse range of activation mechanisms (7).

Members of the σ^{70} family are modular proteins composed of up to 4 core domains (σ_{1-4}) that are

classified into 4 groups. While group 1 represents the essential, full-length version of the σ^{70} family, σ^{70}

groups 2-4, also known as alternative σ factors, represent truncations of this general structure that are

usually non-essential (8, 9). Members of group 2 lack the $\sigma_{1.1}$ region and members of group 3 lack the

full σ_1 domain (8). Compared to other members of the σ^{70} family, ECFs (or group 4 σ^{70}s) are truly

minimalistic regulators, since they only contain the σ_2 and σ_4 domains, essential for contact with RNAP

and transcription initiation (8). The functions of both domains are well differentiated: while σ_4 initiates

the first step of promoter recognition by binding to the -35 element (10), σ_2 is responsible for

recognition and melting of the -10 element (11). Given that bacteria typically contain several alternative

σ factors that compete for the binding to the RNAP core complex, it is key to regulate the activity of

ECFs in response to changing conditions. The most common mode of regulation of ECFs is their

sequestration by anti-σ factors (3). Anti-σ factors typically contain a cytoplasmic anti-σ domain (ASD)

that sequesters the cognate ECF into an inactive conformation. Upon the onset of the inducing signal,

anti-σ factors undergo conformational changes or get degraded, thereby releasing their cognate ECFs,

which can then guide RNAP to their specific alternative promoter signatures to initiate target gene

expression (2). While ECFs are often activated in response to external stimuli involved in cell envelope

homeostasis and stress adaptation (8, 12, 13), some ECFs also detect cytoplasmic stimuli and

regulate functions such as detoxification of reactive oxidative species, stationary phase survival (14,

15), metal resistance and homeostasis (12), morphological changes throughout the cell-cycle (1, 8)

and virulence (16, 17). Besides the regulation via anti-σ factors, ECFs may also be regulated by

various other mechanisms, such as conformational changes (18, 19), two-component systems (15,

20), C-terminal extensions (21, 22) and the proposed, but never proven, phosphorylation by

serine/threonine kinases, many of which were first predicted by genomics approaches (see (7) for a

review).

Research of the ECF family is heavily dependent on bioinformatic tools from the date of their first

description (24). Indeed, ECFs are especially suitable for the application of prediction tools since they

usually autoregulate their own expression, facilitating the identification of their target promoter, and

because they are co-encoded with regulators of their activity and/or with genes regulated by the ECF.

Accordingly, the first bioinformatic classification of the ECF σ factor family grouped ECFs from less

than 400 genomes into 67 phylogenetic groups, based on sequence similarity, and revealed that

conservation at protein level is often accompanied by conservation of the target promoter motif and a

conserved genomic neighborhood (1). Altogether, this previous work proposed that it is possible to

predict the target promoter of an ECF, its regulatory mechanism and its target genes from sequence

information alone. Following studies expanded the number of phylogenetic groups by focusing on nine

planctomycetal (4) and 100 actinobacterial (5) genomes, again identifying correlations between primary

protein sequence and function.

Although these initial ECF classification studies helped to understand the large diversity of ECFs
across the tree of life, they so far addressed ECFs from a limited number of genomes and/or focused
on specific phyla (1, 4, 5). Based on the relatively sparse sequence basis, some of the initially defined
ECF groups featured natural limitations in that they either defined groups with only very few (less than
10) proteins – so-called "minor" groups (ECF100, ECF102, etc.) (1) – or in that they clustered divergent
sequences into groups that share only few unifying characteristics (e.g. ECF01, ECF10, ECF20) (1, 4).
However, the explosion of annotated sequences in databases, not only from re-sequenced species but
also from new species of underrepresented phyla, nowadays enables a more comprehensive view of
the ECF family. Moreover, the availability of more ECF variants now allows the application of advanced
bioinformatic tools, such as co-variation-based methods. One of the possible outcomes of these
studies are detailed predictions of conserved protein-protein contacts (25) that may guide experimental
analyses of the interplay between ECFs and they regulatory partners.

In the light of the above-mentioned limitations of the initial studies, a comprehensive and robust
expansion of the ECF classification that reflects the massive increase in bacterial genomic sequence
space is overdue. In this work we searched for ECF σ factors in all available genomes and
metagenomes of the National Center for Biotechnology Information (NCBI) database, thereby
expanding the number of identified ECF proteins 50-fold. We clustered the new ECFs into 2,380
subgroups with a high degree of sequence conservation. Subgroups were further grouped into 157
ECF groups according to genetic context conservation and their putative mode of regulation. As a
result, we defined 22 novel ECF groups with no significant similarity to previously described ECF
groups. Furthermore, 62 original groups were preserved and expanded with previously unclassified
sequences. The conservation of the subgroups facilitated downstream *in silico* analyses such as
prediction of conserved target promoter elements, conserved protein domains in the genetic
neighborhood and putative anti-σ factors. Even though the large amount of information collected for
each of the 157 ECF groups only allows us to focus on the most interesting findings, we provide an
extensive compendium of all the information gathered for each group in the supplementary text. The
possibilities offered by this hierarchical, comprehensive classification are illustrated by a recent study
that predicts the functional role of C-terminal extensions in two ECF groups, only possible due to the
expansion of the number of proteins assigned to these groups (26). This wealth of data represents a
comprehensive resource to both computational and experimental researchers and helps guiding the
characterization of ECF σ factors of unknown function.

Results

Rationale of this study
Previous ECF classification efforts (1, 4, 5) were based on 495 genomes and identified a total of 3,554
ECFs, which were classified into 94 ECF groups (summarized in (6)). Upon initiation of this work
(February, 2017), the NCBI database contained 180,909 genomes, of which 4,106 were bacterial
genomes tagged as "reference" or "representative", discarding GenBank assemblies when a RefSeq

assembly was available for the same genome (see
https://www.ncbi.nlm.nih.gov/refseq/about/prokaryotes/). This indicates that the presently available
number and diversity of ECF sequences should be much larger than previously observed. To expand
the library of ECFs, we here performed a comprehensive ECF search in the NCBI database and re-
classified the resulting protein sequences in a hierarchical approach. First, we clustered protein
sequences into fine-grained ECF subgroups with a high degree of sequence similarity, and then
aggregated subgroups into coarse-grained groups that share a common genetic neighborhood and a
putative type of anti-σ factor. Whenever appropriate, the ECF group names were preserved from the
original groups with which they share the same characteristics, e.g. groups ECF41 and ECF42. New
ECF groups that are product of redistributions of original groups or that have no significant similarity to
any of the original groups were designated with running numbers from 201 onwards, e.g. ECF201,
ECF202, etc. Similarly, ECF subgroups were named by adding an 's' to their group's name and a
running number indicating the rank in number of sequences contained in the subgroup, e.g.
ECF201s1, ECF201s2, etc. (for more details see Methods). The similarity among the ECFs contained
in groups allows the identification of common putative target promoter motifs and ECF regulators, as
will be described in detail in the following sections. These hypotheses are confirmed whenever
experimental reports on members of the group exist.

The number of identified ECFs is 50-fold larger than in the founding ECF classification
The number of protein sequences in public databases has expanded extensively since the founding
ECF classification efforts (1) and we expected a proportionate increase in the number of ECF
sequences. To identify novel ECFs, we first extracted the sequences from all previous ECF
classification efforts (1, 4, 5), aligned them and created a general Hidden Markov Model (HMM) for the
ECF core region, including the linker between σ_2 and σ_4, but excluding any potential protein domains
fused N- or C-terminally to the ECF (Fig. 1A). To discriminate ECFs from other σ factors, we first
scored this generic ECF HMM against two sets of training sequences – true ECFs from the founding
classification and a negative control set of σ factors from groups 1, 2 and 3 that additionally contain
domains σ_3 and σ_1 in some cases. This allowed us to define a threshold score that maximizes true
positive ECFs (Fig. 1B; *green*) while minimizing the number of false positive σ factors (Fig. 1B; *red*).
We then selected the non-redundant protein sequences from the NCBI database (see Methods for
selection of relevant genomes), for which the generic ECF HMM yielded scores higher than this
threshold (Fig. 1C). As further quality controls, we filtered for sequences containing the Pfam domains
σ_2 and σ_4 but lacking the σ_3 domain, and discarded proteins with non-amino acidic characters, such as
X or J. This resulted in a library of 177,910 non-redundant ECF sequences. Some of the candidate
ECF σ factors included in this list cluster together with group 3 σ factors, indicating the presence of a
cryptic σ_3 domain, which prompted us to remove them from the list of ECF σ factors. This left us with
177,341 non-redundant ECFs, accounting for a ~50-fold expansion over the number of ECFs in the
founding ECF classification (1, 4, 5) (Fig. 1C). The full list of ECFs extracted during this study can be
found in Table S1.

Next, we analyzed the taxonomic origin of this expanded ECF library to extract/determine the
typical/average ECF numbers found in individual bacterial phyla. To enable such statistics, we focused
on the subset of complete genomes of non-metagenomic origin, classified as NCBI "reference
genomes" or "representative genomes", thereby mitigating bias towards heavily sequenced species.
Analysis of 12,539 ECFs extracted from 1,234 of these genomes showed that the taxonomic
distribution of species became more diverse than in the original classification efforts (Fig. 2A;
Genomes). In particular, the fraction of the three most abundant phyla – Proteobacteria, Actinobacteria
and Firmicutes – was reduced from 86.9% in the original to 77.6% in the new classification. This
reduction was accompanied by an increase in the number of species from underrepresented phyla,
such as Bacteroidetes and Cyanobacteria (Fig. 2A; *Genomes*). In addition, 19 new ECF-containing
phyla emerged (Table S3, *New phyla*). Yet, these 19 phyla have a limited contribution to the overall
ECF database, given their low number of sequenced genomes. This difference in the taxonomic origin
of the species included in original and new classifications naturally changes the taxonomic origin of
ECFs gathered in each library. For instance, the fraction of ECFs from underrepresented genomes,
such as Bacteroidetes and Planctomycetes, is larger in the new ECF library (Fig. 2A; *ECFs*). This is
not the case for Cyanobacteria and Acidobacteria, which contribute a smaller percentage of ECFs than
in the original library (Fig. 2A; *ECFs*). These differences in taxonomic composition in the ECF library
are reflected in the average number of ECFs per genome, which increases from approx. seven ECFs
per genome in the original ECF libraries (1, 4, 5) to about ten ECFs per genome in the new library (Fig.
2B). Confirming the findings of previous reports (1, 5), the number of ECFs per genome is directly
proportional to genome size (Fig. S1), although the average number of ECFs per genome depends on
the phyla of origin (Fig. 2B). Bacteroidetes and Actinobacteria have the greatest abundance of ECFs,
with an average of 22.5 and 17.7 ECFs per genome, respectively (Fig. 2B). Phyla with a lower
abundance of ECFs include Cyanobacteria and Spirochaetes, with an average of 2.7 and 3.7 ECFs per
genome, respectively (Fig. 2B). Firmicutes and Proteobacteria contain an intermediate number of
ECFs, 7.1 and 7.5, respectively (Fig. 2B). These differences might indicate different dependence on
ECFs as signal-transduction system in different phyla, as previously noticed for Actinobacteria, which
are particularly rich in ECFs, but also in 1CS and 2CS (5).
183

ECF classification 2.0
The wealth of new proteins identified in the initial library expansion prompted us to reclassify ECF σ
factors into groups with common characteristics. To this end, we first subjected the 177,910 protein
sequences of the new ECF library to the rapid MMSeqs2 clustering algorithm (27), followed by a
quality control that bisects the resulting clusters until the maximum pairwise k-tuple distance between
sequences was ≤ 0.60 (Fig. 3A, see Methods for more details). Clusters with ≤ 10 proteins were
discarded to ensure high sequence coherence within clusters, while preventing an explosion of small
clusters with limited statistical relevance (Fig. 3A). This procedure yields a total of 2,380 ECF clusters
(referred to as "subgroups") with a median of 22 non-redundant proteins per subgroup (Fig. 3D).

Subgroups capture 77.3% of the proteins, while 22.7% of the proteins remain unclassified, similar to
the statistics in the original classification (1) (Table S1). Permutation tests on subgroups showed that
the average k-tuple distance is significantly lower (two-tailed Student's t-test; p-value < 1e-16) in our
subgroups as compared to random clusters of the same size distribution, indicating that subgroups are
well defined (Fig. 3F).

Then, we computed a phylogenetic tree based on the consensus sequence of each subgroup.
This tree helps to identify the evolutionary relationship between the ECF subgroups (Fig. 3A, bottom;
Fig. 4). As outgroups we included sequences with a low-scoring σ_3 domain, as well as the consensus
sequence of all σ_3-containing proteins in Pfam, the latter of which we selected as root of the tree. Not
surprisingly, proteins with a low-scoring σ_3 domain clustered at the base of the tree (Fig. 4) and formed
three groups with significant similarity to the sporulation σ factor SigF from Firmicutes and
Actinobacteria, the flagellum biosynthesis σ factor FliA and the stationary phase σ factor SigH from
Bacillus spp. Although they are not part of the ECF classification, these groups constitute the link
between the group 3 and group 4 (ECF) σ factors (9) and account for the quality of our clustering
approach. Other sequences with σ_3 domain remained unclassified (0.18% of the unclassified
sequences).

To identify subgroups with common characteristics, we performed an in-depth analysis of the
genomic context of ECFs in each subgroup, and aggregated subgroups into a total of 157 ECF groups.
For the definition of these ECF groups, the phylogenetic tree was manually split into monophyletic
clades, unless clades shared a similar genetic context and putative anti-σ factor type (Fig. 3B and
methods for more details). Genetic neighborhood and anti-σ factor type were evaluated in the subset of
ECFs that come from "representative" or "reference" genomes, as defined by NCBI
(https://www.ncbi.nlm.nih.gov), including only RefSeq genomes when both RefSeq and GenBank
assemblies exist. This helps to mitigate the bias towards commonly sequenced bacteria. As a result,
76.0% of the ECFs were captured in groups, displaying a median group size of 243 non-redundant
proteins (Fig. 3D). As an example, figure 3B shows a close-up view on 19 ECF subgroups within the
ECF tree, together with the proteins in their genetic neighborhood that feature >50% domain
architecture conservation (i.e., a combination of their Pfam domains). Here, it is evident that ECFs in
subgroups ECF02s1, ECF02s2 and ECF02s3 share a conserved genomic context with the anti-σ
factor RseA, and the regulators RseB and RseC for the former two subgroups, suggesting that ECFs in
these subgroups feature the same mode of regulation as RpoE from *E. coli* (belonging to ECF02
subgroup 1). Likewise, the subgroups aggregated into group ECF32 display strong conservation with a
2CS and a large number of genes encoding a type III secretion system (T3SS) (Fig. 3B). The
assignment of ECFs to ECF groups and subgroups can be found in table S1. These results underline
the previous notion that ECFs with close phylogenetic distance often share a conserved genomic
context, the gene products of which are typically involved in the regulation of ECF activity and/or direct
transcriptional targets of the ECF (1). This not only provides the basis for the definition of an ECF

group, but also helps to predict putative functions and regulatory mechanisms to ECF groups with no
experimentally described members (Table S2).

To provide a systematic overview on the conserved genomic context in each ECF group, we
analyzed the frequencies of genes with a conserved protein domain architecture encoded up- and
downstream of the ECF (Fig. 3C). For group ECF02, for instance, this reveals that downstream of the
regulators RseA-C there is enrichment of genes encoding translation regulators (e.g. EF-Tu), even
though the specific position of individual genes is less conserved (Fig. 3C). However, despite the
overall conservation of the genomic context within an ECF group, we often find subgroup-specific traits
with respect to the positioning and the specific type of conserved genes (Supplementary Text; Table
S2), clearly indicating that the definition of ECF subgroups is highly relevant to the biological function of
an ECF.

In addition to the conserved genomic context, ECFs often auto-regulate the expression of their
own genes, allowing bioinformatic prediction of their putative (sub)group-specific target promoters from
conserved bi-partite DNA elements upstream of the ECF-encoding operon (1, 28). When applying a
similar analysis to ECF subgroups and groups (Fig. 3B and C), we found overrepresented promoter
motifs in many groups, e.g. ECF02, while others did not show significant motifs, e.g. ECF32, consistent
with observations that the latter are not auto-regulated (20). Interestingly, even though predicted target
promoter motifs were not used in the definition of the ECF groups, split points that define ECF groups
(based on conserved genomic context) usually agree well with similar promoter elements (Fig. 3B).
However, as for the conservation of the genomic context, we sometimes find subgroup-specific
putative target promoters (e.g. in group ECF30 and others in supplementary text), highlighting the
added value of the fine-grained clustering approach taken here.

The definition of ECF groups based on genomic context conservation is further supported by the
statistical properties of the ECF subgroup tree, which typically displays high bootstrap support scores
at the rooting branches of ECF groups (Fig. 3B and E), indicating that these groups are robust with
respect to re-sampling of the original data set. To further check the performance of the new
classification approach, we created profile Hidden Markov models (HMMs) from the conserved σ_2 and
σ_4 domains of all sequences at the ECF group and subgroup level and tested whether these models
were capable of faithfully classifying ECF sequences from their own groups (Fig. 3G and Fig. S2). This
analysis showed that sequences were assigned to the correct ECF group in 99.3% of cases, while
assignment to the correct subgroup was successful in 94% of the cases. The lower performance of
subgroup-assignment was not surprising, given that neighboring ECF subgroups share higher
sequence similarity than neighboring ECF groups. These results confirm that the definition of ECF
groups and subgroups is based on a rational statistical approach and our findings suggest that the
used HMMs are specific and sensitive to allow for the classification of novel ECF σ factors in the future.
265

ECF classification 2.0 refines original and identifies novel ECF groups

As a proof of concept, we compared the original ECF classification and the classification presented in
this work. To this end, we created HMMs from the original ECF groups and used them to classify the
new ECFs gathered here. We saw that many of the sequences classified into a particular new group
were also classified into a particular original group (Fig. 4, ring #1), indicating that there is a broad
degree of correlation between the different classification approaches. Accordingly, for these groups of
high coherence we maintained the original group names to label ECF groups presented in this work.
Further in-depth analysis of the composition of the new groups revealed that 62 out of the 94 original
groups are preserved, 21 are merged into larger groups, five remain mainly ungrouped, three are
scattered across several subgroups, and three are present only in small percentages in some groups
(Table 1 and Fig. S3).

One case of an extremely scattered original group is ECF01 (Fig. S3). This group was already
considered highly diverse in the first ECF classification (1) and, based on the relatively unspecific HMM
model of this group, it acquired more sequences in subsequent classification efforts (4, 5). As a result,
we did not consider the proteins from ECF01 for the nomenclature of the ECF groups in this work.
Another highly scattered original group is ECF20 (Fig. S3). ECF20 is present in four main groups of our
classification: ECF281, ECF289, ECF290 and ECF291 (Table S2). ECF281, ECF290 and ECF291
seem to be related to heavy-metal stress, since their genetic neighborhoods contain a conserved
heavy-metal resistance protein in position +2 downstream of the ECF-encoding sequence in ECF281
and ECF290, and the full operon of a metal efflux pump in ECF291. This function of ECF291 has been
experimentally confirmed for CnrH in *Cupriavidus metallidurans* (ECF291s9) (29). Nevertheless, the
anti-σ factors encoded in the genetic context of members of these groups differ. ECF281 features a
zinc finger-containing anti-σ factor in position +1, while in the case of ECF289 this protein contains a
DUF3520 domain fused to a von Willebrand factor; ECF290 contains a RskA-like anti-σ factor, and,
lastly, ECF291 contains a CnrY-like anti-σ factor in position -2 (Supplementary Text, Table S2). Based
on this anti-σ factor diversity, it seems likely that the cognate ECFs are regulated in response to
different input stimuli, thereby warranting the definition of different ECF groups.

The last scattered group is ECF10. Even though minor parts of the original group ECF10 appear
across the new ECF classification, only groups ECF239 and ECF240 receive most of the proteins of
the original ECF10. Even though these two groups are both located within the large clade of FecI-like
ECFs (Fig. 4), members of ECF239 do not contain genes with a conserved carbohydrate-binding
domain in their neighborhood, a characteristic described for members of the original ECF10 (1). These
proteins have been suggested to be involved in carbohydrate degradation. However, experimental
confirmation for this biological function is still pending.

Even though scattered original groups are interesting, group merging events are more common.
Its high occurrence is probably due to the incorporation of new sequences that bridge previously
isolated ECF groups. Indeed, this possibility was considered in the founding ECF classification (1).
One example of a merged group is ECF243, which constitutes the largest group of the new
classification and contains the proteins previously associated to original FecI-like groups ECF05 to

ECF09 (Fig. S3). The reasons for merging were 1) members of these original groups form a
monophyletic clade in the ECF tree (Fig. 4) and 2) they contain a common genetic neighborhood with a
FecI-like anti-σ factor typically in position +1 from their coding sequence and a TonB-dependent
receptor in position +2 (Supplementary Text, Table S2). Another example of a merged group is
ECF238, which contains sequences from the original groups ECF24 and ECF44 (Fig. S3, Table S2).
Members of ECF238 contain a cysteine-rich C-terminal extension of approximately 20 amino acids
(Supplementary text), which is likely required for the activation of members of ECF238 when the
appropriate metal in the right redox state is present in the cytoplasm, as found for CorE2 from
Myxococcus xanthus (ECF238s15) (30).

What is likely the most interesting contribution of the new classification are the entirely new
groups. We found 22 new groups that could not be assigned to any original group (Table 2). Six of
these groups contain less than ten proteins from representative/reference organisms and, therefore,
were not further analyzed. From the analyzed groups, 10 share a conserved genetic neighborhood with
putative anti-σ factors. A special case of these is ECF241, which is located in the FecI-like clade and
represents an evolutionary intermediate between groups ECF240 (derived from original ECF10) and
the iron uptake FecI-like group ECF242, although ECF241 shows no significant similarity to any
original FecI-like group. Instead of the canonical FecR-like anti-σ factor from FecI-like groups,
members of ECF241 contain a conserved two-transmembrane helix protein in position +1 or -1 from
their coding sequence that in some cases hits the Pfam model for heavy-metal resistance proteins
(Pfam: PF13801). Given the lack of an anti-σ factor in a group within the FecI-like clade, we further
analyzed this conserved protein. Its N-terminus, the region that typically contains the ASD in anti-σ
factors, is not long enough to feature a typical ASD. However, a multiple sequence alignment of this
protein with the ASDs of canonical FecR-like anti-σ factors revealed that a putative, divergent ASD
might be located in the C-terminal cytoplasmic part of the conserved protein. To our knowledge, this is
the first time an anti-σ domain has been predicted C-terminally from transmembrane helices. The
second most common regulators of ECF activity in these new ECF groups are C-terminal extensions
(four out of 22), with groups ECF287 and ECF288, from Actinobacteria and Firmicutes, respectively,
containing cysteine-rich C-terminal extensions, and group ECF294 with a SnoaL-like extension (Table
S2, supplementary text). A potential regulator was not found for members of ECF201 and ECF282. In
the case of ECF282, the regulation could be carried out by a novel mechanism that involves
transcriptional regulation and ClpXP proteolysis, as explained below.

Taken together, the ECF groups presented in this work preserve many of the original groups,
expanding them with more proteins, and splitting or merging them in some cases. Here, we described
the new findings concerning the 22 new ECF groups with no significant similarity to any original group.
However, a full overview of all the ECF groups and their occurrence in different bacterial phyla is
shown in Fig. 5 and the full description of the groups is available in Supplementary Text.

ECF σ factors feature diverse, often multi-layered, modes of regulation

Given the large diversity of the ECF σ factor family, it is essential to focus on individual groups in order
to extract conclusions concerning their biological function, regulation and DNA binding site. Genetic
neighborhoods of ECF σ factors typically contain an anti-σ factor with a single transmembrane helix,
encoded in position +1 downstream of the ECF coding sequence. However, it is well known that other
regulatory elements might be substituting it, ranging from fused C-terminal extensions, to two-
component systems and serine/threonine protein kinases (1, 7, 23). Here, we provide an overview of
the different modes of regulation present across the groups in the present ECF classification. A
comprehensive description of all ECF groups can be found in Supplementary Text and in Table S2.

Most of the ECF groups (114 out of 157) contain a putative anti-σ factor, as defined by 1) the
presence of Pfam domains of known anti-σ factors, 2) detectable similarity to anti-σ factors of the
founding classification (1) and 3) presence of transmembrane helices (see Methods for details). This
anti-σ factor is typically encoded in position +1 from the ECF coding sequence. A list of putative anti-σ
factors identified in this study can be found in Table S4. In most of the cases, the putative anti-σ factor
does not match any Pfam domain of experimentally addressed anti-σ factors. In order to decipher
common types of anti-σ factors present across the ECF tree, we built HMMs from the cytoplasmic area
(when it harbors a conserved region representing a putative ASD) of the extracted putative anti-σ
factors that did not fit any Pfam domain. With these models, we searched the proteins encoded by 10
genes up- and downstream of the ECF coding sequence in all ECF groups (Fig. 6F). Interestingly, we
found that most of the anti-σ factors are ECF group-specific, in agreement with previous experimental
observations showing orthogonality between anti-σ factors of different groups (28). However,
exceptions are the clade that contains ECF222, ECF51, ECF38 and ECF39, which share the same
type of one-transmembrane helix anti-σ factor, groups regulated by FecR-like anti-σ factors, mostly
located in the FecI-like clade of the ECF tree (ECF239, ECF240, ECF242 and ECF243), groups
regulated by RskA-like anti-σ factors, anti-σ factors with a putative zinc-finger, and anti-σ factors with a
DUF4179, almost exclusively present in Firmicutes (Fig. 6F). The number of transmembrane helices
on putative anti-σ factors is usually one (82 ECF groups), followed by two (14 groups), six (five
groups), four (three groups) and three (one group) (Fig. 6E). Soluble anti-σ factors, as defined by the
absence of a predicted TM helix, are present in ten ECF groups (Fig. 6E). However, given that our
analysis can only identify soluble anti-σ factors with detectable similarity to existing anti-σ factors, it is
likely that other soluble anti-σ factor variants may exist. Additionally, even though we evaluated the
similarity of new ECF σ factors to known ASDs, it is not guaranteed that all the new putative anti-σ
factors function as such, given the vast diversity, lack of sequence conservation and lack of studies
confirming their function.

ECF107 contains two putative anti-σ factors, which could be part of the same protein complex or
compete for binding the ECF (6), thereby illustrating the complexity and diversity of anti-σ factor
mediated regulation. A second example is ECF102, whose only described member, SigX from *P.
aeruginosa*, has been suggested to have a role in mechanosensing (31). SigX is part of a seven-gene
operon which includes a mechanosensitive ion channel (CmpX) encoded in position -1, a putative anti-

σ factor (CfrX) encoded in position -2 and an outer membrane porin (OprF) encoded in position +1
(31). Even though original reports hypothesized that the regulation of SigX is carried out by the putative
anti-σ factor CfrX (6), new reports suggest that its regulatory mechanism is more complex and involves
also CmpX and OprF (31). We observed that these proteins are conserved in ECF102s1. Moreover,
the mechanosensitive ion channel is conserved in subgroups 2 and 5, which indicates a similar
regulation as members of subgroup 1. A similar case, in which proteins in addition to the anti-σ factor
are required for ECF regulation, is ECF31. The only characterized member of ECF31, SigY from *B.*
subtilis (subgroup 1), requires both the anti-σ factor YxlC, encoded in +1, and YxlD, encoded in +2, for
its regulation, presumably forming a protein complex with the ECF (32). YxlCD homologs are
conserved across ECF31.

The second most common regulatory mode of ECF σ factors is the presence of C-terminal protein
extensions to the ECF (19 groups) (Fig. 6B), which is typically correlated with the lack of putative anti-σ
factors (Fig. 6F). This agrees with the idea that the extension is substituting the anti-σ factor in the
regulation of the ECF (33). ECFs with the same type of C-terminal extension cluster together in the
same group, i.e., members of ECF42, with tetratricopeptide repeats in their extension, or in
neighboring groups, i.e. members of ECF41, ECF56, ECF115, ECF294, and ECF295, with SnoaL-like
C-terminal extensions. Given that only the core ECF domains were inputs of the ECF classification
process, this supports the notion that the extension interacts with the core ECF regions in a unique
manner depending on the type of domain that it bears (26). An interesting exception is ECF205, which
also has a SnoaL-like extension but is located in proximity to the base of the ECF tree (Fig. 4),
indicating that more factors, in addition to its C-terminal extension, determine the sequence
conservation of this group. Aside from the Pfam domains identified in C-terminal extensions of the
founding ECF classification (1, 4, 5), we identified a domain of unknown function (DUF1835) in
ECF264, an extension with five or seven transmembrane helices in ECF263, and a CGxxGxGxCxC
motif in ECF288.

Canonical C-terminal extensions are usually longer than 50aa, but we found that some groups
contain short C-terminal extensions difficult to identify when only looking at protein length. These
groups usually lack any other discernable means of regulation, which points towards the short
extension as a modulator of ECF activity. One of these groups is ECF238, which merges the original
groups ECF44 and ECF24. Members of ECF238 contain conserved cysteine residues in their short
(~20aa) C-terminal extension and also in the linker in some instances. One of the described members
of ECF238, CorE2 from *Myxococcus xanthus* (subgroup 15), is known to be activated by Cd^{2+} and Zn^{2+}
via this cysteine-rich C-terminal domain (30, 34). Another characterized member of ECF238 is SigZ
from *Bacillus subtilis* (subgroup 9). SigZ is not regulated by any anti-σ factor and the studies about its
function are very limited, since its deletion is not linked to any important phenotype (35). However,
association of SigZ with group ECF238 suggests that the two cysteine residues present in its C-
terminus could have a functional role. Another group with a short C-terminal extension is ECF29, which
contains a conserved RCE/D motif in its ~30 extra amino acids. Unfortunately, no member of ECF29

has been experimentally addressed, but the absence of a putative anti-σ factor similarly suggests a
regulatory role of this short extension.

N-terminal extensions of the ECF core regions occur less often, they are generally shorter than
canonical C-terminal extensions and they are prone to be overlooked whenever the ECF is translated
from non-canonical start codons. The only well-described N-terminal extension appears in ECF121
(Fig. 6C). This extension has been studied in BldN (subgroup 1) from *Streptomyces coelicolor*, where it
has to be proteolytically degraded to process the proprotein to its mature ECF, which then is subject to
anti-σ factor regulation (36). Nonetheless, subgroups from several groups contain N-terminal
extensions (Table S2). For instance, in ECF36s4, represented by SigC from *M. tuberculosis,* the N-
terminal extension has been proposed to inhibit the DNA contact in the uninduced state, since
members of this subgroup lack of an obvious anti-σ factor (37). Alternatively, the N-terminal extension
of two members of ECF12s1, σ^R from *S. coelicolor* and SigH from *Mycobacterium smegmatis*,
generates an unstable isoform produced from an earlier start codon upon exposure to thiol oxidants
(38). This makes σ^R susceptible towards σ^R-activated ClpP1/P2 proteases and thus implements a
negative feedback loop that contributes to turning off the stress response (38).

Other putative regulators of σ factor activity that we often found in the conserved genetic
neighborhood of ECFs were serine/threonine protein kinases (Fig. 6D). ECF σ factors of five original
groups have been hypothesized to be directly phosphorylated by a protein kinase (ECF43 and ECF59-
ECF62 (1, 4)). We add to the list of protein kinase-associated groups ECF217, ECF267 and ECF283
(Fig. 6D). Other groups such as ECF40, ECF27 or ECF210 contain protein kinases only in certain
subgroups. Proteins from original group ECF60 were not classified by the pipeline since only eight
members of ECF60 were extracted. Protein kinase-related ECF groups typically lack co-encoded anti-
σ factors (Fig. 6D), consistent with the notion that direct phosphorylation of the σ factor regulates its
ability to interact with RNAP and/or the promoter DNA. The only exception is group ECF267, which
contains a putative FecR-like anti-σ factor with tetratricopeptide repeats in position +1. Given that
ECFs from group ECF267 are very distant from the FecI-like clade (ECF239-ECF243) (Fig. 4), it
seems possible that this anti-σ factor does not target members of ECF267, but other FecI-like ECFs.
However, none of the organisms that contain members of ECF267 contain any FecI-like ECF σ factor.
Whether the anti-σ factor and/or the protein kinase regulate the activity of members of ECF267 is
unclear.

A total of four groups contain two-component systems in their genetic context. These regulators
can co-occur in conjunction with anti-σ factors, as in the case of ECF15 and ECF246, or not, as in
ECF32, ECF234 and subgroups 1, 2 and 3 of ECF39. These possibilities reflect the different regulatory
mechanisms exerted by two-component systems. On one hand, members of ECF15, the main general
stress response σ factors in alphaproteobacteria, are activated by a partner-switching mechanism
where an anti-anti-σ factor sequesters the cognate anti-σ factor, thereby releasing the ECF (15). The
anti-anti-σ factor is fused to the response regulator of a two-component system, also encoded in the
same genetic context, and is able to bind the anti-σ factor when phosphorylated (15). This is unlikely

the case for members of ECF246, since their response regulator is fused to a transcriptional regulator.
Future analysis of members ECF246 could determine whether the putative anti-σ factor and/or the two-
component system encoded in their genetic context regulates ECF activity. For members of ECF32, it
was indeed shown that the two-component system indirectly regulates transcription of the ECF σ factor
by inducing the expression of the transcription factors HrpSR (20, 40, 41). Members of ECF32 in turn
activate the synthesis of the type III secretion system *hrp*, required for plant infection (20, 41). In the
case of ECF39, 2CSs are directly regulating the transcription of the ECF, as described for SigE from *S.*
coelicolor and σ^{25} from *Streptomyces avermitilis* (42, 43). Members of this group are involved in
antibiotic synthesis and cell-wall stress resistance (42, 43). This direct regulation of the 2CS over the
ECF expression could also occur in members of ECF234, given the absence of a putative anti-σ factor
and the fusion of the response regulator to a transcriptional regulator. The physiological function of
ECF234 seems to be related to an ABC transporter present in its genetic context, but the substrate of
the transporter is unknown.

On top of these proteins, we found that some ECF groups contain conserved transcriptional
regulators encoded in their genetic contexts, such as TetR-like repressors, which appear in groups with
anti-σ factors (ECF125) and, remarkably, in ECF203, which lacks any obvious regulator (Fig. 6D).
Given the lack of characterized members of ECF203, it is unclear whether this TetR repressor
regulates the expression of members of ECF203 or is part of their response. In favor of the former,
members of ECF203 do not seem to be auto-regulated, given the lack of a conserved (predicted)
target promoter motif (Table S2). Other transcriptional regulators include LysR- and MerR-like
repressors, which appear in several ECF groups associated with anti-σ factors (Fig. 6D).

A total of 16 ECF groups are not linked to any of the above-mentioned regulators (Fig. 7), inspiring
us to predict novel, putative regulators of ECF activity. So far, only three of the 16 groups have
experimentally addressed members, namely ECF228, ECF282 and ECF114. SigP from
Porphyromonas gingivalis (ECF228s7) is only present in measurable concentrations when stabilized
by direct interaction with the response regulator PorX from the two-component system PorXY (44).
Even though response regulators are not conserved in the genetic context of members of ECF228
(Fig. 7), it is possible that members of ECF228 are unstable and require other proteins as chaperons.
In the case of the novel group ECF282, σ^{AntA} from *Streptomyces albus* (subgroup 2) is regulated at the
level of transcription and might be target of ClpXP proteolysis (45). Indeed, homologs of σ^{AntA} have
been considered a new group of ECF σ factors that control the expression of antimycins (45). Even
though the C-terminal AA dipeptide, suggested as target of ClpXP proteolysis (45), is only present in
members of subgroup 2, members of other ECF282 subgroups could be regulated in a similar manner,
since different ClpXP proteases have different binding specificities (46). In ECF114, SigH from
Porphyromonas gingivalis (subgroup 4) plays a role in aerotolerance. SigH it is induced upon exposure
to O_2 and promotes oxidative stress protection and hemin uptake (47). Even though it is speculated
that SigH is transcriptionally activated (47), no transcription factor in charge of this task has been
identified.

Given that the genetic neighborhood of the remaining 13 groups does not feature canonical ECF
regulators, but other conserved elements (Fig. 7), we speculated about their putative function.
However, a general issue of this analysis is that it is hard to discriminate whether these elements are
regulators and/or targets of ECF activity, suggesting that both options should be considered in
downstream experimental analyses. Interestingly, we found new putative regulators/targets of
regulation of the original groups ECF54 and ECF130. ECF54 is encoded in close proximity to a protein
with a 4Fe-4S cluster, whereas ECF130 is encoded in proximity to a helix-turn-helix (HTH) containing
protein (Fig. 7). Since HTH motifs are usually related to DNA binding, this protein could be a new type
of transcriptional factor involved in transcriptional control of members of ECF130. A similar case is
found in the new group ECF201, which is usually co-encoded with HTH proteins in position -1 (Fig. 7).
Another interesting case is found in members of ECF237, which contain several "killing trait" proteins
(Pfam: PF11757) in their vicinity (Fig. 7). These domains were described for RebB, one of the three
proteins necessary for the assembly of R-bodies in the Paramecium endosymbiont *Caedibacter*
taeniospiralis (48). Given the absence of conservation for the rest of the proteins from the R-body and
the presence of several copies of the killer domain in members of ECF237 (which are mainly present in
Bacteroidetes unrelated to the alphaproteobacterium *C. taeniospirails*), it is possible that proteins with
this domain have an alternative function, not related to R-body assembly, but potentially involved in
controlling ECF activity. Lastly, members of ECF286 and ECF292 share genetic neighborhood with
several copies of Asp23 proteins (Pfam: PF03780) (Fig. 7). Asp23 is one of the most abundant
proteins of *Staphylococcus aureus* and its deletion leads to upregulation of the cell wall stress
response (49). Therefore, Asp23 proteins could be acting as a new type of anti-σ factor that regulate
the activity of members of ECF286 and ECF292.

Discussion

This work unifies, refines and greatly expands previous ECF classification efforts. Thanks to its two-
tiered clustering approach, it provides a high-resolution view of the ECF family. ECF subgroups,
composed of closely related proteins, are further hierarchically clustered into 157 ECF groups, defined
on the basis of a common genetic neighborhood, which indicates a similar mode of regulation. As part
of the *in silico* characterization of ECF groups, we predicted their putative regulators, their target
promoter motifs and their most likely function (Supplementary Text 1 and Table S2). As already
observed for the previous classification, these predictions are biologically meaningful in that they
correctly reflect results of experimentally studied members, whenever available. The comprehensive
description of the ECF groups serves as a source of testable hypotheses that will support the
experimental description of new ECFs, which will lead, in turn, to more precise and detailed group
descriptions.

Furthermore, the fine-grained resolution of the ECF family and the great enrichment in
phylogenetically diverse proteins allows for the application of *in silico* prediction tools for individual
groups. These types of analyses have led to the finding of functional differences between the C-

terminal extensions from ECF41 and ECF42 (26), where C-termini from ECF41 had an inhibitory role over ECF activity, while ECF42's were essential for transcriptional activation (26). These results were only possible due to the large number of sequences present in these groups, with >12,000 and >10,000 in ECF41 and ECF42, respectively (26).

The new ECF classification presented in this work has changes with respect to the founding classifications. Even though 62 of the 94 original groups are preserved, 21 are merged, five were ungrouped, and three each were scattered or present in the new classification but composing only small parts of their new group (Table 1). The new ECF groups are monophyletic clades of the ECF phylogenetic tree, that are subdivided into hierarchically-distributed ECF subgroups. This high-resolution, comprehensive classification provides advantages with respect to partial updates. One example comes from ECF54 and ECF58, identified in two different works (4, 5) and in two phyla, Actinobacteria and Planctomycetes, respectively. Within our ECF tree, these two groups are direct neighbors with a bootstrap support value of 17, indicating a significant protein similarity between them. None of them has a putative anti-σ factor or any other clear regulator, and they contain different elements in their genetic context (Fig. 7). These results suggest that ECF58 and ECF54 have the same origin, but they evolved independently in Actinobacteria and Planctomycetes, acquiring the regulation of different genes in their genetic neighborhood. What remains unclear is whether the regulation of members of ECF54 and ECF58 has common features, as expected for ECFs with a common origin.

As part of the description of ECF groups, we analyzed their most likely regulators and the types of putative anti-σ factors encoded in their genetic neighborhood (Fig. 6). Most of the predicted anti-σ factors are highly specific for their own groups (Fig. 6F). Exceptions occur in neighboring ECF groups, e.g. in the FecI-like clade (ECF239 to ECF243) or in the clade formed by groups ECF214, ECF18 and ECF19, indicating co-evolution between ECF and anti-σ factor sequences. The general lack of the same type of anti-σ factors in neighboring groups reflects their large diversity and their specificity, which has been exploited for the construction of orthogonal genetic circuits (28). Anti-σ factors are not the only genes conserved in the genetic context of ECF σ factors. In this study, we identified the ECF groups associated to other known ECF regulators such a C-terminal and N-terminal extensions, two-component systems, serine/threonine kinases (7), and other regulators such as TetR repressors (Fig. 6).

One important insight of this work is that ECF groups controlled by several regulatory layers are more common than originally thought. For instance, members of ECF121 are dually regulated by anti-σ factors and N-terminal extensions, some members of ECF12 are regulated by both anti-σ factors and alternative promoters that generate an unstable longer versions of the ECF σ factor (38) and members of ECF18 and ECF19 are not only regulated by RskA-like anti-σ factors, but also by a pair of conserved cysteine residues known to form a disulphide bridge that senses oxidative stress in SigK from *M. tuberculosis* (ECF19s1) (50). While these regulatory layers have only been deciphered for a few well-studied ECF σ factors, they might point towards a broader means of regulation also

571   implemented in additional ECF groups. For instance, several ECF groups feature conserved cysteine
572   residues potentially able to form disulphide bridges (Table S2), and members of ECF267 contain both
573   a FecR-like anti-σ factor and a conserved protein kinase in their genomic neighborhood. Given their
574   multi-layered regulation, abundance and diversity, we suggest that ECF σ factors have higher signal
575   integration capabilities than previously anticipated.

576   With an average of approx. 10 ECFs per genome, these regulators are more abundant than
previously thought (1). Confirming previous reports (1, 5, 51), we find that the number of ECFs is
proportional to genome size (Fig. S1), with species thriving in diverse environments typically featuring
larger genomes that provide them with the ability to sense and respond to a variety of external signals.
One example is the bacterium *Sorangium cellulosum* So0157-2, which features a genome that is more
than 1Mbp larger than its close relative *S. cellulosum* So ce56, allowing the former to adapt to alkaline
conditions (51). Accordingly, the number of ECFs in *S. cellulosum* So0157-2 (82 ECFs) is significantly
larger than in *S. cellulosum* So ce56 (70 ECFs), emphasizing the increased regulatory capacity
incurred by genome expansion. Among the ECF groups acquired exclusively in *S. cellulosum* So0157-
2, we found ECF03 (one extra member), ECF26 (one extra member), ECF41 (two extra members) and
ECF56 (one extra member). ECF03 and ECF26 are novel acquisitions present in *S. cellulosum*
So0157-2 but not in *S. cellulosum* So ce56. Indeed, members of ECF03 are mainly present in
Bacteroidetes (Table S2), and could have been acquired by horizontal gene transfer. However, this
protein is not overexpressed under alkaline conditions (51), indicating that this ECF is either not
autoregulated, or not responsible for alkaline resistance in *S. cellulosum* So0157-2. In contrast, the
additional member of ECF26 contained in *S. cellulosum* So0157-2 is overexpressed at pH 10 (51) and
could therefore be part of the alkaline resistance observed for *S. cellulosum* So0157-2. This ECF
belongs to ECF26s1, which shares a conserved genetic neighborhood with a catalase (-1 from the
ECF coding sequence) and a cytochrome b561 (position -2). Whether ECF26 or any other of these
ECFs provides *S. cellulosum* So0157-2 with alkaline resistance needs further investigation.

596   The search of ECF σ factors presented in this work has some limitations related to the quality
filters that we applied during the ECF retrieval. These filters limit the diversity of the extracted
sequences, while ensuring that the collected proteins function as real ECF σ factors. In particular, we
noticed that two main types of ECF σ factors could not be captured, namely, ECF σ factors from
phages and ECFs whose conserved σ_2 and σ_4 domains are divergent. σ factors of phage origin have
been described in literature (see review (52)); nevertheless, they are usually divergent from traditional
σ factors (52), in some cases incorporating alternative domains replacing σ core domains. For
instance, in *Bacillus* phage vB_BceM-HSE3, the ECF Gp17 contains a double zinc ribbon domain
(Pfam: PF12773) in the position of the σ_2 domain, while the generic Pfam domain for σ_4 is not found at
all (53). Similarly, the σ factors Gp01 and Gp103 contain only σ_2 domain or no Pfam domain,
respectively (53). Another reason for the lack of phage proteins in the present work is that viral
genomes are usually not annotated in NCBI (54) and did not enter the ECF search in most of the
cases. Other types of ECF-like σ factors not included in the current version are ECFs whose σ_4 (e.g.

SigI from *Bacillus subtilis*) or σ_2 domain (such as VP0055 from *Vibrio parahaemolyticus* or ComX from

Streptococcus pneumoniae) do not hit their Pfam models. A special example of this are σ^I-like ECFs,

which contain a σ_{I-C} domain instead of a canonical σ_4 domain (55). These ECFs are involved in the

synthesis of components of the cellulosome in cellulolytic clostridia (55). Attempts to classify these

proteins against the current ECF classification were unsuccessful. The group with the highest

probability of containing σ^I-like ECFs is ECF201 (probability = 1.12e-19), the outermost group of the

ECF classification, indicating that σ^I-like ECFs are distant from canonical ECFs and might have

evolved in parallel to them from group 3 σ^{70} factors. Lowering the stringency in our extraction pipeline

would allow to study these non-canonical ECFs in further depth.

The average number of identified ECF σ factors per genome varies for different groups and

bacterial phyla (Fig. 5). In this analysis it turned out that the number of unclassified ECFs per organism

is larger in bacterial phyla underrepresented in biological databases (Fig. 5). In the future, clustering

strategies could specifically target these proteins, which are likely too diverse and scarce to be

clustered with the currently available dataset. We also found that some ECF groups are particularly

enriched in certain phyla. For instance, we observed an average of 5.3 copies of ECF57 per

planctomycetal genome, and an average of 3.4 copies of ECF240 per Bacteroidetes genome (Fig. 5).

In these cases, a question that remains unsolved is whether the function of members of the same

group is redundant in the same organism, or they rather hold specialized functions. In favor of the

latter, members of ECF240, which inherits most of its characteristics from the original group ECF10,

have been associated to carbohydrate scavenging in Bacteroidetes (1). Even though no member has

been experimentally addressed to date, it is possible that the different members of ECF240 present in

the same genome are involved in the uptake of different carbohydrates. A similar case occurs in the

proteobacterial group ECF243, which merges original FecI-like groups ECF05-09, and is in charge of

iron uptake (1, 56). We found an average of 1.13 members of ECF243 per proteobacterial genome.

However, under closer inspection only 33% of the proteobacterial genomes contain members of

ECF243, indicating that, when present, members of ECF243s are duplicated and appear in 3.4

proteins per organism on average. Interestingly, only 8.9% of the organisms contain ECF243s from the

same subgroup, suggesting that different subgroups fulfill different physiological functions. One

possibility is that members of different subgroups detect signals from different FecR-like anti-σ factors,

which in turn, detect the presence of iron-siderophore complexes from different FecA transporters (see

(56) for a review). Future analyses have to answer whether the different members of the same ECF

group in the same genome have acquired different specificities and whether this specificity is a general

feature of ECF σ factors.

In summary, the updated ECF classification presented in this work serves as a detailed source of

testable hypotheses to guide the experimental characterization of this important class of bacterial

regulators. The ECF classification comes together with a full description of ECF groups, including the

putative group-specific regulators of ECF activity, conserved proteins encoded in the same genetic

neighborhood, and predicted target promoter motifs (Supplementary Text and Table S2). Collectively,

this information allows prediction of the potential function of the members of the group, which is verified
by experimentally described members whenever they are available (Supplementary Text). Moreover,
our hierarchical two-level classification provides a broad sequence collection with an appropriate
degree of similarity (or variability) required for *in silico* prediction tools that employ sequence variation-
based algorithms, such as co-variation-based prediction of protein-protein interactions or structural
predictions.

Methods

General bioinformatic tools

The data was processed with custom scripts written in Python when nothing else is stated (available
upon request from the corresponding author). Multiple Sequence Alignments (MSAs) were generated
by Clustal Omega 1.2.3 (57) and were visualized in CLC Main Workbench 7.7.2 (QUIAGEN®).
HMMER suite 3.1b2 (58) was utilized for generating and employing Hidden Markov Models (HMMs).
HMMs were produced by hmmbuild (58) and proteins were searched against HMMs using hmmscan.
The maximum-likelihood phylogenetic tree of ECF subgroups and bootstrap values were retrieved by
IQ-TREE 1.5.5 with default options and automatic model selection (59). Phylogenetic trees were
visualized using iTOL (60). Transmembrane helix predictions were carried out with TopCons (61) and
PRED_TMBB2 for outer membrane proteins (62).

ECF retrieval

The full retrieval process is depicted in Figure 1. The sequence between the start of σ_2 and the end of
σ_4 domains (core ECF region) was extracted from the MSA of the 3,755 ECFs from the original library
(1, 4, 5) (position 912 and 2059 in the MSA) and was used to build the general HMM for ECF
sequence retrieval. Then, ECFs from the original library and proteins with a σ_3 domain (Pfam:
PF04539) retrieved from Pfam (release 31.0 (63)) were scored against this model. The resulting bit-
scores were used to construct a Receiver Operator Characteristic (ROC) curve (using the function
"roc_curve" from the "scikit-learn" Python package (64)) in order to select the threshold score able to
capture the greatest number of ECFs (highest sensitivity), while minimizing the number of σ_3-
containing proteins (highest specificity). This resulted in an optimum bit-score threshold of 60.8, a
sensitivity of 0.92, a specificity of 0.98 and an Area Under the Curve (AUC) of 0.98, accounting for a
robust performance of the classifier. Then, proteins with a score higher than the optimum threshold
were selected as putative ECF σ factors for the next steps. For this search, we considered all protein
sequences from genomes with annotation (.gff), protein (.faa) and genome (.fna) file available in NCBI
(as of February 2017), including RefSeq and GenBank entries, accounting for a total of 156,241
genomes and 554,108,437 proteins. Subsequent quality controls were applied to ensure that all the
putative ECFs contain σ_2 (Pfam: PF04542) and σ_4 (Pfam: PF04545 and PF08281) domains and lack
the σ_3 domain using Pfam HMM profiles (hmmsearch with default settings). Only the non-redundant
proteins without ambiguous amino acids were considered further. Sequence redundancy was

assessed with Cd-hit (65) at 100% sequence identity. In summary, from the 554,108,437 annotated
proteins, 714,848 had a positive score against the ECF model. Of those, 177,910 had σ_2 and σ_4
domains, lacked σ_3, lacked non-amino acidic symbols and were non-redundant, constituting the
extended ECF library used in the next steps.

Within this dataset we found that 1,217 proteins generated low-scoring hits against the model of
the σ_3 domain (Pfam: PF04539) when hmmscan was applied for σ_3's HMM alone. This is likely due to
the size of the ECF database, where E-values are less significant than for smaller datasets (see
HMMER documentation (58)). Given that the Pfam HMM of σ_3 wrongly hits the σ_2 or σ_4 domain in
some cases, only the proteins where the highest scoring σ_2, σ_3 and σ_4 domains are not overlapping are
considered as σ_3-containing proteins. After their identification, the 1,217 σ_3-containing proteins were
used as outliers for the clustering, since they are the closest σ factors to ECFs.

The average number of ECFs per genome was computed considering only the 12,539 ECFs from
the 1,234 complete genomes tagged as 'representative' or 'reference' in NCBI, giving priority to
RefSeq genomes over GenBank if both exist for the same organism, unless stated otherwise.

Clustering and ECF group definition

Non-redundant ECF sequences were stripped to σ_2 and σ_4 regions using hmmscan (HMMER suite
3.1b2 (58)) and Pfam models for σ_2 and σ_4. We selected the 'envelope' region of the hit with the lowest
(i.e. best) E-value. For the first step of clustering, we applied MMseqs2 (27) with default parameters.
However, the phylogenetic distance between pairs of sequences within clusters, as calculated from the
k-tuple distance (66) obtained from Clustal Omega 1.2.3. with options '--full, --full-iter and --distmat-out'
(57), was large in some cases (Fig. S4), indicating imperfect clustering at this stage. To solve this
issue, MMseqs2 clusters were split using a bisecting k-Means algorithm until the maximum pairwise
distance between sequences of the clusters was ≤ 0.6. This threshold was chosen since it was the
largest maximum k-tuple distance whose associated clusters contained proteins similar enough to
produce homogeneous MSAs. The resulting 2,380 clusters with more than 10 sequences are defined
as ECF subgroups. A total of 137,452 ECFs (77.26%) were classified into subgroups. Then, the
consensus sequences of subgroups (computed with Biopython (67)) were used for the construction of
a phylogenetic tree. As outliers of the tree we included the consensus of a MSA calculated from all
proteins included in Pfam (release 31.0 (63)) that contain all of the three domains σ_2, σ_3 and σ_4. We
also included the 1,217 closest proteins to ECF σ factors with σ_3 domain. The subsequent phylogenetic
tree was manually split into monophyletic clades using a divisive strategy, where two clades were kept
together in the same ECF group unless the genetic context or the putative anti-σ factor (when present)
differed. ECF groups featuring only a single subgroup were only assigned if they have significant
similarity to an original ECF group, while all other single subgroups were maintained as singleton
subgroups (for nomenclature see next paragraph). This strategy resulted in 157 ECF groups that
contained 135,259 ECF σ factors, corresponding to 76.03% of the new ECF library. For a graphical
summary of the clustering strategy, see Fig. 3.
721

Nomenclature

Names of original ECF groups are maintained for groups with the same characteristics. When several original groups are represented in an ECF group or the ECF group has no significant similarity to any original group, the name of this ECF group follows the pattern ECF2XX, standing for ECF classification 2.0, where XX is a running number assigned according to the position in the phylogenetic tree (Fig. 4). For instance, ECF201 is closer to the base of the tree than ECF260. Subgroups are referred with the name of the ECF group they are part of, followed by 's', standing for subgroup, followed by a running number that increases for decreasing subgroup size. For instance, subgroups ECF02s1 (ECF02 subgroup 1) and ECF02s2 (ECF02 subgroup 2) are both part of group ECF02, and s1 contains more non-redundant proteins than s2. Subgroups that are not part of any ECF group are named 'ECFs' followed by a running number according to their position in the phylogenetic tree.

Clustering validation

Groups and subgroups were evaluated according to the performance of their HMMs, built from the concatenated σ_2 and σ_4 domains of the constituting protein sequences. These HMMs were used to score proteins from all groups or subgroups. Bit-scores below the reporting threshold were considered to be equal to 0. The average score of members of each group against each HMM was normalized by dividing by the score of the group against its own HMM. The average normalized scores are plotted in a heatmap (Fig. S2A). We validated the subgroups by generating 100 randomly permutated sets of proteins with the same size distribution as the subgroups (Fig. 3F). The mean average k-tuple distance in the permutated data was 0.79 ± 0.01, whereas this value was 0.29 ± 0.11 for ECF subgroups. The difference between the distributions of average pairwise k-tuple distances of ECF subgroups and permutated clusters was statistically significant (two-tailed Student's t-test p-value < 1e-16). Furthermore, we evaluated the support of the branches of the phylogenetic tree by running 100 bootstrap replicates, as implemented in IQ-TREE (59). As a further plausibility test, we verified the agreement between original and new classification (Fig. 4).

Classification of proteins against ECF clusters

In order to classify proteins against original groups and new groups/subgroups, we derived two bit-score cut-offs for each group/subgroup: (i) The trusted cut-off is defined as the minimum bit-score achieved by a true member of a group/subgroup, whereas (ii) the noise cut-off is the maximum bit-score achieved by all ECFs that are non-members of this group/subgroup. Trusted and noise cut-offs provide the first step in the evaluation of the membership of a protein against a group/subgroup. Then, we obtained the probability that a protein belongs to a cluster using a logistic fit, as described in (68). First, we represented in the x-axis the bit-score and in the y-axis the probability that a bit-score is produced by a real member of the group/subgroup under evaluation, this is 1 for members and 0 for non-members. We fitted the data to a logistic regression describing the transition of the bit scores from non-members to members of a cluster (Fig. S5). The logistic function describing the probability $P(q, i)$ of protein sequence q belonging to group/subgroup i reads

759
$$P(q,i) = \frac{1}{1+e^{-k_i \times \frac{B_{qi}-\overline{B_i}}{\sigma(B_i)} - c_i}}$$
(1)

with B_{qi} being the bit-score obtained for the protein sequence q scored against the HMM model of

group/subgroup i and B_i and $\sigma(B_i)$ being the mean and standard deviation of all bit-scores obtained for

member sequences of group/subgroup i, respectively. The parameters k_i and c_i were fitted using the

function 'curve_fit' (with a Levenberg-Marquardt algorithm) from the Python package 'scipy.optimise'.

These parameter values are available for groups, subgroups and original groups (Table S5). Some

small subgroups could not to be fitted. In that case, the probability is considered 1 if the protein has a

bit score larger that the trusted cut-off and it does not match any other subgroup, or 0 otherwise. The

selection of a unique probability threshold was carried out with a ROC curve, using the members of the

group/subgroup as true positives and members of all other groups/subgroups as true negatives. This

probability threshold, together with trusted and noise cut-offs, and fitting parameters of the logistic

curve are available for groups, subgroups and original groups (Table S5).

In the classification process itself only protein sequences stripped to their σ_2 and σ_4 domains were

used. The regions of these domains were defined by the HMMER envelope region of the highest

scoring hit of their respective Pfam models. In a first step of the classification only the

groups/subgroups against which the bit-score of a protein is higher than the trusted cut-off (or the noise

cut-off if none exceeds the trusted cut-off) are considered as putative groups/subgroups. In a second

step, the protein is assigned to the cluster with the highest probability, calculated with the logistic curve

of each cluster, only if this probability is higher than the ROC-optimized probability threshold.

Promoter motif prediction

Since ECFs often auto-regulate their own transcription, their putative target promoters can be predicted

from the regions upstream of the operon in which the ECF is encoded. To this end we searched for

conserved bipartite nucleotide motifs in the 200 base pairs (bp) upstream of the ECF operon, by

executing BioProspector (69) with the parameters '-W 7 -w 5 -G 18 -g 15 -n 150 -d 1', as previously

described (28). Operons were considered to be the set of coding sequences transcribed in the same

direction as the ECF and with an intergenic distance shorter than 50 bp. Only the highest scoring motif

of each input sequences was further considered. The region containing the -35 and -10 motifs in

addition to ±10 bp up- and downstream is represented in a sequence logo produced by WebLogo 3

(70).

ECF group analysis

For the analysis of ECF group characteristics we included only proteins from 'representative' and

'reference' genomes as defined by NCBI (https://www.ncbi.nlm.nih.gov), thereby reducing the bias

towards frequently sequenced organisms. Only RefSeq assemblies were considered when both

RefSeq and GenBank assemblies are available. To define coherent ECF groups and to elucidate the

putative function of members of each group, we analyzed the protein domain composition of the

proteins encoded in a distance of ±10 coding sequences from the ECFs coding sequence. First, we

queried these proteins against the HMMs of Pfam 31.0 (63). For every protein, we only considered the
non-overlapping Pfam domains with the lowest (best) E-value, leading to a set of specific domains
(which we define as the 'domain architecture') for each protein in the genomic neighborhood of the
ECF. For each ECF subgroup we analyzed the conservation of the domain architecture in specific
positions up- and downstream of the ECF. A domain architecture is defined as conserved if it appears
in more than 75% of the genomic contexts of an ECF subgroup. To avoid biases due to low number of
ECFs, we only analyzed the genetic context of subgroups with more than 10 ECFs.

For anti-σ factor identification we used (i) Pfam domains of known anti-σ factors, (ii) detectable
sequence similarity to anti-σ factors of the founding classification (1) and (iii) presence of
transmembrane helices, as described in the following. Most of the anti-σ factors cannot be predicted
due to the lack of Pfam domains that describe them. Therefore, we used the anti-σ factors retrieved by
(1) as the database to query candidate anti-σ factors using BLAST (71) with an E-value < 0.01.
Moreover, since anti-σ factors are usually transmembrane proteins, we predicted the presence of
transmembrane helices using the consensus prediction of TopCons (61). In cases where the presence
of the transmembrane helix was not clear, we aligned the sequences of the putative anti-σ factors to
determine the presence of conserved hydrophobic regions likely to be transmembrane domains. Since
anti-σ factors are usually located in positions ±2 from the ECF coding sequence, those were the main
positions we focused our search on.

References

1. Staroń,A., Sofia,H.J., Dietrich,S., Ulrich,L.E., Liesegang,H. and Mascher,T. (2009) The third pillar of
bacterial signal transduction: classification of the extracytoplasmic function (ECF) σ factor protein
family. *Mol. Microbiol.*, **74**, 557–581.

2. Helmann,J.D. (2002) The extracytoplasmic function (ECF) sigma factors. *Adv. Microb. Physiol.*,
10.1016/S0065-2911(02)46002-X.

3. Paget,M.S.B. and Helmann,J.D. (2003) The σ70 family of sigma factors. *Genome Biol.*, 10.1186/gb-
2003-4-1-203.

4. Jogler,C., Waldmann,J., Huang,X., Jogler,M., Glöckner,F.O., Mascher,T. and Kolter,R. (2012)
Identification of proteins likely to be involved in morphogenesis, cell division, and signal
transduction in Planctomycetes by comparative genomics. *J. Bacteriol.*, **194**, 6419–30.

5. Huang,X., Pinto,D., Fritz,G. and Mascher,T. (2015) Environmental Sensing in Actinobacteria: a
Comprehensive Survey on the Signaling Capacity of This Phylum. *J. Bacteriol.*, **197**, 2517–2535.

6. Pinto,D. and Mascher,T. (2016) The ECF Classification: A Phylogenetic Reflection of the Regulatory
Diversity in the Extracytoplasmic Function σ Factor Protein Family. In *Stress and Environmental*
Regulation of Gene Expression and Adaptation in Bacteria.

7. Mascher,T. (2013) Signaling diversity and evolution of extracytoplasmic function (ECF) σ factors.
Curr. Opin. Microbiol., **16**, 148–155.

8. Paget,M.S. (2015) Bacterial Sigma Factors and Anti-Sigma Factors: Structure, Function and

Distribution. *Biomolecules*, **5**, 1245–65.

9. Lonetto,M., Gribskov,M. and Gross,C.A. (1992) The sigma 70 family: sequence conservation and
evolutionary relationships. *J. Bacteriol.*, **174**, 3843–9.

10. Lane,W.J. and Darst,S.A. (2006) The structural basis for promoter -35 element recognition by the
group IV sigma factors. *PLoS Biol.*, **4**, e269.

11. Campagne,S., Marsh,M.E., Capitani,G., Vorholt,J.A. and Allain,F.H.T. (2014) Structural basis for -
10 promoter element melting by environmentally induced sigma factors. *Nat. Struct. Mol. Biol.*,
10.1038/nsmb.2777.

12. Grosse,C., Friedrich,S. and Nies,D.H. (2007) Contribution of extracytoplasmic function sigma
factors to transition metal homeostasis in Cupriavidus metallidurans strain CH34. *J. Mol.*
Microbiol. Biotechnol., **12**, 227–40.

13. Mascher,T., Hachmann,A.B. and Helmann,J.D. (2007) Regulatory overlap and functional
redundancy among Bacillus subtilis extracytoplasmic function σ factors. *J. Bacteriol.*,
10.1128/JB.00904-07.

14. Staroń,A. and Mascher,T. (2010) General stress response in a-proteobacteria: PhyR and beyond.
Mol. Microbiol., 10.1111/j.1365-2958.2010.07336.x.

15. Francez-Charlot,A., Frunzke,J., Reichen,C., Ebneter,J.Z., Gourion,B. and Vorholt,J.A. (2009)
Sigma factor mimicry involved in regulation of general stress response. *Proc. Natl. Acad. Sci.*,
10.1073/pnas.0810291106.

16. Llamas,M.A., Van Der Sar,A., Chu,B.C.H., Sparrius,M., Vogel,H.J. and Bitter,W. (2009) A novel
extracytoplasmic function (ECF) sigma factor regulates virulence in Pseudomonas aeruginosa.
PLoS Pathog., 10.1371/journal.ppat.1000572.

17. Kazmierczak,M.J., Wiedmann,M. and Boor,K.J. (2005) Alternative Sigma Factors and Their Roles
in Bacterial Virulence. *Microbiol. Mol. Biol. Rev.*, 10.1128/mmbr.69.4.527-543.2005.

18. Campbell,E.A., Greenwell,R., Anthony,J.R., Wang,S., Lim,L., Das,K., Sofia,H.J., Donohue,T.J. and
Darst,S.A. (2007) A conserved structural module regulates transcriptional responses to diverse
stress signals in bacteria. *Mol. Cell*, **27**, 793–805.

19. Li,W., Stevenson,C.E.M., Burton,N., Jakimowicz,P., Paget,M.S.B., Buttner,M.J., Lawson,D.M. and
Kleanthous,C. (2002) Identification and structure of the anti-sigma factor-binding domain of the
disulphide-stress regulated sigma factor σR from Streptomyces coelicolor. *J. Mol. Biol.*,
10.1016/S0022-2836(02)00948-8.

20. Nizan-Koren,R., Manulis,S., Mor,H., Iraki,N.M. and Barash,I. (2003) The Regulatory Cascade That
Activates the Hrp Regulon in Erwinia herbicola pv. gypsophilae. *Mol. Plant-Microbe Interact.*,
10.1094/mpmi.2003.16.3.249.

21. Wecke,T., Halang,P., Staroń,A., Dufour,Y.S., Donohue,T.J. and Mascher,T. (2012)
Extracytoplasmic function σ factors of the widely distributed group ECF41 contain a fused
regulatory domain. *Microbiologyopen*, 10.1002/mbo3.22.

22. Liu,Q., Pinto,D. and Mascher,T. (2018) Characterization of the Widely Distributed Novel ECF42

Group of Extracytoplasmic Function σ Factors in Streptomyces venezuelae. *J. Bacteriol.*,
10.1128/jb.00437-18.

23. Chandrashekar Iyer,S., Casas-Pastor,D., Kraus,D., Mann,P., Schirner,K., Glatter,T., Fritz,G. and
Ringgaard,S. (2019) Transcriptional regulation by σ factor phosphorylation in bacteria. *Nat.*
Microbiol.

24. Lonetto,M.A., Brown,K.L., Rudd,K.E. and Buttner,M.J. (1994) Analysis of the Streptomyces
coelicolor sigE gene reveals the existence of a subfamily of eubacterial RNA polymerase sigma
factors involved in the regulation of extracytoplasmic functions. *Proc. Natl. Acad. Sci.*, **91**, 7573–
7577.

25. Weigt,M., White,R.A., Szurmant,H., Hoch,J.A. and Hwa,T. (2009) Identification of direct residue
contacts in protein-protein interaction by message passing. *Proc. Natl. Acad. Sci. U. S. A.*,
10.1073/pnas.0805923106.

26. Wu,H., Liu,Q., Casas☐Pastor,D., Dürr,F., Mascher,T. and Fritz,G. (2019) The role of C☐terminal
extensions in controlling ECF σ factor activity in the widely conserved groups ECF 41 and ECF
42. *Mol. Microbiol.*, 10.1111/mmi.14261.

27. Steinegger,M. and Söding,J. (2017) MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. *Nat. Biotechnol.*, **35**, 1026–1028.

28. Rhodius,V.A., Segall-Shapiro,T.H., Sharon,B.D., Ghodasara,A., Orlova,E., Tabakh,H.,
Burkhardt,D.H., Clancy,K., Peterson,T.C., Gross,C.A., *et al.* (2013) Design of orthogonal genetic
switches based on a crosstalk map of σs, anti-σs, and promoters. *Mol. Syst. Biol.*,
10.1038/msb.2013.58.

29. Grass,G., Fricke,B. and Nies,D.H. (2005) Control of expression of a periplasmic nickel efflux pump
by periplasmic nickel concentrations. In *BioMetals*.

30. Marcos-Torres,F.J., Perez,J., Gomez-Santos,N., Moraleda-Munoz,A. and Munoz-Dorado,J. (2016)
In depth analysis of the mechanism of action of metal-dependent sigma factors: Characterization
of CorE2 from Myxococcus xanthus. *Nucleic Acids Res.*, 10.1093/nar/gkw150.

31. Chevalier,S., Bouffartigues,E., Bazire,A., Tahrioui,A., Duchesne,R., Tortuel,D., Maillot,O.,
Clamens,T., Orange,N., Feuilloley,M.G.J., *et al.* (2019) Extracytoplasmic function sigma factors in
Pseudomonas aeruginosa. *Biochim. Biophys. acta. Gene Regul. Mech.*, **1862**, 706–721.

32. Yoshimura,M., Asai,K., Sadaie,Y. and Yoshikawa,H. (2004) Interaction of Bacillus subtilis
extracytoplasmic function (ECF) sigma factors with the N-terminal regions of their potential anti-
sigma factors. *Microbiology*, **150**, 591–9.

33. Pinto,D., Liu,Q. and Mascher,T. (2019) ECF σ factors with regulatory extensions: The
one☐component systems of the σ universe. *Mol. Microbiol.*, 10.1111/mmi.14323.

34. Pérez,J., Muñoz-Dorado,J. and Moraleda-Muñoz,A. (2018) The complex global response to copper
in the multicellular bacterium Myxococcus xanthus. *Metallomics*, **10**, 876–886.

35. Luo,Y., Asai,K., Sadaie,Y. and Helmann,J.D. (2010) Transcriptomic and phenotypic
characterization of a Bacillus subtilis strain without extracytoplasmic function σ factors. *J.*

909  *Bacteriol.*, **192**, 5736–45.

910 36. Bibb,M.J. and Buttner,M.J. (2003) The streptomyces coelicolor developmental transcription factor
911   σBldN is synthesized as a proprotein. *J. Bacteriol.*, 10.1128/JB.185.7.2338-2345.2003.

912 37. Thakur,K.G., Joshi,A.M. and Gopal,B. (2007) Structural and biophysical studies on two promoter
913   recognition domains of the extra-cytoplasmic function σ factor σC from Mycobacterium
914   tuberculosis. *J. Biol. Chem.*, 10.1074/jbc.M606283200.

915 38. Kim,M.S., Hahn,M.Y., Cho,Y., Cho,S.N. and Roe,J.H. (2009) Positive and negative feedback
916   regulatory loops of thiol-oxidative stress response mediated by an unstable isoform of σR in
917   actinomycetes. *Mol. Microbiol.*, 10.1111/j.1365-2958.2009.06824.x.

918 39. Bayer-Santos,E., Lima,L.D.P., Ceseti,L. de M., Ratagami,C.Y., de Santana,E.S., da Silva,A.M.,
919   Farah,C.S. and Alvarez-Martinez,C.E. (2018) Xanthomonas citri T6SS mediates resistance to
920   Dictyostelium predation and is regulated by an ECF σ factor and cognate Ser/Thr kinase.
921   *Environ. Microbiol.*, **20**, 1562–1575.

922 40. Lan,L., Deng,X., Zhou,J. and Tang,X. (2006) Genome-wide gene expression analysis of
923   Pseudomonas syringae pv. tomato DC3000 reveals overlapping and distinct pathways regulated
924   by hrpL and hrpRS. *Mol. Plant. Microbe. Interact.*, **19**, 976–87.

925 41. Merighi,M., Majerczak,D.R., Stover,E.H. and Coplin,D.L. (2003) The HrpX/HrpY two-component
926   system activates hrpS expression, the first step in the regulatory cascade controlling the Hrp
927   regulon in Pantoea stewartii subsp. stewartii. *Mol. Plant. Microbe. Interact.*, **16**, 238–48.

928 42. Luo,S., Sun,D., Zhu,J., Chen,Z., Wen,Y. and Li,J. (2014) An extracytoplasmic function sigma
929   factor, σ(25), differentially regulates avermectin and oligomycin biosynthesis in Streptomyces
930   avermitilis. *Appl. Microbiol. Biotechnol.*, **98**, 7097–112.

931 43. Tran,N.T., Huang,X., Hong,H., Bush,M.J., Chandra,G., Pinto,D., Bibb,M.J., Hutchings,M.I.,
932   Mascher,T. and Buttner,M.J. (2019) Defining the regulon of genes controlled by σ E , a key
933   regulator of the cell envelope stress response in Streptomyces coelicolor. *Mol. Microbiol.*,
934   10.1111/mmi.14250.

935 44. Kadowaki,T., Yukitake,H., Naito,M., Sato,K., Kikuchi,Y., Kondo,Y., Shoji,M. and Nakayama,K.
936   (2016) A two-component system regulates gene expression of the type IX secretion component
937   proteins via an ECF sigma factor. *Sci. Rep.*, 10.1038/srep23288.

938 45. Seipke,R.F., Patrick,E. and Hutchings,M.I. (2014) Regulation of antimycin biosynthesis by the
939   orphan ECF RNA polymerase sigma factor σ AntA. *PeerJ*, 10.7717/peerj.253.

940 46. Balogh,D., Dahmen,M., Stahl,M., Poreba,M., Gersch,M., Drag,M. and Sieber,S.A. (2017) Insights
941   into ClpXP proteolysis: heterooligomerization and partial deactivation enhance chaperone affinity
942   and substrate turnover in Listeria monocytogenes. *Chem. Sci.*, 10.1039/c6sc03438a.

943 47. Yanamandra,S.S., Sarrafee,S.S., Anaya-Bergman,C., Jones,K. and Lewis,J.P. (2012) Role of the
944   Porphyromonas gingivalis extracytoplasmic function sigma factor, SigH. *Mol. Oral Microbiol.*,
945   10.1111/j.2041-1014.2012.00643.x.

946 48. Heruth,D.P., Pond,F.R., Dilts,J.A. and Quackenbush,R.L. (1994) Characterization of genetic

determinants for R body synthesis and assembly in Caedibacter taeniospiralis 47 and 116. *J. Bacteriol.*, 10.1128/jb.176.12.3559-3567.1994.

49. Müller,M., Reiß,S., Schlüter,R., Mäder,U., Beyer,A., Reiß,W., Marles-Wright,J., Lewis,R.J., Pförtner,H., Völker,U., *et al.* (2014) Deletion of membrane-associated Asp23 leads to upregulation of cell wall stress genes in Staphylococcus aureus. *Mol. Microbiol.*, 10.1111/mmi.12733.

50. Shukla,J., Gupta,R., Thakur,K.G., Gokhale,R. and Gopal,B. (2014) Structural basis for the redox sensitivity of the Mycobacterium tuberculosis SigK-RskA σ-anti-σ complex. *Acta Crystallogr. D. Biol. Crystallogr.*, **70**, 1026–36.

51. Han,K., Li,Z., Peng,R., Zhu,L., Zhou,T., Wang,L., Li,S., Zhang,X., Hu,W., Wu,Z., *et al.* (2013) Extraordinary expansion of a Sorangium cellulosum genome from an alkaline milieu. *Sci. Rep.*, **3**, 2101.

52. Nechaev,S. and Severinov,K. (2003) Bacteriophage-induced modifications of host RNA polymerase. *Annu. Rev. Microbiol.*, **57**, 301–22.

53. Peng,Q. and Yuan,Y. (2018) Characterization of a novel phage infecting the pathogenic multidrug-resistant Bacillus cereus and functional analysis of its endolysin. *Appl. Microbiol. Biotechnol.*, **102**, 7901–7912.

54. Brister,J.R., Ako-Adjei,D., Bao,Y. and Blinkova,O. (2015) NCBI viral Genomes resource. *Nucleic Acids Res.*, 10.1093/nar/gku1207.

55. Ortiz de Ora,L., Lamed,R., Liu,Y.J., Xu,J., Cui,Q., Feng,Y., Shoham,Y., Bayer,E.A. and Muñoz-Gutiérrez,I. (2018) Regulation of biomass degradation by alternative σ factors in cellulolytic clostridia. *Sci. Rep.*, 10.1038/s41598-018-29245-5.

56. Braun,V., Mahren,S. and Ogierman,M. (2003) Regulation of the FecI-type ECF sigma factor by transmembrane signalling. *Curr. Opin. Microbiol.*, **6**, 173–80.

57. Sievers,F., Wilm,A., Dineen,D., Gibson,T.J., Karplus,K., Li,W., Lopez,R., McWilliam,H., Remmert,M., Söding,J., *et al.* (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.*, 10.1038/msb.2011.75.

58. Finn,R.D., Clements,J. and Eddy,S.R. (2011) HMMER web server: Interactive sequence similarity searching. *Nucleic Acids Res.*, 10.1093/nar/gkr367.

59. Nguyen,L.-T., Schmidt,H.A., von Haeseler,A. and Minh,B.Q. (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.*, **32**, 268–74.

60. Letunic,I. and Bork,P. (2016) Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res.*, 10.1093/nar/gkw290.

61. Tsirigos,K.D., Peters,C., Shu,N., Käll,L. and Elofsson,A. (2015) The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides. *Nucleic Acids Res.*, 10.1093/nar/gkv485.

62. Tsirigos,K.D., Elofsson,A. and Bagos,P.G. (2016) PRED-TMBB2: Improved topology prediction and detection of beta-barrel outer membrane proteins. In *Bioinformatics*.

63.   El-Gebali,S.,   Mistry,J.,   Bateman,A.,   Eddy,S.R.,   Luciani,A.,   Potter,S.C.,   Qureshi,M.,
Richardson,L.J., Salazar,G.A., Smart,A., *et al.* (2019) The Pfam protein families database in
2019. *Nucleic Acids Res.*, **47**, D427–D432.
64.   Pedregosa,F., Varoquaux,G., Gramfort,A., Michel,V., Thirion,B., Grisel,O., Blondel,M., Müller,A.,
Nothman,J., Louppe,G., *et al.* (2012) Scikit-learn: Machine Learning in Python.
65. Li,W. and Godzik,A. (2006) Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. *Bioinformatics*, **22**, 1658–9.
66. Wilbur,W.J. and Lipman,D.J. (1983) Rapid similarity searches of nucleic acid and protein data
banks. *Proc. Natl. Acad. Sci. U. S. A.*, **80**, 726–30.
67. Cock,P.J.A., Antao,T., Chang,J.T., Chapman,B.A., Cox,C.J., Dalke,A., Friedberg,I., Hamelryck,T.,
Kauff,F., Wilczynski,B., *et al.* (2009) Biopython: Freely available Python tools for computational
molecular biology and bioinformatics. *Bioinformatics*, 10.1093/bioinformatics/btp163.
68. Brown,D.P., Krishnamurthy,N. and Sjölander,K. (2007) Automated protein subfamily identification
and classification. *PLoS Comput. Biol.*, **3**, e160.
69. Liu,X., Brutlag,D.L. and Liu,J.S. (2001) BioProspector: discovering conserved DNA motifs in
upstream regulatory regions of co-expressed genes. *Pac. Symp. Biocomput.*
70. Crooks,G.E., Hon,G., Chandonia,J.M. and Brenner,S.E. (2004) WebLogo: A sequence logo
generator. *Genome Res.*, 10.1101/gr.849004.
71. Altschul,S.F., Gish,W., Miller,W., Myers,E.W. and Lipman,D.J. (1990) Basic local alignment search
tool. *J. Mol. Biol.*, 10.1016/S0022-2836(05)80360-2.
1005