Differences in Activation of Intracellular Signalling in Primary Human Retinal Endothelial Cells Between Isoforms of VEGFA-165.

Wendelin Dailey[1], Roberto Shunemann[1,2], Fang Yang[1,3], Megan Moore[1], Austen Knapp[1], Peter Chen[1], Mrinalini Deshpande[1], Brandon Metcalf[1], Quentin Tompkins[1], Alvaro E. Guzman[1], Jennifer Felisky[1], Kenneth P. Mitton[1]*

1. Eye Research Institute, Oakland University, Rochester Michigan, USA

2. Currenty, Clinica Opthtalmus, Joinville – SC, Brazil

3. Currently Department of Ophthalmology, Renmin Hospital, Hubei University of Medicine, Shiyan, Hubei 442000,P.R. China.

Supported by National Eye Institute / National Institutes of Health (USA) grant NIH EY025089 (KPM).

* Corresponding author.

Kenneth P. Mitton, PhD FARVO

Associate Professor of Biomedical Sciences

Oakland University

mitton@oakland.edu

1-248-370-2079

Keywords: blood-retinal barrier, VEGFA$_{165}$a, VEGFA$_{165}$b, MAPK, AKT, Endothelial blood-retinal barrier, retinal cell culture

ABSTRACT

Purpose: There are reports that a b-isoform of Vascular Endothelial Growth Factor-A-165 (VEGFA$_{165}$b) is predominant in normal human vitreous, switching to the a-isoform (VEGFA$_{165}$a) in the vitreous of some diseased eyes. While these isoforms appear to have a different ability to activate the VEGF-Receptor-2 (VEGFR2) in various endothelial cells, the nature of their ability to activate intracellular signalling pathways is not fully characterized, especially in retinal endothelial cells. We determined their activation potential for two key intracellular signalling pathways (MAPK, AKT) over complete dose-response curves and compared potential effects on the expression of several VEGFA$_{165}$ target genes in primary human retinal microvascular endothelial cells (HRMECs).

Methods: To determine full dose-response curves for the activation of MAPK (ERK1/2), AKT and VEGFR2, direct in-cell western assays were developed using primary Human Retinal Microvascular Endothelial Cells (HRMECs). Potential differences in dose-response effects on gene expression markers related to endothelial cell / leukocyte adhesion (*ICAM1*, *VCAM1* and *SELE*) and tight-junctions (*CLDN5* and *OCLN*) were tested by quantitative-PCR.

Results: Activation dose-response analysis revealed much stronger activation of MAPK, AKT and VEGFR2 by the a-isoform at lower doses. MAPK activation in primary HRMECs displayed a sigmoidal dose-response to a range of VEGFA**$_{165}$**a concentrations spanning 10-250 pM, which shifted higher into the 100-5,000 pM range with VEGFA**$_{165}$**b. Similar maximum activation of MAPK was achieved by both isoforms at high concentration. Maximum activation of AKT by VEGFA**$_{165}$**b was only half of the maximum activation from VEGFA**$_{165}$**a. At a lower intermediate dose, where VEGFA**$_{165}$**a activated intracellular signalling stronger than VEGFA**$_{165}$**b, the changes to VEGFA target gene expression was generally greater with VEGFA**$_{165}$**a.

Conclusions: In primary HRMECs, VEGFA$_{165}$a could maximally activate MAPK and AKT at lower concentrations where VEGFA$_{165}$b had relatively little effect. The timing for maximal activation of MAPK was similar for both isoforms, which is different than reprorted for non-retinal endothelial cells. While VEGFA$_{165}$a and VEGFA$_{165}$b are limited to the sequence of their six C-terminal six amino acids, this results in a large difference in their ablility to activate at least two key intracellular signalling pathways and potentially VEGF target gene expression in primary human retinal endothelial cells.

Introduction

Vascular Endothelial Growth Factor-A-165 (VEGFA$_{165}$) is the isoform of VEGF that is primarily responsible for driving retinal vascular pathology in diabetic retinopathy (DR), age-related macular degeneration (AMD) and retinopathy of prematurity (ROP), through activation of VEGF Receptor-2 (VEGFR2). A seminal analysis of the vitreous fluids of patients with diabetic retinopathy established the presence of elevated VEGF concentrations associated with this and other conditions [1]. That study contributed to the eventual development of intravitreal VEGF-blocking drugs to treat neovascularization and edema in AMD, DR, and the current exploration of their use for ROP [2–5].

   Blockade of VEGFA activity is provided by the use of intravitreal drugs including ranibizumab (Lucentis, Genentech), bevacizumab (Avastin, Genentech), pegaptanib (Macugen, Bausch & Lomb), and aflibercept (Eylea, Regeneron). Some VEGF-blocking drugs injected into the vitreous can enter the systemic circulation and, in the case of bevicizumab and aflibercept they can substantially decrease serum VEGF concentration for several days [6]. There is interest to advance VEGF-regulating therapies by more precise titration of VEGF concentration or

modulating specific VEGF-mediated signalling rather than the full blockade provided by current drugs. This field will continue to benefit from a more complete knowledge of VEGF's signaling mechanisms within retinal endothelial cells. Unfortunately, much previous VEGF research does not technically differentiate between VEGFA's isoforms and in the case of the retina there is far less investigation using retinal endothelial cells, and even less from human retina.

In addition to endothelial cells, VEGF receptors are expressed by cells of the immune system, including early and late hematopoietic progenitor cells, dendritic cells, T-lymphocytes and macrophages [7]. Among the three VEGF receptor tyrosine-kinases (R1, R2, R3), VEGFR2 is required for angiogenesis and vasculogenesis, and its expression is most abundant in endothelial and endothlial progenitor cells [8–10]. VEGFR2 binds several isoforms of the VEGFA that are produced from alternative-splicing [11–14]. The most frequently detected isoforms are VEGFA$_{121}$, VEGFA$_{165}$ and VEGFA$_{189}$, with 121, 165 and 189 amino acids respectively. VEGFA$_{121}$ is most diffusible, lacking the heparin-binding domains of VEGFA$_{165}$ and VEGFA$_{189}$. VEGFA$_{189}$ has even more affinity for heparin than VEGFA$_{165}$ with an additional heparin-binding domain encoded by exon-6 [11,12]. VEGFA$_{165}$ bind as a dimer to activate VEGFR2, and it also binds Neuropilin-1 as a co-receptor. VEGFA$_{121}$ lacks two regions required to binding Neuropilin-1.

A vast knowledge of VEGF-mediated signalling is mostly derived from the study of non-retinal cell types, and is beyond the scope of this paper but readers can refer to the excellent review by Koch and Claesson-Welch [15]. In endothelial cells, several pathways are activated by VEGFA$_{165}$ that affect proliferation, migration, survival and endothelial permeability. Two of

these pathways include MAPK (ERK1/2) and the AKT (PKB), where MAPK is a dominant regulator of cell proliferation and AKT modulates cell survival and permeability [16–19]. Auto-phosphorylation of VEGFR2 (Y1175) leads to RAS-independent activation of the PLCγ (Phospholipase-Cγ) / Protein Kinase C / MAPK pathway [20,21]. Activation of VEGFR2 also leads to activation of the TSAD / PI3K (Phosphotidylinositol-3 Kinase) / PDK (Phosphoinositide-dependent protein kinase) / AKT pathway [22,23]. AKT phosphorylates the BCL-2-associated death protein (BAB) and Caspase-9 to block apoptosis and increase EC-survival [24]. AKT also activates eNOS (endothelial Nitric-Oxide Synthase, *NOS3*) to modulate vascular permeability [25].

More recently b-isoforms of VEGFA were described that differ from the previously known a-isoforms in the sequence of their C-terminal six amino acids: CDKRPP in VEGFA$_{165}$a becomes SLTRDKD in VEGFA$_{165}$b [13,14]. This difference renders VEGFA$_{165}$b incapable of bringing Neuropilin-1 into a receptor/ligand complex with VEGFR2 [26–28]. Similar to VEGFA$_{121}$, VEGFA$_{165}$b is reported to be less angiogenic than VEGFA$_{165}$a [13,29,30]. VEGFA$_{165}$b's binding affinity for VEGFR2 itself is similar to that of VEGFA$_{165}$a in Human Umbilical Vein Endothelial Cells (HUVECs) and less than VEGFA$_{165}$a in direct binding anlaysis in transformed HEK293 cell assays [31,32]. VEGFA$_{165}$b also displays weaker activation of VEGFR2 and ERK1/2 (MAPK) than VEGFA$_{165}$a in transfected porcine aortic endothelial (PAE) cells [31]. Perrin et al. (2005), reported a switch from mostly VEGFA$_{165}$b to mostly VEGFA$_{165}$a in the vitreous of patients with active diabetic retinopathy [33]. A similar shift, from VEGFA$_{165}$b to VEGFA$_{165}$a, was reported in vitreous humor from young patients with ROP [34]. A murine equivalent of isoform switching (VEGFA$_{164}$b to VEGFA$_{164}$a) was also reported in a mouse model

of oxygen-induced retinopathy [35]. Several publications using isoform specific antibodies have detected VEGFA$_{165}$b in human and mouse tissues, but their presense from RNA analysis in sheep hypothalamus is controversial [36].

While single-dose studies using transfected cells or non-retinal endothelial cells indicate that VEGFA$_{165}$b activates VEGFR2 and MAPK less than VEGFA$_{165}$a, we do not know if this is true throughout a full dose range. We also wanted to determine full-dose response curves for activation of intracellular signaling in primary human retinal microvascular endothelial cells. We hypothesized that there is a significant difference in the full dose-reponse curves for activation of these pathways in human retinal endothelial cells between VEGFA$_{165}$a and VEGFA$_{165}$b. To address this, we used *in situ* assays to determined full dose-reponse curves for the activation of MAPK (ERK1/2), AKT and VEGFR2 in primary HRMECs for VEGFA$_{165}$a and VEGFA$_{165}$b. Our results demonstrate that there is a significant dose-response difference in the ability of these two isoforms to activate intracellular signalling kinases and that these dose-response differences propagate to differences in VEGF-target gene expression in primary human retinal endothelial cells.

Methods

Certification of Primary Human Retinal Microvascular Endothelial Cells: Primary Human Retinal Microvascular Endothelial Cells (HRMECs, genotype XY) were obtained from Cell Systems as passage-3 cells (Kirkland, WA, Catalog number: ACBRI-181). Cells used for experimentation were not used past passage-7. Endothelial character of the cell line was established by Cell Systems immunofluorescence testing: > 95% positive by immunofluorescence

for Cytoplasmic VWF / Factor VIII, Cytoplasmic uptake of Di-I-Ac-LDL, Cytoplasmic CD31, and < 1% by immunofluorescence for GFAP (Glial Fibrillary Acidic Protein), Glutamine Synthetase, NG2 (Neural/Glial Antigen 2), and PDGFR-beta (Platelet Derived Growth Factor Receptor beta). The cell line was also subjected to short tandem repeat (STR) Profile Testing by Cell Systems at the Master Level (P1), performed by the laboratory analysis service of the American Type Culture Collection (ATCC). Seventeen STR-loci plus the gender determining locus, Amelogenin, were amplified using the commercially available PowerPlex® 18D Kit from Promega. The cell line sample was processed using the ABI Prism® 3500xl Genetic Analyzer. Data were analyzed using the GeneMapper® IDX v1.2 software (Applied Biosystems). Appropriate positive and negative controls were used throughout the test procedure. DNA Profile (STR #12625 Analysis) was as follows: TH01: 9, 9.3; D5S818: 9, 13; D13S317: 8, 10; D7S820: 11, 12; D16S539: 11, 12; CSF1PO: 10, 12; Amelogenin: X, Y; vWA: 16, 19; TPOX: 8, 11. The ATCC test conclusions were that the submitted sample profile is human, but not a match any profile in the ATCC STR database, as would be expected for primary HRMECs. Additionally, routine testing for retinal endothelial character in the author's laboratory (Eye Research Institute, Oakland University) confirmed that the HRMECs express all three of the human Norrin receptor-complex genes *FZD4*, *LRP5* and *TSPAN12,* using Taqman real-time PCR analysis (data not shown). HRMECs also demonstrated VEGF-medidated regulation of several VEGF target genes, as shown in the results.

Cell Culture and Antibodies: EndoGRO-MV Complete media kit was obtained from Millipore (Burlington, MA). EndoGro basal medium was supplemented with rhEGF, L-Glutamine, Heparin Sulfate and Ascorbic Acid according to the kit instructions. Additionally,

supplementation with FBS and Hydrocortisone Hemisuccinate (1µg/ml) was dependent upon the

assay requirements. Recombinant human VEGFA$_{165}$a and VEGFA$_{165}$b were obtained from R&D

Systems (Minneapolis, MN). Odyssey Blocking Buffer and an Odyssey Infrared Imager were

purchased from LI-COR Biosciences (Lincoln, NE). The antibodies used for in situ labeling of

HMRECs (In Cell Westerns, ICW) and Western blotting are summarized in Table 1. Mini

Protean TGX gels were purchased from Bio-Rad (Hercules, CA).

Table -1. Specific Antibodies used and Sources

Antibody	IgG Species	Company	Cat. No.
Phospho-AKT (P-Ser473)	Rabbit	Cell Signaling Technologies	4058
AKT(pan)	Rabbit	Cell Signaling Technologies	4691
Phospho-p44/42 MAPK (Thr202/204)	Rabbit	Cell Signaling Technologies	4370
p44/42 MAPK (pan)	Rabbit	Cell Signaling Technologies	4695
Phospho-VEGFR2 (Tyr1175)	Rabbit	Cell Signaling Technologies	2478
Beta-Actin	Mouse	Millipore	MAB1501
Anti-mouse IgG IRDye 800CW	Goat	LI-COR	32210
Anti-rabbit IgG IRDye 680RD	Goat	LI-COR	68071
Anti-rabbit IgG IRDye 800CW	Goat	LI-COR	32211
Anti-mouse IgG IRDye 680RD	Goat	LI-COR	68070

Immunoblotting of activated MAPK and AKT in HRMECs: HRMECs were grown in 100mm

dishes that had been pre-coated with Attachment Factor (Cell Systems, WA). Media was

EndoGRO (No VEGF, Millipore Sigma, Burlington, MA) with 5% FBS to establish confluency.

When cells were confluent, the media was replaced with fresh media with or without VEGFA**$_{165}$**

isoforms. After incubation for 10 minutes, the cells were washed with ice cold PBS and the

dishes were scraped to detach the cells. The cell suspension was centrifuged at 1000 X g for 5

minutes and the cells were reconstituted in RIPA cell-lysis buffer (150 mM NaCl, 1% Triton-X-100, 0.1% SDS, 50 mM Tris, pH 8.0, 10 mM Sodium Fluoride, 1 mM Sodium Orthovanadate and complete Protease inhibitor cocktail (1 tablet/10 ml)). The cells were sonicated or vortexed every 10 minutes while kept on ice for 30 minutes. The cell lysate was collected after centrifuging at 14,000 X g for 15 minutes at 4°C. The protein concentration was measured using Pierce BCA Protein Assay (Thermo Scientific). The samples were prepared with Laemmli sample buffer and loaded onto a (4-15%) gradient gels for SDS-PAGE electrophoresis. After electrophoresis the gels were equilibrated in cold transfer buffer and transferred to PDVF membranes overnight. The membranes were blocked with Odyssey blocking buffer and incubated with appropriates rabbit primary antibodies (Table-1) along with mouse monoclonal Actin antibody which was used for normalization. After washing with PBS, 0.1% tween-20 (4 X 5 min), the membranes were incubated with secondary antibodies (Goat Anti-rabbit IRDye680 and Goat Anti-mouse IRDye800CW) for 30-60 minutes. Membranes were washed and scanned on an Odyssey Infrared imager (Li-Cor, Lincoln, NE).

Dose-response analysis of intracellular signaling in primary HMRECs: HRMECs were seeded into black 96 well plates (5000/well) that had been coated with Attachment Factor. The cells were grown to confluence using fully supplemented EndoGRO-MV media for 4-5 days. The cells were serum starved overnight using EndoGRO-MV without Hydrocortisone. The cells were treated with $VEGFA_{165}a$ or $VEGFA_{165}b$ for varied lengths of time after which the treatments were immediately removed and replaced with of 4% Paraformaldehyde. The cells were fixed for 20 minutes at room temperature followed by permeabilization with PBS, 0.1% Triton X-100 (10 minutes). Cells were blocked by incubation with Odyssey Blocking Buffer (Li-Cor) for 1.5 hours

at room temperature and incubated with primary antibodies (1:200) for either 2 hours at room

temperature or over night at 4 °C. Rabbit antibodies were used against the proteins of interest and

a mouse monoclonal anti-Beta-Actin antibody was used for normalization. The cells were washed

with PBS, 0.1% Tween 20 (5x5 minutes) and incubated with secondary antibodies, Goat anti-

Rabbit IRDye 800CW & Goat anti-Mouse IRDye 680RD (1:750) for 45 minutes at room

temperature. After washing with PBS 0.1% Tween 20 (5x5 minutes) the plates were scanned on

an Odyssey Imager (Li-Cor). For time-response and dose-response experiments, the doses of

VEGFA$_{165}$a and VEGFA$_{165}$b were assayed in quadruplicate wells and dose-response experiments

were repeated three times to confirm reproducibility of relative dose-response curves for

activation of MAPK (phospho-Thr202/Tyr204), AKT (phospho-Ser473) and VEGFR2 (phospho-

Tyr1175). Data were fit to a 4-parameter log-logistic response function Eq (1) at the 95%

confidence level for each dose x using the Drc package for R. [37]. The parameters fit were: b,

the steepness of the curve at e the effective dose ED50, with c and d the lower and upper limits of

the response. The 95% confidence level for fitting was used to produce curves.

$$f(x,(b,c,d,e)) \;=\; c + \frac{(d\text{-}c)}{(1+ exp(b(log(x) - log(e))))} \qquad\qquad Eq\ (1)$$

Quantitative PCR analysis of HMREC Gene Expression: Primary HMRECs were grown to

confluence in six well plates. After the desired treatment, cells were trypsinized. Total RNA was

isolated using the Absolutely RNA miniprep kit (Agilent, Santa Clara, CA). Cells were

homogenized in 200µL of lysis buffer. Homogenization was accomplished using conical pellet-

pestles in 1.5 mL microfuge tubes, with a hand-held rotary tool (Bel-Art,Wayne, NJ). First-strand

cDNA was synthesized by reverse transcribing 500 ng of total RNA per sample using either the

AffinityScript qPCR DNA Synthesis kit (Agilent), or the LunaScript RT Super Mix Kit (NEB #

E3010(S/L), Ipswich, MA) with Oligo-dT priming. The reaction conditions were according to the

manufacturer's instructions. For AffinityScript: 25ºC for 5 min, 42ºC for 20 min, 95ºC for 5 min,

and 10ºC for 10 min. For LunaScript RT: 25ºC for 2 min, 55ºC for 10 min, and 95ºC for 1 min.

All compared samples were processed using the same reagent set. Stock first strand cDNA

preparations were stored at -70°C, and were not used for analysis after a maximum of three

freeze-thaws. Duplex reaction format was utilized with FAM-labeled probe/primer pairs for the

gene of interest and VIC-labeled probe/primer-limited pairs for TBP (Tata-Binding Protein) as

the normalizer gene (ThermoFisher, Waltham, MA). For real-time PCR reactions, sample first-

strand cDNA was diluted 5-fold with deionized water and 2µL added to 18µL of master mix for

20µL PCR reactions. Triplicate reactions were used for each sample. Master Mix chemistries

were either 2x Gene Expression Master Mix (ThermoFisher, Applied Biosystems # 4369016,

Waltham, MA), or the Luna Universal Probe qPCR Master Mix (2x) Gene Expression (New

England BioLabs # M3004L, Ipswich, MA), both mixes with the Rox reference dye option.

Reactions were run on either an Mx3000P real-time PCR system using the MxPro software, or an

AriaMx Real-time PCR System using the AriaMx HRM QPCR Software (Agilent, *Santa Clara,*

CA). Gene expression assays were evaluated for high PCR efficiency using a dilution series of

HMREC cDNA to ensure validity of using the delta-delta Ct method for comparing relative gene

expression. Each replicate reaction was internally normalized relative to endogenous TBP gene

expression. The specific assay probe sets used for gene expression analysis are listed in Table 2

Table 2. Taqman gene expression probes for HRMEC gene expression.

Gene	Probe Set #	Spans Exons
CLDN5	Hs01561351_m1	1-2
ICAM1	Hs00164932_m1	2-3
OCLN	Hs00170162_m1	5-6
SELE	Hs00174057_m1	4-5
TBP	Hs00427620_m1	3-4
VCAM1	Hs01003372_m1	6-7

RESULTS

Preliminary observations:

The cell-based experimental results reported here were inspired by observations made during preliminary testing for a different project to develop an intravitreal injection model using VEGFA$_{165}$a to cause dilation of primary retinal veins, which was reported to occur in the Long Evans rat [38]. That model typically involves a high dose of VEGFA. Using fluoresceine angiography, and SD-OCT imaging, it was noted that similar effects on the vasculature could be induced with VEGFA$_{165}$b or VEGFA$_{165}$a. While the observational studies did not use large numbers of rats, they are included as supplimental observations because they brought our attention that the dose-responses for intracellular pathway activation were not fully known in primary human retinal endothelial cells.

Activation of MAPK and AKT in primary HRMECs by VEGFA$_{165}$a and VEGFA$_{165}$b:

To examine the effect of VEGF$_{165}$ isoforms on intracellular signaling pathways we chose primary human retinal microvascular endothelial cells (HRMECs) and initially used immunoblotting to

test activation of MAPK and AKT pathways by both isoforms. These pathways were examined as two of the intracellular signaling kinases implicated in mediating the effects of VEGFA on endothelial proliferation and blood-retinal barrier. Using a high dose (100 ng/mL, 5300 pM) for maximal activation, both isoforms of VEGFA$_{165}$ increased the amounts of the active form of MAPK (Figure-1a) and the active form of AKT (Figure-1b). From three separate full experimental repeats, the average activation was about 2.7-fold and 1.6-fold (Figure-1c) for MAPK and AKT respectively. While VEGFA$_{165}$a and VEGFA$_{165}$b treatments always increased the activation of both kinases, the magnitude of the increase was variable between experiments as illustrated by comparing two of the three experiments on the same gel (Figure 1a,b).

Dose-Response for the Activation of MAPK by VEGFA$_{165}$a and VEGFA$_{165}$b:

While repeated immunoblotting experiments confirmed activation of MAPK and AKT by both VEGFA$_{165}$ isoforms, the experimental variability introduced by protein extraction, protein assays, electro-transfer, band-shape and image analysis was judged to be unsuitable for dose-response analysis. To address the variability, new *in situ* assays, or In-Cell Western (ICW) assays, were developed using the same kinase-specific antibodies while removing the sample processing required for immunoblotting. This also permitted faster fixation of cells at the desired time, the use of many doses, and the use of several biological replicates per dose.

Activation of the MAPK (ERK1/2) pathway in primary HRMECs occurred maximally at 10 minutes for both VEGFA$_{165}$a and VEGFA$_{165}$b treatment, and returned to control levels by 90 minutes, using a dose of 2ng/mL (100 pM). See Figure-2a. The activation of VEFGA$_{165}$a was significantly stronger than that of VEGFA$_{165}$b at 10 minutes (t-test, P<0.01).

We next carried out dose-response analysis for the activation of the MAPK at 10 minutes with cells pre-adapted to lower serum. Eleven doses of each isoform, with 4 biological replicates per dose, were evaluated and dose-response data for MAPK activation were fit to the Log-Logistic 4-parameter dose-response function using the Drc package in R [37]. (Figure-2b) Activation of MAPK by VEGFA$_{165}$b showed a typical sigmoidal dose-response. Compared to VEGFA$_{165}$b, the dose-response curve for VEGFA$_{165}$a displayed a strong allosteric shift to a more binary-like activation response. VEGFA$_{165}$a was more potent for activation of MAPK at lower doses. A similar maximum level of MAPK activation could be achieved with the highest dose tested (10,000 pM) using either isoform. Activation of MAPK by VEGFA$_{165}$b was only 10% of that generated by VEGFA$_{165}$a using a dose of 250 pM. The ED$_{50}$ was 73 pM for VEGFA$_{165}$a and 1015 pM for VEGFA$_{165}$b for the experiment shown. The experiment was repeated three times confirming the large difference in ED$_{50}$ between the two isoforms. Differences between the ED$_{50}$ values were typically in the 800 to 1000 pM range.

Dose-Response for the Activation of AKT Activation by VEGFA$_{165}$a and VEGFA$_{165}$b:

Maximum activation of the pAKT pathway in HRMECs occurred 30 minutes after VEGFA$_{165}$a treatment and 15 minutes after VEGFA$_{165}$b treatment (Figure-3a) using what was thought to be a maximally activating dose (100 ng/mL 5,300 pM). Dose response analysis was carried out for AKT activation using these respective maximum time points and the data were fit to the 4-parameter log-logistic response function. (Figure-3b). With VEGFA$_{165}$a, AKT activation in HRMECs displayed a much steeper dose-response curve than for VEGFA$_{165}$b. The ED$_{50}$ values were 53 pM for VEGFA$_{165}$a compared to 126 pM for VEGFA$_{165}$b. The maximum activation of

AKT generated by the b-isoform was less than half that obtained from treatment with the a-isoform. Repetition of the experiment confirmed similar relative responses.

Dose-Response for the Activation of VEGFR2 by VEGFA$_{165}$a and VEGFA$_{165}$b:

To test for relative difference in dose-response at the level of the receptor itself we examined the dose-response for activation of VEGFR2. Maximum activation of the receptor was fast, within less than a few minutes, as quickly as cells could be processed for treatment and fixation. Five and ten minute experiments were similar in relative response, so ten minutes were used for the dose-response experiments and data were fit to the 4-parameter log-logistic response function as shown in Figure-4a. VEGFA$_{165}$a treatment achieved maximum activation by about 500 pM whereas VEGFA$_{165}$b caused little if any activation at that dose. The ED$_{50}$ for activation by VEGFA$_{165}$a was 254 pM compared to 1192 pM for the VEGFA$_{165}$b. Furthermore, the maximum activation by VEGFA$_{165}$b was 73% of that obtained with VEGFA$_{165}$a. Again, the relative pattern was confirmed by a repeated experiment (data not shown).

Isoform Effects on Expression Leukocyte Docking Protein Genes:

We first examined gene expression from 1 to 24 hours after treatment with high doses (100 ng/mL, 5300 pM) of VEGFA$_{165}$a and VEGFA$_{165}$b to determine an optimal time-point for monitoring any effects on the expression of *ICAM1*, *Selectin-E*, and *VCAM1*. A dose of 5300 pM was likely a saturating concentration and more than what would be experienced *in vivo*, to find a suitable time-point where gene expression could be subsequently compared using physiologically relevant doses. All three of these genes displayed increased expression after treatment with both isoforms of VEGFA$_{165}$ (Figure-5 a, c, e). These trends were confirmed by repeated experiments

(data not shown). *Of the genes examined here, ICAM1'*s increasing expression trend was more variable in timing, not always maximum at 6 hours, sometimes later by 24 hours (data not shown).

Selecting a fixed treatment time of 6-hours, and guided by the activation dose-response curves for MAPK and AKT, we tested a low (2ng/mL, 100 pM), intermediate (19 ng/mL, 1000 pM) and high (95 ng/mL, 5000 pM) dose to test for differences in the effects on leukocyte-docking gene expression. For all three leukocyte docking protein genes (*ICAM1, SELE, VCAM1*), VEGFA$_{165}$a increased their expression at the intermediate dose of 1000 pM where VEGFA$_{165}$b still had very little effect (Figure-5 b, d, f).

Isoform Effects on Expression of Tight Junction Protein Genes:

The effects of VEGFA$_{165}$a and VEGFA$_{165}$b on the expression of the two key tight-junction protein genes, CLDNS and OCLN, were also compared. These genes encoding the tight-junction proteins Claudin-5 and Occludin. (See Figure-6) Both isoforms of VEGFA$_{165}$ reduced the expression of these genes by the 6 hour time-point (Figure-6 a, c). We next comparing a low (2ng/mL, 100 pM), intermediate (19 ng/mL, 1000 pM) and high saturating dose (95 ng/mL, 5000 pM) at the fixed 6-hour time point. VEGFA$_{165}$b was less effective at suppressing the expression of *Claudin-5* and *Occludin* compared to VEGFA$_{165}$a (Figure-6 b, d)

DISCUSSION

While the role of VEGFA in vascular development, tumorgenesis and hypoxia has been studied for almost three decades, less is known about the functional differences between VEGFA isoforms. More recently, isoform-specific analysis reported that most VEGFA$_{165}$ in normal vitreous was VEGFA$_{165}$b, with the ratio changing to mostly VEGFA$_{165}$a in eyes with active diabetic retinopathy and ROP [33,34]. The relative expression of VEGFA$_{165}$a and VEGFA$_{165}$b can be regulated by alternative splicing of the VEGFA$_{165}$ pre-mRNA, which is regulated by phosphorylation of the ASF/SF2 splice-factor by SR-protein kinase (SRPK1/2) [39]. This raised the possibility that modulation of this isoform ratio might be exploited for therapeutic purposes. Modulation of the splicing between the VEGFA$_{165}$a and VEGFA$_{165}$b mRNAs was demonstrated using SRPK inhibitors in human retinal pigment epithelial (RPE) cells, and topical infusion of these inhibitors reduced neovasularization in a mouse laser-induced CNV model [40]. Another group used intravitreal injection of a different SRPK inhibitor to inhibit neovascularization in the mouse CNV model [41]. An antibody specific for the C-terminus of VEGFA$_{165}$a, which does not bind VEGFA$_{165}$b, can also inhibit the proangiogenic effects of VEGFA$_{165}$a [42].

Specific blockade of the VEGFA$_{165}$a protein, or supressing VEGFA$_{165}$a expression in favor of VEGFA$_{165}$b, could be a therapeutic strategy if there are significant differences in the ability of these isoforms to activate intracellular signalling in the retinal endothelium. The preference for examining signaling differences using primary HRMECs is supported by studies that revealed differences between endothelial cells derived from different organs. For example, viral delivery of a murine recombinant-VEGF$_{164}$b attenuated inflammatory-response damage in a murine model of ulcerative colitis, yet it exacerbated blood-brain-barrier damage in a model of

focal cerebral ischemia [43–45]. Extensive gene expression profiling has also demonstrated that

HUVECs have a greater expression of embryonic genes, and other differences, compared to

endothelial cells from the choroid and neural retina. Significant differences also exist between

human retinal endothelial cells and human choroidal endothelial cells in the expression of genes

that are related to endothelial function and neovascularization [46].

We began with immunoblotting to examine the activation of MAPK and AKT in primary

HRMECs. While VEGFA$_{165}$b does not bind the Neuropilin-1 co-receptor, it does bind to

VEGFR2 with similar or less affinity as VEGFA$_{165}$a depending on the cell model and method

used [31,32]. Irregardless of relative binding affinity, VEGFA$_{165}$b is reported to bind and activate

VEGFR2 based on immunoblotting studies [31,47,48]. Using immunoblotting, we found that

both MAPK and AKT were activated to a similar extent by VEGFA$_{165}$a or VEGFA$_{165}$b, using a

dose of 100 ng/ml (5,300 pM) in primary HRMECs. Previous single-dose immunoblotting

studies in different cell-types have reported that VEGFA$_{165}$b activated VEGFR2 and MAPK less

than VEGFA$_{165}$a, and that the maximum activation of MAPK by VEGFA$_{165}$b occurred later than

VEGFA$_{165}$a in porcine aortic endothelial cells (PAECs) and bovine aortic endothelial cells using

a dose of 2,600 pM [31,47]. Another study of human pulmonary microvascular endothelial cells

(HPMECs) found that the a-isoform was a stronger activator of VEGFR2 and MAPK (ERK1/2)

than the b-isoform at a dose of 20 ng/ml (1000 pM) [48].

We found that both MAPK and AKT were activated less by VEGFA$_{165}$b at the lower dose

(20 ng/ml, 1000 pM) (data not shown). However, it was noted that while activation was detected

in repeated experiments, the relative manitude of activation was variable using immunoblotting.

Immunublotting does involvenumerous processing steps and a subtantial number of cells are required for a single sample, therefore we decided that *in situ* assays would be supperior for generating full dose-response curve data.

To determine the dose-response for activation of MAPK and AKT, *in situ* ICW assays were developed that permitted the use of multiple replicate doses with a limited supply of primary cells in a 96 well plate format. The resulting dose-response curves confirmed that there were substantial differences in the activation of both MAPK and AKT between VEGFA$_{165}$a and VEGFA$_{165}$b in primary HRMECs. In the case of MAPK, the ED$_{50}$ of VEGFA$_{165}$a 900 pM less than the ED$_{50}$ for VEGFA$_{165}$b. VEGFA$_{165}$a caused near maximal activation of MAPK at a concentration of 250 pM, whileVEGFA$_{165}$b had little effect at 250 pM. Higher doses of VEGFA$_{165}$b could activate MAPK to a similar maximum level, and the timing for maximum activation of MAPK was similar for both isoforms in HRMECs, occuring by 10 minutes and decreasing by 30 minutes. This relative timing was different from that reported for HUVECs where the maximum concentration of active MAPK occured 30 minutes after treatment with VEGFA$_{165}$b [31].

For AKT, the difference in ED$_{50}$ between VEGFA$_{165}$a and VEGFA$_{165}$b was not as large as seen for MAPK, just two-fold greater for VEGFA$_{165}$b compared to VEGFA$_{165}$a. However, the activation of AKT by VEGFA$_{165}$b was subtanially less than from VEGFA$_{165}$a over the entire dose-response range. In contrast to MAPK activation, the time to maximum activation of AKT was earlier with VEGFA$_{165}$b (15 minutes) compared to the VEGFA$_{165}$a (30 minutes). While the differences in ED50 values were not as large as seen for MAPK activation, the maximum

activation of AKT obtained with VEGFA$_{165}$b was only 50% of the maximum activation generated by VEGFA$_{165}$a. This was another difference from that reported in a non-retinal endothelial cell-type, human pulmonary endothelial cells (HPECs), whereAKT activation is the same for VEGFA$_{165}$a and VEGFA$_{165}$b at a dose of 20 ng/ml (1000pM) in HPECs [48].

The differences noted above in the activation of MAPK and AKT between HRMECs and non-retinal endothelial cells suggest that VEGF-mediated intracellular signaling patterns are not completely universal between endothelial cells from different organs. An evaluation of primary endothelial cells from the human retina should be utilized whenever possible to study the retinal context. Altogether, our dose-response curves for the activation of VEGFR2, MAPK, and AKT in primary HRMECs were consistent with the model proposed by Whitaker et al. (2001), wherein VEGFA$_{165}$a activates VEGFR2 more intensely when it can bind the co-receptor Neuropilin-1, to form a larger activation complex [49].

Based on the primary HRMEC dose-response data for MAPK activation, VEGFA$_{165}$a was more potent for activating intracellular signalling than VEGFA$_{165}$b at lower doses, which fall into a range of total VEGFA$_{165}$a concentrations previously reported in human vitreous. Vitreous concentrations for VEGFA have been measured by various techniques in several disease conditions. Most elevated VEGFA is the VEGFA$_{165}$ isoform, which diffuses slowly, and the vitreous has protease activity, therefore it is possible that the concentrations measured in the vitreous are somewhat lower than in the neural retina itself. With that stated, vitreal concentrations of 25 pM to over 400 pM have been reported for proliferative diabetic retinopathy from several studies [1,50–54]. Other studies have reported 10 pM in diabetic macular edema

[55], 100 to 450 pM in retinopathy of prematurity [34,56] and an average of 430 pM, with as high as 580 pM, in central retinal vein occlusion [57]. It is possible that elevations of retinal VEGFA$_{165}$a concentration to several-hundred pico-Molar could increase the activation of endothelial intracellular signalling above its normal baseline. In contrast, the much lower activation potential of VEGFA$_{165}$b in this concentration range would leave MAPK activation close to normal if most of the VEGFA$_{165}$ existed as the b-isoform.

We also found that VEGFA$_{165}$b could affect the expression of the same target genes that are also regulated by VEGFA$_{165}$a in primary HRMECs. These included genes involved in leukocyte-endothelial cell adhesion and tight-junction formation. Even though the 5,000 pM dose of VEGFA$_{165}$b could activate MAPK and AKT more strongly than the intermediate 1,000 pM dose, it is unlikely that the 5,000 pM concentration would be experienced *in vivo*. For that reason we suggest that the effects on gene expression at 1,000 pM were most informative. We found that the *CLDN5* gene was particulary succeptible to repression by both isoforms of VEGFA$_{165}$ compared to *OCLN* in this cell type. VEGFA-mediated activation of the VEGFR2 receptor is known to disrupt adherin-junctions and causes ß-Catenin mediated repression of *Claudin-5* gene expression in endothelial cells [58]. For all three leukocyte docking protein genes (*ICAM1, SELE, VCAM1*), VEGFA$_{165}$a increased their expression at the intermediate dose of 1000 pM where VEGFA$_{165}$b had less effect. We concluded that there were significant dose-response differences in the regulation of VEGFA-target genes, with VEGFA$_{165}$a proving to be the stronger affector of gene expression overall.

Summary: Our results indicated that differences in the last 6 amino acids between VEGFA$_{165}$a and VEGFA$_{165}$b have a significant affect on the relative activation of MAPK and AKT within primary HRMECs. VEGFA$_{165}$a activated both pathways at lower concentrations where VEGF$_{165}$b had little effect. The greater potential of VEGFA$_{165}$a to affect HRMECs also extended to changes in the expression of genes required for leukocyte-docking and tight-junction structure. Large dose-response differences in activation of these pathways exist in ranges of elevated VEGFA$_{165}$a concentrations that have been reported in the vitreous of diseased eyes. Key aspects of MAPK and AKT activation, such as timing and maximal activation, were different in HRMECs compared to those reported for some other non-retinal endothelial cell-types. These results would support the concept that specific blockade of VEGFA$_{165}$a, or modulating the VEGFA$_{165}$b / VEGFA$_{165}$a ratio, could be useful thereapeutic strategies for retinal diseases involving elevated VEGFA concentration.

Acknowledgements

The authors would like to thank Oakland University students Regan Miller and Anju Thomas for assistance with some cell culture and some testing of gene expression probesets used for these studies. Research supported by National Eye Institute / National Institutes of Health (USA) grant NIH EY025089 (KPM).

References

1. Aiello LP, Avery RL, Arrigg PG, Keyt BA, Jampel HD, Shah ST, Pasquale LR, Thieme H, Iwamoto MA, Park JE. Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. N Engl J Med 1994; 331:1480–7.

2. Friedman DS. 2012 Fifth Edition Vision Problems in the U.S. Vis Probl US 2012; .

3. Miller JW, Adamis AP, Aiello LP. Vascular endothelial growth factor in ocular neovascularization and proliferative diabetic retinopathy. Diabetes Metab Rev 1997; 13:37–50.

4. Robbins SG, Conaway JR, Ford BL, Roberto KA, Penn JS. Detection of vascular endothelial growth factor (VEGF) protein in vascular and non-vascular cells of the normal and oxygen-injured rat retina. Growth Factors 1997; 14:229–241.

5. Amadio M, Govoni S, Pascale A. Targeting VEGF in eye neovascularization: What's new?: A comprehensive review on current therapies and oligonucleotide-based interventions under development. Pharmacol Res 2016; 103:253–269.

6. Avery RL, Castellarin AA, Steinle NC, Dhoot DS, Pieramici DJ, See R, Couvillion S, Nasir MA, Rabena MD, Le K, Maia M, Visich JE. Systemic pharmacokinetics following intravitreal injections of ranibizumab, bevacizumab or aflibercept in patients with neovascular amd. Br J Ophthalmol 2014; 98:1636–1641.

7. Li Y-L, Zhao H, Ren X-B. Relationship of VEGF/VEGFR with immune and cancer cells: staggering or forward? Cancer Biol Med 2016; 13:206–14.

8. Quinn TP, Peters KG, De Vries C, Ferrara N, Williams LT. Fetal liver kinase 1 is a receptor for vascular endothelial growth factor and is selectively expressed in vascular endothelium. Proc Natl Acad Sci U S A 1993; 90:7533–7.

9. Millauer B, Wizigmann-Voos S, Schnürch H, Martinez R, Møller NP, Risau W, Ullrich A. High affinity VEGF binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis. Cell 1993; 72:835–46.

10. Oelrichs RB, Reid HH, Bernard O, Ziemiecki A, Wilks AF. NYK/FLK-1: a putative receptor protein tyrosine kinase isolated from E10 embryonic neuroepithelium is expressed in endothelial cells of the developing embryo. Oncogene 1993; 8:11–8.

11. Park JE, Keller GA, Ferrara N. The vascular endothelial growth factor (VEGF) isoforms:

differential deposition into the subepithelial extracellular matrix and bioactivity of extracellular matrix-bound VEGF. Mol Biol Cell 1993; 4:1317–1326.

12. Ferrara N. Binding to the Extracellular Matrix and Proteolytic Processing: Two Key Mechanisms Regulating Vascular Endothelial Growth Factor Action. Mol Biol Cell 2010; 21:687–690.

13. Bates DO, Cui TG, Doughty JM, Winkler M, Sugiono M, Shields JD, Peat D, Gillatt D, Harper SJ. VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, is down-regulated in renal cell carcinoma. Cancer Res 2002; 62:4123–4131.

14. Qiu Y, Hoareau-Aveilla C, Oltean S, Harper SJ, Bates DO. The anti-angiogenic isoforms of VEGF in health and disease. Biochem Soc Trans 2009; 37:1207–1213.

15. Koch S, Claesson-Welsh L. Signal transduction by vascular endothelial growth factor receptors. Cold Spring Harb Perspect Med 2012; 2:1–21.

16. Im E, Kazlauskas A. Regulating angiogenesis at the level of PtdIns-4,5-P2. EMBO J 2006; 25:2075–2082.

17. Gerber HP, McMurtrey A, Kowalski J, Yan M, Keyt BA, Dixit V, Ferrara N. Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation. J Biol Chem 1998; 273:30336–43.

18. Adini I. RhoB controls Akt trafficking and stage-specific survival of endothelial cells during vascular development. Genes Dev 2003; 17:2721–2732.

19. Gille H, Kowalski J, Yu L, Chen H, Pisabarro MT, Davis-Smyth T, Ferrara N. A repressor sequence in the juxtamembrane domain of Flt-1 (VEGFR-1) constitutively inhibits vascular endothelial growth factor-dependent phosphatidylinositol 3'-kinase activation and endothelial cell migration. EMBO J 2000; 19:4064–73.

20. Takahashi T, Yamaguchi S, Chida K, Shibuya M. A single autophosphorylation site on KDR/Flk-1 is essential for VEGF-A-dependent activation of PLC-γ and DNA synthesis in vascular endothelial cells. EMBO J 2001; 20:2768–2778.

21. Takahashi T, Ueno H, Shibuya M. VEGF activates protein kinase C-dependent, but Ras-independent Raf-MEK-MAP kinase pathway for DNA synthesis in primary endothelial cells. Oncogene 1999; 18:2221–30.

22. Ruan G-X, Kazlauskas A. Axl is essential for VEGF-A-dependent activation of PI3K/Akt.

EMBO J 2012; 31:1692–703.

23. Cantley LC. The phosphoinositide 3-kinase pathway. Science 2002; 296:1655–7.

24. Cardone MH, Roy N, Stennicke HR, Salvesen GS, Franke TF, Stanbridge E, Frisch S, Reed JC. Regulation of cell death protease caspase-9 by phosphorylation. Science 1998; 282:1318–21.

25. Fulton D, Gratton JP, McCabe TJ, Fontana J, Fujio Y, Walsh K, Franke TF, Papapetropoulos A, Sessa WC. Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. Nature 1999; .

26. Woolard J, Wang WY, Bevan HS, Qiu Y, Morbidelli L, Pritchard-Jones RO, Cui TG, Sugiono M, Waine E, Perrin R, Foster R, Digby-Bell J, Shields JD, Whittles CE, Mushens RE, Gillatt DA, Ziche M, Harper SJ, Bates DO. VEGF165b, an inhibitory vascular endothelial growth factor splice variant: mechanism of action, in vivo effect on angiogenesis and endogenous protein expression. Cancer Res 2004; 64:7822–7835.

27. Soker S, Takashima S, Miao HQ, Neufeld G, Klagsbrun M. Neuropilin-1 is expressed by endothelial and tumor cells as an isoform- specific receptor for vascular endothelial growth factor. Cell 1998; 92:735–745.

28. Prahst C, Heroult M, Lanahan AA, Uziel N, Kessler O, Shraga-Heled N, Simons M, Neufeld G, Augustin HG. Neuropilin-1-VEGFR-2 complexing requires the PDZ-binding domain of neuropilin-1. J Biol Chem 2008; 283:25110–25114.

29. Magnussen AL, Rennel ES, Hua J, Bevan HS, Beazley Long N, Lehrling C, Gammons M, Floege J, Harper SJ, Agostini HT, Bates DO, Churchill AJ, Long NB, Lehrling C, Gammons M, Floege J, Harper SJ, Agostini HT, Bates DO, Churchill AJ. VEGF-A165b is cytoprotective and antiangiogenic in the retina. Invest Ophthalmol Vis Sci 2010; 51:4273–4281.

30. Konopatskaya O, Churchill AJ, Harper SJ, Bates DO, Gardiner TA. VEGF165b, an endogenous C-terminal splice variant of VEGF, inhibits retinal neovascularization in mice. Mol Vis 2006; 12:626–632.

31. Cébe Suarez S, Pieren M, Cariolato L, Arn S, Hoffmann U, Bogucki A, Manlius C, Wood J, Ballmer-Hofer K. A VEGF-A splice variant defective for heparan sulfate and neuropilin-1 binding shows attenuated signaling through VEGFR-2. Cell Mol Life Sci 2006; 63:2067–77.

32. Peach CJ, Kilpatrick LE, Woolard J, Hill SJ. Comparison of the ligand-binding properties of fluorescent VEGF-A isoforms to VEGF receptor 2 in living cells and membrane preparations using NanoBRET. Br J Pharmacol 2019; .

33. Perrin RM, Konopatskaya O, Qiu Y, Harper S, Bates DO, Churchill AJ. Diabetic retinopathy is associated with a switch in splicing from anti- to pro-angiogenic isoforms of vascular endothelial growth factor. Diabetologia 2005; 48:2422–2427.

34. Zhao M, Xie WK, Bai YJ, Huang LZ, Wang B, Liang JH, Yin H, Li XX, Shi X. Expression of total vascular endothelial growth factor and the anti☐angiogenic VEGF165b isoform in the vitreous of patients with retinopathy of prematurity. Chin Med J (Engl) 2015; 128:2505–2509.

35. Zhao M, Shi X, Liang J, Miao Y, Xie W, Zhang Y, Li X. Expression of pro- and anti-angiogenic isoforms of VEGF in the mouse model of oxygen-induced retinopathy. Exp Eye Res 2011; 93:921–926.

36. Lomet D, Piégu B, Wood SH, Dardente H. Anti-angiogenic VEGFAxxxb transcripts are not expressed in the medio-basal hypothalamus of the seasonal sheep. PLoS One 2018; 13:e0197123.

37. Ritz C, Baty F, Streibig JC, Gerhard D. Dose-Response Analysis Using R. PLoS One 2015; 10:e0146021.

38. Miyamoto K, Khosrof S, Bursell SE, Moromizato Y, Aiello LP, Ogura Y, Adamis AP. Vascular endothelial growth factor (VEGF)-induced retinal vascular permeability is mediated by intercellular adhesion molecule-1 (ICAM-1). Am J Pathol 2000; 156:1733–1739.

39. Nowak DG, Amin EM, Rennel ES, Hoareau-Aveilla C, Gammons M, Damodoran G, Hagiwara M, Harper SJ, Woolard J, Ladomery MR, Bates DO. Regulation of vascular endothelial growth factor (VEGF) splicing from pro-angiogenic to anti-angiogenic isoforms: a novel therapeutic strategy for angiogenesis. J Biol Chem 2010; 285:5532–40.

40. Gammons M V., Fedorov O, Ivison D, Du C, Clark T, Hopkins C, Hagiwara M, Dick AD, Cox R, Harper SJ, Hancox JC, Knapp S, Bates DO. Topical antiangiogenic SRPK1 inhibitors reduce choroidal neovascularization in rodent models of exudative AMD. Invest Ophthalmol Vis Sci 2013; 54:6052–62.

41. Hatcher JM, Wu G, Zeng C, Zhu J, Meng F, Patel S, Wang W, Ficarro SB, Leggett AL,

Powell CE, Marto JA, Zhang K, Ki Ngo JC, Fu X-D, Zhang T, Gray NS. SRPKIN-1: A Covalent SRPK1/2 Inhibitor that Potently Converts VEGF from Pro-angiogenic to Anti-angiogenic Isoform. Cell Chem Biol 2018; 25:460-470.e6.

42. Carter JG, Gammons MVR, Damodaran G, Churchill AJ, Harper SJ, Bates DO. The carboxyl terminus of VEGF-A is a potential target for anti-angiogenic therapy. Angiogenesis 2015; 18:23–30.

43. Cromer W, Jennings MH, Odaka Y, Michael Mathis J, Steven Alexander J. MURINE rVEGF164b, AN INHIBITORY VEGF REDUCES VEGF-A DEPENDENT ENDOTHELIAL PROLIFERATION AND BARRIER DYSFUNCTION. Microcirculation 2010; 17:no-no.

44. Cromer WE, Ganta C V, Patel M, Traylor J, Kevil CG, Alexander J, Mathis J. VEGF-A isoform modulation in an preclinical TNBS model of ulcerative colitis: protective effects of a VEGF164b therapy. J Transl Med 2013; 11:207.

45. Chaitanya G V., Cromer WE, Parker CP, Couraud PO, Romero IA, Weksler B, Mathis JM, Minagar A, Alexander JS. A Recombinant Inhibitory Isoform of Vascular Endothelial Growth Factor164/165 Aggravates Ischemic Brain Damage in a Mouse Model of Focal Cerebral Ischemia. Am J Pathol 2013; 183:1010–1024.

46. Browning AC, Halligan EP, Stewart EA, Swan DC, Dove R, Samaranayake GJ, Amoaku WM. Comparative gene expression profiling of human umbilical vein endothelial cells and ocular vascular endothelial cells. Br J Ophthalmol 2012; 96:128–132.

47. Kawamura H, Li X, Harper SJ, Bates DO, Claesson-Welsh L. Vascular endothelial growth factor (VEGF)-A165b is a weak in vitro agonist for VEGF receptor-2 due to lack of coreceptor binding and deficient regulation of kinase activity. Cancer Res 2008; 68:4683–4692.

48. Ourradi K, Blythe T, Jarrett C, Barratt SL, Welsh GI, Millar AB. VEGF isoforms have differential effects on permeability of human pulmonary microvascular endothelial cells. Respir Res 2017; 18:116.

49. Whitaker GB, Limberg BJ, Rosenbaum JS. Vascular Endothelial Growth Factor Receptor-2 and Neuropilin-1 Form a Receptor Complex that is Responsible for the Differential Signaling Potency of VEGF165 and VEGF121. J Biol Chem 2001; 276:25520–25531.

50. Li B, Li M-D, Ye J-J, Chen Z, Guo Z-J, Di Y. Vascular endothelial growth factor

concentration in vitreous humor of patients with severe proliferative diabetic retinopathy after intravitreal injection of conbercept as an adjunctive therapy for vitrectomy. Chin Med J (Engl) 2020; NA;664–669.

51. Rezzola S, Nawaz MI, Cancarini A, Semeraro F, Presta M. Vascular Endothelial Growth Factor in the Vitreous of Proliferative Diabetic Retinopathy Patients: Chasing a Hiding Prey? Diabetes Care 2019; 42:e105–e106.

52. Wang J, Chen S, Jiang F, You C, Mao C, Yu J, Han J, Zhang Z, Yan H. Vitreous and plasma VEGF levels as predictive factors in the progression of proliferative diabetic retinopathy after vitrectomy. PLoS One 2014; .

53. Baharivand N, Zarghami N, Panahi F, Yazdan DGM, Mahdavi Fard A, Abas M. Relationship between vitreous and serum vascular endothelial growth factor levels, control of diabetes and microalbuminuria in proliferative diabetic retinopathy. Clin Ophthalmol 2012; 185.

54. Funatsu H. Risk evaluation of outcome of vitreous surgery for proliferative diabetic retinopathy based on vitreous level of vascular endothelial growth factor and angiotensin II. Br J Ophthalmol 2004; 88:1064–1068.

55. Yenihayat F, Özkan B, Kasap M, Karabaş VL, Güzel N, Akpınar G, Pirhan D. Vitreous IL-8 and VEGF levels in diabetic macular edema with or without subretinal fluid. Int Ophthalmol 2019; 39:821–828.

56. Sonmez K, Drenser KA, Capone A, Trese MT, Capone Jr. A, Trese MT. Vitreous levels of stromal cell-derived factor 1 and vascular endothelial growth factor in patients with retinopathy of prematurity. Ophthalmology 2008; 115:1065-1070 e1.

57. Ehlken C, Rennel ES, Michels D, Grundel B, Pielen A, Junker B, Stahl A, Hansen LL, Feltgen N, Agostini HTHT, Martin G. Levels of VEGF but not VEGF165b are Increased in the Vitreous of Patients With Retinal Vein Occlusion. Am J Ophthalmol 2011; 152:298-303.e1.

58. Taddei A, Giampietro C, Conti A, Orsenigo F, Breviario F, Pirazzoli V, Potente M, Daly C, Dimmeler S, Dejana E. Endothelial adherens junctions control tight junctions by VE-cadherin-mediated upregulation of claudin-5. Nat Cell Biol 2008; 10:923–934.

FIGURE LEGENDS

Figure 1. MAPK and AKT are activated in HRMECs by VEGFA$_{165}$a and VEGFA$_{165}$b

A) Immunoblots of total MAPK and activated MAPK (Phospho-MAPK) 10 minutes after treatment with VEGFA$_{165}$a (Va), VEGFA$_{165}$b (Vb), or media control (C). Two representative experiments are shown. **B)** Immunoblots of total AKT and activated AKT (Phospho-AKT) 10 minutes after treatment with VEGFA$_{165}$a (Va), VEGFA$_{165}$b (Vb), or media control (C). Two representative experiments are shown. **C)** Fold activation of MAPK and AKT from three immunoblotting experiments. P-values are shown, t-test relative to media controls (1-fold).

Figure 2. VEGFA$_{165}$a strongly activates MAPK in primary HRMECs at doses where VEGFA$_{165}$b has little effect.

Activation of MAPK was measured by *in-situ* immunofluorescence with a phospho-specific antibody (phospho-Thr202/Tyr204) for p44/42. **A)** Activation of MAPK at three time-points after the addition of 100 pM VEGFA$_{165}$a or VEGFA$_{165}$b. Bar shows standard deviation. (t-test, *** $p<0.001$ relative to untreated control, n=8 biological replicates.) **B)** Dose-response curves for activation of MAPK. The ED$_{50}$ was 73 pM for VEGFA$_{165}$a and 1015 pM for VEGFA$_{165}$b. Bars indicate the 95% confidence interval for data fit to the 4-parameter log-logistic function. N=4 biological replicates per dose.

Figure 3. VEGFA$_{165}$b is a weaker activator of AKT in primary HRMECs compared to VEGFA$_{165}$a.

A) Activation of AKT was measured by *in-situ* immunofluorescence with a phospho-specific antibody for Ser473-AKT at five time-points after the addition of 20 ng/mL (1,250 pM) VEGFA$_{165}$a or VEGFA$_{165}$b. Bar shows standard deviation. (t-test, ***$p<0.001$ relative to untreated control, n=4 biological replicates.) **B)** Dose response curves for activation of AKT. The ED$_{50}$ value for AKT activation by VEGFA$_{165}$a was 53 pM compared to 126 pM for VEGFA$_{165}$b. Bars indicate the 95% confidence interval for data fit to the 4-parameter log-logistic function. N=4 biological replicates per dose.

Figure 4. Differences in the activation of intracellular signalling between VEGFA$_{165}$a and VEGFA$_{165}$b in primary HRMECs begin at the receptor VEGFR2.

VEGFA$_{165}$a was a stronger activator of VEGFR2 compared to VEGFA$_{165}$b. The ED$_{50}$ for activation by VEGFA$_{165}$a was 254 pM compared to 1192 pM for VEGFA$_{165}$b. Bars indicate the 95% confidence interval for data fit to the 4-parameter log-logistic function. N=4 biological replicates per dose.

Figure 5. VEGFA$_{165}$a has a stronger effect on leukocyte-endothelial adhesion gene expression in primary HRMECs than VEGFA$_{165}$b.

Relative gene expression measured by qPCR. **A)** *ICAM1* gene expression at 1, 3, 6 and 24 hours after VEGFA$_{165}$ treatment. Confluent HRMECs were treated with a saturating high-dose of VEGFA$_{165}$a or VEGFA$_{165}$b (100 ng/mL, 5300 pM). **B)** *ICAM1* gene expression comparing VEGFA$_{165}$a and VEGFA$_{165}$b at a low, intermediate and high dose measured after 6-hours. **C)** *Selectin-E* gene expression at 1, 3, 6 and 24 hours after VEGFA$_{165}$ treatment. **D)** *Selectin-E* gene expression comparing VEGFA$_{165}$a and VEGFA$_{165}$b at a low, intermediate and high doses measured after 6-hours. **E)** *VCAM1* gene expression at 1, 3, 6 and 24 hours after VEGFA$_{165}$ treatment. **F)** *VCAM1* gene expression comparing VEGFA$_{165}$a and VEGFA$_{165}$b at a low, intermediate and high dose measured after 6-hours. (Triplicate assays, error bars show standard deviation. Anova: compared to media control *p<0.05, **p<0.01; compared to VEGFA$_{165}$a † p<0.05, ††p<0.01)

Figure 6. VEGFA$_{165}$a has a stronger effect on tight-junction gene expression in primary HRMECs than VEGFA$_{165}$b.

Relative gene expression measured by qPCR . **A)** *Claudin-5* gene expression at 1, 3, 6 and 24 hours after VEGFA$_{165}$ treatment. Confluent HRMECs were treated with a saturating high-dose of VEGFA$_{165}$a or VEGFA$_{165}$b (100 ng/mL, 5300 pM). **B)** *Claudin-5* gene expression comparing VEGFA$_{165}$a and VEGFA$_{165}$b at a low, intermediate and high doses measured after 6-hours. **C)** *Occludin* gene expression at 1, 3, 6 and 24 hours after VEGFA$_{165}$ treatment. **D)** *Occludin* gene expression comparing VEGFA$_{165}$a and VEGFA$_{165}$b at a low, intermediate and high dose measured after 6-hours. (Triplicate assays, error bars show standard deviation. Anova: compared to media control *p<0.05, **p<0.01; compared to VEGFA$_{165}$a †† p<0.01)