Direct antiviral activity of interferon stimulated genes is responsible for resistance to

paramyxoviruses in ISG15-deficient cells

David Holthaus[1]*‡, Andri Vasou[1]*, Connor G. G. Bamford[2]‡, Jelena Andrejeva[1], Christina Paulus[1],

Richard E. Randall[1], John McLauchlan[2], David J. Hughes[1]†

[1]Biomedical Sciences Research Complex, School of Biology, University of St Andrews, St Andrews,

United Kingdom. [2]MRC-University of Glasgow Centre for Virus Research, Glasgow, United Kingdom.

*These authors contributed equally.

‡ Present addresses: (DH) Unit 16 Mycotic and Parasitic Agents and Mycobacteria, Robert Koch-

Institute, Berlin, Germany. (CGGB) Wellcome-Wolfson Institute for Experimental Medicine,

Queen's University, Belfast UK.

†Corresponding author: djh25@st-andrews.ac.uk; Tel: +44 1334 467197.

Abstract

Interferons (IFNs), produced during viral infections, induce the expression of hundreds of IFN-

stimulated genes (ISGs). Some ISGs have specific antiviral activity while others regulate the cellular

response. In addition to functioning as an antiviral effector, IFN-stimulated gene 15 (ISG15) is a

negative regulator of IFN signalling and inherited ISG15-deficiency leads to autoinflammatory

interferonopathies where individuals exhibit elevated ISG expression in the absence of infection. We

have recapitulated these effects in cultured human A549-ISG15$^{-/-}$ cells and (using A549-UBA7$^{-/-}$ cells)

confirmed that posttranslational modification by ISG15 (ISGylation) is not required for regulation of

the type-I IFN response. ISG15-deficient cells pre-treated with IFN-α were resistant to paramyxovirus

infection. We also showed that IFN-α treatment of ISG15-deficient cells led to significant inhibition

of global protein synthesis leading us to ask whether resistance was due to the direct antiviral

activity of ISGs or whether cells were non-permissive due to translation defects. We took advantage

of the knowledge that IFN-induced protein with tetratricopeptide repeats 1 (IFIT1) is the principal

antiviral ISG for parainfluenza virus 5 (PIV5). Knockdown of IFIT1 restored PIV5 infection in IFN-α-

pre-treated ISG15-deficient cells, confirming that resistance was due to the direct antiviral activity of

the IFN response. However, resistance could be induced if cells were pre-treated with IFN-α for

longer times, presumably due to inhibition of protein synthesis. These data show that the cause of

virus resistance is two-fold; ISG15-deficiency leads to the 'early' over-expression of specific antiviral

ISGs, but the later response is dominated by an unanticipated, ISG15-dependent, loss of

translational control.

Introduction

The innate immune response against pathogens is underpinned by the evolutionary conserved

interferon (IFN) system. All cells express pathogen recognition receptors (PRRs) that sense the

products of infection and establish a signalling cascade leading to the production of cytokines,

including type I IFN (IFN-α/β) (1, 2). IFN is secreted from cells and binds to cell surface receptors

expressed on both infected and non-infected cells, initiating a JAK/STAT signalling cascade,

culminating in the expression of hundreds of interferon stimulated genes (ISGs) (3). The biological

effects of ISGs are extensive and their principle role is to generate an unfavourable environment for

the replication of viruses. Many ISGs have broad antiviral activity, such as double-stranded RNA

dependent protein kinase (PKR) that, upon recognition of viral dsRNA, dampens general protein

synthesis and prevents the translation of viral mRNAs (4). Other antiviral ISGs, such as IFN-induced

protein with tetratricopeptide repeats (IFIT) proteins, inhibit specific viruses, but for many, they are

inconsequential (5). Additionally, multiple ISGs are generally required to limit infection because the

majority of ISGs result in low to moderate levels of inhibition (6); however, ISGs with specific

antiviral properties for a given virus are usually not known. Nevertheless, the nature of the innate

immune response necessitates the production of the complete spectrum of ISGs, albeit with a high

degree of redundancy, as during a natural infection, the identity of the infecting virus is not known.

This response is inevitably tightly regulated, as a dysregulated response leads to a suite of

autoinflammatory diseases (7).

The ubiquitin-like protein (Ubl) ISG15 is strongly induced by IFN and is critical for regulating how cells

respond to infection. As a posttranslational modification (PTM), it can covalently modify proteins in a

process known as ISGylation, and in many cases, modification of viral proteins forms part of the

antiviral response (8). Covalently bound ISG15 can also be removed from proteins by the ubiquitin

specific protease USP18 (9). Importantly, loss-of-function mutations in ISG15 have been identified in

human patients with subsets of autoinflammatory interferonopathies and typically these individuals

demonstrate elevated ISG expression in the absence of infection (10). Mechanistically, it was shown

that ISG15 functions as a negative regulator of type I IFN signalling by stabilising USP18, a known

inhibitor of JAK/STAT signalling (11-13). Intriguingly, despite the known functions of ISG15 and

USP18 in the ISGylation process, the regulation of type I IFN signalling was entirely independent of

ISGylation (10). Interestingly, mouse Isg15 is not required to stabilise Usp18 and appears not to be

needed to regulate IFN signalling, suggesting a species-specific, gain-of-function for human ISG15

(although other species have not yet been tested) (14).

Previous work has shown that cells from ISG15-deficient patients expressed higher levels of ISGs

compared to normal controls when treated with recombinant IFN-α and these cells were resistant to

several viruses (14); however, it was not clear at what stage of infection viruses were blocked nor

how. Furthermore, cells were treated with IFN-α followed by washing (to remove IFN) and rested for

36 hours prior to infection. Since ISG15 is involved in regulating the cell cycle (15) and protein

synthesis (shown in this report), an over-amplified IFN response (due to lack of ISG15 and reduced

levels of USP18) may have led to virus resistance simply because cells were no longer permissive to

infection. This has implications for our understanding as to why ISG15-deficient patients are not

more susceptible to viral infections; these observations have led to the suggestion that, unlike in

mice, human ISG15 is not an antiviral effector (14, 16).

In this study, we recapitulated the phenotype observed in ISG15-deficient patient cells upon

treatment with recombinant IFN-α2b in a cell culture model and dissected the mechanisms that

result in virus resistance during an antiviral state. We showed that resistance was due to the direct

antiviral activity of the type I IFN response and discuss the implications of ISG15-loss-of-function

during the innate immune response. Based on our findings, we conclude that observations from

ISG15-deficient patients alone cannot be used to infer that ISG15 does not possess antiviral effector

functions, as has been proposed (14, 16).

Materials and methods

Cells

Vero cells (African green monkey kidney epithelial cells) and A549 cells (human adenocarcinoma

alveolar basal epithelial cells), and derivatives, were maintained in Dulbecco's modified Eagles's

medium (DMEM; Sigma) supplemented with 10% (v/v) heat-inactivated foetal bovine serum (FBS,

Biowest) and incubated in 5% (v/v) CO_2 at 37°C in a humidified incubator. A549-shIFIT1 have been

described elsewhere (17) and were maintained in blasticidin (10 µg/ml). A549-ISG15$^{-/-}$ cells were

generated by CRISPR/Cas9n system that utilises the D10A dual 'nickase' mutant of Cas9 (Cas9n) that

ostensibly limits off-target effects. Briefly, to disrupt exon 2 of the ISG15 gene, single guide RNA

(sgRNA) sequences were cloned using the pPX460 and transfected into A549 cells as previously

described (18). Transfectants were enriched after 48 h by treating cells with puromycin (1 µg/ml) for

2 d and then diluted to single cells in 96-well plates. Correctly edited cell clones were verified by

immunoblot analysis. A549-ISG15$^{-/-}$-shIFIT1 cells were generated as previously described using A549-

ISG15$^{-/-}$ (B8) and maintained in media with blasticidin (10 µg/ml) (17). To generate A549-UBA7$^{-/-}$

cells, A549 cells were first made to stably express *Streptococcus pyogenes* Cas9 following blasticidin

selection of cells transduced with lentiCas9-Blast (gift from Feng Zhang, Addgene plasmid # 52962

(19)). The sgRNA sequence that targeted exon 3 of UBA7 was chosen computationally

(https://www.deskgen.com) and complementary oligonucleotides (sense:

caccGCACACGGGTGACATCACTG; antisense: aaacCAGTGATGTCACCCGTGTGC) were hybridised and

ligated into the *Bsm* BI site of pLentiGuide-Puro (gift from Feng Zhang, Addgene # 52963 (20)). Cas9-

expressing A549s were transduced with UBA7 sgRNA-expressing lentiGuide-Puro and selected with

puromycin. Puromycin-resistant cells were single-cell cloned by FACS and successful knockout cells

were validated by immunoblot analysis. A549-Npro cells have been described previously (21).

Virus infections and treatments.

Viruses used were human parainfluenza virus 2 (HPIV2) strain Colindale (HPIV2-Co), HPIV3 strain

Washington/47885/57 (HPIV3-Wash) (20), PIV5 strain W3 (PIV5-W3) (22) and PIV5 strain CPI- (PIV5-

CPI-) (23). Virus stocks were prepared by inoculating Vero cells at a multiplicity of infection (MOI) of

0.001 with continual rocking at 37°C. Supernatants were harvested at 2 d p.i., clarified by

centrifugation at 3,000 xg for 15 min, aliquoted and snap-frozen. Titres were estimated by standard

plaque assay on Vero cells in 6-well plates.

For infection studies, cell monolayers were infected in 6-well plates with virus diluted in medium to

achieve a MOI of 10, unless stated otherwise. Virus adsorption was for 1 h, after which the viral

inoculum was removed and replaced with media supplemented with 2% (v/v) FBS and incubated in

5% (v/v) CO_2 at 37°C until harvested. When cells were treated with IFN-α prior to infection (pre-

treated) this was done with 1000 IU/ml IFN-α2b (IntronA, Merck Sharp & Dohme Ltd.) 18 h prior to

infection, unless otherwise stated. IFN-α remained on cells for the duration of experiments. Cells

were either processed for immunoblot analysis or (if infecting with rPIV5-mCherry, kind gift of Dr He,

University of Georgia, USA) imaged using an IncuCyte Zoom imaging system (Sartorius).

For plaque assays 30-40 PFU PIV5-CPI- in 1 ml DMEM, 2% FBS were adsorbed for 1 h onto confluent

monolayers of cells in 6-well plates while rocking at 37°C. Following adsorption, 2 ml overlay

(DMEM, 2% FBS, Avicel) was added to wells and incubated for 6 d. Cells were fixed with 5%

formaldehyde (10 min), washed in PBS and permeabilised for 10 min (PBS, 1% Triton X-100, 3% FBS)

washed again and incubated for 1 h with a pool of PIV5-specific antibodies (24) diluted in PBS, 3%

FBS (1:1000). Following PBS washes, cells were incubated for 1 h with goat anti-mouse IgG

antibodies conjugated to alkaline phosphatase (Abcam Cat# ab97020) diluted 1:1000 in PBS, 3%FBS.

Cells were washed in PBS and signals were detected using SIGMAFAST BCIP/NBT (Sigma).

Reverse transcription quantitative PCR.

Total cellular RNA was purified from cells that had been treated with 1000 IU/ml IFN-α2b (Intron A)

for 18 h, or left untreated, using TRIzol reagent (ThermoFisher Scientific) and Direct-zol RNA

Miniprep Plus kits followed by on-column DNase I treatment for the removal of contaminating DNA

(Zymo Research). Complementary DNA (cDNA) was synthesised in 20 µl reaction volumes with 500

ng total RNA and oligo(dT) using GoScript reverse transcriptase (Promega) according to the

manufacturer's recommendations. Quantitative PCR reaction mixes (20 µl) included 1x PerfeCTa

SYBR green SuperMix (Quanta BioScience), 0.5 µM each primer and 1 µl cDNA reaction mix. Cycling

was performed in a Mx3005P real time PCR machine (Stratagene) and included an initial 3 min

enzyme activation step at 95°C, followed by 40 cycles of 10 s at 95°C, 10 s at 58°C and 20 s at 72°C.

Melting curve analysis was performed to verify amplicon specificity. Quantification of *β-ACTIN* mRNA

was used to normalize between samples and the average cycle threshold (CT) was determined from

three independent cDNA samples from independent cultures. Relative expression compared to non-

treated control cells was calculated using the ΔΔCT method. Primer sequences were: *HERC5*

5'GACGAACTCTTGCACCGTCTC and 5'GCGTCCACAGTCATTTTCCAC, *USP18*

5'ATGCTGCCCAACTGTACCTC and 5'CCTGCAGTCTCTCCACCAAG, *MxA* 5'GCCTGCTGACATTGGGTATAA

and 5'CCCTGAAATATGGGTGGTTCTC, *IFIT1* 5'CCTGGAGTACTATGAGCGGGC and

5'TGGGTGCCTAAGGACCTTGTC, *β-ACTIN* 5'AGCGAGCATCCCCCAAAGTT and

5'AGGGCACGAAGGCTCATCATT.

Immunoblotting.

Confluent monolayers in 6-well dishes were lysed with 250 µl 2 x Laemmli sample buffer (4% w/v

SDS, 20% v/v glycerol, 0.004% w/v bromophenol blue and 0.125 M Tris-HCl, pH 6.8 with 10% v/v β-

mercaptoethanol) for 10 min, incubated at 95°C for 10 min, sonicated at 4°C with 3 cycles of 30 s on

30 s off in a Bioruptor Pico (Diagenode) and clarified by centrifugation at 12,000 xg, 4°C for 10 min.

SDS-PAGE in Tris-glycine-SDS running buffer and immunoblotting followed standard techniques

using the following antibodies: mouse monoclonal anti-ISG15 F-9 (Santa Cruz Biotechnology Cat#

sc166755), rabbit polyclonal anti-MxA (Proteintech Cat# 13750-1-AP), goat polyclonal anti-IFIT1 N-16

(Santa Cruz Biotechnology Cat# sc82946), mouse monoclonal anti-β-ACTIN, UBA7 (anti-UBE1L B-7;

Santa Cruz Biotechnology Cat# sc-390097), mouse monoclonal anti-PIV5 NP 125 (24), mouse

monoclonal anti-HPIV2 and anti-PIV5 P 161 (antibody cross-reacts with P of both viruses(24)), mouse

monoclonal anti-HPIV3 NP (25). Quantitative immunoblots were analysed using an Odyssey CLx

scanner and of membranes incubated with IRDye secondary antibodies (LiCOR). Data were

processed using Image Studio software (LiCOR).

^{35}S-methionine labelling.

Subconfluent A549 and A549-ISG15$^{-/-}$ (B8) cells in 6-well plates were treated with 1000 IU/ml IFN-α

or left untreated. At 24 h, 48 h and 72 h following treatment cells were pulse-labelled with 500

Ci/mmol ^{35}S-Methionine (^{35}S-Met; MP Biomedical) in Met-free media (Sigma) for 1 h. Cells were

washed in PBS, lysed in 2 x Laemmli sample buffer and equal amounts of protein were separated by

SDS-PAGE. Gels were stained with Coomassie (and imaged to ensure equal loading), dried under

vacuum, exposed to a storage phosphor screen and analysed by phoshoimager analysis.

Results

ISG15-knockout A549 cells recapitulate ISG15-deficient patient cells

Among the several immune modulatory roles of ISG15 (8), intracellular ISG15 expression, at least in human cells, is critical for regulating the magnitude of the type I IFN response (10, 14). To investigate the pleotropic nature of human ISG15 we developed cell lines that lack ISG15 expression. Because of our interest in respiratory viruses, including paramyxoviruses, we chose to knockout ISG15 expression in the lung adenocarcinoma cell line A549 by CRISPR/Cas9 genome editing as described previously (18). Furthermore, A549 cells have proved to be a very useful model for understanding virus-IFN interactions. The resulting culture was single cell cloned and ISG15 expression was assessed by immunoblotting three clones (B8, B6 and C4). We also selected a clone that had gone through the CRISPR/Cas9 process but retained ISG15 expression (C4+) (Fig. 1a). In addition to control A549 cells, all clones were treated with IFN-α for 24 h, 48 h or left untreated. Immunoblot analysis showed that, compared to control cells, expression of the ISG MxA was higher in A549-ISG15$^{-/-}$ cells (Fig. 1a). We also tested the impact of ISG15-deficiency on the expression of various ISG mRNAs. For this, we selected clone B8 for further analyses; these cells were, in addition to control A549 cells, treated with IFN-α, or left untreated, for 24 h and the expression of various ISGs were examined by RT-qPCR. Whilst IFN-α treatment enhanced the expression of all ISGs tested, this increase was larger in ISG15-deficient cells compared to control A549 cells (between 5- and 10-fold, depending on the ISG) (Fig. 1b). Importantly, the expression of ISGs in non-stimulated cells was equivalent to control cells suggesting that ISG15-dependent regulation is specific to the IFN response and not required for the regulation of basal gene expression. Further experiments showed that lack of ISG15 prolonged the longevity of ISG protein expression, which presumably has an impact on patients with autoinflammatory diseases associated with ISG15 loss-of-function. Here, control A549 and knockout cells were treated with IFN-α for 24 h. The cells were washed and media (without IFN-α) was then added. Cells were harvested every 24 h for 96 h and MxA expression was assessed by

immunoblotting (Fig. 1c). In control A549 cells MxA expression peaked at 24 h (the point at which

IFN was removed) and had returned to basal levels between 72 and 96 h. In knockout cells MxA

expression was clearly higher than in control cells, corroborating our mRNA analyses. Furthermore,

while MxA expression in A549-ISG15[-/-] did recede between 72 and 96 h, high protein levels remained

at 96 h. A dysregulated IFN response in ISG15-deficient cells is thought to be due to destabilisation

of USP18, a known negative regulator of JAK/STAT signalling (10). To determine if USP18 is similarly

affected in our cell lines, A549-ISG15[-/-] cells were treated with IFN-α for 24 or 48 h (or left untreated)

and whole cell lysates were probed for USP18 by immunoblotting. USP18 was robustly induced in

A549 cells following IFN-α treatment; however, levels of USP18 were much lower in IFN-α-treated

ISG15-deficient cells (Fig. 1d). *USP18* mRNA levels were approximately 10-fold higher in IFN-treated

ISG15-deficient cells compared to control A549s, demonstrating that reduced USP18 in A549-ISG15[-/-]

cells was not due to reduced transcription (Fig. 1b). Together, these data show that ISG15 is critical

for the regulated expression of ISGs. Moreover, they demonstrate that the effects of IFN treatment

on our ISG15 knockout A549 cell lines recapitulate the findings in cells derived from ISG15-deficent,

patient cells.

ISG15-deficiency leads to translational repression following IFN treatment

During our studies we observed that IFN-α-treatment of ISG15-knockout cells led to a reduction in

protein synthesis and reasoned that this was a likely contributor to the reported virus resistance in

ISG15-deficient cells (14). To investigate this we treated, or left untreated, A549 and A549-ISG15[-/-]

(B8) cells with IFN-α. At 24 h, 48 h and 72 h following treatment cells were pulse labelled with [35]S-

Methionine ([35]S-Met) for 1 h and the incorporation of [35]S-Met was analysed by phoshoimager

analysis. These data showed, compared to control cells, that there was a pronounced decrease in

protein synthesis in ISG15[-/-] cells between 24 h and 48 h (Fig. 2a). Furthermore, when cells were pre-

treated with IFN-α for 8 h, or left untreated, infected with the orthorubulavirus PIV5 (family

Paramyxoviridae, sub-family *Orthorubulavirinae*) at a MOI of 10 and then labelled for 1 h with ^{35}S-

Met at 24 h and 48 h p.i. (32 h and 56 h post IFN-α treatment, respectively), there was no evidence

of viral protein synthesis in treated A549-ISG15$^{-/-}$ cells, whereas in treated control cells, viral protein

synthesis was evident at 48 h p.i. (Fig. 2b). These data demonstrate that IFN-treatment of A549-

ISG15$^{-/-}$ cells led to inhibition of protein synthesis that was associated with viral resistance, at least at

later times.

A pre-existing antiviral state in ISG15-deficient cells renders them resistant to parainfluenza virus

infection

Previous studies have shown that IFN treatment of ISG15-deficient patient cells renders them

resistant to viral infection, including the murine respirovirus (family *Paramyxoviridae*, sub-family

Orthoparamyxovirinae) Sendai virus (14), and this seems to extend to PIV5 with our *in vitro* system

(Fig. 2b). To investigate this in A549-ISG15$^{-/-}$ cells, control A549 cells and the ISG15 knockout clones

described above were either untreated or treated with 1000 IU/ml IFN-α2b (the same concentration

and IFN-α type used in (14)) for 18 h. Cells were then infected with PIV5 (strain W3) (22) for 24 and

48 h and analysed by immunoblotting. In all cell lines, the levels of PIV5 nucleoprotein (NP)

expression was equivalent at 24 and 48 h in unstimulated cells (Fig. 3a). In IFN-α pre-treated control

cells, including C4+ that retained ISG15 expression, the level of NP expression was markedly reduced

at 24 h. By 48 h, the level of NP increased showing that infection had progressed even in the

presence of IFN-α (Fig. 3a). This is because the PIV5-V protein targets STAT1 for proteasomal

degradation, and once sufficient V is expressed, the IFN response is dismantled allowing the virus to

replicate (23). However, all A549-ISG15$^{-/-}$ cell lines that had been pre-treated with IFN-α were

resistant to PIV5 infection as shown by dramatically reduced, or even absent, NP expression at both

time points (Fig. 3a).

Previous reports have shown that the ISG15 regulation of IFN signalling is independent of its ability

to covalently modify proteins by ISGylation (10). To confirm this, we again applied CRISPR/Cas9

genome engineering technology and knocked out expression of UBA7, the E1 enzyme required for

ISGylation. For this we took a different approach compared to generating our ISG15 knockout cells

(19). Here, we introduced constitutive expression of Cas9 by lentiviral transduction of A549 cells and

transduced A549-Cas9 cells with lentiGuide-Puro lentivirus carrying a guide RNA specific for UBA7,

followed by single-cell cloning. We confirmed that all clones were UBA7-deficient by immunoblot

analysis, which demonstrated that they retained expression of ISG15 but had lost the ability to

ISGylate proteins (Fig. 3b). Additionally, following the scheme used in Fig. 3a, these cells were

infected with PIV5-W3. These data showed that, compared to ISG15 knockout cells that were

resistant to infection, all IFN-α-pre-treated UBA7-knockout cells were infected as efficiently as

control cells (Fig. 3b), confirming reports that ISG15-dependent regulation of type I IFN signalling

does not require ISGylation (10).

The direct antiviral activity of ISGs is responsible for virus resistance

Virus resistance can be induced following 8 h IFN-α treatment (shorter times were not tested), well

before any obvious effect on global protein synthesis (Fig. 2). Therefore, shutdown of translation is

unlikely to be the sole contributor to virus resistance at early time points and so we wished to

determine whether the direct antiviral activity of ISGs was responsible. Addressing this question is

complex since, for most viruses, the specific ISG(s) responsible for blocking replication is not known.

However, for PIV5, it has been established that IFIT1 is the principle ISG responsible for most of the

IFN-dependent antiviral activity (17, 26). We therefore hypothesised that if virus resistance was

caused by the direct antiviral activity of ISGs, knockdown of IFIT1 in ISG15-deficient cells would

permit PIV5 replication during an antiviral response. We reduced IFIT1 (according to (17)) in A549

and A549-ISG15$^{-/-}$ cells and all four cell lines (A549, A549-ISG15$^{-/-}$ and the respective shIFIT1 cells)

were pre-treated, or left untreated, with IFN-α and then infected with PIV5-W3 (MOI 10) for 24 and

48 h. Expression of PIV5 NP, analysed by quantitative immunoblotting, was used to measure virus

infection (Fig. 4a). IFIT1 levels and expression of ISG15 were likewise tested. Typically, pre-treatment

of naïve cells with IFN-α reduces infection, as shown by a reduction in NP levels, compared to non-

treated cells (Fig. 2b & 3a); nevertheless, because PIV5 expresses the IFN antagonist V protein, NP

levels reach similar levels to untreated cells by 48 h p.i. However, this IFN-dependent reduction in

virus infection is diminished when IFIT1 is knocked down, confirming earlier reports of IFIT1's

antiviral activity against PIV5 (17, 26). While IFN-α pre-treatment of A549-ISG15$^{-/-}$ cells renders them

resistant to infection, when IFIT1 was also knocked down, PIV5 infection was restored (Fig. 4a).

Because we performed quantitative immunoblotting of NP and β-Actin, we were able to quantify NP

levels, allowing us to analyse these changes statistically (Fig. 4b). These data show that in IFN-α-pre-

treated cells, knocking IFIT1 down restored NP to similar levels to those seen in untreated cells,

regardless of ISG15 status.

Rather than solely relying on viral protein expression as a surrogate for virus infection, we also

tested virus replication using a biologically relevant plaque assay (Fig. 4c). Cells were infected with

approximately 30-40 PFU of PIV5 stain CPI- (PIV5-CPI-) (27), a strain unable to block IFN signalling

due to a mutation in its V protein. Infected cells were fixed 6 d p.i. and stained for viral antigen. As

previously demonstrated (17), PIV5-CPI- was unable to efficiently form plaques in IFN-competent

A549 cells. However, PIV5-CPI- did replicate when cells were unable to produce IFN, such as in A549-

Npro cells that constitutively express bovine viral diarrhea virus (BVDV) Npro that cleaves IRF3 (a

transcription factor critical for IFN induction (21)). Furthermore, when IFIT1 was knocked down,

PIV5-CPI- was able to replicate (albeit less efficiently), further highlighting the major role of IFIT1 as

an anti-PIV5 protein. As expected, and like A549 cells, there was very little virus replication in A549-

ISG15$^{-/-}$ cells; however, when IFIT1 was knocked down, cells were able to support virus replication. It

must be noted however that virus replication in A549-ISG15$^{-/-}$/shIFIT1 cells did not recover to the

same degree as A549-shIFIT1 cells. We propose that the reason for this will be complex and may

include the likelihood that additional, yet to be identified, anti-PIV5 ISGs exist which are expressed at

higher levels in ISG15-deficient cells. Another possible explanation is the inhibition of protein

synthesis, including that of viral proteins, in ISG15-deficent cells; cells were infected for 6 days prior

to performing the plaque assays, a time point beyond that required to observe a significant effect on

protein synthesis (Fig. 2a). Therefore, the plaques observed in A549-ISG15$^{-/-}$/shIFIT1 cells likely result

from virus that replicated prior to the inhibition of global protein synthesis.

We also investigated infection of these cell lines with other paramyxoviruses whose sensitivity to

IFIT1 has been previously reported. Cells were treated with IFN-α and then infected with HPIV2

strain Colindale (MOI 10; family *Paramyxoviridae*, sub-family *Orthorubulavirinae*), which is reported

to be moderately sensitive to IFIT1-restriction (26), for 24 and 48 h (untreated cells were not

analysed because of high cytopathic effect in the absence of IFN). To investigate infection, we

detected expression of HPIV2 phosphoprotein by quantitative immunoblotting (Fig. 5a), which

showed that IFN-α-pre-treated A549-ISG15$^{-/-}$ cells were largely resistant to infection, although by 48

h p.i. there was some, albeit low level, evidence of viral protein accumulation. Nevertheless,

infection of A549-ISG15$^{-/-}$/shIFIT1 did allow significantly more viral protein expression. Quantitative

analyses demonstrated that viral protein accumulation in A549-ISG15$^{-/-}$/shIFIT1 cells was significantly

higher than in A549-ISG15$^{-/-}$ cells, but this was not as high as in A549 control cells, which agrees with

the reported partial sensitivity of HPIV2 to IFIT1 restriction indicating that additional ISGs target

HPIV2 (Fig. 5b). We performed a similar analysis with HPIV3 strain Washington (20) (family

Paramyxoviridae, sub-family *Orthoparamyxovirinae*), a virus reported to have limited sensitivity to

IFIT1 (26). Interestingly, pre-treatment of A549 and A549-shIFIT1 cells with IFN-α had less of an

effect on virus protein accumulation compared to the effects on PIV5 infection (Fig. 5c).

Furthermore, while infection of IFN-α-pre-treated ISG15 knockout cells significantly reduced

infection compared to control cells, virus infection in these cells was still more robust compared to

PIV5 and HPIV2-infected cells. Nevertheless, knockdown of IFIT1 only slightly increased HPIV3

protein expression in both ISG15-competent and ISG15-deficient cells (Fig. 5d), supporting reports of

a minor role of IFIT1 during the antiviral response to HPIV3 (26).

Our data have so far suggested that early virus resistance is mediated by the direct antiviral activity

of the IFN response. However, protein synthesis is reduced at later times post-IFN treatment and

this is likely to cause resistance; therefore, we investigated whether PIV5 resistance could be

induced independently of the direct antiviral activity of IFIT1. To do this we pre-treated the four cell

lines (A549, A549-shIFIT1, A549-ISG15$^{-/-}$ and A549-ISG15$^{-/-}$-shIFIT1) with IFN-α for different periods of

time, infected with a recombinant PIV5 that expresses the fluorescent protein mCherry (rPIV5-

mCherry) for 48 h (MOI 10) and measured fluorescence as a marker of virus replication (Fig. 6a).

Virus replication in A549 cells was equivalent regardless of the time cells had been pre-treated with

IFN-α and, as expected, A549-ISG15$^{-/-}$ cells were resistant to infection at any time post IFN-α

treatment (Fig. 6b). Any advantage to PIV5 replication as a result of IFIT1 knockdown in A549-shIFIT1

cells was lost when cells had been pre-treated for 16 h or more, as longer periods of pre-treatment

resulted in replication equivalent to IFN-pre-treated A549 cells. Similarly, PIV5 replication in A549-

ISG15$^{-/-}$-shIFIT1 cells was higher than A549 control cells, and equivalent to A549-shIFIT1 cells,

following 8 and 16 h pre-treatment; however, when cells were pre-treated for 24 h, replication was

lower than in A549 and A549-shIFIT1 cells. Interestingly, as the time of pre-treatment of A549-ISG15^{-}

$^{/-}$-shIFIT1 cells extended, virus replication reduced further until cells became resistant (e.g. at 60 h

and 72 h pre-treatment, Fig. 6b), which was not observed in A549 or A549-shIFIT1 cells. These data

suggest that cell permissiveness progressively reduced with longer times of IFN-α pre-treatment,

which correlated with the effects of IFN-α treatment on protein synthesis in ISG15-deficient cells

(Fig. 2).

A previous report demonstrated that ISG15-dependent stabilisation of USP18 was required to bring

about regulation of the type I IFN response and this was sufficient for these cells to once again be

infected (14). However, what aspects of the antiviral response was responsible for resistance was

not investigated. Taken together, these data strongly suggest that virus resistance in early IFN-

treated ISG15-deficient cells was caused by the direct antiviral activity of ISGs and not due to a lack

of permissiveness as a result of IFN-dependent inhibition of protein synthesis. Nevertheless, because

of the reduced protein synthesis in IFN-α-treated ISG15-deficient cells, cells later become non-

permissive to infection, even when key ISGs are eliminated.

Discussion

Previous work had shown that virus resistance was observed in cells that had been treated with IFN-α and then left to rest for 36 h prior to challenge (14). We had observed that IFN-α treatment of A549-ISG15$^{-/-}$ cells led to dramatic decreases in protein synthesis, particularly between 24 and 48 h; therefore, it was not clear whether the initially reported virus resistance was due to defects in translation (including of viral mRNAs) at the timepoint used in (14) or due to the direct antiviral activity of the IFN response. For most viruses, the specific ISG(s) with antiviral activity for a given virus is not known, making the latter difficult to discern; however, for PIV5, it is well established that IFIT1 is responsible for the majority of the antiviral response (17). To study this we generated A549-ISG15$^{-/-}$ cells and showed these cells recapitulated the effects observed in ISG15-deficient patient cells following treatment with IFN-α which included dysregulated ISG expression and reduced USP18 protein levels following IFN-α treatment (Fig. 1). Additionally, by knocking-out UBA7, the first enzyme in the ISGylation cascade, we showed that ISGylation is not required for a regulated response (Fig. 3b), confirming previous reports that 'free' ISG15 is required for regulation (10).

Using these cell lines in combination with a PIV5 infection model, we showed that infection of IFN-α-pre-treated ISG15-deficient cells in which IFIT1 had been knocked down restored infection, thus confirming that at early times post infection, resistance was indeed due to the direct antiviral activity of the IFN response. Furthermore, because IFIT1 blocks the translation of viral transcripts, our data suggest that IFN-treated A549-ISG15$^{-/-}$ cells were still susceptible to infection, allowing viral transcription to take place prior to IFIT1 restriction, and that ISG15 was unlikely to significantly regulate processes involved in entry. Nevertheless, if ISG15-deficient cells were treated for longer periods with IFN-α prior to infection they did become resistant, even when IFIT1 was knocked down, suggesting that at later times the inhibition of protein synthesis was the principal cause of resistance. These data suggest that the virus resistance reported by Speer et al. (14) was due to a

lack of permissiveness and not a result of the direct antiviral activity of the IFN response, although

different cells were used in that study.

The data here demonstrate that the mechanism of resistance is likely two-fold, depending on the

duration that cells are exposed to IFN-α. It is not currently possible to know which mechanism is

dominant in ISG15-deicient patients, but it is likely to be a combination of both. Nevertheless, virus

resistance results from a lack of IFN signalling control - as a consequence of ISG15-loss-of-function -

which would explain why ISG15-deficient patients were not more susceptible to severe infection.

This observation, therefore, cannot be used to support the notion that human ISG15 does not

possess direct antiviral activity, as proposed (14, 16). It is likely that many viruses will not be

sensitive to ISG15-dependent antiviral activity; however, this is true of many antiviral effectors. For

example, and as confirmed in this study, IFIT1 strongly restricts PIV5 infection, yet it has reduced

activity against HPIV2 and likely no activity against HPIV3 or human respiratory syncytial virus (26). It

is also true that several ISGs are often required to limit infection (6); therefore, if one antiviral

effector mechanism is absent (such as ISGylation), there is sufficient redundancy to avoid severe

effects of infection (redundancy that can complicate the investigation of specific antiviral

mechanisms in *in vitro* studies). Nevertheless, several human viruses have been shown to be

sensitive to ISGylation and many have evolved specific mechanisms to counteract antiviral

ISGylation, adding further weight to the argument that human ISG15 does have antiviral activity

(reviewed in (8)). Indeed, other than the handful of patients that have been found to lack ISG15

expression (10, 28), individuals will possess an intact IFN response where the antiviral activity of

ISG15 (and other effectors) will function, if the infecting virus is sensitive to it.

It was surprising that protein synthesis was so affected in ISG15-deficient cells following IFN

treatment. It is well established that inhibition of general protein translation is a key feature of the

antiviral response and this is through the actions of proteins such as PKR or PERK (PKR-like ER kinase)

(4). However, for PKR to be activated it must recognise dsRNA, which was absent in IFN-α-treated

cells. Similarly, PERK is activated upon endoplasmic reticulum stress which might be expected during a viral infection, but not following treatment with IFN alone. Previous reports have shown that carcinoembryonic antigen-related cell adhesion molecule 1 (CEACAM1) has antiviral activity against human cytomegalovirus, influenza virus and metapneumovirus by supressing mTOR-mediated protein synthesis (29, 30). The membrane protein CEACAM1 is induced by innate sensors such as TLR-4 (31) and IFI16 (30) and delivers inhibitory signals via SHP1 (haematopoietic cells) or SHP2 (epithelial and endothelial cells) phosphatase activity through CEACAM1 immunoreceptor tyrosine-based inhibitory motifs (ITIMs) (32). CEACAM1 expression is rapidly induced following activation of NF-κB and IRF1, but whether IFN-α alone (as used here) can induce it expression is not clear. The *IRF1* promoter possesses a single GAS element, but no IRES, and so its expression is induced by STAT1 homodimers (33). Type I IFN signalling predominantly leads to the formation of STAT1-STAT2 heterodimers that associate with IRF9 (to form the ISGF3 transcription factor) to drive expression of ISGs that possess IRES elements in their promoters; however, STAT1 homodimers are formed after type I IFN treatment, but these are at lower concentrations. It is possible that 'late' inhibition of protein synthesis in ISG15-deficient cells (compared to the swifter antiviral activity of IRES-containing genes such as IFIT1) may relate to the kinetics of CEACAM1 expression as the accumulation of STAT1 homodimers is required to drive the expression of *IRF1*, that itself needs to be translated before it induces *CEACAM1*. Of course, the accumulation of STAT1 homodimers may be higher in ISG15-deficient cells because of a dysregulated type I IFN response. Nevertheless, it is plausible that the overamplified type I IFN response in ISG15-deficient cells led to high levels of CEACAM1 (compared to control cells) resulting in inhibition of protein synthesis. Moreover, ISG15 may have yet-to-be characterised functions in regulating the cellular response to stressors that lead to inhibition of protein synthesis.

It has been reported that ISG15 has a role in regulating the cell cycle through its interactions with SKP2 and USP18, although experiments in that study were not performed in IFN-treated cells, nor were ISG15 knockout cells tested (15). While rates of protein synthesis differ during different stages

of the cell cycle, translation is thought to be lowest during mitosis (34). Perturbation of the ISG15-

SKP2-USP18 axis following ablation of USP18 led to a delayed progression from G1 to S phase which

is not generally thought to be associated with translational repression (35). Of note, we have not

observed any obvious differences in cell growth in non-treated A549-ISG15$^{-/-}$ cells. Further work is

required to dissect the mechanism responsible for ISG15's effects on general protein translation

during an antiviral response.

ISG15 has emerged as a central regulator of immunity. It is a pleotropic protein that is strongly

expressed following activation of innate immune sensors and connects innate and adaptive

immunity. In this study, we have shown that a lack of ISG15 leads to virus resistance by two

kinetically distinct mechanisms; the rapid induction of antiviral ISGs and the unexpected effects on

protein synthesis. Our newly developed cell lines and infection model will pave the way for further

studies investigating the regulatory mechanisms of ISG15 during the antiviral response.

Acknowledgments

This work was supported by grants from the Academy of Medical Sciences (SBF003/1028 to DJH),

Wellcome Trust Institutional Strategic Support Fund (to DJH), Wellcome Trust (101788/Z/13/Z to

RER) and the UK Medical Research Council (MC_UU_12014/1 to JM and CGGB). We are grateful to

ERASMUS+ for supporting DH.

 Figure Legends

Fig. 1. Functional characterisation of A549 ISG15 knockout cell lines. (a) CRSIPR/Cas9 genome editing

was used to knockout ISG15 expression in A549 cells followed by single-cell cloning (following

previously reported procedures (18)). Four independent clones were treated with 1000 IU ml-1 IFN-

α for 24 and 48 h, or left untreated, and protein expression was tested by immunoblot analysis of

ISG15, MxA and β-Actin. 'Control' cells were naïve A549 cells. Representative image from two

independent experiments. (b) A549 and A5549-ISG15$^{-/-}$ (clone B8) were treated with 1000 IU/ml for

24 h. Expression of interferon stimulated genes was tested using RT-qPCR with primers specific for

HERC5, USP18, IFIT1 and *MxA*. Relative expression was determined following SYBR green qPCR using

ΔΔCt method. *B-Actin* expression was used to normalise between samples. Error bars represent the

standard deviation of the mean from three independent RNA samples. (c) A549 and A5549-ISG15$^{-/-}$

(clone B8) were treated with 1000 IU/ml for 24 h. Cells were washed and fresh media (without IFN-

α) was replaced. Cells were processed for immunoblot analysis using antibodies specific for MxA and

β-Actin at 24 h post IFN-α and every 24 h thereafter for 96 h. Controls were cells without IFN-α.

Representative image from two independent experiments. (d) A549 and A5549-ISG15$^{-/-}$ (clone B8)

were treated with 1000 IU/ml for 24 h and 48 h (or left untreated). Whole cell lysates were analysed

by immunoblotting with antibodies specific for USP18, ISG15 and β-Actin. Image is representative of

>3 independent experiments.

Fig. 2. Analysis of cellular and viral protein synthesis in ISG15-deficient cells during an antiviral state.

(a) Sub-confluent A549 and A549-ISG15$^{-/-}$ (B8) cells were treated with 1000 IU ml^{-1} IFN-α or left

untreated. At 24 h, 48 h and 72 h cells were pulsed for 1 h with L-[^{35}S]-Methionine (^{35}S-Met) in Met-

free media to metabolically label nascent proteins. Proteins were resolved by SDS-PAGE and stained

with Coomassie to ensure equal loading. Labelled proteins were visualised by phophoimager

analysis. (b) A549 and A549-ISG15$^{-/-}$ (B8) cells were treated with 1000 IU ml^{-1} IFN-α for 8 h or left

untreated and then infected with PIV5 strain W3 (MOI = 10). At 24 or 48 h post infection cells pulsed

and processed as in (a). Both experiments were performed independently at least twice.

Fig. 3. A pre-existing antiviral state in ISG15-deficient cells leads to virus resistance which is

independent of ISGylation. (a) Control (naïve A549) and 4 independent clones of A549-ISG15$^{-/-}$ cells

generated by CRISPR/Cas9 genome editing were treated with 1000 IU/ml IFN-α for 16 h or left

untreated and then infected with PIV5 strain W3 (MOI = 10). Cells were harvested at 24 h and 48 h

p.i. and processed for immunoblot analysis using antibodies specific for PIV5 nucleoprotein (NP),

ISG15, MxA and β-Actin. This experiment was independently performed twice. (b) UBA7 knockout

cells were generated using CRISPR/Cas9 genome editing; Cas9-expressing A549 cells were first

generated (following transduction with lentiCas9-Blast) and then transduced with lentiGuide-Puro

expressing a single guide RNA that targeted exon 3 of the UBA7 gene. Knockout cells were single-cell

cloned and three were selected for further analysis. These cells were treated with IFN-α, infected

and processed as in (a) using antibodies specific for PIV5 NP, ISG15, UBA7 and β-Actin. This

experiment was independently performed twice.

Fig. 4. Direct antiviral activity of ISGs is responsible for virus resistance due to ISG15-loss-of-function.

(a) IFIT1 was constitutively knocked down in A549 or A549-ISG15$^{-/-}$ (B8) cells following a previously

described method (17). A549, A549-ISG15$^{-/-}$ (B8) and the corresponding IFIT1 knockdown cells were

treated with IFN-α, infected with PIV5 and processed as in Fig. 3a. Following immunoblotting with

specific antibodies, PIV5 NP and β-Actin were detected using near-infrared (NIR) dye-conjugated

secondary antibodies to facilitate quantification. IFIT1 and ISG15 proteins were detected using

chemiluminescence following incubation with horseradish peroxidase (HRP)-conjugated secondary

antibodies. (b) Experiments described in (a) were performed independently three times (infections

were performed on three separate occasions) and NP and β-Actin levels were quantified using Image

Studio software (LiCOR). Signals were relative to those generated from IFN-α-treated A549 cells

infected for 48 h p.i. (set to 100%). Error bars represent the standard deviation of the mean from the

three experiments. ** indicates P < 0.01 using Student's t test. (c) Indicated cells were infected for 1

h with 30 – 40 plaque forming units (PFU) of PIV5 (CPI-), a strain unable to block the IFN response

due to mutation in V. Monolayers were fixed 6 d p.i. Plaques were detected using a pool of anti-PIV5

antibodies specific for hemagglutinin (HN), nucleoprotein (NP), phosphoprotein (P) and matrix

protein (M) (see (24)). Plaque assays were performed on 3 independent occasions.

Fig. 5. Restoration of paramyxovirus infection in IFN-α-pretreated ISG15-/- cells reflects their

reported sensitivity to IFIT1. (a-b) Experiments were performed as in (Fig. 4a-b). (a) HPIV2 proteins

were detected with antibodies specific for HPIV2 P (clone 161; (24)) and NIR-conjugated secondary

antibodies. Asterisks denotes non-specific band, arrow denotes HPIV2 P. Samples not treated with

IFN-α were omitted due to the highly lytic nature of HPIV2 which hampered their accurate

quantification. (b) Quantification of normalised NP signals and compared to the 48 h p.i. sample that

was set to 100%. (c) HPIV3 NP proteins were detected using antibodies specific for HPIV3 NP and

NIR-conjugated secondary antibodies. (d) Normalised signals were quantified as in (b) and compared

to IFN-α-treated, 48 h p.i. samples (set to 100%). Means and standard deviations were derived from

three independent experiments performed on different occasions. Asterisks denote statistical

significance (* P < 0.05, ** P < 0.01, *** P < 0.001 using Student's t test).

Fig. 6. Virus resistance is induced in ISG15-deficient cells following longer periods of IFN-α pre-

treatment. (a) Experimental workflow. (b) Cells were treated 1000 IU ml^{-1} IFN-α in 6-well plates for

the indicated times prior to infection. Pre-treated cells were infected with rPIV5-mCherry (MOI 10)

for 48 h and mCherry fluorescence was measured using an IncuCyte ZOOM. Background

fluorescence from mock-infected wells was subtracted. Data are representative to two independent

experiments.

References

1. McFadden, M.J., N.S. Gokhale, and S.M. Horner, Protect this house: cytosolic sensing of
viruses. *Curr Opin Virol*, 2017. **22**: p. 36-43.
2. Kawai, T. and S. Akira, The role of pattern-recognition receptors in innate immunity: update
on Toll-like receptors. *Nat Immunol*, 2010. **11**(5): p. 373-84.
3. Randall, R.E. and S. Goodbourn, Interferons and viruses: an interplay between induction,
signalling, antiviral responses and virus countermeasures. *J Gen Virol*, 2008. **89**(Pt 1): p. 1-47.
4. Dalet, A., E. Gatti, and P. Pierre, Integration of PKR-dependent translation inhibition with
innate immunity is required for a coordinated anti-viral response. *FEBS Lett*, 2015. **589**(14):
p. 1539-45.
5. Mears, H.V. and T.R. Sweeney, Better together: the role of IFIT protein-protein interactions
in the antiviral response. *J Gen Virol*, 2018. **99**(11): p. 1463-1477.
6. Schoggins, J.W., S.J. Wilson, M. Panis, M.Y. Murphy, C.T. Jones, P. Bieniasz, and C.M. Rice, A
diverse range of gene products are effectors of the type I interferon antiviral response.
Nature, 2011. **472**(7344): p. 481-5.
7. Yu, Z.X. and H.M. Song, Toward a better understanding of type I interferonopathies: a brief
summary, update and beyond. *World J Pediatr*, 2019.
8. Perng, Y.C. and D.J. Lenschow, ISG15 in antiviral immunity and beyond. *Nat Rev Microbiol*,
2018. **16**(7): p. 423-439.
9. Malakhov, M.P., O.A. Malakhova, K.I. Kim, K.J. Ritchie, and D.E. Zhang, UBP43 (USP18)
specifically removes ISG15 from conjugated proteins. *J Biol Chem*, 2002. **277**(12): p. 9976-81.
10. Zhang, X., D. Bogunovic, B. Payelle-Brogard, V. Francois-Newton, S.D. Speer, C. Yuan, S.
Volpi, Z. Li, O. Sanal, D. Mansouri, I. Tezcan, G.I. Rice, C. Chen, N. Mansouri, S.A. Mahdaviani,
Y. Itan, B. Boisson, S. Okada, L. Zeng, X. Wang, H. Jiang, W. Liu, T. Han, D. Liu, T. Ma, B. Wang,
M. Liu, J.Y. Liu, Q.K. Wang, D. Yalnizoglu, L. Radoshevich, G. Uze, P. Gros, F. Rozenberg, S.Y.
Zhang, E. Jouanguy, J. Bustamante, A. Garcia-Sastre, L. Abel, P. Lebon, L.D. Notarangelo, Y.J.
Crow, S. Boisson-Dupuis, J.L. Casanova, and S. Pellegrini, Human intracellular ISG15 prevents
interferon-alpha/beta over-amplification and auto-inflammation. *Nature*, 2015. **517**(7532):
p. 89-93.
11. Francois-Newton, V., G. Magno de Freitas Almeida, B. Payelle-Brogard, D. Monneron, L.
Pichard-Garcia, J. Piehler, S. Pellegrini, and G. Uze, USP18-based negative feedback control is
induced by type I and type III interferons and specifically inactivates interferon alpha
response. *PLoS One*, 2011. **6**(7): p. e22200.
12. Malakhova, O.A., K.I. Kim, J.K. Luo, W. Zou, K.G. Kumar, S.Y. Fuchs, K. Shuai, and D.E. Zhang,
UBP43 is a novel regulator of interferon signaling independent of its ISG15 isopeptidase
activity. *EMBO J*, 2006. **25**(11): p. 2358-67.
13. Sarasin-Filipowicz, M., X. Wang, M. Yan, F.H. Duong, V. Poli, D.J. Hilton, D.E. Zhang, and M.H.
Heim, Alpha interferon induces long-lasting refractoriness of JAK-STAT signaling in the
mouse liver through induction of USP18/UBP43. *Mol Cell Biol*, 2009. **29**(17): p. 4841-51.
14. Speer, S.D., Z. Li, S. Buta, B. Payelle-Brogard, L. Qian, F. Vigant, E. Rubino, T.J. Gardner, T.
Wedeking, M. Hermann, J. Duehr, O. Sanal, I. Tezcan, N. Mansouri, P. Tabarsi, D. Mansouri,
V. Francois-Newton, C.F. Daussy, M.R. Rodriguez, D.J. Lenschow, A.N. Freiberg, D. Tortorella,
J. Piehler, B. Lee, A. Garcia-Sastre, S. Pellegrini, and D. Bogunovic, ISG15 deficiency and
increased viral resistance in humans but not mice. *Nat Commun*, 2016. **7**: p. 11496.
15. Vuillier, F., Z. Li, P.H. Commere, L.T. Dynesen, and S. Pellegrini, USP18 and ISG15
coordinately impact on SKP2 and cell cycle progression. *Sci Rep*, 2019. **9**(1): p. 4066.
16. Hermann, M. and D. Bogunovic, ISG15: In Sickness and in Health. *Trends Immunol*, 2017.
38(2): p. 79-93.

17. Andrejeva, J., H. Norsted, M. Habjan, V. Thiel, S. Goodbourn, and R.E. Randall, ISG56/IFIT1 is
primarily responsible for interferon-induced changes to patterns of parainfluenza virus type
5 transcription and protein synthesis. *J Gen Virol*, 2013. **94**(Pt 1): p. 59-68.
18. Domingues, P., C.G. Bamford, C. Boutell, and J. McLauchlan, Inhibition of hepatitis C virus
RNA replication by ISG15 does not require its conjugation to protein substrates by the
HERC5 E3 ligase. *J Gen Virol*, 2015. **96**(11): p. 3236-42.
19. Sanjana, N.E., O. Shalem, and F. Zhang, Improved vectors and genome-wide libraries for
CRISPR screening. *Nat Methods*, 2014. **11**(8): p. 783-784.
20. Durbin, A.P., J.M. McAuliffe, P.L. Collins, and B.R. Murphy, Mutations in the C, D, and V open
reading frames of human parainfluenza virus type 3 attenuate replication in rodents and
primates. *Virology*, 1999. **261**(2): p. 319-30.
21. Hilton, L., K. Moganeradj, G. Zhang, Y.H. Chen, R.E. Randall, J.W. McCauley, and S.
Goodbourn, The NPro product of bovine viral diarrhea virus inhibits DNA binding by
interferon regulatory factor 3 and targets it for proteasomal degradation. *J Virol*, 2006.
80(23): p. 11723-32.
22. Choppin, P.W., Multiplication of a Myxovirus (Sv5) with Minimal Cytopathic Effects and
without Interference. *Virology*, 1964. **23**: p. 224-33.
23. Didcock, L., D.F. Young, S. Goodbourn, and R.E. Randall, The V protein of simian virus 5
inhibits interferon signalling by targeting STAT1 for proteasome-mediated degradation. *J
Virol*, 1999. **73**(12): p. 9928-33.
24. Randall, R.E., D.F. Young, K.K. Goswami, and W.C. Russell, Isolation and characterization of
monoclonal antibodies to simian virus 5 and their use in revealing antigenic differences
between human, canine and simian isolates. *J Gen Virol*, 1987. **68 (Pt 11)**: p. 2769-80.
25. Rydbeck, R., C. Orvell, A. Love, and E. Norrby, Characterization of four parainfluenza virus
type 3 proteins by use of monoclonal antibodies. *J Gen Virol*, 1986. **67 (Pt 8)**: p. 1531-42.
26. Young, D.F., J. Andrejeva, X. Li, F. Inesta-Vaquera, C. Dong, V.H. Cowling, S. Goodbourn, and
R.E. Randall, Human IFIT1 Inhibits mRNA Translation of Rubulaviruses but Not Other
Members of the Paramyxoviridae Family. *J Virol*, 2016. **90**(20): p. 9446-56.
27. Chen, C., R.W. Compans, and P.W. Choppin, Parainfluenza virus surface projections:
glycoproteins with haemagglutinin and neuraminidase activities. *J Gen Virol*, 1971. **11**(1): p.
53-8.
28. Bogunovic, D., M. Byun, L.A. Durfee, A. Abhyankar, O. Sanal, D. Mansouri, S. Salem, I.
Radovanovic, A.V. Grant, P. Adimi, N. Mansouri, S. Okada, V.L. Bryant, X.F. Kong, A. Kreins,
M.M. Velez, B. Boisson, S. Khalilzadeh, U. Ozcelik, I.A. Darazam, J.W. Schoggins, C.M. Rice, S.
Al-Muhsen, M. Behr, G. Vogt, A. Puel, J. Bustamante, P. Gros, J.M. Huibregtse, L. Abel, S.
Boisson-Dupuis, and J.L. Casanova, Mycobacterial disease and impaired IFN-gamma
immunity in humans with inherited ISG15 deficiency. *Science*, 2012. **337**(6102): p. 1684-8.
29. Diab, M., A. Vitenshtein, Y. Drori, R. Yamin, O. Danziger, R. Zamostiano, M. Mandelboim, E.
Bacharach, and O. Mandelboim, Suppression of human metapneumovirus (HMPV) infection
by the innate sensing gene CEACAM1. *Oncotarget*, 2016. **7**(41): p. 66468-66479.
30. Vitenshtein, A., Y. Weisblum, S. Hauka, A. Halenius, E. Oiknine-Djian, P. Tsukerman, Y.
Bauman, Y. Bar-On, N. Stern-Ginossar, J. Enk, R. Ortenberg, J. Tai, G. Markel, R.S. Blumberg,
H. Hengel, S. Jonjic, D.G. Wolf, H. Adler, R. Kammerer, and O. Mandelboim, CEACAM1-
Mediated Inhibition of Virus Production. *Cell Rep*, 2016. **15**(11): p. 2331-9.
31. Muenzner, P., M. Naumann, T.F. Meyer, and S.D. Gray-Owen, Pathogenic Neisseria trigger
expression of their carcinoembryonic antigen-related cellular adhesion molecule 1
(CEACAM1; previously CD66a) receptor on primary endothelial cells by activating the
immediate early response transcription factor, nuclear factor-kappaB. *J Biol Chem*, 2001.
276(26): p. 24331-40.
32. Gray-Owen, S.D. and R.S. Blumberg, CEACAM1: contact-dependent control of immunity. *Nat
Rev Immunol*, 2006. **6**(6): p. 433-46.

33.   Michalska, A., K. Blaszczyk, J. Wesoly, and H.A.R. Bluyssen, A Positive Feedback Amplifier
Circuit That Regulates Interferon (IFN)-Stimulated Gene Expression and Controls Type I and
Type II IFN Responses. *Front Immunol*, 2018. **9**: p. 1135.
34.   Polymenis, M. and R. Aramayo, Translate to divide: small es, Cyrillicontrol of the cell cycle by
protein synthesis. *Microb Cell*, 2015. **2**(4): p. 94-104.
35.   Stumpf, C.R., M.V. Moreno, A.B. Olshen, B.S. Taylor, and D. Ruggero, The translational
landscape of the mammalian cell cycle. *Mol Cell*, 2013. **52**(4): p. 574-82.

Figure 1. Functional characterisation of A549 ISG15 knockout cell lines. (a) CRSIPR/Cas9 genome

editing was used to knockout ISG15 expression in A549 cells followed by single-cell cloning

(following previously reported procedures (18)). Four independent clones were treated with 1000 IU

ml-1 IFN-α for 24 and 48 h, or left untreated, and protein expression was tested by immunoblot

analysis of ISG15, MxA and β-Actin. 'Control' cells were naïve A549 cells. Representative image from

two independent experiments. (b) A549 and A5549-ISG15$^{-/-}$ (clone B8) were treated with 1000 IU/ml

for 24 h. Expression of interferon stimulated genes was tested using RT-qPCR with primers specific

for *HERC5, USP18, IFIT1* and *MxA*. Relative expression was determined following SYBR green qPCR

using ΔΔCt method. *B-Actin* expression was used to normalise between samples. Error bars

represent the standard deviation of the mean from three independent RNA samples. (c) A549 and

A5549-ISG15$^{-/-}$ (clone B8) were treated with 1000 IU ml^{-1} for 24 h. Cells were washed and fresh

media (without IFN-α) was replaced. Cells were processed for immunoblot analysis using antibodies

specific for MxA and β-Actin at 24 h post IFN-α and every 24 h thereafter for 96 h. Controls were

cells without IFN-α. Representative image from two independent experiments. (d) A549 and A5549-

ISG15$^{-/-}$ (clone B8) were treated with 1000 IU ml^{-1} for 24 h and 48 h (or left untreated). Whole cell

lysates were analysed by immunoblotting with antibodies specific for USP18, ISG15 and β-Actin.

Image is representative of >3 independent experiments.

Figure 2. Analysis of cellular and viral protein synthesis in ISG15-deficient cells during an antiviral

state. (a) Sub-confluent A549 and A549-ISG15[-/-] (B8) cells were treated with 1000 IU ml-1 IFN-α or

left untreated. At 24 h, 48 h and 72 h cells were pulsed for 1 h with L-[^{35}S]-Methionine (^{35}S-Met) in

Met-free media to metabolically label nascent proteins. Proteins were resolved by SDS-PAGE and

stained with Coomassie to ensure equal loading. Labelled proteins were visualised by phophoimager

analysis. (b) A549 and A549-ISG15[-/-] (B8) cells were treated with 1000 IU/ml IFN-α for 8 h or left

untreated and then infected with PIV5 strain W3 (MOI = 10). At 24 or 48 h post infection cells pulsed

and processed as in (a). Both experiments were performed independently at least twice.

Figure 3. A pre-existing antiviral state in ISG15-deficient cells leads to virus resistance which is

independent of ISGylation. (a) Control (naïve A549) and 4 independent clones of A549-ISG15$^{-/-}$ cells

generated by CRISPR/Cas9 genome editing were treated with 1000 IU/ml IFN-α for 16 h or left

untreated and then infected with PIV5 strain W3 (MOI = 10). Cells were harvested at 24 h and 48 h

p.i. and processed for immunoblot analysis using antibodies specific for PIV5 nucleoprotein (NP),

ISG15, MxA and β-Actin. This experiment was independently performed twice. (b) UBA7 knockout

cells were generated using CRISPR/Cas9 genome editing; Cas9-expressing A549 cells were first

generated (following transduction with lentiCas9-Blast) and then transduced with lentiGuide-Puro

expressing a single guide RNA that targeted exon 3 of the UBA7 gene. Knockout cells were single-cell

cloned and three were selected for further analysis. These cells were treated with IFN-α, infected

and processed as in (a) using antibodies specific for PIV5 NP, ISG15, UBA7 and β-Actin. This

experiment was independently performed twice.

Figure 4. Direct antiviral activity of ISGs is responsible for virus resistance due to ISG15-loss-of-

function. (a) IFIT1 was constitutively knocked down in A549 or A549-ISG15[-/-] (B8) cells following a

previously described method [17]. A549, A549-ISG15[-/-] (B8) and the corresponding IFIT1 knockdown

cells were treated with IFN-α, infected with PIV5 and processed as in Fig. 3a. Following

immunoblotting with specific antibodies, PIV5 NP and β-Actin were detected using near-infrared

(NIR) dye-conjugated secondary antibodies to facilitate quantification. IFIT1 and ISG15 proteins were

detected using chemiluminescence following incubation with horseradish peroxidase (HRP)-

conjugated secondary antibodies. (b) Experiments described in (a) were performed independently

three times (infections were performed on three separate occasions) and NP and β-Actin levels were

quantified using Image Studio software (LiCOR). Signals were relative to those generated from IFN-α-

treated A549 cells infected for 48 h p.i. (set to 100%). Error bars represent the standard deviation of

the mean from the three experiments. ** indicates P < 0.01 using Student's t test. (c) Indicated cells

were infected for 1 h with 30 – 40 plaque forming units (PFU) of PIV5 (CPI-), a strain unable to block

the IFN response due to mutation in V. Monolayers were fixed 6 d p.i. Plaques were detected using a

pool of anti-PIV5 antibodies specific for hemagglutinin (HN), nucleoprotein (NP), phosphoprotein (P)

and matrix protein (M) (see [24]). Plaque assays were performed on 3 independent occasions.

Figure 5. Restoration of paramyxovirus infection in IFN-α-pretreated ISG15[-/-] cells reflects their

reported sensitivity to IFIT1. (a-b) Experiments were performed as in (Fig. 4a-b). (a) HPIV2 proteins

were detected with antibodies specific for HPIV2 P (clone 161; [24]) and NIR-conjugated secondary

antibodies. Asterisks denotes non-specific band, arrow denotes HPIV2 P. Samples not treated with

IFN-α were omitted due to the highly lytic nature of HPIV2 which hampered their accurate

quantification. (b) Quantification of normalised NP signals and compared to the 48 h p.i. sample that

was set to 100%. (c) HPIV3 NP proteins were detected using antibodies specific for HPIV3 NP and

NIR-conjugated secondary antibodies. (d) Normalised signals were quantified as in (b) and compared

to IFN-α-treated, 48 h p.i. samples (set to 100%). Means and standard deviations were derived from

three independent experiments performed on different occasions. Asterisks denote statistical

significance (* P < 0.05, ** P < 0.01, *** P < 0.001 using Student's t test).

Figure 6. Virus resistance is induced in ISG15-deficient cells following longer periods of IFN-α pre-

treatment. (a) Experimental workflow. (b) Cells were treated 1000 IU ml-1 IFN-α in 6-well plates for

the indicated times prior to infection. Pre-treated cells were infected with rPIV5-mCherry (MOI 10)

for 48 h and mCherry fluorescence was measured using an IncuCyte ZOOM. Background

fluorescence from mock-infected wells was subtracted. Data are representative to two independent

experiments.