Evolutionary reticulation makes an important contribution to Amazonian tree diversity

Rowan J. Schley[1,2*], R. Toby Pennington[3,4], Oscar Alejandro Pérez-Escobar[1], Andrew J. Helmstetter[5], Manuel de la Estrella[6], Isabel Larridon[1,7], Izai Alberto Bruno Sabino Kikuchi[1,8], Timothy Barraclough[2], Félix Forest[1], and Bente Klitgård[1]

[1]*Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK*

[2]*Department of Life Sciences, Imperial College London, Silwood Park, Ascot, Berkshire, UK, SL5 7PY*

[3] *Geography, University of Exeter, Laver Building, North Park Road, Exeter EX4 4QE*

[4]*Royal Botanic Garden Edinburgh, 20a Inverleith Row, Edinburgh EH3 5LR*

[5]*Institut de Recherche pour le Développement (IRD), UMR-DIADE, BP 64501, F-34394 Montpellier cedex 5, France.*

[6]*Departamento de Botánica, Ecología y Fisiología Vegetal, Facultad de Ciencias, Campus de Rabanales, Universidad de Córdoba, 14071, Córdoba, Spain*

[7] *Systematic and Evolutionary Botany Lab, Department of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000 Gent, Belgium*

[8]*Universiteit Leiden, Hortus botanicus Leiden, PO Box 9500, Leiden, 2300 RA, The Netherlands*

*Corresponding author at: *Room E.2.6, Herbarium, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AE, UK*

E-mail address: *r.schley@kew.org* *(R. J. Schley).*

Abstract

Hybridization has the potential to generate or homogenize biodiversity and is a particularly common phenomenon in plants, with an estimated 25% of species undergoing inter-specific gene flow. However, hybridization has rarely been demonstrated among tree species in Amazonia, the world's largest rainforest, and its role in diversification remains enigmatic. We show that within *Brownea*, a characteristic tree genus of Amazonia, there is extensive evidence of hybridization. Using both phylogenomic and population genomic approaches we find multiple historical hybridization events within *Brownea,* along with contemporary hybridization among co-occurring species. Finally, we infer homogeneous rates of gene flow among different genomic loci, indicating a lack of selection against hybrids, reflecting their persistence over time. These results demonstrate that gene flow between Amazonian tree species has occurred across temporal scales, which contrasts with previous

work on Amazonian trees, and may have contributed to the evolution of the most diverse tree flora on

Earth.

<u>Main</u>

Reproductive isolation is often seen as a prerequisite for speciation and as a defining feature of

species [1, 2]. Despite this, hybridization between species is known to occur, and has several different

outcomes, from the erosion of evolutionary divergence [3, 4] to the formation of entirely new 'hybrid

species' [5]. Neotropical rainforests harbour the highest levels of plant diversity on Earth [6], and to date

there has been little convincing evidence that hybridization consistently occurs between tree species

therein. Indeed, the prevailing view has been that hybridization between tropical tree species is an

exceptionally rare event [7-9]. Although a few observations of reproductive biology support this, such as

the intersterility between lineages of *Inga*, a species-rich genus in the legume family [10], evidence for

hybridization has been poorly tested empirically in tropical floras. The only potential example of

hybridization among neotropical trees which has been documented using DNA sequence data

involves two species of *Carapa* [11] found in Amazonia, the largest expanse of rainforest in the world

which contains at least 6,800 tree species [12].

One factor that suggests hybridization might occur more frequently in neotropical rainforest tree

species, if their reproductive isolation is not absolute, is the remarkable level of sympatry found for

closely related species. In several neotropical lineages, including *Inga*, *Guatteria* and *Swartzia*, many

recently diverged species co-occur [13], often to a remarkable degree. One such example of this is the

co-occurrence of 19 *Inga* species in a single hectare of the Ecuadorian Amazon [14]. As such, for many

rainforest taxa the opportunity for hybridization is constantly present.

Hybridization can have a range of evolutionary consequences. In many cases, hybridization simply

results in the formation of sterile, maladapted offspring with poor reproductive fitness, allowing

genetic isolation between species to be maintained (i.e., reinforcement [15]). In other cases, there may be

a permanent movement of genetic material from one lineage to another [16], which is known as

'introgression'. This transfer of genetic material through hybridization may confer a selective

advantage to resultant offspring [17], in which case it is referred to as 'adaptive introgression'. Adaptive

introgression is most often observed between closely-related taxa during the invasion of new habitats

[18, 19]. Furthermore, hybridization can lead to rapid evolutionary radiations. This occurs through the re-

assembly of standing genetic variation which has accumulated between diverging lineages, and which

has already been subject to selection. This 'combinatorial' process is much more rapid than the

gradual accumulation of variation through mutation, and the passage of these variants through

hybridization often fuels rapid diversification events [20]. This has been demonstrated in a wide range of

taxa, including sunflowers, African cichlid fish and Darwin's finches [21-23].

Introgression can occur at different rates in different regions of the genome [24]. Regions under

divergent selection may remain distinct due to reduced fitness of hybrid genotypes at such loci,

resulting in a low rate of introgression. Conversely, regions under little or no selection tend to

introgress more freely, becoming homogenised between species. Moreover, if there is selection for

hybrid genotypes (as in adaptive introgression), the rate of introgression for a region may be further

increased relative to the rest of the genome [25]. Such a process has been demonstrated in temperate-

zone tree species, where divergence between hybridizing lineages is maintained through

environmentally-driven selection [26, 27]. This explains why species that hybridize can remain as

biologically distinct (and taxonomically identifiable) groups despite undergoing genetic exchange

with other species [28, 29].

Brownea, a member of the pea and bean family (Fabaceae/Leguminosae), is a characteristic tree

genus of lowland neotropical rainforests, and contains around 27 species distributed across northern

South America. Previous work indicates that there is a broad degree of phylogenetic incongruence

evident in this genus [30] which might indicate hybridization, although this could also result from

incomplete lineage sorting (the differential inheritance of genetic variation from a common ancestral

species in different descendent lineages). However, there are numerous *Brownea* hybrids in

cultivation (e.g. *B. x crawfordii* [31] and *B. hybrida* [32]), as well as several instances of putative

hybridization among *Brownea* lineages in the wild. The most notable of these instances is that

proposed between the range-restricted *Brownea jaramilloi* [33] and the wide-ranging *B. grandiceps*,

which co-occur in the Ecuadorian Amazon (Fig. 1). There are multiple morphological distinctions

between these two species, including differences in inflorescence colour and structure, growth habit,

tree height and leaf morphology [33]. Although they co-occur, these species favour different habitats: *B.*

jaramilloi grows on ridge tops and hillsides, whereas *B. grandiceps* shows a slight preference for

swamps and valleys but is more evenly distributed [33, 34]. Despite this, hybridization appears to occur,

as evidenced by the existence of a putative hybrid between these two species known as *B.* "rosada"

(Fig. 1). *Brownea "rosada"* displays an intermediate morphology between its two parental species,

producing pink flowers. The hypothesis of a *B. jaramilloi* x *B. grandiceps* hybrid has not yet been

tested, however, using molecular data.

Figure 1: Two co-occurring *Brownea* lineages (*Brownea grandiceps* (photograph © Rowan Schley)

and *Brownea jaramilloi* (photograph © Xavier Cornejo)) as well as their putative hybrid *Brownea*

"rosada" (photograph © J. L. Clark).

As a member of the legume family, which dominates Amazonian forests [35], and with its apparent

propensity for hybridization, *Brownea* is an excellent system with which to study the phylogenetic

patterns and genomic architecture of introgression in neotropical rainforest trees. Systematically

documenting hybridization at a range of time scales and taxonomic levels within this group could

reveal how admixture has contributed to the assembly of one of the world's richest floras.

Accordingly, this investigation aims to answer the following questions:

1. Is there evidence of hybridization at a deep phylogenetic level in this lineage of neotropical rainforest tree?

2. Is there evidence of more recent gene flow, and if so, does this occur evenly across most of the genome?

Results

Phylogenetic analysis

Using DNA sequence data from 226 nuclear gene regions, we produced a 'species' tree using *ASTRAL* [36] (Fig. 2) based on gene trees inferred with the program *RAxML* [37]. This resulted in well-supported relationships between major subclades (>0.9 local posterior probability (PP)), with lower support for inter-specific relationships. *ASTRAL* inferred a high degree of discordance among gene tree topologies at many nodes (quartet score = 0.52), with multiple alternative bipartitions reconstructed at most nodes. This is evident from the presence of many more conflicting gene trees (numbers below branches) than congruent gene trees (numbers above branches) in Supplementary Fig. 1.

Figure 2: *ASTRAL* species tree inferred from *RAxML* gene trees using the multi-species coalescent model. Numbers at nodes of the tree signify local posterior probability (LPP) values, a measure of support for each quadripartition. Taxa within the red box belong to the 'Grandiceps' subclade, taxa within the green box belong to the 'Coccinea' subclade, taxa within the pink box represent the 'Stenantha' subclade and those within the blue box belong to the 'Chocóan' subclade. Images show inflorescences of species within *Brownea*- species shown are, from top to bottom: *B. grandiceps* (photograph © Rowan Schley), *B. jaramilloi* (photograph © Xavier Cornejo), *B. macrophylla* (photograph © Bente Klitgaard), *B. coccinea* subsp. *capitella* (photograph © Xavier Cornejo), *B. longipedicellata* (photograph © Domingos Cardoso) and *B. multijuga* (photograph © Bente Klitgaard).

Ancient introgression in Brownea species

Phylogenetic networks were estimated using concordance factors (CF) generated from *RAxML* gene

trees to ascertain whether the historical pattern of diversification in *Brownea* was tree-like or

reticulate. Networks with different numbers of hybridization events (h) were compared using -log

pseudolikelihoods. These comparisons pinpointed two hybridization events within *Brownea* (i.e., the

best-fit network model was $h = 2$, as indicated by the -log pseudolikelihood values in Supplementary

Fig. 2). Negative log pseudolikelihoods increased steadily between $h = 0$ and $h = 2$, after which the

increasing number of hybridization events only make minimal improvements to -log

pseudolikelihood.

Figure 3: Phylogenetic network with two hybridization events ($h = 2$), estimated using *SNaQ!* in the

Julia package *PhyloNetworks*. Light blue horizontal branches indicate inferred hybridization events,

and numbers next to the branches show the estimated proportion of genes contributed by each lineage

in the hybridization event. Red branches signify the ancestral lineage and what proportion of the

modern lineage's genes were contributed from it. Taxa were chosen in order to represent one

accession per lineage, as per the assumptions of *PhyloNetworks*, and the individual with the highest

gene recovery was used for each lineage. Coloured boxes containing taxa are described in the legend

of Fig. 2.

The inferred phylogenetic network (Fig. 3) showed a broadly similar topology to the species tree in

Fig. 2, with the addition of two hybridization events. The first hybridization event occurred between

the lineage leading to the Venezuelan accessions of *B. grandiceps/B. leucantha* and *B. birschelli*,

suggesting that the lineage leading to *B. birschelli* has in the past contributed around 34% of the

genetic material present in the common ancestor of the Venezuelan *B. grandiceps* and *B. leucantha*.

The second inferred occurrence of hybrid ancestry occurs between the ancestors of the subclade

containing the Colombian accessions of *B. coccinea/B. santanderensis* and the lineage leading to *B.*

enricii, which contributed around 37% of the genetic material belonging to the ancestor of the

aforementioned subclade.

TICR (Tree Incongruence Checking in R) analysis indicated that a network-like model best described

the patterns of incongruence in the single-accession-per-species gene trees inferred during this study.

This method suggested an excess of outlier quartets ($P= 1.240 \times 10^{-19}$, $X^2 = 91.149$), which differed

significantly from the CF values expected under a model containing only incomplete lineage sorting.

As such, a tree-like model was able to be rejected as an explanation for the observed relationships

between taxa in *Brownea*, suggesting that hybridization occurred over the course of its evolutionary

history.

Introgression dynamics across loci

Having examined the degree of historical reticulation among *Brownea* lineages, population genomic

data were generated with ddRAD sequencing and used to investigate recent introgression at a finer

taxonomic scale. More specifically, the degree of shared genetic variation was visualised for co-

occurring individuals of *B. grandiceps* and *B. jaramilloi*, which putatively hybridize in the wild.

Following this, in order to make inferences about the potential evolutionary significance of recent

introgression, the rates at which different loci introgress relative to the rest of the genome were

estimated for these taxa using genomic clines.

In total, ddRAD sequencing resulted in 640,898,950 reads between 350-550bp in length for 171

individuals. Among these reads, 28,503,796 (4.45%) were discarded due to poor recovery. There was

a mean of 3,747,947 reads per individual, with an average coverage depth across all samples of 27.5x.

Relatively high levels of shared genetic variation were observed among taxa, along with low levels of

genetic differentiation and only marginal differences in the amount of genetic variation, as shown by

the population genetic statistics calculated by the Stacks pipeline for the two *Brownea* species

(including *B.* "rosada", which is grouped with *B. grandiceps*) (Supplementary Table 3). The F_{st}

calculated between *B. grandiceps/B.* "rosada" and *B. jaramilloi* was 0.111, representing a low degree

of fixation, and so a high amount of shared variation.

Patterns of shared genetic variation were visualised using principal component analysis (PCA) and a

genetic distance network inferred with the program *SPLITSTREE* [38]. These analyses indicated a

distinct signature of admixture between *B. grandiceps* and *B. jaramilloi*. The PCA of SNP variation

inferred using the R package *ggplot2* (Fig. 4) indicated that the first principal component (PC1)

explained the largest proportion of the genetic variation among all the principal components (20.4%).

Individuals of *B. grandiceps* are tightly clustered along PC1, where the individuals of *B. jaramilloi*

show much more variability, with many individuals forming an intergradation between the two main

species clusters. Additionally, the *B.* "rosada" accessions appear to have clustered more closely to *B.*

grandiceps than to *B. jaramilloi* along PC1. PC2, which explained 10.7% of the variation in the SNP

data, shows a similar degree of variability in both *B. grandiceps* and *B. jaramilloi*, with two

accessions of *B.* "rosada" shown to cluster in between both species. This pattern is reflected by the

implicit network built using *SPLITSTREE* (Supplementary Fig. 3). *SPLITSTREE* recovered a

clustering of individuals into two main groups, largely representing *B. grandiceps* and *B. jaramilloi,*

with four putative hybrid individuals displaying an intermediate relationship between the two species

clusters.

Figure 4: Principal component analysis of genotype data for all SNPs inferred at all loci. Individuals

are coded by colour and point shape: red circles denote individuals identified as *B. grandiceps*, yellow

circles denote individuals identified as *B. jaramilloi,* and blue triangles denote putative hybrid

individuals (*B.* "rosada"). The amount of the genetic variation explained by each principle component

is shown next to axes, which are labelled PC1 and PC2.

A *fastSTRUCTURE* analysis [39] was performed to further examine the degree of shared genetic

variation, as well as providing information on the population genetic structure of the two species. This

analysis indicated that there was a large amount of shared ancestry between the genotyped individuals

(Fig. 5), with evidence of extensive backcrossing due to the widely differing proportions of ancestry

in different individuals. Marginal likelihood comparison indicated that the best value of K (i.e., the

number of populations) was two ($\Delta_{\text{marginal likelihood}} = 0.085$, Supplementary Fig. 4A), which was the

value that resulted in the largest increase in marginal likelihood. Since the 'best' value of K is an

estimate, f*astSTRUCTURE* plots generated with other K values are shown in Supplementary Fig. 4B.

Figure 5 shows that most individuals of *B. jaramilloi* have varying fractions of *B. grandiceps* ancestry

in their genome. In addition, the individuals identified as *B.* "rosada" appear to have inherited most of

their ancestry from *B. grandiceps*, with only a minimal contribution from *B. jaramilloi*. The same

pattern was recovered from a *fastSTRUCTURE* run incorporating 40 random individuals from each

species, performed to account for any bias which may have been incurred by differences in sample

size (Supplementary Fig. 4C).

Figure 5: *fastSTRUCTURE* plot, indicating ancestry proportions for two populations ($K = 2$). The

'populations' correspond to two different species: *B. grandiceps* (shown in red) and *B. jaramilloi*

(shown in yellow). Each individual accession is represented by a column, and the proportion of

ancestry from either species (mean Q) is proportional to the length of different coloured bars in each

column. Individuals identified as *B.* "rosada" are marked with a blue box and a blue triangle beneath

their column. All taxa to the left and right of these boxes were identified as *B. jaramilloi* and *B.*

grandiceps, respectively.

A *NEWHYBRIDS* analysis [40] (https://github.com/eriqande/newhybrids) was used to categorize 169 of

171 genotyped individuals into different hybrid classes (pure, F1, F2, and backcrosses). This was

performed on three subsets of 500 loci due to the computational limitations of the program.

NEWHYBRIDS revealed that most hybrid individuals were the result of continued hybridization, with

pure *B. grandiceps* making up 73.9% of the genotyped population, and pure *B. jaramilloi* making up

21.9%. There were no F1 (first generation) hybrids identified by the subset of loci analysed, and all

hybrid individuals were either F2 hybrids (0.592%) or had a broad distribution of probabilities across

hybrid classes (3.55%). In the latter, the probability of any one hybrid class did not exceed 90%,

which was the threshold used to categorize individuals as belonging to a certain class.

Bayesian estimation of genomic clines (*bgc*) was utilised to determine whether any loci showed

outlying rates of admixture relative to the genome-wide average among all genotyped individuals.

This method estimates two parameters for each locus: α (the 'direction' of introgression) and β (the

'rate' of introgression), which were used to infer the impact of selection on introgressing loci. *bgc*

recovered a signal of asymmetric introgression but did not detect any differential rates of

introgression between loci. Of the 19,130 loci under study, *bgc* recovered 251 loci (1.3%) with mostly

B. jaramilloi alleles among all individuals (i.e. with positive α estimates) and 1089 loci (5.69%) with

mostly *B. grandiceps* alleles among all individuals (i.e., with negative α estimates). However, no loci

displayed extreme rates of introgression relative to the average rate across the genome (i.e., there were

no statistically outlying β parameter estimates). The MCMC runs from which these results were

drawn showed adequate mixing (Supplementary Fig. 5A-C).

Discussion

Our study represents the first clear case of reticulated evolution among Amazonian trees documented

at multiple phylogenetic levels. This contrasts with previous work, which suggested that hybridization

is an extremely rare phenomenon in Amazonian trees. We demonstrate that within *Brownea*, a

characteristic Amazonian tree genus, reticulated evolution has occurred over the course of its

evolutionary history, with evidence of introgression deep in the history of the genus and more

recently, between the closely related *B. grandiceps* and *B. jaramilloi*.

Phylogenetic network analysis suggested that introgression has taken place between *Brownea* lineages

in the past, with two separate hybridization events inferred and displayed in Fig. 3. The concordance

factors obtained from gene trees were not adequately explained by a tree-like model (i.e., one

accounting only for incomplete sorting of ancestral variation), suggesting that a network-like model

(i.e., one including introgression) best describes the diversification patterns within *Brownea* ($P=$

1.240×10^{-19}, $X^2 = 91.149$). Hybridization mainly occurs between individuals which are

geographically overlapping, or have been so at some point in the past [41]. Our findings are thus further

corroborated by the fact that both historical introgression events were between taxa that co-occur

(e.g., *B. grandiceps, B. leucantha* and *B. birschelli* on the Northern coast of Venezuela, and *B.*

coccinea, B. santanderensis and *B. enricii* within the Colombian cordilleras). This is also in

agreement with previous work [30], which indicated that the 'stem' lineages within *Brownea* shared a

broad distribution across northern South America, which may have further facilitated earlier

hybridization. Our study also recovered a low degree of support for bifurcating inter-specific

relationships (Fig. 2) and a high degree of incongruence among gene trees (Supplementary Fig. 1).

These results can be partially explained by introgression, rather than being caused purely by the

stochasticity of lineage sorting in large populations, which is suggested to be a common phenomenon

in many rainforest trees with large population sizes[42]. It seems that amongst closely-related species of

Brownea it is likely that divergence has not progressed to the point of complete reproductive isolation,

as has been shown in temperate *Quercus* species[43]. Moreover, the low quartet scores and minimal

gene-tree concordance found at the species level in this study mirror those observed in *Lachemilla*, a

montane neotropical plant genus, in which gene tree discordance was shown to be explained by both

historical and recent hybridization [44].

Since the inferred events of ancient introgression both involved the common ancestors of two sets of

present-day species, they are likely to have occurred several million years in the past, before the

divergence of the descendent species. The inferred introgression events are likely to have occurred

since the Miocene period (~23Ma), as date estimates from previous work suggest that *Brownea*

originated around this time [30]. Accordingly, it is unlikely that the inferred hybridization events are due

to 'accidental' hybridization between co-occurring species, merely producing maladapted F1 hybrids

which will eventually be outcompeted, fortifying the boundaries between species (a process known as

'reinforcement' [15]). Rather, our results suggest a persistence of introgression over evolutionary time

within *Brownea*.

This investigation also uncovered a substantial signal of recent introgression between *B. grandiceps*

and *B. jaramilloi* in the Yasuní National Park 50ha plot located in the Ecuadorian Amazon, with

evidence of multi-generational 'backcrossing', further suggesting that hybridization does not always

result in reinforcement. All hybrid individuals exhibited some degree of backcrossing in the

NEWHYBRIDS analysis, which implies that hybrids persist over time. The low F_{st} estimate for the *B.*

grandiceps/B. "rosada" and *B. jaramilloi* populations (0.11), in addition to the principal component

analysis (Fig. 4), the *fastSTRUCTURE* analysis (Fig. 5) and the *SPLITSTREE* plot (Supplementary

Fig. 3) indicated a high degree of shared variation between these lineages. All these approaches show

that individuals cluster into two main groups, largely representing *B. grandiceps* and *B. jaramilloi*, as

well as a set of individuals forming an intergradation between the two main clusters. The individuals

which form a part of this intergradation were mostly identified as *B. jaramilloi*, although hybrid

individuals are also observed in the *B. grandiceps* cluster. The higher number of variant sites and the

higher nucleotide diversity (π) recovered for *B. jaramilloi* (Supplementary Table 3) could also reflect

the hybrid ancestry of many individuals identified as this species. A similar introgression-driven

increase in nucleotide diversity has been shown in closely-related species of *Mimulus* which undergo

asymmetric introgression [45].

The *bgc* analysis indicated that most introgression was asymmetric, with more loci containing mostly

B. grandiceps alleles (5.96% of all loci) than loci containing mostly *B. jaramilloi* alleles (1.3% of all

loci). Importantly, this *bgc* analysis also indicated that gene flow occurs largely at the same rate

across loci, since there were no outlying estimates of the β parameter. The excess of *B. grandiceps*

ancestry mirrors the asymmetry of introgression suggested by Fig. 5 and is likely driven by the

uneven population sizes of the two species. *Brownea jaramilloi* has only ever been observed in a

small part of the western Amazon [33] and is much less populous than *B. grandiceps*, which occurs

across northern South America. More specifically, the disproportionate donation of alleles from *B.*

grandiceps may be due to 'pollen swamping', whereby pollen transfer from another, more populous

species is more likely than pollen transfer from less numerous conspecifics [46]. Pollen swamping can

serve as a mechanism of invasion (e.g., in *Quercus* [47]), and the associated asymmetric introgression

may lead to the extinction of the rarer species, especially when hybrids are inviable or sterile [48].

However, due to the ongoing introgression evident between *B. grandiceps* and *B. jaramilloi* it appears

that hybrids are not always sterile, or at least that 'foreign' alleles are not always subject to strong

purifying selection. The best evidence of this is the absence of loci with extreme values of the β

parameter recovered by our *bgc* analysis. Extreme deviations in β are mainly expected in the presence

of gene flow when selection against hybrids is strong, resulting in underdominance and a paucity of

heterozygous sites [25].

Further to this, the observed asymmetry in introgression could be caused by selection favouring

hybrid genotypes. The transfer of genomic regions which confer a selective advantage from a donor to

a recipient species (i.e. adaptive introgression) can result in a disproportionate amount of one species'

genome being present in hybrid individuals [49]. This occurs when viable hybrids only tend to backcross

with one of the parental species, thereby introducing the other species' genetic material into the first

in an asymmetrical fashion. Selection has been suggested as a driver for asymmetric introgression in

multiple plant genera, including *Helianthus* [50] and *Iris* [51]. Accordingly, it is possible that selectively

advantageous alleles are passing between species (e.g., between *B. jaramilloi* and *B. grandiceps*)

through backcrossing events over many generations as observed by the *fastSTRUCTURE* and

NEWHYBRIDS analyses in this study [52]. This may facilitate adaptation to new niches due to the

widening of the pool of variation upon which selection can act [18]. However, it is difficult to ascertain

whether adaptive introgression has occurred without measuring the impact of introgression on

variation in the phenotype and its fitness effects [19].

Studies such as ours that document hybridization between tree species within Amazonia, and within

tropical rainforests in general, are rare. Indeed, it was previously suggested that inter-specific hybrids

between rainforest tree species are poor competitors, and that fertile hybrid populations are nearly

non-existent [8]. While there is some evidence of introgression in tropical trees[11], most available studies

substantiating this are based on trees from other tropical regions or habitats (e.g., *Shorea* in Asia [53, 54],

or *Rhizophora* in Indo-Pacific mangroves [55]). Many of these other instances appear to occur only in

degraded habitats, or involve infertile first-generation hybrids with minimal backcrossing, which

contrasts with the findings of our study. Within *Brownea* we found evidence of introgression across

taxonomic, spatial and temporal scales, with evidence of backcrossing and a lack of selection against

hybrid genotypes. It is possible that a lack of selection against hybrids allows the passage of variation

between *Brownea* species, which may persist over evolutionary time and could help explain why we

found evidence of introgression both at macroevolutionary and microevolutionary scales. While it is

difficult to determine whether hybridization between Amazonian tree lineages is a more common

occurrence than previously thought, or whether it is a tendency unique to certain rainforest genera

such as *Brownea*, it may be prudent to review how relationships among tropical tree lineages are

inferred. Accordingly, in order to understand the evolutionary relationships within such groups using

phylogenetic networks may provide additional insight into whether reticulate evolution has

contributed to diversification within Amazonian rainforest, which is among the most species-rich

environments on Earth [41, 56].

Methods

Phylogenetic analysis

Sequences from 226 nuclear genes were used to elucidate the evolutionary relationships between

Brownea species using a phylogenomic approach. DNA sequence data were generated from leaf

material collected from herbarium specimens and silica-gel dried accessions using targeted bait

capture. Targeted bait capture uses RNA probes to enable the sequencing of fragmented DNA, such as

that collected from historical natural history collections. Details of library preparation, hybridization

and sequencing are detailed in Supplementary Methods 1. Twenty-three of twenty-seven lineages

were sampled within the genus *Brownea*, including the three subspecies of *B. coccinea* and one

putative hybrid lineage (*Brownea* "rosada", a putative hybrid of *B. grandiceps* and *B. jaramilloi*). In

total, 59 accessions were sampled within *Brownea*, as well as an additional 13 outgroup taxa from the

genera *Macrolobium*, *Heterostemon, Paloue* and *Browneopsis*, which form part of the 'Brownea

clade' (Leguminosae, subfamily Detarioideae [57]). The list of accessions and their associated

information can be found in Supplementary Table 1.

DNA sequencing reads were quality-checked with the program *FASTQC* v0.11.3 [58], and were

subsequently trimmed using *Trimmomatic* v.0.3.6 [59]. This was done to remove adapter sequences and

to quality-filter reads, by removing reads that were too short or of poor quality, as assessed by their

phred-33 score. Following quality-filtering, loci were assembled using *SPAdes* v3.11.1 [16] by the

HybPiper pipeline v1.2 [60], and potentially paralogous loci were removed using the Python [61] script

'*paralog_investigator.py*', distributed with the HybPiper pipeline. All sequences were aligned by gene

region using *MAFFT* v7.215 [62]. In order to infer gene trees for phylogenetic network analysis, the 226

gene regions were further refined to include only 20 taxa, representing a single accession per lineage

in *Brownea*, using *Macrolobium colombianum* as the outgroup taxon. Where applicable, samples were

chosen by comparing the sequence recovery of conspecific accessions and choosing the individual

with the best gene recovery. This resulted in 220 single-accession-per-lineage gene alignments.

Further details of quality filtering, read assembly and alignment of DNA sequencing reads are given

in Supplementary Methods 1.

Gene trees were generated for each of the 226 gene alignments using *RAxML* v.8.0.26 [37] with 1,000

rapid bootstrap replicates and the GTRCAT model of nucleotide substitution, which is a parameter-

rich model and hence is most suitable for large datasets. *Macrolobium colombianum* was used to root

both the full and single-accession dataset analyses since it was the *Macrolobium* accession that was

present in the most gene alignments. The gene trees from both datasets were used to generate species

trees under the multi-species coalescent model in *ASTRAL* v.5.6.1 [36]. The heuristic version of *ASTRAL*

was used because gene trees were estimated for >30 taxa, and the default parameters for the program

were used for species tree inference. Monophyly was not enforced for individuals belonging to the

same species (the '-*a*' option in *ASTRAL*). Finally, discordance between gene trees was calculated and

visualised for the full dataset using the Java program *PhyParts* v0.0.1 [63]

(https://bitbucket.org/blackrim/phyparts). The pattern of incongruence between gene trees for each

node was then mapped onto the *ASTRAL* species tree using the Python script *PhyPartsPieCharts* v1.0

(https://github.com/mossmatters/MJPythonNotebooks).

Inferring ancient introgression

Phylogenetic networks were inferred for 220 gene trees from the single-accession-per-lineage dataset

to understand whether introgression occurred over the course of the evolutionary history of *Brownea*.

Networks were inferred with the program *SNaQ!,* implemented in the *Julia* v0.6.4[64] package

PhyloNetworks v0.11.0 [56]. Networks were estimated using gene trees from the single-accession-per-

lineage dataset as per the assumptions of the program, which requires that each tip of the phylogenetic

trees represent a single lineage. This dataset contained 20 taxa from the genera *Brownea, Browneopsis*

and *Macrolobium.* The network with the number of hybridization events (*h*) best describing the data

was chosen using negative log pseudolikelihood comparison. Finally, the topologies of the gene trees

were compared with those of the best-fit phylogenetic network and those expected under a coalescent

model (i.e., a 'tree-like' model which accounts for incomplete lineage sorting but not hybridization).

The fit of the observed gene tree topologies to either the 'network-like' or the 'tree-like' model was

assessed using the Tree Incongruence Checking in R (TICR) test [65], implemented in the R v3.4.4 [66]

package *PHYLOLM* [67]. Proportions of genes contributed between lineages by hybridization events

were taken from the 'Gamma value' output of *PhyloNetworks*. Further details of phylogenetic

network analysis are contained in Supplementary Methods 1.

 Population-level introgression

The architecture and rates of introgression across the genome were examined using two ecologically

divergent *Brownea* species which co-occur in the Ecuadorian Amazon. In order to do this we used

ddRADseq [68], a reduced-representation genotyping approach which uses restriction enzymes to

fragment genomic DNA to generate many thousands of sequence markers *de-novo* for SNP discovery.

One-hundred and seventy-one specimens in total were genotyped using dried leaf material. Sampling

consisted of 128 individuals of *Brownea grandiceps*, 40 individuals of *B. jaramilloi* and three

individuals of *B.* "rosada" (the putative hybrid of *B. grandiceps* and *B. jaramilloi*), representing the

relative abundance of each species in the forest plot from which they were sampled. Leaf material was

collected from *Brownea* trees in the Yasuní National Park 50ha forest plot in Napo, Ecuador, and

dried in a herbarium press. The sample list is shown in Supplementary Table 2. Sampling, library

preparation and sequencing protocols are explained further in Supplementary Methods 2.

DNA sequencing reads from the ddRADseq genotyping were processed *de novo* (i.e., without the use

of a reference genome) using the Stacks pipeline v2.1 [69]. Reads were quality filtered using Stacks by

removing reads which were of poor quality (i.e., had a phred score <10), following which 'stacks' of

reads were created, and SNPs were identified among all *de novo* 'loci' and across individuals. This

was done using a minimum coverage depth (the '*-m'* flag in Stacks) of three and a within-individual

mismatch distance (*-M*) of seven nucleotides. Individuals with sequencing coverage under 7.5x were

removed. Loci found in fewer than 40% of individuals and sites with a minor allele frequency

threshold of 5% were filtered out using the *'populations'* module of Stacks.

This resulted in a dataset containing 22,046 loci with 120,085 SNPs for 171 individuals. A dataset

containing only one SNP per locus was also extracted using the Stacks *populations* module for use in

analyses which assumed no linkage disequilibrium. This subsetting resulted in a dataset containing

19,130 loci with 19,130 SNPs for 171 individuals. Details of read assembly and filtering are shown in

Supplementary Methods 2.

In order to understand the patterns of introgression at the population level, the full ddRAD dataset was

used to visualise the degree of shared variation between *B. grandiceps* and *B. jaramilloi*. This was

performed using principal component analysis implemented in R v3.4.4, followed by plotting with

ggplot2 [70], and by producing a neighbour net plot in *SPLITSTREE* v4.14.6 [38]. Furthermore, the single-

SNP-per-locus dataset containing 19,130 RAD loci was used to infer population structure with the

program *fastSTRUCTURE* v1.0 [39]. The number of populations (*K*) was chosen using the value which

provided the largest improvement in marginal likelihood. Finally, *NEWHYBRIDS* v1.1 [40]

(https://github.com/eriqande/newhybrids) was used to categorise individuals into different hybrid

classes, using three runs of 500 randomly subsetting ddRAD loci. Runs were performed with 50,000

MCMC sweeps following 50,000 burn-in sweeps under the default parameters of the program.

The relative 'rate' and 'direction' of introgression for each locus between the two *Brownea* species

was estimated using Bayesian estimation of genomic clines (*bgc*) v1.03 [71]. For each locus, *bgc*

compares the probability of ancestry at the locus relative to an individual's genome wide ancestry,

thereby allowing it to estimate two parameters for each locus. These parameters are α, which roughly

equates to the 'direction' of introgression, and β, which may be summarised as the 'rate' of

introgression for a locus [25,71]. In order to estimate these parameters from the single-SNP dataset

consisting of 19,130 RAD loci, 50,000 MCMC generations with a 50% burn-in were used in *bgc*.

Admixture proportions (i.e., mean *Q* values) generated by *fastSTRUCTURE* were used to assign each

individual to three populations (*Brownea grandiceps*, *B. jaramilloi* and admixed). Convergence was

checked for the MCMC output from *bgc* in Tracer v1.6 [72] and with the R package *coda* [73] using

Geweke's diagnostic [74]. Loci with significant 'excess ancestry' were identified by ascertaining

whether the 99% posterior probability estimates of the α and β parameters included zero (i.e., by

identifying positive or negative non-zero estimates of the parameters). In addition, loci which were

extreme 'introgression outliers' were identified for both parameters by identifying loci whose median

estimates were not included in the 99% posterior probability credible intervals [71]. Further details of

visualising shared variation, hybrid category assignment and *bgc* analysis are shown in

Supplementary Methods 2.

Data Availability

The datasets that support the findings of this study are available from online repositories. All raw

reads generated with the targeted bait capture and ddRADseq methods are available on the NCBI

Sequence Read Archive with the accession numbers SAMN13439069- SAMN13439140 and

SAMN13441804- SAMN13441974, respectively, under the Bioproject number PRJNA592723. All

full phylogenomic sequence alignments, single-accession-per-species alignments and tree files, bgc

input files, Stacks output files and the Detarioideae bait kit sequence file are found on Dryad

(https://doi.org/10.5061/dryad.k3j9kd53w). The full phylogenomic sequence alignments underlie the

phylogeny in Fig. 2, while the single-accession-per-species alignments and tree files underlie Fig. 3.

The variant call format (VCF) and Structure files provided on Dryad underlie Fig. 4 and Fig. 5. Data

are under embargo until publication, and any further data required are available from the

corresponding author upon request. A reporting summary for this article is available as a

Supplementary Information file.

494   References

1. Mayr, E. in *Systematics and the Origin of Species* (Columbia University Press, New York, New
York, USA., 1942).

2. Barraclough, T. G. in *The Evolutionary Biology of Species* (Oxford University Press, Oxford,
2019).

3. Kearns, A. M. *et al*. Genomic evidence of speciation reversal in ravens. *Nat. Comm.* **9**, 906 (2018).

4. Vonlanthen, P. *et al*. Eutrophication causes speciation reversal in whitefish adaptive radiations.
Nature **482**, 357-362 (2012).

5. Mallet, J. Hybrid speciation. *Nature* **446**, 279-283 (2007).

6. Antonelli, A. & Sanmartín, I. Why are there so many plant species in the Neotropics? *Taxon* **60**,
403-414 (2011).

7. Ehrendorfer, F. Evolutionary patterns and strategies in seed plants. *Taxon* **19**, 185-195 (1970).

8. Ashton, P. S. Speciation among tropical forest trees: some deductions in the light of recent
evidence. *Biol. J. Linn. Soc.* **1**, 155-196 (1969).

9. Gentry, A. H. Neotropical floristic diversity: phytogeographical connections between Central and
South America, Pleistocene climatic fluctuations, or an accident of the Andean orogeny? *Ann. Mo.*
Bot. Gard. **69**, 557-593 (1982).

10. Koptur, S. Outcrossing and pollinator limitation of fruit set: breeding systems of neotropical Inga
trees (Fabaceae: Mimosoideae). *Evolution* **38**, 1130-1143 (1984).

11. Scotti-Saintagne, C. *et al*. Phylogeography of a species complex of lowland Neotropical rain
forest trees (Carapa, Meliaceae). *J. Biogeogr.* **40**, 676-692 (2013).

12. Cardoso, D. *et al*. Amazon plant diversity revealed by a taxonomically verified species list. *Proc.*
Natl. Acad. Sci. U. S. A. **114**, 10695-10700 (2017).

13. Dexter, K. G. *et al*. Dispersal assembly of rain forest tree communities across the Amazon basin.
Proc. Natl. Acad. Sci. U. S. A. **114**, 2645-2650 (2017).

14. Valencia, R., Balslev, H. & Miño, G. P. Y. High tree alpha-diversity in Amazonian Ecuador.
Biodiversity & Conservation **3**, 21-28 (1994).

15. Hopkins, R. Reinforcement in plants. *New Phytol.* **197**, 1095-1103 (2013).

16. Rieseberg, L. H. & Wendel, J. F. Introgression and its consequences in plants. *Hybrid Zones and*
the Evolutionary Process **70**, 70-109 (1993).

17. Taylor, S. A. & Larson, E. L. Insights from genomes into the evolutionary importance and
prevalence of hybridization in nature. *Nature Ecology & Evolution* **3**, 170-177 (2019).

18. Whitney, K. D., Randell, R. A. & Rieseberg, L. H. Adaptive introgression of abiotic tolerance
traits in the sunflower *Helianthus annuus*. *New Phytol.* **187**, 230-239 (2010).

19. Suarez-Gonzalez, A., Lexer, C. & Cronk, Q. C. B. Adaptive introgression: a plant perspective.
Biol. Lett. **14**, 10.1098/rsbl.2017.0688 (2018).

20. Marques, D. A., Meier, J. I. & Seehausen, O. A combinatorial view on speciation and adaptive
radiation. *Trends in Ecology & Evolution* **34**, 531-544 (2019).

21. Lamichhaney, S. *et al*. Rapid hybrid speciation in Darwin's finches. *Science* **359**, 224-228 (2018).

22. Rieseberg, L. H. *et al*. Major ecological transitions in wild sunflowers facilitated by hybridization.
Science **301**, 1211-1216 (2003).

23. Meier, J. I. *et al*. Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nature*
Communications **8**, 14363 (2017).

24. Payseur, B. A. Using differential introgression in hybrid zones to identify genomic regions
involved in speciation. *Molecular Ecology Resources* **10**, 806-820 (2010).

25. Gompert, Z., Parchman, T. L. & Buerkle, C. A. Genomics of isolation in hybrids. *Philos. Trans.*
R. Soc. Lond. B. Biol. Sci. **367**, 439-450 (2012).

26. Hamilton, J. A. & Aitken, S. N. Genetic and morphological structure of a spruce hybrid (*Picea*
sitchensis × *P. glauca*) zone along a climatic gradient. *Am. J. Bot.* **100**, 1651-1662 (2013).

27. Sullivan, A. R., Owusu, S. A., Weber, J. A., Hipp, A. L. & Gailing, O. Hybridization and
divergence in multi-species oak (*Quercus*) communities. *Bot. J. Linn. Soc.* **181**, 99-114 (2016).

28. Abbott, R. J. *et al*. Hybridization and speciation. *J. Evol. Biol.* **26**, 229-246 (2013).

29. Seehausen, O. *et al*. Genomics and the origin of species. *Nature Reviews Genetics* **15**, 176-192
(2014).

30. Schley, R. J. *et al*. Is Amazonia a 'museum'for Neotropical trees? The evolution of the Brownea
clade (Detarioideae, Leguminosae). *Mol. Phylogenet. Evol.* **126**, 279-292 (2018).

31. Crawford, T. & Nelson, E. Irish Horticulturists. I: WH Crawford. *Garden History* **7**, 23-26 (1979).

32. Backer, C. A. in *Schoolflora voor Java* 418 (Visser & Co., Weltevreden, Java, Dutch East Indies,
1911).

33. Pérez, Á J., Klitgård, B. B., Saslis-Lagoudakis, C. & Valencia, R. *Brownea jaramilloi*
(Leguminosae: Caesalpinioideae), a new, over-looked species endemic to the Ecuadorian Amazon .
Kew Bull. **68**, 157-162 (2013).

34. Klitgaard, B. B. Ecuadorian *Brownea* and *Browneopsis* (Leguminosae-Caesalpinioideae):
taxonomy, palynology, and morphology. *Nord. J. Bot.* **11**, 433-449 (1991).

35. ter Steege, H. *et al*. Hyperdominance in the Amazonian tree flora. *Science* **342**, 1243092 (2013).

36. Zhang, C., Rabiee, M., Sayyari, E. & Mirarab, S. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**, 153 (2018).

37. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).

38. Huson, D. H. & Bryant, D. Application of phylogenetic networks in evolutionary studies. *Mol.*
Biol. Evol. **23**, 254-267 (2005).

39. Raj, A., Stephens, M. & Pritchard, J. K. fastSTRUCTURE: variational inference of population
structure in large SNP data sets. *Genetics* **197**, 573-589 (2014).

40. Anderson, E. C. & Thompson, E. A. A model-based method for identifying species hybrids using
multilocus genetic data. *Genetics* **160**, 1217-1229 (2002).

41. Burbrink, F. T. & Gehara, M. The biogeography of deep time phylogenetic reticulation. *Syst. Biol.*
67, 743-744 (2018).

42. Pennington, R. T. & Lavin, M. The contrasting nature of woody plant species in different
neotropical forest biomes reflects differences in ecological stability. *New Phytol.* **210**, 25-37 (2016).

43. Lepais, O. & Gerber, S. Reproductive patterns shape introgression dynamics and species
succession within the European white oak species complex. *Evolution* **65**, 156-170 (2011).

44. Morales-Briones, D. F., Liston, A. & Tank, D. C. Phylogenomic analyses reveal a deep history of
hybridization and polyploidy in the Neotropical genus Lachemilla (Rosaceae). *New Phytol.* **218**,
1668-1684 (2018).

45. Sweigart, A. L. & Willis, J. H. Patterns of nucleotide diversity in two species of *Mimulus* are
affected by mating system and asymmetric introgression. *Evolution* **57**, 2490-2506 (2003).

46. Buggs, R. J. & Pannell, J. R. Rapid displacement of a monoecious plant lineage is due to pollen
swamping by a dioecious relative. *Current Biology* **16**, 996-1000 (2006).

47. Petit, R. J., Bodénès, C., Ducousso, A., Roussel, G. & Kremer, A. Hybridization as a mechanism
of invasion in oaks. *New Phytol.* **161**, 151-164 (2004).

48. Balao, F., Casimiro-Soriguer, R., García-Castaño, J. L., Terrab, A. & Talavera, S. Big thistle eats
the little thistle: does unidirectional introgressive hybridization endanger the conservation of
Onopordum hinojense? *New Phytol.* **206**, 448-458 (2015).

49. Yang, W. *et al*. Genomic evidence for asymmetric introgression by sexual selection in the
common wall lizard. *Mol. Ecol.* **27**, 4213-4224 (2018).

50. Scascitelli, M. *et al*. Genome scan of hybridizing sunflowers from Texas (*Helianthus annuus* and
H. debilis) reveals asymmetric patterns of introgression and small islands of genomic differentiation.
Mol. Ecol. **19**, 521-541 (2010).

51. Arnold, M. L., Tang, S., Knapp, S. J. & Martin, N. H. Asymmetric introgressive hybridization
among Louisiana Iris species. *Genes* **1**, 9-22 (2010).

52. Mallet, J. Hybridization as an invasion of the genome. *Trends in Ecology & Evolution* **20**, 229-237
(2005).

53. Kenzo, T. *et al*. Overlapping flowering periods among *Shorea* species and high growth
performance of hybrid seedlings promote hybridization and introgression in a tropical rainforest of
Singapore. *For. Ecol. Manage.* **435**, 38-44 (2019).

54. Kamiya, K. *et al*. Morphological and molecular evidence of natural hybridization in *Shorea*
(Dipterocarpaceae). *Tree Genetics & Genomes* **7**, 297-306 (2011).

55. Lo, E. Testing hybridization hypotheses and evaluating the evolutionary potential of hybrids in
mangrove plant species. *J. Evol. Biol.* **23**, 2249-2261 (2010).

56. Solís-Lemus, C., Bastide, P. & Ané, C. PhyloNetworks: a package for phylogenetic networks.
Mol. Biol. Evol. **34**, 3292-3298 (2017).

57. Legume Phylogeny Working Group. A new subfamily classification of the Leguminosae based on
a taxonomically comprehensive phylogeny: The Legume Phylogeny Working Group (LPWG). *Taxon*
66, 44-77 (2017).

58. Andrews, S. FastQC: a quality control tool for high throughput sequence data. Available online
at http://www.bioinformatics.babraham.ac.uk/projects/fastqc. (2010).

59. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* **30**, 2114-2120 (2014).

60. Johnson, M. G. *et al*. HybPiper: Extracting coding sequence and introns for phylogenetics from
high-throughput sequencing reads using target enrichment. *App. Plant. Sci.* **4**, 1600016 (2016).

61. Python Software Foundation. Python Language Reference, version 2.7. *http://www.python.org*
(2010).

62. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772-780 (2013).

63. Smith, S. A., Moore, M. J., Brown, J. W. & Yang, Y. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants. *BMC*
Evolutionary Biology **15**, Published online: DOI 10.1186/s12862-015-0423-0 (2015).

64. Bezanson, J., Edelman, A., Karpinski, S. & Shah, V. B. Julia: A fresh approach to numerical
computing. *SIAM Rev* **59**, 65-98 (2017).

65. Stenz, N. W., Larget, B., Baum, D. A. & Ané, C. Exploring tree-like and non-tree-like patterns
using genome sequences: an example using the inbreeding plant species *Arabidopsis thaliana* (L.)
Heynh. *Syst. Biol.* **64**, 809-823 (2015).

66. R Development Core Team. R: A language and environment for statistical computing. Available
at: http://www.R-project.org/. (2013).

67. Ho, L. S. T. & Ané, C. A linear-time algorithm for Gaussian and non-Gaussian trait evolution
models. *Syst. Biol.* **63**, 397-408 (2014).

68. Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S. & Hoekstra, H. E. Double digest RADseq:
an inexpensive method for de novo SNP discovery and genotyping in model and non-model species.
PloS one **7**, e37135 (2012).

69. Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W. & Postlethwait, J. H. Stacks: building and
genotyping Loci de novo from short-read sequences. *G3 (Bethesda)* **1**, 171-182 (2011).

70. Wickham, H. in *ggplot2: elegant graphics for data analysis* (Springer, 2016).

71. Gompert, Z. & Buerkle, C. A. Bayesian estimation of genomic clines. *Mol. Ecol.* **20**, 2111-2127
(2011).

72. Rambaut, A., Suchard, M. A., Xie, D. & Drummond, A. J. Tracer.
http://tree.bio.ed.ac.uk/software/tracer/ **1.6** (2015).

73. Plummer, M., Best, N., Cowles, K. & Vines, K. CODA: convergence diagnosis and output
analysis for MCMC. *R news* **6**, 7-11 (2006).

74. Geweke, J. in *Evaluating the accuracy of sampling-based approaches to the calculation of*
posterior moments (Federal Reserve Bank of Minneapolis, Research Department, Minneapolis, MN,
USA, 1991).

Acknowledgements

Laboratory work, fieldwork and herbarium visits to NY and US were funded by the NERC SSCP

DTP, and a Genetics society Heredity Fieldwork grant funded part of the fieldwork in Ecuador.

Phylogenomic sequencing was also part-funded by a grant from the Royal Botanic Gardens, Kew.

Thank you to Dario Ojeda for baits and associated help, to Laszlo Csiba, Penny Malakasi, Niroshini

Epitawalage, Dion Devey, Robyn Cowan and Steven Dodsworth for help in the laboratory, and to

Gregor Gilfillan at the Norwegian sequencing centre for help with ddRAD sequencing. Thank you to

Andres Melo Burbano for his help during fieldwork, as well as to Renato Valencia and Álvaro Pérez

for helping to coordinate fieldwork in Ecuador. Thanks to Colin Hughes, Jeff Doyle and Matteo

Fumagalli for comments on analytical methods, and to Peter Raven and Benjamin Torke for their

comments on the rarity of hybridization in tropical trees.

Collection, transport and extraction of genetic material from plant accessions collected in Yasuní

National Park were authorised by Ecuador's Environment Ministry. Permits were issued with the

following authorization numbers:

• **Collection permit:** 021-2016-IC-FAU-FLO-DPAO-PNY;

• **Mobilization permit**: 037-2016-MOV-FLO-MAE-DPAO;

• **Export permit**: 208-2019-EXP-CM-FAU-DNB/MA; and

• **Genetic research permit (Contrato Marco)**: MAE-DNB-CM-2018-0082

Author Contributions

R.J.S. conceived this study and performed all analyses, supervised by T.B., F.F. and B.K. Population

genomic data were generated by R.J.S., and phylogenomic data were generated by R.J.S., I.K. and I.L.

Baits for hybrid capture were provided by M.d.l.E. R.J.S. wrote the first draft of the manuscript with

contributions from R.T.P., O.A.P.E., A.J.H., M.d.l.E., I.L., T.B., F.F. and B.K.

<u>Competing Interests</u>

The authors declare no competing financial interest.