Cavefish cope with environmental hypoxia by developing more erythrocytes and

overexpression of hypoxia inducible genes

Running Title: Cavefish response to hypoxia

Corine M. van der Weele and William R. Jeffery

Department of Biology, University of Maryland, College Park, MD 20742 USA

Correspondence: Jeffery@umd.edu

Keywords

Hypoxia, *Astyanax mexicanus*, Cavefish, Hematopoiesis, Erythrocytes, Hypoxia-inducible genes,

Cave colonization.

Summary

Astyanax mexicanus cavefish cope with hypoxic environments by expanding embryonic

hematopoietic domains, increasing the capacity for erythrocyte development, and constitutive

overexpression of hypoxia-inducible genes.

Abstract

Dark caves lacking primary productivity can expose subterranean animals to hypoxia. We used the surface-dwelling (surface fish) and cave-dwelling (cavefish) morphs of *Astyanax mexicanus* as a model for understanding the mechanisms of hypoxia tolerance in the cave environment. Primitive hematopoiesis, which is restricted to the posterior lateral mesoderm in other teleosts, also occurs in the anterior lateral mesoderm in *Astyanax*, potentially pre-adapting surface fish for hypoxic cave colonization. Cavefish have enlarged both hematopoietic domains and develop more erythrocytes than surface fish, which are required for normal development in both morphs. Laboratory induced hypoxia suppresses growth in surface fish but not in cavefish. Both morphs respond to hypoxia by overexpressing Hypoxia-Inducible Factor (HIF) pathway genes, but some *hif* genes are constitutively upregulated in normoxic cavefish to similar levels as hypoxic surface fish. We conclude that cavefish cope with hypoxia by increasing erythrocyte development and constitutive HIF gene overexpression.

Introduction

A unifying theme of cave habitats is complete darkness (Culver and Pipan, 2009) and the absence of primary productivity can subject cave-dwelling animals to oxygen deficiency (Malard and Hervant, 1999). We have used the teleost *Astyanax mexicanus*, which consists of surface-dwelling (surface fish) and cave-dwelling (cavefish) morphs, as a model to study adaptation to hypoxia in caves. Surface fish colonized subterranean waters in the Sierra de El Abra region of Mexico (Mitchell et al., 1977; Gross, 2012) during the late Pleistocene (Fumey et al., 2018;

Herman et al., 2018), and their cavefish descendants evolved numerous traits for adaptation to

perpetual darkness. Cavefish adaptations include more taste buds and cranial neuromasts,

increased olfactory capacity, and fat deposits, which could facilitate feeding and survival during

times of low food input, and decreased metabolic rates and loss of eyes, which may be important

in energy conservation (Yamamoto et al., 2004; Moran et al., 2014; 2015, Varatharasan et al.,

2009; Blin et al., 2018; Yoshizawa et al., 2014; Hüppop, 1986; Xiong et al., 2018). Hypoxia

may be a deterrent to the survival of colonizing teleosts in the Sierra de El Abra, where oxygen

reductions of 50% or more have been measured in cave waters (Boggs and Gross, 2021; Rohner

et al., 2013, Ornelas-Garcia et al., 2018). *Astyanax* cavefish thrive under these conditions, but

little is known about how they have adapted to hypo-oxygenated waters.

57   Oxygen is bound to hemoglobin and transported through the body by red blood cells.

Increased erythrocyte production is a well-documented strategy for coping with hypoxia at high

altitude (Haase, 2013), but the role of red blood cells in adaptation to hypoxic caves is unknown.

Hematopoiesis, the process in which erythrocytes and other blood cells develop during

embryogenesis, occurs in two waves in vertebrates: the primitive and definitive waves (Davidson

and Zon, 2004; Paik and Zon, 2010). In the zebrafish *Danio rerio* and other teleost embryos, the

first or primitive wave occurs in the anterior and posterior lateral mesoderm beginning during the

tailbud stage. The anterior lateral mesoderm (ALM) undergoes myelopoiesis to form

macrophages, and some of these later differentiate into neutrophils and microglia (Herbomel et

al., 1999; Herbomel et al., 2001; Le Guyader et al., 2008), whereas the posterior lateral

mesoderm (PLM) undergoes erythropoiesis to form primitive erythrocytes. Subsequently, the

PLM converges into the intermediate cell mass, where the precursors of endothelial cells lining

the first blood vessels, primitive erythrocytes, and myeloid cells will ultimately differentiate.

The second or definitive wave of hematopoiesis begins later in development when hematopoietic

stem and progenitor cells are formed in the aorta-gonad-mesonephros (AGM), which seed the

caudal hematopoietic tissue and form the posterior blood islands, where definitive erythrocytes

begin to differentiate (Paik and Zon 2010; Gore et al., 2018). Eventually, definitive

hematopoiesis moves to the thymus and head kidneys in adults. The primitive and definitive

waves of hematopoiesis are potential targets for adaptive changes induced by hypoxia.

Hematopoiesis is directed by a series of transcription factors producing different types of

blood cells (Davidson and Zon, 2004; Carroll and North, 2014). The *growth factor*

independence 1 (*gfi1aa*) gene, which encodes a transcriptional repressor of many different genes

(Möröy et al., 2015), is important for primitive hematopoiesis at several developmental stages

(Wei et al., 2008; Cooney et al., 2013; Moore et al., 2018). In zebrafish, *gfi1aa* expression occurs

throughout the period of embryonic segmentation, first in the PLM, then in the intermediate cell

mass, and finally in the AGM (Wei et al., 2008; Cooney et al., 2013), where it directs the

development of hemogenic endothelia into hematopoietic stem and progenitor cells, which

ultimately differentiate into red blood cells (Moore et al., 2018). The *gfi1aa* gene is also

expressed in the ALM, albeit weakly and only for a short time (Wei et al., 2008). Its expression

at different stages of hematopoiesis makes *gfi1aa* an excellent marker for investigating

differences between surface fish and cavefish hematopoiesis. The *LIM domain only 2* gene

encodes another transcription factor essential for determination of hematopoietic domains early

in primitive hematopoiesis (Patterson et al., 2007) and is also an excellent marker for differences

during the initial stages of hematopoiesis.

Oxygen is required in aerobic organisms to produce energy, and insufficient oxygen leads

to the activation of an evolutionarily-conserved transcriptional response (Majmundar et al.,

2010). The responses to oxygen deprivation are coordinated by activation of the hypoxia-

inducible factors (HIF1, HIF2 and HIF3) (Semenza and Wang, 1992), a family of transcription

factors controlling a large number of downstream target genes involved in promoting different

responses to hypoxia (Pelster and Egg, 2018; Rashid et al., 2017). Increased expression of the α

subunit of HIF1 and the HIF regulated genes are indicators of hypoxia (Mandic et al., 2021).

Notable responses include the improvement of oxygen supply through increasing red blood cell

production and function (Haase, 2013; Kulkarni et al., 2010; Cai et al., 2020), metabolic

homeostasis balancing ATP production with oxygen consumption, management of Reactive

Oxygen Species (ROS) via glycolysis, modulating anaerobic respiration and the pentose

phosphate pathway (PPP) (Singh et al., 2017; Wheaton and Chandel, 2011; Stincone et al.,

2015), and conservation of energy by growth inhibition (Kamei, 2020). *Astyanax* cavefish

(Hüppop, 1986; Moran et al., 2014, Aspiras et al., 2015) and other cave-adapted animals (Bishop

et al., 2004) have evolved reduced metabolic rates to conserve energy in the cave environment.

However, the strategies used to conserve energy in hypoxic cave environments are poorly

understood.

The present investigation shows that cavefish embryos develop significantly more

erythrocytes than surface fish. The results also revealed that cavefish larvae have constitutively

high expression of some of the HIF1 pathway genes and show no effects of oxygen depletion on

growth, traits that could promote larval survival in hypoxic environments. These studies reveal

that changes in development of the circulatory system may underlie adaptation of cavefish to

hypo-oxygenated environments.

Results

Cavefish have more erythrocytes than surface fish

To test the hypothesis that cavefish develop more red blood cells as a response to

environmental hypoxia, we compared erythrocytes between surface fish and cavefish embryos in

several different ways. First, we conducted direct observations of blood cell numbers after

circulation begins in the Duct of Cuvier (DC) at 36 hpf (Fig. 1A-D). At this time, the circulatory

system is still partially open, and blood cells can be imaged or stained and quantified as they

flow across the yolk (Videos 1, 2 and Fig. 1B-D, F). The number of circulating blood cells was

quantified in sequential video frames using Image-J analysis software, and the results showed

that cavefish have more blood cells than surface fish (Fig. 1B-D, F). Second, the number of

erythrocytes was quantified in the DC by staining with o-dianisidine (Iuchi and Yamamoto,

1983), and the results revealed that cavefish have more erythrocytes than surface fish (Fig. 1E,

G). Third, expression of the *hbb2* and *hbbe2 ß-hemoglobin* genes was followed during surface

fish and cavefish development by qPCR and *in situ* hybridization (Fig. 2A, B, D-F). The qPCR

results revealed significantly higher levels of *hbb2* mRNA at 24 hpf and 60 hpf and *hbbe2*

mRNA at 60 hpf (Fig. 2 A, B). *In situ* hybridization also showed more extensive *hbb2* staining

in cavefish compared to surface fish (Fig. 2D-F), which was particularly evident in the yolk mass

at 36 hpf (Fig. 2E). Fourth, the expression of *gfi1aa*, which encodes a transcription factor

essential for hematopoiesis (Cooney et al., 2013), was compared by qPCR and *in situ*

hybridization (Fig. 2C, G). The results indicated that *gfi1aa* mRNA levels were significantly

increased in cavefish compared to surface fish at 24 hpf (Fig. 2C, G), prior to detection of the

highest levels of hemoglobin transcription (Fig. 2A, B). These results provide multiple lines of

evidence indicating that red blood cells are increased during cavefish development.

Hematopoietic domains in surface fish and cavefish embryos

To determine the developmental basis of erythrocyte enhancement, we compared the

timing and spatial expression of *hbb2*, *gfi1aa,* and other hematopoietic marker genes during

primitive hematopoiesis in surface fish and cavefish embryos (Fig. 3). As in zebrafish (Brownlie

et al., 2003; Wei et al., 2008), *hbb2* was expressed in the intermediate cell mass of the PLM

during primitive hematopoiesis in *Astyanax* surface fish and cavefish embryos (Fig. 3A). In

striking contrast to zebrafish, in which erythropoiesis occurs exclusively in the PLM, *hbb2*

expression was detected in both the ALM and PLM of *Astyanax* surface fish and cavefish

embryos (Fig. 3A). The expression of *gfi1aa* was also present in the ALM and PLM during

Astyanax embryogenesis (Fig. 3A). Furthermore, *gfi1aa* and *hbb2* staining were stronger in the

cavefish ALM and PLM compared to surface fish (Fig. 3A). To further evaluate the primitive

erythropoietic domains, we examined expression of the *lim domain only 2* (*lmo2*) gene, a marker

of developing blood cells (Patterson et al., 2007), by *in situ* hybridization. The results confirmed

blood cell development in both the ALM and PLM in *Astyanax* embryos and revealed that both

primitive erythropoietic domains are expanded in cavefish embryos (Fig. 3C).

In zebrafish, the ALM undergoes myelopoiesis, rather than erythropoiesis, producing

macrophages that disperse throughout the embryo (Herbomel et al., 1999). Therefore, we asked

if macrophage development in *Astyanax* is compromised by the expansion of primitive

erythropoiesis into the ALM. To address this possibility, we examined expression of the *l-*

plastin1 (*lcp1*) gene, a marker for macrophage differentiation (Herbomel et al., 2001). *In situ*

hybridization showed that *lcp1* expressing cells were distributed throughout the ventral body and

yolk masses of surface fish and cavefish embryos, implying that macrophages migrated from the

ALM and differentiated (Fig. 2B). Therefore, macrophage development does not appear to be

compromised by the expansion of erythropoiesis into the ALM.

In summary, the results reveal the presence of a novel domain of primitive erythropoiesis

in the anterior region of *Astyanax* embryos and suggest that red blood cells are increased in

cavefish due to expansion of both the anterior and posterior primitive erythropoiesis domains.

Blood cell enhancement is a maternally controlled trait

Hematopoietic domains develop in the lateral mesoderm of zebrafish embryos during

gastrulation (Kimmel et al., 1990), and changes in gastrulation and axis determination are under

maternal control in cavefish (Ma et al., 2018; Torres-Paz et al., 2019). Therefore, to determine

whether the increase in cavefish blood cells is controlled by maternal or zygotic changes, we

conducted reciprocal hybrid crosses (cavefish female x surface male and surface fish female x

cavefish male) and quantified blood cells in the DC of F1 hybrid embryos by video imaging (Fig.

4). As controls, crosses were also done and blood cell numbers quantified in the offspring of

parental cavefish x cavefish and surface fish x surface fish crosses. The F1 hybrid progeny of

the cavefish female x surface fish male and the cavefish x cavefish crosses (Fig. 4C, D)

developed more blood cells than the progeny of surface fish female x cavefish male and surface

fish x surface fish crosses (Fig. 4A, B), showing that the extent of erythrocyte development is

dependent on the source of eggs. These results reveal that the increase in cavefish blood cells is

a maternally controlled trait.

Erythrocytes are required for Astyanax development

Erythrocytes also develop early in zebrafish embryogenesis, but are not required for

oxygen delivery until much later in development (Pelster and Burggren, 1996). Accordingly, we

performed experiments to determine whether red blood cells have a role in early *Astyanax*

development. To address this question, we compared the effects of hemolytic anemia in surface

fish and cavefish embryos (Fig. 5). Hemolysis of red blood cells was induced by treatment with

phenylhydrazine (PHZ) (Houston et al., 1988) beginning at 14 hpf (13-17 somite stage), and

hbb2 staining, red blood cell number, and post-anal tail length (used as a proxy for normal

growth and development) were compared in surface fish and cavefish 14 hrs. later (Fig. 5A-C).

We found that *hbb2* staining and red blood cells were reduced at increasing PHZ concentrations

in both surface fish and cavefish embryos (Fig. 5A, B). However, at the same PHZ concentration

both erythrocyte phenotypes were more strongly affected in surface fish than in cavefish,

consistent with more red blood cells in the latter. The results also showed that growth decreased

in concert with hemoglobin transcript and erythrocyte reduction in surface fish, and concordant

with more red blood cells, no effects on growth were seen in PHZ-treated cavefish (Fig. 5B).

The PHZ-treated embryos developed axial defects, most notably edemas, twisted notochords,

and shortened and ventrally bent tails (Fig. 5C), which were more prevalent at lower PHZ

concentrations in surface fish than cavefish embryos (Table 1). To address the possibility that

reduced tail length was a consequence of abnormal notochord morphogenesis rather than growth

suppression, tail length and notochord morphology were investigated at 36 hpf in surface fish

exposed to a PHZ concentration that virtually abolished red blood cells (Fig. 5D). Under these

conditions, post-anal tail growth was significantly reduced, but notochord defects were not

detected, suggesting that hemolytic effects on axial growth are related to the absence of red

blood cells. The results show that erythrocytes are required during early development in

Astyanax and more red blood cells may reduce the sensitivity of cavefish to hemolytic anemia.

Cavefish but not surface fish maintain growth in a hypoxic laboratory environment

Subterranean waters harboring *Astyanax* cavefish are low in oxygen (Rohner et al., 2013;

Ornelas-Garcia et al., 2018; Boggs and Gross, 2021), which is essential for normal development

and growth (Giaccia et al., 2004). Therefore, we hypothesized that cavefish may be more

tolerant to the effects of hypoxia on growth than surface fish. To test this hypothesis, 36 hpf

surface fish and cavefish were exposed to a hypoxic laboratory environment for 18 hours (Fig.

6A, B). The effects on growth were determined by measuring post-anal tail length before and

after hypoxia treatment, and the number of erythrocytes was quantified by o-dianisidine staining

at the end of the treatment period. The results showed that hypoxia reduced the number of red

blood cells in the DC and suppressed growth in surface fish but not cavefish (Fig. 6B), showing

that cavefish are more resistant to oxygen depletion. Reasoning that eye growth would be

especially sensitive to hypoxia because of the energy cost associated with retinal differentiation

and maintenance (Wong-Riley, 2010; Moran et al., 2015), eye size was also examined in surface

fish and cavefish exposed to normoxia or hypoxia for 18 hrs. The results showed that eye size

was more sensitive to hypoxia relative to normoxia in surface fish compared to cavefish (Fig.

6C). The results suggest that cavefish growth is resistant to hypoxia and development of more

erythrocytes may contribute to hypoxia tolerance.

Differential expression of oxygen-sensitive genes during hypoxia

To determine the molecular responses of surface fish and cavefish to hypoxia, we

quantified some of the key oxygen sensitive genes by qPCR (Fig. 7). Exposure of surface fish or

cavefish larvae to hypoxia for 2 hrs. resulted in increased transcript levels of the *insulin growth*

factor binding protein 1a (*igfbp1a*) gene, the HIF1 family genes *hif1aa* and *hif1a-like2*, and the

HIF1-regulated downstream gene *hexokinase 1* (*hk1*) (Fig. 7). In contrast, the *hifab* and *hif1a-*

like genes were expressed at similar levels in normoxic cavefish and hypoxic surface fish (Fig.

7), implying constitutive expression in cavefish. The expression of the *glucose-6-phosphate*

dehydrogenase (*g6pd*), *lactate dehydrogenase a* (*ldha*), and *pyruvate dehydrogenase kinase 1*

(*pdk1*) genes were not significantly increased by hypoxia in surface fish or cavefish (Fig. 7). The

results indicate that *igfbp1*, *hif1a-like2* and *hk1* are upregulated by hypoxia in both surface fish

and cavefish, whereas *hif1aa* and *hif1a-like* are expressed at higher levels in cavefish than in

surface fish in either normoxic or hypoxic laboratory conditions. Thus, cavefish exhibit a

complex molecular response to hypoxia: some oxygen sensing genes are upregulated, others are

not changed, and some of the latter genes may be insensitive to oxygen depletion because they

are already expressed at maximal levels in normoxic cavefish.

Discussion

We show that cavefish embryos develop more red blood cells than surface fish embryos,

a classic response to hypoxia (Martinez, 2004; Timmerman and Chapman, 2004; Rutjes et al.,

2007; Haase, 2013), when they are raised under normoxic laboratory conditions. This conclusion

is based on direct counts of blood cells, including erythrocytes, increased expression of multiple

erythropoietic marker genes, and hyper-expression of the *gfi1aa* gene, a crucial regulator of

hematopoiesis. In addition, we report that the increase in erythrocytes may be caused by

expansion of embryonic hematopoietic domains in cavefish compared to surface fish. Finally,

we provide evidence that cavefish have evolved a permanent response to hypoxia involving the

constitutive overexpression of the some of the *hif1* genes, which persists under normoxic

laboratory conditions.

Hematopoiesis occurs in the primitive and definitive waves in vertebrates (Davidson and

Zon, 2004; Paik and Zon, 2010). Increased *hbb2, hbbe2,* and *gfi1aa* expression was observed

during embryogenesis and after larval hatching in cavefish. In zebrafish, transfusion experiments

indicate that blood cells circulating during the first four days of development are primitive

erythrocytes (Weinstein et al., 1996). Accordingly, our findings suggest that *Astyanax* larval

blood cells are primitive erythrocytes derived from the first wave of hematopoiesis.

Nevertheless, we also found stronger expression of erythropoietic marker genes in the AGM and

caudal hematopoietic tissue, which are involved in definitive hematopoiesis, implying that the

second wave may also be expanded in cavefish. Further investigation will be required to

determine whether erythrocytes are also increased in adults, although this possibility seems

likely because adult cavefish have increased levels of hemoglobin transcripts (Sears et al., 2020)

and more erythrocytes in their head kidneys compared to surface fish (Peuß et al., 2020). We

conclude that cavefish have evolved a permanent enhancement in red blood cells as a potential

adaptation for survival in hypoxic cave waters.

The results showed that surface fish and cavefish primitive erythrocytes are derived from

two different hematopoietic domains, one located anteriorly in the ALM and the other posteriorly

in the PLM. Importantly, both the anterior and posterior embryonic domains show expanded

marker gene expression in cavefish compared to surface fish. It is noteworthy that expansion of

the posterior hematopoietic domain can also be induced in zebrafish by overexpressing LMO2

together with its partner transcription factor SCL/TAL1 (Gering et al., 2003), suggesting that

teleosts have the capacity to modulate blood cell numbers by changing the size of primitive

erythropoietic domains during development. The existence of erythrocyte forming capacity in the

ALM of surface fish, albeit smaller than in cavefish, suggests that the expansion of primitive

erythropoiesis may pre-adapt *Astyanax* for colonizing hypoxic cave environments.

The expansion of erythropoiesis into the ALM during primitive hematopoiesis in

Astyanax contrasts strikingly to zebrafish, medaka, and other teleosts that have been investigated,

in which primitive erythropoiesis is confined to the PLM, and the ALM is devoted only to

myelopoiesis (Detrich et al., 1995; Govoni et al., 2005, Moriyama et al., 2010). Expression of

the macrophage marker *lcp1* (Herbomel et al., 2001) indicated that myelopoietic precursors had

already differentiated into macrophages as erythrocyte progenitors were being formed in the

ALM during early *Astyanax* development. Cell tracing studies have shown that blood precursor

cells can produce either erythrocytes or macrophages, but not both cell types (Warga et al.,

2009), indicating an opposing relationship between the erythroid and myeloid cell populations.

However, this relationship can be reversed by knockdown of the myeloid-controlling

transcription factor PU.1, which induces blood precursors of the ALM to form erythrocytes

instead of macrophages (Rhodes et al., 2005). Apparently, teleosts are potentially able to

undergo erythropoiesis in the ALM and PLM, but in most species this capacity is normally

confined to the PLM. In *Astyanax,* the balanced relationship between myeloid and erythroid cell

specification appears to have tilted toward the production of erythrocytes. However, the

development of erythrocytes in the ALM may not be at the expense of fewer myeloid cells

because there is also an overall expansion of the ALM (and PLM), which may promote the

specification of more blood precursor cells. In future studies, it will be interesting to determine

how the expansion of erythropoietic potential is controlled in cavefish.

Specification of cells in the primitive blood lineage can be traced to their positions during

gastrulation in zebrafish (Kimmel et al., 1990): ALM macrophages are originally derived from

progenitors on the dorsal side of the embryo, and PLM erythrocytes originate from the ventral

side of the embryo (Davidson and Zon, 2004; Warga et al., 2009). Erythrocyte progenitors

corresponding to the ALM and PLM are also likely to be positioned on the dorsal or ventral sides

respectively in gastrulating *Astyanax* embryos. Therefore, increased erythropoiesis in the ALM

suggests that a relative ventralization of hematopoietic development has evolved in cavefish

compared to surface fish. This possibility is supported by expanded erythroid development

induced by overexpression of the ventralizing factors *bmp2b* and *bmp7* in zebrafish gastrulae

(Lengerke et al., 2008) and is in line with evidence implicating the importance of maternal

factors in dorsal-ventral axis determination of cavefish embryos (Ren et al; 2018; Ma et al.,

2018; Torres-Paz et al., 2019). Maternal control of primitive erythropoiesis may promote

maximal oxygen distribution at the early stages of development in hypoxic environments. This

finding extends our understanding of evolutionary differences in early hematopoiesis to events

beginning during oogenesis.

In the present investigation, we have examined primitive hematopoiesis and red blood

cells in the Pachón cavefish population. About 30 different *Astyanax* cavefish populations have

been described in northeastern Mexico, and some of these cavefish are thought to have been

derived independently from surface fish ancestors (Gross, 2012; Herman et al., 2018). It is

conceivable that diverse cavefish lineages inhabit cave pools with different levels of

oxygenation. Therefore, in the future it would be interesting to determine the status of

erythrocytes in additional cavefish populations. It will also be important to determine if other

aquatic cave animals modify their oxygen delivery systems to cope with hypoxia.

322   Teleosts show considerable diversity in utilizing hemoglobin and red blood cells for

oxygen transport during development: some species begin hemoglobin synthesis during

embryogenesis, whereas others postpone hemoglobin and erythrocyte production until much

later in larval development (Wells, 2009). Our studies on the induction of hemolytic anemia

with PHZ confirm the functional significance of red blood cells produced during early *Astyanax*

development. At the same PHZ concentrations, surface fish, but not cavefish with residual red

blood cells, show negative effects on growth, and hypoxia treatment inhibited growth in surface

fish but not in cavefish. These results suggest that red blood cells are necessary for oxygen

transport at early stages of *A. mexicanus* development and that increased erythrocytes may be

advantageous in cavefish. This is contrary to zebrafish, where PHZ induced anemia showed that

red blood cells are not required during the first few weeks of development under normoxic

conditions (Pelster and Burggren, 1996), although under severe hypoxia zebrafish embryos also

use red blood cells to increase the supply of oxygen (Rombough and Drader, 2009).

335   Teleosts have evolved different strategies to cope with environments depleted in oxygen,

including changes in gill morphology (Sollid et al., 2003; Dhillon et al., 2013), behavioral

avoidance of hypoxic waters (Congleton, 1980; Mandic et al., 2009), and altered metabolism

(Torres et al., 2012). *Astyanax* surface fish larvae appear to deal with hypoxia by conserving

energy and reducing growth. It is likely that growth is regulated similarly to zebrafish via

upregulation of *igfbp1a* (Kajimura et al., 2005). The upregulation of *igfbp1* under hypoxia is

regulated by HIF1 (Kajimura et al., 2006). *Astyanax* cavefish, which are presumably unable to

avoid hypoxic waters in their natural environment, show increased expression of some of the *hif1*

genes in either hypoxic or normoxic conditions. The constitutive increase in *hif1* related gene

expression may have conferred growth resistance to hypoxia, even under conditions in which

igfbp1 is upregulated, and may be responsible for increased red blood cell number.

The increased expression of the metabolic gene *hk1*in cavefish compared to surface fish

indicates an increase in glycolysis, consistent with the observations of Krishnan et al. (2019).

The increase in glycolysis does not seem to affect the TCA cycle since *ldha* and *pdk1* are not

differentially expressed between cavefish and surface fish (Medley et al., 2020). Wild captured

cavefish also have higher expression of the genes encoding enzymes involved in glutathione

metabolism compared to wild caught surface fish (Krishnan et al., 2019; Medley et al., 2020).

Glutathione is an important player in the mitigation of ROS, and the Pentose Phosphate Pathway

produces a large portion of the NADPH used in the production of glutathione (Stincone et al.,

2015). Therefore, the constitutive expression of *hif1*-genes in cavefish could play a role in ROS

mitigation via glutathione during hypoxia (Birnie-Gauvin et al., 2017).

In conclusion, the present investigation has revealed two potential adaptations that may

have permitted *Astyanax* cavefish to cope with hypoxic cave waters. First, cavefish have

enhanced erythrocyte development, and as a maternal effect this phenotype may have increased

the capacity of cavefish to carry and distribute essential oxygen to tissues and organs early in

development. Second, cavefish have adjusted to hypoxic conditions by evolving constitutively

high expression levels of some of the *hif1* genes. Together, these changes may be important for

the successful adaptation of *Astyanax* to hypoxic subterranean environments.

Materials and methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Astyanax mexicanus* surface fish)	*hemoglobin subunit beta-2-like*, *hbb2*	NCBI:GeneID 111196758, Ensembl:ENSAMXG000000031275		
gene (*Astyanax mexicanus* surface fish)	*growth factor independent 1A transcription repressor a, gfi1aa*	NCBI:GeneID103029320 Ensembl:ENSAMXG00000006669		
gene (*Astyanax mexicanus* surface fish)	*LIM domain only 2 (rhombotin-like 1), lmo2*	NCBI:GeneID 111190647 Ensembl:ENSAMXG000000032986		
gene (*Astyanax mexicanus* surface fish)	*lymphocyte cytosolic protein, lcp1*	NCBI:GeneID 103042114 Ensembl:ENSAMXG000000012855		
gene (*Astyanax mexicanus* surface fish)	*hbbe2*	Ensembl:ENSAMXG000005017210		
gene (*Astyanax mexicanus* surface fish)	*ribosomal protein L13a, rpl13a*	NCBI:geneID: 103025160 Ensembl:ENSAMXG00000033532		
gene (*Astyanax mexicanus* surface fish)	*guanine nucleotide binding protein (G protein) beta polypeptide 1b, gnb1b*	NCBI:GeneID 103023031 Ensembl:ENSAMXG000000040710		

gene (*Astyanax mexicanus* surface fish)	*hypoxia inducible factor 1 subunit alpha a, hif1aa*	NCBI:GeneID 103022448 Ensembl:ENS AMXG000000 39550		
gene (*Astyanax mexicanus* surface fish)	*hypoxia inducible factor 1 subunit alpha b, hif1ab*	NCBI:GeneID 103033873 Ensembl:ENS AMXG000000 19342		
gene (*Astyanax mexicanus* surface fish)	*hypoxia inducible factor 1 subunit alpha like, hif1alike, hif1al*	NCBI:GeneID 103027586 Ensembl:ENS AMXG000000 08564		
gene (*Astyanax mexicanus* surface fish)	*hypoxia inducible factor 1 subunit alpha like 2, hif1alike2, hif1al2*	NCBI:GeneID 103041845 Ensembl:ENS AMXG000000 07272		
gene (*Astyanax mexicanus* surface fish)	*hexokinase 1, hk1*	NCBI:GeneID 103028521 Ensembl:ENS AMXG000000 12670		
gene (*Astyanax mexicanus* surface fish)	*glucose-6-phosphate dehydrogenase, g6pd*	NCBI:GeneID 103035433 Ensembl:ENS AMXG000000 17509		
gene (*Astyanax mexicanus* surface fish)	*lactate dehydrogenase A4, ldha*	NCBI:GeneID 103047177 Ensembl:ENS AMXG000000 32467		
gene (*Astyanax mexicanus* surface fish)	*pyruvate dehydrogenase kinase 1, pdk1*	NCBI:GeneID 103033744 Ensembl:ENS AMXG000000 39808		

gene (*Astyanax mexicanus* surface fish)	*insulin-like growth factor binding protein 1a, igfbp1a*	NCBI:GeneID 103033920 Ensembl:ENSAMXG00000009512		
strain, strain background (Escherichia coli)	One Shot Top10 chemically competent cells, *E.coli*	Invitrogen	Cat# 404003	
biological sample (*Astyanax mexicanus* surface fish)	Surface fish, SF, Texas	Jeffery lab		
biological sample (*Astyanax mexicanus* cave fish)	Cave fish, CF, Pachón,	Jeffery lab		
antibody	Anti-Dig-AP (sheep polyclonal)	Roche	Cat# 11093274910 RRID:AB_514497	(1:5000)
recombinant DNA reagent	pCRII-TOPO dual promotor vector	Invitrogen	Cat# 45-0640	
peptide, recombinant protein	Proteinase K	Roche	Cat# 03115887001	
peptide, recombinant protein	DNaseI I, RNase-free	Thermo-Scientific	Cat# EN0521	
commercial assay or kit	SYBR Premix Ex Taq (Tli RNaseH Plus)	Takara	Cat# RR420L	
commercial assay or kit	T7 RNA polymerase	Roche	Cat# 10881767001	
commercial assay or kit	SP6 RNA polymerase	Roche	Cat# 10810274001	

commercial assay or kit	ReadyMix Taq PCR Reaction Mix	Sigma	Cat# P4600	
commercial assay or kit	Dig RNA labelling Mix	Roche	Cat# 11277073910	
commercial assay or kit	Blocking reagent	Roche	Cat# 11096176001	
commercial assay or kit	BM Purple AP Substrate, precipitating	Roche	Cat# 11442074001	
commercial assay or kit	SuperScript III First strand synthesis supermix	Invitrogen	Cat# 18080-400	
commercial assay or kit	SuperScript IV VILO mastermix with ezDNase	Invitrogen	Cat# 11766050	
chemical compound, drug	paraformaldehyde	Electron Microscopy Sciences	Cat# 15710	
chemical compound, drug	o-dianisidine	Sigma	Cat# D9143	
chemical compound, drug	phenylhydrazine hydrochloride	Sigma	Cat# 114715	
chemical compound, drug	Trizol	Life Technologies	Cat# 15596018	

software, algorithm	Image-J	Imagj.nih.gov	RRID:<u>SCR 003070</u>	
software, algorithm	JMP Pro 14	SAS Institute Inc.		
other	CytoOne 24 well plate	USA Scientific	Cat# CC76727424	
other	Cellstar 12 well cell culture plate	Greiner	Cat# 665180	
other	Netwell inserts	Corning	Cat# 3478	
other	Hypoxia chamber, ProOx Model P110	BioSpherix		

Biological materials

Astyanax mexicanus surface fish and cavefish were obtained from laboratory stocks

descended from collections in Balmorhea Springs State Park, Texas and Cueva de El Pachón,

Tamaulipas, Mexico respectively. Fish were raised in a constant flow culture system as described

previously (Jeffery et. al. 2000; Ma et al., 2021). Embryos were obtained by natural spawning

and reared at 23°C. Fish handling and husbandry protocols were approved by the University of

Maryland Animal Care and Use Committee (IACUC #R-NOV-18-59) (Project 1241065-1) and

conformed to National Institutes of Health guidelines.

Generation of reciprocal crosses

The offspring of reciprocal crosses were generated by placing a female cavefish and a

male surface fish or a female surface fish and a male cavefish together in a tank and inducing

spawning by increased feeding and water temperature (Ma et al., 2021). Controls were generated

by crossing a surface fish male and surface fish female and a cavefish male and cavefish female.

Quantification of circulating blood cells

The number of circulating blood cells was analyzed in surface fish and cavefish larvae

immobilized by treatment with 2 μg ml^{-1} tricaine buffered with Tris to pH 7.0 (Western

Chemical, Inc, Ferndale, WA, USA) at 36 hrs. post-fertilization (hpf) and placed in water on a

concave microscope slide. Blood flow was imaged as cells flowed through the Duct of Cuvier

(Isogai, 2001) for 10 seconds using a stereoscope (Olympus SZX12) or a compound microscope

(Zeiss, Axioskop2) with 50x magnification and a 5 MP, Color, AmScope Microscope Eyepiece

Camera (MD500). Recordings were captured with Photobooth on a Macbook computer via USB

(OS 10.13.6). Videos were rendered into separate frames in Adobe Photoshop (CC2017, Adobe

Inc., San Jose, CA, USA) and processed and analyzed in Image-J (Schindelin et al.,

2012), (RRID:SCR_003070). After conversion to 32 bit, a data image showing the movement of

particles was created by subtracting two frames 0.15 seconds apart. This data image was used in

the plugin TrackMate (Tinevez et al., 2017) to analyze the number of moving blood cells in a

region of interest (ROI) covering 50% or more of the blood stream over the yolk (150x150

pixels). Using the LoG detector, moving blood cells were automatically detected by setting the

"blob diameter" to 5 and the threshold to 2. An automatic quality threshold was used to obtain

the number of circulating blood cells and checked against the recording for accuracy. The blood

flow of larvae from at least 8 different clutches of eggs per morph were analyzed.

Staining and quantification of erythrocytes

Red blood cells were stained at 36 hpf with 0.6 mg ml^{-1} o-dianisidine (Sigma-Aldrich, St.

Louis, MO, USA), 0.01 M sodium acetate (pH 4.5), 0.65 % H_2O_2 for 15 min in the dark (Iuchi

and Yamamoto, 1983). The stained embryos were rinsed in phosphate buffered saline (PBS),

fixed in 4% paraformaldehyde (PFA), and imaged as described below for *in situ* hybridization

50x magnification. The number of blood cells were counted manually in an ROI (150x150

pixels) by hand. Larvae from 3 separate clutches of eggs were used per morph.

In situ hybridization

*In sit*u hybridizations were performed using probes to the *hemoglobin subunit beta-2-like*

(*hbb2*), *growth factor independent 1A transcription repressor* (*gfi1aa*), *LIM domain only 2*

(*lmo2*), or *lymphocyte cytosolic protein 1* (*lcp1*) genes. Genes were cloned from a 24 hpf surface

fish cDNA library and 10 hpf, 24 hpf and 30 hpf cavefish cDNA libraries using the pCRII TOPO

dual promoter vector (ThermoFisher Scientific, Waltham, MA, USA) transformed into One shot

Top10 cells and the following primers: *hbb2* (24 hpf SF; ENSAMXG00000031275: 5'-

gcaggacaagtagaaacctcaaagtc-3' and 5'-tttcgtaagggcagagcctaca-3'), *gfi1aa* (24 hpf CF;

ENSAMXG00000006669:5'-gaaggtctgcgctcgtgatatt-3' and 5'- agttatccgcggtgtgaacag-3'), *lmo2*

(10 hpf CF; ENSAMXG00000032986: 5'-ggcctctacaatcgagaggaaa-3' and 5'

taccaagttgccgtttagtttgg-3'), and *lcp1* (36 hpf CF; ENSAMXG00000012855: 5'-

aggccttcagcaaagttgatgtg-3' and 5'-ttcaggtcctctgcaccgatatt-3'). DIG labeled probes were made

using SP6 or T7 transcription kits (Roche, Mannheim, Germany) from linearized plasmid or, in

the case of *gfi1aa,* from a PCR product made with the cloned cDNA as template and the above

mentioned primers, after a RNA-polymerase promoter site was added to the 5' end.

In situ hybridizations were performed as previously described (Ma et al., 2014). Briefly,

stored (-20^{o}C) 4% PFA fixed and methanol dehydrated embryos were rehydrated stepwise into

PBS, fixed with 4% PFA, digested with proteinase K, fixed with 4% PFA, and hybridized with

probes at 60^{o}C for 16 hours. Un-hybridized probe was removed by 2 X and 0.2 X SSCT (150

mM sodium chloride; 15 mM sodium citrate; 0.1% Tween 2) stringency washes followed by

incubation in MABT blocking solution (Roche) and subsequently with anti-DIG-AP Fab

fragments (Roche), (RRID:AB_514497). Embryos were rinsed in MABT buffer and PBS,

equilibrated in AP buffer, and stained with BM-Purple (Roche). The stained embryos were

imaged and photographed using a Zeiss Discovery V20 stereoscope with a Zeiss AxioCam HRc

camera.

Induction of hemolytic anemia with phenylhydrazine

Phenylhydrazine hydrochloride (PHZ; Sigma-Aldrich, St. Louis, MO, USA) was used to

induce hemolytic anemia (Houston et al., 1988). At 14 hpf, 20 manually de-chorionated embryos

were placed in clean fish system-water containing a particular concentration of PHZ or clean fish

system water as a control. Embryos were incubated with PHZ for 28 hours when blood

circulation was clearly visible over the yolk. Embryos were fixed and used for *in situ*

hybridization with a *hbb2* probe. In a separate experiment, embryos were imaged on a Zeiss

Axioskop2 microscope with a Axiocam 503 camera and measurements of tail length (the axial

length from the posterior end of the yolk mass to the tip of the tail) were made in Image-J. After

imaging, PHZ treated embryos and controls were stained with o-dianisidine and imaged for

blood cell quantification as described above. Because low numbers of blood cells were present

after PHZ treatment, blood cells were counted in a larger area covering the whole anterior side of

the yolk in lateral view (400 pixels wide). Larvae of two separate clutches of eggs were used per

morph for the latter experiment.

Surface fish larvae were also treated with PHZ at 34 hpf for 1 hr., rinsed extensively for 1

hr., imaged with a microscope as described above, and left for 16 hrs. before imaging again. At

the end of the experiment, larvae were stained with o-dianisidine. The length of the post-anal tail

was measured using Image-J and growth determined as the differential between the two time

points. Blood cell number was determined by manual counting as described above.

Laboratory hypoxia treatment and growth determinations

A hypoxia chamber (ProOx Model P110, BioSpherix, Parish, NY, USA) was used to

create a hypoxic laboratory environment in which oxygen was reduced by nitrogen gas (HP,

Airgas, Hyattsville, MD, USA). At 30 hpf larvae were exposed to hypoxia (1 mg/L oxygen) for

18 hours or to normoxia (outside of the chamber). Larvae were placed individually in wells of a

24 well plate (CytoOne, USA scientific, Ocala, FL, USA) containing 2 ml clean fish system-

water. Twenty-four larvae were used per treatment for surface fish and 48 larvae for cavefish

from two separate clutches of eggs per morph. To measure post-anal tail growth, each larva was

imaged before placement in a well and again at the end of the treatment using a microscope as

described above. Image-J was used to measure post-anal tail length (Fig. 6A) and growth

determined as the differential between the two time points. To measure eye size, eye diameter

was measured after the 18 hr. hypoxia or normoxia treatment with image -J. After imaging,

larvae were stained with o-dianisidine and blood cells were counted from images as described

above.

RNA isolation and quantitative real time polymerase chain reactions

To compare the extent of early erythropoiesis between cavefish and surface fish, the

expression of *hemoglobin beta embryonic 2* (*hbb2*) and *hbbe2*, (*hbbe2* encodes an *A. mexicanus*

embryonic β-hemoglobin most similar to zebrafish *hbbe1*; Ganis et al., 2012) genes and the

gfi1aa gene were quantified at 10, 24 and 60 hpf. Three potential reference genes were

evaluated: *ribosomal protein L13a* (*rpl13a*), *actin alpha 1* (*acta1b*), and *lsm12a.* Only *rpl13a*

did not vary over time or between surface and cavefish and was therefore used as the final

reference gene. RNA was extracted from 30 embryos with Trizol (ThermoFisher), treated with

RNase-free DNase I (ThermoFisher), cleaned and concentrated by phenol/chloroform extraction

and precipitated with ammonium acetate and ethanol. Poly(A)-primed cDNA was made with

SuperScript III First Strand Synthesis SuperMix (ThermoFisher) and used in qPCR with Takara

SYBR Premix Ex Taq (Tli RNaseH Plus) (Takara Bio USA Inc., Mountain View, CA, USA) and

LC480 (Roche).

In the laboratory hypoxia experiments, RNA was isolated from 24 larvae placed in the 15

mm Netwell insert (Corning, Corning, New York, USA) in a well of a 12 well plate. Two hours

after the start of the hypoxia treatment the larvae were collected in Trizol (ThermoFisher) within

1 min after taking them out the hypoxia chamber by removing the Netwell inserts from the wells,

decanting excess water, and the larvae were immersed in Trizol. RNA was isolated for 6 samples

of 24 larvae each with 3 samples per clutch of eggs. Two separate clutches of eggs were used per

morph.

The amounts of *hypoxia inducible factor 1(hif1aa)* and its duplicate gene *hif1ab, hypoxia*

inducible factor 1-like (hif1alike) and its duplicate gene *hif1alike2* (equivalent to *hif3a*), *insulin*

growth factor binding protein 1a (igfbp1a), hexokinase 1 (hk1), glucose-6-phosphate

dehydrogenase (G6PD), lactate dehydrogenase a (ldha) and *pyruvate dehydrogenase kinase 1*

(*pdk1*) were quantified by qPCR in surface fish and cavefish larvae exposed to hypoxic or

normoxic conditions. We evaluated *rpl13a, acta1b, endoplasmic reticulum protein 44 (erp44)*

and *guanine nucleotide binding protein ß polypeptide 1b (gnb1b)* as potential reference genes:

gnb1b had Ct values closest to the Ct values of the queried genes, did not differ between

normoxic and hypoxic surface fish or cavefish, and was chosen as a reference gene. RNA was

extracted from 24 embryos and treated with ezDNase and cDNA was made with SuperScript IV

VILO mastermix (ThermoFisher). The primers used in qPCR analysis are shown in Table 2.

A ΔCt for each gene was calculated by subtracting the average Ct value of each reference

gene. For comparison of gene expression in cavefish to surface fish over time, ΔΔCt was

calculated by subtracting the average ΔCt of surface fish at 10 hpf from each ΔCt for each gene.

For comparison of gene expression under normoxic and hypoxic conditions, the ΔCt of surface

fish under normoxic conditions was subtracted from the ΔCt for each gene. For graphical

representation, the fold change was calculated as $2^{-(\Delta\Delta Ct)}$, where values >1 show an increase and

values <1 a decrease. Variation was expressed as the range of fold change $2^{-(\Delta\Delta Ct+stdev\Delta\Delta Ct)}$ for the

upper value or $2^{-(\Delta\Delta Ct-stdev\Delta\Delta Ct)}$ for the lowest value.

Statistics

Statistics were done using JMP pro 14 (SAS Institute Inc.). Normal distribution of data

was determined using the Shapiro Wilk test. Comparison of blood cell number was done using

Wilcoxon/Kruskal-Wallis Rank Sums test (1F N = 72, 1G N=36) (p< 0.05). Expression as $\Delta\Delta$Ct

of *hbb2, hbbe2* and *gfi1aa* over time was analyzed per gene with a two-way ANOVA followed

by pairwise comparison with Student's t-test (*hbb2* N=8, *hbbe2* N=4, *gfi1aa* N=3), (p<0.05).

Comparisons of blood cell number in reciprocal hybridization experiments were analyzed using a

Wilcoxon/Kruskal-Wallis Rank Sums test followed by Wilcoxon for each pair (N=20), (p<0.05).

Differences in tail length between surface and cavefish at different concentrations of PHZ were

analyzed using Wilcoxon/Kruskal-Wallis Rank Sums test followed by Wilcoxon for surface fish

(SF) or cavefish (CF) for each pair (SF 0, 2 mg L^{-1} N=23, SF 1, 3 mg L^{-1} N=24, CF 0, 1, 2 mg L^{-1}

$^{-1}$ N=24, CF 3 mg L^{-1}), (p<0.05). Comparison of blood cell number at different concentrations of

PHZ were analyzed with Wilcoxon/Kruskal-Wallis Rank Sums test followed by Wilcoxon for

each pair (SF 0, 1, 2 and CF 3 mg L^{-1} N=20, SF 3 mg L^{-1} N=10, CF 0 mg L^{-1} N=19, CF 1 mg L^{-1}

N=21, CF 2 mg L^{-1} N=22), (p<0.05). Post-anal tail growth at 0 and 5 mg L^{-1} PHZ was analyzed

with a Student's t-test (0 mg L^{-1} N=19, 5 mg L^{-1} N=20), (p<0.05) and cell number with

Wilcoxon/Kruskal-Wallis Rank Sums test (N=10), (p<0.05). Post-anal tail growth under

normoxic and hypoxic conditions was analyzed with Wilcoxon/Kruskal-Wallis Rank Sums test

followed by Wilcoxon for each SF and CF pair (SF-N and SF-H N=23, CF-N N=46 and CF-H

N=47), (p<0.05). Cell number under normoxic and hypoxic conditions was analyzed with one-

way ANOVA followed by Student's t-test (SF-N N=22, SF-H N=21, CF-N N=47, CF-H N=45),

(p<0.05). The diameter of eye was analyzed by one-way ANOVA followed by Student's t-test

(SF-N N=18, SF-H N=14, CF-N N=30, CF-H N=28), (p<0.05). Expression as $\Delta\Delta$Ct of genes

under normoxic and hypoxic conditions was analyzed per gene by one-way ANOVA followed

by Student's t-test or when variances where unequal with Wilcoxon/Kruskal-Wallis Rank Sums

test followed by Wilcoxon for each pair (N = 6), (p<0.05).

Acknowledgments

539   This research was supported by NIH grant EY024941 to WRJ. We thank Ruby

540 Dessiatoun and Karina Lacroix for maintenance of the *A. mexicanus* colony and Mandy Ng for

541 technical assistance.

Competing interests

Funding

548   Funding for this study was provided by the National Institutes of Health (EY024941 to

549 WRJ)

References

Aspiras AC, Rohner N, Martineau B, Borowsky RL, Tabin CJ. 2015. Melanocortin 4

receptor mutations contribute to the adaptation of cavefish to nutrient-poor conditions.

Proceedings of the National Academy of Sciences of the United Stated of America **112**: 9668-

9673. doi:10.1073/pnas.1510802112

Birnie-Gauvin K, Costantini D, Cooke SJ, Willmore WG. 2017. A comparative and

evolutionary approach to oxidative stress in fish: A review. *Fish and Fisheries* **18**:928-942.

doi.org/10.1111.faf.12215.

Bishop RE, Kakuk B, Torres JJ. 2004. Metabolism and proximate composition of Caribbean

troglobitic crustaceans with observations on the water chemistry of two anchialine caves.

Journal of Crustacean Biology. **24**:379-392. https://doi.org/10.1651/C-2459

Blin M, Tine E, Meister L, Elipot Y, Bibliowicz J, Espinasa L, Rétaux, S. 2018**.**

Developmental evolution and developmental plasticity of the olfactory epithelium and olfactory

skills in Mexican cavefish. *Developmental Biology* **441**:242-251**.**

Boggs T, Gross J. 2021. Reduced oxygen as an environmental pressure in the evolution of the

blind Mexican cavefish. *Diversity* **13**:26. doi.org/10.3390/d13010026

Brownlie A, Hersey C, Oates AC, Paw BH, Falick AM, Witkowska HE, Flint J, Higgs D,

Jessen J, Bahary N, Zhu H, Lin S, Zon L. 2003. Characterization of embryonic globin genes of

the zebrafish. *Developmental Biology*. **255**:48-61. doi: 10.1016/s0012-1606(02)00041-6

Cai X, Zhou Z, Zhu J, Liao Q, Zhang D, Liu X, Wang J, Ouyang G, Xiao W. 2020.

Zebrafish Hif3α modulates erythropoiesis via regulation of gata1 to facilitate hypoxia tolerance.

Development. **147**: dev185116. doi: 10.1242/dev.185116.

Carroll KJ, North TE. 2014.. Oceans of opportunity: exploring vertebrate hematopoiesis in

zebrafish. *Experimental Hematology*. **42**:684-696. doi:10.1016/j.exphem.2014.05.002

Congleton JL. 1980. Observations on the responses of some southern California tide pool fishes

to nocturnal hypoxic stress. *Comparative Biochemical Physiology*. **66A**:719-122.

https://doi.org/10.1016/0300-9629(80)90026-2

Cooney J D, Hildick-Smith GJ, Shafizadeh E, McBride PF, Carroll KJ, Anderson H, Shaw

GC, Tamplin OJ, Branco DS, Dalton AJ, Shah DI, Wong C, Gallagher PG, Zon LI, North

TE, Paw BH. 2013. Teleost *growth factor independence* (*gfi*) genes differentially regulate

successive waves of hematopoiesis. *Developmental Biology* 373:431-441. doi:

10.1016/j.ydbio.2012.08.015

Cui XG, Han ZT, He SH, Wu XD, Chen TR, Shao CH, Chen DL, Su N, Chen YM, Wang T,

Wang J, Song D-W, Yan W-J, Yang X-H, Liu T, Wei H-F, Xiao J. 2017. HIF1/2α mediates

hypoxia-induced LDHA expression in human pancreatic cancer cells. *Oncotarget* 8:24840-

24852. doi:10.18632/oncotarget.15266.

Culver DC, Pipan T. 2009. *The Biology of Caves and Other Subterranean Habitats.* Oxford

UK: Oxford University Press.

Davidson AJ, Zon LI. 2004. The 'definitive' (and 'primitive') guide to zebrafish hematopoiesis.

Oncogene **23**:7233-7246. doi: 10.1038/sj.onc.1207943.

Detrich HW, Kieran M W, Chan FY, Barone LM, Yee K, Rundstadler J, Pratt S, Ransom

D, Zon LI. 1995. Intraembryonic hematopoietic cell migration during vertebrate development.

Proceedings of the National Academy of Sciences of the United States of America **92**:10713-

10717. doi: 10.1073/pnas.92.23.10713

Dhillon RS, Yao L, Matey V, Chen B-J, Zhang A-J, Cao Z-D, Fu S-J, Brauner CJ, Wang

YS, Richards JG. 2013. Interspecific differences in hypoxia-induced gill remodeling in carp.

Physiology Biochemisty Zoology 86:727-739. doi: 10.1086/673180

Fumey J, Hinaux H, Noirot C, Thermes C, Rétaux S, Casane D. 2018. Evidence for late

Pleistocene origin of *Astyanax mexicanus* cavefish. *BioMed Central Evolutionary Biology*.

18(1):43. doi: 10.1186/s12862-018-1156-7.

Ganis J J, Hsia N, Trompouki E, de Jong JLO, DiBiase A, Lambert J S, Zhiying J, Sabo

PJ, Weaver M, Sandstrom R, Stamatoyannopoulos JA, Zhou Y, Zon LI. 2012. Zebrafish

globin switching occurs in two developmental stages and is controlled by the LCR.

Developmental Biology **366**:185-194. doi: 10.1016/j.ydbio.2012.03.021

Gering M, Yamada Y, Rabbitts TH, Patient RK. 2003. Lmo2 and Scl/Tal1 convert non-axial

mesoderm into haemangioblasts which differentiate into endothelial cells in the absence of

Gata1. *Development* **130**:6187-6199. doi:10.1242/dev.00875

Giaccia A J, Simon MC, Johnson R. 2004. The biology of hypoxia: the role of oxygen

sensing in development, normal function, and disease. *Genes & De*velopment **18**: 2183-2194.

Gore AV, Pillay L M, Galanternik MV, Weinstein BM. 2018. The zebrafish: A fintastic

model for hematopoietic development and disease. *Wiley Interdisciplinary Reviews in*

Developmental Biology **7**:e312-328. doi: 10.1002/wdev.312.

Govoni JJ, West MA, Bonaventura J, Goddette G, Jenkins TE. 2005. The ontogeny of

hematopoiesis in the marine teleost *Leiostomus xanthurus* and a comparison of the site of initial

hematopoiesis with *Opsanus tau*. *Journal of Fish Biology*. **67**:696-712.

https://doi.org/10.1111/j.0022-1112.2005.00770.x

Gross JB. 2012. The complex origin of *Astyanax* cavefish. *BioMed Central Evolutionary*

Biology. **12**:105-116. doi: 10.1186/1471-2148-12-105.

Haase VH. 2013**.** Regulation of erythropoiesis by hypoxia-inducible factors. *Blood Review*

27:41-53**.** doi: 10.1016/j.blre.2012.12.003

Herbomel P, Thisse B, Thisse C. 1999. Ontogeny and behaviour of early macrophages in the

zebrafish embryo. *Development* **126**:3735-3745.

Herbomel P, Thisse B, Thisse C. 2001. Zebrafish early macrophages colonize cephalic

mesenchyme and developing brain, retina and epidermis through a M-CSF receptor-dependent

invasive process. *Develomental Biology*. **238**:274-288. doi: 10.1006/dbio.2001.0393

Herman A, Brandvain Y, Weagley J, Jeffery WR, Keene AC, Kono TJY, Bilandžija H,

Borowsky R, Espinasa L, O'Quin K, Ornelas-García CP, Yoshizawa M, Carlson B,

Maldonado E, Gross JB, Cartwright RA, Rohner N, Warren WC, McGaugh SE. 2018. The

role of gene flow in rapid and repeated evolution of cave related traits in Mexican tetra, *Astyanax*

mexicanus. *Molecular Ecology* **27**: 4397-4416. doi: 10.1111/mec.14877

Houston AH, Murad A, Gray JD. 1988. Induction of anemia in goldfish, *Carassius auratus* L.,

by immersion in phenylhydrazine hydrochloride. *Canadian Journal of Zool*ogy **66**:729-

736. https://doi.org/10.1139/z88-108.

Hüppop K. 1986. Oxygen consumption of *Astyanax fasciatus* (Characidae, Pisces): a

comparison of epigean and hypogean populations. *Environmental Biology of Fishes* **17**: 299-308.

Isogai S, Horiguchi M, Weinstein BM. 2001. The vascular anatomy of the developing

zebrafish: an atlas of embryonic and early larval development. *Developmental Biology* 230: 278-

301. doi: 10.1006/dbio.2000.9995

Iuchi I, Yamamoto M. 1983. Erythropoiesis in the developing rainbow trout, *Salmo gairdneri*

irideus: histochemical and immunochemical detection of erythropoietic organs. *Journal of*

Experimental Zoology. **226**:409-417. doi: 10.1002/jez.1402260311

Jeffery WR, Strickler A., Guiney S, Heyser D, Tomarev SI. 2000**.** *Prox 1* in eye degeneration

and sensory compensation during development and evolution of the cavefish *Astyanax*.

Development Genes & Evolution. **210**:223-230**.** doi:10.1007/s004270050308.

Kajimura S, Aida K, Duan C. 2005. Insulin-like growth factor-binding protein-1 (IGFBP-1)

mediates hypoxia-induced embryonic growth and developmental retardation. *Proceedings of the*

National Academy of Science of the United States of America **102**: 1240-1245. doi:

10.1073/pnas.0407443102.

Kajimura S, Aida K, Duan C. 2006. Understanding hypoxia-induced gene expression in early

development: in vitro and in vivo analysis of hypoxia-inducible factor 1-regulated zebra fish

insulin-like growth factor binding protein 1 gene expression. *Molecular Cell Biology.* **26:**1142-

1155. doi: 10.1128/MCB.26.3.1142-1155.2006

Kamei H. 2020. Oxygen and embryonic growth: the role of insulin-like growth factor signaling.

General & Comparative Endocrinology. **294**:113473. doi: 10.1016/j.ygcen.2020.113473.

Kimmel CB, Warga RM, Schilling TF. 1990. Origin and organization of the zebrafish fate

map. *Development* **108**:581-594.

Krishnan J, Persons JL, Peuß R, Hassan H, Kenzior A, Xiong S, Olsen L, Maldonado E,

Kowalko JE., Rohner N. 2019. Comparative transcriptome analysis of wild and lab populations

of *Astyanax mexicanus* uncovers differential effects of environment and morphotype on gene

expression. *Journal of Experimental Zoology B. Molecular & Developmental Evolution* **334**:530-

539. doi: 10.1002/jez.b.22933

Kulkarni RP, Tohari S, Ho A, Brenner S, Venkatesh B. 2010. Characterization of a hypoxia-

response element in the *Epo* locus of the pufferfish, *Takifugu rubripes*. Marine Genomics **3**:63-

70. doi: 10.1016/j.margen.2010.05.001.

Le Guyader D, Redd MJ, Colucci-Guyon E, Murayama E, Kissa K, Briolat V, Mordelet E,

Zapata A, Shinomiya H, Herbomel P. 2008. Origins and unconventional behavior of

neutrophils in developing zebrafish. *Blood* **111**:132-141. doi: 10.1182/blood-2007-06-095398

Lengerke C, Schmitt S, Bowman TV, Jang I. H., Maouche-Chretien L, McKinney-

Freeman, S, Davidson, AJ, Hammerschmidt M, Rentzsch F, Green JB, Zon L, et al.. 2008.

BMP and Wnt specify hematopoietic fate by activation of the Cdx-Hox pathway. *Cell Stem*

Cell **2**:72−82. doi:10.1016/j.stem.2007.10.022

Ma L, Parkhurst A, Jeffery W R. 2014. The role of a lens survival pathway including *sox2* and

αA-crystallin in the evolution of cavefish eye degeneration. *Evo Devo*. **5**: 28-41. doi:

10.1186/2041-9139-5-28

Ma L, Strickler AG, Parkhurst A, Yoshizawa M, Shi J, Jeffery WR. 2018. Maternal genetic

effects in *Astyanax* cavefish development. *Developmental Biology*. **441**:209-220. doi:

10.1016/j.ydbio.2018.07.014

Ma L, Dessiatoun R, Shi J, Jeffery WR. 2021. Incremental Temperature Changes for Maximal

Breeding and Spawning in *Astyanax mexicanus. The Journal of Visualized Experiments*

doi: 10.3791/61708

Majmundar A J, Wong WJ, Simon MC. 2010. Hypoxia-inducible factors and the response to

hypoxic stress. *Molecular Cell* **40**:294-309. doi:10.1016/j.molcel.2010.09.022

Malard F, Hervant F 1999. Oxygen supply and the adaptations of animals in groundwater.

Freshwater Biology **41**:1-30. https://doi.org/10.1046/j.1365-2427.1999.00379.x

Mandic M, Sloman KA, Richards JG. 2009. Escaping to the surface: A phylogenetically

independent analysis of hypoxia-induced respiratory behaviors in Sculpins. *Physiology*

Biochemistry Zoology. **82**, 730-738. doi: 10.1086/605932

Mandic M., Joyce, W., Perry, S. F. 2021. The evolutionary and physiological significance of

the Hif pathway in teleost fishes. *Journal of Experimental Biology* **224**, jeb231936.

Doi:10:1242/jeb.231936

Martinez ML, Chapman LJ, Grady JM, Rees BB. 2004. Interdemic variation in heamatocrit

and lactate dehydrogenase in the African cyprinid *Barbus neumayeri. Journal of Fish Biology.*

65:1056-1069.

Medley JK, Person J, Peuß R, Olsen L, Xiong S, Rohner N. 2020. Untargeted metabolomics

of the cavefish *Astyanax mexicanus* reveals the basis of metabolic strategies in adaptation to

extreme conditions**.** *BioRxiv* DOI: https://doi.org/10.1101/2020.10.27.358077

Mitchell RW, Russel WH, Elliott WR. 1977. *Mexican eyeless characin fishes, genus*

Astyanax: environment, distribution and evolution. (Lubbock: Texas Tech Press).

Moore C, Richens J., Hough Y, Ucanok D, Malla S, Sang F, Chen Y, Elworthy S.

Wilkinson RN, Gering M. 2018. Gfi1aa and Gfi1b set the pace for primitive erythroblast

differentiation from hemangioblasts in the zebrafish embryo. *Blood A*dvances **2**: 2589-2606. doi:

10.1182/bloodadvances.2018020156

Moran D, Softley R, Warrant E J. 2014**.** Eyeless Mexican cavefish save energy by eliminating

the circadian rhythm in metabolism**.** *Public Library of Science ONE* **9**:

e107877.https://doi.org/10.1371/journal.pone.0107877

Moran D, Softley R. Warrant EJ. 2015. The energetic cost of vision and the evolution of

eyeless Mexican cavefish. *Scientific Advances* **1**:e1500363. doi:10.1126/sciadv.1500363

Moriyama A, Inohaya K, Maruyama K, Kudo A. 2010. Bef medaka mutant reveals the

essential role of c-myb in both primitive and definitive hematopoiesis. *Developmental Biology*

345:133-143. doi: 10.1016/j.ydbio.2010.06.031

Möröy T, Vassen L, Wilkes B, Khandanpour C. 2015. From cytopenia to leukemia: the role

of Gfi1 and Gfi1b in blood formation. *Blood* **126**:2561-2569. doi:10.1182/blood-2015-06-

655043

Ornelas-García P, Pajares S, Sosa-Jiménez VM, Rétaux S, Miranda-Gamboa RA. 2018.

Microbiome differences between river-dwelling and cave-adapted populations of the fish

Astyanax mexicanus (De Filippi, 1853). *Peer Journal* **6**: e5906-5926. doi: 10.7717/peerj.5906

Paik E J, Zon LI. 2010. Hematopoietic development in the zebrafish. *International Journal of*

Developmental Biology **54:**1127-1137. doi: 10.1387/ijdb.093042ep.

Patterson LJ, Gering M, Eckfeldt CE, Green AR, Verfaillie CM, Ekker SC, Patient R.

2007. The transcription factors Scl and Lmo2 act together during development of the

hemangioblast in zebrafish. *Blood* **109**:2389-2398. doi:10.1182/blood-2006-02-003087

Pelster B, Burggren WW. 1996. Disruption of hemoglobin oxygen transport does not impact

oxygen-dependent physiological processes in developing embryos of zebra fish (*Danio rerio*).

Circulation Research. **7**:358-362. doi:10.1161/01.res.79.2.358

Pelster B, Egg M. 2018. Hypoxia-inducible transcription factors in fish: expression, function

and interconnection with the circadian clock. *Journal of Experimental Biology*. **221**:jeb163709.

doi: 10.1242/jeb.163709

Peuß R, Box AC, Chen S, Wang, Y, Tsuchiya D, Persons JL, Kenzior A, Maldonado E,

Krishnan J, Scharsack JP, Slaughter B, Rohner N. 2020. Adaptation to low parasite

abundance affects immune investment and immunopathological responses of cavefish. *Nature*

Ecology Evolution **4,** 1416–1430 https://doi.org/10.1038/s41559-020-1234-2

Rashid I, Nagpure NS, Srivastava P, Kumar R, Pathak AK, Singh M, Kushwaha B. 2017.

HRGFish: A database of hypoxia responsive genes in fishes. *Scientific Reports* **7**:42346. Doi:

10.1038/srep42346.

Ren X, Hamilton N, Müller F, Yamamoto Y. 2018. Cellular rearrangement of the prechordal

plate contributes to eye degeneration in the cavefish. *Developmental Biol*ogy **441**:221-234. doi:

10.1016/j.ydbio.2018.07.017

Rhodes J, Hagen A, Hsu K, Deng M, Ting XL, Look T, Kanki JP. 2005**.** Interplay of Pu.1

and gata1 determines myelo-erythroid progenitor cell fate in zebrafish. *Developmental. Cell*

8:97-108**.** doi: 10.1016/j.devcel.2004.11.014

Rohner N, Jarosz DF, Kowalko JE, Yoshizawa M, Jeffery WR, Borowsky RL, Lindquist S,

Tabin CJ. 2013. Cryptic variation in morphological evolution: HSP90 as a capacitor for loss of

eyes in cavefish. *Science* **342**:1372-1375. doi:10.1126/science.1240276.

Rombough R, Drader H. 2009. Hemoglobin enhances oxygen uptake in larval zebrafish (*Danio*

rerio) but only under conditions of extreme hypoxia. *Journal of Experimental Biology* **212**:778-

784. doi: 10.1242/jeb.026575

Rutjes HA, Nieveen MC, Weber RE, Witte F, Van den Thillart GEEJM. 2007. Multiple

strategies of Lake Victoria cichlids to cope with lifelong hypoxia include hemoglobin switching.

American Journal of Physiology. Regulatory, Integrative and Comparative. Physiology **293**:

R1376-R1383. doi: 10.1152/ajpregu.00536.2006

Sears CR, Boggs TE, Gross JB. 2020. Dark-rearing uncovers novel gene expression patterns in

an obligate cave-dwelling fish. *Journal of Experimental Zoology Part B Molecular Development*

Evolution. **334**:518–529.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.

*Nature Meth*ods **9**:676-682. 10.1038/nmeth.2019

Semenza GL, Wang GL. 1992. A nuclear factor induced by hypoxia via de novo protein

synthesis binds to the human erythropoietin gene enhancer at a site required for transcriptional

activation. *Molecular Cell Biology* **12**:5447-54. doi: 10.1128/mcb.12.12.5447

Singh D, Arora R, Kaur P, Singh B, Mannan R, Arora S. 2017. Overexpression of hypoxia-

inducible factor and metabolic pathways: possible targets of cancer. *Cell Bioscience.* **7**: 62. doi:

10.1186/s13578-017-0190-2

Stincone A, Prigione A, Cramer T, Wamelink MMC, Campbell K, Cheung E, Olin-

Sandoval V, Grüning N-M, Krüger A, Alam MT, Keller MA, Breitenbach M, Brindle K,

Rabinowitz JD, Ralser M. 2015. The return of metabolism: biochemistry and physiology of the

pentose phosphate pathway. *Biological Reviews of the Cambridge Philosophical Society* **90**:927-

963. doi: 10.1111/brv.12140.

Sollid J, De Angelis P, Gundersen K, Nilsson GE. 2003. Hypoxia induces adaptive and

reversible gross morphological changes in crucian carp gills. *Journal of Experimental Biology*

206:3667-3673. doi: 10.1242/jeb.00594

Torres-Paz J, Leclercq J, Rétaux S. 2019. Maternally regulated gastrulation as a source of

variation contributing to cavefish forebrain evolution. *eLIFE* **8**:e50160. doi:

https://doi.org/10.7554/eLife.50160

Timmerman CM, Chapman LJ 2004. Behavioral and physiological compensation for chronic

hypoxia in the sailfin molly (*Poecilia latipinna*). *Physiological Biochemical Zoology* **7**:601-610.

doi: 10.1086/421754

Tinevez J-Y, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek

SY, Shorte SL, Eliceiri KW. 2017. TrackMate: An open and extensible platform for single-

particle tracking. *Methods* **115**:80-90. doi:10.1016/j.ymeth.2016.09.016.

Torres JJ, Grigsby MD, Clarke ME. 2012. Aerobic and anaerobic metabolism in oxygen

minimum layer fishes: the role of alcohol dehydrogenase. *Journal of Experimental Biology*

215:1905-1914. doi:10.1242/jeb.060236

Varatharasan N, Croll RP, Franz-Odendaal T. 2009. Taste bud development and patterning

in sighted and blind morphs of *Astyanax mexicanus*. *Developmental Dynamics* **238**:3056-3064.

doi: 10.1002/dvdy.22144

Warga RM, Kane DA, Ho RK. 2009. Fate mapping embryonic blood in zebrafish: multi- and

unipotential lineages are segregated at gastrulation. *Developmental Cell* **16**:744-755. doi:

10.1016/j.devcel.2009.04.007

Wheaton WW, Chandel NS. 2011. Hypoxia. 2. Hypoxia regulates cellular metabolism.

American Journal of Physiology-Cell Physiology **300**:C385-C393.

doi:10.1152/ajpcell.00485.2010

Wei W, Wen L, Huang P, Zhang Z, Chen,Y, Xiao A, Huang, H, Zhu Z, Zhang B, Lin S.

2008. GFi1.1 regulates hematopoietic lineage differentiation during zebrafish embryogenesis.

Cell Research **18**:677-685. doi: 10.1038/cr.2008.60

Weinstein BM, Schier AF, Abdelilah S, Malicki J, Solnica-Krezel L, Stemple, D, Stainier,

DY, Zwartkruis F, Driever W, Fishman MC. 1996. Hematopoietic mutations in the zebrafish.

Development **123:**303-309.

Wells RMG. 2009. Blood-gas transport and hemoglobin function: adaptations for functional and

environmental hypoxia. In *Fish Physiology, Vol. 27* (ed. J.G. Richards, A.P. Farrell and C.J.

Brauner). pp. 255–299. New York: Elsevier.

Wong-Riley, M. 2010. Energy metabolism of the visual system. *Eye Brain* 2: 99-116.

Xiong S, Krishnan J, Peuβ R, Rohner N. 2018. Early adipogenesis contributes to excess fat

accumulation in cave populations of *Astyanax mexicanus*. *Dev. Biol*. **441**, 297-304. doi:

10.1016/j.ydbio.2018.06.003

Yamamoto Y, Stock DW, Jeffery WR. 2004. Hedgehog signalling controls eye degeneration in

blind cavefish. *Nature* **431**: 844-847. doi:10.1038/nature02864.

Yoshizawa M, Jeffery WR, van Netten S, McHenry MJ. 2014. The sensitivity of lateral line

receptors and their role in the behavior of Mexican blind cavefish (*Astyanax mexicanus). Journal

of Experimental Biology* **217**, 886-895. doi: 10.1242/jeb.094599

Figures and Legends

Figure 1. Cavefish have more erythrocytes than surface fish. (A) At 36 hpf blood

circulates from the heart (H) through the lateral dorsal aorta (LDA), the dorsal aorta (DA) and

caudal artery (CA) (blood flow shown in red), and back to the heart through the caudal vein

(CV), the posterior cardinal vein (PCV), and the Duct of Cuvier (DC) (blood flow shown in

blue). Scale bar is 200 μm. (B-D). Video analysis of circulating blood cells (arrowheads) in a

region of interest (ROI, boxes) in the DC of surface fish (SF) and cavefish (CF) larvae at 36 hpf

visualized by subtraction of two video frames 0.15 seconds apart (C, D) and quantified using the

plugin TrackMate (ImageJ) (D, purple dots). E. o-dianisidine staining of red blood cells in the

DC of surface fish (SF) and cavefish (CF) larvae at 36 hpf. Boxes: erythrocyte quantification

regions. Scale bar in E is 100μm, B-E are the same magnifications. (F, G). Quantification of

blood cells in the DC by video analysis (F) and o-dianisidine staining. Box plots show the

median, quartiles, min-max values and outliers (dots). Asterisks: p < 0.05, N = 72 (in F) and p <

0.0001, N = 36 (in G). Statistics by Wilcoxon/Kruskal-Wallis Rank Sums test.

Figure 2. Expression of hematopoietic genes during surface fish and cavefish

development. (A-C). qPCR quantification of *hhb2, hbbe2,* and *gfi1aa* transcripts in surface fish

(SF) and cavefish (CF) larvae at 24 hpf and 60 hpf. Error bars: range of fold change. Asterisks:

(A) N = 8. p = 0.0045 for SF vs CF at 24 hpf and p = 0.0254 for SF vs CF at 60 hpf; (B) N = 4. P

= 0.0271 for SF vs CF at 60 hpf.; (C) N = 3. P = 0.0413 for SF vs CF at 24 hpf. Statistics by

two-way ANOVA followed by Student's t test. (D-G). *In situ* hybridizations showing *hbb2*

staining at 24, 36, and 60 hpf, and (D-F) and *gfi1aa* staining (G) at 24 hpf in CF and SF. AGM:

aorta-gonad-mesonephros. PBI: posterior blood island. Scale bars are 200 μm in each frame.

Magnifications are the same in SF and CF.

Figure 3. Hematopoietic domains in 14 hpf surface fish and cavefish embryos. (A). *In*

situ hybridizations showing expression of the *gfi1aa* (top frames) and *hbb2* (bottom frames)

genes in the anterior lateral mesoderm (ALM, arrowheads) and posterior lateral mesoderm

(PLM, arrows) in surface fish (SF) and cavefish (CF). (B) Expression of the macrophage marker

gene *lcp1* in SF and CF embryos. (C). Expression of the hematopoietic marker gene *lmo2* in the

ALM and PLM of CF and SF embryos, and expansion of expression in CF embryos. Scale bar

in A is 200μm; magnification is the same in all frames.

Figure 4. Maternal control of increased blood cells in cavefish determined by reciprocal

hybridizations. Video imaging in the Duct of Cuvier and blood cell quantification at 34 hpf in

the F1 progeny of a (A) surface fish (SF) X SF control cross, (B) a SF female X cavefish (CF)

male cross, (C) a CF) female X SF male cross, and (D) a CF X CF control cross. (A-D) Top

row. Representative images of blood cell number aligned with box plots below. Scale Bar is

100μm; magnifications are the same in each frame. Bottom row. Boxplots of blood cell numbers

showing medians, quartiles, min-max values, and outliers (dots). Asterisks: $p<0.05$. ns: not

significant. N=20 for each box plot. Statistics by Wilcoxon/Kruskal-Wallis Rank Sums test

followed by Wilcoxon for each pair.

Figure 5. Differential effects of hemolytic anemia on surface fish and cavefish development. (A).

In situ hybridization with the *hbb2* gene marker in 40 hpf larvae treated with different

phenylhydrazine (PHZ) concentrations. Scale bar is 200 μm. Magnification is the same in all

frames. (B). Larval tail length (top) and erythrocyte number in the Duct of Cuvier (DC) (bottom)

in SF and CF embryos as a function of PHZ concentration. Asterisks in B (top) $p < 0.0001$. N =

23 for SF 0, 2 mg L^{-1} and CF 3 mg L^{-1}. N = 24 for SF 1 and 3 mg L^{-1} and CF 0, 1, and 2 mg L^{-1}.

Asterisks in B (bottom) consecutively from left to right: $p = 0.0197$, $p < 0.0001$, $p = 0.0002$, $p <$

0.0001, and p 0.0102. $N = 20$ for SF 0, 1, 2 and CF 3 mg L^{-1}. $N = 10$ for SF 3 mg L^{-1}. $N = 19$

for CF 0 mg L^{-1}. $N = 21$ for CF 1 mg L^{-1}. $N = 22$ CF 2 mg L^{-1}. Statistics by Wilcoxon/Kruskal-

Wallis Rank Sums test followed by Wilcoxon for each pair. (C). Axial defects induced by

hemolytic anemia in SF and CF embryo. Scale bars are 200 μm; magnifications are the same in

top four frames and bottom two frames. Data from B and C were obtained from the same

embryos. (D). Effects of a 1 hr. treatment with 5 mg/L PZH on post-anal tail growth (top),

erythrocyte number measured in the DC (middle), and axial development (bottom). Scale bar is

μm; magnification is the same in both frames. Asterisk (top): $p = 0.0115$, $N = 20$. Statistics by

Student's t test. Asterisk (middle): $p = 0.0001$, $N = 10$. Statistics by Wilcoxon/Kruskal-Wallis

Rank Sums test followed by Wilcoxon for each pair. Box plots in C and D show whiskers

(=5%), mean (line) and outlier (dot). SF: surface fish. CF: cavefish. Box plots show the median,

quartiles, min-max values and outliers (dots).

Figure 6. Effects of laboratory induced hypoxia on surface fish and cavefish growth. (A).

Image of a 36 hpf surface fish larval showing post-anal tail length. Scale Bar is 200 μM. (B).

Relative post-anal tail growth (top) and erythrocyte numbers (bottom) of surface fish and

cavefish over an 18 hr. period under normoxic and hypoxic conditions. Top frame asterisks from

left to right: p < 0.0001, p = 0.002; p < 0.0001, and p = 0.0021. N = 23 for SF-N and SF-H. N =

46 for CF-N. N = 47 for CF-H. Statistics by Wilcoxon/Kruskal-Wallis Rank Sums test followed

by Wilcoxon for each pair. Middle frame asterisks from left to right: p = 0.0116, and p = 0.0078.

N = 22 for SF-N. N = 21 for SF-H. N = 47 for CF-N, and N = 45 for CF-H. Statistics by one-

way ANOVA followed by Student's t-test. (C) Difference in eye diameters in surface fish and

cavefish after 18 hr. of hypoxia or normoxia. Bottom frame asterisks: p = 0.0002 and p < 0.0001.

N = 18 for SF-N. N = 14 for SF-H. N = 30 for CF-N. N = 28 for CF-H. Statistics by one-way

ANOVA followed by Student's t-test. Box plots show median, quartiles and min-max values.

Figure 7. Changes in expression of some of the key HIF pathway genes in surface fish

and cavefish exposed to hypoxia or normoxia. Bars indicate qPCR fold difference over surface

fish in normoxic condition. Error bars: range of fold change. Asterisks from left to right: p <

0.0001, p < 0.0001, p = 0.0044, p = 0.0306, p = 0.453, p = 0.0155, p = 0.0083, p = 0.0306, p =

0.0002, and p = 0.0003. N = 6 for each determination. Statistics by one-way ANOVA followed

by Student's t-test (*igfbp1a*, *hif1ab, hif1a-like*, *hif1a-like2*, *hk1*, *g6pd*, *ldha*, *pdk1*) or with

unequal variance by Wilcoxon/Kruskal-Wallis Rank Sums test followed by Wilcoxon for each

pair (*hif1aa*).

Tables

Table 1. Penetrance of abnormalities in phenylhydrazine-treated surface fish and cavefish
embryos.
990

991
992

Morph	PHZ concentration	% edema	% defective notochord
SF	0	0	0
CF	0	0	0
SF	1	4.1	87.5
CF	1	0	41.7
SF	2	75.0	100
CF	2	8.3	87.5
SF	3	95.8	100
CF	3	25.0	100

1002   PHZ: phenylhydrazine. SF: surface fish. CF: cavefish. N = 24 for each morph treatment.
1003
1004
1005   Table 2: Primer sequences used for gene expression analysis with qPCR

1006

rpl13a	GeneID: 103025160	ENSAMXG00000033532	caagtactgctgggccacaaag	aggaaagccaggtacttcaatttgtt
gfi1aa	GeneID: 103029320	ENSAMXG00000006669	agtgtgtgtgatcgaccttcaga	ggacattcttcattgtctggtgacg
hbbe2		ENSAMXG00005017210*	taaatccctctgcagggctctgat	cctgatcacctccggattagccataata

hbb2	GeneID: 111196758	ENSAMXG00000031275	gctcacggtgtagttgttctc	ggatccacgtgcagtttctc
gnb1b	GeneID: 103023031	ENSAMXG00000040710	ctctgctaaactgtgggatgtg	ccgttagggaagaaacagatgg
hif1aa	GeneID: 103022448	ENSAMXG00000039550	cagcaccaacacacacactcaa	gtcactgaccaccagtcctaca
hif1ab	GeneID: 103033873	ENSAMXG00000019342	gcatgggccttacacagttt	gcaccagcatttccctcatt
hif1alike	GeneID: 103027586	ENSAMXG00000008564	tgcctcacctgcttctaactct	agctgtattctcctctggcttga
hif1alike2	GeneID: 103041845	ENSAMXG00000007272	cattctaagttccagcccatcc	cattggctgcaccatctctc
hk1	GeneID: 103028521	ENSAMXG00000012670	ctcaatcggctgaaggacaacaa	agccgtcgagaatactgtggat
g6pd	GeneID: 103035433	ENSAMXG00000017509	tcctactctgtggtggttgtt	gagacggtctgcttcagtatct
ldha	GeneID: 103047177	ENSAMXG00000032467	tgtggtgtccaacccagttgata	agcgagctgagtccaagttagt
pdk1	GeneID: 103033744	ENSAMXG00000039808	tcctcaaccagcacactcttct	agtgacacgacagtgaggatcaa

igfbp1a	GeneID: 103033920	ENSAMXG00000009512	cccaacagaagctggaagataag	ctgcccatccagagttgattc

1007

1008   *blasts to the cavefish genome but also cloned from surface fish in this study

1009

1010   **Videos**

1019

1020

1028   Videos 1 and 2. Blood cells flowing from the cardinal vein through the Duct of Cuvier to the

1029   heart in (1) surface fish and (2) cavefish