Integration of immunome with disease gene network reveals pleiotropy and novel drug repurposing targets

Abhinandan Devaprasad[1], Timothy RDJ Radstake[1], Aridaman Pandit[1*]

[1] Center for Translation Immunology, University Medical Center Utrecht, Utrecht, The Netherlands

*Correspondence: Aridaman Pandit: a.pandit@umcutrecht.nl

Abstract

Immune system is crucial for the development and progression of immune-mediated and non-immune mediated complex diseases. Studies have shown that multiple complex diseases are associated with several immunologically relevant genes. Despite such growing evidence, the effect of disease associated genes on immune functions has not been well explored. Here, we curated the largest *immunome* (transcriptome profiles of 40 different immune cells) and integrated it with disease gene networks and drug-gene database, to generate a Disease-gene IMmune cell Expression network (DIME). We used the DIME network to: (1) study 13,510 genes and identify disease associated genes and immune cells for >15,000 complex diseases; (2) study pleiotropy between various phenotypically distinct rheumatic and other non-rheumatic diseases; and (3) identify novel targets for drug repurposing and discovery. We implemented DIME as a tool (https://bitbucket.org/systemsimmunology/dime) that allows users to explore disease-immune-cell associations and disease drug networks to pave way for future (pre-) clinical research.

Introduction

The genetic and epigenetic heterogeneity has been known to play a major role in the development and progression of complex diseases. The past two decades have seen a major surge in studies that characterize genes and loci associated with disease. The use of high-throughput omics technology and functional screenings have boosted our knowledge about genetic, epigenetic and metabolic factors underlying complex diseases[1]. As a result of these genetic and epigenetic screenings, we now know that the majority of complex diseases and genes/loci have a many-to-many relationship meaning that a complex disease is linked to many different genes and a gene/loci is associated with many different genes[2].

Large high-throughput screening studies have typically used bulk tissue or whole blood to study disease associated genes (DAGs). However, the expression of each gene is known to vary between tissues and cell types[3,4]. Thus, bulk tissue- or blood-based studies on DAGs do not consider the role played by different cells and tissues in the disease biology. To improve the understanding and molecular basis of complex diseases, a large number of research groups and consortiums have started to functionally identify disease associated cells (DACs) or tissue types[3–7]. The Genotype-Tissue Expression (GTEx) is one such valuable project, which maps gene expression profiles of 54 different human tissue types and the corresponding expression quantitative trait loci (eQTLs)[5–7]. Furthermore, the growth of single cell technologies have advanced our understanding of DACs and have helped in identifying cell types associated with complex diseases including cancer[8], Alzheimer's[9], rheumatoid arthritis[10], among others.

The immune system is known to play a key role in the development and progression of immune-mediated as well as non-immune mediated chronic diseases. A large number of association and functional studies have shown that multiple DAGs are expressed in immune cells and perturbing these DAGs can modulate immune cell functions[11]. However, very few studies have explored the impact of DAGs on specific cell types and even fewer on immune cells, many of which focus on limited number of cell subsets[12–16]. Recently Schmiedel *et al.* studied the effect of genetic variants on the expression of genes in 13 different immune cell types[17]. However, this study largely focused on the analysis of genetic variants and their impact on a total of 13 immune cell types: monocytes (classical and non-classical), NK cells, naïve B-cells and nine sub-populations of T-cells.

In this study, we mapped the largest available and expert curated disease-gene network (from the DisGeNet curated from 16 different databases) on the largest *immunome* data curated by us comprising gene expression profiles of 40 different immune cell types. We then quantified the effects of 13,510 DAGs on the *immunome*, to identify DACs for 15,367 different diseases in the DisGeNet. Using the DACs and the DAGs, we constructed the Disease-gene IMmune cell Expression (DIME) network. We use the DIME network to: (1) study the underlying cell-specific mechanisms of complex diseases; (2) identify cell-specific targets for complex disease; (3) identify networks of genes and cells that are commonly associated with different pairs of diseases; and (4) predict drug repurposing targets towards identified disease mechanisms shared between different diseases. We further built a user-friendly shinyapp called DIME (https://bitbucket.org/systemsimmunology/dime), which can be used to identify DACs and construct DIME network for: (1) diseases from the DisGeNet, (2) diseases from the EBI genome wide association study (GWAS) catalogue, or (3) custom set of genes defined by the user.

Methods

Transcriptome data - *Immunome*

The transcriptome data consists of RNA-sequencing datasets of 40 different immune cell types curated using 316 samples from a total of 27 publicly available datasets (see **Supplementary Table 1** for list of GEO datasets and samples used). The 40 different immune cells cover the entire hematopoietic stem cell differentiation tree comprising of 9 progenitors, 19 lymphoid, and 12 myeloid cell types. The samples used here were manually curated considering only the unstimulated (except for macrophages, that were monocyte derived) immune cells that were sorted using Fluorescence-activated cell sorting (FACS) and were isolated from either blood, bone marrow or cord blood from healthy donors. All the selected datasets were downloaded as FASTQ files using the fastq-dump tool from sratoolkit[18]. The "—split-files" option was given if the library type was paired end sequencing. FASTQ files were then aligned to reference genome (GRCH.Hg38.79) using STAR aligner[19]. The result is a SAM file which was then converted into a sorted BAM file using the samtools program[20]. These were then used to calculate the count of aligned reads using the HTSeq program[21] with the mode option "intersection non-empty". HTSeq was run

for all possible stranded mode options, the count file with the maximum counts was chosen as
the respective count file for the sample.

The data was then filtered by removing all genes that had less than 20 read counts in 95
percent of the samples using R programming. The filtered data was then lane normalized
using the "betweenLaneNormalization" function from the RUVSeq package[22]. The RUVr
method from RUVSeq was used to identify residual factors contributing to the batch effect.
The resulting filtered, batch corrected and normalized data had expression for 34,906 genes
that was void of any observable batch effect. We calculated counts per million (cpm) for the
filtered genes and used cpm as the gene expression measure. We then used the median gene
expression for each cell type for the rest of the analysis. This processed, batch corrected,
normalized and median representative data of 40 immune cells is referred to as the
immunome.

Disease gene network from DisGeNet

The full disease gene association network from DisGeNet[23] was downloaded from the
DisGeNet database (www.disgenet.org/downloads). All HLA associated genes was removed
from the network, this was done to ensure that bias towards myeloid cells and B cells are
removed, since the HLA genes are largely expressed by these cells. The resulting network
was further filtered to include only those genes that were present in the *immunome*. The final
network comprised of 15367 diseases and 13510 DAGs.

The DisGeNet consists of expert curated disease-gene interactions from 16 different
databases: UNIPROT, CGI, ClinGen, Genomics England, CTD, PsyGeNET, Orphanet, RGD,
MGD, CTD, Human Phenotype Ontology, Clinvar, GWAS catalogue, GWAS DB, LHDGN
and BeFree. The DisGeNet is the largest and most comprehensive disease-gene association
network available in the literature that was known to us. We also tested our methods on more
specific disease networks such as those from the EBI GWAS database.

Other disease gene networks - EBI GWAS data

In addition to the DisGeNet, we also used a refined GWAS based dataset from the EBI[24]. The
GWAS catalogue of Version 1.0, e89, was downloaded from the EBI website, which

contained information on the disease associated SNP for about 1900 diseases/traits. The

reported p-value of all the disease associated SNP in the catalogue was ≤ 0.05. The catalogue

also provided the corresponding mapped gene information for all the SNP which was used to

construct the disease to gene association network. We further filtered this network using the

same filtering criteria that was used for the DisGeNet. The EBI GWAS dataset was used to

infer SNP based disease cell associations.

Mapping disease gene network to *Immunome* data

For a given disease D and its DAG_D, we first extracted the corresponding *Immunome*

expression matrix. This expression matrix (X_D) comprised the gene expression of the DAG_D

across the 40 cells forms as the input data upon which further analysis was performed. Thus,

the dimension of each X_D was given as:

$$dim(X_D) = length(DAG_D) \times 40,$$

where, $length(DAG_D)$ is the number of DAGs in disease D and 40 corresponds to the number

of cell types in the *immunome* data.

Using NMF to cluster X_D into k classes

We used the NMF package[25] in R and applied the non-negative matrix factorization method

using Brunet's[26] algorithm to the expression matrix (X_D) to factor it into two matrices namely

W_D and H_D such that.

$$X_D \approx W_D\, H_D,$$

where, W_D and H_D are the basis and coefficient matrices computed by the NMF. The

dimensions of W_D and H_D are given as:

$$dim(W_D) = length(DAG_D) \times k,$$

$$dim(H_D) = k \times 40,$$

where, k is the number of classes/clusters that splits the data, such that it satisfies the above

NMF equation. The W_D matrix comprises of the weights of the DAGs across the k clusters (in

each column) and the H_D matrix comprises of the weights of the cells in the corresponding k

clusters (in each row). We used Brunet *et al.* method to identify the ideal k value using the

cophenetic correlation coefficient method[26].

Identifying the key DAG_D and DAC_D from W_D and H_D

The NMF algorithm clusters the data into k clusters such that, in each cluster 'i', where $i \in$

$(1, \ldots, k)$, the genes that have high values in w_D^i are constitutively expressed by the cells that

have high values in h_D^i. Where, w_D^i is the i^{th} column of W_D and h_D^i is the i^{th} row of H_D. For

each cluster i, we chose the cell-gene pairs that were in the top 25^{th} percentile range of their

corresponding w_D^i and h_D^i values. These cells and their corresponding gene pairs are regarded

as the key DAC_D and DAG_D respectively. The cell-gene pairs were extracted from all the

clusters and were compiled together. The resulting cell-gene pairs of all the clusters form the

edges of the **D**isease-gene to **IM**mune cell **E**xpression network, hereby referred as the **DIME**

network.

Identifying the key cluster

157   We then identified the largest weighted cluster among the k clusters identified by the NMF.

158   That is, the subset of genes and cells of X_D that can capture most of its expression pattern. We

159   did this by using the following approach.

160   Since X_D, W_D and H_D can be represented as below:

161
$$X_D \approx W_D H_D = \begin{bmatrix} | & | & | & | \\ w_D^1 & w_D^2 & \ldots & w_D^k \\ | & | & | & | \end{bmatrix} \begin{bmatrix} - & h_D^1 & - \\ - & h_D^2 & - \\ - & \vdots & - \\ - & h_D^k & - \end{bmatrix} = \sum_{i=1}^{k} w_D^i \, h_D^i.$$

163   We calculated the Frobenius norm of each $w_D^i h_D^i$ for all values of i. We then identified the

164   cluster (represented as c) for which $||w_D^i h_D^i||_F$ is the maximum. This can be mathematically

165   represented as:

166
$$c = \text{argmax}\left(||w_D^i h_D^i||_F\right); \; i \in \{1, \ldots, k\},$$

where, the c^{th} cluster represents that cluster which maximally captures/represents the

expression matrix X_D. We used the w_D^c as the scores for the DAG_D and h_D^c as the scores for the

DAC_D.

$$DAG_D \text{ score} = w_D^c,$$

$$DAC_D \text{ score} = h_D^c,$$

The scores were scaled between 0 and 1, with 1 representing the cell or gene with the highest

score.

Pleiotropic associations

To identify common mechanisms between two diseases, we looked at their overlapping cell-

gene connections in their corresponding DIME networks. Jaccards' index (JI) was used to

measure the overlap between the two diseases with Fisher's exact test (FET) used to obtain

confidence p-value for the given overlap. The overlapping genes were used to calculate JI and

statistical significance of overlap using FET. The pleiotropy based overlapping cell-gene

network between the two diseases is referred to as the DIME-pleiotropy network.

Integrating drug-gene network

The drug to gene target database from DGIdb was downloaded[27]. The data was filtered to

contain only the CHEMBL interactions and only those pertaining to the drugs approved by

the food and drug administration (FDA) of USA. This FDA approved drug to gene target

interaction serves as the drug-gene network in this study. To identify potential drug

candidates from the drug-gene network for disease D, we choose its key DAG_D as identified

in the previous step and extracted its corresponding target drugs from the drug-gene network.

This network between the drugs and the key DAG$_D$ is referred to as the DIME-drug network

for disease D. The DIME-drug network represents all the drugs that target the key DAG$_D$

identified in the given DIME network. To identify potential common acting drugs between

different diseases, we used their corresponding DIME-pleiotropy network and extracted the

drug-gene connections between the cell-gene and the drugs in drug-gene network. The drugs

identified using this approach represent those that act on the common mechanisms between

the two diseases and can be potential candidates for drug repurposing.

Rationale for restricting analysis to only diseases with ≥ 100 DAGs

The filtered disease-gene network of the DisGeNet comprised of DAGs for about 15,367 diseases. This could make the pleiotropy analysis between diseases extremely cumbersome, with the number of disease-to-disease comparisons reaching as large as 118,064,661. Additionally, to have sufficient number of DAC-DAG associations in the DIME-pleiotropy network and also have sufficient number of drug-target gene associations in the DIME-drug network, we used a smaller subset of diseases but with larger DAGs associated to them. We found by looking at the distribution of DAGs across the diseases, that the number of diseases with ≥ 100 DAGs were fewer in number, i.e., 613 diseases. This was sufficiently large DAC-DAG associations for the pleiotropy analysis and less cumbersome to analyze the comparisons (187,578). Hence, we choose those diseases with ≥ 100 DAGs. All analysis presented in this study have been performed on this disease subset.

DIME shinyapp

To construct the DIME and DIME-drug network for other diseases not mentioned in this study or using a custom gene input, we built a tool in R/Bioconductor called DIME that is available as a shiny app (https://bitbucket.org/systemsimmunology/dime). The app can be used to identify the key DAC, DAG and the DIME-drug network for the diseases from the DisGeNet, the EBI-GWAS catalogue or for custom set of genes from the user.

Results

The aim of this study is to identify disease associated cell types (DACs) based on the existing disease gene network, and to further identify disease associated gene (DAG) subsets that may perturb the associated immune cells. To achieve this, we integrated our *immunome* data with the disease network from DisGeNet and used the non-negative matrix factorization (NMF) method to identify those subsets of genes and cells that maximally represent the **D**isease gene **IM**mune cell **E**xpression (DIME) network, as illustrated in **Figure 1A**. The constructed immunome dataset comprises 40 different cell types and is the largest bulk RNASeq meta-dataset of the immune cell types known to us. The filtered disease-gene network from

DisGeNet consists of associations between 15,367 diseases and 13,510 DAGs. In this
massive disease network, numerous DAGs were found to be common between both
phenotypically similar and distinct diseases. We explore these common DAG patterns in
more detail in the next section.

Common DAGs of phenotypically distinct diseases

The 15,367 diseases in the DisGeNet belong to 29 different disease MeSH (Medical Subject
Headings) terms (**Figure 1B**). The MeSH-MeSH network in **Figure 1B** depicts the
connections between the different MeSH terms, where the thickness of the connections
represents the number of DAGs common between the different MeSH terms. The neoplasm
MeSH term was the most well connected disease category in the network. The highest
number of common DAGs (6,959 DAGs) between two different MeSH terms was observed
between neoplasm and digestive system diseases. Other top MeSH-MeSH connections that
had more than 5,000 common DAGs include those between neoplasm and that of the skin and
connective tissue, nervous system, congenital, endocrine system, and female urogenital
diseases and pregnancy complications. Thus, the MeSH-MeSH gene network revealed the
shared DAGs across phenotypically distinct diseases belonging to different MeSH terms.

We further studied the DAG patterns across MeSH terms and found that TP53 was
preferentially associated with diseases categorized under the neoplasm MeSH term (**Figure
1C**). Interestingly, TP53 was also associated with numerous diseases from various other
MeSH terms such as immune system diseases, nervous system diseases, and skin and
connective tissue diseases. Similarly, TNF was associated with numerous diseases from
various MeSH terms including immune system diseases (**Figure 1D**). APOE was largely
associated with nervous system diseases, and ACE was largely associated with cardiovascular
diseases (**Figure 1E** and **F**). TLR4 and CXCL8 were prevalently associated to several MeSH
terms (**Figure 1G** and **H**). The above-mentioned genes (**Figure 1C-1F**) are those that are
either the top represented (degree) genes across all diseases or for disease within specific
MeSH terms. More examples of DAGs associated with specific MeSH terms and their degree
can be found in the **Supplementary Figure 1**. Majority of the DAGs mapped to two or more
MeSH terms (**Figure 1I**) or diseases (**Figure 1J**) also hinting towards shared disease
mechanisms between phenotypically distinct diseases. These genes that are associated to
many diseases/MeSH terms (i.e. having high degree) were regarded as the hub genes of the

disease-gene network. We ranked the DAGs based on their degree and found TP53, TNF, VEGFA, BCL2, IL1B as the top 5 DAGs each being associated with >750 diseases (**Figure 1K** and **Supplementary Figure 1**).

Relevance of the immune system

We further evaluated the expression of the hub genes identified in the previous step, in the *immunome* data. The rationale was to assess if the hub genes representing the core of the disease-gene network were important to the immune system. Indeed, we found that many of the hub genes of the disease-gene network were expressed constitutively by either all immune cells or by some specific immune cells, as seen in **Figure 1K**. For example, the nervous system associated gene, APOE, important to many of neurological diseases such as Alzheimer's and Parkinson[28] were found to be expressed specifically by macrophages, as seen in **Figure 1L**. Studies have shown that genetic polymorphisms in APOE protein leads to defective clearance of the Aβ plaques by macrophages[29–31]. Thus, conferring macrophages as one of the key players of the disease. Such links between DAGs, to observing altered function in a cell type, questions the need to study DAGs by integrating cell-specific expression information.

Identifying disease associated cell types

We further identified such disease associated cells (DACs) based on the DAGs and report DACs for about 600 diseases in the disease-gene network (see Methods). The immune cells can be broadly categorized as myeloids, lymphoids and progenitors. Using the DAC profiles of about 600 diseases, we observed that in most diseases, DAGs do not associate with all immune cells but tend to associate with specific immune cells or with specific category of immune cells (**Supplementary Figure 2**). We observed from these DAC profiles that several phenotypically different diseases had similar DAC profiles. For example, diseases such as skin carcinoma, muscle degeneration, juvenile arthritis, epilepsy, etc. clustered together, showing progenitors as their top DACs. Similarly, peripheral T-cell lymphoma, celiac disease, atopic dermatitis, malignant glioma, basal cell carcinoma, etc. clustered together, with lymphoid cells as the top DACs. Interestingly, systemic lupus erythematosus (SLE), heart failure, myocardial infarction, colitis, type 1 diabetes, etc. cluster together showing

myeloid cells as their top DACs. Thus the DAC profiles like the DAG profiles showed that
phenotypically different diseases had similar DAC pattern.

We constructed the DIME network for different diseases individually, the DIME network
consists of the top DACs and their respective DAGs, (see methods). The DIME network
represents the key cell-gene mechanisms that can be drawn from the given set of DAGs. As
proof of concept, we present here the DIME network of lymphoid leukemia (**Figure 2A**), a
group of blood cancer that typically affects the lymphocytes. The DIME network revealed 4
DAC clusters for lymphoid leukemia. The key DAC cluster comprised of lymphoid
progenitor cells as well as myeloid progenitor cells. The key DAGs contributing to this
cluster included genes associated with hematopoiesis such as RPS14, HSP90AA1, MPO,
ETV6, ATF4 and TAL1. The other key DAC cluster comprised of all the subsets of T cells,
primarily the CD4$^+$ T cells. The key DAGs contributing to this cluster included ETS1,
CXCR4, IKZF1, ATM, LCK, and KMT2A. The pathway enrichment (**Supplementary**
Figure 4B) of these genes revealed pathways associated to TCR and BCR signaling and
PI3K/AKT signaling. Interestingly, these pathways have been shown to be important for
survival of cancer cells and are targets for anti-cancer drugs in acute and chronic lymphoid
leukemia[32]. Thus the DIME network of lymphoid leukemia revealed the key DAGs and
DACs implicated in the disease.

We then explored the DIME network of different kinds of rheumatic and/or fibrotic diseases,
such as systemic scleroderma (MeSH: skin disease), pulmonary fibrosis (MeSH: respiratory
tract disease) and SLE (MeSH: immune system disease). Moreover, we aimed to characterize
the diseases on the basis of the DAGs, DACs and the accompanying DIME networks. The
DIME network of systemic scleroderma, an autoimmune rheumatic condition with chronic
inflammation and fibrotic phenotype, revealed a complex relationship between their DAGs
and their DACs (**Figure 2B**). For systemic scleroderma, myeloid cells (neutrophils,
granulocytes, BDCA1$^+$CD14$^+$, and CD11c myeloid dendritic cells), and lymphoid cells (NK,
CD4$^+$ T regulatory, ILC3, and ILC2) were identified as key cluster of DACs (see methods).
The key DAGs contributing to this cluster included PTPRC, FOS, SRRM2, MSN, JUNB,
CXCR4, ITGB2, and TNFAIP3 genes. For systemic scleroderma, the other key DAC cluster
comprised myeloid cells (macrophages and pDCs) and myeloid progenitor cells. The key
DAGs contributing to this cluster included HSP90AB1, SPP1, HSP90AA1, MMP9, WNK1,
HIF1A and IRF8. The two systemic scleroderma associated DAC clusters together were
enriched in DAGs from interleukin signaling, TGF beta signaling, TLR signaling, ECM

organization and neutrophil degranulation pathways (**Supplementary Figure 3A**). Of which, evidently, TGF beta is known to play an essential role in the pathogenesis of fibrosis[33]. Likewise, the neutrophil degranulation pathway or otherwise known as NETosis, is the mechanism by which neutrophils exhibit defensive mechanisms to trap and kill foreign bodies[34]. And this pathway has been found to be implicated in other autoimmune diseases, and is now being currently investigated in clinical trials[35]. Interestingly, we found neutrophils and its NETosis associated genes to be in the top cluster of the DIME network for systemic scleroderma.

We then constructed the DIME network of another fibrotic disease, pulmonary fibrosis. The DIME network revealed 4 DAC clusters for pulmonary fibrosis (**Figure 2C**). Similar to systemic scleroderma, we found a DAC cluster comprising of T-cells and NK cells enriched in DAGs associated with TGF beta signaling (**Supplementary Figure 3B**). Interestingly, we also found that for pulmonary fibrosis neutrophils-granulocytes, macrophages, BDCA1$^+$CD14$^+$ cells, pDCs and myeloid progenitors were among the top DACs similar to that of systemic scleroderma, (**Supplementary Figure 3B**). We also found an enrichment of NLRP3 pathway genes, which are known to play a role in the collagen deposition mechanisms commonly dysregulated in fibrosis[36].

We then studied the DIME network of SLE, an autoimmune disease. The DIME network revealed 2 DAC clusters for SLE (**Figure 2D**). The key cluster comprised of myeloid cells like neutrophils-granulocytes, macrophage M1, BDCA1$^+$CD14$^+$ and monocytes as the DACs. The key DAGs contributing to this cluster included CD74, FOS, LYZ, SOD2 and HSP90AB1. The other cluster comprised of CD4$^+$ TEMRA, CD4$^+$ TEM, CD4$^+$ TCM and CD4$^+$ TH1 as their DACs. The key DAGs contributing to this cluster included B2M, IGHG3, IL7R, ETS1, RPS19, and TNFAIP3. The pathway enrichment for the SLE DIME network includes pathways that are heavily described in the literature for SLE, such as the neutrophil degranulation pathways or NETosis[37–41], interleukin 4 and interleukin 13 signaling[42,43], TLR signaling pathway[44], translocation of ZAP70 to immunological synapsis[45], and immune-regulatory interactions between lymphoid and non-lymphoid cells (**Supplementary Figure 4A**).

The DIME networks revealed immune system mediated mechanisms for the different diseases. As observed in the **Figure 2**, **Supplementary Figure 3** and **4**, the rheumatic and/or fibrotic diseases such as SLE and SSc, had overlap in specific cell-gene mechanisms

356 including those related to the neutrophil degranulation, degradation of ECM, interleukin and
357 TGF beta signaling pathways. These pathways and common mechanisms observed in the
358 DIME networks could serve as additional layers to understand the pathogenesis of these type
359 I interferon driven diseases[46]. We further explored such common mechanisms based on the
360 cell-gene relationships in the next section.

362 Pleiotropy based on DIME

Pleiotropy is when one gene affects two or more diseases[47]. Pleiotropy has been observed in
several studies for many different diseases based on gene mutation to phenotype
associations[48]. Unlike most pleiotropy studies performed in the past that looks at only the
common DAGs between two diseases, we used a different approach. Since the DIME
networks from the previous analysis (for example, between SSc and pulmonary fibrosis)
revealed several overlapping cell-gene connections, we extended this approach to look for
cell-gene connections that are common between the DIME networks of all possible pairs of
diseases. Hence, we define pleiotropy in this study as the cell-gene connections that are found
in one or more diseases. Using this approach, we constructed the pleiotropy network as
shown in **Figure 3A**. The pleiotropy network consists of diseases as nodes. The nodes are
connected if there exists a significant (JI $\geq$ 0.1and FET p-value $\leq$ 0.01) number of common
cell-gene connections between their corresponding DIME networks, (see methods). The
network shown in **Figure 3A** is trimmed to contain only those nodes with degree $\geq$ 2. The
node colors represent the MeSH term (as shown in **Figure 1B**) of the disease. The grey
colored nodes represent those diseases for which the MeSH term ontology was not available
from the DisGeNet. As seen from **Figure 3A**, several cancer related diseases (shown by
orange color for neoplasm MeSH term) cluster together. Same can be observed for the eye
related disease (shown by green color MeSH term) and the chemically induced disorders
(shown by pink and yellow color MeSH terms) seen in the bottom of the network **Figure 3A**.
These clusters together highlight the similar mechanisms (cell-gene connections) between
these diseases.

Within this pleiotropy network, we further looked at diseases that belong to different MeSH
terms and searched for common cell-gene mechanistic patterns between them. To do so, we
constructed the DIME-pleiotropy network of several pairs of phenotypically distinct
rheumatic diseases within the pleiotropy network. We present here some examples of the

DIME-pleiotropy network that had a JI similarity of ≥ 0.1. In the DIME-pleiotropy network

(**Figure 3B**) of Crohn's disease (MeSH: digestive system disease) and psoriasis (MeSH: skin

disease), we found cell-gene networks of lymphoid (CD4$^+$, NK and ILC's) and myeloid cells.

The pathway enrichment of the DAGs in this DIME-pleiotropy network revealed pathways

related to TCR signaling, interleukin signaling, neutrophil degranulation and regulation of

TLR (**Supplementary Figure 5A**). As observed in the DIME-pleiotropy network (**Figure

3B**), CD4$^+$ TH1 cells have been implicated in the pathogenesis of both Crohn's disease and

psoriasis[49,50]. Likewise, in the DIME-pleiotropy network (**Figure 3C**) of SSc (MeSH: skin

disease) and pulmonary fibrosis (MeSH: respiratory tract disease), we found cell-gene

networks between progenitors and pDCs, NK cells, CD4$^+$ TEMRA and CD4$^+$ T regulatory,

and other myeloid cells. The pathway enrichment of the DAGs in this DIME-pleiotropy

network revealed pathways (**Supplementary Figure 5B**) related to TGF beta receptor

signaling, interleukin, ECM and integrin cell surface interactions. Both of these being fibrotic

diseases, the involvement of TGF beta signaling and ECM is well represented in the DIME-

pleiotropy network and known to be implicated in literature[51–53]. Interestingly, we also

observed cell-gene network between macrophages and genes like TGF-B1, MMP9 and

TIMP1. Evidently, macrophages may exacerbate pulmonary fibrosis by TGF beta production

or cause ECM degradation via matrix metalloproteinase (MMP) activities[54]. The DIME-

pleiotropy network has captured the intricate network of these key DAGs and the cells

(macrophages) accurately.

Patients with rheumatoid arthritis (RA) have an increased risk of cancer due to the severe

regimen of disease modifying anti-rheumatic drugs[55]. However, RA patients seem to have

lower risk of colon cancer in comparison to the general population[55,56]. To explore the factors

responsible for the protective effect against colon cancer in RA patients, we constructed the

DIME-pleiotropy network (**Figure 3D**) of RA (MeSH: immune system) and colon carcinoma

(MeSH: neoplasm). We found cell-gene networks of CD4$^+$ T cells and many of the myeloid

cells. The pathway enrichment of the DAGs in this DIME-pleiotropy network revealed

pathways (**Supplementary Figure 5C**) related to TLR signaling, interleukin signaling,

neutrophil degranulation, ECM organization, FCER1 and EGFR signaling. Although clear

signatures of the protective effect were not observed, we did find the presence of PTGS1 and

PTGS2 (also known as COX1 and COX2) in the DIME-pleiotropy network of RA and colon

carcinoma. These genes are targets of the non-steroidal anti-inflammatory drugs (NSAIDs)

that are frequently taken by RA patients. Evidently, NSAIDs like aspirin have been shown to

confer protective affect against colorectal cancer even in lower doses[57]. Perhaps the missing

link in the protective effect of RA in colon cancer lies in the anti-inflammatory role of the

NSAIDs taken by the RA patients that target the pro-inflammatory mediators[58] such as

PTGS1 and PTGS2 in both diseases, thus conferring protection.

These DIME-pleiotropy analyses highlighted the common cell-gene relationships between

the different diseases (**Figure 3B-D**). Thus, making such pleiotropic relationships less

"ubiquitous"[59] but rather based on the similar cell-gene mechanism implicated between the

two diseases. We used this to further capture the plausible drug targets based on the immune

system mechanism as we identified with RA and colon cancer.

Immunome mediated drug repurposing

Using our immunome data, the DisGeNet and the NMF, we constructed the DIME network

and the DIME-pleiotropy network (as described before and as depicted in **Figure 4A**, part 1

and 2). We extended this approach to construct drug-gene network to identify drug targets

based on the DIME network (**Figure 4A**, part 3), which are referred to as the DIME-drug

network. We then used the pleiotropy method used before to then identify common drug

targets between the diseases (**Figure 4A**, part 4) based on the DIME-drug networks, which

forms the basis of the immunome mediated drug repurposing, (see methods). **Figure 4B**

shows an example of the DIME-drug network of Crohn's disease that represents the

connections between the top DAGs of Crohn's disease as the potential drug targets and their

associated drugs. The DIME-drug network of Crohn's disease identified some known drugs

like the corticosteroids (such as prednisone, methylprednisolone, hydrocortisone and

budesonide, targets of NR3C1) and the aminosalicylates (such as sulfasalazine and

mesalamine, targets of PTGS2 and ALOX5) that are current line of drugs used in treatment of

Crohn's disease[60]. In addition to the known drugs and drug targets, the DIME-drug network

of Crohn's disease also revealed some novel and potentially interesting drugs and drug

targets, such as lifitegrast that target the integrins, ITGB2 and ITGAL. This is particularly

interesting because integrin based therapies (such as natalizumab and vedolizumab) are

already in use for Crohn's disease[61]. Exploring other integrin based therapies for Crohn's

disease may be beneficial since both ITGAL and ITGB2 show in the top DAGs and are also

implicated in Crohn's disease[62,63].

As we discovered common cell-gene mechanisms in the DIME-pleiotropy network in the previous analysis (**Figure 3**), we extended this approach to discover plausible drugs and drug targets between the disease pair comparison to find new candidates for drug repurposing, we refer to these analysis as the pleiotropy-drug network analysis (**Figure 4A**, part 4). In addition to the drug targets revealed from the DIME-drug network of Crohn's disease, the pleiotropy drug network of Crohn's disease and psoriasis (**Figure 4C**), comprised some known drug targets such as IL6R for psoriasis and IL1B for Crohn's disease. The associated drug of IL6R, tocilizumab is known to be used in the treatment for psoriasis[64]. Interestingly, anti IL6 therapy has been shown to have promising clinical response for Crohn's disease as well[65]. Similarly, anti-IL1 therapies have also been explored for psoriasis and have shown beneficial clinical outcome[64]. Thus, DIME-network identifies established and novel potential targets for drug repurposing.

Studying the pleiotropy-drug network of fibrotic disease systemic scleroderma and pulmonary fibrosis (**Figure 4D**), we found the DNMT1 (a DNA methyl transferase enzyme) targeting drugs decitabine and azacitidine. Epigenetic modulation as a therapy has been explored as a treatment option in SSc[66]. Similarly, in the pleiotropy drug network of SSc and myocardial infarction (MeSH: Cardiovascular), **Figure 4E**, we found canakinumab an IL1B targeting drug. Anti-IL drugs have also been tested on patients with myocardial infarction and were shown to have fewer cardiovascular events than placebo[67].The pleiotropy drug network of rheumatoid arthritis and colon carcinoma was found to be larger (**Figure 4F**), due to the larger overlap between the two diseases, see **Figure 3D**. The network shows possibilities of using anti-inflammatory drugs for treatment or prevention of colon carcinoma as discussed before. With drugs that target genes such as IL2RG, TYK2, PTGS1, PTGS2, etc. that are widely used to treat anti-inflammatory or immune mediated diseases, these can be used as preventive care drugs or for use along with chemotherapy[68].

We also looked at the known drug targets that are present across the DIME networks of the 613 diseases analyzed by us. We found the top 5 drug targets to be BCL2 (**Figure 4B, C, E, and F**), PTGS2 (**Figure 4B, 4E-4F**), PIK3CD (**Figure 4D, 4E**), CXCR4 (**Figure 4B-4F**) and IL1B (**Figure 4B, C, E, and F**) that were implicated in the DIME networks of more than 200 out of the 613 diseases analyzed by us. The CXCR4 targetable by plerixafor is present in all of the pleiotropy-drug networks shown here (**Figure 4B-4F**). Plerixafor is a drug intended for use in cancer after stem cell transplantation to initiate migration of stem cells in the bloodstream[69]. This drug is now in clinical trials (NCT01413100) to be evaluated for use

after autologous transplant in patients with scleroderma[70]. Such trials may potentially be
extended to other immune diseases like psoriasis, Crohn's disease, rheumatoid arthritis, and
potentially for a variety of other phenotypically distinct rheumatic diseases that are driven by
CXCR4 mediated dysregulation of immune system. Many such potential drug repurposing
targets (enlisted in **Supplementary Table 2**) can be similarly evaluated in future studies and
trials.

Discussion

Despite decades of experimental data on the understanding of the molecular mechanism of
diseases, we know little about the perturbations in the niche cell compartments that are
specific to the disease. To address this gap several efforts at the tissue[5–7] and immune cell[16,17]
level have been performed to identify these disease specific compartments. However,
previous studies have concentrated at whole tissues, not distinguishing different immune cell
subsets or, in contrast, focused on a few immune cell subsets thereby likely missing the
complex molecular network underpinning immune mediated diseases. Additionally, it is
important to understand which of the gene subsets contribute to a mechanism within the
different cell populations. A disease may have perturbations occurring at a single cell type or
at multiple cell types with completely different or similar genes being involved. To truly
understand a disease, it is essential to capture these cell-gene networks as holistically as
possible.

To address the above-mentioned gap, we used the systems immunology approach of
dissecting diseases using the *immunome* comprising of 40 immune cells, the vast literature on
the available disease network and computational methods to compute and construct DIME
networks. The unique integration of these parts resulted in the novel mechanisms being
captured by our method. In this report, we highlight some of the known mechanisms we
capture from the DIME networks. For example, the role of NETosis and granulocytes in
many of the immune diseases, the role of TGF beta signaling in fibrotic diseases, the role of
BCR signaling and PI3K/AKT pathways in cancer, etc. We further outlined methods to
capture pleiotropy between diseases using the combination of cell-gene commonalities
between the DIME networks of the diseases, to ensure the robust capture of common
mechanism between diseases. We further extend this approach to identify immune
mechanism based drug targets that provides additional support and rationale for drug

repurposing. For example, we found from the DIME drug network that several specific

interleukins are involved in Crohn's disease, psoriasis and myocardial infarction. Given the

success of anti-IL therapies in many of the immune related diseases, our results indicate that

the anti-IL therapies might also be a promising option for these immune related diseases.

Likewise, our results indicate that using NSAIDs might be a promising option for cancer

prevention and treatment. However, additional functional experiments and extensive clinical

trials have to be done to support this approach.

The DACs and DAGs of the DIME network make it more robust to pinpoint which

mechanism and in which cell type. And this makes it a useful tool to dissect diseases. Hence

we built a shiny app to identify and construct DIME networks for all the diseases in the

DisGeNet, the GWAS network and also for user defined set of genes. The tool also identifies

potential drug targets based on the DIME networks. A caveat is that we focused on 40

different immune cells in this study, certain diseases may have manifestations that may or

may not be perturbed in the immune system or the progenitor cells that we have looked at.

Such diseases should be analyzed with additional data to identify the DACs and the

underlying mechanism. We have built the tool to accommodate and plugin such futuristic or

existing data consisting of gene expression information (coming from single cell or bulk) on

the different cell types apart from those found in the immune system. We believe that this tool

will aid scientist to increase the understanding of disease pathology and facilitate drug

development by better determining drug targets, thereby mitigating risk of failure in late

clinical development.

REFERENCES

1. Sun, Y. V & Hu, Y.-J. Integrative Analysis of Multi-omics Data for Discovery and
Functional Studies of Complex Human Diseases. *Adv. Genet.* **93**, 147–90 (2016).

2. Li, Y. & Agarwal, P. A Pathway-Based View of Human Diseases and Disease
Relationships. *PLoS One* **4**, e4346 (2009).

3. Camp, J. G., Platt, R. & Treutlein, B. Mapping human cell phenotypes to genotypes
with single-cell genomics. *Science (80-.).* **365**, 1401–1405 (2019).

4. Rozenblatt-Rosen, O., Stubbington, M. J. T., Regev, A. & Teichmann, S. A. The
Human Cell Atlas: from vision to reality. *Nature* **550**, 451–453 (2017).

5. Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**,
580–585 (2013).

6.   Ardlie, K. G. *et al.* The Genotype-Tissue Expression (GTEx) pilot analysis:
Multitissue gene regulation in humans. *Science (80-.).* **348**, 648–660 (2015).

7.   GTEx Consortium *et al.* Genetic effects on gene expression across human tissues.
Nature **550**, 204–213 (2017).

8.   Tirosh, I. & Suvà, M. L. Deciphering Human Tumor Biology by Single-Cell
Expression Profiling. *Annu. Rev. Cancer Biol.* **3**, 151–166 (2019).

9.   Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting
Development of Alzheimer's Disease. *Cell* **169**, 1276-1290.e17 (2017).

10.   Kuo, D. *et al.* HBEGF+ macrophages in rheumatoid arthritis induce fibroblast
invasiveness. *Sci. Transl. Med.* **11**, eaau8587 (2019).

11.   Knight, J. C. Genomic modulators of the immune response. *Trends Genet.* **29**, 74–83
(2013).

12.   Raj, T. *et al.* Polarization of the Effects of Autoimmune and Neurodegenerative Risk
Alleles in Leukocytes. *Science (80-.).* **344**, 519–523 (2014).

13.   Quach, H. *et al.* Genetic Adaptation and Neandertal Admixture Shaped the Immune
System of Human Populations. *Cell* **167**, 643-656.e17 (2016).

14.   Lee, M. N. *et al.* Common Genetic Variants Modulate Pathogen-Sensing Responses in
Human Dendritic Cells. *Science (80-.).* **343**, 1246980–1246980 (2014).

15.   Fairfax, B. P. *et al.* Innate Immune Activity Conditions the Effect of Regulatory
Variants upon Monocyte Gene Expression. *Science (80-.).* **343**, 1246949–1246949
(2014).

16.   Chen, L. *et al.* Genetic Drivers of Epigenetic and Transcriptional Variation in Human
Immune Cells. *Cell* **167**, 1398-1414.e24 (2016).

17.   Schmiedel, B. J. *et al.* Impact of Genetic Polymorphisms on Human Immune Cell
Gene Expression. *Cell* **175**, 1701-1715.e16 (2018).

18.   SRA-Tools - NCBI. http://ncbi.github.io/sra-tools/.

19.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

20.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

21.   Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

22.   Risso, D., Ngai, J., Speed, T. P. & Dudoit, S. Normalization of RNA-seq data using
factor analysis of control genes or samples. *Nat. Biotechnol.* **32**, 896–902 (2014).

23.   Piñero, J. *et al.* DisGeNET: a comprehensive platform integrating information on
human disease-associated genes and variants. *Nucleic Acids Res.* **45**, D833–D839
(2017).

24.   Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide
association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* **47**,
D1005–D1012 (2019).

25.   Gaujoux, R. & Seoighe, C. A flexible R package for nonnegative matrix factorization.
591   *BMC Bioinformatics* **11**, 367 (2010).

592   26.   Brunet, J.-P., Tamayo, P., Golub, T. R. & Mesirov, J. P. Metagenes and molecular
593   pattern discovery using matrix factorization. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 4164–
594   9 (2004).

595   27.   Cotto, K. C. *et al.* DGIdb 3.0: a redesign and expansion of the drug–gene interaction
596   database. *Nucleic Acids Res.* **46**, D1068–D1073 (2018).

597   28.   Verghese, P. B., Castellano, J. M. & Holtzman, D. M. Roles of Apolipoprotein E in
598   Alzheimer's Disease and Other Neurological Disorders. *Lancet Neurol.* **10**, 241
599   (2011).

600   29.   Costarelli, L., Malavolta, M., Giacconi, R. & Provinciali, M. Dysfunctional
601   macrophages in Alzheimer Disease: another piece of the "macroph-aging"
602   puzzle? *Aging (Albany. NY).* **9**, 1865–1866 (2017).

603   30.   Strittmatter, W. J. *et al.* Apolipoprotein E: high-avidity binding to beta-amyloid and
604   increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc.*
605   *Natl. Acad. Sci. U. S. A.* **90**, 1977–81 (1993).

606   31.   Bell, R. D. *et al.* Transport Pathways for Clearance of Human Alzheimer's Amyloid β-
607   Peptide and Apolipoproteins E and J in the Mouse Central Nervous System. *J. Cereb.*
608   *Blood Flow Metab.* **27**, 909–918 (2007).

609   32.   Sanchez, V. E., Nichols, C., Kim, H. N., Gang, E. J. & Kim, Y.-M. Targeting PI3K
610   Signaling in Acute Lymphoblastic Leukemia. *Int. J. Mol. Sci.* **20**, (2019).

611   33.   Biernacka, A., Dobaczewski, M. & Frangogiannis, N. G. TGF-β signaling in fibrosis.
612   *Growth Factors* **29**, 196–202 (2011).

613   34.   Yipp, B. G. & Kubes, P. NETosis: how vital is it? *Blood* **122**, 2784–94 (2013).

614   35.   Neutrophil Extracellular Traps in Systemic Sclerosis - Full Text View -
615   ClinicalTrials.gov. https://clinicaltrials.gov/ct2/show/NCT03374618.

616   36.   M. Artlett, C. The Role of the NLRP3 Inflammasome in Fibrosis. *Open Rheumatol. J.*
617   **6**, 80–86 (2012).

618   37.   Yu, Y. & Su, K. Neutrophil Extracellular Traps and Systemic Lupus Erythematosus. *J.*
619   *Clin. Cell. Immunol.* **4**, (2013).

620   38.   Lande, R. *et al.* Neutrophils Activate Plasmacytoid Dendritic Cells by Releasing Self-
621   DNA–Peptide Complexes in Systemic Lupus Erythematosus. *Sci. Transl. Med.* **3**,
622   73ra19-73ra19 (2011).

623   39.   Garcia-Romo, G. S. *et al.* Netting Neutrophils Are Major Inducers of Type I IFN
624   Production in Pediatric Systemic Lupus Erythematosus. *Sci. Transl. Med.* **3**, 73ra20-
625   73ra20 (2011).

626   40.   Villanueva, E. *et al.* Netting Neutrophils Induce Endothelial Damage, Infiltrate
627   Tissues, and Expose Immunostimulatory Molecules in Systemic Lupus Erythematosus.
628   *J. Immunol.* **187**, 538–552 (2011).

629   41.   Leffler, J. *et al.* Neutrophil Extracellular Traps That Are Not Degraded in Systemic
630   Lupus Erythematosus Activate Complement Exacerbating the Disease. *J. Immunol.*

631 **188**, 3522–3531 (2012).

632 42. Yang, W.-C. *et al.* Interleukin-4 Supports the Suppressive Immune Responses Elicited
633   by Regulatory T Cells. *Front. Immunol.* **8**, 1508 (2017).

634 43. Wang, R. *et al.* Association of interleukin 13 gene polymorphisms and plasma IL 13
635   level with risk of systemic lupus erythematosus. *Cytokine* **104**, 92–97 (2018).

636 44. Gao, W. *et al.* Notch signalling pathways mediate synovial angiogenesis in response to
637   vascular endothelial growth factor and angiopoietin 2. *Ann. Rheum. Dis.* **72**, 1080–8
638   (2013).

639 45. Moulton, V. R. & Tsokos, G. C. Abnormalities of T cell signaling in systemic lupus
640   erythematosus. *Arthritis Res. Ther.* **13**, 207 (2011).

641 46. Crow, M. K. & Rönnblom, L. Report of the inaugural Interferon Research Summit:
642   interferon in inflammatory diseases. *Lupus Sci. Med.* **5**, e000276 (2018).

643 47. Wagner, G. P. & Zhang, J. The pleiotropic structure of the genotype–phenotype map:
644   the evolvability of complex organisms. *Nat. Rev. Genet.* **12**, 204–213 (2011).

645 48. Yang, C., Li, C., Wang, Q., Chung, D. & Zhao, H. Implications of pleiotropy:
646   challenges and opportunities for mining Big Data in biomedicine. *Front. Genet.* **6**, 229
647   (2015).

648 49. Casciano, F., Pigatto, P. D., Secchiero, P., Gambari, R. & Reali, E. T Cell Hierarchy in
649   the Pathogenesis of Psoriasis and Associated Cardiovascular Comorbidities. *Front.*
650   *Immunol.* **9**, 1390 (2018).

651 50. Li, N. & Shi, R.-H. Updated review on immune factors in pathogenesis of Crohn's
652   disease. *World J. Gastroenterol.* **24**, 15–22 (2018).

653 51. Todd, N. W., Luzina, I. G. & Atamas, S. P. Molecular and cellular mechanisms of
654   pulmonary fibrosis. *Fibrogenesis Tissue Repair* **5**, 11 (2012).

655 52. Wynn, T. A. Integrating mechanisms of pulmonary fibrosis. *J. Exp. Med.* **208**, 1339–50
656   (2011).

657 53. Pattanaik, D., Brown, M., Postlethwaite, B. C. & Postlethwaite, A. E. Pathogenesis of
658   Systemic Sclerosis. *Front. Immunol.* **6**, 272 (2015).

659 54. Kolahian, S., Fernandez, I. E., Eickelberg, O. & Hartl, D. Immune Mechanisms in
660   Pulmonary Fibrosis. *Am. J. Respir. Cell Mol. Biol.* **55**, 309–322 (2016).

661 55. Wilton, K. M. & Matteson, E. L. Malignancy Incidence, Management, and Prevention
662   in Patients with Rheumatoid Arthritis. *Rheumatol. Ther.* **4**, 333–347 (2017).

663 56. Simon, T. A., Thompson, A., Gandhi, K. K., Hochberg, M. C. & Suissa, S. Incidence
664   of malignancy in adult patients with rheumatoid arthritis: a meta-analysis. *Arthritis*
665   *Res. Ther.* **17**, 212 (2015).

666 57. Din, F. V. N. *et al.* Effect of aspirin and NSAIDs on risk and survival from colorectal
667   cancer. *Gut* **59**, 1670–1679 (2010).

668 58. Dubois, R. N. Role of inflammation and inflammatory mediators in colorectal cancer.
669   *Trans. Am. Clin. Climatol. Assoc.* **125**, 358–72; discussion 372-3 (2014).

670 59. Chesmore, K., Bartlett, J. & Williams, S. M. The ubiquity of pleiotropy in human

671   disease. *Hum. Genet.* **137**, 39–44 (2018).

672   60.   Akobeng, A. K. Crohn's disease: current treatment options. *Arch. Dis. Child.* **93**, 787–
673   92 (2008).

674   61.   Ley, K., Rivera-Nieves, J., Sandborn, W. J. & Shattil, S. Integrin-based therapeutics:
675   biological basis, clinical use and new drugs. *Nat. Rev. Drug Discov.* **15**, 173–83
676   (2016).

677   62.   Ellinghaus, D. *et al.* Analysis of five chronic inflammatory diseases identifies 27 new
678   associations and highlights disease-specific patterns at shared loci. *Nat. Genet.* **48**,
679   510–518 (2016).

680   63.   Bernstein, C. N., Sargent, M. & Rector, E. Alteration in expression of beta 2 integrins
681   on lamina propria lymphocytes in ulcerative colitis and Crohn's disease. *Clin.*
682   *Immunol.* **104**, 67–72 (2002).

683   64.   Tsai, Y.-C. & Tsai, T.-F. Anti-interleukin and interleukin therapies for psoriasis: current
684   evidence and clinical usefulness. *Ther. Adv. Musculoskelet. Dis.* **9**, 277–294 (2017).

685   65.   Ito, H. *et al.* A pilot randomized trial of a human anti-interleukin-6 receptor
686   monoclonal antibody in active Crohn's disease. *Gastroenterology* **126**, 989–996
687   (2004).

688   66.   O'Reilly, S. Epigenetic modulation as a therapy in systemic sclerosis. *Rheumatology*
689   **58**, 191–196 (2019).

690   67.   Ridker, P. M. *et al.* Antiinflammatory Therapy with Canakinumab for Atherosclerotic
691   Disease. *N. Engl. J. Med.* **377**, 1119–1131 (2017).

692   68.   Terzić, J., Grivennikov, S., Karin, E. & Karin, M. Inflammation and Colon Cancer.
693   *Gastroenterology* **138**, 2101-2114.e5 (2010).

694   69.   Hübel, K. *et al.* Plerixafor with and without chemotherapy in poor mobilizers: results
695   from the German compassionate use program. *Bone Marrow Transplant.* **46**, 1045–
696   1052 (2011).

697   70.   Scleroderma Treatment With Autologous Transplant (STAT) Study - Full Text View -
698   ClinicalTrials.gov. https://clinicaltrials.gov/ct2/show/study/NCT01413100.

FIGURE LEGENDS

Figure 1: A Brief overview of methods, highlighting the integration of the 3 layers. **B** MeSH-MeSH network highlighting different MeSH (disease category) terms as nodes and the edges represent the genes common between the MeSH terms. Node names in network represent first two letters of the complete MeSH term as shown in the legend on right. **C-H** Gene prevalence over different MeSH terms. The degree distribution of genes across **I** MeSH terms and **J** diseases shown as histogram. **K** heatmap of gene expression across immunome of high degree genes in the DisGeNet. **L** Gene expression of APOE in the immunome. In **B, E-J**, size of node represents number of diseases in MeSH term.

Figure 2: DIME network of **A** lymphoid leukemia, **B** systemic scleroderma, **C** pulmonary fibrosis and **D** systemic lupus erythematosus. Green nodes represent genes, blue represents cell types and red represents diseases. Size of nodes is proportional to DAG score in genes and DAC score in cell types.

Figure 3: **A** Top pleiotropy network of the disease subset of the DisGeNet. Nodes are diseases that have a minimum degree of 2 nodes. Edges between diseases exist if the JI ≥ 0.1 and FET p-value ≤ 0.01 of the common DIME network between the diseases. Pleiotropy analysis between **B** Crohn's disease and psoriasis, **C** systemic scleroderma and pulmonary fibrosis, and **D** rheumatoid arthritis and colon carcinoma. **B, C, D** Venn diagrams represent overlap of cell-gene connections between the disease comparisons. JI and FET p-value are calculated for overlap in the genes in the DIME network of the diseases.

Figure 4: **A** Schema showing the different analysis performed in this study from building the DIME network (part1) to the pleiotropy (part 2) to the DIME-drug network (part3) and to the pleiotropy-drug network (part 4). **B** DIME-drug network of Crohn's disease. Pleiotropy-drug network for **C** Crohn's disease and psoriasis, **D** systemic scleroderma and pulmonary fibrosis, **E** systemic scleroderma and myocardial infarction, and **F** rheumatoid arthritis and colon carcinoma.

Supplementary Information

Supplementary Figure 1: Shows the degree of the top 10 high degree genes for different MeSH term categories and for all diseases.

Supplementary Figure 2: Heatmap of DAC scores for top weighted feature of 613 diseases from the DisGeNet.

Supplementary Figure 3: Reactome pathway enrichment analysis of top DAGs in the different clusters of the DIME network of **A** systemic scleroderma and **B** pulmonary fibrosis.

Supplementary Figure 4: Reactome pathway enrichment analysis of top DAGs in the different clusters of the DIME network of **A** systemic lupus erythematosus and **B** lymphoid leukemia.

Supplementary Figure 5: Reactome pathway enrichment analysis of pleiotropy DIME network of **A** Crohn's disease and psoriasis, **B** systemic scleroderma and pulmonary fibrosis, and **C** rheumatoid arthritis and colon carcinoma.

Supplementary Table 1: Represents all GEO datasets and samples used to construct the *immunome*.

Supplementary Table 2: List of genes as potential drug targets identified from the pleiotropy-drug network analysis. The degree represents the number of diseases in which the gene was found to be in the DIME-drug network. A total of 613 diseases were analyzed.

A — Lymphoid leukemia
B — Systemic scleroderma
C — Pulmonary Fibrosis
D — Systemic lupus erythematosus