Chromatin remodeling in bovine embryos indicates species-specific regulation of genome activation

Michelle M Halstead[1], Xin Ma[1,2], Richard M Schultz[3,4,†], and Pablo J Ross[1, †] *

[1] Department of Animal Science, University of California, Davis, CA, USA

[2] College of Animal Science and Technology, Jilin Agricultural University, Changchun, China

[3] Department of Anatomy, Physiology and Cell Biology, School of Veterinary Medicine, University of

California, Davis, CA, USA

[4] Department of Biology, University of Pennsylvania, Philadelphia, PA, USA

* Correspondence: Department of Animal Science, University of California, 2239 Meyer Hall, Davis, CA

95616, USA. Tel: +530-771-7225; E-mail: pross@ucdavis.edu

[†]Grant support: NIH-NICHD R01HD070044

Running Title: Open chromatin in mammalian embryos

Key words: chromatin accessibility / cow / genome activation / maternal-to-zygotic transition /

preimplantation embryo

Abstract

The maternal-to-zygotic transition (MZT) is underpinned by wide-spread transcriptomic and epigenomic

remodeling that facilitates totipotency acquisition. Although the MZT is a universal feature of mammalian

preimplantation development, the factors that regulate this transition remain unclear because the timing of

embryonic genome activation (EGA) is highly species-specific, suggesting divergence of regulatory

circuitry. Indeed, open chromatin during mouse and human EGA shows differential motif enrichment,

implicating distinct regulators. To better assess chromatin remodeling during mammalian development,

chromatin accessibility was profiled in bovine oocytes and *in vitro* produced embryos. Accessibility that

was established pre-EGA marked maternal factor recognition sites, i.e. CTCF, KLFs, NFY, and SP1,

echoing observations in humans and mice, and suggesting that a conserved set of maternal factors

regulate chromatin remodeling prior to EGA. Subsequently, open chromatin established during EGA was

primarily enriched for homeobox motifs, including that of DUX, a potential pioneer factor in mice and

humans. Remarkably, the open chromatin landscape during bovine EGA clearly resembled that of

humans', revealing that cattle could be a more relevant model for human development than mouse.

Introduction

Preimplantation development encompasses several critical milestones as embryos progress following

fertilization to blastocyst implantation. Fusion of the transcriptionally quiescent oocyte and sperm results

in a zygote with two haploid pronuclei. Gametic syngamy is completed during the first round of

replication, when pronuclear membranes dissolve, allowing maternal and paternal chromosomes to

intermingle on the metaphase plate. Subsequent rounds of cleavage ultimately lead to formation of a

blastocyst. However, the cleavage-stage embryo must first complete the maternal-to-zygotic transition

(MZT), wherein the embryo assumes control over its own continued development by degrading oocyte-

derived products and initiating its own transcriptional program. This dramatic change in gene expression

proceeds gradually; minor embryonic genome activation (EGA) results in low levels of transcription in

early cleavage-stage embryos (Abe *et al*, 2018; Ram & Schultz, 1993), and leads to major EGA, which

involves wide-spread activation of embryonic transcription (Schulz & Harrison, 2019; Vastenhouw *et al*, 2019). This shift from maternal dependence to self-sufficiency serves at least three functions: elimination of oocyte-specific messages, replenishment of transcripts that are common to both the oocyte and the embryo, and generation of novel embryonic-specific transcripts (Latham *et al*, 1991).

While embryos undergo this massive shift in transcriptional program during the MZT, wide-spread changes in chromatin structure (Eckersley-Maslin *et al*, 2018; Xu & Xie, 2018) and other epigenetic marks are simultaneously restructuring the embryonic epigenome. Chromatin remodeling is necessary to eradicate gamete-specific signatures and establish an open chromatin landscape that supports embryonic transcriptional programs. Specifically, chromatin structure defines the genomic context within which transcriptional machinery can operate, thereby determining the cell-specific gene expression patterns that confer cell identity and function.

Following fertilization, the zygotic genome is globally demethylated (Iurlaro *et al*, 2017; White *et al*, 2016), and this loss of DNA methylation coincides with global decreases in repressive histone modifications (Eckersley-Maslin *et al*, 2018; Xu & Xie, 2018; Ross *et al*, 2008). Overall, epigenetic factors linked to relaxed chromatin are more abundant in mouse zygotes, whereas factors implicated in chromatin compaction become more prevalent during EGA (Funaya & Aoki, 2017), pointing to a more permissive chromatin landscape in pre-EGA embryos. Indeed, mouse zygotes demonstrate elevated histone mobility (Ooga *et al*, 2016), highly dispersed chromatin(Ahmed *et al*, 2010), and lack chromocenters (congregations of pericentromeric heterochromatin) (Martin *et al*, 2006; Probst *et al*, 2007, 2010). Moreover, in mouse, 3-D chromatin architecture is largely absent after fertilization, and is gradually established throughout preimplantation development, which facilitates long-distance chromatin interactions in later stage embryos (Du *et al*, 2017), and indicates increasing chromatin compaction and organization. The co-occurrence of genome activation and chromatin remodeling raises an interesting causality dilemma, namely, whether chromatin remodeling is necessary for transcription activation, or

whether transcription activation leads to chromatin accessibility. Several maternal products prompt

transcription initiation by altering chromatin structure (Bultman *et al*, 2006; Jimenez *et al*, 2015;

Hernandez *et al*, 2018), and some chromatin compaction occurs in the absence of embryonic transcription

(Du *et al*, 2017). However, inhibiting embryonic transcription pervasively disrupts the establishment of

open chromatin during EGA(Wu *et al*, 2016, 2018), suggesting that EGA and chromatin remodeling are

likely interdependent.

In mouse and human preimplantation embryos, accessible sites are gradually established as development

progresses (Li *et al*, 2018; Wu *et al*, 2018; Gao *et al*, 2018; Liu *et al*, 2019; Lu *et al*, 2016; Wu *et al*,

2016); however, these open regions demonstrate different motif enrichment patterns, implicating distinct

sets of transcription factors (TFs) in either murine (RARG, NR5A2, ESRRB), (Wu *et al*, 2016) or human

EGA (OTX2, GSC, POU5F1) (Wu *et al*, 2018; Gao *et al*, 2018). Although some TFs appear to regulate

EGA in multiple species, i.e. Krüppel-like factors (KLFs) (Bogliotti *et al*, 2019; Wu *et al*, 2016, 2018),

double homeobox (DUX) (Whiddon *et al*, 2017; Hendrickson *et al*, 2017; De Iaco *et al*, 2017; Young *et*

al, 2013; Kruse *et al*, 2019; Iturbide & Torres-Padilla, 2017), ZSCAN4 (Falco *et al*, 2007; Takahashi *et*

al, 2019), and CTCF (Wu *et al*, 2016, 2018), it remains unclear if there is any mechanistic conservation

across mammals. In fact, the timing of genome activation is highly species-specific: major EGA in mice

occurs during the 2-cell stage (Latham *et al*, 1991; Ram & Schultz, 1993), in humans (Braude *et al*, 1988)

and pigs (Davis, 1985) during the 4- to 8-cell stage, and in sheep (Crosby *et al*, 1988) and cattle (Frei *et*

al, 1989; Barnes & First, 1991) between the 8- and 16-cell stages. The relative timing with which mice

activate their genomes could indicate that the mechanism behind murine EGA differs significantly from

other species', which would have significant implications for modeling human preimplantation

development. In particular, the timing of bovine EGA more closely resembles that of human EGA, as do

global changes to histone PTMs. For instance, the active mark trimethylation of lysine 4 on histone 3

(H3K4me3) decreases in global abundance during human(Zhang *et al*, 2012) and bovine EGA(Wu *et al*,

2011), but increases during murine EGA(Liu *et al*, 2016; Zhang *et al*, 2016). However, the chromatin

remodeling events that underscore bovine preimplantation development have yet to be catalogued, and it

remains unclear whether the regulation and execution of the MZT in cattle resembles that which has been

observed in humans.

To this end, we here describe the chromatin accessibility landscapes of bovine oocytes and

preimplantation embryos. We find that open chromatin is gained progressively throughout development,

with promoters enriched for CTCF and NFY motifs gaining accessibility in earlier stages, and distal open

chromatin becoming accessible at later stages. Moreover, embryonic transcription was not required for

the appearance of promoter open chromatin in 2- and 4-cell embryos, but was absolutely necessary to

establish stage-specific and distal open chromatin, especially in 8-cell embryos, indicating that maternal

and embryonic products both participate in chromatin remodeling in a complementary fashion. Sequence

enrichment in open chromatin revealed that several TFs likely play roles in both bovine and human EGA

(OTX2, SP1), whereas regulators specific to murine EGA(Wu *et al*, 2016) demonstrate no enrichment in

bovine (RARG, NR5A2, ESRRB), suggesting that cattle may be a more informative model for human

preimplantation development. Nevertheless, several TFs (DUX, KLFs, CTCF, NFY) seem to play a role

in preimplantation development in all three species, raising the possibility that genome activation may be

mechanistically conserved across mammals, whereas the specific transcriptional program that is activated

may differ between species.

Results and Discussion

Global dynamics of open chromatin in bovine preimplantation embryos

For each developmental stage, ATAC-seq libraries were prepared from cells derived from three separate

oocyte collections. A subset of embryos from each collection was also cultured in the presence of the

transcriptional inhibitor, α-amanitin, to interrogate the relationship between embryonic transcription and

chromatin remodeling (Figure 1a). Between 30 and 87 million non-mitochondrial monoclonal uniquely

mapping reads were collectively obtained for each developmental stage, and at least 20 million non-

mitochondrial monoclonal reads were collectively obtained for transcription blocked embryos (TBE) at

each stage (Table S1). Genome-wide normalized ATAC-seq coverage demonstrated a striking shift in the

open chromatin landscape between 4- and 8-cell embryos (Figure 1b), suggesting that large-scale

chromatin remodeling coincided with the main transcriptomic shift observed during major EGA (Figure

S1). Similarity between replicates (Table S2) indicated that both the technique and embryo production

were robust, generating comparable chromatin accessibility profiles across different rounds of oocyte

collection and embryo production (Figure 1c). Reads from replicates were pooled together to obtain

greater sequencing depth and maximize power for identifying regions of open chromatin. To gauge

changes in chromatin accessibility throughout development, regions of open chromatin, or peaks, were

called for each stage of development. To minimize bias from sequencing depth, peaks were called from

either 30 million monoclonal uniquely mapped reads when comparing controls, or 20 million reads when

comparing TBE with control embryos.

As observed in humans and mice (Lu *et al*, 2016; Wu *et al*, 2016, 2018; Gao *et al*, 2018; Liu *et al*, 2019),

regions of open chromatin were gradually established throughout bovine preimplantation development,

with the lowest enrichment for accessible sites in 2-cell embryos (Figure 1d). Rather than reflecting

chromatin inaccessibility in 2-cell embryos, this dearth of canonical "open chromatin" probably results

from a highly permissive chromatin structure. Indeed, chromocenters are absent from bovine embryos

until the early 8-cell stage (Martin *et al*, 2006), indicating a relaxed chromatin configuration in early

cleavage-stage embryos. Because assays that employ endonucleases – e.g., ATAC-seq and DNase-seq –

depend on increased cutting events at consistently accessible loci, we speculate that genome-wide

chromatin relaxation would lead to random cutting events genome-wide, resulting in the observed high

background and low enrichment in 2-cell embryos. Attempts to use endonuclease-based methods to

profile open chromatin in mouse zygotes (Lu *et al*, 2016; Wu *et al*, 2016) and human 2-cell embryos (Liu

et al, 2019; Gao *et al*, 2018) have encountered similar difficulties with low enrichment. As in bovine

embryos, chromocenters are also conspicuously absent in pre-EGA mouse embryos (Martin *et al*, 2006;

Probst *et al*, 2007, 2010), and electron microscopy (Ahmed *et al*, 2010) and fluorescence recovery after

photobleaching analysis (Ooga *et al*, 2016) also indicate highly dispersed chromatin. Furthermore, a

recent study that detected open chromatin based on methylation of accessible GpC sites, rather than

endonucleases, found that genome-wide accessibility in human embryos actually decreased from the

zygote stage onward (Li *et al*, 2018). Global chromatin relaxation appears to be a shared characteristic of

human, bovine, and murine pre-EGA embryos. Thus, it is tempting to speculate that this "naïve"

chromatin state acts as a blank epigenetic slate, which is then gradually compacted and structured to meet

the needs of the growing embryo. Indeed, accessible sites were progressively established in 4-cell, 8-cell,

and morula stage embryos (Figure 1d). Interestingly, many of these regions were only open transiently at

a specific stage, whereas others maintained their accessibility throughout later stages (Figure 1e),

suggesting that chromatin remodeling serves two functions: progressive establishment of a "totipotent"

chromatin landscape, and transient stage-specific regulation.

Stepwise remodeling yields stage-specific open chromatin with distinct functionality

To delve further into the potential function of regions that lost or gained accessibility between

consecutive stages (Figure 2a), intergenic regions of open chromatin were evaluated for sequence

enrichment, revealing that the binding motifs of distinct groups of TFs were gaining accessibility at each

developmental stage (Table S3). These enriched sequences corresponded to the known binding motifs of

several TFs that were either maternally provided or expressed in embryos (Figure S2), demonstrating that

changing chromatin structure subjects each stage of development to distinct regulatory circuitry (Figure

2b).

The first major transition in chromatin structure mostly involved loss of hyperaccessible sites in 2-cell

embryos following fertilization (Figure 2a). Intergenic loci that lost accessibility during this transition

(n=54,264) were enriched for the binding motifs of CTCF – an insulator protein implicated in 3D

chromatin organization (Ghirlando & Felsenfeld, 2016) – and several RFX factors – a highly conserved

family of transcriptional repressors (Lubelsky *et al*, 2005) (Figure 2b). Interestingly, one quarter of the

intergenic regions that closed in 2-cell embryos regained accessibility at the 4-cell stage (n=14,859), and

many of these remained open all the way through to the morula stage (n=6,186). Most intergenic regions

that close after fertilization likely contribute to oocyte-specific regulation, considering that they remain

inaccessible in embryos, whereas regions that re-opened in 4-cell embryos could be implicated in general

house-keeping functions.

The first major gains in accessibility occurred in 4-cell embryos (Figure 2a). About a quarter of the

regions that opened during the 4-cell stage maintained their accessibility in both 8-cell embryos and

morulae (Figure 1e), although 74% of these had been previously open in GV oocytes (n=15,466).

Surprisingly, nearly half of the regions that opened during the 4-cell stage were only transiently accessible

(Figure 1e). Four-cell-specific open chromatin was most significantly enriched for binding motifs of

NFkB family members (Figure 2c), which was particularly intriguing because NFkB activation is

specifically necessary in 1-cell mouse embryos for development to progress past EGA (Nishikimi *et al*,

1999; Yang *et al*, 2017). Although NFkB factors are maternally provided (Figure 2c), they are initially

sequestered in the cytoplasm until they translocate into the nucleus at the early 1-cell stage in mice

(Nishikimi *et al*, 1999; Yang *et al*, 2017) and the 4-to-8 cell stage in cattle (Paciolla *et al*, 2011). In

particular, one of the NFkB subunits capable of activating gene expression – RELA – binds DNA with

increased frequency in cattle embryos compared to oocytes (Paciolla *et al*, 2011), suggesting that NFkB

activation of target gene expression could be one of the first events in a cascade leading to major EGA. In

fact, one of the few genes with upregulated expression in 4-cell embryos, as compared to MII oocytes,

was TRIM8 – a positive regulator of NFkB activity (Figure S3). Additionally, RELA binding sites that

were accessible in 4- and 8-cell embryos mark genes that encode key regulators of early preimplantation

development, raising the possibility that NFkB binding is involved in their transcription initiation (Figure

S4), although the contribution of NFkB signaling to gene expression during minor EGA in cattle has not

yet been established.

In contrast to open chromatin in pre-EGA embryos, regions that gained accessibility at the 8-cell stage

(Figure 2a) were considerably enriched for the binding motifs of several homeobox TFs, including OTX2,

GSC, CRX, PHOX2A, PAX7, and PITX1 (Figure 2b,e), although only OTX1, OTX2, and PITX2 were

appreciably expressed during the 8-cell stage (Figure 2f). More so than other homeobox recognition

motifs, 8-cell open chromatin was most enriched for binding motifs of the double homeobox (DUX)

family, which has been extensively implicated in EGA regulation (Whiddon *et al*, 2017; Hendrickson *et*

al, 2017; De Iaco *et al*, 2017; Young *et al*, 2013; Kruse *et al*, 2019; Iturbide & Torres-Padilla, 2017).

More than a third of the accessible intergenic loci in 8-cell embryos harbored a sequence that most closely

matched double homeobox (DUX) binding motifs (Figure 2g). DUX factors are increasingly recognized

as potential developmental pioneer factors, participating in chromatin remodeling(Kruse *et al*, 2019;

Hendrickson *et al*, 2017), and expression of repetitive elements(Whiddon *et al*, 2017; Hendrickson *et al*,

2017; De Iaco *et al*, 2017; Young *et al*, 2013) and cleavage-stage genes(Hendrickson *et al*, 2017; Iturbide

& Torres-Padilla, 2017). In cattle, DUXA is expressed transiently and strongly during the 8- and 16-cell

stages (Figure 2h), similar to DUX4 and Dux expression in human and mouse embryos, respectively, for

a specific window prior to EGA(Hendrickson *et al*, 2017; De Iaco *et al*, 2017). Although the maternal

factors DPPA2 and DPPA4 induce Dux expression in mouse embryonic stem cells (ESC) (Eckersley-

Maslin *et al*, 2019), only DPPA3 is maternally provided in cattle (Figure S5). It remains unclear if

DPPA3 is capable of inducing DUX expression; however, DPPA3 is maternally provided in mice, cattle,

and humans (Boroviak *et al*, 2018; Bakhtari & Ross, 2014), and knockdown DPPA3 decreases the

developmental competency of mouse (Payer *et al*, 2003) and bovine (Bakhtari & Ross, 2014) embryos.

Although the maternal factors that induce DUXA expression in cattle remain unclear, the synchronized

upregulation of DUXA and increased accessibility of its binding sites during bovine development

strongly indicates a conserved role for DUX in mammalian preimplantation development, especially

considering that DUX family members are highly conserved and specific to placental mammals (Young *et*

al, 2013; Leidenroth & Hewitt, 2010). Further functional validation will be necessary to determine if

DUXA is required for bovine embryogenesis, or if is an important but non-essential regulator of EGA, as

in mouse (Chen & Zhang, 2019; Guo *et al*, 2019; Iaco *et al*, 2019).

The expression profile of DUXA in bovine preimplantation development closely mirrored that of another

TF implicated in EGA: ZSCAN4 (Figure S6a) (Ko, 2016). ZSCAN4 is a downstream target of DUX

factors in humans (Hendrickson *et al*, 2017) and mice (De Iaco *et al*, 2017; Whiddon *et al*, 2017), and the

coordinated expression of these two factors in cattle certainly suggests a similar mechanism at play.

Although ZSCAN4 depletion disrupts development past EGA in mice (Falco *et al*, 2007) and cattle

(Takahashi *et al*, 2019), its binding motifs were not enriched in open chromatin during bovine

preimplantation development (Figure S6b-d), suggesting that the binding motif of bovine ZSCAN4

differs from those in mice and humans.

Finally, the transition from 8-cell to morula also incurred extensive chromatin remodeling, with even

more sites gaining accessibility than during the 8-cell stage (Figure 2a). Compared to earlier stages,

intergenic loci that gained accessibility in morulae were primarily enriched for GATA factor binding

motifs (Figure 2a; Table S3) – key regulators of trophectoderm establishment and maintenance (Bai *et al*,

2013) – bolstering the idea that morulae are already initiating differentiation programs necessary for

blastocyst formation.

Progressive establishment of maintained open chromatin sets the stage for genome activation

Although many regions only experienced stage-specific gains in accessibility, a stable open chromatin

landscape was also progressively established after fertilization. As early as the 2-cell stage, regions began

to gain accessibility that was maintained until the morula stage. These regions of maintained accessibility

were enriched for CTCF motifs (Figure 3a; Table S4b), especially those that were first established during

the 4-cell stage. CTCF binding delineates chromatin loop boundaries, thus determining the genomic space

within which genes can interact with their regulatory elements (Ghirlando & Felsenfeld, 2016). Therefore,

enrichment of CTCF motifs in maintained peaks could point to a gradual establishment of a stable 3-D

chromatin architecture in preparation for major EGA. This proposal is consistent with 3-D chromatin

architecture in mouse embryos, which is greatly diminished after fertilization and then gradually re-

established throughout preimplantation development, facilitating long-distance chromatin contacts, i.e.,

promoter-enhancer interactions, in later stage embryos (Du *et al*, 2017). Indeed, transcription during

minor EGA in mouse is primarily driven by proximal promoters, whereas enhancers are dispensable for

transcription until major EGA (Abe *et al*, 2015; Wiekowski *et al*, 1993). The global reorganization of 3D

chromatin architecture in mouse has also been observed in bovine embryos, wherein gene-rich regions

switch from a random distribution to a chromosome-specific distribution during major EGA (Koehler *et*

al, 2009). Collectively, these results suggest that gradual establishment of 3D chromatin architecture is a

conserved feature of pre-EGA embryos, although the mechanisms regulating this restructuring remain

unknown.

Other than CTCF, maintained peaks were also more enriched for KLF motifs than transiently open sites

(Figure 3b). Although most KLFs were not expressed until the 8-cell stage, several KLFs were maternally

provided (Figure 3c), including KLF4, a master regulator of induced pluripotent stem cell (iPSC)

reprogramming (Takahashi & Yamanaka, 2006). Recent evidence suggests that KLF4 contributes to

reprogramming iPSC by mediating pluripotency-associated enhancer-promoter contacts (Di Giammartino

et al, 2019); thus, the concurrent establishment of open chromatin corresponding to CTCF and KLF

binding motifs suggests that pre-EGA embryos are priming a similar mechanism for use during major

EGA, especially because about 50% of activated genes during bovine major EGA contain KLF motifs in

their promoters (Bogliotti *et al*, 2019).

In fact, loci that remained open from the 2- and 4-cell stages onward occurred in genic regions more often

than stage-specific open chromatin (Figure 3d) and marked the promoters of genes that were functionally

enriched for housekeeping functions (Figure 3e), including 33 of the 51 genes that were upregulated in 4-

cell embryos compared to MII oocytes (Figure S3). Moreover, these regions of maintained accessibility

were strongly enriched for NFY and SP1 binding sites (Table S4a,b), which is highly reminiscent of

chromatin remodeling in mouse embryos, as proximal promoters enriched for NFY are the first regions to

gain accessibility (Lu *et al*, 2016). Of note, NFY enhances binding of the pluripotency factors POU5F1,

SOX2, and KL4 to their recognition motifs (Oldfield *et al*, 2014; Soufi *et al*, 2015), and is clearly

involved in murine EGA, as NFY knockdown embryos demonstrated impaired open chromatin

establishment and downregulation of gene expression (Lu *et al*, 2016). Similarly, SP1 is essential for

early mouse development, with knockout embryos arresting at day 11 of gestation (Marin *et al*, 1997).

Intriguingly, human zygotes also demonstrate KLF, SP1, and NFY motif enrichment in open chromatin

(Liu *et al*, 2019), overall suggesting that a conserved set of maternal regulators participates in chromatin

remodeling and transcription activation in pre-EGA embryos.

Both maternal products and embryonic transcription drive chromatin reorganization

Considering that embryonic transcription is extremely limited before the 8-cell stage in cattle, the

appearance of open chromatin in pre-EGA embryos suggests that maternal factors participate in

chromatin remodeling. To further dissect the maternal contribution to epigenetic reprogramming and

EGA, embryos were cultured in the presence of α-amanitin to inhibit POLR2-dependent transcription

elongation. Loss of embryonic transcription had a drastic inhibitory effect on the appearance of open

chromatin and intensified as development progressed (Figure 4a); 64% of loci that should have opened

during the 4-cell stage failed to become accessible without embryonic transcription, and in 8-cell TBE

embryos, 96% of loci that should have opened remained closed (Figure 4b), disrupting the chromatin state

of key genes, such as KLF4 (Figure 4c), and coinciding with developmental arrest.

Interestingly, inhibiting embryonic transcription did not uniformly affect chromatin remodeling genome-

wide, but instead preferentially disrupted establishment of stage-specific open chromatin, whereas

maintained open chromatin established in 2- and 4-cell embryos appeared despite embryonic transcription

inhibition (Figure 4d). Transcription-independent maintained open chromatin marked the promoters of

nearly 60% of embryonically--expressed genes (n=1,038/1,784 genes identified from data from Bogliotti

et al 2019; Figure S7), suggesting that maternal factors actively remodel the local chromatin structure of

target genes, possibly priming them for expression later on in development. Nevertheless, the appearance

of stage-specific open chromatin almost completely depended on embryonic transcription, indicating that

maternal and embryonic factors cooperate in a complementary fashion to establish the appropriate

chromatin landscape for activation of embryonic transcriptional programs.

310 Open chromatin in preimplantation embryos is enriched for repetitive elements

Evaluation of changes in chromatin structure as they relate to gene expression gives an incomplete

perspective of the genome-wide changes that occur during preimplantation development, because the

MZT is not just characterized by a shift in the transcriptome, but also in the repeatome. Similar to other

mammalian species, 44% of the bovine genome is comprised of repeats derived from retrotransposons

(Adelson *et al*, 2009) – interspersed repeats that are increasingly thought to play major roles in cellular

processes and development. Retrotransposons propagate through a 'copy and paste' mechanism, and their

expression is generally suppressed to avoid deleterious integrations (Friedli & Trono, 2015; Hancks &

Kazazian Jr, 2016). However, retrotransposons are often actively transcribed in early embryos, and

although this phenomenon was recently thought to be nothing more than opportunistic expression by

repetitive elements due to an unusually permissive chromatin state in the developing embryo, the activity

of some retrotransposons is actually crucial for development (Spadafora, 2004). Although the specific

mechanisms behind this necessity are still being investigated, transposable elements have been implicated

at several regulatory levels, as they can provide binding sites for TFs, allowing them to be co-opted as

alternative promoters and enhancers (Chuong *et al*, 2017) and participate in 3-D chromatin architecture

(Raviram *et al*, 2018).

Although repetitive element expression in bovine early embryos was reported a decade ago using a cDNA

array (Bui *et al*, 2009), a complete catalogue of repeat transcription throughout bovine preimplantation

development was lacking. To address this gap in knowledge, available RNA-seq data (Graf *et al*, 2014)

were assessed for expression of repetitive elements. Importantly, these libraries were not subjected to

polyA selection. As has been observed in the mouse and human (Yandım & Karakülah, 2019; Franke *et*

al, 2017), expression and accessibility of repetitive elements throughout bovine preimplantation

development was highly stage-specific and dynamic (Figure 5a; Figure S8).

Non-long terminal repeat (non-LTR) retrotransposons – long interspersed elements (LINEs) and short

interspersed elements (SINEs) – are increasingly transcribed during human preimplantation development

(Yandım & Karakülah, 2019; Li *et al*, 2018). In cattle, LINEs also demonstrated increasing expression

(Figure 5a) and accessibility (Figure 5b) starting during the 8-cell stage. Although the function of LINE

elements in bovine preimplantation embryos has yet to be established, their activation is crucial for mouse

development: perturbing LINE expression in mouse preimplantation embryos causes developmental arrest

at the 2-cell stage and perturbs gene expression (Beraldi *et al*, 2006). Moreover, LINE-1 activation may

regulate global chromatin accessibility in mouse embryos (Jachowicz *et al*, 2017). Of the SINE families,

MIR element expression and accessibility patterns most echoed those of L2 LINE elements, with

increased expression and accessibility starting at the 8-cell stage, suggesting that these elements could be

acting as enhancers or promoters.

In particular, LTR activation is a key feature in human (Yandım & Karakülah, 2019), mouse (Franke *et*

al, 2017), and bovine preimplantation development (Bui *et al*, 2009). Of these, mammalian LTRs and

ERVLs were increasingly enriched in open chromatin starting during the 8-cell stage (Figure 5b),

although their transcript abundance dropped throughout development (Figure 5a), indicating that other

mechanisms likely regulate repeat expression, e.g., DNA methylation or histone modifications (Bestor &

Bourc'his, 2004; Martens *et al*, 2005; Shirai *et al*, 2017). Only endogenous retroviral K (ERVK) elements

demonstrated both increasing expression (Figure 5a) and moderately increased prevalence in distal open

chromatin during the 8-cell stage (Figure 5b), suggesting that ERVK elements function as regulatory

elements, as observed in human preimplantation embryos(Pontis *et al*, 2019).

Mounting evidence suggests that specific types of LTR retrotransposons, especially intact elements, play

pivotal roles in early development. During bovine preimplantation development, the most expressed

retrotransposons were ERV1-1_BT and ERV1-2_BT (Figure 6a), as has been observed previously (Bui *et*

al, 2009). Upon further inspection, intact ERV1-1_BT elements often co-occurred with MER41_BT

repeats in a specific sequence, which demonstrated a highly reproducible pattern of transcription at

ERV1-1_BT elements and chromatin accessibility at MER41_BT elements (Figure S9a). Furthermore,

MER41_BT elements that were accessible in 8-cell embryos were enriched for the binding motifs of

several pluripotency factors, including POU5F1, NFY, KLF4, OTX2, and TEAD (Figure S9b),

suggesting that pluripotency factors are driving transposon expression.

The TF DUX has also been implicated in driving the expression of intact ERVL elements in human and

mouse pre-EGA embryos (Whiddon *et al*, 2017; Hendrickson *et al*, 2017; De Iaco *et al*, 2017; Liu *et al*,

2019). Specifically, in human embryos DUX4 appears to bind MLT2A1 elements – primate-specific

LTRs that flank human ERVL –activating their expression (Liu *et al*, 2019). Considering that intergenic

open chromatin was especially enriched for DUX binding sites in 8-cell bovine embryos, it seemed likely

that these sites would also correspond to retrotransposons. Indeed, several MLT elements were enriched

in 8-cell open chromatin harboring DUX motifs (Table S5), suggesting that bovine DUX binds LTRs

with sequence similarity to primate-specific MLT2A1 elements, regulating their expression. Nearly all

LTRs that were enriched in 8-cell open chromatin with DUX binding sites demonstrated dynamic

expression profiles throughout development (Figure S10). In particular, increasing accessibility at

MLT1A0 elements harboring DUX recognition sites (Figure 6c) was mirrored by a sharp increase in

transcription in 8-cell embryos (Figure 6d).

Whether the transcripts derived from LTRs have a functional role in bovine development, or are simply a

result of opportunistic expression, remains to be established. However, evidence in other species suggests

that retrotransposon-derived regulatory elements are often co-opted by the embryo as promoter and

enhancers(Chuong *et al*, 2017): a phenomenon which appears to extend to bovine embryos, as ATAC-seq

and RNA-seq suggest that MLT1A0 elements are co-opted as alternative promoters at several loci,

including CD1 (Figure 6e), ZNF41, and LPIN2. As such, an interesting balance appears to exist between

repetitive elements and the embryo, wherein the repeatome leverages the embryo's existing regulatory

network to drive transposition, while simultaneously providing new regulatory elements and TF binding

sites that the embryo co-opts to drive the expression of its own transcriptional programs.

A model for mammalian genome activation

Several lines of evidence suggest that the regulatory circuitry responsible for the MZT may differ

between mammals. First, EGA occurs in a highly species-specific fashion, with the major wave of

transcription in mice during the 2-cell stage (Latham *et al*, 1991; Ram & Schultz, 1993), in humans

during the 4-to-8-cell stage (Braude *et al*, 1988), and cattle during the 8-to-16-cell stage (Frei *et al*, 1989).

Second, the expression patterns of repetitive elements are not only highly-stage specific, but also species-

specific; primate-specific and murine ERVL elements are strongly expressed during human and mouse

preimplantation development, respectively (Whiddon *et al*, 2017; Hendrickson *et al*, 2017; De Iaco *et al*,

2017; Liu *et al*, 2019), whereas specific ERV1 elements were most prominently activated in bovine

embryos. Finally, the maternal programs in human and mice are divergent; human maternal programs

conspicuously lack the murine maternal effect transcripts POU5F1, HSF1, and DICER1, and are

functionally enriched for translational processing (Boroviak *et al*, 2018), reflecting the need to translate

maternal transcripts during the extended period before human EGA.

Maternal products are essential to support cleavage stage-embryos prior to genome activation, especially

in species where EGA is delayed for several cell divisions, e.g., humans and cattle. The relaxed chromatin

structure in early preimplantation embryos provides a unique regulatory context for maternal factors.

Whereas the condensed chromatin structure in somatic cells generally restricts DNA-binding proteins to

regions of open chromatin, the dispersed chromatin in 2-cell bovine embryos may allow maternal factors

to opportunistically and pervasively bind their recognition motifs. Indeed, several maternal factors appear

to participate in chromatin restructuring in pre-EGA embryos, leading to open chromatin establishment at

promoters enriched for NFY and SP1 binding sites, as well as CTCF and KLF motifs. Although not

maternally provided, the DUX family appears to play a conserved role in genome activation. In humans

and mice, DUX factors have been implicated in chromatin remodeling (Kruse *et al*, 2019; Hendrickson *et*

al, 2017) and transcription activation of cleavage-stage genes (Hendrickson *et al*, 2017; Iturbide &

Torres-Padilla, 2017). We find a similar pattern of DUX expression in bovine embryos, as well as

increased accessibility of DUX binding sites around the 8-cell stage, suggesting that DUX may also

modulate gene expression and chromatin accessibility during bovine EGA.

Although the chromatin landscape changes markedly upon major EGA in bovine, human (Wu *et al*, 2018;

Liu *et al*, 2019; Li *et al*, 2018), and mouse (Lu *et al*, 2016; Wu *et al*, 2016) embryos, the regulatory

circuitry that is active during this stage in development appears to differ significantly between humans

and mice (Wu *et al*, 2018), which suggests that regulation of mammalian EGA is highly species-specific.

To identify and compare putative regulators of mammalian EGA, regions that were accessible during

major EGA in bovine, human, and mouse embryos were evaluated for binding motif enrichment of

actively expressed TFs. Comparing EGA regulatory circuitry between species reflects a critical

divergence in regulatory and transcriptional programs that clearly separates humans and cattle from the

mouse (Figure S11). Compared to mouse 2-cell embryos, open chromatin in bovine and human 8-cell

embryos demonstrated remarkably similar patterns of sequence enrichment (Figure 7). In cattle and

humans, SP1, OTX2, and NFY were particularly implicated during major EGA, whereas NR5A2, RARG,

and ESRRB were solely enriched in mouse embryos. Nevertheless, it is unclear if the regulatory factors

that are enriched in open chromatin during major EGA are regulators of EGA or simply products of it.

For instance, although NR5A2is enriched in 2-cell mouse embryos, it is an early regulator of inner cell

mass and trophectoderm programs and is not essential for genome activation (Wu *et al*, 2016). Similarly,

OTX2 is essential for neuronal lineage specification in mice (Perea-Gomez *et al*, 2001), and has been

implicated in the transition from naïve to primed pluripotency (Acampora *et al*, 2013; Buecker *et al*,

2014; Schoorlemmer *et al*, 2014). However, OTX2 protein is clearly present in human and marmoset

zygotes (Boroviak *et al*, 2018), suggesting that this homeobox TF may play an as-of-yet undetermined

role in regulation of human and bovine EGA. Although several TFs appeared to only be important in

either mouse, or in cattle and humans, KLF factors were substantially enriched during EGA in all three

species. Considering the well-established role of KLF factors in somatic cell reprogramming and

establishment of pluripotency in multiple species (Takahashi & Yamanaka, 2006; Graf, 2011), KLFs may

play a conserved role in the MZT. Although future research will be necessary to elucidate the function of

specific regulatory factors, the high consistency between cattle and humans, both with respect to the

timing of EGA and the regulatory circuitry that accompanies it, strongly suggests that cattle are a more

appropriate model system for human preimplantation development than mouse.

Conclusions

Sweeping changes to chromatin structure during bovine preimplantation development suggest that 2-cell

embryos are characterized by globally decondensed chromatin, which is gradually compacted as

development progresses, which echoes similar observations in humans and mice. In particular, it is

tempting to speculate that a conserved set of maternal factors establish basal promoter accessibility and

the necessary chromatin architecture for enhancer-promoter interactions that will drive gene expression

during major EGA (Figure 8). However, the open chromatin landscape during major EGA clearly

distinguished mice from cattle and humans, suggesting that whereas maternal regulation of EGA may be

conserved across mammals, the transcriptional programs that are subsequently activated have diverged

substantially. Practically, this difference suggests that human development may be better modeled in

cattle than in mice. Nevertheless, the factors appear to regulate the MZT in cattle, humans, and mice

certainly warrant further investigation and validation, which will provide invaluable insight into the

regulatory framework that governs successful preimplantation development

Materials and Methods

Oocyte collection and maturation

Ovaries were procured from a local abattoir and transported to the laboratory in a warm saline solution.

Follicles measuring 2-10 mm were aspirated to obtain cumulus oocyte complexes (COCs). Only COCs

with healthy layers of cumulus cells were selected for maturation. These were washed in collection

medium (6:4 M199 (Sigma M7653): SOF-Hepes, supplemented with 2% fetal bovine serum (FBS;

Hyclone/Thermo Scientific) and transferred to maturation medium (modified M199 medium (Sigma

M2154) supplemented with ALA-glutamine (0.1 mM), sodium pyruvate (0.2 mM), gentamicin (5 μg/ml),

EGF (50 ng/ml), oFSH (50 ng/ml), bLH (3 μg/ml), cysteamine (0.1 mM), and 10% FBS.

In vitro fertilization and embryo culture

After COCs matured for 24 h, MII oocytes were washed in SOF-IVF medium (107.7 mM NaCl, 7.16 mM

KCl, 1.19 mM KH_2PO_4, 0.49 mM, $MgCl_2$, 1.17 mM $CaCl_2$, 5.3 mM sodium lactate, 25.07 mM $NaHCO_3$,

0.20 mM sodium pyruvate, 0.5 mM fructose, 1X non-essential amino acids (NEAA), 5 μg/ml gentamicin,

10 μg/ml heparin, 6 mg/ml fatty acid-free (FFA) BSA) and transferred to drops of SOF-IVF medium

under mineral oil. Frozen semen from a Holstein bull was thawed, and 10^6 sperm/ml were added to drops

with MII oocytes, which were incubated at 38.5°C for 12-18 h. Zygotes were then removed from

fertilization medium, and cumulus cells were removed by vortexing for 5 min in SOF-Hepes medium.

Zygotes were then transferred to culture media (KSOMaa supplemented with 4mg/mL BSA) under

mineral oil, and incubated at 38.5°C in 5% CO_2, 5% O_2, and 90% N_2. If embryos were to be

transcriptionally inhibited, the culture medium was supplemented with α-amanitin (50 μg/ml) on day-one.

On day-three, the culture medium was supplemented with 5% stem-cell qualified FBS (Gemini Bio 100-

525). Blastocyst development was evaluated at 7 days post-insemination (dpi).

Collection of oocytes and preimplantation embryos for ATAC-seq

Oocytes and embryos were collected for ATAC-seq library preparation from three separate collections

per developmental stage. Embryos intended for collection at the 2-cell, 4-cell, or 8-cell stages were

divided into two groups, one of which was supplemented with α-amanitin, and cultured simultaneously.

Germinal vesicle-stage oocytes were collected for ATAC-seq prior to maturation. Preimplantation

embryos were collected at the 2-cell (30-32 h post-insemination), 4-cell (2 dpi), 8-cell (3 dpi), and morula

stages (5 dpi).

ATAC-seq library preparation

Oocytes or embryos (a minimum of 500 cells) were treated with pronase (10 mg/ml) to completely

remove the *zona pellucida* and washed with SOF-Hepes on a warming plate. Cells were then transferred

to 1 ml cold SOF-Hepes, and centrifuged at 500 rcf, 4°C, for 5 min. Morulae were vortexed for 3 min in

cold ATAC-seq lysis buffer (10 mM Tris-Cl, pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$ and 0.1% IGEPAL CA-

630). Cell pellets were then resuspended in 1 ml cold ATAC-seq cell lysis buffer and centrifuged at 500

rcf, 4°C, for 10 min. Nuclear pellets were then resuspended in 50 µl transposition reaction mix (25 µl TD

buffer (Nextera DNA Library Prep Kit, Illumina), 2.5 µl TDE1 enzyme (Nextera DNA Library Prep Kit,

Illumina), 22.5 µl nuclease-free H_2O) and incubated for 60 min at 37°C, shaking at 300 rpm. The

transposase, which is loaded with Illumina sequencing adapters, cuts DNA where it is not sterically

hindered and simultaneously ligates adapters, effectively producing a library in one incubation step.

Transposed DNA was purified with the MinElute PCR purification kit (Qiagen, Hilden, Germany) and

eluted in 10 µl buffer EB. Libraries were then PCR amplified: 50 µl reactions (25 µl SsoFast Evagreen

supermix with low ROX (Bio-Rad, Hercules, CA), 0.6 µl 25 µM custom Nextera PCR primer 1, 0.6 µl 25

µM custom Nextera PCR primer 2 (with barcode; Supplementary Table 2), 13 µl nuclease-free H_2O, and

10 µl eluted DNA) were cycled as follows: 72°C for 5 min, 98°C for 30 s, and then thermocycling at

98°C for 10 s, 63°C for 30 s and 72°C for 1 min. Libraries from GV oocytes, 2-cell, and 4-cell embryos

were thermocycled for 13 cycles, and 8-cell and morulae libraries were thermocycled for 11 cycles. PCR-

amplified libraries were again purified with the MinElute PCR purification kit (Qiagen, Hilden, Germany)

and eluted in 10 µl buffer EB. Libraries were then evaluated for DNA concentration and nucleosomal

laddering patterns using the Bioanalyzer 2100 DNA High Sensitivity chip (Agilent, Palo Alto, CA).

Expected nucleosomal laddering was evidenced by the presence of both small fragments, corresponding

to hyper-accessible DNA that was frequently transposed, and larger fragments, corresponding to DNA

that was wrapped around one or more nucleosomes. This study focused on mapping open chromatin;

therefore, the sub-nucleosomal length fraction of each library (150-250 bp) was size selected using the

PippinHT system (Sage Science, Beverly, MA) with 3% agarose cassettes. Size-selected libraries were

run on a Bioanalyzer DNA High Sensitivity chip to confirm size-selection and determine DNA

concentration. Final libraries were then pooled for sequencing on the Illumina NextSeq platform to

generate 40 bp paired end reads.

ATAC-seq read alignment and peak calling

Raw sequencing reads were trimmed with Trim_Galore, a wrapper around Cutadapt (v0.4.0) (Martin,

2011), to remove residual Illumina adapter sequences and low quality (q<20) ends, keeping unpaired

reads and reads 10 bp or longer after trimming. Trimmed reads were then aligned to either the GRCm38

(mouse), GRCh38 (human), or ARS-UCD1.2 (cattle) assemblies using BWA aln (-q 15) and sampe (Li &

Durbin, 2010). PCR duplicates were removed with PicardTools, and mitochondrial and low-quality

alignments (q<15) were removed with SAMtools (Li *et al*, 2009). Alignments from biological replicates

from each stage were merged and randomly subsampled to equivalent depth with SAMtools for detection

of open chromatin. To determine which regions of the genome demonstrated significant enrichment of

ATAC-seq signal, broad peaks were called with MACS2 (Zhang *et al*, 2008), using a q-value cutoff of

0.05, and settings *--nomodel --shift -100 --extsize 200*.

RNA-seq alignment and gene expression quantification

Raw sequencing reads were trimmed with Trimmomatic (v0.33) (Bolger *et al*, 2014). Low-quality leading

and trailing bases (3 bases) were clipped, and illumine adapter sequences were removed, allowing 2 seed

mismatches, a palindrome clip threshold of 30, and a simple clip threshold of 10. Sliding window

trimming was conducted with a window size of 4 bases, and a quality threshold of 15. Reads 36 bases or

longer were retained after trimming. Trimmed reads were aligned to either the GRCm38 (mouse),

GRCh38 (human), or ARS-UCD1.2 (cattle) assemblies with STAR (v2.7.2a) with options –

outFilterScoreMinOverLread 0.85 and –seedSearchStartLmax 30. Low quality alignments (q<5) were

removed with SAMtools. Raw counts were calculated for genes in the Ensembl 96 annotations for each

species with summarizeOverlaps, from the R package GenomicAlignments (v1.18.1) (Lawrence *et al*,

2013), using 'Union' mode and allowing fragments for paired end data. Gene counts were MLE-

normalized using the DESeq2 R package (v1.22.2) (Anders & Huber, 2010) and submitted to the variance

stabilizing transformation for some analyses. DESeq2 was also used for differential expression analysis,

with genes demonstrating a logFC > 2 and an adjusted p-value < 0.05 considered differentially expressed.

Repeat expression quantification

Trimmed reads were aligned to the ARS-UCD1.2 genome assembly with STAR (v2.7.0) (Dobin *et al*,

2013) with options –outFilterMultimapNmax 100, --winAnchorMultimapNmax 100, and --twopassMode

Basic. Raw expression values for individual repetitive elements were calculated for repeats in the

RepeatMasker annotation for the ARS-UCD1.2 assembly (download from the UCSC Genome Browser)

using TEtoolkit (Lerat *et al*, 2017) in 'multi' mode, which improves quantification of transposable

elements transcripts by including ambiguously mapped reads. Raw expression values were MLE-

normalized using the DESeq2 R package (v1.22.2).

Comparison of replicate libraries and ATAC-seq and RNA-seq signal at regions of interest

For both ATAC-seq and RNA-seq data, alignments were converted to bigwig format using bamCoverage

from the DeepTools suite (Ramírez *et al*, 2014), which binned the genome into 50 bp windows and

calculated normalized signal (reads per kilobase million; RPKM) in each window. The plotPCA function

from DeepTools was then used to generate the PCA plot, with options *--transpose and --log2.* The

plotCorrelation function from DeepTools was used to calculate the Spearman correlation coefficient

between replicate libraries, based on genome-wide normalized coverage. To assess average accessibility

or expression at genomic intervals of interest, average ATAC-seq or RNA-seq normalized signal was

visualized using the Deeptools plotHeatmap function

Comparison and classification of ATAC-seq peaks

Peak sets from different stages were compared using the BEDtools intersect function (Quinlan & Hall,

2010), requiring a minimum of 1bp overlap to consider a peak shared by both sets. Similarly, peaks were

classified as genic if they overlapped either the 2 kb region upstream of a transcription start site (TSS),

exons, or introns by 1 bp. Otherwise, peaks were considered intergenic.

Motif enrichment

Genomic regions were evaluated for binding motif enrichment using the findMotifsGenome.pl script from

HOMER (v4.8) (Heinz *et al*, 2010), using the exact sizes of the input genomic intervals (-size given). The

most significant known or *de novo* motifs were reported, based on p-value. Known motifs that matched

significantly enriched *de novo* motifs were reported if their match score exceeded 0.6.

Genome-wide motif location prediction

Position-weight matrices were downloaded from the JASPAR database for TFs of interest (Fornes *et al*,

2019). Using the FIMO tool from the MEME suite (v5.0.4) (Bailey *et al*, 2009), TF motif locations (p <

1e-4) were predicted genome-wide in the ARS-UCD1.2 genome assembly.

Repeat class enrichment in genomic intervals

To determine if repetitive elements (either individual elements, families, or classes) were enriched in open

chromatin, the number of ATAC-seq peaks overlapping a set of repetitive elements was compared to

randomized intervals (ATAC-seq peak locations shuffled with BEDtools shuffle function) overlapping

the same set of repetitive elements, yielding a log ratio of random to observed.

Functional annotation enrichment analysis

Gene sets were submitted to DAVID (v6.8) (Huang *et al*, 2009a, 2009b) to identify enriched biological

functions using medium stringency. Gene ontology terms with an FDR < 0.05 were reported.

Acknowledgements

This work was supported by a grant from NIH-NICHD (R01HD070044) to P.J.R. and R.M.S. Semen

used for *in vitro* embryo production was provided by Semex (Madison, WI). M.M.H. was supported by

USDA NIFA National Need Fellowship #20143842021796 and an Austin Eugene Lyons Fellowship. The

authors thank the entire Ross laboratory for contributions to embryo production and lively discussion.

Author contributions

Conceived and designed the experiments: M.M.H., R.M.S. and P.J.R. Performed the experiments: X.M.

and M.M.H. Wrote the manuscript: M.M.H., R.M.S. and P.J.R.

Conflict of interest

The authors declare no competing financial interests.

Data availability

The following published NGS data sets were used, and accessed through the NCBI GEO repository: for

bovine oocytes and *in vitro* produced embryos, raw RNA-seq data were downloaded from accession

number GSE52415 (Graf *et al*, 2014). For mouse and human preimplantation embryo ATAC-seq and

RNA-seq data, raw sequencing files were downloaded from accession numbers GSE66390 (Wu *et al*,

2016) and GSE101571 (Wu *et al*, 2018), respectively. The ATAC-seq data produced in this study is

available via the NCBI SRA repository.

References

Abe K-I, Funaya S, Tsukioka D, Kawamura M, Suzuki Y, Suzuki MG, Schultz RM & Aoki F (2018)

Minor zygotic gene activation is essential for mouse preimplantation development. *Proc. Natl.*

Acad. Sci. U. S. A. **115:** E6780–E6788

Abe K, Yamamoto R, Franke V, Cao M, Suzuki Y, Suzuki MG, Vlahovicek K, Svoboda P, Schultz RM

& Aoki F (2015) The first murine zygotic transcription is promiscuous and uncoupled from splicing

and 3' processing. *Embo j* **34:** 1523–1537

Acampora D, Di Giovannantonio LG & Simeone A (2013) Otx2 is an intrinsic determinant of the

embryonic stem cell state and is required for transition to a stable epiblast stem cell condition.

Development **140:** 43–55 Available at: https://www.ncbi.nlm.nih.gov/pubmed/23154415

Adelson DL, Raison JM & Edgar RC (2009) Characterization and distribution of retrotransposons and

simple sequence repeats in the bovine genome

Ahmed K, Dehghani H, Rugg-Gunn P, Fussner E, Rossant J & Bazett-Jones DP (2010) Global chromatin

architecture reflects pluripotency and lineage commitment in the early mouse embryo. *PLoS One* **5:**

e10531–e10531 Available at: https://www.ncbi.nlm.nih.gov/pubmed/20479880

Anders S & Huber W (2010) Differential expression analysis for sequence count data. *Genome Biol.* **11:**

R106 Available at: https://doi.org/10.1186/gb-2010-11-10-r106

Bai H, Sakurai T, Godkin JD & Imakawa K (2013) Expression and potential role of GATA factors in

trophoblast development. *J. Reprod. Dev.* **59:** 1–6 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/23428586

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW & Noble WS (2009)

MEME Suite: tools for motif discovery and searching. *Nucleic Acids Res.* **37:** W202–W208

Available at: https://doi.org/10.1093/nar/gkp335

Bakhtari A & Ross PJ (2014) DPPA3 prevents cytosine hydroxymethylation of the maternal pronucleus

and is required for normal development in bovine embryos. *Epigenetics* **9:** 1271–1279 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/25147917

Barnes FL & First NL (1991) Embryonic transcription in in vitro cultured bovine embryos. *Mol. Reprod.*

Dev. **29:** 117–123 Available at: https://www.ncbi.nlm.nih.gov/pubmed/1878221

Beraldi R, Pittoggi C, Sciamanna I, Mattei E & Spadafora C (2006) Expression of LINE-1 retroposons is

essential for murine preimplantation development. *Mol Reprod Dev* **73:** 279–287

Bestor TH & Bourc'his D (2004) Transposon silencing and imprint establishment in mammalian germ

cells. *Cold Spring Harb. Symp. Quant. Biol.* **69:** 381–387 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/16117671

Bogliotti YS, Chung N, Paulson E, Chitwood J, Halstead M, Kern C, Schultz RM & Ross PJ (2019)

Transcript profiling of bovine embryos implicates specific transcription factors in the maternal-to-

embryo transition. *Biol. Reprod.* Available at: https://doi.org/10.1093/biolre/ioz209

Bolger AM, Lohse M & Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **30:** 2114–2120

Boroviak T, Stirparo GG, Dietmann S, Hernando-Herraez I, Mohammed H, Reik W, Smith A, Sasaki E,

Nichols J & Bertone P (2018) Single cell transcriptome analysis of human, marmoset and mouse

embryos reveals common and divergent features of preimplantation development. *Development* **145:**

dev167833 Available at: https://www.ncbi.nlm.nih.gov/pubmed/30413530

Braude P, Bolton V & Moore S (1988) Human gene expression first occurs between the four- and eight-

cell stages of preimplantation development. *Nature* **332:** 459–461

Buecker C, Srinivasan R, Wu Z, Calo E, Acampora D, Faial T, Simeone A, Tan M, Swigut T & Wysocka

J (2014) Reorganization of enhancer patterns in transition from naive to primed pluripotency. *Cell*

Stem Cell **14:** 838–853 Available at: https://www.ncbi.nlm.nih.gov/pubmed/24905168

Bui LC, Evsikov A V., Khan DR, Archilla C, Peynot N, Hénaut A, Le Bourhis D, Vignon X, Renard JP

& Duranthon V (2009) Retrotransposon expression as a defining event of genome reprograming in

fertilized and cloned bovine embryos. *Reproduction* **138:** 289–299

Bultman SJ, Gebuhr TC, Pan H, Svoboda P, Schultz RM & Magnuson T (2006) Maternal BRG1

regulates zygotic genome activation in the mouse. *Genes Dev* **20:** 1744–1754

Chen Z & Zhang Y (2019) Loss of DUX causes minor defects in zygotic genome activation and is

compatible with mouse development. *Nat. Genet.* **51:** 1

Chuong EB, Elde NC & Feschotte C (2017) Regulatory activities of transposable elements: from conflicts

to benefits. *Nat. Rev. Genet.* **18:** 71–86 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/27867194

Crosby IM, Gandolfi F & Moor RM (1988) Control of protein synthesis during early cleavage of sheep

embryos. *J. Reprod. Fertil.* **82:** 769–775

Davis DL (1985) Culture and storage of pig embryos. *J. Reprod. Fertil. Suppl.* **33:** 115–124 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/3910819

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M & Gingeras TR

(2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29:** 15–21

Du Z, Zheng H, Huang B, Ma R, Wu J, Zhang X, He J, Xiang Y, Wang Q, Li Y, Ma J, Zhang X, Zhang

K, Wang Y, Zhang MQ, Gao J, Dixon JR, Wang X, Zeng J & Xie W (2017) Allelic reprogramming

of 3D chromatin architecture during early mammalian development. *Nature* **547:** 232–235 Available

at: https://www.ncbi.nlm.nih.gov/pubmed/28703188

Eckersley-Maslin M, Alda-Catalinas C, Blotenburg M, Kreibich E, Krueger C & Reik W (2019) Dppa2

and Dppa4 directly regulate the Dux-driven zygotic transcriptional program. *Genes Dev.* **33:** 194–

208

Eckersley-Maslin MA, Alda-Catalinas C & Reik W (2018) Dynamics of the epigenetic landscape during

the maternal-to-zygotic transition. *Nat. Rev. Mol. Cell Biol.* **19:** 436–450

Falco G, Lee S-L, Stanghellini I, Bassey UC, Hamatani T & Ko MSH (2007) Zscan4: a novel gene

expressed exclusively in late 2-cell embryos and embryonic stem cells. *Dev. Biol.* **307:** 539–550

Available at: https://www.ncbi.nlm.nih.gov/pubmed/17553482

Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, Modi BP, Correard S,

Gheorghe M, Baranašić D, Santana-Garcia W, Tan G, Chèneby J, Ballester B, Parcy F, Sandelin A,

Lenhard B, Wasserman WW & Mathelier A (2019) JASPAR 2020: update of the open-access

database of transcription factor binding profiles. *Nucleic Acids Res.* Available at:

https://doi.org/10.1093/nar/gkz1001

Franke V, Ganesh S, Karlic R, Malik R, Pasulka J, Horvat F, Kuzman M, Fulka H, Cernohorska M,

Urbanova J, Svobodova E, Ma J, Suzuki Y, Aoki F, Schultz RM, Vlahovicek K & Svoboda P (2017)

Long terminal repeats power evolution of genes and gene expression programs in mammalian

oocytes and zygotes. *Genome Res.* **27:** 1384–1394

Frei RE, Schultz GA & Church RB (1989) Qualitative and quantitative changes in protein synthesis occur

at the 8-16-cell stage of embryogenesis in the cow. *J. Reprod. Fertil.* **86:** 637–641 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/2760892

Friedli M & Trono D (2015) The developmental control of transposable elements and the evolution of

higher species. *Annu. Rev. Cell Dev. Biol.* **31:** 429–451 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/26393776

Funaya S & Aoki F (2017) Regulation of zygotic gene activation by chromatin structure and epigenetic

factors. *J. Reprod. Dev.* **63:** 359–363 Available at: https://www.ncbi.nlm.nih.gov/pubmed/28579579

Gao L, Wu K, Liu Z, Yao X, Yuan S, Tao W, Yi L, Yu G, Hou Z, Fan D, Tian Y, Liu J, Chen Z-J & Liu J

(2018) Chromatin Accessibility Landscape in Human Early Embryos and Its Association with

Evolution. *Cell* **173:** 248-259.e15

Ghirlando R & Felsenfeld G (2016) CTCF: making the right connections. *Genes Dev* **30:** 881–891

Di Giammartino DC, Kloetgen A, Polyzos A, Liu Y, Kim D, Murphy D, Abuhashem A, Cavaliere P,

Aronson B, Shah V, Dephoure N, Stadtfeld M, Tsirigos A & Apostolou E (2019) KLF4 is involved

in the organization and regulation of pluripotency-associated three-dimensional enhancer networks.

Nat. Cell Biol. **21:** 1179–1190 Available at: https://doi.org/10.1038/s41556-019-0390-6

Graf A, Krebs S, Heininen-Brown M, Zakhartchenko V, Blum H & Wolf E (2014) Genome activation in

bovine embryos: review of the literature and new insights from RNA sequencing experiments. *Anim*

Reprod Sci **149:** 46–58

Graf T (2011) Historical origins of transdifferentiation and reprogramming. *Cell Stem Cell* **9:** 504–516

Available at: https://www.ncbi.nlm.nih.gov/pubmed/22136926

Guo M, Zhang Y, Zhou J, Bi Y, Xu J, Xu C, Kou X, Zhao Y, Li Y, Tu Z, Liu K, Lin J, Yang P, Gao S &

Wang Y (2019) Precise temporal regulation of Dux is important for embryo development. *Cell Res.*

Available at: https://doi.org/10.1038/s41422-019-0238-4

Hancks DC & Kazazian Jr HH (2016) Roles for retrotransposon insertions in human disease. *Mob. DNA*

7: 9 Available at: https://www.ncbi.nlm.nih.gov/pubmed/27158268

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H & Glass CK

(2010) Simple combinations of lineage-determining transcription factors prime cis-regulatory

elements required for macrophage and B cell identities. *Mol Cell* **38:** 576–589

Hendrickson PG, Doráis JA, Grow EJ, Whiddon JL, Lim J-W, Wike CL, Weaver BD, Pflueger C, Emery

BR, Wilcox AL, Nix DA, Peterson CM, Tapscott SJ, Carrell DT & Cairns BR (2017) Conserved

roles of mouse DUX and human DUX4 in activating cleavage-stage genes and MERVL/HERVL

retrotransposons. *Nat. Genet.* **49:** 925–934 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28459457

Hernandez C, Wang Z, Ramazanov B, Tang Y, Mehta S, Dambrot C, Lee Y-W, Tessema K, Kumar I,

Astudillo M, Neubert TA, Guo S & Ivanova NB (2018) Dppa2/4 Facilitate Epigenetic Remodeling

during Reprogramming to Pluripotency. *Cell Stem Cell* **23:** 396-411.e8 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/30146411

Huang DW, Sherman BT & Lempicki RA (2009a) Systematic and integrative analysis of large gene lists

using DAVID bioinformatics resources. *Nat. Protoc.* **4:** 44–57 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/19131956

Huang DW, Sherman BT & Lempicki RA (2009b) Bioinformatics enrichment tools: paths toward the

comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37:** 1–13 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/19033363

De Iaco A, Planet E, Coluccio A, Verp S, Duc J & Trono D (2017) DUX-family transcription factors

regulate zygotic genome activation in placental mammals. *Nat. Genet.* **49:** 941–945 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28459456

Iaco A De, Verp S, Offner S & Trono D (2019) DUX is a non-essential synchronizer of zygotic genome

activation. *bioRxiv***:** 569434 Available at:

http://biorxiv.org/content/early/2019/03/06/569434.abstract

Iturbide A & Torres-Padilla M-E (2017) Starting embryonic transcription for the first time. *Nat. Genet.*

49: 820–821 Available at: https://www.ncbi.nlm.nih.gov/pubmed/28546569

Iurlaro M, von Meyenn F & Reik W (2017) DNA methylation homeostasis in human and mouse

development. *Curr. Opin. Genet. Dev.* **43:** 101–109

Jachowicz JW, Bing X, Pontabry J, Bošković A, Rando OJ & Torres-Padilla M-E (2017) LINE-1

activation after fertilization regulates global chromatin accessibility in the early mouse embryo. *Nat.*

Genet. **49:** 1502 Available at: https://doi.org/10.1038/ng.3945

Jimenez R, Melo EO, Davydenko O, Ma J, Mainigi M, Franke V & Schultz RM (2015) Maternal SIN3A

regulates reprogramming of gene expression during mouse preimplantation development. *Biol.*

Reprod. **93:** 89 Available at: https://www.ncbi.nlm.nih.gov/pubmed/26353893

Ko MSH (2016) Zygotic Genome Activation Revisited: Looking Through the Expression and Function of

Zscan4. *Curr. Top. Dev. Biol.* **120:** 103–124 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/27475850

Koehler D, Zakhartchenko V, Froenicke L, Stone G, Stanyon R, Wolf E, Cremer T & Brero A (2009)

Changes of higher order chromatin arrangements during major genome activation in bovine

preimplantation embryos. *Exp. Cell Res.* **315:** 2053–2063 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/19254712

Kruse K, Díaz N, Enriquez-Gasca R, Gaume X, Torres-Padilla M-E & Vaquerizas JM (2019)

Transposable elements drive reorganisation of 3D chromatin during early embryogenesis. *bioRxiv*:

523712 Available at: http://biorxiv.org/content/early/2019/01/17/523712.abstract

Latham KE, Garrels JI, Chang C & Solter D (1991) Quantitative analysis of protein synthesis in mouse

embryos. I. Extensive reprogramming at the one- and two-cell stages. *Development* **112:** 921 LP –

932 Available at: http://dev.biologists.org/content/112/4/921.abstract

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT & Carey VJ (2013)

Software for computing and annotating genomic ranges. *PLoS Comput. Biol.* **9:** e1003118–

e1003118 Available at: https://www.ncbi.nlm.nih.gov/pubmed/23950696

Leidenroth A & Hewitt JE (2010) A family history of DUX4: phylogenetic analysis of DUXA, B, C and

Duxbl reveals the ancestral DUX gene. *BMC Evol. Biol.* **10:** 364 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/21110847

Lerat E, Fablet M, Modolo L, Lopez-Maestre H & Vieira C (2017) TEtools facilitates big data expression

analysis of transposable elements and reveals an antagonism between their activity and that of

piRNA genes. *Nucleic Acids Res.* **45:** e17–e17 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28204592

Li H & Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics **26:** 589–595

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R &

Subgroup 1000 Genome Project Data Processing (2009) The Sequence Alignment/Map format and

SAMtools. *Bioinformatics* **25:** 2078–2079 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/19505943

Li L, Guo F, Gao Y, Ren Y, Yuan P, Yan L, Li R, Lian Y, Li J, Hu B, Gao J, Wen L, Tang F & Qiao J

(2018) Single-cell multi-omics sequencing of human early embryos. *Nat. Cell Biol.* **20:** 847–858

Available at: https://doi.org/10.1038/s41556-018-0123-2

Liu L, Leng L, Liu C, Lu C, Yuan Y, Wu L, Gong F, Zhang S, Wei X, Wang M, Zhao L, Hu L, Wang J,

Yang H, Zhu S, Chen F, Lu G, Shang Z & Lin G (2019) An integrated chromatin accessibility and

transcriptome landscape of human pre-implantation embryos. *Nat. Commun.* **10:** 364 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/30664750

Liu X, Wang C, Liu W, Li J, Li C, Kou X, Chen J, Zhao Y, Gao H, Wang H, Zhang Y, Gao Y & Gao S

(2016) Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation

embryos. *Nature* **537:** 558–562

Lu F, Liu Y, Inoue A, Suzuki T, Zhao K & Zhang Y (2016) Establishing Chromatin Regulatory

Landscape during Mouse Preimplantation Development. *Cell* **165:** 1375–1388

Lubelsky Y, Reuven N & Shaul Y (2005) Autorepression of Rfx1 Gene Expression: Functional

Conservation from Yeast to Humans in Response to DNA Replication Arrest. *Mol. Cell. Biol.* **25:**

10665–10673 Available at: http://mcb.asm.org/

Marin M, Karis A, Visser P, Grosveld F & Philipsen S (1997) Transcription Factor Sp1 Is Essential for

Early Embryonic Development but Dispensable for Cell Growth and Differentiation. *Cell* **89:** 619–

628 Available at: http://www.sciencedirect.com/science/article/pii/S0092867400802433

Martens JH, O'Sullivan RJ, Braunschweig U, Opravil S, Radolf M, Steinlein P & Jenuwein T (2005) The

profile of repeat-associated histone lysine methylation states in the mouse epigenome. *Embo j* **24:**

800–812

Martin C, Beaujean N, Brochard V, Audouard C, Zink D & Debey P (2006) Genome restructuring in

mouse embryos during reprogramming and early development. *Dev. Biol.* **292:** 317–332 Available

at: http://www.sciencedirect.com/science/article/pii/S0012160606000157

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011* **17:**

Available at: http://journal.embnet.org/index.php/embnetjournal/article/view/200

Nishikimi A, Mukai J & Yamada M (1999) Nuclear translocation of nuclear factor kappa B in early 1-cell

mouse embryos. *Biol. Reprod.* **60:** 1536–1541 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/10330116

Oldfield AJ, Yang P, Conway AE, Cinghu S, Freudenberg JM, Yellaboina S & Jothi R (2014) Histone-

821  fold domain protein NF-Y promotes chromatin accessibility for cell type-specific master

822  transcription factors. *Mol. Cell* **55:** 708–722 Available at:

823  https://www.ncbi.nlm.nih.gov/pubmed/25132174

824 Ooga M, Fulka H, Hashimoto S, Suzuki MG & Aoki F (2016) Analysis of chromatin structure in mouse

825  preimplantation embryos by fluorescent recovery after photobleaching. *Epigenetics* **11:** 85–94

826  Available at: https://www.ncbi.nlm.nih.gov/pubmed/26901819

827 Paciolla M, Boni R, Fusco F, Pescatore A, Poeta L, Ursini M V, Lioi MB & Miano MG (2011) Nuclear

828  factor-kappa-B-inhibitor alpha (NFKBIA) is a developmental marker of NF-kappaB/p65 activation

829  during in vitro oocyte maturation and early embryogenesis. *Hum. Reprod.* **26:** 1191–1201

830 Payer B, Saitou M, Barton SC, Thresher R, Dixon JPC, Zahn D, Colledge WH, Carlton MBL, Nakano T

831  & Surani MA (2003) Stella is a maternal effect gene required for normal early development in mice.

832  *Curr. Biol.* **13:** 2110–2117 Available at: https://www.ncbi.nlm.nih.gov/pubmed/14654002

833 Perea-Gomez A, Lawson KA, Rhinn M, Zakin L, Brûlet P, Mazan S & Ang SL (2001) Otx2 is required

834  for visceral endoderm movement and for the restriction of posterior signals in the epiblast of the

835  mouse embryo. *Development* **128:** 753–765 Available at:

836  https://www.ncbi.nlm.nih.gov/pubmed/11171400

837 Pontis J, Planet E, Offner S, Turelli P, Duc J, Coudray A, Theunissen TW, Jaenisch R & Trono D (2019)

838  Hominoid-Specific Transposable Elements and KZFPs Facilitate Human Embryonic Genome

839  Activation and Control Transcription in Naive Human ESCs. *Cell Stem Cell* **24:** 724-735.e5

840 Probst A V, Okamoto I, Casanova M, El Marjou F, Le Baccon P & Almouzni G (2010) A strand-specific

841  burst in transcription of pericentric satellites is required for chromocenter formation and early mouse

842  development. *Dev Cell* **19:** 625–638

843 Probst A V, Santos F, Reik W, Almouzni G & Dean W (2007) Structural differences in centromeric

844  heterochromatin are spatially reconciled on fertilisation in the mouse zygote. *Chromosoma* **116:**

845  403–415 Available at: https://www.ncbi.nlm.nih.gov/pubmed/17447080

846 Quinlan AR & Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features.

Bioinformatics **26:** 841–842 Available at: https://www.ncbi.nlm.nih.gov/pubmed/20110278

Ram PT & Schultz RM (1993) Reporter gene expression in G2 of the 1-cell mouse embryo. *Dev. Biol.*

156: 552–556 Available at: https://www.ncbi.nlm.nih.gov/pubmed/8462750

Ramírez F, Dündar F, Diehl S, Grüning BA & Manke T (2014) deepTools: a flexible platform for

exploring deep-sequencing data. *Nucleic Acids Res.* **42:** W187–W191 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/24799436

Raviram R, Rocha PP, Luo VM, Swanzey E, Miraldi ER, Chuong EB, Feschotte C, Bonneau R & Skok

JA (2018) Analysis of 3D genomic interactions identifies candidate host genes that transposable

elements potentially regulate. *Genome Biol.* **19:** 216 Available at: https://doi.org/10.1186/s13059-

018-1598-7

Ross PJ, Ragina NP, Rodriguez RM, Iager AE, Siripattarapravat K, Lopez-Corrales N & Cibelli JB

(2008) Polycomb gene expression and histone H3 lysine 27 trimethylation changes during bovine

preimplantation development. *Reproduction* **136:** 777–785

Schoorlemmer J, Pérez-Palacios R, Climent M, Guallar D & Muniesa P (2014) Regulation of Mouse

Retroelement MuERV-L/MERVL Expression by REX1 and Epigenetic Control of Stem Cell

Potency. *Front. Oncol.* **4:** 14 Available at: https://www.ncbi.nlm.nih.gov/pubmed/24567914

Schulz KN & Harrison MM (2019) Mechanisms regulating zygotic genome activation. *Nat. Rev. Genet.*

20: 221–234

Shirai A, Kawaguchi T, Shimojo H, Muramatsu D, Ishida-Yonetani M, Nishimura Y, Kimura H,

Nakayama J-I & Shinkai Y (2017) Impact of nucleic acid and methylated H3K9 binding activities of

Suv39h1 on its heterochromatin assembly. *Elife* **6:** e25317 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28760201

Soufi A, Garcia MF, Jaroszewicz A, Osman N, Pellegrini M & Zaret KS (2015) Pioneer transcription

factors target partial DNA motifs on nucleosomes to initiate reprogramming. *Cell* **161:** 555–568

Available at: https://www.ncbi.nlm.nih.gov/pubmed/25892221

Spadafora C (2004) Endogenous reverse transcriptase: a mediator of cell proliferation and differentiation.

Cytogenet. Genome Res. **105:** 346–350 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/15237222

Takahashi K, Ross PJ & Sawai K (2019) The necessity of ZSCAN4 for preimplantation development and

gene expression of bovine embryos. *J. Reprod. Dev.* **65:** 319–326 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/31019155

Takahashi K & Yamanaka S (2006) Induction of pluripotent stem cells from mouse embryonic and adult

fibroblast cultures by defined factors. *Cell* **126:** 663–676 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/16904174

Vastenhouw NL, Cao WX & Lipshitz HD (2019) The maternal-to-zygotic transition revisited.

Development **146:**

Whiddon JL, Langford AT, Wong C-J, Zhong JW & Tapscott SJ (2017) Conservation and innovation in

the DUX4-family gene network. *Nat. Genet.* **49:** 935–940 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28459454

White CR, MacDonald WA & Mann MRW (2016) Conservation of DNA Methylation Programming

Between Mouse and Human Gametes and Preimplantation Embryos. *Biol. Reprod.* **95:** 61

Wiekowski M, Miranda M & DePamphilis ML (1993) Requirements for promoter activity in mouse

oocytes and embryos distinguish paternal pronuclei from maternal and zygotic nuclei. *Dev. Biol.*

159: 366–378 Available at: https://www.ncbi.nlm.nih.gov/pubmed/8365573

Wu J, Huang B, Chen H, Yin Q, Liu Y, Xiang Y, Zhang B, Liu B, Wang Q, Xia W, Li W, Li Y, Ma J,

Peng X, Zheng H, Ming J, Zhang W, Zhang J, Tian G, Xu F, et al (2016) The landscape of

accessible chromatin in mammalian preimplantation embryos. *Nature* **534:** 652–657

Wu J, Xu J, Liu B, Yao G, Wang P, Lin Z, Huang B, Wang X, Li tong, Shi S, Zhang N, Duan F, Ming J,

Zhang X, Niu W, Song W, Jin H, Guo Y, Dai S, Hu L, et al (2018) Chromatin analysis in human

early development reveals epigenetic transition during ZGA. Available at:

https://doi.org/10.1038/s41586-018-0080-8

Wu X, Li Y, Xue L, Wang L, Yue Y, Li K, Bou S, Li G-P & Yu H (2011) Multiple histone site epigenetic

modifications in nuclear transfer and in vitro fertilized bovine embryos. *Zygote* **19:** 31–45 Available

at: https://www.ncbi.nlm.nih.gov/pubmed/20609268

Xu Q & Xie W (2018) Epigenome in Early Mammalian Development: Inheritance, Reprogramming and

Establishment. *Trends Cell Biol.* **28:** 237–253 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/29217127

Yandım C & Karakülah G (2019) Expression dynamics of repetitive DNA in early human embryonic

development. *BMC Genomics* **20:** 439 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/31151386

Yang Y, Zhou C, Wang Y, Liu W, Liu C, Wang L, Liu Y, Shang Y, Li M, Zhou S, Wang Y, Zeng W,

Zhou J, Huo R & Li W (2017) The E3 ubiquitin ligase RNF114 and TAB1 degradation are required

for maternal-to-zygotic transition. *EMBO Rep.* **18:** 205–216 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/28073917

Young JM, Whiddon JL, Yao Z, Kasinathan B, Snider L, Geng LN, Balog J, Tawil R, van der Maarel SM

& Tapscott SJ (2013) DUX4 binding to retroelements creates promoters that are active in FSHD

muscle and testis. *PLoS Genet.* **9:** e1003947–e1003947 Available at:

https://www.ncbi.nlm.nih.gov/pubmed/24278031

Zhang A, Xu B, Sun Y, Lu X, Gu R, Wu L, Feng Y & Xu C (2012) Dynamic changes of histone H3

trimethylated at positions K4 and K27 in human oocytes and preimplantation embryos. *Fertil. Steril.*

98: 1009–1016

Zhang B, Zheng H, Huang B, Li W, Xiang Y, Peng X, Ming J, Wu X, Zhang Y, Xu Q, Liu W, Kou X,

Zhao Y, He W, Li C, Chen B, Li Y, Wang Q, Ma J, Yin Q, et al (2016) Allelic reprogramming of

the histone modification H3K4me3 in early mammalian development. *Nature* **537:** 553–557

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M,

Li W & Liu XS (2008) Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9:** R137

Figure legends

Figure 1. Chromatin accessibility in bovine oocytes and in vitro preimplantation embryos. A) Schematic

of in vitro embryo production and ATAC-seq library preparation. B) Principal components analysis

(PCA) of ATAC-seq read depth normalized by reads per kilobase million (RPKM) in 50 bp windows

covering the whole genome. C) Normalized coverage (RPKM) of replicate ATAC-seq libraries for each

stage of development. D) Proportion of genome that is covered by genic and intergenic ATAC-seq peaks,

called from 30 million reads at each developmental stage. E) Categorization of accessible regions in 2-, 4-

, and 8-cell embryos into stage-specific and maintained peaks (accessibility maintained up until the

morula stage). Maintained peaks are carried over from latter stages to show cumulative maintained peaks.

Figure 2. Gradual establishment of open chromatin enriched in regulatory motifs. A) Regions that

significantly lost accessibility from the GV oocyte to 2-cell stage, and regions that significantly gained

accessibility from the 2- to 4-cell, 4- to 8-cell, and 8-cell to morula stages, according to 30 million reads

per stage. Accessibility at each region, scaled to average width $\pm$ 1kb, was determined by normalized

ATAC-seq read depth (reads per kilobase million; RPKM), based on 20 million reads per stage. B) Top

five enriched *de novo* motifs enriched in intergenic peaks with matching known motifs (match score >

0.6) of TFs that were expressed at the given stage. C) Top seven known motifs enriched in 4-cell specific

peaks. D) DESeq2 normalized expression of TFs corresponding to enriched motifs in 4-cell specific open

chromatin. E) Top seven known motifs enriched in 8-cell specific peaks. F) DESeq2 normalized

expression of TFs corresponding to enriched motifs in 8-cell specific open chromatin. G) Proportion of

intergenic 8-cell peaks overlapping the de novo motif most closely matching the DUXA motif, relative to

background. H) DESeq2-normalized expression of DUXA throughout development. RNA-seq data from

Graf *et al* (2014).

Figure 3. Binding motif enrichment in open chromatin opened at the 2-, 4-, or 8-cell stages and

maintained up to morula stage. Proportion of peaks with A) CTCF or B) KLF5, binding motifs. C)

DESeq2-normalized expression of KLF genes. RNA-seq data from Graf *et al* (2014). D) Proportion of

maintained and stage-specific peaks that were genic or intergenic. Maintained peaks were established at

the indicated stage and maintained up to the morula stage. E) Gene ontology term enrichment of the 9,456

genes that were marked by maintained genic open chromatin that was first established in 2- or 4-cell

embryos.

Figure 4. Effect of transcription inhibition on chromatin remodeling. A) Transcription inhibition effect on

loci that should have opened or closed between consecutive stages. Peaks called based on 20 million

reads per stage and categorized as either genic or intergenic. B) Normalized read depth (RPKM) at loci

which opened during the 4- to 8-cell transition in control 8-cell embryos and transcriptionally inhibited 8-

cell embryos. Regions scaled to average width ± 1 kb. C) Normalized ATAC-seq and RNA-seq signal

(RPKM) in 8-cell and 8-cell TBE at the KLF4 locus. RNA-seq data from Bogliotti *et al* (2019). D)

Proportion of stage-specific and maintained peaks that should appear at each stage, but which fail to open

in TBE.

Figure 5. Expression and accessibility dynamics of repeat families during bovine preimplantation

development. A) DESeq2 normalized expression profiles of select LTR, LINE, and SINE repeat families.

RNA-seq data from Graf *et al* (2014). B) Enrichment of several transposable element (TE) families in

ATAC-seq peaks, called from 30M reads. Promoter peaks fell within the 2kb region upstream of TSS.

Intergenic peaks did not overlap the 2kb regions upstream of TSS, exons, or introns.

Figure 6. Activity of LTR elements. A) Open chromatin enrichment of LTR elements with increased

accessibility in 4-cell, 8-cell, and/or morulae, and VST-normalized expression of the same LTR elements;

three replicates per stage. B) ATAC-seq and RNA-seq read coverage at a highly expressed ERV1-2 repeat

sequence. ATAC-seq tracks show coverage from 30 million reads per library; RNA-seq tracks show

combined coverage from three replicates. TF motifs predicted from JASPAR binding motifs MA0039.3

(KLF4) and MA0712.1 (OTX2), respectively. C) Average normalized ATAC-seq and D) RNA-seq signal

(RPKM) at MLT1A0 repeats overlapped by 8-cell intergenic open chromatin harboring DUXA motifs,

predicted based on JASPAR motif MA0468.1. E) Co-option of an accessible MLT1A0 element with a

DUXA binding motif as an alternative promoter upstream of the C1D locus. RNA-seq data from Graf *et*

al (2014).

Figure 7. Inference of key regulatory factors during EGA in cattle, human, and mouse, based on

enrichment for TF binding factors in open chromatin and expression of the corresponding TF. Bovine

RNA-seq data from Graf *et al* (2014); human ATAC-seq and RNA-seq data from Wu *et al* (2018); mouse

ATAC-seq and RNA-seq from Wu *et al* (2016).

Figure 8. Potential mechanistic model depicting events leading to major EGA. Chromatin structure is

globally decondensed following fertilization, allowing opportunistic binding of maternal factors, which

initiate a minor wave of transcription and begin to establish 3D chromatin architecture. This sets the stage

for major EGA, wherein maternal products, minor EGA products, and promoter-enhancer contacts

mediate the first major wave of gene expression and continue to refine 3D chromatin structure.

1000

1001

1002

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Figure 7.

Figure 8.