Honeybees adjust colour preferences in response to concurrent social information

from conspecifics and heterospecifics

Jose E Romero-Gonzalez*, Cwyn J Perry, Lars Chittka

Biological and Experimental Psychology, School of Biological and Chemical Sciences,

Queen Mary University London, London, U.K.

*Correspondence: J.E. Romero-Gonzalez, School of Biological and Chemical Sciences,

Queen Mary University of London, London E1 4NS, UK.

E-mail address: j.e.romerogonzalez@qmul.ac.uk

Abstract

Bees efficiently learn asocial and social cues to optimize foraging from fluctuating

floral resources. However, it remains unclear how bees respond to divergent sources of

social information, and whether such social cues might modify bees' natural preferences

for asocial cues (e.g. flower colour), hence affecting foraging decisions. Here, we

investigated honeybees' (*Apis mellifera*) inspection and choices of unfamiliar flowers

based on both natural colour preferences and simultaneous foraging information from

conspecifics and heterospecifics. Individual honeybees' preferences for flowers were

recorded when the reward levels of a learned flower type have declined and novel-

coloured flowers were available where they would find either no social information or

one conspecific and one heterospecific (*Bombus terrestris*), each foraging from a

different coloured flower (either magenta or yellow). Honeybees were found to have a

natural preference for magenta flowers. Social information affected honeybees'

inspection time of both types of flowers, i.e., honeybees modified their approaching

flights to yellow and magenta flowers in response to conspecific and heterospecific

social information. The presence of either demonstrator on the less-preferred yellow

flower increased honeybees' inspection time of yellow flowers. Conspecific social

information influenced observers' foraging choices of yellow flowers, thus outweighing

their original preference for magenta flowers, such an influence was not elicited by

heterospecific social information. Our results indicate that flower colour preferences of

honeybees are rapidly adjusted in response to conspecific social information, which in

turn is preferred over heterospecific information, thus favouring the transmission of

adaptive foraging information within species.

Keywords: behavioural flexibility, decision making, innate colour bias, insects,

intraspecific, interspecific, social information

Foraging decisions are central to an animal's survival and reproduction; deciding

where to forage in unpredictably changing environments is a major challenge that

animals constantly encounter (Stephens et al. 2007). Foraging decisions can be

influenced in a context-dependant manner (Laland 2004; Kendal et al. 2009) by innate

preferences (Raine et al. 2006), previous individual experience (Sclafani 1995), and the

observation or interaction with another animals at a foraging resource, i.e., social

information (Heyes 1994; Hoppitt and Laland 2013). For example, naïve individuals

tend to rely more on social than individual information to gain familiarity about

foraging sources (Galef and Giraldeau 2001; Galef and Laland 2005). Conversely,

experienced individuals that have information on an advantageous foraging resource,

will often ignore social cues. Thus, social information is only draw upon when

individual information has become outdated and acquiring up to date information may

be costly (Laland 2004; Galef and Laland 2005; Kendal et al. 2009). The influence of

social information on foraging decisions is taxonomically widespread (Galef and

Giraldeau 2001; Valone and Templeton 2002; Grüter and Leadbeater 2014). Social

information allows animals to find profitable foraging resources efficiently, instead of

iteratively sampling the environment through trial and error (Galef and Laland 2005).

In most animal communities, multiple species frequently share the same

foraging resources; thus members of the same (conspecific) and different

(heterospecific) species can potentially act as sources and users of social information

(Seppänen et al. 2007; Goodale et al. 2010; Avarguès-Weber et al. 2013; Parejo and

Avilés 2016). Using social information indiscriminately is not adaptive but animals

should be selective when acquiring information from other individuals (Laland 2004;

Kendal et al. 2018). In a multi-species context, animals have access to social

information from different sources, which may give rise to a trade-off between selecting

sources with a close ecological similarity, implying high competition, or sources whose

ecological distance may involve less competition but a lower informative value

(Seppänen et al. 2007). Furthermore, in a foraging context, social information has to be

integrated with unlearned preferences (Laland and Plotkin 1993; Leadbeater and Chittka

2007a; Jones et al. 2015) and existing individual information (van Bergen Yfke et al.

2004; Jones et al. 2015) to determine foraging decisions in a context-dependant manner

(Laland 2004; Kendal et al. 2009).

82   Social bee workers forage for nectar and pollen from rapidly changing floral

resources (Heinrich 1979; Chittka et al. 1999) within multi-species communities (Fægri

and Van der Pijl 1979; Kevan and Baker 1983) that might offer a wide spectrum of

social information. This makes bees a valuable model to explore how different sources

of social information can affect learned and unlearned preferences of individuals to

shape foraging decisions. Traditionally, the laboratory paradigms that investigate the

use of social information in bees tend to oversimplify the real field contexts where

animals naturally acquire information from others. For example, they test bees in

relatively unnatural settings where a single source of social information is presented,

e.g., a dead demonstrator pinned to simulated flowers (reviewed in Leadbeater and

Dawson 2017). Contrastingly, bees seeking nectar and pollen in the wild might

encounter far more complex circumstances where they have access to multiple sources

of social information (Fægri and Van der pijl 1979; Kevan and Baker 1983) that may

concur in time and space and diverge in their intrinsic relevance.

96   Honeybees and bumblebees are sympatric species that typically forage upon

similar flowers due to their generalist diet (Rogers et al. 2013; Xie et al. 2016). Previous

evidence indicates that bumblebees can acquire foraging information from dead

demonstrator honeybees (Dawson and Chittka 2012). However, in more realistic

conditions, a bee forager whose known floral resources have decreased in reward levels,

will likely concur with other foragers, of their own and different species, feeding from

different types of flowers (Fægri and Van der Pijl 1979; Kevan and Baker 1983). It

remains unclear how bees respond to such divergent social information, and whether in

this context, social cues might modify bees' natural preferences for particular colours in

flowers (Chittka et al. 2004; Raine et al. 2006; Raine and Chittka 2007), hence affecting

foraging decisions. We address these questions by testing whether honeybees might

adjust their colour preferences in response to simultaneous sources of social

information, i.e., a conspecific and heterospecific, each foraging from either a preferred

or non-preferred flower type.

MATERIALS AND METHODS

(a) Set-up

We used free-flying honeybee foragers (*Apis mellifera*) of two different colonies

(Colonies A and B) for experiments. Over two consecutive summers (2017-2018), we

trained foragers (one colony per year) to collect 30% sucrose solution (w/w) from a

gravity feeder located 2 m away from their hive. The gravity feeder attracted foragers

from neighboring hives, thus individuals for experiments could have come from at least

three additional hives. In addition, we used bumblebee (*Bombus terrestris*) foragers

from two colonies (Biobest, Belgium N.V.) for the experiments. Bumblebee nests were

housed in bipartite wooden nest-boxes ($l = 29.5 \times w = 11.5 \times h = 9.5$ cm) connected to a

wooden flight arena ($l = 77$ cm, $w = 52$ cm, $h = 30$ cm) by a Plexiglas tunnel ($l = 25$ cm,

3.5×3.5 cm). The floor of the flight arena was covered in white laminated paper. Three

plastic sliding doors located along the corridor allowed controlled access to the arena.

Before and after experiments, bumblebees could freely feed upon 30% (w/w) sucrose

solution from a mass feeder in the middle of the arena. Bumblebee colonies were

provided with 7 g of frozen pollen (Koppert B.V., The Netherlands) every two days.

(b) Training of honeybees

Training of demonstrators and observers took place over 18 daily sessions. Each

day, five foraging honeybees (demonstrators) from the gravity feeder were lured to and

trained to enter the wooden flight arena located outdoors. The five honeybees were

group trained, i.e. allowed to enter the arena together at various times, to learn to forage

upon an array of twelve plastic chips (2.4 × 2.4 cm, henceforth "flowers") placed on the

top of transparent glass vials (h = 4 cm), positioned on the floor of the arena. The array

was arranged in a rectangular grid formation (3 × 4) with a separation of 7.5 cm

between the edges of the flowers. We filled four yellow flowers with 10 µL of

rewarding 50% (w/w) sucrose solution, and four magenta and four transparent flowers

with 10 µL of unrewarding water (Figure 1A). Flowers were refilled after they were

depleted by foragers and the foragers left the flower. Half of the foragers experienced

the opposite color-reinforcement contingency. We identified trained individuals by

marking their thorax with a white paint mark (Posca Pen, Worcester, UK). Three hours

after concluding the demonstrator's training, we selected a separate batch of five

foragers (observers) that we trained them on a different set-up. Every day, observers

were group trained to enter the same arena and forage upon a rectangular grid array of

twelve transparent flowers (3 × 4) (Figure 1C). To encourage honeybee foragers to

sample multiple flowers during their foraging trips, only half of the flowers contained

40 µL of 50% (w/w) sucrose solution, whereas the other half contained 40 µL of water.

Rewarding flowers were replenished after emptied and the forager left the flower, and

the positions of all the flowers were changed every 10 minutes over 1 hour of training.

We marked trained observers with a green dot to distinguish them from the

demonstrators. After each training session, we cleaned the flowers and the arena floor

with 70% ethanol.

(c) Training of bumblebees

In the same flight arena where honeybees were trained, bumblebee foragers

(demonstrators) from two colonies were separately trained in group. In this training, the

colour-reinforcement contingency was reversed so that bumblebees learned to forage on

flowers of the opposite colour that honeybees were trained on. Half of the bumblebees

experienced four magenta flowers with 40 μL of 50% (w/w) sucrose solution, and four

yellow and four transparent flowers with 40 μL of water (Figure 1B). The other half of

the bumblebees experienced the reverse color-reinforcement contingency. Rewarding

flowers were refilled after depleted by the foragers and the foragers left the flower.

Training took place for one day over 2 hours. We also carried out a refreshment bout

(20 min) each day before testing. We tracked demonstrator bees' identity with

individual number tags (Opalithplättchen, Warnholz & Bienenvoigt, Ellerau, Germany)

glued to the top of the thorax by means of Loctite Super Glue Gel (Loctite, Ohio, US).

The floor of the arena and flowers were cleaned with 70% ethanol after completing

training.

(d) Testing the effect of color preference on honeybees' foraging decisions

After each training session, we selected two honeybees trained on transparent

flowers, twenty-one subjects overall, and evaluate them individually in a control test

where transparent flowers were unrewarding to assess whether they might show a

preference to explore and forage from one flower type between two unfamiliar

alternatives, i.e., yellow and magenta. One honeybee was let in the arena to explore a

rectangular grid array of twelve flowers (section b, above), consisting of four familiar

transparent flowers containing 20 μL of water and eight unfamiliar flowers, four yellow

and four magenta, all filled with a scentless reward of 20 μL of 50% (w/w) sucrose

solution, (Figure 1D). The test began once the individual inspected any flower. That is,

the honeybee approached the flower with its head oriented towards it and displayed a

slow side-to-side hover within at least one body length (Ings et al. 2009). Individuals

were given up to 3 min to land and forage upon any unfamiliar flower, before the test

concluded. As this test was designed to evaluate the influence of honeybees' colour

preferences on an actual foraging decision, rather than measuring their innate colour

preferences, we regarded the flower type where the individual landed and foraged as

preferred over the alternative type. To prevent re-testing the same individuals, we

captured honeybees, after concluding the test, to give them a distinctive red paint mark.

The flowers and arena floor were cleaned with 70% ethanol between tests. To evaluate

honeybees' inspection of flowers before they chose a flower to forage (foraging

decision), we recorded the test with a sport camera (Yi, Xiaomi Inc. China) featuring a

recording frame rate of 30 fps and a resolution of 720 p (1,280 × 720 pixels). The

camera was positioned 20 cm above the entrance of the bumblebee nest (Figure 1). Its

field of view was adjusted such that it looked down into the arena at ~50° from a

horizontal angle.

(e) Testing the influence of social information on foraging decisions

To evaluate the influence of simultaneous sources (conspecific and

heterospecific) of social information on honeybee's inspection of either yellow or

magenta flowers and their subsequent foraging decisions, we tested fifty-three

honeybees in the same context as individuals in the control group (section d), but in the

presence of one bumblebee and one honeybee demonstrator, each foraging upon either a

yellow or magenta flower. We introduced the demonstrators and observer in the arena

in the following order: 1) a honeybee demonstrator was let in through a sliding door on

the back wall of the arena (Figure 1D). Once she began to feed from a yellow flower

and started to forage, a bumblebee demonstrator was released from the Plexiglas tunnel

that connected the arena to the nest-box (Figure 1D). When she landed and began to

feed from a magenta flower, a honeybee observer was let into the arena through the

sliding door on the back wall. This color-reinforcement contingency was

counterbalanced for thirty-three observers. The test began once the observer inspected

(see section d) any occupied or unoccupied flower. Observers were given up to 3 min to

select and forage upon any unfamiliar flower before the test concluded. In the test,

demonstrators moved freely between flowers after depleting them, as they naturally do

within inflorescences. To prevent re-testing the same individuals, we caught tested

honeybees and marked them with a distinctive red paint mark. The flowers and arena

floor were cleaned with 70% ethanol between tests. To evaluate honeybees' inspection

of flowers and their interactions with the demonstrators before they made a foraging

decision, the test was recorded as described in section (d) above.

Figure 1. Training and testing protocols in the flight arena. (**A**) Honeybee foragers (demonstrators) were trained to find 50% sucrose solution on either four yellow or magenta flowers and water on four flowers of the alternative colour and four transparent flowers. Bumblebee foragers (demonstrators) were group trained using the same protocol (**C**) A different batch of honeybee foragers (observers) was group trained to find 50% sucrose solution on six of twelve transparent flowers and unrewarding water on the other six transparent flowers. (**D**) Honeybee observers were tested on their preparedness to forage on an unfamiliar coloured flower type, depending on its colour (yellow or magenta), once their learned flower type (transparent) yielded no reward. (**E**) Two demonstrators (one honeybee and one bumblebee) and one observer honeybee were sequentially introduced in the flight arena: (1) A honeybee demonstrator was let in the flight arena, once she foraged from a flower of her trained colour, (2) a bumblebee demonstrator was released into the arena and let to forage from a flower of her trained colour (different colour from the honeybee demonstrator), (3) an observer honeybee was then introduced to test her in a context where the flowers she was trained to associate with reward (transparent) were unrewarding and unfamiliar colored flowers were demonstrated by a conspecific and a heterospecific (bumblebee) demonstrator, each foraging from non-preferred (yellow) and preferred (magenta) flowers, this design was counterbalanced.

(d) Analyses

We analysed the behaviour of tested honeybees from video recordings, using the

BORIS behavioural observation software (Friard and Gamba, 2016). To assess

individuals' inspection of either yellow or magenta flowers and their preference to

forage upon one type of flower over the other, we analysed two main behavioural

categories, the frequency of inspecting transparent, yellow and magenta flowers (Ings et

al. 2009: 200; Balamurali et al. 2018) and individual's foraging choice, i.e., the yellow

or magenta flower where individuals landed and foraged. For honeybees in the control

group (no social information), we used a Chi-squared test to compare the proportion of

honeybees that foraged upon either a magenta or yellow flower, as well as the

likelihood of first inspecting one of these types of flower. Further, we compared the

proportion of instances that honeybees inspected a transparent, yellow or magenta

flower, using a Friedman test and a Wilcoxon signed rank test for *post hoc* comparisons.

For honeybees exposed to social information, we considered inspection of

occupied flowers as an indicator that observers detected the demonstrator's presence.

Thereby, four different scenarios were possible before honeybees made a foraging

decision: they could have detected both demonstrators, either the honeybee or

bumblebee demonstrator, or none of them. We compared the likelihood of each

situation against the expected probability with a Chi-Square Goodness of Fit Test. Nine

individuals that did not detect the presence of the demonstrators were not considered for

all analyses.

To contrast the influence of individual's color preferences and simultaneous,

conspecific and heterospecific, social information on foraging decisions of honeybees,

we compared the proportion of individuals that foraged on an unfamiliar flower

between the control group and the group exposed to social information. Further, to

evaluate the influence of social information on honeybees' readiness to forage on an

unfamiliar flower, we compared two measurements between the control group and the

group exposed to social information – the total number of flowers that individuals

inspected before choosing a flower to forage, and the time it took them to make such a

decision (latency to forage). To explore whether such readiness could be affected by the

species of the first demonstrator that the observers detected, we compared, with a

Wilcoxon rank sum test, the aforementioned measurements between the observers that

first detected the presence of either a honeybee or bumblebee foraging on a flower,

irrespective of whether they detected both or only one demonstrator during the test.

To determine whether observers that detected both demonstrators inspected the

flowers occupied by a honeybee and bumblebee at a similar frequency, we compared,

with a Wilcoxon signed rank test, the proportion of flowers occupied by each

demonstrator that observers inspected. We also analysed, with a Wilcoxon rank sum

test, whether the honeybees that only detected either a conspecific or heterospecific

demonstrator, differed in the proportion of occupied flowers that they inspected, in

relation to all the flowers they inspected during the test. To compare the specific

influence of conspecific and heterospecific social information on honeybees' foraging

decisions, we analysed, with a Chi-Squared test, the proportion of individuals that

selected the flower type demonstrated by either a honeybee or bumblebee, as well as the

likelihood that observers would select a flower occupied by either a honeybee or

bumblebee after inspecting it for the first time.

The instances in which observers inspected an occupied flower that did not

progress into landing and foraging were recorded as rejections. We evaluated whether

observers rejected the flowers occupied by a honeybee or bumblebee demonstrator at a

similar frequency. For observers that detected both demonstrators, we compared the

proportion of times that they rejected the flowers occupied by each demonstrator, using

a Wilcoxon signed rank test. We also compared, with a Wilcoxon rank sum test, the

total number of occupied flowers that observers rejected when they only detected either

the honeybee or bumblebee demonstrator.

292   To assess whether conspecific and heterospecific social information affected

293   honeybees' inspection of yellow or magenta flowers, we compared the proportion of

294   inspecting transparent, yellow and magenta flowers between honeybees in the control

295   group and the group that had access to social information.

296   To estimate whether the observers altered their inspection of flowers, after they

297   detected the presence of either a foraging honeybee or bumblebee, we adjusted the

298   frequency of observers' inspection of flowers by normalising the data through an index

299   calculation, with the following equation:

300
$$\text{Inspecting Index (II)} = \frac{i_m - i_y}{i_m + i_y}$$

302   , where, i_m and i_y are the frequency that the observer inspected magenta and yellow

303   flowers, respectively.

304   In the indices, a negative value (minimum -1) equates to a preference to inspect

yellow flowers, whereas a positive value (maximum +1) equates to a preference to

inspect magenta flowers, an index near equal to zero implies that the observers either

inspected both types of flowers equally, or they did not inspect the flowers at all. We

calculated inspecting indices (II) to represent the events that preceded the honeybees'

foraging decisions. That is, each approaching index (AI) represented observers'

inspection of flowers, before and after they detected either demonstrator (honeybee and

bumblebee) foraging on a flower . We compared indices against chance expectation

(Index = 0) with a Wilcoxon signed-rank test. A significant switch from a negative or

positive value, before the observer detected a demonstrator, to a positive or negative

value, after this occurred, indicates that observers modified their inspection of yellow or

magenta flowers in response to conspecific or heterospecific social information. All

analyses were conducted using R statistical software (R Core Team, 2019).

RESULTS

Effect of color preference on honeybees' foraging decisions

In a context where a familiar flower type ceases to yield reward, how does

colour preference influence honeybees' exploration of contiguous, novel flowers and

ultimate move to foraging from a new flower type? We carried out a control test to

assess whether honeybees preferentially inspect one unfamiliar flower type over

another, and whether this might be reflected in their ultimate foraging decision (move to

foraging on a new flower type). We trained honeybees (observers) to find either a food

reward (50% sucrose solution) or unrewarding water on transparent flowers (Materials

and Methods), then we tested them in a context where transparent flowers were

unrewarding, and two unfamiliar yellow and magenta flower types, delivered an

undetectable food reward. The majority of individuals (77%) did not inspect the familiar

transparent flowers at all; rather, they inspected the magenta flowers more frequently

than the other types of flower (Friedman; X^2 (2) = 19.07, df = 2, N = 18, p = 7.231e-05;

post hoc, Wilcoxon signed rank; comparison between flower types: magenta *versus*

transparent: p = 0.002; magenta *versus* yellow: p = 0.0008; yellow *versus* transparent: p

= 0.67; figure 2A).

Similarly, honeybees were more likely to first inspect a magenta than a yellow

flower (Chi-square; chi = 14.23, N = 18, p = 1.6132e-04; figure 2B), and a higher

proportion of individuals selected magenta over yellow flowers to forage (Chi-square;

chi = 28.44, N = 18, p = 9.6426e-08; figure 2C). These results indicate that tested

individuals had a common preference for magenta over yellow flowers, which

influenced their inspection of flowers and foraging decisions.

Figure 2. Honeybees prefer magenta over yellow flowers **(A)** Honeybees inspected magenta flowers

more frequently than yellow and transparent flowers. **(B)** The likelihood of honeybees inspecting an

unfamiliar flower for the first time was higher for magenta compared to yellow flowers. **(C)** A higher

proportion of honeybees preferred to forage on magenta than yellow flowers. Bars = mean. Horizontal

lines = standard error. Asterisks = p < 0.05. n.s. = not significantly different.

Influence of social information on honeybees' foraging decisions

In a field-like context, including simultaneous and divergent conspecific and

heterospecific social information, we investigated how this social information might

affect honeybees' inspection of flower types and ultimate foraging decisions. We

considered that observers detected the demonstrators' presence once they inspected an

occupied flower for the first time. We found no difference in the proportion of observers

that detected both demonstrators, only the honeybee or bumblebee demonstrator, or

none of them (Chi-Square Goodness of Fit Test; χ^2= 2.2, df= 3, N = 45, p = 0.312;

Figure 3A). The proportion of observer honeybees that detected the foraging

demonstrators and then selected an unfamiliar flower to forage was not different from

individuals that selected unfamiliar flowers in the control group (Chi-square; $\chi^2 =$

1.022, N = 56, p = 0.312; Figure 3B). Similar to the control group, the vast majority of

observers (90%) did not inspect the transparent flowers at all. These results suggest that

both individual colour bias and social information similarly influenced honeybees when

locating a new foraging resource. Interestingly, honeybees in the control group whose

foraging decisions were solely influenced by their color preference, inspected a similar

number of flowers (Wilcoxon rank-sum; W = 125, N = 30, p= 0.447; Figure 3D and

made equally swift foraging decisions (Wilcoxon rank-sum; W = 133, N = 30, p=

0.299; Figure 3C) as honeybees exposed to social information that in the test, initially

detected the conspecific demonstrator foraging. However, when observer honeybees

first detected the heterospecific demonstrator foraging, they made slower foraging

decisions and inspected more flowers compared to both the control group (Wilcoxon

rank-sum; latency: W = 71, N = 35, p = 0.007; Figure 3C, number of flowers inspected:

W = 77, N = 35, p = 0.011; Figure 3D) and those honeybees that detected the

conspecific demonstrator first (Wilcoxon rank-sum; latency: W = 184, N = 29, p =

0.0003; Figure 3C, number of flowers inspected: W = 176.5, N = 29, p = 0.0009; Figure

3D). These results suggest that individual preference for magenta flowers and

conspecific social information similarly influenced honeybees' foraging choices which

allowed them to make fast decisions. This did not occur in response to heterospecific

social information as observers inspected the flowers extensively before making a

foraging decision.

Observer honeybees similarly inspected flowers occupied by a honeybee and

bumblebee demonstrator, regardless whether in the test they detected only one

demonstrator (Wilcoxon rank-sum; W = 34.5, N= 16; p-value = 0.449; Figure 3F) or

both demonstrators (Wilcoxon signed-rank; V = 22.5, N = 13, p = 1; Figure 3E).

However, a significantly higher proportion of honeybees selected and foraged on the

type of flowers demonstrated by conspecifics over the flowers demonstrated by

heterospecifics (Chi-square; χ^2 = 17.15, N = 29, p = 3.4516e-05; Figure 3G). Further,

observer honeybees that inspected an occupied flower for the first time, were

considerably more likely to forage on it if the occupier was a conspecific demonstrator

(Chi-square; χ^2 = 15.6508, N = 41, p = 7.6181e-05; Figure 3H). This evidence

demonstrates that foraging conspecifics had a strong influence on honeybees' foraging

decisions. Contrastingly, observer honeybees rejected flowers occupied by a bumblebee

more frequently than the flowers occupied by a conspecific in both situations, when

they detected either one (Wilcoxon rank-sum; W = 2, N = 16, p = 0.002; Figure 3I) or

both demonstrators (Wilcoxon signed-rank; V = 3, N = 13, p = 0.008; Figure 3J).

Figure 3. Concurrent conspecific and heterospecific social information influences foraging behaviour of

honeybees differently **(A)** Observer honeybees were equally likely to detect both demonstrators, only the

honeybee or bumblebee demonstrator, and none of them. **(B)** The proportion of honeybees that foraged

on unfamiliar flowers was not different between observers that detected the presence of demonstrators

and individuals in the control group (no social information). **(C)** Honeybees in the control group and

those that first detected a foraging conspecific made faster foraging decisions than observers that first

detected a foraging heterospecific. **(D)** Honeybees in the control group and those that first detected a

foraging conspecific inspected fewer flowers before making a foraging decision than observers that first

detected the presence of a foraging heterospecific. **(E)** Observers that detected both demonstrators during

the test inspected occupied flowers irrespective of the demonstrator's species. **(F)** Observers that only

detected either the conspecific or heterospecific demonstrator showed a similar proportion of inspecting

occupied flowers. **(G)** A higher proportion of observers preferred to forage on the type of flowers

demonstrated by conspecifics compared to the type of flowers demonstrated by heterospecifics. **(H)** Once

observers detected the presence of a conspecific demonstrator foraging on an unfamiliar flower type, they

were more likely to land and forage on this flower, but they never chose a flower occupied by a

heterospecific demonstrator. **(I)** Observers that only detected either a conspecific or heterospecific

demonstrator, rejected flowers occupied by a bumblebee more frequently than flowers occupied by a

conspecific. **(J)** Observers that detected both demonstrators rejected the flowers occupied by a bumblebee

at a higher proportion than the flowers occupied by a conspecific. HB = honeybee, BB = bumblebee. Bars

= mean. Horizontal lines = standard error. Asterisks = p < 0.05. n.s. = not significantly different.

Adjustment of honeybees' colour preference in response to social information

431    Our results indicate that social information can influence honeybee's foraging

decisions. But how does such information interact with an individual's colour

preference to shape honeybees' flower choices? Compared to individuals in the control

group, observers that detected a honeybee foraging on magenta flowers, and a

bumblebee on the yellow ones, inspected a similar proportion of magenta flowers

(Wilcoxon rank-sum test; W = 96, N = 29, p = 0.9029; Figure 4A) but they inspected

more yellow flowers (Wilcoxon rank-sum; W = 137.5, N = 29, p = 0.002; Figure 4B).

In the alternative scenario, where honeybees observed a conspecific and heterospecific

demonstrator, respectively foraging on a yellow and magenta flower, the proportion of

yellow flowers that observers inspected was markedly higher than individuals in the

control group (Wilcoxon rank-sum; W = 221.5, N = 29, p = 1.499e-05; Figure 4B);

hence they tended to inspect fewer magenta flowers, compared to the control group

(Wilcoxon rank-sum; $W = 49$, $N = 29$, $p = 0.095$; Figure 4A). These results

demonstrate that individuals in the control group and those exposed to social

information similarly inspected magenta flowers, regardless the species of the

demonstrator foraging on these flowers. Contrastingly, the presence of either

demonstrator foraging on yellow flowers, scarcely explored by honeybees in the control

group, attracted observers' attention to such type of flowers. Particularly, observers that

detected a conspecific foraging on a yellow flower, focused on inspecting these flowers,

so that their preference for magenta flowers was overridden. This evidence confirms

that even though honeybees' foraging decisions was determined by their preference for

magenta flowers in the absence of social information, this colour preference was clearly

outweighed by conspecific social information.

454   The inspecting indexes (II), described in (d), allowed us analysing honeybees'

inspection of flowers as a flexible process that culminated in a foraging decision, and

whose variation in response to social information was measurable in terms of an index

value. Observer honeybees showed no preference to inspect magenta or yellow flowers

before they detected the bumblebee demonstrator foraging on either a magenta

(Wilcoxon signed-rank test; $V = 20$, $N = 9$, $p = 0.149$; Figure 4D) or yellow flower

(Wilcoxon signed-rank test; $V = 12$, $N = 9$, $p = 0.112$; Figure 4D). After honeybees

detected a bumblebee demonstrator foraging on a magenta flower, they preferentially

inspected this type of flower (Wilcoxon signed-rank; $V = 26$, $N = 9$, $p = 0.026$; Figure

4A). However, after honeybees observed a bumblebee foraging on a yellow flower, they

showed no preference to inspect either type of flower (Wilcoxon signed-rank; $V = 18.5$,

$N = 9$, $p = 0.5$; Figure 4D). Honeybees did not prefer to inspect neither type of flower

before they detected a conspecific foraging upon either a magenta (Wilcoxon signed-

rank; $V = 16.5$, $N = 11$, $p = 0.612$; Figure 4E) or yellow flower (Wilcoxon signed-rank;

V = 23, N = 11, p = 0.261; Figure 4E). Honeybees responded to the presence of a

conspecific foraging on either a magenta or yellow flower by increasing their inspection

of demonstrated flowers (Wilcoxon signed-rank; magenta: V = 52, N = 11, p =

0.006471; yellow: V = 1, N = 13, p = 0.002; Figure 4E). These results show that

honeybees' inspection of the different types of flowers underwent a sequential

adjustment in response to conspecific and heterospecific social information, which

ultimately affected foraging decisions.

Figure 4. Honeybees adjust their exploration of flowers and foraging preferences in response to

simultaneous conspecific and heterospecific social information **(A)** Observer honeybees, previously

trained on transparent flowers, similarly inspected unfamiliar magenta flowers in the absence of social

information (control group) and when these flowers were demonstrated by either a conspecific or a

heterospecific (bumblebee) demonstrator. **(B)** In comparison to individuals in the control group, observers

inspected more yellow flowers in response to the presence of either a conspecific or heterospecific

demonstrator foraging on this flower type. **(C)** and **(D)** Inspecting indices (II) compare to chance level

(Index = 0) the inspection of magenta (dark gray bars) and yellow (light gray bars) flowers that observers

displayed before and after detecting either a conspecific or heterospecific demonstrator foraging on either

a magenta or yellow flower. **(C)** Before observers detected a bumblebee demonstrator foraging on either

a magenta or yellow flower, they showed no preference to inspect neither of these flower types. After

observers detected a bumblebee demonstrator foraging on a magenta flower, they inspected more

frequently this type of flowers, but when observers detected the bumblebee demonstrator foraging on a

yellow flower, they did not alter their inspection of yellow or magenta flowers. **(D)** Before observers

detected a conspecific foraging upon either a yellow or magenta flower, they did not preferentially inspect

magenta or yellow flowers. After observers detected a conspecific demonstrator foraging on either a

yellow or magenta flower, they increased their inspection of the demonstrated flower type. HB =

honeybee, BB = bumblebee. Index = 0. Bars = mean. Horizontal lines = standard error. Asterisks = p <

0.05. n.s. = not significantly different between groups or from chance level.

DISCUSSION

The results presented here provide evidence that honeybees'colour preferences

can be adjusted in response to simultaneous social information from conspecific and

heterospecific sources, with conspicuous effects on honeybees' exploration of flowers

and consequent foraging decisions. Unexpectedly, honeybees payed little attention to

the transparent flowers on which they were trained, instead prior to making a foraging

decision, they inspected more frequently the magenta flowers (control group) or the

type of flower demonstrated by a conspecific.

Despite inspection of flowers provide foragers with no information on the

reward status of a flower, they are important in the process of choosing a flower (Lunau

et al. 1996). In the absence of social information, honeybees' natural preference for

magenta flowers influenced inspection and choices of flowers . Magenta is actually a

mixture of blue and red, since the red component is not fully perceived by bees (Menzel

and Shmida 1993), they perceive magenta flowers as blue (Waser and Chittka 1998).

Honeybees and bumblebees usually have innate biases for colours in the violet to blue

range of the spectrum (Giurfa et al. 1995; Chittka et al. 2004), which adaptively

correlates with the nectar production of local flowers (Giurfa et al. 1995; Chittka et al.

2004). Although colour biases can somewhat be overridden by individual learning

(Raine et al., 2006), they potentially govern foraging decisions in bees when selecting

among novel flower types (Gumbert 2000). In the absence of social information, flower

choices of tested honeybees conceivably resulted from either their innate colour

preferences or previous experience in the field with local flower species as both factors

can be linked with the likelihood of finding a profitable food reward (Giurfa et al. 1995;

Chittka et al. 2004).

In the wild, honeybee foragers are likely to encounter members of the same and

different species foraging concurrently in a flower patch from distinct flower types

(Fægri and Van der Pijl 1979; Kevan and Baker 1983). Our results demonstrate that

such social information is certainly integrated with honeybees' colour preferences

during the process of making a foraging decision. In this realistic scenario, with two

simultaneous sources of conspecific and heterospecific social information, when the

first foraging demonstrator that honeybees detected was a member of the same species,

they promptly responded to social information by joining the demonstrator on the

unfamiliar flower type. This evidence is consistent with previous findings in

bumblebees (reviewed in Leadbeater and Dawson 2017) and supports the notion that the

use of conspecific social information in bees may be mediated by local enhancement,

with foragers being attracted to flowers occupied by members of the same species

(Leadbeater and Chittka 2007a). Such a response potentially optimises the foraging

efficiency of bees by facilitating their movements between different flower types

(Leadbeater and Chittka 2005; Kawaguchi et al. 2007).

Honeybees and bumblebees often forage upon similar floral resources; it is then

conceivable that they may concur in the same flower patches (Rogers et al. 2013; Xie et

al. 2016). In our experiments, we used freely moving bumblebee demonstrators to

elucidate how social information might flow between these species. Our results indicate

that in the likely scenario of a honeybee exploring a flower patch and encountering both

a conspecific and heterospecific (bumblebee) individual foraging from different flower

types, its foraging decisions might depend on the species identity of the first source of

social information that is observed. That is, foraging decisions of observer honeybees

that detected a bumblebee before a conspecific demonstrator were characteristically

preceded by an extensive exploration of the flowers. This evidence suggests that in a

foraging context, honeybees' discrimination of conspecifics and heterospecifics might

determine their readiness to use social information in making foraging decisions.

Theory predicts that using social information from heterospecifics should be favourable

when there is a large niche overlap (Seppänen et al. 2007) yet, on adaptive grounds,

animals should typically favour conspecific over heterospecific social information when

they can select between both sources (Jaakkonen et al. 2015). In line with this,

honeybees consistently selected the type of flowers demonstrated by conspecifics, even

though they attended to the presence of both demonstrators whilst exploring the set-up.

Prioritizing conspecific over heterospecific choices can be explained due to its high

fitness value (Seppänen et al. 2007; Jaakkonen et al. 2015) but it can also serve to the

transmission of novel behavioural traits or preferences that may be adaptively valuable

for a particular species (Laland and Plotkin 1993; Jaakkonen et al. 2015; Alem et al.

2016; Danchin et al. 2018).

Floral reward levels differ strongly among plant species and constantly change

over time in an unpredictable manner (Heinrich 1979). To achieve efficient foraging,

bees can rapidly learn to associate floral traits such as colour, shape and scent with

reward quality in flowers (Chittka et al. 1999). Bees can be initially attracted to forage

from an unfamiliar flower species via either innate and learned colour preferences

(Giurfa et al. 1995; Gumbert 2000; Chittka et al. 2004; Raine and Chittka 2007) or

using social information (Leadbeater and Chittka 2007b; Grüter and Leadbeater 2014;

Leadbeater and Dawson 2017). In the wild, bees seeking floral resources are unlikely to

experience asocial and social cues separately, rather they may frequently be exposed to

a complex combination of cues potentially affecting their flower choices. Our findings

demonstrate a clear interplay between colour preferences and social information

simultaneously delivered by conspecifics and heterospecifics. This may underlie the

process of gathering new information that shapes honeybees' foraging decisions.

Compared to honeybees in the control group (i.e., no social information) the exploration

behaviour of honeybees that observed a conspecific or heterospecific demonstrator

foraging upon the less preferred yellow flowers, reflected a more evenly distributed

inspection of magenta and yellow flowers. That is, the presence of either demonstrator

on a yellow flower increased the 'attractiveness' of such flower type for honeybees,

possibly via stimulus enhancement, an effect widely described in social learning

literature (Heyes 2012). Remarkably, honeybees' inspection of flowers, naturally biased

towards magenta flowers, underwent a sequential and flexible adjustment before they

made a foraging decision. Such adjustment was modulated by visual foraging

information from members of the same and different species, and resulting foraging

decisions reflected the selective use of conspecific over heterospecific social

information. It has been demonstrated that bees are attracted to the presence of foraging

conspecifics when presented with unfamiliar flowers which may lead them to

identifying new rewarding flower species (Leadbeater and Chittka 2007a; Jones et al.

2015). Our results indicate that conspecific social information not only directed

honeybees' foraging choices to flowers that matched their colour preference (magenta),

but it also outweighed this colour preference, such that honeybees selected the

originally less preferred yellow flowers. Whereas intraspecific social transmission of

foraging information may be more stable when it reinforces a prior preference (Laland

and Plotkin 1990), such preferences may be adjusted and potentially overridden in

response to conspecific social information about a novel foraging resource (Jones et al.

2015). Thus, enabling a fast adaptation to varying floral reward levels. If the presence of

a foraging conspecific can reliably be associated with a rewarding outcome, the use of

such social cue should be reinforced to continuously influence bees' flowers choices

across foraging contexts (Leadbeater and Chittka 2009). Interestingly, in our

experiments the presence of a foraging heterospecific attracted honeybees' attention to

the demonstrated flower types, yet this effect did not influence honeybees' foraging

decisions as they selectively used social information from conspecifics. Even though,

heterospecific social information is predictably valuable when there is a large niche

overlap (Seppänen et al. 2007), conspecifics choices might offer a more predictable

social cue that lowers the risk of acquiring maladaptive information (Giraldeau et al.

2002) as conspecifics share the same ecological needs (Seppänen et al. 2007; Goodale

et al. 2010).

Multiple social and asocial cues can potentially affect animals' foraging

decisions in different circumstances (Sclafani 1995; Galef and Giraldeau 2001; Grüter

and Leadbeater 2014). The interaction between social information and previous, innate

or learned preferences possibly not only influences the foraging decision of a particular

individual but may also promote the transmission of adaptive information about food

sources (Laland and Plotkin 1990; Galef and Giraldeau 2001). Despite social

information offers a clear advantage to individuals exploring novel foraging resources

(Galef and Giraldeau 2001), its use should not be indiscriminate but respond to

particular circumstances in order to lead to adaptive choices (Laland 2004; Kendal et al.

2018). Our results extend previous evidence showing that conspecific social information

is commonly used in situations of uncertainty (Kendal et al. 2004; van Bergen Yfke et

al. 2004; Galef et al. 2008; Smolla et al. 2016), such that it can outweigh both

predetermined individual preferences (Dugatkin 1996; Jones et al. 2015) and

heterospecific social information (Jaakkonen et al. 2015) to influence ecologically

relevant decisions in animals.

It is conceivable that natural selection should favour the salience of social

stimuli with high ecological relevance (Seppänen et al. 2007; Leadbeater and Dawson

2017), conspecifics, in our experiments. Whether such salience can similarly operate to

select information from heterospecific sources, based on their relative informative

value, deserves further consideration. By depicting in depth, the plausible ecological

process, thereby multiple sensory inputs may shape foraging decisions in bees, findings

presented here contribute to our current knowledge about the flexibility of individual

preferences and their adjustment in response to different types of social information.

Our results in turn shed light on the selectivity of animals for conspecific over

heterospecific information as a possible mechanism to facilitate the social transmission

of foraging information within species.

Acknowledgments

JERG was supported by CONACyT and QMUL, CVU 446980.

References

636 Alem S, Perry CJ, Zhu X, Loukola OJ, Ingraham T, Søvik E, Chittka L. 2016.
637 Associative Mechanisms Allow for Social Learning and Cultural Transmission of String
638 Pulling in an Insect. PLOS Biol. 14(10):e1002564. doi:10.1371/journal.pbio.1002564.
639
640 Avarguès-Weber A, Dawson EH, Chittka L. 2013. Mechanisms of social learning
641 across species boundaries. J Zool. 290(1):1–11. doi:10.1111/jzo.12015.
642
643 Balamurali GS, Nicholls E, Somanathan H, Hempel de Ibarra N. 2018. A comparative
644 analysis of colour preferences in temperate and tropical social bees. Sci Nat. 105(1):8.
645 doi:10.1007/s00114-017-1531-z.
646
647 Chittka L, Ings TC, Raine NE. 2004. Chance and adaptation in the evolution of island
648 bumblebee behaviour. Popul Ecol. 46(3):243–251. doi:10.1007/s10144-004-0180-1.
649
650 Chittka L, Thomson JD, Waser NM. 1999. Flower Constancy, Insect Psychology, and
651 Plant Evolution. Naturwissenschaften. 86(8):361–377. doi:10.1007/s001140050636.
652
653
654 Dugatkin LA. 1996. Interface between culturally based preferences and genetic
655 preferences: female mate choice in Poecilia reticulata. Proc Natl Acad Sci. 93(7):2770–
656 2773. doi:10.1073/pnas.93.7.2770.
657
658 Danchin E, Nöbel S, Pocheville A, Dagaeff A-C, Demay L, Alphand M, Ranty-Roby S,
659 Renssen L van, Monier M, Gazagne E, et al. 2018. Cultural flies: Conformist social
660 learning in fruitflies predicts long-lasting mate-choice traditions. Science.
661 362(6418):1025–1030. doi:10.1126/science.aat1590.
662
663 Dawson EH, Chittka L. 2012. Conspecific and Heterospecific Information Use in
664 Bumblebees. PLOS ONE. 7(2):e31444. doi:10.1371/journal.pone.0031444.
665
666 Fægri K, Van der Pijl L. 1979. Chapter 11 - Animals as pollinators. In: Fægri K, Van
667 der Pijl L, editors. Principles of Pollination Ecology (Third Edition). Amsterdam:
668 Pergamon. p. 96–133.
669
670 Galef BG, Dudley KE, Whiskin EE. 2008. Social learning of food preferences in
671 'dissatisfied' and 'uncertain' Norway rats. Anim Behav. 75(2):631–637.
672 doi:10.1016/j.anbehav.2007.06.024.
673
674 Galef BG, Giraldeau L-A. 2001. Social influences on foraging in vertebrates: causal
675 mechanisms and adaptive functions. Anim Behav. 61(1):3–15.
676 doi:10.1006/anbe.2000.1557.
677
678 Galef BG, Laland KN. 2005. Social Learning in Animals: Empirical Studies and
679 Theoretical Models. BioScience. 55(6):489–499. doi:10.1641/0006-
680 3568(2005)055[0489:SLIAES]2.0.CO;2.
681
Giraldeau L, Valone TJ, Templeton JJ. 2002. Potential disadvantages of using socially
acquired information. Johnstone RA, Dall SRX, editors. Philos Trans R Soc Lond B

Biol Sci. 357(1427):1559–1566. doi:10.1098/rstb.2002.1065.

Giurfa M, Núñez J, Chittka L, Menzel R. 1995. Colour preferences of flower-naive
honeybees. J Comp Physiol A. 177(3):247–259. doi:10.1007/BF00192415.

Goodale E, Beauchamp G, Magrath RD, Nieh JC, Ruxton GD. 2010. Interspecific
information transfer influences animal community structure. Trends Ecol Evol.
25(6):354–361. doi:10.1016/j.tree.2010.01.002.

Grüter C, Leadbeater E. 2014. Insights from insects about adaptive social information
use. Trends Ecol Evol. 29(3):177–184. doi:10.1016/j.tree.2014.01.004.

Gumbert A. 2000. Color choices by bumble bees (Bombus terrestris): innate preferences
and generalization after learning. Behav Ecol Sociobiol. 48(1):36–43.
doi:10.1007/s002650000213.

Heinrich B. 1979. Resource heterogeneity and patterns of movement in foraging
bumblebees. Oecologia. 40(3):235–245. doi:10.1007/BF00345321.

Heyes C. 2012. What's social about social learning? J Comp Psychol. 126(2):193–202.
doi:10.1037/a0025180.

Heyes CM. 1994 Social-learning in animals–categories and mechanisms. Biol. Rev.
Camb. Phil. Soc. 69, 207–231. doi:10.1111/j.1469- 185X.1994.tb01506.x

Hoppitt W, Laland KN. 2013. Social Learning: An Introduction to Mechanisms,
Methods, and Models. Princeton University Press.

Ings TC, Raine NE, Chittka L. 2009. A population comparison of the strength and
persistence of innate colour preference and learning speed in the bumblebee Bombus
terrestris. Behav Ecol Sociobiol. 63(8):1207–1218. doi:10.1007/s00265-009-0731-8.

Jaakkonen T, Kivelä SM, Meier CM, Forsman JT. 2015. The use and relative
importance of intraspecific and interspecific social information in a bird community.
Behav Ecol. 26(1):55–64. doi:10.1093/beheco/aru144.

Jones PL, Ryan MJ, Chittka L. 2015. The influence of past experience with flower
reward quality on social learning in bumblebees. Anim Behav. 101:11–18.
doi:10.1016/j.anbehav.2014.12.016.

Kawaguchi Lina G, Ohashi Kazuharu, Toquenaga Yukihiko. 2007. Contrasting
responses of bumble bees to feeding conspecifics on their familiar and unfamiliar
flowers. Proc R Soc B Biol Sci. 274(1626):2661–2667. doi:10.1098/rspb.2007.0860.

Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL. 2018. Social
Learning Strategies: Bridge-Building between Fields. Trends Cogn Sci. 22(7):651–665.
doi:10.1016/j.tics.2018.04.003.

Kendal RL, Coolen I, Laland KN. 2004. The role of conformity in foraging when
personal and social information conflict. Behav Ecol. 15(2):269–277.
doi:10.1093/beheco/arh008.
737
738 Kendal, RL, Coolen, I, Laland, KN. Adaptive trade-offs in the use of social and
739 personal information. In: Dukas R, Ratcliffe JM, editors. Cognitive ecology II. Chicago:
740 The University of Chicago Press. p. 249-271.
741
742 Kevan PG, Baker HG. 1983. Insects as Flower Visitors and Pollinators. Annu Rev
743 Entomol. 28(1):407–453. doi:10.1146/annurev.en.28.010183.002203.
744
745 Laland KN. 2004. Social learning strategies. Anim Learn Behav. 32(1):4–14.
746 doi:10.3758/BF03196002.
747
748 Laland KN, Plotkin HC. 1990. Social learning and social transmission of foraging
749 information in Norway rats (Rattus norvegicus). Anim Learn Behav. 18(3):246–251.
750 doi:10.3758/BF03205282.
751
752 Laland KN, Plotkin HC. 1993. Social transmission of food preferences among norway
753 rats by marking of food sites and by gustatory contact. Anim Learn Behav. 21(1):35–41.
754 doi:10.3758/BF03197974.
755
756 Leadbeater E, Chittka L. 2005. A new mode of information transfer in foraging
757 bumblebees? Curr Biol. 15(12):R447–R448. doi:10.1016/j.cub.2005.06.011.
758
759 Leadbeater E, Chittka L. 2007a. The dynamics of social learning in an insect model, the
760 bumblebee (Bombus terrestris). Behav Ecol Sociobiol. 61(11):1789–1796.
761 doi:10.1007/s00265-007-0412-4.
762 Leadbeater E, Chittka L. 2007b. Social Learning in Insects — From Miniature Brains to
763 Consensus Building. Curr Biol. 17(16):R703–R713. doi:10.1016/j.cub.2007.06.012.
764
765 Leadbeater E, Dawson EH. 2017. A social insect perspective on the evolution of social
766 learning mechanisms. Proc Natl Acad Sci. 114(30):7838–7845.
767 doi:10.1073/pnas.1620744114.
768
769 Leadbeater Ellouise, Chittka Lars. 2009. Bumble-bees learn the value of social cues
770 through experience. Biol Lett. 5(3):310–312. doi:10.1098/rsbl.2008.0692.
771
772 Lunau K, Wacht S, Chittka L. 1996. Colour choices of naive bumble bees and their
773 implications for colour perception. J Comp Physiol A. 178(4).
774 doi:10.1007/BF00190178.
775
776 Menzel R, Shmida A. 1993. The ecology of flower colours and the natural colour vision
777 of insect pollinators: the Israeli flora as a study case. Biol Rev. 68(1):81–120.
778 doi:10.1111/j.1469-185X.1993.tb00732.x.
779
780 Parejo D, Avilés JM. 2016. Social information use by competitors: Resolving the
781 enigma of species coexistence in animals? Ecosphere. 7(5):e01295.
782 doi:10.1002/ecs2.1295.
783

784   Raine NE, Chittka L. 2007. The Adaptive Significance of Sensory Bias in a Foraging
785   Context: Floral Colour Preferences in the Bumblebee Bombus terrestris. PLOS ONE.
786   2(6):e556. doi:10.1371/journal.pone.0000556.
787
788   Raine Nigel E., Ings TC, Dornhaus A, Saleh N, Chittka L. 2006. Adaptation, Genetic
789   Drift, Pleiotropy, and History in the Evolution of Bee Foraging Behavior. In: Advances
790   in the Study of Behavior. Vol. 36. Academic Press. p. 305–354.
791
792   Raine Nigel E, Ings TC, Ramos-Rodriguez O, Chittka L. 2006. Unterschiede im
793   Lernverhalten zwischen Kolonien einer freilebenden Britischen Hummelpopulation
794   (Hymenoptera: Apidae: Bombus terrestris audax). Entomol Gen. 28(4):241–256.
795   doi:10.1127/entom.gen/28/2006/241.
796
797   Rogers SR, Cajamarca P, Tarpy DR, Burrack HJ. 2013. Honey bees and bumble bees
798   respond differently to inter- and intra-specific encounters. Apidologie. 44(6):621–629.
799   doi:10.1007/s13592-013-0210-0.
800
801   Sclafani A. 1995. How food preferences are learned: laboratory animal models. Proc
802   Nutr Soc. 54(2):419–427. doi:10.1079/PNS19950011.
803
Seppänen J-T, Forsman JT, Mönkkönen M, Thomson RL. 2007. Social Information Use
Is a Process Across Time, Space, and Ecology, Reaching Heterospecifics. Ecology.
88(7):1622–1633. doi:10.1890/06-1757.1.
Smolla M, Alem S, Chittka L, Shultz S. 2016. Copy-when-uncertain: bumblebees rely
on social information when rewards are highly variable. Biol Lett. 12(6):20160188.
doi:10.1098/rsbl.2016.0188.
810
811   Stephens DW, Brown JS, Ydenberg RC, editors. 2007. Foraging: behavior and ecology.
812   Chicago: University of Chicago Press.
813
814   Valone TJ, Templeton JJ. 2002. Public information for the assessment of quality: a
815   widespread social phenomenon. Johnstone RA, Dall SRX, editors. Philos Trans R Soc
816   Lond B Biol Sci. 357(1427):1549–1557. doi:10.1098/rstb.2002.1064.
817
818   van Bergen Y, Coolen I, Laland KN. 2004. Nine-spined sticklebacks exploit the most
819   reliable source when public and private information conflict. Proc R Soc Lond B Biol
820   Sci. 271(1542):957–962. doi:10.1098/rspb.2004.2684.
821
822   Waser N, Chittka L. 1998. Bedazzled by flowers. Nature. 394(6696):835–836.
823   doi:10.1038/29657.
824
825   Xie Z, Pan D, Teichroew J, An J. 2016. The Potential Influence of Bumble Bee
826   Visitation on Foraging Behaviors and Assemblages of Honey Bees on Squash Flowers
827   in Highland Agricultural Ecosystems. PLOS ONE. 11(1):e0144590.
828   doi:10.1371/journal.pone.0144590.
829