Honeybees adjust colour preferences in response to concurrent social information

from conspecifics and heterospecifics

José E Romero-González[a]*, Cwyn Solvi[a,b], Lars Chittka[a]

[a] Biological and Experimental Psychology, School of Biological and Chemical

Sciences, Queen Mary University of London, London, U.K.

[b] Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109,

AU

*Correspondence: J.E. Romero-Gonzalez, School of Biological and Chemical Sciences,

Queen Mary University of London, London E1 4NS, UK.

E-mail address: j.e.romerogonzalez@qmul.ac.uk

Abstract

Bees efficiently learn asocial and social cues to optimise foraging from fluctuating

floral resources. However, it remains unclear how bees respond to divergent sources of

social information, and whether such social cues might modify bees' natural preferences

for non-social cues (e.g. flower colour), hence affecting foraging decisions. Here, we

investigated honeybees' (Apis mellifera) inspection and choices of unfamiliar flowers

based on both natural colour preferences and simultaneous foraging information from

conspecifics and heterospecifics. Individual honeybees' preferences for flowers were

recorded when the reward levels of a learned flower type had declined and novel-

coloured flowers were available where they would find either no social information or

one conspecific and one heterospecific (Bombus terrestris), each foraging from a

different coloured flower (magenta or yellow). Honeybees showed a natural preference

for magenta flowers. Honeybees modified their inspection of both types of flowers in

response to conspecific and heterospecific social information. The presence of either

demonstrator on the less-preferred yellow flower increased honeybees' inspection of

yellow flowers. Conspecific social information influenced observers' foraging choices

of yellow flowers, thus outweighing their original preference for magenta flowers. This

effect was not elicited by heterospecific social information. Our results indicate that

flower colour preferences of honeybees are rapidly adjusted in response to conspecific

social information, which in turn is preferred over heterospecific information, possibly

favouring the transmission of adaptive foraging information within species.

Keywords: behavioural flexibility, decision making, innate colour bias, insects,

interspecific, intraspecific, social information

Foraging decisions are central to an animal's survival and reproduction; deciding

where to forage in unpredictably changing environments is a major challenge that

animals constantly encounter (Chittka et al. 1999; Stephens et al. 2007). Foraging

decisions can be influenced in a context-dependant manner (Laland 2004; Kendal et al.

2009) by innate preferences (Lunau et al. 1996; Raine et al. 2006), previous individual

experience (Sclafani 1995), and the observation or interaction with another animals at a

foraging resource, i.e., social information (Heyes 1994; Leadbeater and Chittka, 2007b;

Hoppitt and Laland 2013). For example, naïve individuals tend to rely more on social

than individual information to gain familiarity about foraging sources (Galef and

Giraldeau 2001; Galef and Laland 2005). Conversely, experienced individuals that have

information on an advantageous foraging resource, will often ignore social cues. Thus,

social information is only drawn upon when individual information has become

outdated and acquiring up to date information may be costly (Laland 2004; Galef and

Laland 2005; Kendal et al. 2009). The influence of social information on foraging

decisions is taxonomically widespread (Galef and Giraldeau 2001; Valone and

Templeton 2002; Grüter and Leadbeater 2014). Social information allows animals to

find profitable foraging resources efficiently, instead of iteratively sampling the

environment through trial and error (Galef and Laland 2005).

In most animal communities, multiple species frequently share the same

foraging resources; thus members of the same (conspecific) and different

(heterospecific) species can potentially act as sources and users of social information

(Seppänen et al. 2007; Goodale et al. 2010; Avarguès-Weber et al. 2013; Parejo and

Avilés 2016; Loukola et al. 2020). Using social information indiscriminately is not

adaptive but animals should be selective when acquiring information from other

individuals (Laland 2004; Kendal et al. 2018). In a multi-species context, animals have

access to social information from different sources, which may give rise to a trade-off

between selecting sources with a close ecological similarity, implying high competition,

or sources whose ecological distance may involve less competition but a lower

informative value (Seppänen et al. 2007). Furthermore, in a foraging context, social

information has to be integrated with unlearned preferences (Laland and Plotkin 1993;

Leadbeater and Chittka 2007a; Jones et al. 2015) and existing individual information

(van Bergen et al. 2004; Jones et al. 2015) to determine foraging decisions in a context-

dependant manner (Laland 2004; Kendal et al. 2009).

Social bee workers forage for nectar and pollen from rapidly changing floral

resources (Heinrich 1979; Chittka et al. 1999) within multi-species communities (Fægri

and van der Pijl 1979; Kevan and Baker 1983) that might offer a wide spectrum of

social information. This makes bees a valuable model to explore how different sources

of social information can affect learned and unlearned preferences of individuals to

shape foraging decisions. Traditionally, the laboratory paradigms that investigate the

use of social information in bees tend to oversimplify the real field contexts where

animals naturally acquire information from others. For example, they test bees in

relatively unnatural settings where a single source of social information is presented,

e.g., a dead demonstrator pinned to simulated flowers (reviewed in Leadbeater and

Dawson 2017). Contrastingly, bees seeking nectar and pollen in the wild might

encounter far more complex circumstances where they have access to multiple sources

of social information (Fægri and van der Pijl 1979; Kevan and Baker 1983) that may

concur in time and space and diverge in their intrinsic relevance.

Honeybees and bumblebees of various species are in many locations sympatric

and typically forage upon similar flowers due to their generalist diet (Rogers et al. 2013;

Xie et al. 2016). Previous evidence indicates that bumblebees can acquire foraging

information from demonstrator honeybees – in this study, the "demonstrators" were

dead individuals placed on artificial flower types (Dawson and Chittka 2012). However,

in more realistic conditions, a bee forager whose known floral resources have decreased

in reward levels, will likely encounter other foragers, of their own and different species,

feeding from different types of flowers (Fægri and van der Pijl 1979; Kevan and Baker

1983). It remains unclear how bees respond to such divergent social information, and

whether in this context, social cues might modify bees' natural preferences for

particular colours in flowers (Chittka et al. 2004; Raine et al. 2006; Raine and Chittka

2007), hence influencing foraging decisions. We address these questions by testing

whether honeybees might adjust their colour preferences in response to simultaneous

sources of social information, i.e., a conspecific and heterospecific, each foraging from

either a preferred or non-preferred flower colour.

MATERIALS AND METHODS

(a) Set-up

We used free-flying honeybee foragers (*Apis mellifera*) from hives located in an urban area of London, UK. Over two consecutive summers (2017-2018), we trained foragers of one hive to collect 30% sucrose solution (w/w) from a gravity feeder (von Frisch, 1965, Figure 16, p. 18), placed 2 m away from one the hive. The feeder in turn attracted foragers from the other hives, which were then included in the experiments. The feeder was refilled every day at 0800 hours, yet honeybees were also free to forage on local floral resources.

In addition, we used bumblebee foragers from two colonies (Biobest, Belgium N.V.) for the experiments. Bumblebee nests were housed in bipartite wooden nest-boxes ($l = 29.5 \times w = 11.5 \times h = 9.5$ cm) connected to a wooden flight arena ($l = 77$ cm, $w = 52$ cm, $h = 30$ cm) by a Plexiglas tunnel ($l = 25$ cm, 3.5×3.5 cm). The floor of the flight arena was covered in white laminated paper. Three plastic sliding doors located along the corridor allowed controlled access to the arena. Before and after experiments, bumblebees could freely feed upon 30% (w/w) sucrose solution from a mass feeder in the middle of the arena. Bumblebee colonies were provided with 7 g of frozen pollen (Koppert B.V., The Netherlands) every two days.

(b) Training of bumblebees

We trained 30 bumblebee foragers (demonstrators) from two colonies (one colony per year) in a flight arena. In this group-training, foragers were allowed to enter the arena together at various times, and bumblebees learned to forage on an array of twelve plastic chips (2.4×2.4 cm, henceforth "flowers") placed on the top of transparent glass vials (h = 4 cm), positioned on the floor of the arena. The array was arranged in a rectangular grid formation (3×4) with a separation of 7.5 cm between the

edges of the flowers. We filled four yellow flowers with 10 μL of rewarding 50% (w/w)

sucrose solution, and four magenta and four transparent flowers with 10 μL of

unrewarding water (Figure 1A). Flowers were refilled after they were depleted by

foragers and the foragers left the flower. Half of the foragers experienced the opposite

colour-reinforcement contingency. Training took place for one day over 2 hours. We

carried out refreshment bouts (20 min) each day prior to testing. We tracked

demonstrator bees' identity with individual number tags (Opalithplättchen, Warnholz &

Bienenvoigt, Ellerau, Germany) glued to the top of the thorax by means of Loctite

Super Glue Gel (Loctite, Ohio, US). The floor of the arena and flowers were cleaned

with 70% ethanol after completing training.

(c) Training of honeybees

Once we completed bumblebees' training, we proceed to train honeybee

demonstrators and observers, which took place over 18 daily sessions. Each day, we

trained a set of five honeybees (demonstrators) foraging from the gravity feeder. In this

group-training, honeybees learned to enter the same flight arena (located outdoors)

where bumblebees were trained. We reversed the colour-reinforcement contingency so

that honeybees learned to forage on flowers of the opposite colour that bumblebees

were trained on. Half of the honeybees experienced four magenta flowers with 40 μL of

50% (w/w) sucrose solution, and four yellow and four transparent flowers with 40 μL of

water (Figure 1A). The other half of the honeybees experienced the reverse colour-

reinforcement contingency. Rewarding flowers were refilled after depletion by the

foragers and the foragers left the flower. Training lasted 1 hour, consisting of six bouts

(10 min). We identified trained individuals by marking their thorax with a white paint

mark (Posca Pen, Worcester, UK). We cleaned the floor of the arena and flowers with

70% ethanol after completing training.

Three hours after finalising the demonstrator's training, we selected a separate

batch of five foragers (observers) to train them on a different set-up. Every day,

observers were trained in group to enter the same flight arena and forage upon a

rectangular grid array of twelve transparent flowers (3 × 4) (Figure 1B). To encourage

foragers to sample multiple flowers during their foraging trips, only half of the flowers

contained 40 µL of 50% (w/w) sucrose solution, whereas the other half contained 40 µL

of water. Rewarding flowers were replenished after emptied and the forager left the

flower. The positions of all the flowers were changed every bout (10 min) over 1 hour

of training. We marked trained observers with a green dot to distinguish them from the

demonstrators. Original honeybees trained (demonstrators) were allowed to visit the

setup during this training, which did not interfere with the previous colour training, as

shown in the results section. After training observers, we cleaned the flowers and the

arena floor with 70% ethanol.

(d) Testing the effect of colour preference on honeybees' foraging decisions

We carried out this control test every day after completing training. Only two to

three individuals from the batch of five observer honeybees (trained on transparent

flowers) regularly showed up at the setup at the time of testing. Thus, we only tested

two daily individuals. Overall, we tested 20 honeybees individually (control group) in a

context where transparent flowers were unrewarding. We assessed whether they might

show a natural preference to inspect and forage from one flower type between two

unfamiliar alternatives, i.e., yellow and magenta. One honeybee was let in the arena to

explore a rectangular grid array of twelve flowers (section b, above), consisting of four

familiar transparent flowers containing 20 µL of water and eight unfamiliar flowers,

four yellow and four magenta, all filled with a scentless reward of 20 µL of 50% (w/w)

sucrose solution, (Figure 1D). The test began once the individual inspected any flower.

That is, the honeybee approached the flower by displaying a slow side-to-side hover,

with its head oriented towards it and within at least one body length (Ings et al. 2009).

The test concluded once the honeybee landed and foraged upon any unfamiliar flower,

or 3 minutes after the test started. As this test was designed to evaluate the influence of

honeybees' colour preferences on an actual foraging decision, rather than measuring

their innate colour preferences, we regarded the flower type where the individual landed

and foraged as preferred over the alternative type. To prevent re-testing the same

individuals, we captured honeybees, after concluding the test, to give them a distinctive

red paint mark. The flowers and arena floor were cleaned with 70% ethanol between

tests. To evaluate honeybees' inspection of flowers before they chose a flower to forage

(foraging decision), we recorded the test with a sport camera (Yi, Xiaomi Inc. China)

featuring a recording frame rate of 30 fps and a resolution of 720 p (1,280 × 720 pixels).

The camera was positioned 20 cm above the entrance of the bumblebee nest (Figure 1).

Its field of view was adjusted such that it looked down into the arena at ~50° from a

horizontal angle.

(e) Testing the influence of social information on foraging decisions

194   To evaluate the influence of simultaneous sources (conspecific and

heterospecific) of social information on honeybee's inspection of either yellow or

magenta flowers and their subsequent foraging decisions, we tested 45 honeybees in the

same context as individuals in the control group (section d) but in the presence of one

bumblebee and one honeybee demonstrator, each foraging upon either a yellow or

magenta flower. We introduced the demonstrators and observer in the arena in the

following order: 1) a honeybee demonstrator was let in through a sliding door on the

back wall of the arena (Figure 1D). Once she began to feed from a flower of her trained

colour, 2) a bumblebee demonstrator was released from the Plexiglas tunnel that

connected the arena to the nest-box (Figure 1D). When she started to forage from a

flower of her trained colour (opposite colour as the honeybee demonstrator), 3) a

honeybee observer was let into the arena through the sliding door on the back wall. Due

to training, both demonstrators swiftly landed and foraged exclusively on flowers of

their trained colour. We tested 19 observer honeybees with a conspecific demonstrator

foraging on magenta flowers and a heterospectific foraging on yellow flowers and 26

observers with the reversed colour-reinforcement contingency.

The test began once the observer honeybee inspected (see section d) any

occupied or unoccupied flower. The test concluded once the observer landed and

foraged upon any unfamiliar flower, or 3 minutes after the test started. In the test,

demonstrators moved freely between flowers after depleting them, as they naturally do

within inflorescences. Thus, we only considered that observers foraged on an unfamiliar

flower when they actually fed upon the sucrose solution reward from unemptied

flowers. To prevent re-testing the same individuals, we caught tested honeybees and

marked them with a distinctive red paint mark. The flowers and arena floor were

cleaned with 70% ethanol between tests. To evaluate honeybees' inspection of flowers

and their interactions with the demonstrators before they made a foraging decision, the

test was recorded as described in section (d) above.

Figure 1. Training and testing protocols in the flight arena. (**A**) Honeybee foragers (demonstrators) were trained to find 50% sucrose solution on either four yellow or magenta flowers and water on four flowers of the alternative colour and four transparent flowers. Bumblebee foragers (demonstrators) were group trained using the same protocol (**B**) A different batch of honeybee foragers (observers) was group trained to find 50% sucrose solution on six of twelve transparent flowers and unrewarding water on the other six transparent flowers. (**C**) Honeybee observers were tested on their preparedness to forage on an unfamiliar coloured flower type, depending on its colour (yellow or magenta), once their learned flower type (transparent) yielded no reward. (**D**) Two demonstrators (one honeybee and one bumblebee) and one observer honeybee were sequentially introduced in the flight arena: (1) A honeybee demonstrator was let in the flight arena, once she foraged from a flower of her trained colour, (2) a bumblebee demonstrator was released into the arena and let to forage from a flower of her trained colour (different colour from the honeybee demonstrator), (3) an observer honeybee was then introduced to test her in a context where the flowers she previously associated with reward (transparent) were unrewarding and unfamiliar coloured

238 flowers were demonstrated by a conspecific and a heterospecific (bumblebee) demonstrator, each

239 foraging from non-preferred (yellow) and preferred (magenta) flowers, this design was counterbalanced.

241 *(d) Analyses*

242   We analysed the behaviour of tested honeybees from video recordings, using the

BORIS behavioural observation software (Friard and Gamba, 2016). To assess

individuals' inspection of either yellow or magenta flowers and their preference to

forage upon one type of flower over the other, we analysed two main behavioural

categories. That is, the frequency of inspecting transparent, yellow and magenta flowers

(Ings et al. 2009; Balamurali et al. 2018) and individual's foraging choice, i.e., the

yellow or magenta flower where individuals landed and foraged. We used logistic

analysis to explore the influence of flower colour on the likelihood of honeybees

inspecting a flower for the first time, the likelihood of foraging on a flower type, as well

as the proportion of transparent, yellow and magenta flowers they inspected. For the

latter two variables, we took underdispersion into account via a quasinomial model.

253   For honeybees exposed to social information, we considered inspection of

occupied flowers as an indicator that observers detected the demonstrator's presence.

Thus, four different scenarios were possible before honeybees made a foraging

decision: they could have detected both demonstrators, either the honeybee or

bumblebee demonstrator, or none of them. We compared the likelihood of each

situation against the expected probability with a Chi-Square Goodness of Fit Test. Nine

individuals that did not detect the presence of the demonstrators were not considered for

all analyses.

261   We used logistic analysis to determine whether colour preferences and

concurrent conspecific and heterospecific social information had a similar influence on

foraging decisions of honeybees. Social information was included as a predictor

variable (binary: 1= present, 0= absent) for the likelihood of honeybees landing on

either an unfamiliar yellow or magenta flower.

Further, to evaluate the influence of social information on honeybees' readiness

to forage on an unfamiliar flower, we compared two measurements between the control

group and the group exposed to social information. That is, the total number of flowers

that individuals inspected before they chose a flower to forage, and the time it took

them to make such decision (latency to forage). We analysed both measurements to

explore whether observers' readiness to forage was influenced by the first foraging

demonstrator they detected in the test (i.e., honeybee or bumblebee). This was

irrespective of whether they detected only one or both demonstrators. These analyses

were conducted with a Wilcoxon rank sum test.

To determine whether observers that detected both demonstrators inspected the

flowers occupied by a honeybee and bumblebee at a similar frequency, we compared,

with a Wilcoxon signed rank test, the proportion of flowers occupied by each

demonstrator that observers inspected. We also analysed, with a Wilcoxon rank sum

test, whether honeybees that only detected either a conspecific or heterospecific

demonstrator, differed in the proportion of occupied flowers that they inspected, relative

to all the flowers they inspected during the test. We used logistic analysis to explore the

influence of flower colour and demonstrator's species on the likelihood of honeybee

observers selecting the demonstrated type of flower and the likelihood that observers

would forage on a flower occupied by either a honeybee or bumblebee, after

approaching it for the first time. For the latter variable, we took underdispersion into

account via a quasinomial model.

The instances in which observers inspected an occupied flower that did not

progress into landing and foraging were recorded as rejections. We evaluated whether

observers rejected the flowers occupied by a honeybee or bumblebee demonstrator at a

similar frequency. For observers that detected both demonstrators, we compared the

proportion of times that they rejected the flowers occupied by each demonstrator, using

a Wilcoxon signed rank test. We also compared, with a Wilcoxon rank sum test, the

total number of occupied flowers that observers rejected when they only detected either

the honeybee or bumblebee demonstrator.

To assess whether observers altered their inspection of flowers after they

initially detected the presence of either a foraging honeybee or bumblebee, we adjusted

the frequency of observers' inspection of flowers by normalising the data through an

index calculation, with the following equation:

299
$$\text{Inspecting Index (II)} = \frac{i_m - i_y}{i_m + i_y}$$

, where, i_m and i_y are the frequency that the observer inspected magenta and yellow

flowers, respectively.

In the indices, a negative value (minimum -1) equates to a preference to inspect

yellow flowers, whereas a positive value (maximum +1) equates to a preference to

inspect magenta flowers, an index near equal to zero implies that the observers either

inspected both types of flowers equally, or they did not inspect the flowers at all. We

calculated inspecting indices (II) based on the sequence of events that preceded the

honeybees' foraging decisions. That is, each inspecting index (II) represented

observers' inspection of flowers, before and after they detected either demonstrator

(honeybee or bumblebee) foraging on a flower. We compared indices against chance

expectation (Index = 0) with a Wilcoxon signed-rank test. A significant switch from a

negative or positive value, before the observer detected a demonstrator, to a positive or

negative value, after this occurred, indicates that observers modified their inspection of

yellow or magenta flowers in response to conspecific or heterospecific social

information. All analyses were conducted using R statistical software (R Core Team,

2019).

Ethical Note

Bees were kept in their natural colony environment in unaltered dark conditions. The

gravity feeder where honeybees foraged from, was only filled when no individuals were

present to avoid disturbance. Bumblebees were fed with a minimum disturbance under

red light, which is poorly visible to bees. Colonies were not food deprived during

experiments. Only bee foragers that freely engaged in foraging behaviour were trained

and tested. Honeybees were tagged with a dot of paint on the thorax while feeding and

bumblebees were carefully handled with dissection forceps to glue number tags on their

thoraxes.

RESULTS

Effect of colour preference on honeybees' foraging decisions

In a context where a familiar flower type ceases to yield reward, how does

colour preference influence honeybees' exploration of contiguous, novel flowers and

ultimate move to foraging upon a new flower type? We carried out a control test to

assess whether honeybees preferentially inspect one unfamiliar flower type over

another, and whether this might affect their ultimate foraging decision (move to

foraging upon a new flower type). We trained honeybees (observers) to find either a

food reward (50% sucrose solution) or unrewarding water on transparent flowers

(Materials and Methods). We then we tested bees in a context where transparent flowers

were unrewarding and two unfamiliar yellow and magenta flower types delivered a food

reward (scentless 50% sucrose solution). The majority of individuals (77%) did not

inspect the familiar transparent flowers at all; rather honeybees preferred to inspect and

forage on magenta flowers. The proportion of flowers inspected by honeybees was

influenced by flower colour (Logistic regression: $F = 35.86$, $N = 20$, P < 0.001; Fig.

3.2A). Flower colour also influenced the likelihood of honeybees first exploring a

flower type (Logistic regression: $\chi^2_1 = 15.44$, $N = 20$, P < 0.001; Fig.3.2B) and the

likelihood of selecting a flower to forage (Logistic regression: $F = 30.04$, $N = 20$, P <

0.001; Fig. 3.2C). This shows that honeybees had a colour preference for magenta over

yellow flowers.

Figure 3.2. Honeybees prefer magenta over yellow flowers **(A)** Honeybees inspected magenta flowers at

a higher proportion than yellow and transparent flowers. **(B)** The likelihood of honeybees inspecting an

unfamiliar flower for the first time was higher for magenta compared to yellow flowers. **(C)** A higher

proportion of honeybees preferred to forage on magenta than yellow flowers. Bars = mean. Horizontal

lines = standard error. Asterisks = p < 0.05.

Influence of social information on honeybees' foraging decisions

In a field-like context, including simultaneous and divergent conspecific and

heterospecific social information, we investigated how this social information might

affect honeybees' inspection of flower types and ultimate foraging decisions. We

considered that observers detected the demonstrators' presence once they inspected an

occupied flower for the first time. We found no difference in the proportion of observers

that detected both demonstrators, only the honeybee or bumblebee demonstrator, or

none of them (Chi-Square Goodness of Fit Test: $\chi^2_3 = 2.2$, $N = 45$, $P = 0.31$; Fig. 3A).

The likelihood of foraging on an unfamiliar flower type was not influenced by the

presence of social information (Logistic regression: $\chi^2_1 = 0.9$, $N = 56$, $P = 0.34$; Fig.

3.2B). Similar to the control group, the vast majority of observers (90%) did not inspect

the transparent flowers at all. These results suggest that both individual colour

preference and social information similarly influenced honeybees when locating a new

foraging resource. Compared to observer honeybees that first detected a foraging

bumblebee in the test, individuals in the control group, whose foraging decisions were

solely influenced by colour preference, made faster choices (Wilcoxon rank-sum: $W =$

71, $N = 35$, $P = 0.007$; Fig. 3.3C) and inspected fewer flowers (Wilcoxon rank-sum: W

$= 77$, $N = 35$, $P = 0.011$; Fig. 3.3D). However there was no difference between the

control group and those observers that first detected the presence of a foraging

conspecific (Wilcoxon rank-sum: latency: $W = 133$, $N = 30$, $P = 0.29$; Fig. 3.3C;

number of flowers: $W = 125$, $N = 30$, $P = 0.45$; Fig. 3.3D). Further, when observers

first detected the conspecific demonstrator, they made faster foraging decisions

(Wilcoxon rank-sum: $W = 184$, $N = 29$, $P < 0.001$; Fig. 3.3C), preceded by less

inspection of flowers (Wilcoxon rank-sum: $W = 176.5$, $N = 29$, $P < 0.001$; Fig. 3.3D),

than when they first detected the heterospecific bumblebee. These results suggest that

both individual preference for magenta flowers and conspecific social information

similarly affected honeybees' foraging choices, enabling swift decisions. Conversely,

observers that initially detected a foraging heterospecific, explored the flowers

extensively before making a foraging decision.

Honeybee observers that detected either one or both demonstrators, similarly

inspected flowers occupied by either the honeybee or bumblebee demonstrator

(Wilcoxon rank-sum; one demonstrator: $W = 34.5$, $N = 16$; $P = 0.45$; both

demonstrators: Wilcoxon signed-rank: $V = 22.5$, $N = 13$, $P = 1$; Fig. 3.3E). However, the

likelihood that an observer would forage on an unfamiliar flower type was influenced

by the demonstrators' species (Logistic regression: $\chi^2_1 = 11.78$, $N = 45$, $P < 0.001$; Fig.

3.3F) and flower colour (Logistic regression: $\chi^2_1 = 5.08$, $N = 45$, $P = 0.024$; Fig. 3.3F)

but was only marginally influenced by the statistical interaction between these main

effects (Logistic regression: $\chi^2_1 = 3.17$, $N = 45$, $P = 0.07$; Fig. 3.3F). Further,

demonstrators' species influenced the likelihood of observers foraging on an occupied

flower after approaching it for the first time (Logistic regression: $\chi^2_1 = 20.51$, $N = 51$, P

< 0.001; Fig. 3.3G). That is, observers readily responded to the presence of a foraging

conspecific by joining such demonstrator on the unfamiliar flower type. This response

was not affected by flower colour (Logistic regression: $\chi^2_1 = 0.03$, $N = 51$, $P = 0.82$;

Fig. 3.3G) nor by any statistical interaction between main effects (Logistic regression:

$\chi^2_1 = 0$, $N = 51$, $P = 1$; Fig. 3.3G). In contrast, observer honeybees rejected the flowers

occupied by a bumblebee demonstrator at a higher proportion (Wilcoxon signed-rank: V

$= 3$, $N = 13$, P $= 0.008$; Fig. 3.3H) than the flowers occupied by a conspecific.

Figure 3. Concurrent conspecific and heterospecific social information influences foraging behaviour of

honeybees **(A)** Observer honeybees were equally likely to detect both demonstrators, only the honeybee

or bumblebee demonstrator, and none of them. **(B)** The proportion of honeybees that foraged on

unfamiliar flowers was not different between observers that detected the presence of demonstrators and

individuals in the control group (no social information). **(C)** Honeybees in the control group and those

that first detected a foraging conspecific made faster foraging decisions than observers that first detected

a foraging heterospecific. **(D)** Honeybees in the control group and those that first detected a foraging

conspecific inspected fewer flowers before making a foraging decision than observers that first detected

the presence of a foraging heterospecific. **(E)** Honeybee observers that detected only one or both

demonstrators during the test; similarly inspected flowers occupied by the honeybee and bumblebee

demonstrator. **(F)** Demonstrators' species and flower colour had an effect on the proportion of honeybees

that foraged on unfamiliar flowers, yet no interaction of main effects was found. **(G)** Once observers

detected a conspecific demonstrator foraging on an unfamiliar flower type, they were more likely to land

and forage on such a flower, but they never foraged on a flower occupied by a heterospecific

demonstrator. **(H)** Observers rejected flowers occupied by a bumblebee at a higher proportion than

flowers occupied by a conspecific. HB = honeybee, BB = bumblebee. Bars = mean. Horizontal lines =

standard error. Asterisks = $p < 0.05$. n.s. = not significantly different.

Adjustment of honeybees' colour preference in response to social information

The inspecting indexes (II), described in (d), allowed us to analyse honeybees'

inspection of flowers as a flexible process that culminated in a foraging decision, and

whose variation in response to social information was measurable in terms of an index

value. Observer honeybees showed no preference to inspect magenta or yellow flowers

before they detected the bumblebee demonstrator foraging on either a magenta

(Wilcoxon signed-rank test: $V = 20$, $N = 9$, $P = 0.15$; Fig.4A) or yellow flower

(Wilcoxon signed-rank test: $V = 12$, $N = 9$, $P = 0.11$; Fig. 4A). After honeybees detected

a bumblebee foraging from a magenta flower, they preferentially inspected this type of

flower (Wilcoxon signed-rank: $V = 26$, $N = 9$, $P = 0.026$; Fig. 3.4A). This preference did

not occur after they observed a bumblebee demonstrator foraging on a yellow flower,

and instead they showed no preference to inspect either type of flower (Wilcoxon

signed-rank: $V = 18.5$, $N = 9$, $P = 0.5$; Fig. 3.4A). Honeybees showed no preference to

inspect either type of flower before they detected a conspecific foraging upon either a

magenta (Wilcoxon signed-rank: $V = 16.5$, $N = 11$, $P = 0.61$; Fig. 3.4B) or yellow

flower (Wilcoxon signed-rank: $V = 23$, $N = 11$, $P = 0.26$; Fig. 3.4B). However, observer

honeybees did respond to the presence of a conspecific foraging on either flower type

by increasing their inspection of demonstrated flowers (Wilcoxon signed-rank:

magenta: $V = 52$, $N = 11$, $P = 0.006$; yellow: $V = 1$, $N = 13$, $P = 0.002$; Fig. 4B). These

results show that honeybees' inspection of unfamiliar flowers underwent a sequential

adjustment in response to conspecific and heterospecific social information, which

ultimately affected their foraging decisions.

Figure 4. Honeybees adjust their exploration of flowers and foraging preferences in response to

concurrent conspecific and heterospecific social information. Inspecting indices (II) compare to chance

level (Index = 0) the inspection of magenta (dark grey bars) and yellow (light grey bars) flowers that

honeybee observers had before and after they detected either a conspecific or heterospecific demonstrator

foraging on either type of flower. **(A)** Before observers detected a bumblebee demonstrator foraging on

either a magenta or yellow flower, they showed no preference to inspect neither of these flower types.

After observers detected a bumblebee demonstrator foraging on a magenta flower, they inspected this

type of flowers more frequently. However, when observers detected the bumblebee demonstrator foraging

on a yellow flower, this did not affect their inspection of yellow or magenta flowers. **(B)** Before observers

detected a conspecific foraging upon either a yellow or magenta flower, they showed no preference to

inspect magenta or yellow flowers. After observers detected a conspecific demonstrator foraging on either

a yellow or magenta flower, they increased their inspection of the demonstrated type of flower. HB =

honeybee, BB = bumblebee. Index = 0. Bars = mean. Horizontal lines = standard error. Asterisks = $p <$

0.05. n.s. = not significantly different between groups or from chance level.

DISCUSSION

The results presented here provide evidence that honeybees' colour preferences

can be adjusted in response to simultaneous social information from conspecifics and

heterospecifics. Thus, specific social information differently influences honeybees'

exploration of flowers and consequent foraging decisions. Unexpectedly, honeybees

payed little attention to the transparent flowers on which they were trained. Instead,

prior to making a foraging decision, they inspected more frequently the magenta flowers

(control group) or the type of flower demonstrated by a conspecific.

Even though inspecting flowers from a close distance does not provide foragers

with information on the reward status of a flower, it is important in the process of

choosing a flower (Lunau et al. 1996). In the absence of social information, honeybees'

natural preference for magenta flowers influenced inspection and choices of flowers.

Magenta is actually a mixture of blue and red, since the red component is not fully

perceived by bees (Menzel and Shmida 1993), they perceive magenta flowers as blue

(Chittka and Waser, 1997; Waser and Chittka 1998). Honeybees and bumblebees

usually have innate biases for colours in the violet to blue range of the spectrum (Giurfa

et al. 1995; Chittka et al. 2004), which adaptively correlates with the nectar production

of local flowers (Giurfa et al. 1995; Chittka et al. 2004). Although colour biases may be

overridden by individual learning (Raine et al., 2006), they potentially govern foraging

decisions in bees when selecting among novel flower types (Gumbert 2000). In the

absence of social information, flower choices of tested honeybees conceivably resulted

from either their innate colour preferences or previous experience in the field with local

flower species as both factors can be linked with the likelihood of finding a profitable

food reward (Giurfa et al. 1995; Chittka et al. 2004).

In the wild, honeybee foragers are likely to encounter members of the same and

different species foraging concurrently in a flower patch including distinct flower types

(Fægri and van der Pijl 1979; Kevan and Baker 1983). Our results demonstrate that

such social information is integrated with honeybees' colour preferences during the

process of making a foraging decision. In this realistic scenario, with two simultaneous

sources of conspecific and heterospecific social information, when the first foraging

demonstrator that honeybees detected was a member of the same species, they promptly

responded to social information by joining the demonstrator on the unfamiliar flower

type. This evidence is consistent with previous findings in bumblebees (reviewed in

Leadbeater and Dawson 2017) and supports the notion that the use of conspecific social

information in bees may be mediated by local enhancement, with foragers being

attracted to flowers occupied by members of the same species (Leadbeater and Chittka

2007a). Such a response potentially optimises the foraging efficiency of bees by

facilitating their movements between different flower types (Leadbeater and Chittka

2005; Kawaguchi et al. 2007).

Honeybees and bumblebees often forage upon similar floral resources; it is then

conceivable that they may concur in the same flower patches (Rogers et al. 2013; Xie et

al. 2016). In our experiments, we used freely moving bumblebee demonstrators to

elucidate how social information might flow between these species. Our results indicate

that in the likely scenario of a honeybee exploring a flower patch and encountering both

a conspecific and heterospecific (bumblebee) individual foraging from different flower

types, its readiness to use social information and make a foraging decision might

depend on the first observed species providing social information. That is, observer

honeybees that detected a conspecific demonstrator before a heterospecific bumblebee

made swifter foraging decisions than those observers that first detected the bumblebee

demonstrator, which spent more time exploring the flowers before making a foraging

decision. This evidence suggests that in a foraging context, honeybees' discrimination

of conspecifics and heterospecifics might determine the selective use of social

information. Theory predicts that using social information from heterospecifics should

be favourable when there is a large niche overlap (Seppänen et al. 2007) yet, on

adaptive grounds, when animals can select between social information from

conspecifics and heterospecifics, they should typically favour the former as this

naturally reflects its ecological needs (Jaakkonen et al. 2015). In line with this,

honeybees consistently selected the type of flowers demonstrated by conspecifics, even

though they attended to the presence of both demonstrators whilst exploring the set-up.

Prioritising conspecific over heterospecific choices can be explained due to its high

fitness value (Seppänen et al. 2007; Jaakkonen et al. 2015) but it can also serve to the

transmission of novel behavioural traits or preferences that may be adaptively valuable

for a particular species (Laland and Plotkin 1993; Jaakkonen et al. 2015; Alem et al.

2016; Danchin et al. 2018).

Floral reward levels differ strongly among plant species and constantly change

over time in an unpredictable manner (Heinrich 1979). To achieve efficient foraging,

bees can rapidly learn to associate floral traits such as colour, shape and scent with

reward quality in flowers (Chittka et al. 1999). Bees can be initially attracted to forage

from an unfamiliar flower species via either innate and learned colour preferences

(Giurfa et al. 1995; Gumbert 2000; Chittka et al. 2004; Raine and Chittka 2007) or

using social information (Leadbeater and Chittka 2007b; Grüter and Leadbeater 2014;

Leadbeater and Dawson 2017). In the wild, bees seeking floral resources are unlikely to

experience asocial and social cues separately, rather they may frequently be exposed to

a complex combination of cues potentially affecting their flower choices. Our findings

demonstrate that simultaneous conspecific and heterospecific social information

influences honeybees' colour preferences, which may in turn shape the process of

acquiring new information underlying foraging decisions.

Compared to honeybees in the control group (i.e., no social information), the

exploration behaviour of honeybees that observed a conspecific or heterospecific

demonstrator foraging upon the less preferred yellow flowers, reflected a more evenly

distributed inspection of magenta and yellow flowers. That is, the presence of either

demonstrator on a yellow flower increased the 'attractiveness' of such flower type for

honeybees, possibly via stimulus enhancement, an effect widely described in social

learning literature (Heyes 2012). Remarkably, honeybees' inspection of flowers,

naturally biased towards magenta flowers, underwent a sequential and flexible

adjustment prior to make a foraging decision. Such adjustment was modulated by visual

foraging information from members of the same and different species.

It has been demonstrated that bees are attracted to the presence of foraging

conspecifics when presented with unfamiliar flowers which may lead them to identify

new rewarding flower species (Leadbeater and Chittka 2007a; Jones et al. 2015). Our

results indicate that the presence of a foraging conspecific not only influenced honeybee

observers to select flowers that matched their colour preference (magenta) but such

conspecific social information also outweighed observers' colour preference so that

they selected the normally non-preferred yellow flowers. Whereas intraspecific social

transmission of foraging information may be more stable when it reinforces a prior

preference (Laland and Plotkin 1990), such preferences may be adjusted and potentially

overridden in response to conspecific social information about a novel foraging resource

(Jones et al. 2015). This may enable bees to reasonably adapt to wildly varying floral

reward levels. Thus, if the presence of a foraging conspecific can reliably be associated

with a rewarding outcome, the use of such social cue should be reinforced to

consistently influence bees' flowers choices across foraging contexts (Leadbeater and

Chittka 2009). Interestingly, in our experiments the presence of a foraging

heterospecific also increased honeybees' attraction to the demonstrated flower types, yet

heterospecific social information did not influence honeybees' foraging choices. In fact,

honeybee observers never landed on the flowers occupied by the bumblebee

demonstrator, in contrast to the flowers occupied by the conspecific demonstrator. Even

though, heterospecific social information is predictably valuable when there is a large

niche overlap (Seppänen et al. 2007), conspecifics choices might offer a more

predictable social cue, decreasing the risk of acquiring maladaptive information

(Giraldeau et al. 2002) as conspecifics share the same ecological needs (Seppänen et al.

2007; Goodale et al. 2010).

Multiple social and asocial cues can potentially affect animals' foraging

decisions in different circumstances (Sclafani 1995; Galef and Giraldeau 2001; Grüter

and Leadbeater 2014). The effect of social information on previous, innate or learned

preferences influences the foraging decision of a particular individual and may also

promote the transmission of adaptive information about food sources (Laland and

Plotkin 1990; Galef and Giraldeau 2001). Despite the fact that social information offers

a clear advantage to individuals exploring novel foraging resources (Galef and

Giraldeau 2001), its use should not be indiscriminate but respond to particular

circumstances in order to lead to adaptive choices (Laland 2004; Kendal et al. 2018).

Our results extend previous evidence showing that conspecific social information is

commonly used in situations of uncertainty (Kendal et al. 2004; van Bergen Yfke et al.

2004; Galef et al. 2008; Smolla et al. 2016), such that it can outweigh both

predetermined individual preferences (Dugatkin 1996; Jones et al. 2015) and

heterospecific social information (Jaakkonen et al. 2015) to influence ecologically

relevant decisions in animals.

It is conceivable that natural selection should favour the salience of social

stimuli with high ecological relevance (Seppänen et al. 2007; Leadbeater and Dawson

2017), conspecifics in our experiments. Whether such salience can similarly operate to

select information from heterospecific sources, based on their relative informative

value, deserves further consideration. We provide an ecologically relevant picture of the

process thereby multiple non-social and social cues may shape foraging decisions of

bees. Thus, findings presented here contribute to our understanding of the flexibility of

individual preferences and their adjustment in response to different sources of social

information. Our results in turn shed light on the selectivity of animals for conspecific

over heterospecific information as a possible mechanism to facilitate the social

transmission of foraging information within species.

611   **References**

612   Alem S, Perry CJ, Zhu X, Loukola OJ, Ingraham T, Søvik E, Chittka L. 2016.
613   Associative Mechanisms Allow for Social Learning and Cultural Transmission of String
614   Pulling in an Insect. PLOS Biol. 14(10):e1002564. doi:10.1371/journal.pbio.1002564.
615
616   Avarguès-Weber A, Dawson EH, Chittka L. 2013. Mechanisms of social learning
617   across species boundaries. J Zool. 290(1):1–11. doi:10.1111/jzo.12015.
618
619   Balamurali GS, Nicholls E, Somanathan H, Hempel de Ibarra N. 2018. A comparative
620   analysis of colour preferences in temperate and tropical social bees. Sci Nat. 105(1):8.
621   doi:10.1007/s00114-017-1531-z.
622
623   Chittka L, Waser NM. 1997. Why red flowers are not invisible to bees. Israel Journal of
624   Plant Sciences. 45(2–3):169–183. doi:10.1080/07929978.1997.10676682.
625
626   Chittka L, Thomson JD, Waser NM. 1999. Flower Constancy, Insect Psychology, and
627   Plant Evolution. Naturwissenschaften. 86(8):361–377. doi:10.1007/s001140050636.
628
629   Chittka L, Ings TC, Raine NE. 2004. Chance and adaptation in the evolution of island
630   bumblebee behaviour. Popul Ecol. 46(3):243–251. doi:10.1007/s10144-004-0180-1.
631
632   Chittka L, Thomson JD, Waser NM. 1999. Flower Constancy, Insect Psychology, and
633   Plant Evolution. Naturwissenschaften. 86(8):361–377. doi:10.1007/s001140050636.
634
635   Dugatkin LA. 1996. Interface between culturally based preferences and genetic
636   preferences: female mate choice in Poecilia reticulata. Proc Natl Acad Sci. 93(7):2770–
637   2773. doi:10.1073/pnas.93.7.2770.
638
639   Danchin E, Nöbel S, Pocheville A, Dagaeff A-C, Demay L, Alphand M, Ranty-Roby S,
640   Renssen L van, Monier M, Gazagne E, et al. 2018. Cultural flies: Conformist social
641   learning in fruitflies predicts long-lasting mate-choice traditions. Science.
642   362(6418):1025–1030. doi:10.1126/science.aat1590.
643
644   Dawson EH, Chittka L. 2012. Conspecific and Heterospecific Information Use in
645   Bumblebees. PLOS ONE. 7(2):e31444. doi:10.1371/journal.pone.0031444.
646
647   Fægri K, van der Pijl L. 1979. Chapter 11 - Animals as pollinators. In: Fægri K, van der
648   Pijl L, editors. Principles of Pollination Ecology (Third Edition). Amsterdam:
649   Pergamon. p. 96–133.
650
651   Galef BG, Dudley KE, Whiskin EE. 2008. Social learning of food preferences in
652   'dissatisfied' and 'uncertain' Norway rats. Anim Behav. 75(2):631–637.
653   doi:10.1016/j.anbehav.2007.06.024.
654
655   Galef BG, Giraldeau L-A. 2001. Social influences on foraging in vertebrates: causal
656   mechanisms and adaptive functions. Anim Behav. 61(1):3–15.
657   doi:10.1006/anbe.2000.1557.
658
Galef BG, Laland KN. 2005. Social Learning in Animals: Empirical Studies and

Theoretical Models. BioScience. 55(6):489–499. doi:10.1641/0006-
3568(2005)055[0489:SLIAES]2.0.CO;2.

Giraldeau L, Valone TJ, Templeton JJ. 2002. Potential disadvantages of using socially
acquired information. Johnstone RA, Dall SRX, editors. Philos Trans R Soc Lond B
Biol Sci. 357(1427):1559–1566. doi:10.1098/rstb.2002.1065.

Giurfa M, Núñez J, Chittka L, Menzel R. 1995. Colour preferences of flower-naive
honeybees. J Comp Physiol A. 177(3):247–259. doi:10.1007/BF00192415.

Goodale E, Beauchamp G, Magrath RD, Nieh JC, Ruxton GD. 2010. Interspecific
information transfer influences animal community structure. Trends Ecol Evol.
25(6):354–361. doi:10.1016/j.tree.2010.01.002.

Grüter C, Leadbeater E. 2014. Insights from insects about adaptive social information
use. Trends Ecol Evol. 29(3):177–184. doi:10.1016/j.tree.2014.01.004.

Gumbert A. 2000. Color choices by bumble bees (Bombus terrestris): innate preferences
and generalization after learning. Behav Ecol Sociobiol. 48(1):36–43.
doi:10.1007/s002650000213.

Heinrich B. 1979. Resource heterogeneity and patterns of movement in foraging
bumblebees. Oecologia. 40(3):235–245. doi:10.1007/BF00345321.

Heyes C. 2012. What's social about social learning? J Comp Psychol. 126(2):193–202.
doi:10.1037/a0025180.

Heyes CM. 1994 Social-learning in animals–categories and mechanisms. Biol. Rev.
Camb. Phil. Soc. 69, 207–231. doi:10.1111/j.1469- 185X.1994.tb01506.x

Hoppitt W, Laland KN. 2013. Social Learning: An Introduction to Mechanisms,
Methods, and Models. Princeton University Press.

Ings TC, Raine NE, Chittka L. 2009. A population comparison of the strength and
persistence of innate colour preference and learning speed in the bumblebee Bombus
terrestris. Behav Ecol Sociobiol. 63(8):1207–1218. doi:10.1007/s00265-009-0731-8.

Jaakkonen T, Kivelä SM, Meier CM, Forsman JT. 2015. The use and relative
importance of intraspecific and interspecific social information in a bird community.
Behav Ecol. 26(1):55–64. doi:10.1093/beheco/aru144.

Jones PL, Ryan MJ, Chittka L. 2015. The influence of past experience with flower
reward quality on social learning in bumblebees. Anim Behav. 101:11–18.
doi:10.1016/j.anbehav.2014.12.016.

Kawaguchi LG, Ohashi K, Toquenaga Y. 2007. Contrasting responses of bumble bees
to feeding conspecifics on their familiar and unfamiliar flowers. Proc R Soc B Biol Sci.
274(1626):2661–2667. doi:10.1098/rspb.2007.0860.

Kendal RL, Boogert NJ, Rendell L, Laland KN, Webster M, Jones PL. 2018. Social

Learning Strategies: Bridge-Building between Fields. Trends Cogn Sci. 22(7):651–665. doi:10.1016/j.tics.2018.04.003.

Kendal RL, Coolen I, Laland KN. 2004. The role of conformity in foraging when personal and social information conflict. Behav Ecol. 15(2):269–277. doi:10.1093/beheco/arh008.

Kendal, RL, Coolen, I, Laland, KN. Adaptive trade-offs in the use of social and personal information. In: Dukas R, Ratcliffe JM, editors. Cognitive ecology II. Chicago: The University of Chicago Press. p. 249-271.

Kevan PG, Baker HG. 1983. Insects as Flower Visitors and Pollinators. Annu Rev Entomol. 28(1):407–453. doi:10.1146/annurev.en.28.010183.002203.

Laland KN. 2004. Social learning strategies. Anim Learn Behav. 32(1):4–14. doi:10.3758/BF03196002.

Laland KN, Plotkin HC. 1990. Social learning and social transmission of foraging information in Norway rats (Rattus norvegicus). Anim Learn Behav. 18(3):246–251. doi:10.3758/BF03205282.

Laland KN, Plotkin HC. 1993. Social transmission of food preferences among norway rats by marking of food sites and by gustatory contact. Anim Learn Behav. 21(1):35–41. doi:10.3758/BF03197974.

Leadbeater E, Chittka L. 2005. A new mode of information transfer in foraging bumblebees? Curr Biol. 15(12):R447–R448. doi:10.1016/j.cub.2005.06.011.

Leadbeater E, Chittka L. 2007a. The dynamics of social learning in an insect model, the bumblebee (Bombus terrestris). Behav Ecol Sociobiol. 61(11):1789–1796. doi:10.1007/s00265-007-0412-4.

Leadbeater E, Chittka L. 2007b. Social Learning in Insects — From Miniature Brains to Consensus Building. Curr Biol. 17(16):R703–R713. doi:10.1016/j.cub.2007.06.012.

Leadbeater E, Dawson EH. 2017. A social insect perspective on the evolution of social learning mechanisms. Proc Natl Acad Sci. 114(30):7838–7845. doi:10.1073/pnas.1620744114.

Leadbeater E, Chittka L. 2009. Bumble-bees learn the value of social cues through experience. Biol Lett. 5(3):310–312. doi:10.1098/rsbl.2008.0692.

Loukola OJ, Gatto E, Híjar-Islas AC, Chittka L. 2020. Selective interspecific information use in the nest choice of solitary bees. Animal Biology. 70(2):215–225. doi:10.1163/15707563-20191233.

Lunau K, Wacht S, Chittka L. 1996. Colour choices of naive bumble bees and their implications for colour perception. J Comp Physiol A. 178(4). doi:10.1007/BF00190178.

Menzel R, Shmida A. 1993. The ecology of flower colours and the natural colour vision of insect pollinators: the Israeli flora as a study case. Biol Rev. 68(1):81–120. doi:10.1111/j.1469-185X.1993.tb00732.x.

Parejo D, Avilés JM. 2016. Social information use by competitors: Resolving the enigma of species coexistence in animals? Ecosphere. 7(5):e01295. doi:10.1002/ecs2.1295.

Raine NE, Chittka L. 2007. The Adaptive Significance of Sensory Bias in a Foraging Context: Floral Colour Preferences in the Bumblebee Bombus terrestris. PLOS ONE. 2(6):e556. doi:10.1371/journal.pone.0000556.

Raine Nigel E., Ings TC, Dornhaus A, Saleh N, Chittka L. 2006. Adaptation, Genetic Drift, Pleiotropy, and History in the Evolution of Bee Foraging Behavior. In: Advances in the Study of Behavior. Vol. 36. Academic Press. p. 305–354.

Raine Nigel E, Ings TC, Ramos-Rodriguez O, Chittka L. 2006. Unterschiede im Lernverhalten zwischen Kolonien einer freilebenden Britischen Hummelpopulation (Hymenoptera: Apidae: Bombus terrestris audax). Entomol Gen. 28(4):241–256. doi:10.1127/entom.gen/28/2006/241.

Rogers SR, Cajamarca P, Tarpy DR, Burrack HJ. 2013. Honey bees and bumble bees respond differently to inter- and intra-specific encounters. Apidologie. 44(6):621–629. doi:10.1007/s13592-013-0210-0.

Sclafani A. 1995. How food preferences are learned: laboratory animal models. Proc Nutr Soc. 54(2):419–427. doi:10.1079/PNS19950011.

Seppänen J-T, Forsman JT, Mönkkönen M, Thomson RL. 2007. Social Information Use Is a Process Across Time, Space, and Ecology, Reaching Heterospecifics. Ecology. 88(7):1622–1633. doi:10.1890/06-1757.1.

Smolla M, Alem S, Chittka L, Shultz S. 2016. Copy-when-uncertain: bumblebees rely on social information when rewards are highly variable. Biol Lett. 12(6):20160188. doi:10.1098/rsbl.2016.0188.

Stephens DW, Brown JS, Ydenberg RC, editors. 2007. Foraging: behavior and ecology. Chicago: University of Chicago Press.

Valone TJ, Templeton JJ. 2002. Public information for the assessment of quality: a widespread social phenomenon. Johnstone RA, Dall SRX, editors. Philos Trans R Soc Lond B Biol Sci. 357(1427):1549–1557. doi:10.1098/rstb.2002.1064.

van Bergen Y, Coolen I, Laland KN. 2004. Nine-spined sticklebacks exploit the most reliable source when public and private information conflict. Proc R Soc Lond B Biol Sci. 271(1542):957–962. doi:10.1098/rspb.2004.2684.

von Frisch K. 1965. Die Tanze der Bienen. In: Tanzsprache und Orientierung der Bienen. Springer, Berlin, Heidelberg. doi: 10.1007/978-3-642-94916-6_2.

Waser N, Chittka L. 1998. Bedazzled by flowers. Nature. 394(6696):835–836.
doi:10.1038/29657.

Xie Z, Pan D, Teichroew J, An J. 2016. The Potential Influence of Bumble Bee
Visitation on Foraging Behaviors and Assemblages of Honey Bees on Squash Flowers
in Highland Agricultural Ecosystems. PLOS ONE. 11(1):e0144590.
doi:10.1371/journal.pone.0144590.

818