CHALLENGED IMMUNE SYSTEM IMPROVES COGNITIVE-BEHAVIORAL RESPONSES IN HOMEOSTASIS AND RECOVERS MALARIA-INDUCED COGNITIVE IMPAIRMENT IN MICE

Luciana Pereira de Sousa*♯, Flávia Lima Ribeiro-Gomes*♯, Roberto Farina de Almeida***+, Tadeu Mello e Souza**, Guilherme Loureiro Werneck***, Diogo Onofre Gomes de Souza** & Cláudio Tadeu Daniel-Ribeiro*1

*Laboratório de Pesquisa em Malária, Instituto Oswaldo Cruz & Centro de Pesquisa, Diagnóstico e Treinamento em Malária (CPD-Mal) of Fundação Oswaldo Cruz (Fiocruz) and of Secretaria de Vigilância em Saúde (SVS), Ministério da Saúde; **Departamento de Bioquímica, Universidade Federal do Rio Grande do Sul and *** Departamento de Epidemiologia of the Instituto de Medicina Social, Universidade do Estado do Rio de Janeiro and Instituto de Estudos de Saúde Coletiva da Universidade Federal do Rio de Janeiro, Brazil.

[1] Corresponding author at : *Laboratório de Pesquisa em Malária, Instituto Oswaldo Cruz, Fiocruz. Av. Brasil 4365, Manguinhos, Rio de janeiro. CEP 2104-360, RJ Brazil. E-mail malaria@fiocruz.br*

♯ These authors contributed equally to the work

+ Present address: *Programa de Pós-Graduação em Ciências Biológicas, Instituto de Ciências Exatas e Biológicas, Universidade Federal de Ouro Preto, Minas Gerais, Brazil*

ABSTRACT

The immune and nervous systems can be categorized as cognitive plastic systems for their ability to know and recognize "real world objects", including microbes, and because of their prerogative of constant self-reorganization as they live learning experiences. After each antigenic or sensory occurrence, vertebrate organisms experience changes in the cellular connections of their immune and nervous systems altering their abilities and structures. Elements of the immune machinery are necessary for neurocognitive function, and the pattern of the immune response triggered by different stimuli may induce regulator or deregulator signals for the nervous functions. Here we show, for the first time, that the immune modulation with anti-inflammatory stimuli can positively regulate the behavior of healthy mice and mitigate the cognitive-behavioral deficits induced by a mild infection of C57BL/6 animals with *Plasmodium berghei* ANKA.

INTRODUCTION

There is considerable evidence for the existence of strong interactions between the immune and nervous cognitive systems[1-7]. Immunomodulation of the nervous system does exist and can either be physiological or pathological. The maturation and homeostasis of the nervous cognitive abilities require the participation of elements of the immune system[6-7]. Exogenous immune stimuli may also have positive or negative effect on the nervous system, depending on the nature and intensity of the immune response elicited[1-4,6]. Different studies, aiming to assess the effect of immune stimuli on brain function, reported i) the influence of maternal immune stimulation impairing the neurocognitive

44 performance of offspring[8-9], ii) both beneficial and harmful effects of neonate vaccination on neuronal
45 plasticity and cognitive function in adulthood[10], and iii) the damaging impact of inflammatory stimuli
46 on the cognitive function of adult mice[11-12]. Some infections may also cause neurocognitive dysfunction
47 in human and experimental models[13-27].

48
Cerebral malaria (CM), the most severe complication of falciparum malaria, can be
accompanied by neurocognitive sequelae, including severe motor deficit, behavioral
alterations and severe learning difficulties[15]. Long-term sequelae, responsible for child
ineptitude, are more common in Africa, where the prevalence of *falciparum* malaria
and of CM is higher[28]. Some of these sequelae are also observed in *Plasmodium*
berghei ANKA (*Pb*A) infected C57BL/6 mice, the most classical model of experimental
CM[27.] In recent years, cognitive impairment, mainly related to learning and memory,
has also been reported in residents of endemic regions presenting non-cerebral
malaria (non-CM) episodes, but has not been recorded in the common murine
experimental models of non-CM[29-31].

In a previous work, we have unprecedentedly adapted the classical experimental
model of CM (*Pb*A-infected C57BL/6 mice) to study neurocognitive alterations after a
short-term episode of non-CM[32]. Using this model, we evaluated here the effect of
immune stimuli on behavioral paradigms, such as memory and anxious phenotype,
during homeostasis or after a mild infection in mice.

Considering the data accumulated so far showing the effect of immune system on the
performance of neurocognitive functions, we hypothesize that some types of immune
stimulation may improve cognitive performance. Our data suggest an inedited
beneficial effect of immunization, with antigens that drive T2 immune responses, on
healthy mice cognitive performance and the recover of the cognitive impairment
associated to plasmodium infectious stress.
70
RESULTS
T2-immune stimuli improve long-term memory in healthy mice
The immune stimuli were categorized according to the profile of cytokine induced by
the specific properties of the studied immunogens. Thus, pro-inflammatory immune
stimuli trigger Th1 cell responses with predominance of IFNγ and TNF-α production
whereas polarization to an anti-inflammatory immune response profile (Th2) is
dependent on T-cell derived IL-4[33-47]. Also, levels of certain antibody isotypes may
indicate targeting of the immune response profile, such as IgG1 and IgG2a production
indicating Th2 and Th1 responses profiles, respectively[42]. Moreover, adjuvants added
to immune stimuli also play an important role in directing the immune response[43]. To
study the effect of immune stimuli on behavioral paradigms, we employed three
immunization strategies according to their pro- and anti-inflammatory immune
response profiles. T1 strategy was used to induce a pro-inflammatory immune
response and T2 to elicit immune responses of type 2[33-47]. The Pool of antigens, for a
combined T1 and T2 strategy, was also used.

86
87 To address the effect of immune responses on locomotion and on long-term
88 habituation, mice were submitted to the open field task (OFT), with a training and a
89 test sessions, 24 hours apart, assessed in a well-established protocol[32]. Locomotion is
90 evaluated when mice access the open field arena for the first time. At the training

session, a high rate of locomotor activity is commonly observed. Surprisingly, Pool and T1 immunized mice showed a reduced total OFT1 locomotion, when compared to non-immunized mice, and T2 immunized animals showed a clear trend towards a decreased locomotion (Extended data, Fig1a) Commonly, after the training session, the exploratory behavior decreases, as the stress related to the novelty disappears, and is significantly lower after 10 minutes of task performance[32,48-49]. Control and all immunized groups of mice presented a decreased locomotion in the test session (OFT2) as compared to the training session (Extended data, Fig1a). These results indicate that the immunization did not affect the long-term habituation memory accessed 24 hours after training.

Mice were then subjected to the novel object recognition test (NORT) in the same Open Field arena. On the training session, a similar exploratory activity of both familiar (FO1 or FO2) objects is expected and was observed in all groups (Control, Pool, T1 and T2) of mice (Fig.1a; Extended data, Fig.1c), exploring them for a mean of 25 seconds (data not shown). Remarkably, as we hypothesized, the groups of mice immunized with the Pool and T2 strategies presented significantly higher recognition memory performance in relation to the control group of non-immunized mice during the test session (Fig.1c; Extended data Fig.1e). These data indicate, for the first time, that an immune response classically categorized as anti-inflammatory (induced by T2 stimuli) may enhance long-term recognition memory in healthy mice.

Immunization of healthy mice did not influence their anxiety-like state
In addition to the exploratory activity, the OFT also allows the evaluation of phenotypes related to anxiety-like behavior through the dwell time or the locomotion rate in the center of the open field, during the first exposition to the apparatus. Immunized mice (Pool, T1 and T2) showed no difference in time (data not show) but presented a significantly reduced locomotion in the center of the open field in relation to non-immunized control group (Fig. 1e). However, these data may have been impacted by the total reduced mobility observed in these animals (Extended data, Fig. 1a), and no conclusion about anxious behavior can be scratched from these data. To address this issue, we used another approach-avoidance conflict test, the classical light-dark specific task, to evaluate the anxiety-like behavior. In this test, immunized mice (Pool, T1 and T2) groups clearly behaved similarly to mice of the control group, remaining during an equal time in the light zone (Fig.1g), implying that immunization did not change the anxiety phenotype.

T2-immune stimuli may recover the cognitive-behavioral damage caused by non-complicated *P. berghei* ANKA infection
About 92% of the world's malaria cases are due to *Plasmodium falciparum* infection, 1 to 2% of which progress to cerebral malaria that accounts for about 80% of deaths due to malaria in the world. Thus, about 90% of malaria cases globally are caused by a lethal species of *Plasmodium*, but occur without clinical complications[28] and may also present with cognitive impairment, indicating that non-complicated malaria can impact not only the children health but also their cognitive development[29-31].

The experimental model we have described uses *Pb*A-infected C57BL/6 treated at day 4 post-infection, before appearance of any clinical sign of CM. The main advantage of such a model is that it best mimics the human situation described above that

corresponds to the largest number of malaria cases in the world (uncomplicated
falciparum malaria treated timely)[28]. Using this model, we have been able to observe
a long-term cognitive-behavioral impairment related to memory and anxious
phenotype[32]. With the evidence of a beneficial effect of immunization on the long-term
memory in healthy mice described above, we decided to evaluate the effect of the
same immune stimuli in mice with behavioral alterations resulting from non-CM.

The behavioral tests were performed 77 days after Chloroquine (CQ) treatment of
infected mice. *Pb*A-infected and treated mice did not present total mobility reduction in
the training session of OFT (Extended data, Fig.1b) when compared to non-infected
mice. However, infected-immunized (Pool, T1 or T2) animals showed a trend towards
a reduction of locomotion in OFT, when compared to infected mice (Extended data,
Fig.1b). This trend becomes a significant reduction when locomotory activity of
infected-immunized mice (Pool and T2) is matched to that of healthy ones (Extended
data, Fig.1b).

As expected, there was no object preference in the NORT training session since all
mice explored similarly (for a mean of 25 seconds - data not show) both familiar objects
(Fig.1b, Extended data, Fig.1d). Control, infected and infected-immunized groups
(Pool and T2, but not T1) of mice presented normal behavior with a significant
decreased locomotion in the test session as compared to the training session of the
OFT (Extended data, Fig. 1b). Consistently, non-CM recovered mice presented long-
term recognition memory sequelae expressed by similar exploration of the FO and NO
in NORT. As hypothesized, this impairment disappeared after the use of the Pool and
T2 strategies of immunization (Fig.1d, Extended data, Fig.1f), pointing to the foreseen
beneficial effect of anti-inflammatory T2 immune responses reversing cognitive deficit
associated to malaria.

P. berghei ANKA infection of mice induces an anxiety-like behavior that is
reverted by the T2 immunization strategy
The distances travelled in the periphery and in the center of the OFT are inversely
related. Since the latter was decreased in *Pb*A infected recovered mice (Fig.1f) and no
OFT1 locomotion change occurred, one can understand this effect as the expression
of an anxiety-like behavior (Extended data, Fig.2b). This anxiety-like behavior was also
detected in the light-dark task, since the infection also reduced the time spent in the
light zone of the apparatus. More importantly, this decrease was reversed by Pool and
T2 immunization (Fig.1h).

This is in accordance with the increased distance travelled in the periphery of the OFT
(data not show). This anxiety-like phenotype was, effectively, confirmed by the
reduction of time spent in the light zone of light-dark task. This decrease was reversed
by the immunization with Pool and T2 strategies (Fig.1h).
177
Immunization procedures and non-complicated _P. berghei_ ANKA malaria do
elicit specific immune responses
Pool and T2 immunized mice showed increased spleen weight and total number of
splenocytes (Extended data, Fig. 2a,b). The specific immune responses triggered by
the immunogens used were evaluated and the effectiveness of the stimuli was
confirmed (Extended data, Fig.3a,b,c,d). We also evaluated the frequencies of B and
T cells, as well as the T cell activation status   in the spleen, at the end of the cognitive

task evaluation. Mice immunized with either the Pool or the T2 strategies (Pool and T2 groups) presented similar patterns of modulation of different immune components analyzed. We observed, in both groups, an increase in the frequency of B cells, CD4 and CD8 T cells with central memory phenotype and CD4 T cells with regulatory function when compared to non-immunized animals (Extended data Fig.4a,b,c,d). A reduction in the frequency of splenic CD8 T cells was also recorded mice immunized with the T2 stimuli, when compared to the control group (Extended data, Fig.5a). Besides, mice immunized with the Pool and T2 strategies presented increased serum levels of IL-4 cytokine, TNFα and IFNγ (Extended data, Fig.6a,b,d).

Alterations in the immune response profile were also evaluated late (77 days following treatment) after *Pb*A infection. *Pb*A-infected and treated mice showed higher frequency of B cells, CD4 and CD8 T cells and increase in the frequency of CD4 and CD8 T cells with naïve and central memory phenotype (Extended data, Fig.4a,b,c,e,f; Fig 5a,b). The frequency of regulatory CD4 T cells, however, was similar between infected and control mice (Extended data Fig.4d). Also, at the time they were immunologically evaluated, infected and treated animals did not present detectable levels of serum cytokines (Extended data, Fig.6a,b,c,d) and only showed an altered shade color, attributable to the hemozoin pigment of the erythrocytic cycle of the *Plasmodium* parasite, with no more difference in the weight and total number of splenocytes in relation to the control group (Extended data, Fig.2a,b,c). It is clear, therefore, that alterations of the immune system are triggered by plasmodium, even though the infection is treated and contained rapidly, and that some of them can be observed several weeks after parasite elimination.

As observed in healthy immunized animals, immunization of *Pb*A-infected and treated mice with Pool or T2 strategies induced splenomegaly (Extended data, Fig2a,b,c), humoral immune response to the vaccine stimuli (Extended Data, Fig.3a,b,c,d) and patterns of splenic B cells, CD4 and CD8 T cells and of their different function subsets comparable for the two stimuli (Extended data, Fig.4a,c,d,e; Fig.5a,b). The marked splenomegaly recorded after immunization could partially explain the reduced total locomotion in OFT (Extended data, Fig.2a,b) and harmed conclusions on the effect of immunization on anxiety-like phenotype in this behavioral task (Fig.1e,f).

It is important to highlight that, in general, the object exploratory pattern in NORT training test was not modified in any feature considering the most rigorous aspects reported in NORT protocol[50], which sustain our interpretation about the beneficial effect of anti-inflammatory T2 immune responses reversing cognitive deficit associated to malaria.

Taken together, our data point to a positive influence of immune responses induced by T2 stimuli on the long-term memory in healthy mice, confirm our previous demonstration of neurocognitive behavioral dysfunction after a single episode of non-cerebral malaria, and indicate a recovering effect of this deficit exerted by T2 immunization procedures.

DISCUSSION

The existence of lymphocyte neuropeptide receptors[51] and cytokine receptors in neurons[52] are commonplace notions that have paved the anatomical and functional basis of the "classical neuroimmunology". On the other hand, as the immune and nervous systems are arrangements with cognitive functions and plastic properties, it is reasonable to assume that learning in one cognitive system may influence the performance of the other.

Different studies have shown that stimulation of the immune system may exert different effects on neurocognitive function depending on the nature and balance of the immune inputs[6,51-52] and the cumulated knowledge points to the notion that, while anti-inflammatory immune responses are responsible for the proper performance of the brain[51,53], pro-inflammatory responses can impair cerebral function and neurocognitive activity[6].

Stimulus such as physical exercise, reading and music are mentioned cognitive enhancement strategy[56-59]. Here, we describe, for the first time, beneficial immunomodulatory effects improving cognition in healthy mice. Our findings show a marked positive effect of immune stimuli specifically triggering type-2 immune response by improving long-term memory of normal adult mice, as verified by the new object recognition task (NORT), a robust and most used behavioral tasks for the analysis of the parameter in rodents[50].

We have previously reported cognitive-behavioral impairment as late sequelae of a single uncomplicated malaria episode using a MC experimental model and treating animals before the appearance of any neurological sign[60]. In our concept, the model is illustrative and adequate for the study of the, by far, most common presentation of malaria; ie: the non-cerebral *Plasmodium falciparum* malaria[61]. In fact, both the experimental model and the human situation correspond to conditions in which we have a *Plasmodium* parasite capable of causing - and a host susceptible of developing - cerebral malaria that is, however, prevented by timely treatment.

As hypothesized, we were able to observe a positive effect of immune stimuli reversing cognitive-behavioral impairment associated with non-complicated malaria (nCM) in the prototype described above. The results show that mice with nCM, treated at D4 of *Pb*A infection and immunized with T2 and Pool strategies for 60 days from D11 on, did not present the long-term memory sequelae - expressed by similar exploration of the FO and NO in NORT - as non-immunized infected mice did. Besides, we reported anxious-like behavior reversal in the light-dark task, also considered a reference tool for specifically analyzing anxiety[62-63], after immunization of infected and treated mice.

These results confirm our conceived idea of an effect of immune stimuli on brain function and downstream consequences reflected in improvement of the cognitive-behavioral performance. It is know that inflammatory stimuli may cause microgial activation and increase of cytokine T1 profile such as TNF-α and IL1-β reported as evil for cognitive function[64] while cytokine T2 profile are beneficial cognitive enhancers such as IL-4 referred to as cytokine to remember[35]. It has also been shown that LPS can cause the basal ganglia to malfunction but the dosages applied to promote this effect are higher than that used in the present study[65].

The inedited data reported here may correspond to a new paradigm and promising alternative for the design of memory improvement strategies in homeostasis and cognitive function impairment, including those associated to malaria and reflected as poor school performance in hundreds of thousands of children in the world. In fact, the relevance of maintaining an updated vaccination calendar is usually claimed for the sustenance of efficient policies for prophylaxis of "immune-preventable" diseases, through individual and collective protection against infections and prevention of epidemics. Our data may suggest that a new thought on the importance of vaccination may arise from its potential use as a strategy for boosting cognition function in health and recovering cognitive dysfunctions in infectious and chronic diseases and ageing. This new perceptiveness of the vaccination benefits could constitute a new weapon to fight the growing anti-vaccine movement in the world.

REFERENCES

1. Jerne NK. Idiotypic networks theory and other preconceived ideas. Immunological Reviews. 1984, 79: 5-24.
2. Cohen, I. The cognitive paradigm and the immunological homunculus. Immunol. Tod. 1992; 13(12):490-4.
3. Cohen I. On autoimmunity. Tending Adam's Garden: evolving the cognitive immune self. Acedemic Press. 2000.
4. Daniel-Ribeiro CT & Martins YC. Imagens, Micróbios e Espelhos: os sistemas imune e nervoso e nossa relação com o ambiente. Ed. Fiocruz. 2017, Cap. 3: pp 161-162.
5. Kivisakk P, et al. Localizing central nervous system immune surveillance: meningeal antigenpresenting cells activate T cells during experimental autoimmune encephalomyelitis. Ann. Neurol. 2009; 65:457–469.
6. Kipnis J, Gadini S, Derecki N. Pro-cognitive properties of T cells. Nat Rev Immunol. 2012; 12 (9): 663-669.
7. Nataf S. Autoimmunity as a Driving Force of Cognitive Evolution. Front. Neurosc. 2017, 11:582.
8. Schwartzer JJ, Careagab M, Coburna MA, Roseb RD, Hughes HK, Paul Ashwood P. Behavioral impact of maternal allergic-asthma in two genetically distinct mouse strains. Bra Beh Immunit. 2017, 63: 99–107.
9. Mueller F, Polesel M, Richetto J, Meyer U, Weber-Stadlbauer U. Mouse models of maternal immune activation: Mind your caging system! Bra Beh and Immunit. 2018, 643-660.
10. Yang J, Qi F, Zou HGJ, Yang Y, Yuan Q, Yao Z. Neonatal BCG vaccination of mice improves neurogenesis and behavior in early life. Brai Resear Bulleet. 2016, 120: 25–33.
11. Bossu P, Cutuli D, Palladino I, Caporali P, Angelucci F, Laricchiuta GF, et. al. A single intraperitoneal injection of endotoxin in rats induces long-lasting modifications in behavior and brain protein levels of TNF-α and IL-18. J. of Neuroimmunofl. 2012, 9:101.
12. Depino AM. Early prenatal exposure to LPS results in anxiety- and depression-related behaviors in adulthood. Neuroscience. 2015, 299: 56-65.

13. Kannan G, Pletnikov MV. Toxoplasma gondii and Cognitive Deficits in Schizophrenia: An Animal Model Perspective. Schizoph Bullet. 2012; 38 (6): 1155–1161.

14. Iqbal J,Mueller U. 2007. Virus infection causes specific learning deficits in honeybee foragers. Proceedings of the Royal Society B-Biological Sciences 274:1517–1521.

15. Odera VM, Snow RW, Newton CRJC. The burden of the neurocognitive impairment associated with Plasmodium falciparum malaria in sub-Saharan Africa. Am J Trop Med Hyg. 2004;71:64–70.

16. Carter JA, Lee JA, Gona JK, Murira G, Rimba K, Neville BG, et al. Severe falciparum malaria and acquired childhood language disorder. Dev Med Child Neurol. 2006;48:51–7.

17. Boivin MJ, Bangirana P, Byarugaba J, Opoka RO, Idro R, Jurek AM, et al. Cognitive impairment after cerebral malaria in children: a prospective study. Pediatrics. 2007;119:e360–6.

18. John CC, Bangirana P, Byarugaba J, Opoka RO, Idro R, Jurek AM, et al. Cerebral malaria in children is associated with long-term cognitive impairment. Pediatrics. 2008; 122:e92-9.

19. Kihara M; Carter J.A; Holding P.A; Khadem F.V; Scott R.C; Idro R; Fegan G.W; Haan M; Neville B.G.R; Newton C.R.J.C. Malaria journal, v. 8, p. 273, 2009.

20. Holmberg D, Franzén- Rohl E, Idro R, Opoka RO, Bangirana P, Sellgren CM et. al. Cerebrospinal fuid kynurenine and kynurenic acid concentrations are associated with coma duration and long-term neurocognitive impairment in Ugandan children with cerebral malaria. Mal J. 2017, 16:303.

21. Thuilliez J, Sissoko MS, Toure OB, Kamate P, Berthelemy JC, Doumbo OK. Malaria and primary education in Mali: a longitudinal study in the village of Donéguébougou. Soc Sci Med. 2010;71:324–34.

22. Bangirana P, Allebeck P, Boivin MJ, John CC, Page C, Ehnvall A, et al. Cognition, behaviour and academic skills after cognitive rehabilitation in Uganda children surviving severe malaria: a randomised trial. BMC Neurol. 2011;11:96.

23. Fink G, Olgiati A, Hawela M, Miller JM, Matafwali B. Association between early childhood exposure to malaria and children's pre-school development evidence from Zambia early childhood development project. Malar J. 2013;12:12.

24. Vorasan N, Pan-Ngum W, Jittamala P, Maneeboonyang W, Rukmanee P, Lawpoolsri S. Long-term impact of the childhood malaria infection on school performance among school children in a malaria endemic area along the Thai-Myanmar border. Malar J. 2015;14:401.

25. Reverchon F, Mortaud S, Sivoyon M, Maillet I, Laugeray A, Palomo J, et. al. IL-33 receptor ST2 regulates the cognitive impairments associated with experimental cerebral malaria. Plos Pathog. 2017, 13(4): e1006322.

26. Desruisseaux MS, Gulinello M, Smith DN, Lee SC, Tsuji M, Weiss LM, et al. Cognitive dysfunction in mice infected with Plasmodium berghei strain ANKA. J Infect Dis. 2008;197:1621–7.

27. Reis PA, Comim CM, Hermani F, Silva B, Barichello T, Portella AC, et al. Cognitive dysfunction is sustained after rescue therapy in experimental cerebral malaria, and is reduced by additive antioxidant therapy. PloS Pathog. 2010;6:e1000963.

28. World Health Organization. World Malaria Report. Geneva: WHO; 2017.

29. Fernando SD, Gunawardena DM, Bandara MR, De Silva D, Carter R, Mendis KN, et al. The impact of repeated malaria attacks on the school performance of children. Am J Trop Med Hyg. 2003;69:582–8.
30. Vitor-Silva S, Reyes-Lecca RC, Pinheiro TR, Lacerda MV. Malaria is associated with poor school performance in endemic area of the Brazilian Amazon. Malar J. 2009;8:230.
31. Tapajós R, Castro D, Gisely Melo G, Balogun S, James M, Pessoa R, Almeida A, Costa M, Pinto R, Albuquerque B, Monteiro W, Braga J, Lacerda M & Mourão MP. Malaria impact on cognitive function of children in a peri-urban community in the Brazilian Amazon. Malar J. 2019; 18:173.
32. de Sousa LP, Almeida RF, Ribeiro-Gomes FL, Carvalho LJM, Souza TM, Souza DOG & Daniel-Ribeiro CT. Long-term effect of uncomplicated Plasmodium berghei ANKA malaria on memory and anxiety-like behaviour in C57BL/6 mice. Parasites & Vectors. 2018; 11:191.
33. Lavigne MV, Castro M, Mateo N, Deluchi S, Atzori C. Whole-cell *Bordetella pertussis* vaccine component modulates the mouse immune response to an unrelated soluble antigen. Microb. and Infect. 2002; 815-820.
34. Castilho SF, Chovel ML, Hernández NG, González LC, Blanco A, Hernández DS, Medina MF, Tito MA, Quiñoy JLP. A Bordetella pertussis proteoliposome induces protection in mice without affecting the immunogenicity of diphtheria and tetanus toxoids in a trivalent formulation. Clinic. and Experiment. Vacc.Reser. 2016; 5:175-178.
35. Goodier MR, Rodriguez-Galan A, Lusa C, Nielsen CM, Darboe A, Moldoveanu AL, White MJ, Behrens R, Riley EM. Influenza Vaccination Generates Cytokine-Induced Memory-like NK Cells: Impact of Human Cytomegalovirus Infection. J. Immunol. 2016; 197(1):313-25.
36. Zhu W, Li S, Wang C, Yu G, Prausnitz MR, Wang BZ. Enhanced immune responses conferring cross-Protection by skin Vaccination With a Tri-component influenza Vaccine Using a Microneedle Patch. Front. in Immunol. 2018; 9:1705.
37. Brahimi K, Pérignon JL, Bossus M, Gras H, Tartar A, Druilhe P. Fast immunopurification of small amounts of specific antibodies on peptides bound to ELISA plates. J. Immunol. Meth. 1993; 162(1):69-75.
38. Oeuvray C, Bouharoun-Tayoun H, Grass-Masse H, Lepers JP, Ralamboranto L, Tartar A, Druilhe P. A novel merozoite surface antigen of Plasmodium falciparum (MSP-3) identified by cellular-antibody cooperative mechanism antigenicity and biological activity of antibodies. Mem. Inst. Oswaldo Cruz. 1994; 89 Suppl 2:77-80.
39. Daher LJ1, Demanga CG, Prieur E, Pérignon JL, Bouharoun-Tayoun H, Druilhe P. Toward the rational design of a malaria vaccine construct using the MSP3 family as an example: contribution of immunogenicity studies in models. Infect. Immun. 2010; 78(1):477-85.
40. Matos DCS, Marcovistz R, Cabello PH, Georgini AR, Sakauchi D, Silva LL. Immunogenicity Test of Tetanus Component in Adsorbed Vaccines by Toxin Binding Inhibition Test. Mem Inst Oswaldo Cruz. 2002, 97(6): 909-913.
41. Lima MCR, Prouvost-Danon A, Patricia MR, Chagas MS, Calheiros AS, Cordeiro RSB. Studies on the mechanisms involved in antigen-evoked pleural. J of Leuk Biol. 1997, 61.

42. Comoy EE, Capron A, Thyphronitis G. In vivo induction of type 1 and 2 immune responses against protein antigens. Int. Immunol. 1997; 9(4):523-31.

43. Terhune TD, Deth RC. Aluminum Adjuvant-Containing Vaccines in the Context of the Hygiene Hypothesis: A Risk Factor for Eosinophilia and Allergy in a Genetically Susceptible Subpopulation?. Internat. J. of Environment. Researc. and Publ. Health. 2018; 15:901.

44. Bosmann M, Russkampa NF, Warda PA. Fingerprinting of the TLR4-induced acute inflammatory response. Exp. Mol. Pathol. 2012; 93(3): 319–323.

45. Kozyreva VS, Subbotovskaia AI, Shilova AN, Karpenko AA. Immunological aspects of formation of restenoses after endothelial lesions. Angiol Sosud Khir. 2014; 20(1):21-6.

46. Couch Y, Trofimov A, Markova3 N, Nikolenko V, Steinbusch HW, Chekhonin V, Schroeter C, Lesch KP, Anthony DC, Strekalova T. Low-dose lipopolysaccharide (LPS) inhibits aggressive and augments depressive behaviours in a chronic mild stress model in mice. J. of Neuroinflamm. 2016; 13:108.

47. Fontana MF, Baccarella A, Kellar D, Oniskey TK, Terinate P, Rosenberg SD et. al. Myeloid expression of the AP-1 transcription factor JUNB modulates outcomes of type 1 and type 2 parasitic infections. 2015, 37(9): 470–478.

48. Almeida RF, Ganzella M, Machado DG, Loureiro SO, Leffa D, Quiconzes Santos A, et al. Olfactory bulbectomy in mice triggers transient and longlasting behavioural impairments and biochemical hippocampal disturbances. Prog Neuropsychopharmacol Biol Psychiatry. 2017;76:1-11.

49. Figueiredo CP, Clarke JR, Ledo JH, Ribeiro FC, Costa CV, Melo HM, et al. Memantine rescues transient cognitive impairment caused by highmolecular-weight aβ oligomers but not the persistent impairment induced by low-molecular-weight oligomers. J Neurosci. 2013;33:9626–34.

50. Leger M, Quiedeville A, Bouet V, Haelewyn B, Boulouard M, Schumann-Bard P, Freret T. Object recognition test in mice. Nat Protoc. 2013; 8(12):2531-7.

51. Gandani SP, Cronk JC, Norris GT, Kipnis J. Interleukin-4: A Cytokine to Remember. J. Immunol. 2012, 189(9): 4213–4219.

52. Brombacher TM, De Gouveia KS, Cruywagen L, Makena N, Booley F, Tamgue O, et al. Nippostrongylus brasiliensis infection leads to impaired reference memory and myeloid cell interference. Nature. 2018; 8:2958.

53. Brynskikh A, Warren T, Zhu Z, Kipnis J. Adaptive immunity affects learning behavior in mice. Bran, Behav and Immun. 2008; 22: 861-869.

54. Zuo Z, Qi F, Yang J, Wang X, YaruWen Y, Yuan Q. Immunization with Bacillus Calmette-Guérin (BCG) alleviates neuroinflammation and cognitive deficits in APP / PS1 mice via the recruitment of inflammation-resolving monocytes to the brain. Neurob of Disea. 2017, 101: 27–39.

55. Gu CJ, Borjabad A, Hadas E, Kelschenbach J, Kim BH, Chao W et. al. EcoHIV infection of mice establishes latent viral reservoirs in T cells and active viral reservoirs in macrophages that are sufficient for induction of neurocognitive impairment. Plos Pathog. 2018; 14(6):e1007061.

56. Voss MW, Weng TB, Narayana-Kumanan K, Cole RC, et. al. Acute Exercise Effects Predict Training Change in Cognition and Connectivity. Med Sci Sports Exerc. 2019.

57. Moreau D, Chou E. The Acute Effect of High-Intensity Exercise on Executive Function: A Meta-Analysis.PerspectPsychol Sci. 2019;14(5): 734-764.

58. Ji Z, Feng T, Mei L, Li Aand Zhang C. Influence of acute combined physical and cognitive exercise on cognitive function:an NIRS study. PeerJ. 2018; 12.

59. Loprinzi PD, Harris F, McRaney K, Chism M, Deming R, Jones T, Zou L, Tan M. Effects of Acute Exercise and Learning Strategy Implementation on Memory Function.Medicina (Kaunas). 2019; 5: 55 (9).

60. Potter S, Chan-Ling T, Ball HJ, Mansour H, Mitchell A, Maluish L, et al. Perforin mediated apoptosis of cerebral microvascular endotelial cells during experimental cerebral malaria. Int J Parasitol. 2006;36:485–96.

61. Worl Malaria Report. 2018. World Health Organization.

62. Almeida RF, Comasseto DD, Ramos DB, Hansel G, Zimmer ER, Loureiro SO, et al. Guanosine anxiolytic-like effect involves adenosinergic and glutamatergic neurotransmitter systems. Mol Neurobiol. 2016;54:423-36.

63. Griebel G, Holmes A. 50 years of hurdles and hope in anxiolytic drug discovery. Nat Rev Drug Discov. 2013; 12(9):667-87.

64. Hoogland ICM, Houbolt C, Westerloo DJV, Gool WAV and Beek D. Systemic inflammation and microglial. Journal of Neuroinflammation. 2015; 12:114.

65. Ramsey CP, Tansey MG. A survey from 2012 of evidence for the role of neuroinflammation in neurotoxin animal models of Parkinson's disease and potential molecular targets. Ex. Neurol. 2014; 256:126-32.

ACKNOWLEDGEMENTS

LPS is grateful to the *Programa de Pós-Graduação em Biologia Parasitária* of the *Instituto Oswaldo Cruz (IOC) of Fundação Oswaldo Cruz (Fiocruz)* for the Doctoral fellowship. The authors are grateful to the *Laboratório de Inflamação* (Dr. Marco Aurélio Martins and Dr. Tatiana Ferreira), *Laboratório de Pesquisa em Malária* (Luana santos Thalita Ferraz), *Laboratório de Pesquisa sobre o Timo* of the *IOC-Fiocruz* (Dr Daniella Mendes Arêas and Dr Dyna Raposo) of *IOC; Biomanguinhos* (Dr. Maria de Lourdes de Sousa Maia, Alessandro Fonseca, Camilla Bayma, and Denise Cristina Matos) and *Farmanguinhos* (Dr. Márcia Coronha Ramos Lima and Dr. Andréa Luca) of *Fiocruz and Instituto Butantan* (Dr. Jorge Kalil and Dr. Paulo Lee Ho and Aline Abrantes) and Vac4all (Dr. Pierre Duiilhe) for reagent supply and study of the immune response to vaccines.

FUNDING

This work is part of LPS's PhD research supported by CAPES (Brazil) and by Faperj (RJ, Brazil) fellowships. GLW, DOS and CTDR are supported by CNPq, Brazil, through a Productivity Research Fellowship and GLW and CTDR are "*Cientistas do Nosso Estado*" recognized by the Faperj. The *Laboratório de Pesquisa em Malária, IOC, Fiocruz* and the *Departamento de Bioquímica* of the *Universidade Federal do Rio Grande do Sul* are National Institutes of Science & Technology (INCT) associated Laboratories.

AUTHOR CONTRIBUTIONS

LPS was responsible for the realization of all experiments (including infection and treatment; immune stimulation, and conduction, observation and data collection/systematisation of cognitive tests and immune response analyses in mice), helped in the analysis and interpretation of tests and drafted the manuscript. FLRG followed all the stages of the experiment, realization of experiments, discussed the protocols and the project, was in charge of the analysis and discussion of immune response data and helped in drafting the manuscript. RFA and TMS helped in systematization of data concerning behavioral tests and analyzed and interpreted the cognitive data. GW proposed the statistical analyzes of the data and was responsible for it. DOS discussed the project since its conception and helped in designing the experiments. CTDR is responsible for conception and design of the study, and helped in data analysis, interpretation and drafting and finalizing the manuscript together with LPSV and FLRG. All authors read, reviewed and approved the final manuscript.

COMPETING INTERESTS

The authors declare that they have no competing interests.

METHODS

Mice and Parasite. The "Instituto de Ciência e Tecnologia em Biomodelos" of the "Fundação Oswaldo Cruz" (ICTB- Fiocruz, Brazil) granted seven week-old C57BL/6 female mice weighing 20-25 g. Mice were conditioned housed in racks with an air filtration system in a room maintained at 25°C and light/dark cycles of 12 hours in cages containing five animals with free acquisition to food and water. All procedures were carried out in accordance with animal welfare approved by the Ethical Committee on the Use of Laboratory Animals of "Instituto Oswaldo Cruz" under CEUA-IOC: L-010/2015 concession. *Plasmodium berghei* ANKA (*Pb*A) infections were carried out using a stable transfected strain of *Pb*A expressing a green fluorescent protein (*Pb*A-GFP) generated as described previously[1].

Infection and treatment of experimental groups. C57BL/6 mice were infected intraperitoneally (ip) with 150 µl of *Pb*A-infected red blood cells, duly cryopreserved and thawed, being called "mice-passage." Five days after infection, the total blood was collected, adjusted to 1 x10^6 parasitized erythrocytes in 100 µl of PBS and injected in the peritoneum of C57BL/6 mice from the experimental groups. Parasitemia was monitored by flow cytometry, based on the percentage of GFP+ erythrocytes. In this experimental model, the evolution to and establishment of cerebral malaria (CM) occur between the fifth and sixth day of infection[2]. In this study, mice were treated on the fourth day of infection (mean parasitaemia 2.5%) with 25 mg/kg of chloroquine (CQ) by gavage for seven days[3], before any clinical signs of CM. All groups were similarly manipulated on procedures. Experiments carried out with groups of uninfected mice treated with CQ or not (control group received PBS) have previously shown, that the CQ treatment did not influence the performance in behavioral tasks and anxiety phenotype[4].

Experimental Description. C57BL/6 mice were divided into groups of *Pb*A infected and Control animals (no infected) and both were treated with chloroquine (CQ) for seven days from the fourth day of infection. Thirteen days after treatment, mice from respective groups were subdivided into Non-Immunized and Immunized groups (Fig. 1). The following vaccines and antigens were used for immunization: Diphtheria and Tetanus toxoids (dT) vaccine for adults, Influenza vaccine, *Plasmodium falciparum* Merozoite Surface Protein 3 (pfMSP-3), White chicken egg ovalbumin (OVA) and Lipopolysaccharide of *Escherichia coli* (ecLPS). Three different immunization strategies were performed: a combination of all antigens and vaccines described above (from here now, called Pool); a combination of antigens and vaccines (Influenza vaccine and ecLPS) that trigger, preferentially, a type 1 pattern of immune response (from here now, denominated T1); and a combination of antigens and vaccines (dT vaccine, pfMSP-3 and OVA) that trigger, preferentially, a type 2 pattern of immune response (from here now, called T2). The groups of mice were denominated: Control (Non-infected/Non-Immunized); Pool, T1 and T2 (Non-infected/Immunized Pool, T1 or T2 of the immunization strategy); Inf (Infected/Non-immunized); and Inf-Pool, T1 or T2 (Infected/Immunized Pool, T1 or T2 of the immunogens inoculation strategy). Subsequently, mice behavioral performance was assessed by Open Field Test (OFT), Novel Object Recognition Test (NORT) and Light-Dark (L/D) (Fig. 2). About 300 mice were used for these experimental strategies in five consecutive sessions.

Immune system stimuli. The immune stimulus was initiated thirteen days after the end of CQ treatment, being performed in the course of the following sixty days (Fig. 2). Antigens and/or vaccines were administered by different routes, in different regions of the animals's body (Table 1). The doses administrated were defined based on dose-response protocols available in the literature capable of stimulating the murine immune system without imparting risk of death to mice immunized[5-12].

Immunization with Plasmodium falciparum Merozoite Surface Protein 3 (pfMSP-3). The mice were challenged with 10 µg of pfMSP-3 recombinant (in collaboration with Clinical Trials of Malaria Vaccines (Vac4All Initiative), Paris, France) adsorbed on 70% adjuvant solution MONTANIDE[TM] ISA 50 V2 W / O (SEPPIC. Air Liquide - Healthcare), in 100 µl of PBS. Three subcutaneous injection were performed at the tail region with a twenty-day interval between immunizations[9-10] (Fig. 2, Table 1).

Immunization with Tetanus- Diphtheria and Influenza Vaccines. The vaccines used in this study were: bacterial double [(Tetanus-Diphtheria (dT): Biological E Limited - BE, Telangana - India, Lot. 34005815. In collaboration with the Division of Health Surveillance - CAP 3.1 of the "Fundação Oswaldo Cruz", Rio de Janeiro, Brazil] and Trivalent Influenza (granted by the Technological Development and Production Division of "Instituto Butantan, São Paulo, Brazil, Lot. 160034).

Mice received 100 µl (1/5 of the human dose) of dT and Influenza vaccines by subcutaneous (dorsal region) and intramuscular (left quadriceps region), routes respectively (Table 1). Three inoculations with a twenty-day interval between immunizations were performed[5-6] (Figure 2).

Immunization with Ovalbumin (allergen). Mice received 50 µg of white chicken egg ovalbumin (SIGMA-ALDERICH, Cod. A5503-50g) adsorbed onto aluminum hydroxide [Al (OH) 3] in a final volume of 200 µl per animal in three inoculations. The first inoculation was performed at the dorsal region by subcutaneous injection and the following (second and third inoculation) by ip route (Table 1) with seven days between them[11] (Fig. 2).

Immunization with Lipopolysaccharide from *Escherichia coli (ecLPS)*. Mice were challenged with 0.1 mg/kg of ecLPS O111: B4 (SIGMA-ALDERICH, L2630-10MG, Lot 025M4040V12140701) diluted in phosphate-buffered saline (PBS). Two ip inoculations were performed (Table 1) with a range of ten days between immunizations[12] (Fig. 2).

Evaluation of the immunization efficacy. Mice were randomly selected and sacrificed for withdrawal of whole blood via cardiac puncture following stimulation of the immune system (twenty days after the last immunization with pfMSP-3 and dT and Influenza vaccines; six days after the last inoculation of Ovalbumin and two days after the last dose of ecLPS), serum samples were preserved at -70 ° C. Total IgG against *pf*MSP-3, Tetanus-Diphtheria toxoids (dT) and Influenza vaccine; the Th1, Th2 and Th17 cytokine profile; lymphocyte subpopulations spleen; and intradermal test for the response against Ovalbumin were evaluated.

The antibody response against *pf*MSP-3 recombinant and Influenza vaccine were determinate by conventional Enzyme-Liked immunosorbent Assay – ELISA[9-10], and

627 the antibody response against dT vaccine was determined by Toxin Binding Inhibition
628 – ToBI[13].
629
Evaluation of the immune response triggered by immunization. Cytokine
production triggered by the Pool, T1 and T2 stimuli was evaluated in the serum of the
different mice groups. Cytokine profile was performed by Cytometric Bead Array
Th1/Th2/Th17 (CBA - BD Biosciences (cat# 560484). The data were collected and
analyzed using FacsCANTO flow cytometer (BD Biosciences) and FACScomp (BD),
respectively. Spleens from the animals of all groups were removed and homogenized
with a syringe plunger above 70-μm strainer, and red blood cells were eliminated by
using ACK lysing buffer (Sigma). The cell suspension was counted and
immunollabeled. For staining, cells were incubated with anti-Fc III/II (CD16/32)
receptor Ab (2.4G2), followed by surface staining with various combination of the
following antibodies for 30 min at 4°C in the dark: anti-CD8 (PE-Cy7, 53-6.7 clone, BD
Pharmingen), anti-CD3 (PerCPCy 5.5, 145-2c11 clone, BD Pharmingen), anti-CD4
(APC-H7, GK1.5 clone, BD Pharmingen), anti-B220 (APC, RA3-6B2 clone, BD
Pharmingen), anti-CD62L (BB515, MEL14 clone, BD Horizon), APC anti-CD44 (APC,
IM7 clone, BD Pharmingen) and Foxp3 (Alexa Flour, 647, R16715, BD Pharmingen).
Acquisition and data analyze by flow cytometry using BD FACSCanto[TM] and FlowJo
®.
647
Behavioral analysis. Mice behavioral performance was assessed according to the
experimental schedule presented in Fig. 2. To note, mice was submitted to different
behavioral paradigms to evaluate exploratory and locomotor activity, cognitive abilities,
and parameters involved in anxiety-like behavioral at days twenty one, seven and three
days after *pf*MSP-3, Tetanus-Diphtheria and Influenza vaccine; Ovalbumin; LPS
inoculation, respectively, and seventy seven days after CQ treatment. The same co-
hort of mice were evaluated on OFT, NORT, Y-maze and L/D, respectively (Fig. 2). All
experiments were performed with incandescent light source of 200 lux of intensity in
the evening period. Animals were acclimatized in the experimental room at least for 2
hours before the experimental sessions. Behavioral performance of mice was recorded
by AnyMaze® software (Stoelting Co., Wood Dale, IL, USA) that captures the videos
and interpret exclusively the mice open field performance and only the locomotor
activity in the novel object recognition training and test sessions. A trained blind-to-
treatment researcher evaluate others behavioral parameters of each mouse
individually by video analysis. In all behavioral tests mice were individually placed on
the apparatus and monitored by a video camera (positioned above and at 90 ° of the
equipment) connected to a monitor to obtain images and data analyze. The apparatus
were properly cleaned with 70% alcohol and dried after each test.
666
Open Field Task (OFT). To address the effect of immune responses on locomotion
and on long-term habituation, mice were submitted to the open field task (OFT) with a
training and a test session 24 hours apart, as described elsewhere[4]. In each OFT
session [training (OFT1) and test, (OFT2)] mice was individually submitted to freely
explore a gray acrylic square box in (50 × 50 × 50 cm, length × width × height) for 10
minutes. In the training session it was evaluated the short-term habituation to novelty
by decreasing the distance travelled during the 1st to 3rd minute, since mice typically
exhibit less exploratory behavior during the first few minutes of testing in a familiar
open field arena; the locomotor activity (the distance travelled during the last 6 minutes

of the test); and anxiety-like behavior by the time and distance travelled in the center
zone. In the test session (OFT2), we evaluate the first 3rd and the last 7th minutes total
distance travelled. Finally, it was analyzed the long-term habituation to novelty,
characterized by a significant decrease in the total distance travelled comparing the
first 3 minutes of the OFT test (OFT2) with the OFT training (OFT1) sessions.
681
Novel Object Recognition Task (NORT).
To evaluate the recognition memory, the NORT was carried out in the same OFT
apparatus, 24 hours after the OFT2 session[4]. NORT starts with a training session, in
which mice are exposed to two familiar objects: FO1 and FO2. On the training session,
a similar exploratory activity (of approximately 20 seconds) of mice for both familiar
objects familiar, is expected[15]. The test session is performed 24 hours later. Mice are
exposed to a New Object (NO) and to one of the previously exposed Familiar Objects
(FO1 or FO2). NORT is based on the tendency of mice to differentiate a novel object
(NO) from a familiar object (FO), thus exploring more the NO in detriment of the FO[4,15].
691
Animals were placed in the periphery of the apparatus with the FO and allowed to
explore them 10 minutes. To the NORT test session, 24 h after the training session,
the animals were placed back in the arena with one of the FO and the NO to measure
the long-term recognition memory. The time spent exploring the objects were again
evaluated for 10 min by a trained blind-to-treatment observer, and an exploration was
only considered when the animal's nose or mouth was in contact with the objects.
During both the training and the test phase, objects are located in opposite and
symmetrical corners of the arena. In the training session both objects are novelty and
time explored on both should be similar. Differently, a successful test session is
indicated by a longer time spent on NO. The time exploration was expressed as
percentage of time exploring each object. Animals that recognized the NO as such
explored it for more than 50% of the total time.
704
Light/Dark Task. The light / dark task was performed as described by Almeida and
collaborators[16] with minor modification to measure anxiety behavior-like phenotype[17].
The apparatus consist in a rectangular box (50 x 30 x 30 cm, height × length × width)
in acrylic with two different sides (colored white and black), separated by a wall (5x5cm)
with an opening at the level of the base of the apparatus joined both sides of the
apparatus. A white 100W lamp, placed 60cm above the center of the apparatus,
illuminated the white side of the apparatus, while the black side was kept closed without
illumination. The mice were individually placed in the light compartment for free
exploration of the apparatus for 5 minutes. The following behavioral parameters were
analyzed: the time spent in the light compartment and the number of transitions
between the compartments (light and dark).
716
Statistical analysis. All statistical analyses were performed using a statistical software
package (Prism 5.0, GraphPad). The data were extracted from the AnyMaze®
software. To analyze OFT and Light / Dark task were used absolute data. The time in
each object in NORT was transformed into a percentage, from which the delta was
extracted based on the subtraction: OF1 - OF2 (training session) and NO - FO (test
session). The two-way ANOVA with Bonferroni correct were used to analyze OFT. The
Student t-test with Mann-Whitney correction were used for to analyze the groups in

NORT, light-dark task and immune response. Data are presented as mean ± standard error. P <0.05 was considered statistically significant.

References

1. Franke-Fayard B, Trueman H, Ramesar J, Mendoza J, Van der Keur M, Van der Linden R, et al. A Plasmodium berghei reference line that constitutively expresses GFP at a high level throughout the complete life cycle. Mol Biochem Parasitol. 2004;137:23–33.
2. Potter S, Chan-Ling T, Ball HJ, Mansour H, Mitchell A, Maluish L, et al. Perforin mediated apoptosis of cerebral microvascular endotelial cells during experimental cerebral malaria. Int J Parasitol. 2006;36:485–96.
3. Reis PA, Comim CM, Hermani F, Silva B, Barichello T, Portella AC, et al. Cognitive dysfunction is sustained after rescue therapy in experimental cerebral malaria, and is reduced by additive antioxidant therapy. PloS Pathog. 2010;6:e1000963.
4. de Sousa LP, Almeida RF, Ribeiro-Gomes FL, Carvalho LJM, Souza TM, Souza DOG & Daniel-Ribeiro CT. Long-term effect of uncomplicated Plasmodium berghei ANKA malaria on memory and anxiety-like behaviour in C57BL/6 mice. Parasites & Vectors. 2018; 11:191.
5. Lavigne MV, Castro M, Mateo N, Deluchi S, Atzori C. Whole-cell *Bordetella pertussis* vaccine component modulates the mouse immune response to an unrelated soluble antigen. Microb. and Infect. 2002; 815-820.
6. Castilho SF, Chovel ML, Hernández NG, González LC, Blanco A, Hernández DS, Medina MF, Tito MA, Quiñoy JLP. A Bordetella pertussis proteoliposome induces protection in mice without affecting the immunogenicity of diphtheria and tetanus toxoids in a trivalent formulation. Clinic. and Experiment. Vacc.Reser. 2016; 5:175-178.
7. Goodier MR, Rodriguez-Galan A, Lusa C, Nielsen CM, Darboe A, Moldoveanu AL, White MJ, Behrens R, Riley EM. Influenza Vaccination Generates Cytokine-Induced Memory-like NK Cells: Impact of Human Cytomegalovirus Infection. J. Immunol. 2016; 197(1):313-25.
8. Zhu W, Li S, Wang C, Yu G, Prausnitz MR, Wang BZ. Enhanced immune responses conferring cross-Protection by skin Vaccination With a Tri-component influenza Vaccine Using a Microneedle Patch. Front. in Immunol. 2018; 9:1705.
9. Oeuvray C, Bouharoun-Tayoun H, Grass-Masse H, Lepers JP, Ralamboranto L, Tartar A, Druilhe P. A novel merozoite surface antigen of Plasmodium falciparum (MSP-3) identified by cellular-antibody cooperative mechanism antigenicity and biological activity of antibodies. Mem. Inst. Oswaldo Cruz. 1994; 89 Suppl 2:77-80.
10. Daher LJ1, Demanga CG, Prieur E, Pérignon JL, Bouharoun-Tayoun H, Druilhe P. Toward the rational design of a malaria vaccine construct using the MSP3 family as an example: contribution of immunogenicity studies in models. Infect. Immun. 2010; 78(1):477-85.
11. Lima MCR, Prouvost-Danon A, Patricia MR, Chagas MS, Calheiros AS, Cordeiro RSB. Studies on the mechanisms involved in antigen-evoked pleural. J of Leuk Biol. 1997, 61.

12. Couch Y, Trofimov A, Markova3 N, Nikolenko V, Steinbusch HW, Chekhonin V, Schroeter C, Lesch KP, Anthony DC, Strekalova T. Low-dose lipopolysaccharide (LPS) inhibits aggressive and augments depressive behaviours in a chronic mild stress model in mice. J. of Neuroinflamm. 2016; 13:108.

13. Matos DCS, Marcovistz R, Cabello PH, Georgini AR, Sakauchi D, Silva LL. Immunogenicity Test of Tetanus Component in Adsorbed Vaccines by Toxin Binding Inhibition Test. Mem Inst Oswaldo Cruz. 2002, 97(6): 909-913.

14. Almeida RF, Ganzella M, Machado DG, Loureiro SO, Leffa D, Quiconzes Santos A, et al. Olfactory bulbectomy in mice triggers transient and longlasting behavioural impairments and biochemical hippocampal disturbances. Prog Neuropsychopharmacol Biol Psychiatry. 2017;76:1-11.

15. Leger M, Quiedeville A, Bouet V, Haelewyn B, Boulouard M, Schumann-Bard P, Freret T. Object recognition test in mice. Nat Protoc. 2013; 8(12):2531-7.

16. Almeida RF, Comasseto DD, Ramos DB, Hansel G, Zimmer ER, Loureiro SO, et al. Guanosine anxiolytic-like effect involves adenosinergic and glutamatergic neurotransmitter systems. Mol Neurobiol. 2016;54:423-36.

17. Griebel G, Holmes A. 50 years of hurdles and hope in anxiolytic drug discovery. Nat Rev Drug Discov. 2013; 12(9):667-87.

LEGENDS FOR FIGURES AND TABLE

Fig. 1. **Behavioral analysis of healthy and *Pb*A-infected mice**. In the New Object Recognition Task (NORT), mice explored similarly the familiar objects in the training session: **a)** in healthy mice and **b)** in infected mice. In the test session, **c)** control (non-infected non-immunized mice) group showed increased new object exploration, as expected. Mice immunized with the Pool and T2 Ag showed increased new object exploration as compared to the Control and the T1 groups and **d)** infected mice showed decreased new object exploration in relation to the mice of the Control and Infected-Immunized groups. In the OFT1, **e)** all groups of healthy immunized mice walked less in the center of the arena when compared to the control group mice and **f),** the infected mice showed a decrease in the distance travelled in the center of the OFT1, as compared to the non-infected mice group. No difference was observed in the light / dark task between **g)** healthy mice groups. In contrast, **h**) *Pb*A-infected mice groups presented anxiety-like behavioral which was attenuated after immunization with the Pool and T2 immune strategies. Healthy mice groups (Control, n = 6; Pool, n = 8; T1, n = 8 and Control, n = 10; Pool, n = 10; T2, n = 10) and infected mice groups (Control, n = 6; Inf, n = 8; Inf-Pool, n = 6; Inf-T1, n = 6 and Control, n = 25; Inf, n = 17; Inf-Pool, n = 20; Inf-T2, n = 18). Data are mean and s.e.m. ***$P < 0.001$; **$P < 0.01$; *$P < 0.05$; Unpaired t-test without and with Mann-Whitney test was used. For the NORT and OFT analysis, data are representative of five (Control, Pool and Inf-Pool groups), one (T1, T2 and Inf-T1 groups) and two (Inf-T2 group) independent experiments; and for the light-dark analysis, data are representative of four (Control, Pool and Inf-Pool groups), one (T1, T2 and Inf-T1 groups) and two (Inf-T2 group) independent experiments.

Extended data, Fig. 1. **Behavioral analysis of healthy and infected mice**. In the open field task (OFT), all groups showed a reduction in the distance travelled in the test session (OF2) when compared to the training session (OF1), **a)** healthy mice groups (Control, n = 6; Pool, n = 8; T1, n = 8 and Control, n = 10; Pool, n = 10; T2) and **b)** infected mice groups (Control, n = 6; Inf, n = 8; Inf-Pool, n = 6; Inf-T1, n = 6 and Control, n = 25; Inf, n = 17; Inf-Pool, n = 20; Inf-T2, n = 18). The (Pool and T1) immunized mice groups **a)** walked less on OFT1 when compared to non-immunized mice and the T2 immunized group showed a tendency (P = 0.0524) towards a decrease in the distance travelled. The same was observed in the infected and Pool and T2 immunized mice groups, **b)** but only when compared to non-infected / non-immunized mice group. Data are representative of five (Pool and Inf-Pool groups), one (T1, T2 and Inf-T1 groups) and two (Inf-T2 group) independent experiments. In the new object recognition task (NORT), all groups of mice explored similarly the familiar objects in the training session: **c)** healthy mice groups and **d)** Infected mice group. **e)** The control group (non-infected non-immunized mice) showed and increased new object exploration, as expected, and the Pool group showed a still increased new object exploration as compared to the Control group. The infected mice **f)** showed a decreased exploration of the new object, when compared to the control group mice; differently from the infected and Pool Immunized mice. Healthy mice group (Control, n = 22 ; Pool, n = 21) and infected mice group (Control, n = 22; Inf, n = 20; Inf-Pool, n = 21). Data are representative of two independent experiments for all groups. Data are mean and s.e.m. ***P < 0.001; **P < 0.01; *P < 0.05; Two-way ANOVA (a, b) and Unpaired t-test (c, d, e, f) was used.

Extended data, Fig. 2. **Spleen weight and size, and total number of spleen cells**. Immunized mice showed increases in the **a)** spleen weight and **b)** total number of splenocytes. **c)** Representative photograph of all studied groups of mice. Groups of infected mice showed a dark color attributed to hemozoin, even two and a half months after infection. All groups of immunized mice showed increased spleen sizes. Samples were collected after the behavioral evaluation. Five mice per group were analyzed. Data are mean and s.e.m. **P < 0.01; *P < 0.05; Unpaired t-test with Mann-Whitney test was used.

Extended data, Fig. 3. **Immune response against dT and influenza vaccines; pfMSP-3 and OVA**. Specific immune responses were triggered by : **a)** dT (n = 7 mice per group); **b)** Influenza vaccine (Control, n = 6; Pool, n = 6; T1, n = 3; Inf, n = 6; Inf-Pool, n = 6; Inf-T1, n = 3); **c)** *pf*MSP-3 (n = 5 mice per group) and **e)** OVA (Control, Pool and T2, n = 5 mice per group; Inf, Inf-Pool and Inf-T2, n = 4 mice per group), confirming the effectiveness of the immune stimuli. Mice were randomly chosen from the experiments, according to the type 1 and 2 immunization strategies, for the analysis of the immune responses. Samples were collected before the behavioral evaluation for analysis of responses against dT and influenza vaccines and were collected after behavior tests for the study of other parameters. Data are mean and s.e.m. **P < 0.01; *P < 0.05; Unpaired t-test with Mann-Whitney test was used.

Extended data, Fig. 4. **Flow cytometry analysis of B and T cell subpopulations in the spleen.** Healthy or infected and treated mice were immunized, or not, with Pool or T2 strategy and the populations of interest analyzed at the end of the cognitive behavioral tasks. **a)** representative gating strategy to identify the populations of B cells (B220+), CD4 T cells (CD3+CD4+), CD8 T cells (CD3+CD8+) and T regulatory cells (CD3+CD4+CD25+Foxp3+); frequency of **b)** B cells; **c)** CD4 T cells and **d)** CD8 T cells per spleen and e**)** frequency of Treg cells among the CD4 T cells. Five mice were randomly chosen per group, after behavior tests, for the analysis. Data are mean and s.e.m. **$P < 0.01$; *$P < 0.05$; Unpaired t-test with Mann-Whitney test was used.

Extended data, Fig. 5. **Analysis of different subsets of splenic CD4 and CD8 T cells.** Healthy or infected and treated mice were immunized, or not, with Pool or T2 strategy and populations of interest analyzed at the end of the cognitive behavioral tasks: **a)** representative gating strategy to identify the Naive (gate: 1; $CD44^- CD64L^+$); Effector/Effector Memory (gate: 2; $CD44^+CD64L^-$) and Central Memory (gate: 3; $CD44^+CD64L^-$) subpopulations of CD4 and CD8 T cells; frequency of **b)** Naïve; **c)** Effector/Effector Memory and **d)** Central Memory subsets of CD4 T cells; frequency of **e)** Naïve; **f)** Effector/Effector Memory and **g)** Central Memory subsets of CD8 T cells. Five mice were randomly chosen per group, after behavior tests, for the analysis. Data are mean and s.e.m. **$P < 0.01$; *$P < 0.05$; Unpaired t-test with Mann-Whitney test was used.

Extended data, Fig. 6. **Serum cytokine profile analysis**. Mice immunized with Pool and T2 strategies presented increased serum levels of TNFα, IFNγ and IL-4 cytokine (**a, b, d**). IL-6 levels were only increased in **c)** infected-Pool and T1 immunized mice groups (Control, n = 5; Pool, n = 5; T1, n = 3; Inf, n = 5; Inf-Pool, n = 5; Inf-T1, n = 3 and Control, n = 3; Pool, n = 4; T2, n = 5; Inf, n = 5; Inf-Pool, n = 5; Inf-T2, n = 5). Data are representative of three (Control, Pool, Inf and Inf-Pool groups) and one (T1, T2, Inf-T1 and Inf-T2 groups) independent experiments. Samples were collected before (Control, Pool, Inf-Pool and Inf-T1) and after (Control, Pool, T2, Inf, Inf-Pool and Inf-T2) the behavioral evaluation. Data are mean and s.e.m. **$P < 0.01$; *$P < 0.05$; Unpaired t-test with Mann-Whitney test was used.

Fig. 1 Material & Methods. **Experimental draw**. Groups of mice, infected or not with *Plasmodium berghei* ANKA (*Pb*A), were treated with chloroquine (25 mg / kg) for seven days via gavage. After 14 days, they were subdivided into groups of immunized (B and D) or non-immunized (A and D) mice, according to the immunization strategies P (Pool), T1 (type 1 stimuli) and T2 (stimuli of type 2). Subsequently, mice were evaluated in behavioral tasks for locomotivity, memory and anxiety phenotype.

Fig. 2 Material & Methods. **Flow chart of immune stimuli and behavioral assessment**. After infection and treatment with CQ, mice were categorized as immunized or non-immunized, according to the immunization (P, T1 and T2)

strategies used. The dT and Influenza vaccines and pfMSP-3 were inoculated at the same day, with three doses in different immunization pathways, with a 20-day interval between inoculations. OVA was inoculated after the third dose of dT and Influenza vaccines and pfMSP-3 protein was administered, three doses with a seven-day interval between the inoculations. The ecLPS was inoculated after the second dose of OVA, being two doses with an interval of ten days between them. Assessment of performance on behavioral tasks started 77 days after chloroquine treatment; 22 days after last inoculation of pfMSP-3 and dT and influenza vaccines; 7 days after last inoculation of Ovalbumin and 3 days after last inoculation with ecLPS. The open field was performed to measure locomotivity, habituation memory and anxiety phenotype, being performed in two sessions [training (OF1) and test (OF2)]. Then, the new object recognition task was performed to measure long-term memory, also in two sessions at consecutive days (training and testing). Finally, the anxious behavior phenotype was specifically evaluated by the light-dark test, performed in an unique session.

Table 1. Immune stimulus inoculation strategy: route, region, concentration, volume and number of injections of immunogens.

Figure 1

Extended data, Figure 1

Extended data, Figure 2

Extended data, Figure 3

Extended data, Figure 4

Extended data, Figure 5

Extended data, Figure 6

Material & Methods, Figure 1

Material & Methods, Figure 2

Chloroquine   Resting

d0   d4   d10   d11   d23   d24   d85   d88   d92

P. berghei ANKA infection

Immune stimuli

Behavioral Evaluation

dT/ Infl. /*pf*MSP-3   dT/ Infl. /*pf*MSP-3   OVA   OVA

d24   d45   d66   d67   d74   d76   d81   d85

dT/ Infl. /*pf*MSP-3   OVA   *ec*LPS   *ec*LPS

Open Field   Light/Dark

d88   d89   d90   d91   d92

Object Recognition

Open Field

Training   Test

d1   d2

Object Recognition

Training   Test

FO-1   FO   NO
FO-2

d3   d4

Light/Dark

d5

Material & Methods, Table I

Immune Stimuli	Route	Region	Concentration	Volume	Inoculation
dT vaccine	Subcutaneous	Back	1/5 human dose	100 µl	3
Influenza vaccine	Intramuscular	Quadriceps	1/5 human dose	100 µl	3
pfMSP-3	Subcutaneous	Base Tail	10 µg/mice	100 µl	3
ecLPS	Intraperitoneal	Abdomen	0,1 mg/kg	100 µl	2
Ovalbumin	Subcutaneous Intraperitoneal	Back and Abdomen	50 µg	200 µl	1 s.b.c. 2 i.p.