Reward produces learning of a consciously inaccessible feature

Xue Dong[1,2], Mingxia Zhang[1,2], Bo Dong[1,3], Yi Jiang[2,4,5], Min Bao[1,2,4,*]

[1] CAS Key Laboratory of Behavioral Science, Institute of Psychology, Chinese Academy of Sciences, Beijing 100101, China

[2] Department of Psychology, University of Chinese Academy of Sciences, Beijing 100049, China

[3] Department of Psychology, Suzhou University of Science and Technology, Suzhou 215009, China

[4] State Key Laboratory of Brain and Cognitive Science, Beijing 100101, China

[5] CAS Center for Excellence in Brain Science and Intelligence Technology, Shanghai 200031, China

* **Corresponding author:** Min Bao

Email: baom@psych.ac.cn

ORCID ID: 0000-0002-5347-9663

Keywords

   reward, attention, eye-of-origin, awareness, inter-ocular suppression.

Abstract

Reward has significant impacts on behavior and perception. Most past work in associative

reward learning has used distinct visual cues to associate with different reward values. Thus, it

remains unknown to what extent the learned associations depend on the consciousness. Here we

resolved this issue by using an inter-ocular suppression paradigm with the monetary rewarding and

non-rewarding cues identical to each other except for the eye-of-origin. Thus, the reward coding

system cannot rely on the consciousness to select the reward-associated cue. Surprisingly, the

targets in the rewarded eye broke into awareness faster than those in the non-rewarded eye. We

further revealed that producing this effect required both attention and inter-ocular suppression.

These findings suggest that the human's reward coding system can produce two different types of

reward-based learning. One is independent of the consciousness yet fairly consuming attentional

resource. The other one results from volitional selections guided by top-down attention.

Introduction

It has long been recognized that reward and punishment strongly modulate behaviors and

perception (Proshansky & Murphy, 1942). In addition, recent studies have brought evidence of

reward processing abnormalities in patients with mental disorders (Whitton, Treadway, & Pizzagalli,

2015). Therefore, reward is becoming an increasingly important topic in both cognitive

neuroscience and clinics.

The effects of associative reward learning in many studies (Anderson, 2017; Della Libera &

Chelazzi, 2006; Marx & Einhauser, 2015; Proshansky & Murphy, 1942; Raymond & O'Brien, 2009;

Serences, 2008; Wilbertz, van Slooten, & Sterzer, 2014) can be epitomized by a phenomenon

known as the law of effect (Thorndike, 1911). The law denotes that a response repetitively followed

by a reward will be more readily to recur when the participants encounter the same stimuli and

context. In other words, actions or perceptions are biased in favor of reward-associated stimuli. In

most cases, participants can consciously realize the rule of reward and thus intentionally direct

attentional resources to the related stimuli. Therefore, reward appears to always work with selective

attention. Indeed, numerous studies have suggested that attention and reward coding are closely

linked processes for stimulus selection. For example, attentional inhibition of distractors in a prime

display is robust for highly (but not poorly) rewarded prime display in a negative priming study (Della

Libera & Chelazzi, 2006). Furthermore, facial stimuli associated with learned reward value can

survive from attentional blink, suggesting privileged processing of reward-associated stimuli

(Raymond & O'Brien, 2009). Interestingly, later studies report that learned reward value can bias

attention independent of strategy and volition (Anderson, Laurent, & Yantis, 2011; Hickey, Chelazzi,

& Theeuwes, 2010). In addition, reward and attention also similarly bias visual perception in
binocular rivalry tasks (Marx & Einhauser, 2015; Wilbertz et al., 2014).

Nevertheless, it is already known that neuromodulatory signals for rewards (and punishments)
are released diffusively throughout the entire brain (Dalley et al., 2001; Schultz, 2000; Vickery,
Chun, & Lee, 2011). Intuitively, the effects of rewards could be free from the constraint of attention
and consciousness. Thus, a lingering possibility to be tested is that reward learning can be
established independent of the consciousness and selective attention. Unfortunately, to our
knowledge, this hypothesis cannot be easily proved by the previous work, because in most of them
perceptually distinguishable visual cues are used to associate with different reward values.
Therefore, the structure of experiments *per se* may be responsible for their observations that
rewards teach or rely on selective attention to choose objects of behavioral significance among a
set of candidates. The present study introduces a novel paradigm in which the participants cannot
consciously differentiate the monetary rewarding and non-rewarding visual cues. This was realized
by rendering the two visual cues identical to each other except for their eye-of-origin information.
With this design, participants should not be able to consciously tell the difference between the two
monocular cues (Wolfe & Franzel, 1988; Zhang, Jiang, & He, 2012). Therefore, they could not
realize which cue was associated with a reward. Selective attention is thus not believed to produce
any eye-specific learning effects in such a paradigm. Any eye-specific learning effects established
over the reward-based training should be contributed from the reward coding system, but
independent of the consciousness and selective attention. We first showed that the eye-specific
reward learning could be observed during inter-ocular suppression using a continuous flash
suppression (CFS) method (Fang & He, 2005; Tsuchiya & Koch, 2005). With five more
experiments, we further examined the role of top-down attention in the learning process and at
what circumstances the eye-specific effects were present or absent.

Results
Our initial experiment was based on a b-CFS paradigm (Jiang, Costello, & He, 2010). Reward was
only associated with the detection of an invisible target in one of the two eyes (Figure 1A-B), which
allowed us to examine whether reward learning can be established independent of selective
attention. Participants completed two reward training sessions; whereas no reward was given in
the pre- and post-test. Since the targets were in the same appearances whether it was presented
to the rewarded eye or non-rewarded eye, selective attention was not likely to modulate the
processing of associative reward learning in an eye-specific way. Any eye-specific learning effect
in this training paradigm should be independent of consciousness. For each session, the

breakthrough ratio was calculated by dividing the number of trials with correct responses by the

total trial count for each eye-of-origin condition, respectively.

Repeated measurements ANOVA and paired t-test were used to statistically analyze the results

(see Table 1 for the detailed statistics of the ANOVA results). In the pre-test, there was no

significant difference between the breakthrough ratios for the rewarded eye and the non-rewarded

eye ($t(35) = 0.193$, $p = 0.848$, Cohen's $d = 0.028$). Therefore, we subtracted the breakthrough ratios

in the pre-test from those in the subsequent sessions to estimate the change of breakthrough ratios

across the sessions. Consistent with the previous finding (Mastropasqua, Tse, & Turatto, 2015),

the targets broke into awareness generally faster in the later sessions than in the pre-test.

Surprisingly, there were more breakthrough trials in the rewarded eye than in the non-rewarded

eye (training 1: $t(35) = 2.815$, $p = 0.008$, $d = 0.463$; training 2: $t(35) = 3.649$, $p < 0.001$, $d = 0.658$,

see Figure 1C). However, this eye specific effect was absent in the post-test ($t(35) = 0.438$, $p =$

0.664, $d = 0.059$) where the participants no longer received monetary rewards.

Figure 1. Stimuli and results of Experiment 1. (A), (B) Trial sequence and four kinds of targets in

Experiment 1a. (C) The improvement of breakthrough ratio relative to pre-test in training and post-

test sessions in Experiment 1a. (D) The stimuli and trial sequence in Experiment 1b. (E) The improvement of breakthrough ratio relative to pre-test in the training and post-test sessions in Experiment 1b. Error bars show ±1 SE. Asterisks indicate the significance level (with **$p < 0.01$, ***$p < 0.001$).

We further examined any potential influences of false alarms. In the present study, a false alarm represents that the participants made a wrong keypress in a trial. This could be due to a wrong judgment of the target position after the target broke into awareness or a cheating response by randomly guessing the location of the invisible target. In the training sessions, the participants might have the motivation to cheat in order to receive more monetary rewards. Considering the four-alternative forced-choice (4AFC) task, the probability of correctly guessed trials was 1/3 of that of wrongly guessed trials. Assuming that all the wrong responses were due to failure guesses, the maximum number of correctly guessed trials in theory could be estimated by dividing the number of trials with wrong responses by three. Therefore, to correct the effect of false alarm, the maximum number of correctly guessed trials was subtracted from the number of trials with correct responses before calculating the breakthrough ratios. After the correction, the breakthrough ratio also showed more increase in the rewarded eye than in the non-rewarded eye (training 1: $t(35) = 2.782$, $p = 0.009$, $d = 0.465$; training 2: $t(35) = 3.562$, $p = 0.001$, $d = 0.653$). The eye specific effect was absent in the post-test ($t(35) = 0.329$, $p = 0.744$, $d = 0.044$). In addition, by comparing the learning effect of the 1st and 2nd block of training 1, we found this eye-specific effect developed very fast in the training.

To test if top-down attention could affect the learning process, another group of participants were asked to perform a RSVP task simultaneously with the b-CFS task. The RSVP stimuli were presented on the center of the screen, and the b-CFS targets were presented 2° above or below the center of the screen. By using the fixation task and moving the b-CFS targets away from the central fixation position, the attention deployed on the b-CFS targets should be greatly reduced as compared to our initial experiment. Participants performed well in the RSVP task in all the sessions (hit rate: 87.23±7.85%, false alarm rate: 1.76±2.49%). The results of the b-CFS task, however, showed no significant difference in the breakthrough ratios between the two eyes in any of the four sessions (pre-test: $t(13) = 0.637$, $p = 0.536$, $d = 0.178$; training 1: $t(13) = 0.764$, $p = 0.458$, $d = 0.145$; training 2: $t(13) = 1.141$, $p = 0.274$, $d = 0.180$; post-test: $t(13) = 1.615$, $p = 0.130$, $d = 0.141$). Similar results were found after the false alarm correction.

Monocular reward learning without inter-ocular suppression. Is inter-ocular suppression necessary for this eye-specific associative reward learning effect? To answer this question, we performed three experiments. We first tested the role of inter-ocular suppression using trials with and without CFS in Experiment 2a. Only in the trials without CFS (i.e. target-only trials) were

participants rewarded for a correct response to the target presented to the rewarded eye. It was
clear that these target-only trials were irrelevant to inter-ocular suppression. We found that in these
trials, participants performed well in the pre-test for both eyes, with no significant difference in the
performance ($t(14) = 0.913$, $p = 0.377$, $d = 0.189$). Training improved the performance for both eyes
slightly by between 1.0% and 1.5% in the target-only trials with no difference across the eyes (all
ps > 0.54). In the with-CFS trials, the break through ratio increased with training. However, no
significant difference was observed between the breakthrough ratios for the rewarded eye and
those for the non-rewarded eye in the pre-test ($t(14) = 0.969$, $p = 0.349$, $d = 0.223$), and the increase
of breakthrough ratios did not show any difference between the two eyes in any of the training
sessions (training 1: $t(14) = 1.007$, $p = 0.331$, $d = 0.293$; training 2: $t(14) = 0.472$, $p = 0.645$, $d =$
0.102; training 3: $t(14) = 0.468$, $p = 0.647$, $d = 0.101$, see Figure 2A). Similar results were found
after the false alarm correction.

Since the performance for the target-only trials was almost perfect, a potential influence of ceiling
effect could not be excluded. We then used a more difficult contrast detection task and an
orientation discrimination task to further examine this issue. Subthreshold stimuli rather than inter-
ocular suppression were used to render the stimuli hard to perceive in these two experiments.

In the contrast detection experiment (Experiment 2b, Figure 2B), no significant difference of the
performance between the eyes was observed (Figure 2C, $F(1, 13) = 1.986$, $p = 0.182$, $\eta^2 = 0.133$),
though the contrast threshold showed a significant change across sessions ($F(1.188, 15.446) =$
7.977, $p = 0.010$, $\eta^2 = 0.380$, Greenhouse-Geisser corrected). The main effect of Session was
predominantly due to the increase of thresholds in the post-test than in other sessions, probably
reflecting reduced motivation in the post-test that lacked incentives as compared to the training
sessions.

Considering that the contrasts of stimuli in Experiment 2b were close to or below the detection
threshold, visual signals to primary visual cortex might be faint. As a result, the eye-specific reward
might not be able to enhance these signals. However, the eye-specific reward learning effect was
still absent in Experiment 2c where high contrast gratings and orientation discrimination task were
used (Figure 2D). For each offset level, the performances in the pre-test were not different between
the two eyes (all ps > 0.80, FDR corrected). After subtracting the correction rates of the pre-test

from those in training sessions, no difference was found between the rewarded and non-rewarded
eyes on any levels in all the training sessions (Figure 2E, all *p*s > 0.19, FDR corrected).

Figure 2. Stimuli and results of Experiment 2. (A) The improvement of breakthrough ratio relative
to pre-test in three training sessions of Experiment 2a. (B) Trial sequence of the contrast detection
task in Experiment 2b. (C) Contrast detection threshold of four testing sessions of Experiment 2b.
(D) Trial sequence of the orientation discrimination task in Experiment 2c. (E) Orientation
discrimination threshold of four testing sessions of Experiment 2c.

When selective attention was involved in the eye-specific rewarding. The results of the
Experiments 1 and 2 indicated that inter-ocular suppression was necessary for eliciting the eye-
specific reward learning effects when participants could not discriminate the rewarding vs. non-
rewarding targets. In Experiment 1a, the eye-of-origin information was the only difference between
the two kinds of targets, though could not be consciously realized. An interesting question is
whether we can still observe the eye-specific learning effects when reward is also related to another
feature that could be consciously distinguished. In Experiment 3, the rewarding target was defined
by a conjunction of two features. Only the bar in one of the two orientations presented to the
rewarded eye was the rewarding target. Although the participants were still not aware of the
manipulation on the eye of origin information, they could quickly learn of the relationship between
orientation and reward over training. We conducted Experiment 3 on two different groups of
participants. Because of a mistake, the first group of participants did not complete the post-test
after training (this was referred to as Experiment 3a hereafter). Therefore, we replicated the
experiment in another group of participants but added the post-test (this was referred to as
Experiment 3b hereafter). No significant differences among conditions were found in the pre-test

of both groups (all ps > 0.18). Surprisingly, we found two different patterns of the learning effect in

the two groups. Results from the first group of participants showed a significant interaction between

Orientation and Session (see Table 1 for detailed statistics). Paired t-test indicated that the reward-

based learning was orientation-specific (Figure 3B, RO-RE vs. NO-RE: training 1: $t(17)$ = 1.336, p

= 0.199, d = 0.176; training 2: $t(17)$ = 2.687, p = 0.016, d = 0.621; training 3: $t(17)$ = 4.205, p <

0.001, d = 0.861. RO-NE vs. NO-NE: training 1: $t(17)$ = 1.559, p = 0.138, d = 0.164; training 2: $t(17)$

= 5.092, p < 0.001, d = 0.457; training 3: $t(17)$ = 6.864, p < 0.001, d = 0.785) but not eye-specific

(ps > 0.20 for RO-RE vs. RO-NE and NO-RE vs. NO-NE). By contrast, results from the second

group of participants showed a significant interaction between Eye-of-origin and Session. Further

analysis revealed that the eye-specific learning effect became obvious in the last training session

and even observable in the post-test where reward has been withdrawn (Figure 3C, RO-RE vs.

RO-NE: training 1: $t(15)$ = 1.307, p = 0.211, d = 0.314; training 2: $t(15)$ = 0.961, p = 0.352, d =

0.302; training 3: $t(15)$ = 2.359, p = 0.032, d = 0.827; post-test: $t(15)$ = 2.587, p = 0.021, d = 0.691.

NO-RE vs. NO-NE: training 1: $t(15)$ = 0.307, p = 0.763, d = 0.077; training 2: $t(15)$ = 0.934, p =

0.365, d = 0.223; training 3: $t(15)$ = 1.343, p = 0.199, d = 0.421; post-test: $t(15)$ = 2.013, p = 0.062,

d = 0.599). Repeated measurements ANOVA disclosed a non-significant trend of interaction

between Orientation and Session. An orientation-specific effect was only observed in the last

training session in the rewarded eye, and was absent in the post-test (RO-RE vs. NO-RE: training

1: $t(15)$ = 1.504, p = 0.153, d = 0.352; training 2: $t(15)$ = 1.312, p = 0.209, d = 0.261; training 3:

$t(15)$ = 3.330, p = 0.005, d = 0.750; post-test: $t(15)$ = 1.833, p = 0.087, d = 0.332; RO-NE vs. NO-

NE: all ps > 0.16). Despite of two different learning patterns, the learning effects of both groups

developed more slowly as compared to that in Experiment 1a. No significant learning effect was

observed in the first training session. Similar results were found after false alarm correction.

Figure 3. Stimuli and results of Experiment 3. (A) The examples of four kinds of trials in each

session: rewarded orientation in the rewarded eye (RO-RE), non-rewarded orientation in the

rewarded eye (NO-RE), rewarded orientation in the non-rewarded eye (RO-NE), non-rewarded

orientation in the non-rewarded eye (NO-NE). The rewarded orientation and rewarded eye were

counter-balanced across participants. (B)-(C) The improvement of breakthrough ratio relative to

pre-test in training and post-test sessions of Experiment 3a and Experiment 3b. Asterisks indicate

the significance level (with *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$).

Discussion

When a monocular pattern is presented to one eye, people have no explicit knowledge of the

pattern's eye of origin (Wolfe & Franzel, 1988; Zhang et al., 2012). We took advantage of this

interesting nature of human vision to imbue the targets' eye-of-origin information with monetary

reward value. Because all the other features (e.g. shape, color, locations, etc.) of the targets were

kept identical between the rewarding and non-rewarding conditions, the participants were not

aware of the existence of two different types of targets. In Experiments 1 and 2, participants

perceived a single kind of target accompanied with reward randomly in part of the trials in which

they made a correct response. The importance of this design is that it dissociates consciousness

and selective attention from reinforcement, thus allowing a close examination of the relationship

between consciousness and reward coding.

The finding of an eye-specific learning effect in Experiment 1a supports our hypothesis that

the reward coding system can rely merely on the eye-of-origin information of the reward-associated

stimuli to produce a non-conscious reward-based learning effect. This result differs from Seitz and

colleagues' findings in their perceptual learning study by several aspects (Seitz, Kim, & Watanabe,

2009; Xue, Zhou, & Li, 2015). First, Seitz et al. used water as rewards for participants who were

deprived of food and water (Seitz et al., 2009); whereas here the effects of monetary reward were

investigated. Second, besides different eye-of-origin, the rewarding and non-rewarding control

stimuli had different orientations in their experiment (Seitz et al., 2009); whereas in our Experiment

1a the only feature difference between the two stimuli was eye-of-origin. Third, the learning effect

Seitz et al. found in their CFS experiment developed over 20 days of training, and could be

observed in a separate sensitivity test where no reward was given (Seitz et al., 2009); whereas the

effect in our Experiment 1a established quickly during the training and vanished immediately in a

post-test that was conducted shortly after the training and with no reward. These distinct

characteristics suggest that Seitz et al.'s findings, as they also proposed (Seitz et al., 2009), should

be considered certain types of perceptual learning effects formed via a classical conditioning

procedure. By contrast, the present findings remind us the brevity of non-conscious fear

conditioning (Raio, Carmel, Carrasco, & Phelps, 2012). To our knowledge, the present study for

the first time suggests that monetary reward, a positive and pleasant rather than a negative and

threatening event (e.g. fear), can also induce non-conscious conditioning and the stimulus-reward

pairing can be established only based on eye-of-origin—a consciously inaccessible feature.

In Raio et al.'s non-conscious fear conditioning study, the non-conscious fear learning was

only significant during the early acquisition and declined quickly in the second half of trials, yet the

conscious fear learning emerged slowly and became stronger during late acquisition (Raio et al.,

2012). Similarly, we found that the reward learning could grow over time in Experiment 3 where the
participants were aware that one orientation was more likely associated with rewards.
Nevertheless, unlike Raio et al.'s finding, the non-conscious reward learning in our Experiment 1a
did not show a rapid forgetting during the training. This might be due to different attentional status
of participants in the training of the two studies. Raio et al. used a CFS paradigm in which
participants did not have a task related to the suppressed stimuli (Raio et al., 2012); whereas we
used a b-CFS paradigm so that participants had to pay attention and report any potential
breakthroughs of the suppressed stimuli. Not only that, attention even seems to play an important
role in generating the eye-specific reward-based learning effect, because the effect was absent
when attention was distracted from the rewarding task. We speculate that top-down attention may
in some way help discern the eye-of-origin information associated with rewards or even likely
participates in reinforcing this association. However, the contribution of top-down attention here
should be clearly different from the more common roles of selective attention in modulating the
actions or perceptions when participants have an explicit knowledge of the rule of reward delivery
in many previous studies (Della Libera & Chelazzi, 2006; Marx & Einhauser, 2015; Raymond &
O'Brien, 2009; Serences, 2008; Wilbertz et al., 2014).

In the past decade, reward has frequently been reported to induce changes in early sensory
regions (Arsenault, Nelissen, Jarraya, & Vanduffel, 2013; Chubykin, Roach, Bear, & Shuler, 2013;
Liu, Coleman, Davoudi, Zhang, & Hussain Shuler, 2015; Persichetti, Aguirre, & Thompson-Schill,
2015; Pleger, Blankenburg, Ruff, Driver, & Dolan, 2008; Serences, 2008; Shuler & Bear, 2006; Zold
& Hussain Shuler, 2015) but see (Rossi et al., 2017). The current work provides a strong case
arguing that reward can induce plasticity in human V1 which is largely independent of
consciousness. Such relatively lower level plasticity should be driven by the co-work of top-down
eye-based attention (Zhang et al., 2012) and diffusively distributed neuromodulatory signals
released by the reward coding system (Dalley et al., 2001; Schultz, 2000; Vickery et al., 2011).
During inter-ocular suppression, the invisible rewarding target activates monocular neurons for the
rewarded eye, while the activities of binocular neurons are mainly dominated by the signals for the
CFS stimuli. Obviously, only the firings of monocular neurons for the rewarded eye are highly
predictive of later rewards, rather than those of binocular neurons (and those of monocular neurons
for the non-rewarded eye). During the training, the reward coding system may soon detect the
reliable association between rewards and responses of monocular neurons for the rewarded eye.
Eye-based attention may then increase the gains particularly for those neurons. Previous work has
revealed that attention and consciousness can be independent to each other in some cases
(Brascamp, Van Boxtel, Knapen, & Blake, 2010; Van Boxtel, Tsuchiya, & Koch, 2010). Therefore,
it is not strange that in the present study attention can work with the reward coding system
independent of consciousness. As a result, the breakthrough was facilitated more for the rewarded
eye than for the non-rewarded eye in Experiment 1a. Once the attentional resource was consumed

by another demanding task, the co-work of attention and reward coding system failed, thus no eye-
specific learning effect was observed in Experiment 1b.

Notably, the above explanation receives further support from the results of our Experiment 2
where we examined whether inter-ocular suppression of the target was necessary for observing
the eye-specific learning effects. The results of Experiment 2a showed no eye-specific learning
effects. To avoid any unwanted influence of the ceiling effect, we used more difficult tasks in
Experiment 2b and Experiment 2c. Again, no eye-specific learning effects were observed.
Therefore, inter-ocular suppression seems to be a necessity for the finding in Experiment 1a.
Without inter-ocular suppression during the training, the targets were represented by the activities
of both monocular and binocular neurons. However, the relationship between rewards and neuronal
responses was different in these two neuronal populations. In case of breakthrough, the firings of
monocular neurons were either 100% (for the rewarded eye) or 0% (for the non-rewarded eye)
predictive of subsequent rewards, yet the firings of binocular neurons were always 50% predictive
of rewards. The absence of the eye-specific learning effects thus indicated that in Experiment 2 the
reward coding system weighted heavily on the activities of binocular neurons and ignored the eye-
of-origin information. This is possible given that binocular neurons greatly outnumber monocular
neurons in the visual cortex.

Is inter-ocular suppression sufficient to observe the eye-specific learning effects? As indicated
by our Experiment 3, this is not the case. As shown in Figure 3, one group of participants showed
only the orientation-specific learning, while the learning of the other group of participants relied on
both the orientation and the eye-of-origin information. In Experiment 3, a rewarding target was
defined as a feature conjunction, so that both the orientation and eye-of-origin had a 50%
probability to be a feature of the rewarding target, though the actual probability for the rewarding
target was 25%. However, over training the participants might only realize that one of the two
orientations was reward-associated, while still unaware of the relationship between rewards and
eye-of-origin. As shown in Figure 3B, the first group of participants only showed the orientation-
specific learning, although rewards were only preceded by half of the targets that were in the
reward-associated orientation (i.e. RO-RE rather than RO-NE), they might treat both the RO-RE
and RO-NE targets as a single type of targets that sometimes (50% probability) brought rewards
at the time of breaking into awareness. As a result, targets in the reward-associated orientation,
which were probably expected by the participants, were more readily to break into awareness than
those in the perpendicular orientation. The increase of breakthrough ratio was thus similar for the
two conditions. In this case, the reward coding system seemed to no longer work in an
unsupervised mode based solely on the eye-of-origin information as in Experiment 1a. Instead, it
worked in a supervised mode, and selectively strengthened the representations in the expected
orientation. This behavior of the reward coding system is very similar to its role in teaching attention
to make selections which is advocated by many other researchers. Importantly, this supervised-

mode mechanism seems to override the unsupervised-mode mechanism, leading to an absence of the eye specificity. By contrast, the learning in the second group showed a more complex pattern. These participants learned even more slowly than the first group, and in the last training session they showed both an orientation-specific and eye-specific learning effect. We presume that the slower development of the orientation-specific learning may indicate a lower efficiency of the supervised-mode mechanism in these participants. Accordingly, the unsupervised-mode mechanism might have the chance to play a role, eventually resulting a significant eye-specific learning effect.

As compared to Experiment 1a, the more complicated reward rule in Experiment 3 might hamper a fast reward-based learning. Consistent with this conjecture, both groups of participants in Experiment 3 learned more slowly than the subjects in Experiment 1a. Alternatively, the sluggish learning in Experiment 3 might result from the additional use of a consciously accessible feature (i.e. orientation), whereas in Experiment 1a only the consciously inaccessible feature was used to associate with rewards. This also agrees with Raio and colleagues' finding that participants learned more slowly when they were aware of the reward-associated stimuli than when not (Raio et al., 2012). Besides different temporal pattern of the learning between Experiments 1a and 3, we also noticed that the learning effect was immediately extinct in the post-test of Experiment 1a, yet still detectable in Experiment 3. Although illuminating this interesting diversity still awaits more systematic future work, we provide a bold surmise that the storage of a learned reward association may be strengthened by the consciousness, thus can be easier or only works when the participants consciously realize the reward-associated feature.

We therefore propose that the reward coding system can produce two different types of reward-based learning. One of them is independent of the consciousness yet fairly consuming attentional resource, likely occurring as early as in V1. We call it the unsupervised learning effect. The other type of learning rests on the close interactions between reward and selective attention, which we call the supervised learning effect. The supervised learning effect can override the unsupervised learning effect when voluntary attention participates in selecting stimuli of behavioral significance. In our Experiment 3a, this is probably because top-down attention selectively enhances the representations in the expected orientation regardless of their eye-of-origin information. With respect to the temporal nature, the eye-specific unsupervised reward learning effect is likely to develop very fast in the training (e.g. reaching the asymptote within the first block of the first training session in Experiment 1a), and highly depends on the context of reward delivery. As soon as the context of reward delivery was absent (e.g. in the post-test), the effect was also absent. Once the supervised reward learning joined, the temporal pattern of the learning changed, hinting distinct timescales between the two types of reward-based learning.

In summary, the present study reports a novel type of reward learning that is established specific to one eye. Physically identical targets were presented either to the rewarded eye or to the non-

rewarded eye, yet rewards were delivered only after the targets in the rewarded eye broke into

awareness. Over training, the breakthrough for the rewarded eye was facilitated more than that for

the non-rewarded eye. This phenomenon was only reliably observed when the targets were

rendered invisible using inter-ocular suppression and the only difference in the visual input signals

between the rewarding and non-rewarding conditions was the eye-of-origin information. Our

findings suggest that human's reward coding system can produce an unsupervised reward learning

effect independent of the consciousness.

Materials and Methods

Participants. Thirty-six participants (17 males and 19 females) who were screened from 105

volunteers participated in Experiment 1a. Fourteen participants (7 males and 7 females) who were

screened from 45 volunteers finished Experiment 1b. Fifteen participants (5 males and 10 females)

who were screened from 57 volunteers finished Experiment 2a. Two groups of fourteen participants

finished Experiment 2b (6 males and 8 females) and Experiment 2c (4 males and 10 females).

Eighty-three volunteers were recruited for Experiment 3 and 34 of them passed the screen test.

Eighteen (12 males and 6 females) participated in Experiment 3a, 16 (5 males and 11 females)

participated in Experiment 3b. The participants ranged in age from 18 to 28 y, all had normal or

corrected-to-normal vision and were naïve to the experimental hypotheses. Experimental

procedures were approved by the Institutional Review Board of the Institute of Psychology, Chinese

Academy of Sciences, and informed consent was obtained from all participants.

Apparatus. Stimuli were presented on a 21-in Dell CRT monitor with a resolution of 1024 × 768

pixels at a refresh rate of 85 Hz, and programmed in Matlab and Psychtoolbox 3 (Brainard, 1997).

The display was calibrated with a Photo Research PR-655 spectrophotometer. To calibrate the

display, we measured the luminance gamma curves and inverted them with a look-up table. The

mean luminance of the screen was 50.9 cd/m^2. A chin-rest was used to help minimize head

movement.

Stimuli and Procedures.

Experiment 1

Experiment 1a. Stimuli were dichoptically presented on a mid-grey background. The target was a

dark gray square frame (1.2° × 1.2°, linewidth: 0.11°) with a horizontal or vertical bar (length: 0.8°,

linewidth: 0.11°) in the center. The target was displayed foveally in one eye, centered 0.25° away

from the central fixation point (0.2°). The central bar could be vertically oriented to the left or right

of the fixation point, or horizontally oriented above or below the fixation point (see Figure 1A). The

CFS stimuli (8° × 8°, flashing at 10 Hz) were displayed foveally in the other eye, which consisted

of 60 Mondrian patterned images created by drawing rectangles of random colors and sizes. A

black-and-white square frame (11° × 11°, linewidth: 0.11°) and the central fixation point were
always presented binocularly to help fusion.

Each trial started with a presentation of the central fixation point for 800 ms. Afterwards, the CFS
stimuli appeared in one eye and kept flashing until the end of the trial. The target appeared in the
other eye after a random interval varied between 100 and 400 ms. The contrast of the target ramped
up to its highest level within the initial 1500 ms, and then remained at the highest contrast level for
500 ms. Participants were required to report the position of the central bar of the target relative to
the fixation once the target broke into awareness. They were told to respond as quickly as possible
on the premise of accuracy. The trial terminated once a response was made, otherwise the target
would be displayed for 2000 ms in total followed by a 600-ms blank interval while the CFS stimuli
were still presented in the other eye. To prevent any afterimage of the target, the target kept drifting
back and forth for 0.08° at 1 Hz along the diagonal.

Each volunteer first participated in a screen test to find individuals with relatively balanced eyes
and the stimuli contrast for non-extreme breakthrough ratio for each eye. Specifically, for targets of
a certain contrast, the difference of break ratio across the eyes should not exceed 20%, while the
break ratio for each eye had to be between 20% and 60%. We used this criterion (around 40%) in
case the estimation of training induced increment of breakthrough ratio was affected by a floor
effect or a ceiling effect from repeatedly performing a b-CFS task. The criterion was decided in a
preliminary stage of the study. Eventually, the break ratios of 29/36 of the participants were within
this range. For the other 7 participants, we used a less stringent criterion (18% - 65%) to have a
larger sample. The contrasts for the target and CFS determined by the screen test were then used
in the subsequent formal experimental sessions. The optimal contrasts used for each participant
were listed in the Table S1, as well as the breakthrough ratios for each eye.

After the pre-test, the participants completed two training sessions, and a post-test. Each session
consisted of two blocks of 160 trials. In the pre- and post-tests, there were no monetary rewards.
The target was presented to the left eye in half of the trials, and to the right eye in the rest of the
trials. The two conditions of trials were randomly interleaved. In the training sessions, for each
participant one eye was assigned to be the rewarded eye. Participants were not aware of this
setting. The selection of the rewarded eye was counter-balanced across the participants. A trial
was called a rewarding trial if the target was presented to the rewarded eye. Immediately after a
correct response for a rewarding trial, a 500-Hz tone would beep for 50 ms, which notified the
participants of winning 0.2 yuan. After each block, a message on the screen showed the
participants the total amount of gain.

Experiment 1b. The stimuli of b-CFS task were similar to those in Experiment 1a, except that the
target was a capital letter "T" in a squared frame. The target was presented at 2° eccentricity above
or below the center of the screen. The letter has four orientations (upright, upside down, right tilt,

left tilt), participants were asked to report the letter orientation by pressing the corresponding arrow

key. Simultaneously with the b-CFS task, participants were required to complete a central RSVP

task. A series of capital letters were binocularly presented in a central white circle (0.6° in diameter)

during the presentation of CFS stimuli. Each letter subtended for 0.5° and was presented for 250

ms. The task was to find "O" in the letter series. A trial lasted for 3700-4000 ms or until the press

of an arrow key was detected. Only one letter "O" was presented in each trial, and it was not

presented at the beginning or the last 200 ms. In each block, 20 trials were planned to be catch

trials without RSVP target. However, since a trial may end before the presentation of "O", there

would be more catch trials. We calculated the hit rate and false alarm rate to measure the

performance of RSVP task. A hit was the response that was made after the presentation of "O" and

before the end of the trial. A false alarm was the response that was made before or without the

presentation of "O" in a trial.

The screen test and procedure were same as those in Experiment 1a. In the training sessions,

participants were told that the reward they could receive was firstly rely on the b-CFS task, while

the hit rate of RSVP task would serve as a discount ratio to the overall reward.

Experiment 2

Experiment 2a. The stimuli and task were similar to Experiment 1a. However, there were two

different types of trials in the training sessions, with-CFS trial and target-only trial. The with-CFS

trial was identical to a typical trial in Experiment 1a. In a target-only trial, no CFS stimuli were

presented, and a target ramped up from 0 to -0.8 contrast (Weber contrast, $C = (L_s – L_b) / L_b$, where

L_s and L_b denoted the luminance of the stimulus and background) within 2 s in one of the two eyes.

The screen test and pre-test were the same as those in Experiment 1a. During the training

sessions, one eye was assigned to be the rewarded eye. However, rewards just occurred in the

target-only trials where the target appeared in the rewarded eye. A correct response in a rewarding

trial would bring a 500-Hz beep and give rise to a reward of 0.31 yuan. In this experiment, all the

participants could finish the task with nearly perfect performance. Therefore, in order to make the

total amount of rewards slightly different across the participants, a random amount (ranging from -

5.00 to 5.00 yuan) was added to the final gain for each block before it was shown on the screen.

Each participant completed a pre-test and three training sessions.

Experiment 2b. The stimuli were sinusoidal gratings (3° in diameter, 1.5 cpd), presented

monocularly on the center of a mid-grey background. The orientation of gratings was fixed for each

participant (either vertical or horizontal), but was counter-balanced across participants.

Participants performed a two-interval forced-choice (2IFC) task. They were required to detect in

which interval the grating was presented. Three practice sessions and four formal test sessions

were completed. The contrasts of test gratings were manipulated by a 2-down-1-up staircase in the

practice sessions. Sixty contrast levels were predetermined for the staircase, ranging
logarithmically from 0.4% (Michelson contrast) to 4% (though 10% for the first practice session for
an easier task). The first practice session included only one block, which was used for the
participants to get familiar with the task. A block contained two interleaved staircases. Each
staircase consisted of 60 trials and started with the highest contrast level. The test contrast
decreased after two successive correct responses and increased after every wrong response. The
step size for the staircase was initially three contrast levels and was reduced to one contrast level
after three reversals. The procedure was similar for the other two practice sessions except that
each session contained two blocks. Contrast threshold for each staircase was calculated by
averaging the contrast levels of the last six reversals. The mean threshold of the latter two practice
sessions was used to determine the seven contrast levels in the formal experiments, which were
designed with the constant stimuli method. Similar to the formal experiments, feedback beeps
(1200 Hz) were given after the response. However, they were delivered randomly in half of the
trials with correct responses and participants were told to ignore the beep.

After the practice, participants finished four formal test sessions, a pre-test session, two training
sessions and a post-test session. The task was same as that used in the practice experiment.
Seven test contrast levels ranged logarithmically from one fourth to three times of the threshold
estimated in the practice sessions were used. For each eye, every contrast level was tested 50
times, resulting in 700 trials per session. The test eye and contrast were randomly selected in each
trial. A session was divided into 4 blocks, allowing the participants to take a break after every block.
The contrast threshold of each eye was estimated by fitting the accuracies at all contrast levels with
a Weibull function (82% correct performance).

Unbeknown to the participants, one eye was selected to be the rewarded eye before the formal
experiments. The selection of rewarded eye was counter-balanced across participants. Every
correct response for a rewarding trial was accompanied with an auditory feedback (1200 Hz).
However, only in the training sessions, participants were informed that the high frequency beep
meant an extra money reward of 0.77 yuan. After each training block, a message on the screen
showed the participants how much they had earned.

Experiment 2c. The stimuli were gratings with the contrast of 80%, and the orientations were about
45°. In each trial, the stimuli were presented monocularly. Participants performed an orientation
discrimination task by judging whether the second grating tilted clockwise or counter-clockwise to
the first one.

After a few practice sessions, participants performed a pilot session where their orientation
discrimination thresholds were measured. The orientation difference between the reference and
test gratings was adjusted according to a staircase procedure. Each practice session included two
interleaved staircases, one for each eye. The pilot session included four interleaved staircases, two

for each eye. Every staircase contained 50 trials. Fifty levels of the orientation offset were
predetermined for the staircase, ranging logarithmically from 0.1° to 10°. The mean orientation
offsets from the last six reversals of each staircase were calculated as the orientation discrimination
threshold for each eye respectively (71% correction threshold).

In the formal experiments, participants finished a pre-test session and 3 training sessions. Each
session consisted of 5 blocks (100 trials per block). For each eye, four subthreshold offset levels
(0.10, 0.23, 0.37, 0.5 × individual threshold) and a threshold offset level were determined. The
threshold level was set to ensure participants could discrimination the orientation difference in some
of the trials so that they would not give up on the task. In each block, 10 trials were tested for each
offset level.

There was no reward in the pre-test. In the training sessions, one eye was selected as the
rewarded eye. For trials in which stimuli with the maximum orientation offset (threshold level) were
presented to the rewarded eye, a beep (1300 Hz) sounded immediately after the participants made
a correct response. For stimuli with subthreshold orientation offset and were presented to the
rewarded eye, the beep was given regardless of whether the responses were correct or not.
Participants were not informed under what circumstance the reward would be given, but were
instructed that each beep meant that they had earned a certain amount of reward (0.09 yuan/trial).
The total gain was presented on the screen after each block. It should be noted that, since
participants were rewarded according to the correction their responses for only 10 trials in each
block, the amount of reward could be nearly equal across the blocks. To increase the variance of
reward over blocks, a random value was either added to or subtracted from the actual money
participants had earned.

Experiment 3. The stimuli were similar to those in Experiment 1a, except that the target was either
a vertical or horizontal bar (length: 1.4°, linewidth: 0.2°) without the square frame (see Figure 3A).
This could make a vertical bar more distinguishable from a horizontal bar.

Like Experiment 1a, participants were required to detect a target in one eye suppressed by the
CFS stimuli in the other eye. Because there were two different targets (vertical or horizontal) and
the target could be presented to one of the two eyes, there were four conditions in this experiment
(vertical target in the left eye, vertical target in the right eye, horizontal target in the left eye, and
horizontal target in the right eye). Only one of the four conditions (counter-balanced across the
participants) was assigned to be the rewarding condition, while the other three conditions were
non-rewarding conditions. A correct response in a rewarding trial would produce a reward of 0.5
yuan accompanied by an audio feedback. The gross of rewards was listed in a message on the
screen after the end of each block. The four types of trials were randomly interleaved within a
session.

Since the eye specific learning effect in Experiment 1a was no longer observed once the reward
was withdrew in the post-test, we first asked a group of participants to complete the experiment
with a screen test, a pre-test, and three training sessions (Experiment 3a). Another group of
participants completed the experiment with an extra post-test without reward after training
(Experiment 3b) to test the persistence of the learning effect.
564

Acknowledgments

This research was supported by the National Natural Science Foundation of China (31871104,
31571112, 31500888, and 31830037) and the Key Research Program of Chinese Academy of
Sciences (XDB02010003 and QYZDB-SSW-SMC030).

Author Contributions

M.B. conceived the study; X.D., M.Z., Y.J. and M.B. designed the experiments; X.D., M.Z. and B.D.
performed the experiments; X.D., M.Z. and M.B. analyzed the data; X.D. and M.B. wrote the paper.

	Conception	Exp. design	Data collection	Data analysis	Paper writing
Exp. 1a		MZ, MB	BD, XD, MZ	MZ, MB	
Exp. 1b		XD, MB, YJ	XD	XD	
Exp. 2a	MB	MZ, MB	MZ	MZ, MB	XD, MB
Exp. 2b		XD, MB, YJ	XD	XD	
Exp. 2c		XD, MB, YJ	XD	XD	
Exp. 3a		MZ, MB	MZ	MZ, MB	
Exp. 3b		XD, MB, YJ	XD	XD	

References

Anderson, B. A. (2017). Reward processing in the value-driven attention network: reward signals
tracking cue identity and location. *Soc Cogn Affect Neurosci, 12*(3), 461-467.
doi:10.1093/scan/nsw141
Anderson, B. A., Laurent, P. A., & Yantis, S. (2011). Value-driven attentional capture. *Proc Natl
Acad Sci U S A, 108*(25), 10367-10371. doi:10.1073/pnas.1104047108
Arsenault, J. T., Nelissen, K., Jarraya, B., & Vanduffel, W. (2013). Dopaminergic reward signals
selectively decrease fMRI activity in primate visual cortex. *Neuron, 77*(6), 1174-1186.
doi:10.1016/j.neuron.2013.01.008
Brainard, D. H. (1997). The Psychophysics Toolbox. *Spat Vis, 10*(4), 433-436.
doi:10.1163/156856897X00357
Brascamp, J. W., Van Boxtel, J. J., Knapen, T. H., & Blake, R. (2010). A dissociation of attention
and awareness in phase-sensitive but not phase-insensitive visual channels. *J Cogn
Neurosci, 22*(10), 2326-2344. doi:10.1162/jocn.2009.21397
Chubykin, A. A., Roach, E. B., Bear, M. F., & Shuler, M. G. (2013). A cholinergic mechanism for
reward timing within primary visual cortex. *Neuron, 77*(4), 723-735.
doi:10.1016/j.neuron.2012.12.039

Dalley, J. W., McGaughy, J., O'Connell, M. T., Cardinal, R. N., Levita, L., & Robbins, T. W. (2001).
Distinct changes in cortical acetylcholine and noradrenaline efflux during contingent and
noncontingent performance of a visual attentional task. *J Neurosci, 21*(13), 4908-4914.
doi:10.1523/JNEUROSCI.21-13-04908.2001
Della Libera, C., & Chelazzi, L. (2006). Visual selective attention and the effects of monetary
rewards. *Psychol Sci, 17*(3), 222-227. doi:10.1111/j.1467-9280.2006.01689.x
Fang, F., & He, S. (2005). Cortical responses to invisible objects in the human dorsal and ventral
pathways. *Nat Neurosci, 8*(10), 1380-1385. doi:10.1038/nn1537
Hickey, C., Chelazzi, L., & Theeuwes, J. (2010). Reward changes salience in human vision via the
anterior cingulate. *J Neurosci, 30*(33), 11096-11103. doi:10.1523/JNEUROSCI.1026-
10.2010
Jiang, Y., Costello, P., & He, S. (2010). Processing of Invisible Stimuli: Advantage of Upright Faces
and Recognizable Words in Overcoming Interocular Suppression. *Psychol Sci, 18*(4), 349-
355. doi:10.1111/j.1467-9280.2007.01902.x
Liu, C. H., Coleman, J. E., Davoudi, H., Zhang, K., & Hussain Shuler, M. G. (2015). Selective
activation of a putative reinforcement signal conditions cued interval timing in primary
visual cortex. *Curr Biol, 25*(12), 1551-1561. doi:10.1016/j.cub.2015.04.028
Marx, S., & Einhauser, W. (2015). Reward modulates perception in binocular rivalry. *J Vis, 15*(1),
15 11 11. doi:10.1167/15.1.11
Mastropasqua, T., Tse, P. U., & Turatto, M. (2015). Learning of monocular information facilitates
breakthrough to awareness during interocular suppression. *Atten Percept Psychophys,*
77(3), 790-803. doi:10.3758/s13414-015-0839-z
Persichetti, A. S., Aguirre, G. K., & Thompson-Schill, S. L. (2015). Value is in the eye of the
beholder: early visual cortex codes monetary value of objects during a diverted
attention task. *J Cogn Neurosci, 27*(5), 893-901. doi:10.1162/jocn_a_00760
Pleger, B., Blankenburg, F., Ruff, C. C., Driver, J., & Dolan, R. J. (2008). Reward facilitates tactile
judgments and modulates hemodynamic responses in human primary somatosensory
cortex. *J Neurosci, 28*(33), 8161-8168. doi:10.1523/JNEUROSCI.1093-08.2008
Proshansky, H., & Murphy, G. (1942). The effects of reward and punishment on perception.
Journal of Psychology: Interdisciplinary and Applied, 13, 295-305.
doi:10.1080/00223980.1942.9917097
Raio, C. M., Carmel, D., Carrasco, M., & Phelps, E. A. (2012). Nonconscious fear is quickly
acquired but swiftly forgotten. *Curr Biol, 22*(12), R477-479.
doi:10.1016/j.cub.2012.04.023
Raymond, J. E., & O'Brien, J. L. (2009). Selective visual attention and motivation: the
consequences of value learning in an attentional blink task. *Psychol Sci, 20*(8), 981-988.
doi:10.1111/j.1467-9280.2009.02391.x
Rossi, V., Vanlessen, N., Bayer, M., Grass, A., Pourtois, G., & Schacht, A. (2017). Motivational
Salience Modulates Early Visual Cortex Responses across Task Sets. *J Cogn Neurosci,*
29(6), 968-979. doi:10.1162/jocn_a_01093
Schultz, W. (2000). Multiple reward signals in the brain. *Nat Rev Neurosci, 1*(3), 199-207.
doi:10.1038/35044563
Seitz, A. R., Kim, D., & Watanabe, T. (2009). Rewards evoke learning of unconsciously processed
visual stimuli in adult humans. *Neuron, 61*(5), 700-707.
doi:10.1016/j.neuron.2009.01.016
Serences, J. T. (2008). Value-based modulations in human visual cortex. *Neuron, 60*(6), 1169-
1181. doi:10.1016/j.neuron.2008.10.051

Shuler, M. G., & Bear, M. F. (2006). Reward timing in the primary visual cortex. *Science, 311*(5767), 1606-1609. doi:10.1126/science.1123513

Thorndike, E. L. (1911). *Animal intelligence*. New York: Macmillan.

Tsuchiya, N., & Koch, C. (2005). Continuous flash suppression reduces negative afterimages. *Nat Neurosci, 8*(8), 1096-1101. doi:10.1038/nn1500

Van Boxtel, J. J., Tsuchiya, N., & Koch, C. (2010). Opposing effects of attention and consciousness on afterimages. *Proceedings of the National Academy of Sciences, 107*(19), 8883-8888. doi:10.1073/pnas.0913292107

Vickery, T. J., Chun, M. M., & Lee, D. (2011). Ubiquity and specificity of reinforcement signals throughout the human brain. *Neuron, 72*(1), 166-177. doi:10.1016/j.neuron.2011.08.011

Whitton, A. E., Treadway, M. T., & Pizzagalli, D. A. (2015). Reward processing dysfunction in major depression, bipolar disorder and schizophrenia. *Curr Opin Psychiatry, 28*(1), 7-12. doi:10.1097/YCO.0000000000000122

Wilbertz, G., van Slooten, J., & Sterzer, P. (2014). Reinforcement of perceptual inference: reward and punishment alter conscious visual perception during binocular rivalry. *Front Psychol, 5*, 1377. doi:10.3389/fpsyg.2014.01377

Wolfe, J. M., & Franzel, S. L. (1988). Binocularity and visual search. *Percept Psychophys, 44*(1), 81-93. doi:10.3758/BF03207480

Xue, X., Zhou, X., & Li, S. (2015). Unconscious reward facilitates motion perceptual learning. *Visual Cognition, 23*, 1-18. doi:10.1080/13506285.2014.981625

Zhang, P., Jiang, Y., & He, S. (2012). Voluntary attention modulates processing of eye-specific visual information. *Psychol Sci, 23*(3), 254-260. doi:10.1177/0956797611424289

Zold, C. L., & Hussain Shuler, M. G. (2015). Theta Oscillations in Visual Cortex Emerge with Experience to Convey Expected Reward Time and Experienced Reward Rate. *J Neurosci, 35*(26), 9603-9614. doi:10.1523/JNEUROSCI.0296-15.2015

Tables

Table 1. Statistics for the results of repeated measurements ANOVA.

		Before FA correction	After FA correction
Exp.1	Session	$F(3, 105) = 49.768$, $p < 0.001$, $\eta^2 = 0.587$	$F(3,105) = 44.471$, $p < 0.001$, $\eta^2 = 0.560$
	Eye-of-origin	$F(1, 35) = 3.785$, $p = 0.060$, $\eta^2 = 0.098$	$F(1, 35) = 4.107$, $p = 0.050$, $\eta^2 = 0.105$
	Session × Eye-of-origin	$F(3,105) = 7.492$, $p < 0.001$, $\eta^2 = 0.176$	$F(3, 105) = 7.348$, $p < 0.001$, $\eta^2 = 0.174$
Exp.2a	Session	$F(3, 42) = 7.795$, $p < 0.001$, $\eta^2 = 0.358$	$F(3, 42) = 6.849$, $p = 0.001$, $\eta^2 = 0.329$
	Eye-of-origin	$F(1, 14) = 1.541$, $p = 0.235$, $\eta^2 = 0.099$	$F(1, 14) = 1.828$, $p = 0.198$, $\eta^2 = 0.115$
	Session × Eye-of-origin	$F(3, 42) = 1.293$, $p = 0.289$, $\eta^2 = 0.085$	$F(3, 42) = 1.171$, $p = 0.332$, $\eta^2 = 0.077$
Exp.3a	Session	$F(2.052, 34.878) = 28.364$, $p < 0.001$, $\eta^2 = 0.625$, Greenhouse-Geisser corrected	$F(2.037, 34.634) = 23.889$, $p < 0.001$, $\eta^2 = 0.584$, Greenhouse-Geisser corrected
	Eye-of-origin	$F(1, 17) = 0.293$, $p = 0.595$, $\eta^2 = 0.017$	$F(1, 17) = 0.159$, $p = 0.695$, $\eta^2 = 0.009$
	Target Orientation	$F(1, 17) = 9.774$, $p = 0.006$, $\eta^2 = 0.365$	$F(1, 17) = 4.562$, $p = 0.048$, $\eta^2 = 0.212$

	Target Orientation × Session	$F(3, 51) = 20.089$, $p < 0.001$, $\eta^2 = 0.542$	$F(2.158, 36.689) = 10.466$, $p < 0.001$, $\eta^2 = 0.381$, Greenhouse-Geisser corrected
	Eye-of-origin × Target Orientation	$F(1, 17) = 3.209$, $p = 0.091$, $\eta^2 = 0.159$	$F(1, 17) = 1.982$, $p = 0.177$, $\eta^2 = 0.104$
	Eye-of-origin × Session	$F(3, 51) = 1.161$, $p = 0.334$, $\eta^2 = 0.064$	$F(2.139, 36.371) = 0.941$, $p = 0.405$, $\eta^2 = 0.052$, Greenhouse-Geisser corrected
	Session × Eye-of-origin × Target Orientation	$F(2.049, 34.831) = 0.115$, $p = 0.896$, $\eta^2 = 0.007$, Greenhouse-Geisser corrected	$F(1.932, 32.847) = 0.173$, $p = 0.834$, $\eta^2 = 0.010$, Greenhouse-Geisser corrected
Exp.3b	Session	$F(4,60) = 16.782$, $p < 0.001$, $\eta^2 = 0.528$	$F(4,60) = 15.438$, $p < 0.001$, $\eta^2 = 0.507$
	Eye-of-origin	$F(1,15) = 1.916$, $p = 0.187$, $\eta^2 = 0.113$	$F(1,15) = 1.676$, $p = 0.215$, $\eta^2 = 0.101$
	Target Orientation	$F(1,15) = 0.073$, $p = 0.791$, $\eta^2 = 0.005$	$F(1,15) = 0.076$, $p = 0.786$, $\eta^2 = 0.005$
	Target Orientation × Session	$F(4,60) = 2.491$, $p = 0.053$, $\eta^2 = 0.142$	$F(4,60) = 1.101$, $p = 0.365$, $\eta^2 = 0.068$
	Eye-of-origin × Target Orientation	$F(1,15) = 1.258$, $p = 0.280$, $\eta^2 = 0.077$	$F(1,15) = 0.954$, $p = 0.344$, $\eta^2 = 0.060$
	Eye-of-origin × Session	$F(4,60) = 4.016$, $p = 0.030$, $\eta^2 = 0.211$, Greenhouse-Geisser corrected	$F(4,60) = 4.021$, $p = 0.029$, $\eta^2 = 0.211$, Greenhouse-Geisser corrected
	Session × Eye-of-origin × Target Orientation	$F(4,60) = 0.913$, $p = 0.462$, $\eta^2 = 0.057$	$F(4,60) = 0.611$, $p = 0.656$, $\eta^2 = 0.039$

668   Here, FA denotes false alarm.