1

Molecular layer interneurons in the cerebellum encode for valence in associative

learning

Ming Ma[1,2], Gregory L. Futia[3], Fabio M. Simoes De Souza[2,4], Baris N. Ozbay[5], Isabel

Llano[6], Emily A. Gibson[1,3] and Diego Restrepo[1,2,*]

[1]Neuroscience Graduate Program, University of Colorado Anschutz Medical Campus,

Aurora, CO 80045, USA

[2]Department of Cell and Developmental Biology, University of Colorado Anschutz

Medical Campus, Aurora, CO 80045, USA

[3]Department of Bioengineering, University of Colorado Anschutz Medical Campus,

Aurora, CO 80045, USA

[4]Center for Mathematics, Computation and Cognition, Federal University of ABC, Sao

Bernardo do Campo, SP, Brazil.

[5]Intelligent Imaging Innovations, Denver, CO, 80216.

[6]Saints Pères Paris Institute for Neurosciences, Université Paris Descartes, 75006 Paris,
France

[*]Corresponding author: Diego Restrepo, diego.restrepo@cuanschutz.edu

We used two-photon microscopy to study the role of ensembles of cerebellar molecular

layer interneurons (MLIs) in a go-no go task where mice obtain a sugar water reward if

they lick a spout in the presence of the rewarded odorant and avoid a time out when they

refrain from licking for the unrewarded odorant. In naïve animals the MLI responses did

not differ between the odorants. With learning, the rewarded odorant elicited a large

increase in MLI calcium responses, and the identity of the odorant could be decoded from

the differential response. Importantly, MLIs switched odorant responses when the valence

of the stimuli was reversed. Finally, mice took a longer time to refrain from licking in the

presence of the unrewarded odorant and had difficulty becoming proficient when MLIs

were inhibited by chemogenetic intervention. Our findings support a role for MLIs in

learning valence in the cerebellum.

The cerebellum plays a pivotal role in coordinating movements through sensorimotor

integration. It receives massive input through mossy fiber (MF) synapses onto granule cells

(GCs) to form a circuit efficient in complex pattern separation[1] (Supplementary Fig. 1). In

addition, the cerebellum contributes to cognition and emotion and is associated with non-

motor conditions such as autism spectrum disorders[2-5]. The Purkinje cells (PCs), the sole

projection neurons of the cerebellar cortex, receive excitatory afferent input from the

parallel fibers (PFs) of GCs and dense feedforward inhibitory inputs from the molecular

layer interneurons (MLIs). Importantly, subtle changes in this PC excitatory-inhibitory

balance generate robust, bidirectional changes in the output of PCs[6].

Plasticity in cerebellar circuit activity plays an important role in generation of adequate

output. Indeed, long term depression (LTD) mediated by dendritic increases in Ca^{2+} in PCs

elicited by motor error signals conveyed by climbing fibers (CFs) is a classical model of

plasticity[7-10]. However, recent studies indicate that CFs also signal reward prediction[11,12]

or decision-making errors[13], and the cerebellum modulates association pathways in the

ventral tegmental area (VTA) contributing to reward-based learning and social behavior[14].

Furthermore, although LTD at the PF-PC synapse is often considered as the substrate for

cerebellar dependent learning[7-10], such learning can occur in the absence of LTD and may

therefore involve other forms of plasticity[15]. A potential substrate for plasticity is the PF-

MLI synapse[16,17] where LTP can be induced in slices by pairing MLI depolarization with

PF stimulation[18] and *in vivo* by conjunctive stimulation of PFs and CFs[19], believed to

underlie changes in the size of cutaneous receptive fields[20,21]. Additionally, high frequency

stimulation of PFs alters subunit composition of AMPA receptors, rendering them calcium-

impermeable[18], a long-lasting change linked to behavioral modifications[22]. Interestingly,

MLIs have been proposed to participate in cerebellar plasticity[21,23,24], and Rowan and co-

workers found graded control of PC plasticity by MLI inhibition[25] suggesting that MLI

inhibition is a gate for learning stimulus valence which conveys information as to whether

the stimulus is rewarded. However, whether there is a causal participation of MLIs in

reward-associated learning is unknown.

Here we explored whether MLI activity plays a role in reward-associated learning in a go-

no go task where the thirsty animal learns to lick to obtain a water reward[26,27]. We applied

two-photon microscopy[28] to record Ca^{2+} changes in ensembles of MLIs and utilized

chemogenetics to explore the functional role of MLI activity in learning.

Results

In order to explore the role of MLIs in associative learning we employed *in vivo* two-photon

microscopy to record neural activity reported by changes in fluorescence emitted by the

Ca^{2+} indicators GCaMP6/7. MLIs were imaged within the superficial 50 μm of the ML in

head-fixed mice through a 2x2 mm glass window implanted above the cerebellar vermis

(lobule VI, Fig. 1, Supplementary Fig. 2), where GCs acquire a predictive feedback signal

or expectation reward[29,30]. We used the go-no go task where thirsty mice initiate the trial

by licking on the spout to elicit odorant delivery 1-1.5 sec after the first lick. Mice received

a water reward when they licked at least once in two lick segments during rewarded odorant

delivery (1% isoamyl acetate, Iso, termed S+) (Fig. 1a, Hit trial, mouse movement shown

in Supplementary Movie 1, quantified in Supplementary Fig. 3). Mice did not receive the

reward if they failed to lick in one of the two lick segments (Miss trial). When the

unrewarded odorant was presented (mineral oil, MO, S- trials) the animals did not receive

a reward when they refrained from licking (Correct Rejection, CR, Fig. 1a), and they

received a timeout of ten seconds if they licked in both segments (False Alarm, FA). A

proficient mouse (percent correct>=80%) licked at least once in each lick segment in the

S+ trials and stopped licking during the segments for the S- trials, see lick trace at the

bottom of Fig. 1d.

An example of an increase in MLI GCaMP6f fluorescence intensity during odorant

application in a Hit trial is shown for a proficient mouse in Figs. 1bi,ii (Supplementary

Movie 2). We extracted regions of interest (ROIs) of the components (Fig. 1biii) and

temporal traces of the normalized change in fluorescence ($\Delta F/F$, Figs. 1c and d) using

constrained nonnegative matrix factorization analysis[31]. The average diameter of the ROIs

in this field of view (FOV) was 10.5$\pm$4 µm (mean$\pm$SD, n=170, Supplementary Fig. 4a),

which falls within the range of diameters reported for MLIs[32]. When the animal was

proficient we observed that MLI $\Delta F/F$s increased when the rewarded odorant was presented

(Fig. 1ci and d, vertical orange lines in 1d indicate odorant on and off times for S+). In

contrast, the unrewarded odorant elicited smaller transient increases in $\Delta F/F$ (Fig. 1cii, Fig.

1d vertical light blue lines and Supplementary Figs. 4b,c,d), with some exceptions where

the increases for $\Delta F/F$ for S- were larger (arrow in Fig. 1d). Finally, the rewarded odorant

elicited increases in $\Delta F/F$ in 89% of the ROIs among a total of 191 (Fig. 1c and

Supplementary Figs. 4c and d).

Cerebellar MLIs develop divergent responses as mice learn to differentiate odorants.

The behavioral performance in a learning session where the animal reached proficient level

after 70 trials is shown in Fig. 2a. When the animal was naïve (<=65% correct) the time

courses for $\Delta F/F$ overlapped between S+ and S- (Fig. 2bi), and when the animal became

proficient the time courses for $\Delta F/F$ diverged (Fig. 2biii). For the proficient animal in S+

trials $\Delta F/F$ started increasing at trial initiation (1-1.5 sec before odorant addition) and

reached a peak when the animal was rewarded (Fig. 2biii, orange trace), while for S- trials,

the response returned to basal levels shortly after the odorant was applied (Fig. 2biii, light

blue trace, Fig. 2b shows the time courses for lick rates). Whereas $\Delta F/F$ does not differ

between S+ and S- before odorant addition (Fig. 2c), it diverges after odorant addition as

the animal learns (Fig. 2d) and $\Delta F/F$ per ROI increased when the animal became proficient

(Fig. 2e). GLM analysis yields a statistically significant difference between S+ and S- and

naïve and proficient, p<0.001, 5550 observations, 5538 d.f., 1 mouse, GLM F-

statistic=267, p<0.001. Finally, the mean $\Delta F/F$ ($\pm$CI, 4 mice) for the last second of odorant

application increases as a function of percent correct performance for the S+ condition

whereas it remains stable for the S- condition (Fig. 2g). A GLM analysis finds a statistically

significant difference between S+ and S- (p<0.01, 47 observations, 44 d.f., n=4 sessions, 4

mice, GLM F-statistic=60, p<0.001) and for the interaction between performance and the

odorant (S+ vs. S-) (p<0.001, 47 observations, 44 d.f., n=4 sessions, 4 mice

(Supplementary Fig. 6 shows learning curves for these mice). Therefore, as the animal

learns, the magnitude and temporal dynamics of the change in Ca^{2+} in MLIs diverges

between S+ and S- odorants, suggesting a possible role for MLIs in associative learning.

Interestingly, the majority of the ROIs responded similarly (Fig. 1c, Supplementary Fig. 4)

and a dimensionality analysis indicated that the dimensionality ranged from 2 to 6 (see

Supplementary Note 1 and Supplementary Fig. 7) indicating that the responses are highly

redundant.

The identity of the stimulus can be decoded from MLI activity in proficient mice. The

finding that the MLIs developed divergent responses for S+ vs. S- trials (Fig. 2g) raised

the question whether ensemble neural activity encodes for the identity of the stimulus. The

learning curve for a session where the mouse achieves >=80% in ~60 trials is shown in Fig.

3a. The first two principal components for a principal component analysis (PCA) of $\Delta F/F$

values for all MLIs in this session showed clear differences between S+ and S- trials for

the reward and odorant periods for trials when the mouse was proficient, in contrast to the

overlap observed between odorants for the naïve mouse (Fig. 3b). This suggested that

odorant identity (S+ vs. S-) could be decoded from ΔF/F values for proficient mice.

We utilized a linear discriminant analysis (LDA) to determine whether a hyperplane placed

in the multidimensional space of ΔF/F values for all MLIs in the ensemble could decode

the stimulus. The LDA was trained with ΔF/F for all trials minus one and was queried to

identify the odorant in the remaining trial (see Methods). The bootstrapped 95% confidence

interval (CI) for an LDA trained after shuffling stimulus was used as a control. The time

course for LDA decoding accuracy for the session with the learning curve in Fig. 3a is

shown in Fig. 3c (Supplementary Fig. 5a shows the corresponding lick frequency time

course). For trials with percent correct <=65%, decoding accuracy raises above the shuffled

95% CI after water reinforcement (Fig. 3ci), while for proficient trials, decoding accuracy

starts rising above CI shortly after odorant addition (Fig. 3cii). When tested in four mice

LDA decoding accuracy for the odorant and reinforcement periods differed from shuffled

LDA (Fig. 3d). A GLM analysis yielded statistically significant interactions between

shuffled vs. reinforcement and for the interactions between (shuffled vs. odorant) x

(proficient vs. naïve) and (shuffled vs. reinforcement) x (proficient vs. naïve) (GLM p

values <0.01 and <0.05, 24 observations, 18 d.f., n=4 sessions, 4 mice, GLM F-statistic

16.4, p<0.001). Post hoc tests corrected for multiple comparison using false discovery rate

(FDR)[33] yielded a statistically significant difference for decoding accuracy for either

reinforcement or odorant vs. shuffled for proficient (p<pFDR=0.007, n=4 sessions, 4

mice). Performing the analysis on subsets of ROIs showed that even analysis with a single

ROI yields decoding accuracy that is significantly different from the shuffled LDA

indicating that the information on the stimulus stored in MLI activity is highly redundant

(see Supplementary Note 2, Figs. 3e and f). The LDA analysis revealed that the accuracy

for decoding the odorant identity from $\Delta F/F$ during the odorant period increases as the

animal learns to differentiate between odorants.

MLI odorant responses switch when stimulus valence is reversed. In order to determine

whether the MLIs responded to the chemical identity of the odorant, as opposed to

responding to the valence (contextual identity: is the stimulus rewarded?), we reversed

odorant reinforcement. When the reward was reversed for a proficient mouse the animal

kept licking for the previous rewarded odorant resulting in a fall in percent correct below

50%, and as the animal learned the new valence the percent correct raised back above 80%

(Fig. 4a and Supplementary Figs. 6e and 6f). We analyzed how odorant-elicited changes

in MLI $\Delta F/F$ varied in this reversal task. Fig. 4b shows single trial examples for mean $\Delta F/F$

odorant responses when the reward is reversed. MLIs respond to Iso (S+) with an increase

in $\Delta F/F$ before reversal (e.g. forward trial 52), maintain the odorant increase in $\Delta F/F$ to

what is now the unrewarded odorant (Iso, S-) immediately following reversal (reverse trial

92) and switch responses with increases to the new rewarded stimulus (MO, S+) when they

become proficient in the reversal task (reverse trial 438) ($\Delta F/F$ for the last second of the

odorant period is shown for all trials in Fig. 4c). When repeated for three different mice the

odorant-induced $\Delta F/F$ changes reversed for proficient mice (>=80% correct) when the

valence was reversed (Fig. 4d). GLM analysis indicates that there is a significant difference

in odorant-induced changes in $\Delta F/F$ for the odorant and the reversal (p<0.001, 12

observations, 8 d.f., n=3 sessions, 3 mice, GLM F-statistic=14.8, p<0.01). Thus, after

successful reversal the ΔF/F time course switched for the two odorants: the stimulus-

induced increase in ΔF/F took place for the reinforced odorant, not for the chemical identity

of the odorant.

Next we computed the accuracy for decoding the reinforced odorant for trials when the

animal was proficient in either the forward or reverse trials using LDA analysis. Fig. 4e

shows for the reversal session in Fig. 4a that for the proficient animal decoding accuracy

rises above the 95% CI calculated with shuffled trials shortly after odorant addition for

both forward and reverse trials. Finally, the results of the forward and reverse LDA analysis

for three mice that are proficient showed that decoding accuracy for the stimuli differed

from shuffled LDA (Fig. 4f). GLM analysis indicates that differences in decoding accuracy

are statistically significant between original and shuffled (P<0.001), but not between

forward and reverse (p>0.05, 12 observations, 8 d.f., n=3 sessions, 3 mice, GLM F-

statistic=16, p<0.001). Finally, decoding accuracy of an LDA analysis for contextual

identity for proficient mice did not differ between correct (Hits and CRs), and incorrect

trials (Miss and FAs) (Supplementary Note 3, Supplementary Figs. 8,9). These results

indicate that for the proficient mouse it is possible to decode contextual identity suggesting

that MLI activity encodes for valence.

Changes in lick rate correlate with changes in MLI activity during reinforcement, but

not during odorant application. To explore whether the observed MLI ensemble activity

recorded in vermis is directly related to licking, as found in Crus II[34,35], we examined the

correlation between ΔF/F and lick rate. When the animal was proficient the animal licked

at least once in each of the two lick segments during application of the S+ odorant,

increased licking after receiving the water reward and refrained from licking for the S-

trials (Fig. 5a). For the proficient mouse the lick frequency diverges between S+ and S-

trials shortly after addition of the odorant (Fig. 5b) and this divergence is evidenced by a

large decrease in the p value calculated with a ranksum test comparing licking between S+

and S- trials (Fig. 5c).

In order to explore the relationship of MLI activity to licking we proceeded to examine the

correlation between $\Delta F/F$ time course and the lick rate (LR) and between the derivative of

the $\Delta F/F$ time course ($D_t\Delta F/F$) and the derivative of the lick rate (D_tLR)[34] and plotted their

relationship during different time periods for proficient mice. Fig. 5d shows examples for

several trial outcomes for the time course for $\Delta F/F$ and the lick rate. Examples of these

correlations for a single session for a proficient mouse are shown in Figs. 5e and f. In this

example the correlation between $\Delta F/F$ and LR was larger during the reinforcement period

($\rho=0.73$, p-value<0.001) compared to the odorant ($\rho=0.19$, p-value<0.001) and pre-odorant

($\rho=-0.06$, p-value<0.05) periods (Fig. 5e). Similarly, the correlation between $D_t\Delta F/F$ and

D_tLR is significant for the reinforcement period ($\rho=0.28$, $p<0.001$), and is smaller for both

the pre-odorant ($\rho=-0.08$, $p<0.05$) and odorant ($\rho=-0.045$, $p>0.05$) periods (Fig. 5f). We

proceeded to compare the correlations between these parameters in several mice. As found

in Crus II, $\Delta F/F$ and LR (Fig. 5g) and $D_t\Delta F/F$ and D_tLR (Fig. 5h) are positively correlated

during the reinforcement period. Interestingly, the correlations are lower during the pre-

odorant and odorant periods (Figs. 5g,h). A t test indicated that there is a statistically

significant difference in the correlation coefficient between the reinforced vs. the odorant

or pre-odorant periods for $\Delta F/F$ vs. LR (p<pFDR=0.05, n=6 sessions, 5 mice) and between

the reinforced and odorant periods for $D_t\Delta F/F$ vs. D_tLR (p<pFDR=0.016, n=6 sessions, 5

mice). These data suggest that changes in $\Delta F/F$ during the reinforcement period reflect

changes in lick activity, while changes in $\Delta F/F$ during the odorant period are less

dependent on licks, and maybe dependent on multiple variables. We performed

complementary studies of $\Delta F/F$ in CR trials when the mouse did not lick and for time-

courses aligned to the beginning of $\Delta F/F$ changes after odorant addition (Supplementary

Note 4 and Supplementary Figs. 10 and 11). Under these conditions, $\Delta F/F$ and lick rate are

uncorrelated, consistent with the potential lack of direct relationship between changes in

$\Delta F/F$ and lick rate during the odorant period.

Do the MLIs respond to reward value? The correlation between $\Delta F/F$ time course and

the lick rate (Fig. 5e) raises the question whether $\Delta F/F$ reflects the reward value as opposed

to the valence of the odorant. Here we define the valence of the odorant as indicating

whether the stimulus is good or bad[36]. Thus, valence is a binary measure of an emotion that

is reflected by the motivation to receive reward, whereas reward value, in the present

paradigm, is related to the amount of sugar water delivered. To evaluate whether $\Delta F/F$

changes are dependent on reward value we recorded changes in $\Delta F/F$ from mice performing

a go-go task where both odorants were rewarded equally and we varied the volume of sugar

water delivered for successful trials. As expected, the mouse responded to both odorants

(Supplementary Fig. 12a). Also, as expected, when the volume of reward was increased,

the lick frequency increased during the delivery of sugar water (wet licks, Supplementary

Fig. 12b), but not during dry licking before reward (dry licks, Supplementary Fig. 12b). A

GLM analysis yielded a statistically significant difference for dry vs. wet licking and for

the interaction between the volume of sugar water delivered and dry vs. wet (p<0.001, 198

observations, 194 d.f., 1 session, 1 mouse, GLM F-statistic=21.8, p<0.001). In contrast, a

GLM analysis of the changes in $\Delta F/F$ as a function of volume delivered in dry and wet lick

conditions did not yield statistically significant changes (Supplementary Fig. 12c, p>0.05,

198 observations, 194 d.f., 1 session, 1 mouse, GLM F-statistic=1.7, p>0.05). Finally, if

MLI activity reflected reinforcement value $\Delta F/F$ should show a positive correlation with

volume of sugar water delivered (see Fig. 3 of [34]). We did not find a significant correlation

between lick frequency and $\Delta F/F$ for either dry (Supplementary Fig. 12di, ρ=0.16, p>0.05)

or wet licking (Supplementary Fig. 12dii, ρ=-0.66, p>0.05). This experiment indicates that

MLI activity does not reflect reward value, and is consistent with MLI activity reflecting

valence.

The main contributor during odorant application to a GLM model fit to MLI activity

is the contextual identity of the odorant. In order to understand the contribution of

different variables to MLI activity we setup a GLM to quantify the dependence of the

average $\Delta F/F$ on the different behavioral and stimulus variables[37]. We included "event"

variables, "whole trial" variables and "continuous" variables (Fig. 6a, Methods).

Continuous variables quantified kinematics including lick rate and the derivative of lick

rate (Fig. 5) and body velocity and body acceleration of movements made by the animal

during the trial (Supplementary Fig. 3). The identity of the odorant (S+/S- odorant) was an

event variable that increased from zero to one during the time for odorant application.

Finally, whole trial variables were accuracy (1 for correct and 0 for incorrect responses),

reinforcement history (1 for reinforcement in the last trial, 0 otherwise) and percent correct

behavior calculated in a window of 20 trials.

Black traces in Fig. 6b show examples of the fit of the GLM model to average ΔF/F time

courses for four example trials with different outcomes. The fit traces in black largely

overlap with the recorded data. When quantified over the different periods within a trial

(pre-odorant, odorant and reinforcement) we found that GLM explained a substantial

percent of the average ΔF/F variance ranging from 23.5% to 95% (Fig. 6c). We next

quantified the relative contribution of the different variables to the ensemble activity in the

three time periods. Fig. 6d shows the contributions to the GLM fit by the variables that

make the largest percent contribution to the fit (body kinematics, licks, reinforcement

history and odorants). Odorant (S+ vs. S-) is the dominant contributor to the activity during

odorant application (Fig. 6d,iii, asterisks show differences with ranksum or t test

p<pFDR=0.033, n=6 sessions, 5 mice).   Furthermore, variables describing the licks

contribute to the GLM fit during the outcome period (Fig. 6d,ii, * p<pFDR=0.016 for a t

test, n=6 sessions, 5 mice). In contrast, the other two variables (body kinematics and

reinforcement history) do not differ in contribution during the different periods

(p>pFDR=0.017, ranksum test). These data indicate that odorant identity contributes to

modeling MLI activity during the odorant application period while licks contribute to MLI

activity during the reinforcement period.

Chemogenetic inhibition of MLI activity leads to impaired go-no go learning. In order

to determine whether activity of MLIs plays a role in behavioral responses in the go-no go

task we used a Cre-dependent AAV virus to express the inhibitory DREADDs receptor

hM4Di in MLIs in six PV-Cre mice (hM4Di group)[38]. To control for off-target effects of

clozapine-N-oxide (CNO)[39] we injected PV-Cre mice with Cre-dependent mCherry AAV

virus in another group of six mice (control group) (see Supplementary Fig. 13a,b for virus

expression). One to two weeks later animals from both groups were trained to differentiate

two odorant mixtures (S+: 0.1% of 60% heptanal+40% ethyl butyrate, S-: 0.1% of 40%

heptanal+60% ethyl butyrate) for 3-4 sessions encompassing 400-500 trials in the go-no

go task. The animals were injected intraperitoneally (IP) with saline 40 minutes before the

start of the sessions (control-saline and hM4Di-saline). One to two weeks later mice were

trained again to differentiate between the same odorant mixtures, but they were injected IP

with CNO (3mg/kg) 40 minutes before the start of the session (control-CNO and hM4Di-

CNO).

Figs. 7a,i and b,i show representative examples of the behavioral performance for the four

conditions. Mice in all groups with the exception of the hM4Di-CNO attained proficiency

(>=80% percent correct) (Figs. 7a,ii and 7b,ii). GLM analysis indicates that there is a

statistically significant interaction between CNO drug treatment and hM4Di expression

($p<0.01$, 24 observations, 20 d.f., n=6 mice, GLM F-statistic=13.7, $p<0.001$) and post hoc

tests indicate that the hM4Di expressing group differs between CNO and saline

($p<pFDR=0.025$, n=6 mice per group), while there are no significant differences between

CNO and saline for control mice ($p>pFDR$, n=6 mice per group), indicating an effect of

CNO-induced inhibition of MLIs expressing hM4Di on behavioral output and the absence

of off-target CNO effects.

We proceeded to compare the time course for lick frequency between different conditions

for trials when the mouse was proficient (>=80% correct, Fig. 7c). Interestingly, the time

course for lick frequency differs between hM4Di mice injected with CNO and the other

conditions (Figs. 7c,i and 7c,ii). With the exception of hM4Di + CNO, shortly after odorant

addition (~0.8 sec) the lick frequency for S- does not increase beyond ~1 Hz while the lick

frequency for S+ keeps increasing beyond 8 Hz (Fig. 7c). In contrast, for hM4Di + CNO

lick frequency for S- keeps increasing beyond 1 Hz and does not diverge from the S+ time

course until it reaches 2.5 Hz at a later time point (~1 sec, Fig. 7c,ii) likely reflecting slow

decision-making. This would explain why the hM4Di + CNO mice accumulated errors in

the go-no go task (Fig. 7bii). In order to quantify this difference in lick frequency we

calculated the lick frequency 0.8 to 1.8 seconds after odorant addition (Fig. 7d). For S+

GLM analysis did not find a statistically significant difference for treatment or hM4Di

expression (or interactions) for the rewarded odorant trials (Fig. 7di, $p > 0.05$, 60

observations, 56 d.f., n=6 mice, GLM F-statistic=2.96, $p > 0.05$). Yet GLM found a

statistically significant difference for S- for the interaction between CNO and hM4Di

expression (Fig. 7dii, $p < 0.001$, 60 observations, 56 d.f., n=6 mice, GLM F-statistic=12.8,

$p < 0.001$). We obtained a similar impairment of performance in a separate set of

experiments with two hM4Di mice and two controls where animals discriminated between

1% Iso and MO, and CNO was applied when the animal was naïve, and we reversed the

reward (Supplementary Fig. 13). These data indicate that inhibition of MLIs causes a

slower differential lick response to odorants leading to impaired behavioral performance.

Discussion

We found that vermal MLIs developed a differential response to odorants in the go-no go

task that switched when the valence was reversed. Decoding analysis revealed that when

the animal was proficient the contextual identity of the odorant could be decoded from MLI

responses. GLM analysis revealed that contextual identity made a large contribution to the

fit of MLI activity during the odorant application period. Chemogenetic inhibition of MLIs

impaired achievement of proficient discrimination of odorants. These data indicate that

MLIs play a role in associative learning by encoding valence.

The cerebellum has been implicated in mediating supervised learning through an iterative

process whereby the response to an input is evaluated against a desired outcome, and errors

are used to adjust adaptive elements within the system[8-10,40]. CFs carrying error signals

make profuse synaptic connections on the dendrites of PCs and elicit powerful excitatory

dendritic Ca^{2+} spikelets[41-44]. Furthermore, CFs also signal reward prediction[11,12] or

decision-making errors[13], and the cerebellum modulates association pathways in VTA

enabling a cerebellar contribution to reward-based learning and social behavior[14]. The

increase in Ca^{2+} mediates LTD in subsets of synapses innervated by co-activated GC

parallel fibers (PFs) carrying sensorimotor information relevant to learning[45,46]. However,

recent studies by Rowan et al.[25] revealed that increasing feed forward inhibition by MLIs

can switch the valence of plasticity from LTD to LTP (also, see[47]). In addition, adaptive

changes in the vestibulo-ocular reflex elicited by CF optogenetic activation switched from

increase to decrease depending on whether MLIs were co-activated[25]. Finally, MLIs gate

supralinear CF-evoked Ca2+ signaling in the PC dendrite[48]. These studies suggest that the

valence of learning is graded by MLI activity.

Here we provide evidence for the involvement of MLIs in conveying information on

contextual identity of a stimulus in associative learning. We do not find that the MLIs

respond to odorants per se. Rather, the reversal experiment (Fig. 4) and the similar $\Delta F/F$

responses and stimulus decoding for correct and incorrect behavioral response trials

(Supplementary Fig. 8) indicate that MLIs respond to contextual odorant identity: "is this

the rewarded odorant?", which is directly related to valence, a binary measure of an

emotion reflected by the motivation to receive reward[36]. Furthermore, our results in the go-

go task where both odorants are rewarded with varying volumes of sugar water

(Supplementary Fig. 12) are consistent with the response reflecting valence (as opposed to

value). Thus, we postulate that the MLI response during the odorant period is related to

valence that reflects reward expectation, consistent with the fact that GCs in lobe VI were

found to respond to reward expectation[29,30].

A question that arises is which circuit mechanism is responsible for the decreased

behavioral performance after chemogenetic inhibition of MLI activity (Fig. 7 and

Supplementary Fig. 13). MLIs receive sensorimotor information from multiple GCs

through PF input and *in vivo* studies have found remodeling of MLI receptive fields upon

repeated electrical stimulation of the skin[49]. In addition, plasticity in PF-MLI synapses are

postulated to increase the information capacity of the MLI-PC network and richness of PC

output dynamics[10,16], and a model of PF-MLI plasticity has been proposed[50]. If long term

plasticity of PF-MLI synapses is responsible for the large change of MLI responsiveness

found here upon reversal of stimulus valence it is likely that the error signal would be

provided by CF spillover resulting in highly redundant stimulation of stellate cells

(SCs)[51,52]. We developed a model of the MLI/PC circuit described in Supplementary Note

5 (Supplementary Figs. 14,15) that suggests that plasticity in SC-PC synapses[53], PF-SC

synapses[50], complemented with CF spillover acting on the feedforward disinhibitory MLI

circuit described recently by Arlt and Hausser[52], complemented with plasticity in SC-PC

synapses[53], PF-SC synapses[50], would explain the changes in behavior we find after

chemogenetic inhibition of MLI activity. Finally, we found that the divergence between

the time courses for lick frequency between S+ and S- took place at a later time when MLIs

were inhibited by chemogenetics (Fig. 7c,ii, Supplementary Fig. 13g) likely reflecting slow

decision-making, consistent with a role for the CF/PC circuit in reward timing prediction[54].

Future studies are necessary to understand the role of plasticity in the PF-MLI-PC circuit

in associative learning.

Interestingly, odorant responses have been reported in the cerebellum. Studies in sexually

trained male rats found that female bedding or almond smell elicited increased cFos

immunoreactive GCs in the vermis compared to rats exposed to clean air[55]. Furthermore,

sexual experience increased the number of cFos positive GCs. In humans odorants induced

significant activation of the cerebellum[56]. In addition, human cerebellar lesions caused

olfactory impairments in the contralesional nostril, and elicited sniffs with lower overall

airflow velocity compared to controls[57]. These findings implicate an olfactocerebellar

pathway prominent in odorant identification and detection that functionally connects each

nostril primarily to the contralateral cerebellum. Our study provides further evidence for

involvement of the cerebellum in olfactory tasks.

Here we found changes in MLI activity during learning in the go-no go associative learning

task in lobule VI where GCs[29,30] and CF[58] activity was proposed to encode aspects of

reward signaling. Recent work on the contribution of cerebellar processing to execution of

reward-driven behaviors indicates that multiple cerebellar regions are involved, including

central and lateral cerebellum[11,29,30,58,59]. As behavioral tests are refined, it is likely that

differences in how these regions process reward will emerge, as suggested by the recent

work on climbing-fiber signaling[11]. Furthermore, we focused our recordings on the more

superficial regions of the molecular layer and therefore, most of our measurements are from

SCs. Recordings from synaptically connected pairs of MLIs and PCs in slices showed that

mean the amplitude of synaptic currents decreases with distance from the PC layer,

suggesting a stronger impact of basket versus SC inhibition on PC firing[60]. Recent

recordings of PC spikes in vivo following genetic deletion of MLIs confirm this

prediction[47] which is in accord with the morphological diversity of MLIs[61]. Finally, the

effect of chemogenetics (Fig. 7) should be on both stellate and basket cells, and future

experiments are necessary to differentiate between the roles of the two cell types.

Our findings show that differential MLI activity develops during learning in an associative

learning task and that MLI activity switches when the rewarded odorant is reversed. We

find that inhibition of MLI activity elicits decreased behavioral performance in the go-no

go task. Our data indicate that MLIs have a role in learning valence. This would likely

increase the information capacity of the MLI-PC network and richness of PC output

dynamics[10,62].

Acknowledgments: We thank Ms. Nicole Arevalo for animal husbandry, Ms. Dnate' Baxter

for laboratory support, Ms. Arianna Gentile-Polese for help with the summary figure and

Mr. Jesse Gilmer with help coding dimensionality. We thank Mr. Matt I. Becker, Mr. Jesse

Gilmer and Dr. Abigail Person for discussions. This research was supported by NIDCD

R01 DC000566, NINDS U01 NS099577, NSF CBET-1631912, NSF BIO-1926676 and

ANR-18-CE16-0010-01.

.

Author contributions: M.M., I.Ll. and D.R. designed all experiments aided with

discussions with E.G., B.O., F.S. and G.F., while M.M. G.F. and D.R. performed them.

E.G., G.F. and B.O. designed, setup and maintained the custom two-photon microscope,

M.M. and D.R. set up the olfactometer. D.R. wrote all programs for data collection and

analysis. F.S. performed circuit modeling, M.M. and D.R. analyzed the data. M.M. and

D.R. generated the figures. All authors discussed the results. M.M. and D.R. wrote the

manuscript, and all authors participated in commentary and revision of the manuscript.

Declaration of interests: The authors declare no competing interests.

Methods

Animals

All animal procedures were performed in accordance with protocols approved by the

Institutional Animal Care and Use Committee of the University of Colorado Anschutz

Medical Campus. Mice were bred in the animal facility. We used both male and female

adult Parvalbumin-Cre (PV-Cre, Stock number 008069, Jackson Laboratory, USA) mice

and wild-type C57BL/6J mice. The animals were housed in a vivarium with a 14/10h

light/dark cycle. Food was available ad libitum. Access to water was restricted in for the

behavioral training sessions according to approved protocols, all mice were weighed daily

and received sufficient water during behavioral training to maintain >=80% of original

body weight.

Immunohistochemistry

To perform immunostaining, mice were sacrificed and transcardially perfused with ice cold

4% paraformaldehyde (Electron Microscopy Sciences, USA), followed by incubation in

30% sucrose (Sigma-Aldrich, USA). After the brain was incubated in the sucrose solution

60 μm thick slices were cut with a cryostat. The slices were imaged using a confocal laser

scanning microscope (Leica TCS SP5II, Germany or Nikon A1R, Japan) to determine the

GCaMP expression patterns in the cerebellum. The slices were counterstained with DAPI

(Thermo Fisher Scientific, USA).

Window implantation

Adult mice (8 weeks or older) were first exposed to isoflurane (2.5%) and then maintained

anesthetized by intraperitoneal ketamine-xylazine injection (100 μg/g and 20 μg/g). A

craniotomy was made over the vermis of cerebellum centered at midline 6.8 mm posterior

to Bregma leaving the dura intact (lobule VI). A square glass window (2 mm x 2 mm) of

No 1 cover glass (0.13 to 0.17 mm thick, Thermo Fisher Scientific, USA) was placed over

the craniotomy and the edges were sealed with cyanoacrylate glue (3M, USA). The window

was further secured with Metabond (Parkell, USA), and a custom-made steel head bracket

was glued to the skull.

Virus expression of GCaMP

In order to express GCaMP6f in cerebellar MLIs we injected the vermis of the cerebellum

in three adult PV-Cre mice with 2.0 μl of AAV1-Syn-Flex-GCaMP6f (Addgene, USA)

(6.8 mm posterior to Bregma, bilaterally ±0.5 mm lateral to midline and 200-400 μm below

the brain surface)[63]. The viral infection method we use has been reported to express

GCaMP in MLIs, and not in PCs[64,65]. Supplementary Fig. 2 shows that indeed expression

of GCaMP6 takes place only in MLIs. Note the absence of fluorescence from GCs as well

as from PC somata and dendrites. Expression in MLIs may reflect the differential activity

of the hSyn promoter, and has been described for a different genetically-encoded Ca^{2+}

indicator[65]. In one C57BL/6 animal each we used AAV5-Syn-GCaMP6s or AAVrg-Syn-

jGCaMP7f (Addgene, USA) with similar results to those obtained with GCaMP6f

(Supplementary Fig. 16a) and therefore the data were pooled. After the injection, the

animals were maintained for at least three weeks before behavior training and imaging.

Go-No Go training

Mice were water deprived by restricting daily consumption of water to 1-1.5 ml. Mice were

monitored for signs of dehydration or a decrease in body weight below 80% of the initial

weight. If either condition occurred the animals received water ad-lib until they recovered.

When the animals were thirsty, they were trained in a head-fixed olfactory go-no go task

with (1% Iso vs MO odorant application, Sigma-Aldrich, USA)[26,66]. Licks were monitored

by an electrical circuit monitoring the resistance between the lick spout and the floor in an

olfactometer that controlled valves to deliver a 1:40 dilution of odorant at a rate of 2 lt/min.

The water-deprived mice started the trial by licking on the water port. The odorant was

delivered after a random time interval ranging from 1 to 1.5 seconds. In S+ trials, the mice

needed to lick at least once in two 2 sec lick segments to obtain a reward (0.1 g/ml sucrose

water) (Fig. 1a). The inter trial interval (ITI) was 22.3-22.8 sec. In S- trials, the mice need

to refrain licking one of the two 2 sec segments to avoid a longer inter-trial interval (22.3-

22.8 + 10 sec). The animal's behavior performance was evaluated in a sliding window of

20 trials and the calculated value was assigned to the last trial in the window. Therefore, it

estimates the performance in the last 20 trials. The percent correct value represents the

percent of trials in which the animal successfully performed appropriate actions, and we

considered the animal proficient if percent correct performance is above 80%. In reverse

go-no go training sessions, the rewarded and un-rewarded odorants were switched.

Movement of the mouse was imaged in the infrared to prevent light interference with the

non-descanned detection in the visible using a 1 Megapixel NIR security camera (ELP-

USB100W05MT-DL36, Amazon.com, USA) at 30 frames per second. Velocity of body

movement was estimated using the Farneback algorithm coded in Matlab[67] (Mathworks,

USA). Measurement of body movement with a single camera gives limited information.

Five mice were used for the go-no go experiments. Four mice were imaged when they were

naïve. The window became opaque for two of the mice preventing MLI imaging for the

reversal experiment.

Go-Go training

For the experiment in Supplementary Figure 12 both odorants were rewarded with the same

volume of sugar water, and the volume of sugar water reward was varied. The two odorants

in this experiment were either 1% Iso and MO, or the same odorant 1% Iso and 1% Iso.

The results were similar for both odorant pairs, and the analysis was performed for all trials

regardless of odorant pair.

Behavioral performance recorded after chemogenetic inhibition of MLI activity

For chemogenetic inhibition of MLI activity[38,39], 1.2 ul of AAV8-hSyn-DIO-hM4D(Gi)-

mCherry virus (Addgene, USA) were bilaterally injected into 6 PV-Cre animals bilaterally

±0.5 mm lateral to midline, 6.8 mm posterior to Bregma and 200-400 μm below the brain

surface. For control an AAV8-hSyn-DIO-mCherry virus (Addgene, USA) was injected in

the same position in 6 PV-Cre animals. This viral infection method results in expression of

the protein in MLIs, and not in PCs[64,65]. We performed two separate experiments: 1) For

the experiments in Fig. 7 four weeks after injection the animals were trained to proficiency

in the go-no go task with 1% Iso (S+) vs MO (S-). When they reached a proficient level,

the animals rested for one to two weeks, and were then injected intraperitoneally (IP) with

saline 40 minutes before starting the session and were trained to discriminate between two

odorant mixtures: 0.1% of 60% heptanal+40% ethyl butyrate (S+) (Sigma-Aldrich, USA)

and 0.1% of 40% heptanal+60% ethyl butyrate (S-) for 3-4 sessions for a total of 400-500

trials. The animals then rested for one to two weeks and were subsequently trained to

discriminate the same odorant mixtures in sessions that took place 40 minutes after IP

injection of 3mg/kg clozapine-N-oxide (CNO, Tocris, USA). 2) For the experiment in

Supplementary Fig. 13 the animals were naïve for the discrimination of Iso vs. MO. The

animals were injected intraperitoneally (IP) with 3mg/kg clozapine-N-oxide (CNO, Tocris,

USA) 40 minutes before starting the session and were trained to discriminate between 1%

Iso (S+) and MO (S-) for 6 sessions for a total of 500-600 trials. The reinforcement was

then reversed and the animals were again injected with CNO 40 minutes before the sessions

and were trained to discriminate between MO (S+) and 1% Iso (S-) for 5 sessions for a

total of 400-500 trials.

Two photon imaging of MLI activity in animals undergoing the Go-No Go task

All the animals were first habituated to the setup to minimize stress during the imaging

experiments. All the imaging sessions started at least 10 minutes after mice had been head-

fixed. We searched for active MLIs while imaging zones in the vermis of lobule VI,

between the midline and the paravermal vein, an area of the cerebellum where GCs acquire

a predictive feedback signal or expectation reward[29,30]. The head fixed two photon imaging

system consisted of a movable objective microscope (MOM, Sutter Instrument Company,

USA) paired with a 80 MHz, ~100 femtosecond laser (Mai-Tai HP DeepSee, Spectra

Physics, USA) centered at 920 nm. The MOM was fitted with a single photon

epifluorescence eGFP filter path (475 nm excitation/500-550 nm emission) used for initial

field targeting followed by switching to the two photon laser scanning path for imaging

GCaMP at the depth of the MLIs. The galvometric laser scanning system was driven by

SlideBook 6.0 (Intelligent Imaging Innovations, USA). The two photon time lapses were

acquired at 256 x 256 pixels using a 1.0 NA/20x water emersion objective (Zeiss,

Germany) at 5.3 Hz. On the day of initial imaging, a FOV was selected to image a large

number of active cerebellar neurons located in the most superficial planes of the molecular

layer (within 50 μm beyond the dorsal surface of the molecular layer) including mostly

SCs, and several batches of 6000 frames ("time series") were collected in each training

session. After two photon imaging a second image of the vasculature was captured with

wide field epifluorescence to reconfirm the field.

Data analysis

Raw imaging data was first surveyed in ImageJ (NIH, USA) to exclude image sequences

exhibiting axial movement. We did not find evidence of axial movement while the animal

was engaged in the go-no go task. In addition, we performed control imaging where we

excited GCaMP6f at 820 nm, a two photon excitation wavelength where fluorescence

emission is Ca^{2+}-independent[68]. Supplementary Fig. 16b,c shows that we did not detect

transient changes in GCaMP6 fluorescence in a mouse engaged in the go-no go task when

the cells were excited at 820 nm. If we found horizontal drift due to motion, we applied

cross correlation-based image alignment using the Turboreg, Image J plugin. The data were

then analyzed with CaImAn Matlab software that uses constrained nonnegative matrix

factorization to define independent spatial and temporal components corresponding to

changes in GCaMP fluorescence in individual MLIs[31]. Baseline of intensity ($F0$) was

defined as the mean fluorescence intensity before trial start, defined as the time when

the animal first licked. This was when fluorescence started to increase above baseline

and the odorant was added at a random time 1-1.5 seconds after trial start. Intensity

traces (F) were normalized according to the formula $\Delta F/F =(F - F0)/F0$. After CaImAn

analysis, the $\Delta F/F$ traces of the spatial components were sorted and we assigned trial traces

to different behavioral events (Hit, CR, Miss and CR) and aligned them to trial start,

odorant onset or water delivery. Finally, the time course for the average $\Delta F/F$ did not differ

greatly between the different GCaMP variants as would be expected for a fast firing

interneuron with small increases in Ca^{2+} per action potential (Supplementary Fig. 16a).

Statistical analysis

Statistical analysis was performed in Matlab (Mathworks, USA). Statistical significance

for changes in measured parameters for factors such as learning and odorant identity (S+

vs. S-) was estimated using a generalized linear model (GLM), with post-hoc tests for all

data pairs corrected for multiple comparisons using false discovery rate[33]. The post hoc

comparisons between pairs of data were performed either with a t-test, or a ranksum test,

depending on the result of an Anderson-Darling test of normality. 95% CIs shown in the

figures as vertical black lines or shading bounding the lines were estimated by bootstrap

analysis of the mean by sampling with replacement 1000 times using the bootci function

in MATLAB.

Principal component analysis was calculated using the Matlab Statistics Toolbox.

Classification of trials using $\Delta F/F$ measured from all components in the FOV was

accomplished via linear discriminant analysis (LDA) in Matlab. We obtained similar result

with perceptron analysis (not shown). $\Delta F/F$ for all components for every trial except one

were used to train the LDA, and the missing trial was classified by its fit into the pre-

existing dataset. This was repeated for all trials and was performed separately for analysis

where the identity of the odorants was shuffled. Fluorescence intensity traces, lick rates

and kinematics in Fig. 6 were low-pass filtered with a hamming window of a time

constant of 0.59 s.

Dimensionality

Following Litwin-Kumar et al.[69] we defined the dimension of the system (dim) with M

inputs as the square of the sum of the eigenvalues of the covariance matrix of the measured

$\Delta F/F$ for all ROIs in the FOV divided by the sum of each eigenvalue squared:

634
$$\text{dim}\,(\Delta F/F) = \left(\sum_{i=1}^{M} \lambda_i\right)^2 \Big/ \left(\sum_{i=1}^{M} \lambda_i^{\,2}\right)$$

where λ_i are the eigenvalues of the covariance matrix of $\Delta F/F$ computed over the

distribution of $\Delta F/F$ signals measured in the FOV. If the components of $\Delta F/F$ are

independent and have the same variance, all the eigenvalues are equal and $\text{dim}(\Delta F/F) = M$.

Conversely, if the $\Delta F/F$ components are correlated so that the data points are distributed

equally in each dimension of an m-dimensional subspace of the full M-dimensional space,

only m eigenvalues will be nonzero and $\text{dim}(\Delta F/F) = m$.

Generalized linear modelling estimate of contribution of different variables to changes in ΔF/F

In order to quantify the contribution of different variables to neural activity, we used GLM as described by Engelhard et al. [37]. We used the Matlab fitglm function to fit the per trial ΔF/F time course for mice proficient in the go-no go task with a GLM. We included "event" variables, "whole trial" variables and "continuous" variables. Continuous variables quantified kinematics including lick rate, the derivative of lick rate and the velocity and acceleration of movements made by base of the tail of the head fixed animal during the trial (body velocity and body acceleration, see Supplementary Fig. 3). The identity of the odorant (S+/S- odorant) was an event variable that increased from zero to one during the time for S+ or S- odorant application. Finally, whole trial variables were accuracy (1 for correct responses and 0 for incorrect responses), reinforcement history (1 for reinforcement in the last trial, 0 otherwise) and percent correct behavior calculated in a window of 20 trials.

Data availability

The data supporting the findings of this study are available in GigaDB (http://gigadb.org/) (we are depositing the data in GigaDB, and this line will be replaced with a link to the data), the model is deposited in ModelDB (https://senselab.med.yale.edu/modeldb/)(we are depositing the data in ModelDB, and this line will be replaced with a link to the data) and the code is available in https://github.com/restrepd/CaImAnDR.

666   **Table 1. Percent of ROIs displaying a change in ΔF/F during odorant application**

Calculation	Percent responsive	Number	Number of ROIs
Percent of ROIs responding in Hit trials	99.9+0.23	21 time series, 5 mice	74-199
Percent of ROIs responding in CR trials	82.3+26	21 time series, 5 mice	74-199
Percent of ROIs responding in Miss trials	97.8+6.5	13 time series, 5 mice	105-199
Percent of ROIs responding in FA trials	78+29	15 time series, 5 mice	74-199
Percent of ROIs responding in Miss trials that also respond in Hit trials	99.96+0.15	13 time series, 5 mice	105-199
Percent of ROIs responding in FA trials that also respond in Hit trials	99.8+0.47	15 time series, 5 mice	74-199

668
669   This calculation was performed in time series that included at least one Miss or FA trial

670   when the animal was proficient (percent correct >=80%). The ROI was classified as

671   responsive when ΔF/F increased above or decreased below baseline ΔF/F by 2.5 x SD.

672   Baseline mean and SD were calculated for the time interval 10 to 2 sec before odorant

673   onset.

Figure 1. Two-photon Ca²⁺ imaging of MLIs in head-fixed mice undergoing the go-no go associative learning task.

a. Go-no go task. Left: Two-photon imaging of a head-fixed mouse responding to odorants

by licking on a water spout in response to the rewarded odorant in the go-no go task. Center:

Scoring of decision making. Right: time course of the trial. For water reward in Hit trials

the animal must lick at least once in each of the two 2 second lick segments.

b. The panels in **i** and **ii** show two photon microscopy images of GCaMP6f fluorescence

recorded from MLIs in a mouse proficient in the go-no go task (basal activity: before trial

start, Hit trial: during reinforced odorant application). The color image in **iii** shows the

regions of interest identified using CaImAn software[31].

c. Pseudocolor plots displaying the average per trial ΔF/F time course for Hit (**i**) and CR

(**ii**) odorants for all ROIs in this example. GLM analysis involving time periods pre-odorant

(1 sec before odorant onset), odorant (last second during odorant application) and

reinforcement (1.5 second after reinforcement) and different events (Hits, Miss, CR and

FA) yielded significant differences between reinforcement and pre-odorant ($p<0.001$),

between odorant and pre-odorant ($p<0.001$), and between all interactions between these

two period pairs and all events ($p<0.01$, 2040 observations, 2028 degrees of freedom,

n=170 ROIs, 1 mouse, GLM F-statistic 234, $p<0.001$). *Post-hoc ranksum $p<pFDR=0.048$.

d. Neural ensemble activity. Black traces are the GCaMP6f fluorescence ($\Delta F/F$) time

courses for a subset of the ROIs identified in the FOV in **b**. This mouse was proficient

(>=80% correct trials). Vertical lines: orange start and end of S+ odorant application, light

blue is S-. The blue trace at the bottom shows the licks. All trials were Hits or CRs with

the exception of the trial identified with the arrow that was a FA. Data shown in panels b-

d are from one session (one mouse).

Figure 2. MLIs developed divergent responses during go-no go learning.

a. Learning curve for a mouse discriminating 1% Iso from MO in the go-no go task. Magenta dots: proficient level (>=80% percent correct), green dots: naïve level (percent correct <=65%).

b. GCaMP6f fluorescence (ΔF/F) time course averaged over all ROIs and trials falling within different proficiency windows shown for the session whose performance is shown in **a**. **i.** <= 65% (naïve), **ii.** 65%-80% and **iii.** >= 80% (proficient). Orange: S+, light blue: S- (the shaded area is the 95% CIs). The vertical black lines are odorant onset and removal and the red lines bound the reinforcement period.

c-d. Per trial ΔF/F averaged over all ROIs for one second before odorant application (**c**,

Pre-odorant**)** and in the last second during odorant application (**d,** Odorant) shown for the

session in **a**. Vertical lines around the mean of each trial are the CI.

e. Violin plot showing per ROI ΔF/Fs for the session shown in **a** for the following time

windows: pre-odorant (one second before odorant application), odorant (the last second of

odorant application) and reinforcement (one and a half seconds after onset of

reinforcement). Per ROI ΔF/Fs are shown for trials falling within different proficiency

windows: <= 65% (naïve) and >= 80% (proficient). A GLM indicates that there are

significant differences between S+ (orange) and S- (light blue), between time windows and

between naïve and proficient mice (p<0.001, 5538 d.f., one mouse, *post-hoc ranksum/t

test p value < pFDR=0.043).

f-g. ΔF/F for four mice averaged over all the ROIs for all trials falling within different

proficiency windows. ΔF/F average was calculated for one second before odorant

application (**f**) and for the last second during odorant application (**g**). For the data during

odorant application (**g**) a generalized linear model (GLM) analysis indicated that there is a

statistical significance for the interaction between percent correct and the identity of the

odorant (p<0.001, 43 d.f., n=4 sessions, 4 mice, GLM F-statistic=60, p<0.001). GLM did

not yield statistically significant differences for pre-odorant data (**f**, p>0.05, 43 d.f., n=4

sessions, 4 mice, GLM F-statistic=0.46, p>0.05)**.** *Post-hoc t-test, p<pFDR=0.018.

Figure 3. The accuracy of decoding the stimulus increases as the animal learns to

discriminate the odorants.

Panels **a-c** show behavioral performance (**a**), PCA analysis (**b**) and LDA analysis (**c**) for

one session (one mouse). **a**. Learning curve for a mouse performing the head-fixed go-no

go task. Magenta dots: proficient level (>=80% percent correct), green dots: percent correct

<=65%.

b. First two principal components for the PCA for the changes in GCaMP6f fluorescence

(ΔF/F) in all the ROIs in the FOV. Top: the mouse was performing =< 65% correct, bottom:

the same mouse was performing >=80% correct. The principal components are shown for

three time periods: Pre-odorant: 1 sec before odorant addition, odorant: 1 sec before

removal of the odorant, and Reinforcement: 1.5 sec after reward. Orange circles: S+ trials,

light blue circles: S- trials.

c. Decoding accuracy for the linear discriminant analysis (LDA) trained to predict odorant

identity using $\Delta F/F$ for all ROIs in the FOV. Shade: 95% CI for LDA trained with shuffled

odorant identity. The vertical black lines are odorant onset and removal and the red lines

bound the reinforcement period.

d. LDA decoding accuracy calculated from the ($\Delta F/F$) for all ROIs in the FOV from data

recoded in four mice. A GLM analysis of decoding accuracy yielded statistically significant

interactions between (shuffled vs. odor) x (proficient vs. naïve) and for (shuffled vs.

reinforcement) x (proficient vs. naïve) (GLM p values <0.01 and <0.05, 24 observations,

18 d.f., n=4 sessions, 4 mice, GLM F-statistic=16.4, p<0.001). The vertical lines show 95%

CIs. *Post hoc t-tests corrected for multiple comparison using false discovery rate (FDR,

p<pFDR=0.007, n=4 sessions, 4 mice).

e. Example of LDA decoding analysis performed with subsets of ROIs ranging from 100

to 1 ROI. The analysis was performed for one session for trials where the mouse was

proficient (percent correct >=80%). **i.** Decoding accuracy (mean$\pm$ 95% CI) as a function of

the number of ROIs. **ii.** Cumulative probability histograms for the decoding accuracy for

the different number of ROIs and for the shuffled LDA. **iii.** Dimensionality as a function

of the number of ROIs.

f. Summary graph showing the decoding accuracy for different numbers of ROIs (mean$\pm$

95% CI, 4 sessions, 4 mice). A GLM analysis indicated that the decoding accuracy for the

shuffled analysis was statistically different from accuracy with the subsets of ROIs

($p<0.001$, 72 observations, 54 d.f., n=4 sessions, 4 mice, GLM F-statistic=23.6, $p<0.001$).

*Post-hoc t-tests or ranksum $p<pFDR=0.03$.

Figure 4. MLI odorant responses switched after reversal.

Panels **a-c and e** show behavioral performance (**a**), ΔF/F (**b** and **c**) and LDA analysis (**e**) for one session where odorant reinforcement was reversed (one mouse).

a. Learning curve showing the animal's behavior after reversal. Green <=65%, magenta: >=80% correct. Forward: S+ 1% Iso and S- MO.

b. Average ΔF/F time courses for example trials before (trials 51 and 52) and shortly after reversal (trials 91 and 92), and after the animal attained proficiency during reversal (trials 438 and 439). The vertical black lines are odorant onset and removal.

c. Per trial average ΔF/F during the last second of odorant application for the reversal task in Fig. 4a. The upper plot shows the responses to Iso and the lower plot shows responses to MO.

d. Odorant-induced changes in ΔF/F recorded from three animals computed for trials when

the animal was behaving >=80% correct. Vertical lines are 95% CIs. Red: Hit, Blue: CR.

A GLM analysis indicates that there is a significant difference in odor-induced changes in

ΔF/F for both odorant and reversal ($p < 0.001$, 12 observations, 8 d.f., n=3 sessions, 3 mice,

GLM F-statistic=14.8, $p < 0.01$).

e. Time course for decoding accuracy computed with LDA analysis of ΔF/F data for all

ROIs in the FOV. The shade is the 95% CI of decoding accuracy calculated with shuffled

odorant identity. The vertical black lines are odorant onset and removal and the red lines

bound the reinforcement period.

f. Decoding accuracy for three animals calculated with odorant-induced changes in ΔF/F

from trials where the animal was performing >=80% correct. Vertical lines are 95% CIs.

Dark-gray: original, light-gray: shuffled. A GLM analysis indicates that differences in

decoding accuracy are statistically significant between original and shuffled ($P < 0.001$), but

not between forward and reverse ($p > 0.05$, n=12, 8 d.f., n=3 sessions, 3 mice, GLM F-

statistic=16, $p < 0.001$).

Figure 5. Changes in lick rate correlate with changes in MLI activity during the

reinforcement period, but not during the odorant period.

Panels **a-d** are examples of lick traces (**a**), lick frequency (**b**), lick ranksum p value (**c**) and

ΔF/F (**d**) for one session (one mouse).

a. Examples of per trial lick traces when the mouse was learning to discriminate the

odorants when the animal was proficient (>=80% correct, orange traces: S+ trials, light

blue traces: S- trials).

b. Average lick frequency for S+ (orange) and S- (light blue) trials shown in **a**.

c. p value for a ranksum test estimating the difference in licks between the S+ and S-

odorants for the example in **a**.

d. Single trial examples of average ΔF/F (±CI, shade, n=105 ROIs) (**i-iv**) and the

corresponding lick frequency and lick traces (**v-viii**).

e. Relationship between the per trial average ΔF/F and the lick frequency shown per time

point for all trials in a go-no go session (one session, one mouse). Time points are

segregated within the last second before odorant addition (Pre-odorant), during the last

second of odorant addition (Odorant) and during the 1.5 seconds after reward

(Reinforcement). Correlation coefficients and p values for these time periods are: Pre-

odorant: 0.058, p<0.05. Odorant: 0.19, p<0.001. Reinforcement: 0.73, p<0.001.

*p<pFDR=0.05, n=6 sessions, 5 mice.

f. Relationship between the derivative of average ΔF/F and the derivative of lick frequency

shown per time point for all trials in a go-no go session (one session, one mouse). Time

points are segregated within the last second before odorant addition (Pre-odorant), during

the last second of odorant addition (Odorant) and during 1.5 seconds after reward

(Reinforcement). Correlation coefficients and p values for these time periods are: Pre-

odorant: -0.076, p<0.01. Odorant: -0.045, p>0.05. Reinforcement: 0.28, p<0.001.

*p<pFDR=0.016, n=6 sessions, 5 mice.

g. Correlation coefficients for the relationship between the average ΔF/F and lick frequency

for six sessions (five mice). The correlation coefficient is significantly different between

reinforcement and odorant (*t test p<pFDR=0.05, n=6 sessions, 5 mice).

h. Correlation coefficients for the relationship between the derivative of average ΔF/F and

the derivative of lick frequency for six sessions (five mice). The correlation coefficient is

significantly different between reinforcement and odorant (*t test p<pFDR=0.016, n=6

sessions, 5 mice).

In b, c and d the vertical black lines are odorant onset and removal and the vertical red lines

bound the reinforcement period.

All data shown in this figure are for proficient mice (percent correct>=80%).

Figure 6. Fit of MLI activity with a generalized linear model

a. Schematic illustration of the variables used by the GLM model to quantify the relationship between average ΔF/F and behavioral variables and odorant valence on a per trial basis.

b. Examples of GLM best-fit (black) and actual ΔF/F Ca^{2+} traces for Hit (**i,** red), Miss (**ii**, cyan), CR (**iii**, blue) and FA (**iv**, magenta) trials. The vertical black lines are odorant onset and removal and the red lines bound the reinforcement period.

c. Percent of the variance explained by GLM in the different time periods (Pre-odorant, Odorant and Outcome). The inset shows the time intervals used for the different periods.

d. Relative contribution of kinematics (**i**), licks (**ii**), odorants (**iii**) and reinforcement history

(**iv**) to the GLM fit for fits in the different time periods (pre-odorant, odorant, outcome).

The results showed that odorants and licks play an important role in contributing to the

activity during odorants and outcome periods respectively, while kinematics contribute to

the activity in all three time periods. Asterisks for Licks (**ii**) and Odorants (**iii**) denote

significant differences determined by ranksum and t test ($p < pFDR = 0.033$, n=6 sessions, 5

mice). There are no significant differences for Kinematics (**i**) or Reinforcement history (**iv**)

($p > pFDR = 0.017$, ranksum test).

Figure 7. Chemogenetic inhibition of MLI activity impairs associative learning.

a. and b. a,i and b,i. Examples of behavioral performance in a go-no go task where mice learned to differentiate between two odorant mixtures (S+: 0.1% of 60% heptanal+40% ethyl butyrate, S-: 0.1% of 40% heptanal+60% ethyl butyrate). Green: <=65% correct, magenta: >=80% correct. The vertical lines are the boundaries between different daily sessions.

a,ii and b,ii. Behavioral performance (mean± 95% CIs, n=6). **a.** Mice expressing mCherry in MLIs (control mice). **b.** Mice expressing hM4Di in MLIs. A GLM analysis indicates that there is a statistically significant interaction between CNO drug treatment and hM4Di expression (p<0.01, 24 observations, 20 d.f., n=6 mice, GLM F-statistic=13.7, p<0.001) and post hoc unpaired t tests indicate that the hM4Di expressing group differs between CNO and saline (asterisks, p<pFDR=0.025, n=12 mice).

c. Average lick frequency time course for the control (**i**) and hM4Di (**ii**) groups (mean± 95% CI, shade). The arrow in the inset shows that for S- trials of CNO treated hM4Di

animals (blue), the animal kept licking for a longer time than S- trials for mice injected

with saline (cyan). The vertical black lines are odorant onset and removal and the red lines

bound the reinforcement period.

d. Mean lick frequency ($\pm$CI) during the initial portion of the odorant application period

(0.8 to 1.8 sec) for mice performing >=80% correct (**i**: S+, **ii**: S-). GLM analysis did not

find a statistically significant difference for treatment or hM4Di expression (or

interactions) for S+ ($p>0.05$, 60 observations, 56 d.f., n=6 mice, GLM F-statistic=2.96,

$p>0.05$), but found a difference for CNO treatment x hM4Di expression for S- ($p<0.001$,

60 observations, 56 d.f., n=6 mice, GLM F-statistic=12.8, $p<0.001$). *$p<pFDR=0.03$ for

post-hoc t-test.

889
890

Cayco-Gajic, N. A. & Silver, R. A. Re-evaluating Circuit Mechanisms Underlying Pattern Separation. *Neuron* **101**, 584-602, doi:10.1016/j.neuron.2019.01.044 (2019).

Strick, P. L., Dum, R. P. & Fiez, J. A. Cerebellum and nonmotor function. *Annu Rev Neurosci* **32**, 413-434, doi:10.1146/annurev.neuro.31.060407.125606 (2009).

Sokolov, A. A., Miall, R. C. & Ivry, R. B. The Cerebellum: Adaptive Prediction for Movement and Cognition. *Trends Cogn Sci* **21**, 313-332, doi:10.1016/j.tics.2017.02.005 (2017).

Adamaszek, M. *et al.* Consensus Paper: Cerebellum and Emotion. *Cerebellum* **16**, 552-576, doi:10.1007/s12311-016-0815-8 (2017).

Badura, A. *et al.* Normal cognitive and social development require posterior cerebellar activity. *Elife* **7**, doi:10.7554/eLife.36401 (2018).

Jelitai, M., Puggioni, P., Ishikawa, T., Rinaldi, A. & Duguid, I. Dendritic excitation-inhibition balance shapes cerebellar output during motor behaviour. *Nat Commun* **7**, 13722, doi:10.1038/ncomms13722 (2016).

Gao, Z., van Beugen, B. J. & De Zeeuw, C. I. Distributed synergistic plasticity and cerebellar learning. *Nat Rev Neurosci* **13**, 619-635, doi:10.1038/nrn3312 (2012).

Marr, D. A theory of cerebellar cortex. *J Physiol* **202**, 437-470 (1969).

910   9   Ito, M. Neural design of the cerebellar motor control system. *Brain Res* **40**, 81-84,
911   doi:10.1016/0006-8993(72)90110-2 (1972).

912   10   Albus, J. S. A theory of cerebellar function. *Mathematical Biosciences* **10**, 25-61
913   (1971).

914   11   Heffley, W. & Hull, C. Classical conditioning drives learned reward prediction
915   signals in climbing fibers across the lateral cerebellum. *Elife* **8**,
916   doi:10.7554/eLife.46764 (2019).

917   12   Tsutsumi, S. *et al.* Modular organization of cerebellar climbing fiber inputs during
918   goal-directed behavior. *Elife* **8**, doi:10.7554/eLife.47021 (2019).

919   13   Deverett, B., Koay, S. A., Oostland, M. & Wang, S. S. Cerebellar involvement in
920   an evidence-accumulation decision-making task. *Elife* **7**, doi:10.7554/eLife.36781
921   (2018).

922   14   Carta, I., Chen, C. H., Schott, A. L., Dorizan, S. & Khodakhah, K. Cerebellar
923   modulation of the reward circuitry and social behavior. *Science* **363**,
924   doi:10.1126/science.aav0581 (2019).

925   15   Schonewille, M. *et al.* Reevaluating the role of LTD in cerebellar motor learning.
926   *Neuron* **70**, 43-50, doi:10.1016/j.neuron.2011.02.044 (2011).

927   16   Dean, P., Porrill, J., Ekerot, C. F. & Jorntell, H. The cerebellar microcircuit as an
928   adaptive filter: experimental and computational evidence. *Nat Rev Neurosci* **11**, 30-
929   43, doi:10.1038/nrn2756 (2010).

930   17   Rancillac, A. & Crepel, F. Synapses between parallel fibres and stellate cells

931   express long-term changes in synaptic efficacy in rat cerebellum. *J Physiol* **554**,

932   707-720, doi:10.1113/jphysiol.2003.055871 (2004).

933   18   Liu, S. Q. & Cull-Candy, S. G. Synaptic activity at calcium-permeable AMPA

934   receptors induces a switch in receptor subtype. *Nature* **405**, 454-458,

935   doi:10.1038/35013064 (2000).

936   19   Jorntell, H. & Ekerot, C. F. Receptive field plasticity profoundly alters the

937   cutaneous parallel fiber synaptic input to cerebellar interneurons in vivo. *J Neurosci*

938   **23**, 9620-9631 (2003).

939   20   Jorntell, H. & Ekerot, C. F. Reciprocal bidirectional plasticity of parallel fiber

940   receptive fields in cerebellar Purkinje cells and their afferent interneurons. *Neuron*

941   **34**, 797-806, doi:10.1016/s0896-6273(02)00713-4 (2002).

942   21   Jorntell, H., Bengtsson, F., Schonewille, M. & De Zeeuw, C. I. Cerebellar

943   molecular layer interneurons - computational properties and roles in learning.

944   *Trends Neurosci* **33**, 524-532, doi:10.1016/j.tins.2010.08.004 (2010).

945   22   Liu, Y. *et al.* A single fear-inducing stimulus induces a transcription-dependent

946   switch in synaptic AMPAR phenotype. *Nat Neurosci* **13**, 223-231,

947   doi:10.1038/nn.2474 (2010).

948   23   Wulff, P. *et al.* Synaptic inhibition of Purkinje cells mediates consolidation of

949   vestibulo-cerebellar motor learning. *Nat Neurosci* **12**, 1042-1049,

950   doi:10.1038/nn.2348 (2009).

24 ten Brinke, M. M. *et al.* Evolving Models of Pavlovian Conditioning: Cerebellar

Cortical Dynamics in Awake Behaving Mice. *Cell Rep* **13**, 1977-1988,

doi:10.1016/j.celrep.2015.10.057 (2015).

25 Rowan, M. J. M. *et al.* Graded Control of Climbing-Fiber-Mediated Plasticity and

Learning by Inhibition in the Cerebellum. *Neuron* **99**, 999-1015 e1016,

doi:10.1016/j.neuron.2018.07.024 (2018).

26 Gire, D. H., Whitesell, J. D., Doucette, W. & Restrepo, D. Information for decision-

making and stimulus identification is multiplexed in sensory cortex. *Nat Neurosci*

16, 991-993, doi:10.1038/nn.3432 (2013).

27 Doucette, W. & Restrepo, D. Profound context-dependent plasticity of mitral cell

responses in olfactory bulb. *PLoS Biol* **6**, e258, doi:08-PLBI-RA-1962 [pii]

10.1371/journal.pbio.0060258 (2008).

28 Helmchen, F. & Denk, W. Deep tissue two-photon microscopy. *Nature methods* **2**,

932-940, doi:10.1038/nmeth818 (2005).

29 Giovannucci, A. *et al.* Cerebellar granule cells acquire a widespread predictive

feedback signal during motor learning. *Nat Neurosci* **20**, 727-734,

doi:10.1038/nn.4531 (2017).

30 Wagner, M. J., Kim, T. H., Savall, J., Schnitzer, M. J. & Luo, L. Cerebellar granule

cells encode the expectation of reward. *Nature* **544**, 96-100,

doi:10.1038/nature21726 (2017).

31   Giovannucci, A. *et al.* CaImAn an open source tool for scalable calcium imaging

data analysis. *Elife* **8**, doi:10.7554/eLife.38173 (2019).

32   Chu, C. P., Bing, Y. H., Liu, H. & Qiu, D. L. Roles of molecular layer interneurons

in sensory information processing in mouse cerebellar cortex Crus II in vivo. *PLoS*

One **7**, e37031, doi:10.1371/journal.pone.0037031 (2012).

33   Curran-Everett, D. Multiple comparisons: philosophies and illustrations.

Am.J.Physiol Regul.Integr.Comp Physiol **279**, R1-R8 (2000).

34   Gaffield, M. A. & Christie, J. M. Movement Rate Is Encoded and Influenced by

Widespread, Coherent Activity of Cerebellar Molecular Layer Interneurons. *J*

Neurosci **37**, 4751-4765, doi:10.1523/JNEUROSCI.0534-17.2017 (2017).

35   Astorga, G. *et al.* Concerted Interneuron Activity in the Cerebellar Molecular Layer

During Rhythmic Oromotor Behaviors. *J Neurosci* **37**, 11455-11468,

doi:10.1523/JNEUROSCI.1091-17.2017 (2017).

36   Tye, K. M. Neural Circuit Motifs in Valence Processing. *Neuron* **100**, 436-452,

doi:10.1016/j.neuron.2018.10.001 (2018).

37   Engelhard, B. *et al.* Specialized coding of sensory, motor and cognitive variables

in VTA dopamine neurons. *Nature* **570**, 509-513, doi:10.1038/s41586-019-1261-9

(2019).

38   Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the

lock to fit the key to create a family of G protein-coupled receptors potently

activated by an inert ligand. *Proc Natl Acad Sci U S A* **104**, 5163-5168,
doi:10.1073/pnas.0700293104 (2007).

39   Gomez, J. L. *et al.* Chemogenetics revealed: DREADD occupancy and activation
via converted clozapine. *Science* **357**, 503-507, doi:10.1126/science.aan2475
(2017).

40   Raymond, J. L. & Medina, J. F. Computational Principles of Supervised Learning
in the Cerebellum. *Annu Rev Neurosci* **41**, 233-253, doi:10.1146/annurev-neuro-
080317-061948 (2018).

41   Davie, J. T., Clark, B. A. & Hausser, M. The origin of the complex spike in
cerebellar   Purkinje   cells.   *J   Neurosci*   **28**,   7599-7609,
doi:10.1523/JNEUROSCI.0559-08.2008 (2008).

42   Otsu, Y. *et al.* Activity-dependent gating of calcium spikes by A-type K+ channels
controls climbing fiber signaling in Purkinje cell dendrites. *Neuron* **84**, 137-151,
doi:10.1016/j.neuron.2014.08.035 (2014).

43   Rancz, E. A. & Hausser, M. Dendritic calcium spikes are tunable triggers of
cannabinoid release and short-term synaptic plasticity in cerebellar Purkinje
neurons. *J Neurosci* **26**, 5428-5437, doi:10.1523/JNEUROSCI.5284-05.2006
(2006).

44   Eccles, J. C., Llinas, R. & Sasaki, K. The excitatory synaptic action of climbing
fibres on the Purkinje cells of the cerebellum. *J Physiol* **182**, 268-296,
doi:10.1113/jphysiol.1966.sp007824 (1966).

45   Finch, E. A., Tanaka, K. & Augustine, G. J. Calcium as a trigger for cerebellar long-
1013   term synaptic depression. *Cerebellum* **11**, 706-717, doi:10.1007/s12311-011-0314-
1014   x (2012).

1015   46   Linden, D. J. & Connor, J. A. Long-term synaptic depression. *Annu Rev Neurosci*
1016   **18**, 319-357, doi:10.1146/annurev.ne.18.030195.001535 (1995).

1017   47   Brown, A. M. *et al.* Molecular layer interneurons shape the spike activity of
1018   cerebellar Purkinje cells. *Sci Rep* **9**, 1742, doi:10.1038/s41598-018-38264-1 (2019).

1019   48   Gaffield, M. A., Rowan, M. J. M., Amat, S. B., Hirai, H. & Christie, J. M. Inhibition
1020   gates supralinear Ca(2+) signaling in Purkinje cell dendrites during practiced
1021   movements. *Elife* **7**, doi:10.7554/eLife.36246 (2018).

1022   49   Jorntell, H. & Ekerot, C. F. Receptive Field Remodeling Induced by Skin
1023   Stimulation in Cerebellar Neurons in vivo. *Front Neural Circuits* **5**, 3,
1024   doi:10.3389/fncir.2011.00003 (2011).

1025   50   Lennon, W., Yamazaki, T. & Hecht-Nielsen, R. A Model of In vitro Plasticity at
1026   the Parallel Fiber-Molecular Layer Interneuron Synapses. *Front Comput Neurosci*
1027   **9**, 150, doi:10.3389/fncom.2015.00150 (2015).

1028   51   Szapiro, G. & Barbour, B. Multiple climbing fibers signal to molecular layer
1029   interneurons exclusively via glutamate spillover. *Nat Neurosci* **10**, 735-742,
1030   doi:10.1038/nn1907 (2007).

52   Arlt, C. & Hausser, M. Microcircuit Rules Governing Impact of Single Interneurons

on Purkinje Cell Output In Vivo. *Cell Rep* **30**, 3020-3035 e3023,

doi:10.1016/j.celrep.2020.02.009 (2020).

53   Bing, Y. H., Wu, M. C., Chu, C. P. & Qiu, D. L. Facial stimulation induces long-

term depression at cerebellar molecular layer interneuron-Purkinje cell synapses in

vivo in mice. *Front Cell Neurosci* **9**, 214, doi:10.3389/fncel.2015.00214 (2015).

54   Chabrol, F. P., Blot, A. & Mrsic-Flogel, T. D. Cerebellar Contribution to

Preparatory Activity in Motor Neocortex. *Neuron* **103**, 506-519 e504,

doi:10.1016/j.neuron.2019.05.022 (2019).

55   Hernandez-Briones, Z. S. *et al.* Olfactory stimulation induces cerebellar vermis

activation during sexual learning in male rats. *Neurobiol Learn Mem* **146**, 31-36,

doi:10.1016/j.nlm.2017.11.003 (2017).

56   Sobel, N. *et al.* Odorant-induced and sniff-induced activation in the cerebellum of

the human. *J Neurosci* **18**, 8990-9001 (1998).

57   Mainland, J. D., Johnson, B. N., Khan, R., Ivry, R. B. & Sobel, N. Olfactory

impairments in patients with unilateral cerebellar lesions are selective to inputs

from the contralesional nostril. *J Neurosci* **25**, 6362-6371,

doi:10.1523/JNEUROSCI.0920-05.2005 (2005).

58   Kostadinov, D., Beau, M., Blanco-Pozo, M. & Hausser, M. Predictive and reactive

reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells. *Nat*

Neurosci **22**, 950-962, doi:10.1038/s41593-019-0381-8 (2019).

1052 59 Heffley, W. *et al.* Coordinated cerebellar climbing fiber activity signals learned

1053   sensorimotor predictions. *Nat Neurosci* **21**, 1431-1441, doi:10.1038/s41593-018-

1054   0228-8 (2018).

1055 60 Vincent, P. & Marty, A. Fluctuations of inhibitory postsynaptic currents in Purkinje

1056   cells from rat cerebellar slices. *J Physiol* **494 (Pt 1)**, 183-199,

1057   doi:10.1113/jphysiol.1996.sp021484 (1996).

1058 61 Palay, S. L. & Chan-Palay, V. *Cerebellar Cortex: Cytology and Organization*.

1059   (Springer Berlin Heidelberg, 1973).

1060 62 Dean, H. L., Hagan, M. A. & Pesaran, B. Only coherent spiking in posterior parietal

1061   cortex coordinates looking and reaching. *Neuron* **73**, 829-841,

1062   doi:10.1016/j.neuron.2011.12.035 (2012).

1063 63 Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity.

1064   *Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).

1065 64 Astorga, G. *et al.* An excitatory GABA loop operating in vivo. *Front Cell Neurosci*

1066   **9**, 275, doi:10.3389/fncel.2015.00275 (2015).

1067 65 Kuhn, B., Ozden, I., Lampi, Y., Hasan, M. T. & Wang, S. S. An amplified promoter

1068   system for targeted expression of calcium indicator proteins in the cerebellar cortex.

1069   *Front Neural Circuits* **6**, 49, doi:10.3389/fncir.2012.00049 (2012).

1070 66 Li, A., Gire, D. H. & Restrepo, D. Y spike-field coherence in a population of

1071   olfactory bulb neurons differentiates between odors irrespective of associated

1072   outcome. *J Neurosci* **35**, 5808-5822, doi:10.1523/JNEUROSCI.4003-14.2015

1073   (2015).

1074   67   Farneback, G. Two-frame motion estimation based on polynomial expansion.

1075   *Proceedings of the 13th Scandinavian Conference on Image Analysis*, 363-370

1076   (2003).

1077   68   Barnett, L. M., Hughes, T. E. & Drobizhev, M. Deciphering the molecular

1078   mechanism responsible for GCaMP6m's Ca2+-dependent change in fluorescence.

1079   *PLoS One* **12**, e0170934, doi:10.1371/journal.pone.0170934 (2017).

1080   69   Litwin-Kumar, A., Harris, K. D., Axel, R., Sompolinsky, H. & Abbott, L. F.

1081   Optimal Degrees of Synaptic Connectivity. *Neuron* **93**, 1153-1164 e1157,

1082   doi:10.1016/j.neuron.2017.01.030 (2017).

1083