# Non-shared dispersal networks with heterogeneity promote species coexistence in hierarchical competitive communities

Helin Zhang[a], Jinbao Liao[a,*]

[a]*Ministry of Education's Key Laboratory of Poyang Lake Wetland and Watershed Research, School of Geography and Environment, Jiangxi Normal University, Ziyang Road 99, 330022 Nanchang, China*

**Corresponding author:** Prof. Dr. Jinbao Liao (jinbaoliao@163.com)

Address: Ziyang Road 99, 330022 Nanchang, Jiangxi Province, China.

Tel.: +86-(0)791-88133622          Fax: +86-(0)791-88120538

**Abstract**

The competition-colonization trade-off has been a classic paradigm to understand the maintenance of biodiversity in natural ecosystems. However, species-specific dispersal heterogeneities are not well integrated into our general understanding of how spatial coexistence emerges between competitors. Combining both network and metapopulation approaches, we construct a spatially explicit, patch-occupancy dynamic model for communities with hierarchically preemptive competition, to explore species coexistence in shared vs. non-shared dispersal networks with contrasting heterogeneities (including regular, random, exponential and scale-free networks). Our model shows that species with the same demography (i.e. identical colonization and extinction rates) cannot coexist stably in shared networks (i.e. the same dispersal pathways), regardless of dispersal heterogeneity. In contrast, increasing dispersal heterogeneity (even at very low levels of heterogeneity) in non-shared networks can greatly promote spatial coexistence, owing to the segregation-aggregation mechanism by which each species is restricted to self-organized clusters with a core of the most connected patches. However, these competitive patterns are largely mediated by species life-history attributes, for example, a unimodal biodiversity response to an increase of species dispersal rate emerges in non-shared heterogeneous networks, with species richness peaking at intermediate dispersal levels. Interestingly, increasing network size can foster species coexistence, leading to a monotonic increase in species-area curves. This strongly suggests that, unexpectedly, many more species can co-occur than the number of

limiting resources. Overall, this modelling study, filling the gap between network

structure and spatial competition, provides new insights into the coexistence

mechanisms of spatial heterogeneity.

*Keywords:* Dispersal heterogeneity, preemptive competition, competitive hierarchy,

spatial coexistence, network theory, segregation-aggregation mechanism.

**Introduction**

Exploring the mechanisms of community maintenance has been a central issue in

ecology. Many mechanisms have been proposed (e.g. niche and neutral theories), and

significant advances have been made in understanding species coexistence and

consequently biodiversity maintenance (Chesson 2000; Hubbell 2001; Levine &

HilleRisLambers 2009; Chu & Adler 2015). Among them, the

competition-colonization trade-off has been a classic paradigm to explain the

maintenance of biodiversity in natural ecosystems (Tilman 1994; Amarasekare 2000;

Yu & Wilson 2001; Yu et al. 2004). But if there is no any tradeoff between species

competition and demographic traits (colonization/extinction), how species can coexist

stably in competitive communities has become a long-standing challenge for

theoretical ecologists. Recently, non-hierarchical competition (i.e. competitive

intransitivity) among species has been proposed as a potential endogenous mechanism

for multispecies coexistence (Laird & Schamp 2006; Allesina & Levine 2011;

Soliveres et al. 2015; Levine et al. 2017). However, a key question remains unsolved

in hierarchical (transitive) competitive systems proposed by Tilman et al. (1994):

whether there exists any other factor fostering species coexistence in such system

without involving the colonization-competition trade-off. To probe the possible

underlying mechanism of such coexistence, we shift our focus to species dispersal in

spatially patchy environments. Most previous models of spatial coexistence assumed

that species dispersal is lattice- or randomly-structured in two-dimensional space,

ignoring the heterogeneity of dispersal networks (i.e. variation in patch connectivities)

across the landscape (but Hanski & Ovaskainen 2000).

In nature, evidence of heterogeneous dispersal networks abounds (Urban & Keitt

2001; Fortuna et al. 2006; Grilli et al. 2015). For example, Kininmonth et al. (2010)

found that the 321 reefs of the Great Barrier Reef constitute a scale-free small-world

dispersal network for the species, where most reefs have only one or a few links and a

very small proportion of reefs are extremely well-connected, following a power-law

degree distribution. Montoya et al. (2008) observed that trees with bird-dispersed

seeds perceive the landscape as a heterogeneous network, as opposed to trees with

wind-dispersed seeds which move through a homogeneous network. Fortuna et al.

(2006) identified a large spatial network of temporary ponds as breeding sites for

amphibian species, observing a shift from a power-law to a truncated power-law

distribution as the level of drought increases. As such, there has been an increasing

interest in characterizing the persistence and dynamics of interacting species with

dispersal heterogeneity using network theory (Urban & Keitt 2001; Bode et al. 2008;

Holland & Hastings 2008; Dale & Fortin 2010; Kininmonth et al. 2010; Gilarranz &

Bascompte 2012; Grilli et al. 2015; Gilarranz et al. 2017). In these representations,

each network is described as a graph consisting of a set of nodes (i.e. colony sites or

patches), and links between these nodes indicate dispersal pathways of individuals or

sub-populations (Fortuna et al. 2006, 2009). These studies found that spatial

heterogeneous networks greatly promote species persistence relative to the

homogeneous networks by increasing local recolonization opportunities,

demonstrating the importance of dispersal network structure for ecological dynamics

(e.g. Holland & Hastings 2008; Gilarranz & Bascompte 2012).

Despite these advances, species-specific dispersal network connectivities have

not been well integrated into our general understanding of how coexistence emerges

among species. Even though a few models have considered spatial dispersal with

different heterogeneities, they still assumed that all species have the same dispersal

pathways (i.e. shared dispersal networks; e.g. Holland & Hastings 2008). This

assumption neglects the fact that different species may perceive the landscape

differently and therefore display distinct dispersal pathways, shaping diverse patterns

of patch connectivity (Bunn et al. 2000; Nicholson & Possingham 2006; Fortuna et al.

2009; Hirt et al. 2018). Furthermore, some species can exhibit markedly different

dispersal patterns even though they disperse at similar times through similar

mechanisms, for example, animal species have distinct movement pathways due to

different feeding preference on different plant seeds (Nathan & Muller-Landau 2000;

Kinlan & Gaines 2003; Becker et al. 2007; Clobert et al. 2009). In fact, such

species-specific dispersal networks not only arise from niche differences (e.g. species

difference in habitat preference), but also result from different vectors for dispersal

among species (e.g. seed dispersal by animals, gravity or wind; Yeaton & Bond 1991;

Monotoya et al. 2008; Germain et al. 2019) or different dispersal traits (e.g. short- or

long-distance dispersal). Using the approach of allometric random walks to predict the

links among patches, Hirt et al. (2018) found that animal movement speed strongly

depends on body mass, and the degree of connectivity of a network follows an

allometric relationship, with medium-sized animals covering longer distances. In

addition, the locomotion mode (flying, running or swimming) can influence the

species-specific network connectivity, with flying animals being able to connect more

distant patches than running ones. Furthermore, Germain et al. (2019) applied satellite

imagery to link dispersal modes to movement of dispersal vectors, and observed that

hydrological networks, animal paths and distance can explain the richness of species

dispersed by gravity, animals and wind, respectively. Thus, there is an urgent need for

spatial coexistence theory to incorporate species-specific dispersal networks with

heterogeneity that are widespread in nature (Amarasekare 2008).

In this study, we construct a spatially explicit patch-dynamic model for multiple

species with hierarchically preemptive competition, in order to make a systematic

comparative analysis of species coexistence in shared *vs.* non-shared dispersal

networks with contrasting heterogeneities (defined as the extent of variation in patch

linking degrees). The shared dispersal networks refer to networks with the same

patches (or colony sites) and dispersal pathways, as opposed to the non-shared

dispersal networks where a system consists of two or more networks with the same

patches but different linking patterns (so-called multilayer networks; see Pilosof et al.

2017). In other words, the non-shared networks mean that each species has their own

linking architectures over the landscape, independent of each other. Note that, it is not

obligatory to require all species to have different characteristics of patch linking

degree distribution (i.e. different dispersal heterogeneities) in the non-shared dispersal

networks, i.e., species can have the same linking degree distribution (i.e. the same

heterogeneity), but their overall dispersal pathways across the landscape should not be

exactly the same (i.e. some links between patches are common for species). To

characterize heterogeneity in patch connectivities, four typical types of spatial

dispersal networks are considered (illustrated in Fig. 1): regular (Bascompte & Solé

1995), random (Erdös & Rényi 1959; Watts & Strogatz 1998), exponential and

scale-free networks (Barabasi & Albert 1999). Using this model, we systematically

explore: (i) whether and how competitors with dispersal heterogeneity can co-occur in

shared vs. non-shared networks when they have the same demography; and (ii) which

properties of dispersal network structure can best maintain species diversity.

**Methods**

*Dispersal networks with heterogeneity*

We represent the landscape as a graph (spatial network) consisting of a set of nodes

(patches or colony sites) connected by links. Each node denotes a patch linked with a

number of other patches (i.e. linking degree $k$), and links between patches represent

species dispersal pathways (i.e. functional connectivity among populations). As such,

each type of dispersal network can be characterized by its linking degree distribution.

Similar to Gilarranz & Bascompte (2012), four typical structures of dispersal

networks are considered (illustrated in Fig. 1a-d):

(i) A regular network where all patches have the same linking degree. For example,

Figure 1a shows a completely homogeneous network where each patch has four links

to other patches ($k$=4), equal to the lattice-structured model with nearest neighbour

dispersal under periodic boundary conditions (Bascompte & Sole 1995; Hiebeler 2000;

Liao et al. 2013a,b).

(ii) A randomly structured network with randomly connected patches (Watts &

Strogatz 1998). In particular, patch linking degrees follow a Poisson distribution with

the variance equal to the average degree per patch (e.g. $\bar{k}$=4 in Fig. 1b), thus there

exists a small variation in patch linking degrees (Erdös & Rényi 1959).

(iii) An exponential network constructed based on the generic algorithm of random

attachment, i.e. the network continuously expands by the addition of new patches that

are randomly connected to the patches already present in the system (see Barabási &

Albert 1999). Figure 1c displays an exponential network with the same average

linking degree $\bar{k}$=4 but more variability in linking degrees than the random network

(i.e. higher heterogeneity with variance approximately equal to 5.8 in our model

networks), following an exponential connectivity distribution (Fortuna et al. 2006).

(iv) A scale-free network constructed according to the algorithm of Barabási & Albert

(1999) with preferential attachment, leading to extremely high heterogeneity (with

variance approximately equal to 27.1 in our model networks). This is a consequence

of two generic mechanisms: (I) networks expand continuously by the addition of new

patches, and (II) new patches attach preferentially to the patches that are already

well-connected (Barabási & Albert 1999). Figure 1d exhibits such a network (keeping

average linking degree at $\bar{k}=4$), where most patches have only a few links, while a

few patches are extremely well-connected, following a power-law degree distribution

(Kininmonth et al. 2010).

In these networks, each patch is reachable for every species, that is, each patch has

at least one link to another patch. Species are assumed to disperse equally in all

directions with no preference, i.e. when patches $i_1$ and $i_2$ are linked, dispersal can

occur from either $i_1$ to $i_2$ or vice versa.

*Competitive dynamics*

We model a competition system structured in a large number of discrete habitat

patches connected by species dispersal. Each patch can be vacant or host a single

species. The system consists of $n$ species with competitive hierarchy (i.e. ranking of

species according to their competitive abilities), with the first species being the best

competitor and the $n$-th species the worst ($1>2>3>\cdots>n$). As colonizing a patch

already occupied by another species may be intrinsically more difficult than

colonizing an empty patch (e.g. plant propagule) because of the priority effect

(Comins & Noble 1985; Calcagno et al. 2006; Fukami 2015), we mainly consider

preemptive competition, i.e. species do not replace others but only compete for empty

patches, with strong competitors having priority to colonize these patches. Thus,

inferior species can colonize an empty patch only if superior species fail to establish

there. All species in the landscape are assumed to have the same colonization (with

rate $c$) and the same extinction (with rate $e$), in order to exclude any coexistence

caused by the colonization-competition tradeoff (Tilman 1994). As such, we can

unambiguously attribute any species coexistence to the explicit structural properties of

our dispersal networks. In each time step, each occupied patch becomes extinct with a

probability $e$ regardless of species identity. At the same time, whether an empty patch

is colonized depends on both species competitive hierarchy and abundances of its

directly connected occupied patches. Therefore, the probability that a given empty

patch $i$ is colonized by the $S$-th competitor ($1 \leq S \leq n$) should be

$$P_i(S) = (1 - c)^{\sum_{j=1}^{S-1} x_j} \cdot [1 - (1 - c)^{x_S}], \tag{1}$$

where $x_j$ ($\geq 0$) is the number of $j$-patches (occupied by species $j$) directly linked to the

empty patch $i$, and $(1 - c)^{\sum_{j=1}^{S-1} x_j}$ is the probability that patch $i$ is unoccupied by those

superior competitors (including species 1, 2, 3…$S$-1). Note that each empty patch is

colonized by one of its directly connected occupied patches with an independent

probability ($c$) regardless of species identity.

***Spatially explicit simulations***

Initially each patch is occupied by a species randomly sampled from the species pool.

For non-shared networks, we regenerate species-specific dispersal pathways

according to their network properties (e.g. variation in patch linking degrees) and

assign to each species randomly. Within each time step, we determine the local

extinction for each occupied patch with a given probability ($e$). Then we calculate the

probability that each empty patch becomes occupied by its directly connected species

according to their competitive hierarchy as well as patch occupancies (see Eq. 1).

Finally we record the patch occupancy for each species at each time step, calculated

as their number of occupied patches divided by the network size (i.e. the total number

of patches).

To determine the steady state, we preliminarily run the system for a long time,

finding that 5000 time steps are sufficient to achieve stability. We eventually run each

case until 10,000 time steps, and estimate patch occupancy for each species at steady

state by averaging their occupancies between $t$=9000~10,000 time steps to avoid

transient dynamics. Each case is explored with 100 replicates, starting from different

dispersal architectures in each replicate but with identical network properties (e.g. the

same network size, total links and their degree distribution). The mean of these 100

replicates ($\pm$ standard deviation SD) yields species abundance at steady state. A broad

range of biologically reasonable parameter combinations are explored and found to

yield qualitatively similar outcomes (Figs S1-S21 in *Appendix*), thus allowing us to

present our general results in Figs 1-5 by choosing one of those parameter

combinations as a reference (cf. Gilarranz & Bascompte 2012; Liao et al. 2020). To

reduce stochastic effects (Figs S1-S2 in *Appendix*), we model patch occupancy

dynamics (via Maltab R2018b) using large networks consisting of 1024 patches and

2048 undirected links (cf. Gilarranz & Bascompte 2012). As such, all types of

network have the same number of patches and links with the same average linking

degree $\bar{k}$=4, allowing us to compare species coexistence in dispersal networks with

contrasting heterogeneities. Note that, in our model, we mainly focus on preemptive

competition (i.e. species only compete for empty patches) without considering

competitive displacement (i.e. the superior competitor can invade into and displace

the inferior competitor; Tilman 1994), as we find that the assumption of competitive

displacement is too strong to keep species coexistence if all species have the same

demography (i.e. excluding the colonization-competition tradeoff) in hierarchical

competitive communities (e.g. the inferior species would be always outcompeted; see

Fig. S22 in *Appendix*).

**Results**

*Two-species system*

To get insights into the competitive dynamics, we first simply analyze two species (*A*

– inferior competitor and *B* – superior competitor) competing for an empty patch *i*

locally, thus the probability of the superior species *B* successfully colonizing the

empty patch should be

$$P_i(B) = 1 - (1 - c)^{x_B},\qquad(2)$$

with 0< *c* <1. But if the species *B* fails to establish in there, then the inferior species *A*

would have the opportunity to occupy this empty patch *i* with

$$P_i(A) = (1 - c)^{x_B} \cdot [1 - (1 - c)^{x_A}],\qquad(3)$$

where $x_A$ (or $x_B$) is the number of species *A* (or *B*) directly linked to the patch *i*.

Now we analyze if there exists the possibility that the inferior species $A$ has higher

opportunity to occupy the focal empty patch $i$ than the superior species $B$. By

setting $P_i(A) > P_i(B)$, we have

$(1 - c)^{x_B} \cdot [1 - (1 - c)^{x_A}] > 1 - (1 - c)^{x_B}.$         (4)

As such, the conditions for $P_i(A) > P_i(B)$ can be derived as

$\begin{cases} x_A > ln[2 - (1 - c)^{-x_B}]/\ln(1 - c) \\ \quad x_B < -\ln 2/\ln(1 - c) \end{cases}$ ,       (5)

otherwise $P_i(A) < P_i(B)$ (see phase diagram at Fig. S23 in *Appendix*). Thus, the

inferior competitor can have local superiority to colonize the vacant patches relative

to the superior competitor when the inferior species has more dispersal links to those

patches, indirectly demonstrating that both inferior and superior species might coexist

regionally in non-shared dispersal networks with heterogeneity.

We then numerically simulate the coexistence of two competitors with the same

demography (i.e. identical colonization and extinction rate) in shared vs. non-shared

dispersal networks with contrasting heterogeneities, including (from most

homogeneous to most heterogeneous) regular, random, exponential and scale-free

networks (Fig. 1). In general, both species cannot coexist regionally in the shared

networks regardless of dispersal heterogeneity, as the superior species eventually

excludes the inferior species (Fig. 1I-IV). In contrast, when their dispersal networks

are non-shared (but with the same heterogeneity; e.g. scale-free networks in Fig. S3),

they can co-occur stably (Fig. 1VI-VIII), except in the regular networks (analogous to

"shared" homogeneous networks) where the poor competitor is excluded (Fig. 1V; see

coexistence pattern in Fig. S4 in *Appendix*). Interestingly, increasing dispersal

heterogeneity in the non-shared networks decreases the amplitude of stochastic

fluctuations in patch dynamics and makes the species abundances converge (increased

abundance of the inferior species and vice versa).

The coexistence patterns described above can, however, be altered by varying the

species' relative extinction or/and colonization rates (Fig. 2; Figs S5-S13 in *Appendix*)

as well as the average linking degree (Fig. S14 in *Appendix*). On the one hand, both

species expectedly display a monotonic decline in global patch occupancy (mean $\pm$

SD of 100 replicates) as the relative extinction rate ($e/c$) increases, thus speeding up

species exclusion. This occurs irrespective of whether the dispersal networks are

shared or not. Yet, species coexistence in the non-shared networks with higher

heterogeneity can tolerate much higher $e/c$-ratios. On the other hand, increasing the

relative colonization rate ($c/e$; i.e. dispersal rate) always leads to higher abundance of

the superior species than the inferior species. In the shared networks, the inferior

species is always outcompeted regardless of heterogeneity. In the non-shared

networks but with the same heterogeneity, the inferior species remains longer in the

system with increasing values of $c/e$ when the network is heterogeneous. In this case,

the patch occupancy of the inferior species initially increases and later declines to

extinction, contrary to regular networks where the inferior species always goes extinct.

Intermediate levels of $c/e$ thus maximize the inferior species' abundance and

consequently promote species coexistence, as opposed to lower or higher dispersal

rate which would speed up species exclusion. This outcome is similar to the case

where the average linking degree is increased (Fig. S14 in *Appendix*). Furthermore,

we find that increasing network heterogeneity increases the parameter space ($c/e$) for

spatial coexistence.

We further explore species coexistence when both competitors (again with the same

demography) use dispersal networks with different heterogeneities (Fig. 3). If the

poor competitor has a regular dispersal network, the strong competitor with higher

dispersal heterogeneity (including random, exponential and scale-free) can

outcompete it (Fig. 3II-IV). In contrast, the inferior species with higher dispersal

heterogeneity surprisingly can exclude the superior species with regular dispersal. In

other words, dispersal heterogeneity can compensate for competitive disadvantage to

some degree, and overturn the competitive outcome (Fig. 3I, V & IX). In other cases

where both competitors exhibit dispersal networks with different heterogeneities

(except regular networks), we observe that both species can co-occur stably, and

higher dispersal heterogeneity decreases the fluctuation amplitude of patch occupancy

dynamics at steady state (Fig. 3VI-VIII vs. X-XII).

*Multispecies system*

Then we investigate how many competitors (with the same demography) can coexist

stably in multispecies communities by varying the relative extinction and/or

colonization rates, again considering shared vs. non-shared dispersal networks (Fig. 4;

Figs S19-S20 in *Appendix*). Generally, increasing the relative extinction rate ($e/c$)

reduces species richness in both shared and non-shared dispersal networks, but

biodiversity maintenance differs depending on dispersal heterogeneity (Fig. 4a & c).

When dispersal networks are shared, only the best competitor survives regardless of

dispersal heterogeneity. In non-shared networks (but with the same heterogeneity),

however, higher dispersal heterogeneity maintains more species (scale-free >

exponential > random > regular). When varying the relative colonization rate ($c/e$),

we find that again, only the best competitor can survive in shared networks regardless

of dispersal heterogeneity, whereas more species can coexist with increasing dispersal

heterogeneity in non-shared networks. Similar to increasing average linking degree

(Fig. S21 in *Appendix*), the biodiversity response to increasing $c/e$ is unimodal in the

non-shared heterogeneous networks: moderate levels of $c/e$ maximize species

diversity, while lower or higher dispersal rates result in more species extinctions.

Finally we examine the effect of network size on biodiversity maintenance (i.e. the

species-area curve), again in both shared and non-shared dispersal networks with

contrasting heterogeneities (Fig. 5). In shared networks, only the best competitor

survives regardless of network size and dispersal heterogeneity. In contrast, an

increasing network size in non-shared heterogeneous networks (except in the regular

network where only the best competitor survives) surprisingly leads to a monotonic

increase in species richness, with greater species richness at higher dispersal

heterogeneity.

**Discussion**

Our spatially explicit model focuses on how species-specific dispersal networks and

hierarchically preemptive competition interact in affecting species coexistence,

assuming that all species have the same demographic traits to exclude the

colonization-competition tradeoff. Yet, current theoretical understanding mostly

comes from models with very regular connections among patches, in contrast with

dispersal heterogeneity in natural systems which is far from regular (Hanski &

Ovaskainen 2000; Fortuna et al. 2006; McIntire et al. 2007). In our study, we thus

concentrate on dispersal heterogeneity, specifically in network structure (including

regular, random, exponential and scale-free), to demonstrate the importance of

dispersal heterogeneity for spatial coexistence. We indeed observe that, regardless of

dispersal heterogeneity, species cannot coexist stably in shared networks if there is no

colonization-competition tradeoff (Figs 1, 2 & 4). Sharing the same dispersal

pathways implies that competitors encounter each other very frequently, so the best

competitor would eventually drive all other species to extinction by quickly seizing

the empty patches. Thus, previous patch-dynamic models focusing only on shared

regular or random networks, might have largely underestimated species diversity, as

species in natural communities always exhibit diverse dispersal patterns with more or

less heterogeneity.

In contrast to shared dispersal networks, non-shared heterogeneous networks (even

at low levels of heterogeneity) greatly promote spatial coexistence and therefore

biodiversity maintenance, especially at higher dispersal heterogeneity (Figs 1, 2 & 4).

We further explore the mechanism underlying these competitive outcomes by

analyzing the spatial distribution for each species subject to its specific dispersal

network (Fig. S4 in *Appendix*), and relating a patch's incidence (i.e. the proportion of

time steps that the node is occupied along the dynamics) with its linking degree as

well as with the average linking degree of the patches it interacts with (Figs S15-S18

in *Appendix*). As observed in Fig. S4, species-specific dispersal networks with

heterogeneity allow species to locally form many self-organized clusters of occupied

patches, with the most connected patches as the core. This can be ascribed to

variability in linking degree across patches, which results in variability across patches'

incidence (Figs S15-S16 in *Appendix*). Obviously, patch incidence grows non-linearly

with linking degree (Eq. 1), that is, patches require a minimum linking degree to stay

occupied in the majority of time steps, above which patch incidence saturates. In turn,

the highly connected patches for a focal species can provide benefit for their directly

linked patches in terms of incidence (Figs S17 & S18 in *Appendix*). For example,

when comparing patches with the same linking degree, those attached to the more

well-connected patches have a higher incidence (positive feedback). Specifically,

conspecifics tend to develop into clusters segregated from other species in space

because of dispersal heterogeneity, which increases the frequency of neighbourhood

dispersal within conspecifics and decreases competition among heterospecifics,

thereby allowing demography-equivalent species to co-occur regionally in spite of

preemptive competition (so-called the segregation-aggregation mechanism; Pacala

1997; Murrell et al. 2001; Holyoak & Loreau 2006). This mechanism can also be

thought of as generating a type of spatial refugia for the poor competitors, i.e.

locations within the clusters favoring persistence of the focal species after the

extinction in surrounding areas. Alternatively, spatially different dispersal network

connectivities can result in local advantages for each species in terms of patch

re-colonization, for example, the inferior species within a local area containing highly

connected patches, can have higher effective colonization rate due to higher number

of links for the core patches. Essentially, such fundamental mechanism that promotes

regional coexistence belongs to spatial heterogeneity in population growth rate.

Therefore, the non-shared heterogeneous dispersal networks offer a new way that the

classic coexistence mechanism of spatial heterogeneity can occur.

When two competitors differ in dispersal heterogeneity, they are able to coexist

stably even though the poor competitor displays lower dispersal heterogeneity than

the strong competitor (Fig. 3), again confirming that dispersal heterogeneity can

weaken interspecific competition and therefore promote species coexistence.

Interestingly, this prediction was observed empirically by Yeaton & Bond (1991),

where two competing shrub species with dispersal differences (one with ant-dispersed

seeds and another with wind-dispersed seeds) can co-occur stably. Furthermore, a

poor competitor with higher dispersal heterogeneity can even exclude a strong

competitor with regular (homogeneous) dispersal, thus altering the species'

competitive rankings. Essentially, relative to the homogeneous dispersal of the

superior competitor, even very low dispersal heterogeneity (e.g. random dispersal) can

greatly promote local recolonization opportunities of the poor competitor, resulting in

a large numerical advantage that overwhelms the competitive superiority and

therefore excludes the strong competitor. This indicates that effects of spatial dispersal

heterogeneity far outweigh the effect of preemptive competition, further confirming

its important role in maintaining species diversity. However, these patterns of regional

coexistence are greatly mediated by species life-history traits. Interestingly, increasing

species dispersal (rates or links) prompts a unimodal biodiversity response in

non-shared heterogeneous networks, demonstrating that effects of species dispersal on

multispecies coexistence can be positive as well as negative (Figs 2 & 4 and Fig. S21

in *Appendix*). It is intuitive that a species is unable to persist locally at very low levels

of dispersal rate. Yet, too much dispersal promotes patch colonization opportunities

for the superior competitors, and consequently leads to the region-wide exclusion of

the poor competitors, similar to increasing average patch linking degree (Figs S14 &

S21 in *Appendix*).

A final observation is that increasing network size monotonically increases species

richness in non-shared heterogeneous networks, with stronger species persistence at

higher dispersal heterogeneity (Fig. 5). Essentially in the non-shared networks with

heterogeneity, there is a much higher chance that species have exclusive access to

specific patches whenever the landscape is composed of a very large number of

patches. The resulting monotonically increasing species-area curves refute the

previous view that the number of species coexisting cannot exceed the number of

limited factors (Levin 1970; Tilman 1982). Instead, we theoretically demonstrate that,

when there is species-specific dispersal heterogeneity, many more species than the

number of limiting resources should be able to coexist, as empirically observed in

several natural systems (Tilman 1982; Kotler & Brown 1988; Wellborn et al. 1996).

Previously, coexistence of an unlimited number of species in a spatial context was

ascribed to the colonization-competition tradeoff (Tilman 1994) rather than to spatial

heterogeneity (Adler & Mosquera 2000). Our model provides an alternative

explanation, if the landscape is large enough, for non-shared dispersal heterogeneities

supporting the coexistence of many more species than expected, based on the

segregation-aggregation mechanism.

By demonstrating that the architecture of dispersal networks strongly governs

species coexistence, mediated by species life-history attributes, our work helps fill the

gap between network structure and spatial competition. We find that incorporating

species-specific dispersal heterogeneities into the traditional hierarchical competitive

systems can greatly promote regional coexistence owing to the formation of

self-organized clusters. This implies that traditional shared lattice- or

randomly-structured models might have severely underestimated biodiversity

maintenance. More importantly, the model suggests significant implications for

biodiversity conservation and management. For instance, as different species often

display diverse patterns of patch connectivity based on their dispersal traits, we

should first construct and analyze dispersal networks independently for multiple target

species, and then overlay or intersect the multiple networks to find locations that are

important for these species, so as to design multispecies conservation planning (e.g.

Bunn et al. 2000; Urban & Keitt 2001; Nicholson & Possingham 2006). However,

two caveats should be addressed when applying our model to terrestrial ecosystems.

Firstly, although there have been a large number of studies on scale-free graphs

(Barabasi & Albert 1999), actual patch mosaics seem to not quite fit the definitions of

such well-studied networks so that they tend to not include the extremely connected

patches that characterize scale-free networks (Urban et al. 2009). Secondly, it may be

inappropriate to apply a graph representation for some landscapes if habitat patches

are poorly resolved spatially (Urban & Keitt 2001). For example, habitat quality

varies continuously and subtly over the landscape, thus aggregating this variability

into discrete patches would be inappropriate (e.g. Liao et al. 2013b). Overall, by

integrating both network and metapopulation approaches, our modelling study

provides a new way to understand the coexistence mechanism of spatial dispersal

heterogeneity, thereby strengthening our comprehension of biodiversity maintenance

in hierarchical competitive communities.

**Acknowledgements**

This study was supported by the National Science Foundation of China (No.

31760172 & 31901175), the Thousand Young Talents Plan of China, the Key Joint

Youth Project of Jiangxi Province (No. 20192ACBL21029), the Jiangxi Provincial

Education Department (No. GJJ160274), and the Doctoral Scientific Research

Foundation of Jiangxi Normal University (No. 12017778).

**Author contributions**

J.L. conceived and wrote this manuscript, and H.Z. conducted simulations and

analyzed results.

**Competing interests:** The author declares no competing interests.

**Supplementary Material**

Appendix accompanying this manuscript is supplied.

**Data accessibility**

This is a theoretical modelling study which does not use data.

**References**

Adler, F.R. & Mosquera, J. 2000. Is space necessary? Interference competition and

limits to biodiversity. *Ecology* 81, 3226-3232.

Allesina, S. & Levine, J.M. 2011. A competitive network theory of species

diversity. *PNAS* 108, 5638-5642.

Amarasekare, P. 2000. Coexistence of competing parasitoids on a patchily distributed

host: local vs. spatial mechanisms. *Ecology* 81, 1286-1296.

Amarasekare, P. 2008. Spatial dynamics of food webs. *Annu. Rev. Ecol. Evol. Syst.* 39,

479-500.

Barab ási, A.L. & Albert, R. 1999. Emergence of scaling in random networks. *Science*

286, 509-512.

Bascompte, J. & Solé, R.V. 1995. Rethinking complexity: modelling spatio-temporal

dynamics in ecology. *Trends Ecol. Evol.* 10, 361-366.

Becker, B.J., Levin, L.A., Fodrie, F.J. & McMillan, P.A. 2007. Complex larval

connectivity patterns among marine invertebrate populations. *PNAS* 104,

3267-3272.

Bode, M., Burrage, K. & Possingham, H.P. 2008.Using complex network metrics to

predict the persistence of metapopulations with asymmetric connectivity patterns.

*Ecol. Model.* 214, 201-209.

Bunn, A.G., Urban, D.L. & Keitt, T.H. 2000. Landscape connectivity: a conservation

application of graph theory. *J. Environ. Manage.* 59, 265-278.

Calcagno, V., Mouquet, N., Jarne, P. & David, P. 2006. Coexistence in a

metacommunity: the competition-colonization trade-off is not dead. *Ecol. Lett.* 9,

897-907.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol.*

*Syst.* 31, 343-366.

Chu, C. & Adler, P.B. 2015. Large niche differences emerge at the recruitment stage

to stabilize grassland coexistence. *Ecol. Monogr.* 85, 373-392.

Clobert, J., Le Galliard, J.F., Cote, J., Meylan, S. & Massot, M. 2009. Informed

dispersal, heterogeneity in animal dispersal syndromes and the dynamics of

spatially structured populations. *Ecol. Lett.* 12, 197-209.

Codeco, C.T. & Grover, J.P. 2001. Competition along a spatial gradient of resource

supply: a microbial experimental model. *Am. Nat.* 157, 300-315.

Comins, H.N. & Noble, I.R. 1985. Dispersal, variability, and transient niches: species

coexistence in a uniformly variable environment. *Am. Nat.* 126, 706-723.

Dale, M. & Fortin, M.J. 2010. From graphs to spatial graphs. *Annu. Rev. Ecol. Evol.*

*Syst.* 41, 21-38.

Erdös, P. & Rényi, A. 1959. On random graphs I. *Publ. Math.* 6, 290-297.

Fortuna, M., Gómez-Rodríguez, C. & Bascompte, J. 2006. Spatial network structure

and amphibian persistence in stochastic environments. *Proc. R. Soc. B* 273,

1429-1434.

Fortuna, M., Albaladejo, R., Fernández, L., Aparicio, A. & Bascompte, J. 2009.

Networks of spatial genetic variation across species. *PNAS* 106, 19044-19049.

Fukami, T. 2015. Historical Contingency in Community Assembly: Integrating Niches,

Species Pools, and Priority Effects. *Ann. Rev. Ecol. Evol. S.* 46, 1, 1-23.

Germain, R.M., Jones, N.T. & Grainger, T.N. 2019. Cryptic dispersal networks shape

biodiversity in an invaded landscape. *Ecology* 100(8), e02738.

Gilarranz, L.J. & Bascompte, J. 2012. Spatial network structure and metapopulation

persistence. *J. Theor. Biol.* 297, 11-16.

Gilarranz, L.J., Rayfield, B., Liňán-Cembrano, G., Bascompte, J. & Gonzalez, A.

2017. Effects of network modularity on the spread of perturbation impact in

experimental metapopulations. *Science* 357, 199-201.

Grilli, J., Barabás, G. & Allesina, S. 2015. Metapopulation Persistence in Random

Fragmented Landscapes. *PLoS Comput. Biol.* 11, e1004251.

Hanski, I. & Ovaskainen, O. 2000. The metapopulation capacity of a fragmented

landscape. *Nature* 404, 755-758.

Hiebeler, D. 2000. Populations on fragmented landscapes with spatially structured

heterogeneities: landscape generation and local dispersal. *Ecology* 81, 1629-1641.

Hirt, M.R., Grimm, V., Li, Y., Rall, B.C., Rosenbaum, B. & Brose, U. 2018. Bridging

scales: allometric random walks link movement and biodiversity research. *Trends*

*Ecol. Evol.* 33(9), 701-712.

Holland, M.D. & Hastings, A. 2008. Strong effect of dispersal network structure on

ecological dynamics. *Nature* 456,792-794.

Holyoak, M. & Loreau, M. 2006. Reconciling empirical ecology with neutral

community models. *Ecology* 87, 1370-1377.

Hubbell, S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*.

Princeton University Press, Princeton, NJ.

Kerr, B., Riley, M.A., Feldman, M.W. & Bohannan, B.J.M. 2002. Local dispersal

promotes biodiversity in a real-life game of rock-paper-scissors. *Nature* 418,

171-174.

Kininmonth, S.J., De'ath, G. & Possingham, H.P. 2010. Graph theoretic topology of

the great but small barrier reef world. *Theor. Ecol.* 3, 75-88.

Kinlan, B.P. & Gaines, S.D. 2003. Propagule dispersal in marine and terrestrial

environments: a community perspective. *Ecology* 84, 2007-2020.

Kotler, B.P. & Brown, J.S. 1988. Environmental heterogeneity and the coexistence of

desert rodents. *Ann. Rev. Ecol. Syst.* 19, 281-307.

Laird, R.A. & Schamp, B.S. 2006. Competitive intransitivity promotes species

coexistence. *Am. Nat.* 168, 182-193.

Levin, S.A. 1970. Community equilibria and stability, and an extension of the

competitive exclusion principle. *Am. Nat.* 104, 413-423.

Levine, J.M. & HilleRisLambers, J. 2009. The importance of niches for the

maintenance of species diversity. *Nature* 461, 254-257.

Levine, J. M., Bascompte, J., Adler, P.B. & Allesina, S. 2017. Beyond pairwise

mechanisms of species coexistence in complex communities. *Nature* 546, 56-64.

Liao, J.B., Li, Z., Hiebeler, D.E., El-Bana, M., Deckmyn, G. & Nijs, I. 2013a.

Modelling plant population size and extinction thresholds from habitat loss and

habitat fragmentation: Effects of neighbouring competition and dispersal strategy.

*Ecol. Model.* 268, 9-17.

Liao, J.B., Li, Z., Hiebeler, D.E., Iwasa, Y., Bogaert, J. & Nijs, I. 2013b. Species

persistence in landscapes with spatial variation in habitat quality: A pair

approximation model. *J. Theor. Biol.* 335, 22-30.

Liao, L.M., Shen, Y. & Liao, J. 2020. Robustness of dispersal network structure to

patch loss. *Ecol. Model.* 424, e109036.

McIntire, E.J.B., Schultz, C.B. & Crone, E.E. 2007. Designing a network for butterfly

habitat restoration: where individuals, populations and landscapes interact. *J. Appl.*

*Ecol.* 44, 725-736.

Montoya, D., Zavala, M.A., Rodríguez, M.A. & Purves, D.W. 2008. Animal versus

wind dispersal and the robustness of tree species to deforestation. *Science* 320,

1502-1504.

Murrell, D.J., Purves, D.W. & Law, R. 2001. Uniting pattern and process in plant

ecology. *Trends Ecol. Evol.* 16, 529-530.

Nathan, R. & Muller-Landau, H.C. 2000. Spatial patterns of seed dispersal, their

determinants and consequences for recruitment. *Trends Ecol. Evol.* 15, 278-285.

Nicholson, E. & Possingham, H.P. 2006. Objectives for multiple-species conservation

planning. *Conserv. Biol.* 20, 871-881.

Pacala, W. 1997. Dynamics of plant communities. In: (Ed. Crawley, M.) *Plant*

*Ecology*. Blackwell Science Ltd., Oxford, pp. 532-555.

Pilosof, S., Porter, M.A., Pascual, M. & Kéfi, S. 2017. The multilayer nature of

ecological networks. *Nat. Ecol. Evol.* 1(4), 0101.

Soliveres, S. *et al.* 2015. Intransitive competition is widespread in plant communities

and maintains their species richness. *Ecol. Lett.* 18, 790-798.

Tilman, D. 1982. *Resource Competition and Community Structure*. Princeton

University Press, Princeton, NJ.

Tilman, D. 1994. Competition and biodiversity in spatially structured habitats.

*Ecology* 75, 2-16.

Urban, D. & Keitt, T. 2001. Landscape connectivity: a graph-theoretic perspective.

*Ecology* 85, 1205-1218.

Urban, D.L., Minor, E.S., Treml, E.A. & Schick, R.S. 2009. Graph models of habitat

mosaics. *Ecol. Lett.* 12(3), 260-273.

Watts, D.J. & Strogatz, S.H. 1998. Collective dynamics of 'small-world' networks.

*Nature* 393, 440-442.

Wellborn, G.A., Skelly, D.K. & Werner, E.E. 1996. Mechanisms creating community

structure across a freshwater habitat gradient. *Ann. Rev. Ecol. Syst.* 27, 337-363.

Yeaton, R.I. & Bond, W.J. 1991. Competition between Two Shrub Species: Dispersal

Differences and Fire Promote Coexistence. *Am. Nat.* 138(2), 328-341.

Yu, D. & Wilson, H.B. 2001. The competition–colonization trade-off is dead; long

live the competition–colonization tradeoff. *Am. Nat.* 158, 49-63.

Yu, D. et al. 2004. Experimental demonstration of species coexistence enabled by

dispersal limitation. *J. Anim. Ecol.* 73(6), 1102-1114.

**Figures**

**Figure 1.** Patch dynamics of two competing species in shared (I-IV) vs. non-shared

(V-VIII) dispersal networks, always containing 1024 patches and 2048 links with

average linking degree $\bar{k}$=4. Four typical networks from the most homogeneous to

the most heterogeneous are included: (a) regular, (b) random, (c) exponential and (d)

scale-free networks. For display purposes, these networks consist of only 36 patches

(nodes) with 72 links. In the shared networks, the dispersal pathways for both

competing species are the same, while they are different in the non-shared networks

but with the same heterogeneity. Parameter values are the same for both species:

colonization rate $c$=0.05 and extinction rate $e$=0.05.

**Figure 2.** Effects of relative extinction (*e*/*c* at fixed *c*=0.05) and colonization (*c*/*e* at

fixed *e*=0.05) rate on patch occupancy (mean ±standard deviation SD of 100

replicates) of both inferior and superior competitors at steady state in shared vs.

non-shared networks with different heterogeneities, including regular, random,

exponential and scale-free networks. These networks consist of 1024 patches and

2048 links, and non-shared networks indicate species-specific dispersal patterns for

both competitors but with the same heterogeneity.

**Figure 3.** Patch dynamics of both inferior and superior competitors with different

heterogeneous networks, consisting of 1024 patches and 2048 links. Four types of

dispersal networks are considered: regular, random, exponential and scale-free

networks. Parameter values for both species are the same: $c=e=0.05$.

**Figure 4.** Effects of relative extinction (*e*/*c* at fixed *c*=0.05) and colonization rate (*c*/*e*

at fixed *e*=0.05) on the number of coexisting species at steady state (mean ±SD of

100 replicates) in hierarchical competitive communities with preemptive competition,

where species compete only for empty patches. All species are assumed to have the

same colonization and extinction rates. Networks are either shared (graphs a & b) or

non-shared but with the same heterogeneity (graphs c & d), and include regular,

random, exponential and scale-free networks (again with 1024 patches and 2048

links).

**Figure 5.** Species-area relationship between network size (i.e. total number of patches)

and the number of coexisting species at steady state in hierarchical competitive

metacommunities with preemptive competition (mean $\pm$ SD of 100 replicates). To

exclude the colonization-competition tradeoff, all species are assumed to have the

same colonization and extinction rate ($c$=$e$=0.05) in (a) shared vs. (b) non-shared

networks (i.e. species-specific dispersal but with the same heterogeneity), with fixed

average patch linking degree ($\bar{k}$=4). Again, four types of dispersal networks with

contrasting heterogeneities are included: regular, random, exponential and scale-free

networks.