Non-shared dispersal networks with heterogeneity promote species

coexistence in hierarchical competitive metacommunities

Helin Zhang, Jinbao Liao[*]

Ministry of Education's Key Laboratory of Poyang Lake Wetland and Watershed

Research, School of Geography and Environment, Jiangxi Normal University, Ziyang

Road 99, 330022 Nanchang, China

*Corresponding author: Prof. Dr. Jinbao Liao (jinbaoliao@163.com)

Address: Ziyang Road 99, 330022 Nanchang, Jiangxi Province, China.

Tel.: +86-(0)791-88133622   Fax: +86-(0)791-88120538

Abstract

Exploring the mechanism of biodiversity maintenance has become a fundamental

issue in ecology. To date, many mechanisms have been proposed subject to the niche

and neutral theories or the colonization-competition tradeoff. Despite these advances,

species-specific dispersal heterogeneities are not well integrated into our general

understanding of how coexistence emerges between competitors. Combining both

network and metapopulation approaches, we construct a discrete, spatially explicit

patch-dynamic model for metacommunities with hierarchically preemptive

competition, to explore species coexistence in the shared vs. non-shared dispersal

networks with contrasting heterogeneities (including regular, random, exponential and

scale-free networks). Our model shows that, in spite of dispersal heterogeneity,

species with the same demography (i.e. identical colonization and extinction rates)

cannot coexist stably in the shared networks (i.e. the same dispersal pathways). In

contrast, increasing dispersal heterogeneity in the non-shared networks greatly

promotes regional coexistence, due to the segregation-aggregation mechanism by

which each species is restricted to the self-organized clusters with the core of the most

connected patches. However, these competitive patterns are largely mediated by

species life-history attributes, for example, a unimodal biodiversity response to an

increase of species dispersal emerges in the non-shared heterogeneous networks, with

species richness peaked at intermediate dispersal levels. Interestingly, increasing

network size can foster more species coexistence, leading to a monotonic increase in

species-area curves. This strongly suggests that unexpectedly, many more species can

co-occur regionally than the number of limiting resources. Overall, this modelling

study, filling the gap between network structure and spatial competition, provides new

insights into the spatial mechanisms of multispecies coexistence, theoretically

demonstrating the importance of species-specific dispersal heterogeneity for

biodiversity maintenance.

Keywords: Dispersal heterogeneity, preemptive competition, competitive hierarchy,

regional coexistence, network theory, segregation-aggregation mechanism.

Introduction

Understanding the mechanisms of how species can coexist in natural communities has

already become a central issue in ecology (Tilman 1982; Chesson 2000; Amarasekare

& Nisbet 2001; Kraft et al. 2015). To date, many mechanisms have been proposed

subject to the niche and neutral theories or the colonization-competition tradeoff, and

significant advances have been made in our understanding of species coexistence and

consequently biodiversity maintenance (Tilman 1994; Hubbell 2001; Yu & Wilson

2001; Levine & HilleRisLambers 2009; Chu & Adler 2015). Yet, these theories suffer

their own shortcomings. For instance, many field studies fail to detect the evidence

for the classic resource-based niche differences or the colonization-competition

tradeoff (Harrison et al. 1995; Amarasekare 2000; Levine & Rees 2002), while the

species equivalence assumption of neutral theory is hard to reconcile with nature

(Turnbull et al. 1999; Yu and Wilson 2001; Jakobsson and Eriksson 2003; Allesina &

Levine 2011). This suggests that there should be a fundamental disconnect between

empirical findings and modelling assumptions (Cadotte et al. 2006). To solve this

mismatch between theory and experiment, non-hierarchical competition (e.g.

intransitivity) among multiple species has been recently proposed to be a potential

mechanism for biodiversity maintenance (Huisman & Weissing 1999; Kerr et al. 2002;

Laird & Schamp 2006; Allesina & Levine 2011). However, there remains one

unsolved question: whether and how species in nature can coexist stably in a spatial

system with hierarchical competition even without involving the

colonization-competition trade-off?

To probe the possible underlying mechanism above, we should switch our focus

on species dispersal in spatial patchy environments. It is well known that most

previous models of spatial coexistence often assumed species dispersal interactions in

a regular or random landscape (i.e. dispersal networks are assumed to be

lattice-structured or randomly-structured), ignoring the more realistic dispersal

heterogeneity across the landscape. In nature, evidence of dispersal networks with

heterogeneity abounds (Fortuna et al. 2006; Grilli et al. 2015). For example,

Kininmonth et al. (2010) found that the GBR (Great Barrier Reef) including 321 reefs

is a scale-free small-world dispersal network for the species, i.e. most of reefs have

one or a few links, and a very small proportion of reefs are extremely well-connected,

following a power-law degree distribution. Furthermore, while trees with

wind-dispersed seeds perceive the landscape as a homogeneous network, trees with

bird-dispersed seeds move through a heterogeneous network (Montoya et al. 2008).

Recently, there has been an increasing interest in characterizing the persistence and

dynamics of interacting species with dispersal heterogeneity using network theory

(Urban & Keitt 2001; Bode et al. 2008; Holland & Hastings 2008; Dale & Fortin 2010;

Kininmonth et al. 2010; Gilarranz & Bascompte 2012; Grilli et al. 2015; Gilarranz et

al. 2017). In these representations, each network is described as a graph consisting of

a set of nodes (i.e. patches), and links between these nodes indicate dispersal

pathways of individuals or sub-populations (Fortuna et al. 2006, 2009). These studies

found that the spatial heterogeneous networks greatly promote species persistence

relative to the homogeneous networks, demonstrating the importance of dispersal

network structure for ecological dynamics (e.g. Holland & Hastings 2008; Gilarranz

& Bascompte 2012).

Despite these advances, species-specific dispersal heterogeneities are not well

integrated into our general understanding of how coexistence emerges between

species. Even though a few models considered spatial dispersal with different

heterogeneities, they still assumed that all species share the same dispersal networks

(i.e. with the same dispersal pathways; e.g. Holland & Hastings 2008). This

assumption greatly neglects the fact that different species may perceive the landscape

differently and therefore display distinct dispersal pathways (i.e. non-shared dispersal

networks due to difference in habitat preference or dispersal traits), shaping different

patterns of patch connectivity (Bunn et al. 2000; Nicholson & Possingham 2006;

Fortuna et al. 2009). For instance, Codeco & Grover (2001) have empirically shown

that spatial variation in the supply ratio of Carbon and Phosphorous can favor

different algal or bacterial species at different spatial locations, therefore avoiding

competitive exclusion via dispersal. In addition, some species often exhibit markedly

different dispersal patterns, even though they disperse at similar times through similar

mechanisms (Nathan & Muller-Landau 2000; Kinlan & Gaines 2003; Becker et al.

2007; Clobert et al. 2009). Thus, there is an urgent need for spatial coexistence theory

to incorporate the non-shared dispersal networks with heterogeneity that is

widespread in nature (Amarasekare 2008).

In this study, we construct a discrete, spatially explicit patch-dynamic model for

multiple species with hierarchically preemptive competition, in order to make a

systematic comparative analysis of species coexistence in shared *vs.* non-shared

dispersal networks with contrasting heterogeneities. Four typical types of spatial

dispersal networks are considered (illustrated in Fig. 1): regular (Bascompte & Solé

1995), random (Erdös & Rényi 1959; Watts & Strogatz 1998), exponential and

scale-free networks (Barabasi & Albert 1999). Using this model, we attempt to

explore: (i) whether and how competitors with dispersal heterogeneity can co-occur in

shared vs. non-shared networks without involving the colonization-competition

tradeoff; and (ii) under what properties of dispersal network structure can best

maintain species diversity.

Methods

Model description

We model a competition system structured in a large number of discrete habitat

patches connected by species dispersal. Each patch can be vacant or host a single

species. The system consists of n species with competitive hierarchy (i.e. ranking

species according to their competitive abilities), in which the first species is the best

competitor while the n-th species being the worst competitor ($1>2>3>\cdots>n$).

According to the more realistic situation that colonizing a patch already occupied by

another species may be intrinsically more difficult than colonizing an empty patch

(e.g. plant propagule) because of preemptive effect (Comins & Noble 1985; Calcagno

et al. 2006), we only consider the preemptive competition (no replacement among

species), i.e. species only compete for empty patches, with strong competitors having

priority to colonize empty patches. Thus, the inferior species can colonize the empty

patch only if those superior species fail to establish there. All species in the landscape

are assumed to have the same colonization (with rate c) and extinction rates (e), in

order to exclude the colonization-competition tradeoff. As such, we can

unambiguously attribute any species coexistence to our explicit structural properties

of dispersal networks. In each time step, each occupied patch becomes extinct with a

probability e regardless of species identity. At the same time, whether the empty patch

is colonized by its directly connected occupied patches depends on species

competitive hierarchy as well as species abundances (i.e. patch occupancy). Therefore,

the probability that a given empty patch i is colonized by the S-th competitor ($1 \leq S \leq n$)

should be

$$P_i(S) = 1 - (1-c)^{\sum_{j=1}^{S} x_j}, \tag{1}$$

where x_j (≥ 0) is the number of j-patches (occupied by species j) directly linked to the

empty patch i, and $(1-c)^{\sum_{j=1}^{S} x_j}$ is the probability of the patch i being unoccupied by

those superior competitors (including species 1,2,3…S). Note that each empty patch is

colonized by one of its directly connected occupied patches with an independent

probability (c) regardless of species identity.

Dispersal networks with heterogeneity

We represent the landscape as a graph (spatial network) consisting of a set of nodes

(patches) connected by links. The link between two nodes indicates the dispersal

potential or frequency between patches. Each node denotes a suitable patch linked

with a number of other patches (i.e. linking degree k), and links between patches

represent species dispersal pathways (i.e. functional connectivity among populations).

As such, each type of dispersal network can be characterized by its linking degree

distribution. Similar to Gilarranz & Bascompte (2012), four structures of dispersal

networks are considered (illustrated in Fig. 1a-d):

(i) A regular network with all patches having the same linking degree. For example,

Fig. 1a shows a completely homogeneous network where each patch owns four links

to other patches (k=4), equal to the lattice-structured model with nearest neighbour

dispersal (Bascompte & Sole 1995; Hiebeler 2000; Liao et al. 2013a,b).

(ii) A randomly structured network with randomly connected patches (Watts &

Strogatz 1998). In particular, patch linking degrees follow a Poisson distribution with

the variance equal to the average degree per patch (e.g. $\bar{k}$=4 in Fig. 1b), thus there

exists a small variation in patch linking degrees (Erdös & Rényi 1959).

(iii) An exponential network constructed based on the algorithm of Barabasi & Albert

(1999) with random attachment. For instance, Fig. 1c displays an exponential network

with the same average linking degree $\bar{k}$=4 but more variability in linking degrees

than the random network (i.e. higher heterogeneity; Fortuna et al. 2006).

(iv) A scale-free network constructed according to the algorithm by Barabasi & Albert

(1999) with preferential attachment. For example, Fig. 1d exhibits a scale-free

network (keeping average patch linking degree at $\bar{k}$=4) with extremely high

heterogeneity, i.e. most patches have a few links, while a few patches are extremely

well-connected, following a power-law degree distribution (Kininmonth et al. 2010).

In these networks, species are assumed to disperse equally in all directions with no

preference, i.e. when both patches i_1 and i_2 are linked, dispersal can occur from either

i_1 to i_2 or vice versa.

Spatially explicit simulations

Initially all patches are assumed to be fully occupied by randomly assigning one of

competitors to each patch. Within each time step, we determine local population

extinction for each occupied patch with a given probability (*e*). Then we calculate the

probability that each empty patch becomes occupied by its directly connected species

according to their competitive hierarchy as well as population abundances (see Eq. 1).

Finally we record patch occupancy for each species at each time step using their

number of occupied patches divided by the network size (i.e. the total number of

patches).

To find the steady state, we preliminarily run the system for a long time, and find

that 5000 time steps are sufficient to achieve system stability. Thus we run each case

until 10,000 time steps, and estimate patch occupancy for each species at steady state

by averaging their patch occupancies between $t=9000\sim10,000$ time steps to avoid

transient dynamics. Each case is explored with 100 replicates, starting from different

dispersal architectures in each replicate but with identical network properties (e.g. the

same network size, total links and their degree distribution). Ultimately, the mean of

these 100 replicates (mean $\pm$ standard deviation SD) yields species abundance at

steady state. A broad range of biologically reasonable parameter combinations are

explored and found to yield qualitatively similar outcomes (Figs S1-S21 in *Appendix*),

thus allowing us to present our general results in Figs 1-5 by choosing one of

parameter combinations as a reference case.

Results

Two-species system

We first explore the coexistence of two competitors with the same demography (i.e.

identical colonization and extinction rates) in shared vs. non-shared dispersal

networks with contrasting heterogeneities, including (from most homogeneous to

most heterogeneous) regular, random, exponential and scale-free networks (Fig. 1).

All types of network have the same number of 1024 patches and 2048 links with the

same average linking degree $\bar{k}$=4, which is large enough to reduce the role of

stochastic effects (see Figs S1-S2 in *Appendix*), thereby allowing us to compare

species coexistence in different heterogeneous networks. In general, both species

cannot coexist regionally in the shared networks in spite of dispersal heterogeneity,

eventually with the superior species excluding the inferior species (Fig. 1I-IV). In

contrast, when their dispersal networks are non-shared, i.e. both competitors have

species-specific dispersal pathways but with the same heterogeneity (e.g. scale-free

networks in Fig. S3), they can co-occur stably (Fig. 1VI-VIII), except in the regular

networks where the poor competitor is competitively excluded (Fig. 1V; see

coexistence pattern in Fig. S4 in *Appendix*). Meanwhile, increasing dispersal

heterogeneity in the non-shared networks interestingly decreases the amplitude of

stochastic fluctuations on patch dynamics and increases the inferior species

abundance.

   The coexistence patterns described above, however, can be altered by varying

species relative extinction or/and colonization rates (Fig. 2; Figs S5-S13 in *Appendix*)

as well as average patch linking degree (Fig. S14 in *Appendix*). On the one hand, both

species intuitively display a monotonic decline in global patch occupancy (mean $\pm$SD

of 100 replicates) as relative extinction rate (e/c) increases irrespective of whether the

heterogeneous networks are shared or not, therefore speeding up species exclusion.

Yet, species coexistence in the non-shared networks with higher heterogeneity can

tolerate much higher ratios of e/c. On the other hand, in the shared networks even

with high heterogeneity, increasing relative colonization rate (c/e; i.e. dispersal rate)

expectedly leads to an increase in the superior species abundance, while the inferior

species is always outcompeted. In the non-shared networks, increasing values of c/e

promotes the superior species occupancy, but the patch occupancy of the inferior

species initially increases but latter declines to extinction, except in the regular

networks where the inferior species always goes extinct. Interestingly, we find that

intermediate levels of c/e can maximize the inferior species abundance and therefore

promote species coexistence, while further decreasing or increasing species dispersal

would speed up species exclusion. This outcome is similar to the case of increasing

average linking degree (Fig. S14 in *Appendix*). Furthermore, we find that increasing

network heterogeneity increases the parameter space (c/e) for regional coexistence.

We further explore species coexistence when both competitors (again with the same

demography) display different heterogeneous networks (Fig. 3). If the poor

competitor shows the regular dispersal network, the strong competitor with higher

dispersal heterogeneity (including random, exponential and scale-free) can

outcompete the poor competitor (Fig. 3II-IV). In contrast, the inferior species with

higher heterogeneity surprisingly can exclude the superior species with the regular

dispersal network. This strongly indicates that dispersal heterogeneity can compensate

the competitive disadvantage, thereby overturning the competitive outcome (Fig. 3I,

V & IX). In other cases where both competitors choose different dispersal networks

with more or less heterogeneity (excluding regular networks), we observe that both

species can co-occur regionally, and higher dispersal heterogeneity decreases the

fluctuation amplitude of patch occupancy dynamics (Fig. 3VI-VIII vs. X-XII).

Multispecies system

Then we investigate how many competitors (with the same demography) can coexist

stably in multispecies metacommunities, again considering shared vs. non-shared

dispersal networks by varying relative extinction or/and colonization rates (Fig. 4;

Figs S19-S20 in *Appendix*). Generally, increasing relative extinction rate (e/c) reduces

species richness in both shared and non-shared dispersal networks, but there exist

differences in biodiversity maintenance among networks with different dispersal

heterogeneities (Fig. 4a & c). For example, if all species have the same dispersal

pathways, we surprisingly observe that more species can co-occur regionally in the

regular networks than in other heterogeneous networks but only at relatively low

values of e/c. In contrast, in the non-shared networks, higher dispersal heterogeneity

can result in more species persistence (i.e. scale-free > exponential > random >

regular). When varying relative colonization rate (c/e), we find that, in most cases,

only the best competitor can survive in the shared networks regardless of dispersal

heterogeneity, whereas more species can coexist with increasing dispersal

heterogeneity in the non-shared networks. Similar to the case of increasing average

linking degree (Fig. S21 in *Appendix*), an increase of c/e interestingly leads to a

unimodal biodiversity response: moderate levels of c/e can maximize species diversity,

while further increasing or decreasing ratios of c/e (i.e. dispersal rate) would result in

more species extinctions.

Finally we examine the effect of network size on biodiversity maintenance (i.e.

species-area curve) again in both shared and non-shared dispersal networks with

contrasting heterogeneities (Fig. 5). In the shared networks, only the best competitor

survives regardless of network size and dispersal heterogeneity. In contrast, increasing

network size in the non-shared heterogeneous networks (except in the regular network

where only the best competitor stays alive) can surprisingly lead to a monotonic

increase in species richness, with more heterogeneity resulting in higher species

diversity.

Discussion

Our spatially explicit model focuses on how species-specific dispersal networks and

hierarchically preemptive competition interact to affect species coexistence, by

assuming that all species have the same demographic traits to exclude the

colonization-competition tradeoff. Yet, most current theoretical understanding comes

from models with very regular connections among patches, by contrast with the

heterogeneity in natural systems that are far from regular (Hanski & Ovaskainen 2000;

Fortuna et al. 2006; McIntire et al. 2007). In our study, we thus concentrate on

dispersal heterogeneity, specifically in network structure (including regular, random,

exponential and scale-free), demonstrating the importance of spatial dispersal

heterogeneity for species coexistence in the non-shared networks. Since we observe

that in spite of dispersal heterogeneity, species cannot coexist stably in the shared

networks if there is no colonization-competition tradeoff (Figs 1, 2 & 4). Intuitively,

sharing the same dispersal pathways indicates that competitors encounter each other

very frequently, and the best competitor would eventually drive all other inferior

species to extinction by quickly seizing the empty patches. Thus, previous

patch-dynamic models only focusing on the shared regular networks, might have

largely underestimated species diversity, as species in natural communities always

exhibit diverse dispersal patterns with more or less heterogeneity.

In contrast to the shared dispersal networks, the non-shared heterogeneous

networks greatly promote species coexistence and therefore biodiversity maintenance,

more obviously at higher dispersal heterogeneity (Figs 1, 2 & 4). We further explore

the mechanism underlying the competitive outcome by analyzing spatial distribution

for each species subject to their specific dispersal networks (Fig. S4 in *Appendix*), and

relating a patch's incidence (i.e. the proportion of time steps that the node is occupied

along the dynamics) with its linking degree as well as with the average degree of the

patches it interacts with (Figs S15-S18 in *Appendix*). As observed in Fig. S4,

species-specific dispersal networks with heterogeneity allow species to shape many

clusters of occupied patches with the core of the most connected patches, locally

forming many self-organized small worlds. This should be ascribed to variability in

linking degree across patches, which results in variability across patches' incidence

(Figs S15-S16 in *Appendix*). Obviously, patch incidence grows non-linearly with

linking degree (Eq. 1), that is, patches require a minimum degree to stay occupied in

the majority of time steps, but patch incidence would saturate after a threshold in

degree. In turn, the highly connected patches for the focal species can provide benefit

for their directly linked patches in terms of incidence (Figs S17 & S18 in *Appendix*).

For example, when comparing patches with the same degree, those attached to the

more well-connected patches have a larger incidence (positive feedback). Essentially,

conspecifics tend to develop into clusters segregated from other species in space

because of dispersal heterogeneity, which increases the frequency of neighbourhood

dispersal within conspecifics but decreases competition among heterospecifics,

thereby allowing demography-equivalent species to co-occur regionally in spite of

preemptive competition (so-called the segregation-aggregation mechanism; Pacala

1997; Murrell et al. 2001; Holyoak & Loreau 2006). This mechanism can also be

thought of as generating a type of spatial refugia for the poor competitors, i.e.

locations within the clusters favoring persistence of the focal species after the

extinction in surrounding areas.

When both competitors exhibit different dispersal heterogeneities, they are able to

coexist stably, even though the poor competitor displays lower dispersal heterogeneity

than the strong competitor (Fig. 3), again demonstrating that dispersal heterogeneity

can weaken interspecific competition and therefore promote species coexistence.

Furthermore, the poor competitor with higher dispersal heterogeneity can exclude the

strong competitor of having the regular (homogeneous) dispersal, therefore altering

the species' competitive rankings. Logically, even very low dispersal heterogeneity

(e.g. random dispersal) can greatly promote population abundance for the poor

competitor relative to the homogeneous dispersal, and such a large numerical

advantage can overwhelm the competitive superiority, thereby excluding the strong

competitor. This indicates that effects of spatial dispersal heterogeneity far outweigh

the effect of preemptive competition, further confirming its important role in

maintaining species diversity. However, these patterns of regional coexistence are

greatly mediated by species life-history traits. Interestingly, a unimodal biodiversity

response to an increase of dispersal rate emerges in the non-shared heterogeneous

networks, showing diverse (positive as well as negative) effects of species dispersal

on multispecies coexistence (Figs 2 & 4). More specifically, intermediate dispersal

levels can maximize species richness, while further increasing or decreasing species

dispersal results in more species exclusion. Intuitively, species is unable to persist

locally at very low levels of dispersal rate. Yet, too much dispersal promotes patch

colonization opportunities for the superior competitors, and consequently leads to the

region-wide exclusion of the poor competitors under surplus dispersal, similar to the

situation of increasing average patch linking degree (Figs S14 & S21 in *Appendix*).

A final observation is that increasing network size leads to a monotonic increase in

species richness in the non-shared heterogeneous networks, with higher dispersal

heterogeneity promoting more species persistence (Fig. 5). The resulting

monotonically increased species-area curves refute the previous view that the number

of species coexisting cannot exceed the number of limited factors (Levin 1970;

Tilman 1982). Instead, we theoretically demonstrate that, when there is

species-specific dispersal heterogeneity, there should be many more species

coexisting regionally than the number of limiting resources, as empirically observed

in several natural systems (Tilman 1982; Kotler & Brown 1988; Wellborn et al. 1996).

Previously, coexistence of an unlimited number of species because of the

colonization-competition tradeoff (Tilman 1994) was ascribed to a stepwise

asymmetric model of competition rather than spatial heterogeneity (Adler &

Mosquera 2000). However, our model provides another potential explanation that if

the landscape is large enough, non-shared dispersal heterogeneities allow many more

species to co-occur regionally than expected due to the segregation-aggregation

mechanism.

Our work helps fill the gap between network structure and spatial competition,

demonstrating that the architecture of dispersal networks largely affects species

coexistence, greatly mediated by species life-history attributes. We find that

incorporating species-specific dispersal heterogeneities into the traditional

hierarchical competitive systems can greatly promote regional coexistence due to the

formation of self-organized clusters. This strongly implies that traditional shared

lattice- or randomly-structured models might have severely underestimated

biodiversity maintenance. More importantly, this model suggests important

implications for biodiversity conservation and management. For instance, as different

species often display diverse patterns of patch connectivity based on their dispersal

traits, we should first construct and analyze dispersal networks independently for

multiple target species, and then overlay or intersect the multiple networks to find

locations that are important for these species, so as to design multispecies

conservation planning (e.g. Bunn et al. 2000; Urban & Keitt 2001; Nicholson &

Possingham 2006). Overall, this modelling study, integrating both network and

metapopulation approaches, takes an important step toward understanding the

coexistence mechanism of multiple species with spatial dispersal heterogeneity,

thereby strengthening our comprehension of biodiversity maintenance in hierarchical

competitive metacommunities.

Acknowledgements

This study was supported by the National Science Foundation of China (No. 31760172 & 31901175), the Thousand Young Talents Plan of China, the Key Joint Youth Project of Jiangxi Province (No. 20192ACBL21029), the Jiangxi Provincial Education Department (No. GJJ160274), and the Doctoral Scientific Research Foundation of Jiangxi Normal University (No. 12017778).

Author contributions

J.L. conceived and wrote this manuscript, and H.Z. conducted simulations and analyzed results.

Competing interests: The author declares no competing interests.

Supplementary Material

Appendix accompanies this manuscript is also supplied.

Data accessibility

This is a theoretical modelling study and does not use data.

References

Adler, F.R. & Mosquera, J. 2000. Is space necessary? Interference competition and

limits to biodiversity. *Ecology* 81, 3226-3232.

Allesina, S. & Levine, J.M. 2011. A competitive network theory of species

diversity. *PNAS* 108, 5638-5642.

Amarasekare, P. 2000. Coexistence of competing parasitoids on a patchily distributed

host: local vs. spatial mechanisms. *Ecology* 81, 1286-1296.

Amarasekare, P. & Nisbet, R.M. 2001. Spatial heterogeneity, source-sink dynamics,

and the local coexistence of competing species. *Am. Nat.* 158, 572-584.

Amarasekare, P. 2008. Spatial dynamics of food webs. *Annu. Rev. Ecol. Evol. Syst.* 39,

479-500.

Barabasi, A.L. & Albert, R. 1999. Emergence of scaling in random networks. *Science*

286, 509-512.

Bascompte, J. & Solé, R.V. 1995. Rethinking complexity: modelling spatio-temporal

dynamics in ecology. *Trends Ecol. Evol.* 10, 361-366.

Becker, B.J., Levin, L.A., Fodrie, F.J. & McMillan, P.A. 2007. Complex larval

connectivity patterns among marine invertebrate populations. *PNAS* 104,

3267-3272.

Bode, M., Burrage, K. & Possingham, H.P. 2008.Using complex network metrics to

predict the persistence of metapopulations with asymmetric connectivity patterns.

Ecol. Model. 214, 201-209.

Bunn, A.G., Urban, D.L. & Keitt, T.H. 2000. Landscape connectivity: a conservation

application of graph theory. *J. Environ. Manage.* 59, 265-278.

Cadotte, M., Mai, D., Jantz, S., Collins, M., Keele, M. & Drake, J. 2006. On testing

the competition-colonization trade-off in a multispecies assemblage. *Am. Nat.* 168,

704-709.

Calcagno, V., Mouquet, N., Jarne, P. & David, P. 2006. Coexistence in a

metacommunity: the competition-colonization trade-off is not dead. *Ecol. Lett.* 9,

897-907.

Chesson, P. 2000. Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol.*

Syst. 31, 343-366.

Chu, C. & Adler, P.B. 2015. Large niche differences emerge at the recruitment stage

to stabilize grassland coexistence. *Ecol. Monogr.* 85, 373-392.

Clobert, J., Le Galliard, J.F., Cote, J., Meylan, S. & Massot, M. 2009. Informed

dispersal, heterogeneity in animal dispersal syndromes and the dynamics of

spatially structured populations. *Ecol. Lett.* 12, 197-209.

Codeco, C.T. & Grover, J.P. 2001. Competition along a spatial gradient of resource

supply: a microbial experimental model. *Am. Nat.* 157, 300-315.

Comins, H.N. & Noble, I.R. 1985. Dispersal, variability, and transient niches: species

coexistence in a uniformly variable environment. *Am. Nat.* 126, 706-723.

Dale, M. & Fortin, M.J. 2010. From graphs to spatial graphs. *Annu. Rev. Ecol. Evol.*

Syst. 41, 21-38.

Erdös, P. & Rényi, A. 1959. On random graphs I. *Publ. Math.* 6, 290-297.

Fortuna, M., Gómez-Rodríguez, C. & Bascompte, J. 2006. Spatial network structure

and amphibian persistence in stochastic environments. *Proc. R. Soc. B* 273,

1429-1434.

Fortuna, M., Albaladejo, R., Fernández, L., Aparicio, A. & Bascompte, J. 2009.

Networks of spatial genetic variation across species. *PNAS* 106, 19044-19049.

Gilarranz, L.J. & Bascompte, J. 2012. Spatial network structure and metapopulation

persistence. *J. Theor. Biol.* 297, 11-16.

Gilarranz, L.J., Rayfield, B., Liňán-Cembrano, G., Bascompte, J. & Gonzalez, A.

2017. Effects of network modularity on the spread of perturbation impact in

experimental metapopulations. *Science* 357, 199-201.

Grilli, J., Barabás, G. & Allesina, S. 2015. Metapopulation Persistence in Random

Fragmented Landscapes. *PLoS Comput. Biol.* 11, e1004251.

Hanski, I. & Ovaskainen, O. 2000. The metapopulation capacity of a fragmented

landscape. *Nature* 404, 755-758.

Harrison, S., Thomas, C.D. & Lewinsohn, T.M. 1995. Testing a metapopulation model

of coexistence in the insect community on ragwort (*Senecio jacobaea*). *Am. Nat.*

145, 546-562.

Hiebeler, D. 2000. Populations on fragmented landscapes with spatially structured

heterogeneities: landscape generation and local dispersal. *Ecology* 81, 1629-1641.

Holland, M.D. & Hastings, A. 2008. Strong effect of dispersal network structure on

ecological dynamics. *Nature* 456,792-794.

Holyoak, M. & Loreau, M. 2006. Reconciling empirical ecology with neutral

community models. *Ecology* 87, 1370-1377.

Hubbell, S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*.

Princeton University Press, Princeton, NJ.

Huisman, J. & Weissing, F.J. 1999. Biodiversity of plankton by species oscillations

and chaos. *Nature* 402, 407-410.

Jakobsson, A. & Eriksson, O. 2003. Trade-offs between dispersal and competitive

ability: a comparative study of wind-dispersed *Asteraceae forbs*. *Evol. Ecol.* 14,

233-246.

Kerr, B., Riley, M.A., Feldman, M.W. & Bohannan, B.J.M. 2002. Local dispersal

promotes biodiversity in a real-life game of rock-paper-scissors. *Nature* 418,

171-174.

Kininmonth, S.J., De'ath, G. & Possingham, H.P. 2010. Graph theoretic topology of

the great but small barrier reef world. *Theor. Ecol.* 3, 75-88.

Kinlan, B.P. & Gaines, S.D. 2003. Propagule dispersal in marine and terrestrial

environments: a community perspective. *Ecology* 84, 2007-2020.

Kotler, B.P. & Brown, J.S. 1988. Environmental heterogeneity and the coexistence of

desert rodents. *Ann. Rev. Ecol. Syst.* 19, 281-307.

Kraft, N.J.B., Godoy, O. & Levine, J.M. 2015. Plant functional traits and the

multidimensional nature of species coexistence. *PNAS* 112, 797-802.

Laird, R.A. & Schamp, B.S. 2006. Competitive intransitivity promotes species

coexistence. *Am. Nat.* 168, 182-193.

Levin, S.A. 1970. Community equilibria and stability, and an extension of the

competitive exclusion principle. *Am. Nat.* 104, 413-423.

Levine, J.M. & Rees, M. 2002. Coexistence and relative abundance in annual plant

assemblages: the roles of competition and colonization. *Am. Nat.* 160, 452-467.

Levine, J.M. & HilleRisLambers, J. 2009. The importance of niches for the

maintenance of species diversity. *Nature* 461, 254-257.

Liao, J.B., Li, Z., Hiebeler, D.E., El-Bana, M., Deckmyn, G. & Nijs, I. 2013a.

Modelling plant population size and extinction thresholds from habitat loss and

habitat fragmentation: Effects of neighbouring competition and dispersal strategy.

Ecol. Model. 268, 9-17.

Liao, J.B., Li, Z., Hiebeler, D.E., Iwasa, Y., Bogaert, J. & Nijs, I. 2013b. Species

persistence in landscapes with spatial variation in habitat quality: A pair

approximation model. *J. Theor. Biol.* 335, 22-30.

McIntire, E.J.B., Schultz, C.B. & Crone, E.E. 2007. Designing a network for butterfly

habitat restoration: where individuals, populations and landscapes interact. *J. Appl.*

Ecol. 44, 725-736.

Montoya, D., Zavala, M.A., Rodríguez, M.A. & Purves, D.W. 2008. Animal versus

wind dispersal and the robustness of tree species to deforestation. *Science* 320,

1502-1504.

Murrell, D.J., Purves, D.W. & Law, R. 2001. Uniting pattern and process in plant

ecology. *Trends Ecol. Evol.* 16, 529-530.

Nathan, R. & Muller-Landau, H.C. 2000. Spatial patterns of seed dispersal, their

determinants and consequences for recruitment. *Trends Ecol. Evol.* 15, 278-285.

Nicholson, E. & Possingham, H.P. 2006. Objectives for multiple-species conservation

planning. *Conserv. Biol.* 20, 871-881.

Pacala, W. 1997. Dynamics of plant communities. In: (Ed. Crawley, M.) *Plant*

Ecology. Blackwell Science Ltd., Oxford, pp. 532-555.

Tilman, D. 1982. *Resource Competition and Community Structure*. Princeton

University Press, Princeton, NJ.

Tilman, D. 1994. Competition and biodiversity in spatially structured habitats.

Ecology 75, 2-16.

Turnbull, L.A., Rees, M. & Crawley, M.J. 1999. Seed mass and the

competition-colonization trade-off: a sowing experiment. *J. Ecol.* 87, 899-912.

Urban, D. & Keitt T. 2001. Landscape connectivity: a graph-theoretic perspective.

Ecology 85, 1205-1218.

Watts, D.J. & Strogatz, S.H. 1998. Collective dynamics of 'small-world' networks.

Nature 393, 440-442.

Wellborn, G.A., Skelly, D.K. & Werner, E.E. 1996. Mechanisms creating community

structure across a freshwater habitat gradient. *Ann. Rev. Ecol. Syst.* 27, 337-363.

Yu, D. & Wilson, H.B. 2001. The competition–colonization trade-off is dead; long

live the competition–colonization tradeoff. *Am. Nat.* 158, 49-63.

Figures & Captions

Figure 1. Patch dynamics of two competing species in shared (I-IV) vs. non-shared

(V-VIII) dispersal networks (containing 1024 patches and 2048 links with average

linking degree $\bar{k}$=4). Four typical networks from the most homogeneous to the most

heterogeneous are included: (a) regular, (b) random, (c) exponential and (d) scale-free

networks. For display purposes, these networks only consist of 36 patches (nodes)

with 72 links. Here the shared networks indicate the same dispersal pathways for both

competing species; while the non-shared networks mean that both competitors show

different dispersal pathways but with the same heterogeneity. Parameter values are the

same for both species: colonization rate c=0.05 and extinction rate e=0.05.

Figure 2. Effects of relative extinction (e/c at fixed c=0.05) and colonization (c/e at

fixed e=0.05) rates on patch occupancy (mean ± standard deviation SD of 100

replicates) of both inferior and superior competitors at steady state in shared vs.

non-shared networks with different heterogeneities, including regular, random,

exponential and scale-free networks. These networks consist of 1024 patches and

2048 links, and non-shared networks indicate species-specific dispersal pathways for

both competitors but with the same network property.

Figure 3. Patch dynamics of both inferior and superior competitors with different

heterogeneous networks, consisting of 1024 patches and 2048 links. Four types of

dispersal networks are considered: regular, random, exponential and scale-free

networks. Parameter values for both species are the same: $c=e=0.05$.

Figure 4. Effects of relative extinction (*e*/*c* at fixed *c*=0.05) and colonization rates (*c*/*e*

at fixed *e*=0.05) on the number of coexisting species at steady state (mean ±SD of

100 replicates) in hierarchical competitive metacommunities, with preemptive

competition that species only compete for empty patches. Initially the competitive

system contain a sufficient number of species, and all species are assumed to have the

same colonization and extinction rates in shared (graphs a & b) vs. non-shared (graphs

c & d) networks, including regular, random, exponential and scale-free networks

(again with 1024 patches and 2048 links). Here non-shared networks mean that all

species have different dispersal pathways while keeping the same network property,

which is realized by regenerating the dispersal network with the same heterogeneity

and assigning them to each species.

Figure 5. Relationship between network size (i.e. total number of patches) and the

number of coexisting species at steady state (i.e. species-area curves) in hierarchical

competitive metacommunities with preemptive competition (mean $\pm$ SD of 100

replicates). To exclude the colonization-competition tradeoff, all species are assumed

to have the same colonization and extinction rates ($c=e=0.05$) in (a) shared vs. (b)

non-shared networks (i.e. species-specific dispersal but with the same network

property), by fixing the average patch linking degree ($\bar{k}=4$). Again four types of

dispersal networks with contrasting heterogeneities are included: regular, random,

exponential and scale-free networks.