Elaboration of host P2X$_7$R-P$_{38}$MAPK axis involvement against intracellular leishmania donovani by Spergulin-A: a comprehensive study

Niladri Mukherjee [a¥], Saswati Banerjee [a¥], Sk. Abdul Amin[b], Tarun Jha[b], Sriparna Datta[c], Krishna Das Saha[a*]

[a]Cancer Biology and Inflammatory Disorder Division, CSIR-Indian Institute of

Chemical Biology, 4 Raja S. C. Mullick Road, Kolkata-700032, India

[b]Natural Science Laboratory, Department of Pharmaceutical Technology, Division of

Medicinal & Pharmaceutical Chemistry, P. O. Box 17020, Jadavpur University,

Kolkata, India.

[c]Department of Chemical Technology, University of Calcutta, Kolkata 700009, India

Email address: krishna@iicb.res.in (Krishna Das Saha); [¥] Equal contribution, *Corresponding Author

Current drugs are inadequate for the treatment of visceral leishmaniasis an

immunosuppressive ailment caused by *Leishmania donovani*. Regrettably, there is no

plant-origin antileishmanial drug present. $P2X_7R$ is constitutively present on

macrophage surfaces and can be a putative therapeutic target in intra-macrophage

pathogens with function attributes towards inflammation, host cell apoptosis, altered

redox and phagolysosomal maturation by activating $p_{38}MAPK$. Here we demonstrated

that the initial interaction of Spergulin-A, a triterpenoid saponin with macrophages was

mediated through $P2X_7R$ involving the signaling cascade intermediates Ca^{++},

$P_{38}MAPK$, and NF-κβ. $P_{38}MAPK$ involvement shown to have specific and stern

importance in leishmanial killing with increased NF-κBp65. Phago-lysosomal

maturation by Spergulin-A also campaigns for another contribution of $P2X_7R$. In vivo

evaluation of the anti-leishmanial activity of Spergulin-A was monitored through

$P2X_7R$, $P_{38}MAPK$, and NF-κβ in murine spleen and bone-marrow macrophages and

advocated Spergulin-A of being a natural compound of leishmanicidal functions which

acted through the $P2X_7R$-$P_{38}MAPK$ axis.

The obligate intra-macrophage parasite *Leishmania donovani* causes the sternest form of leishmaniasis (visceral) and if left untreated can reach casualty as high as 100% within 2 years of infection in developing countries.[1] VL is widespread in tropical and temperate areas and is endemic in 100 countries with an approximate yearly occurrence of 500,000 cases, of which 70,000 people expire every year.[2] Miltefosine, Sodium antimony gluconate, Pentamidine, and Amphotericin B are applied as the choice of chemotherapeutics, however, reported occasions of toxicity or resistance.[3] *Leishmania* parasites have developed various successful strategies for the manipulation of parental activities of macrophage-like antigen presentation, nitric oxide intermediates, and oxygen radical generation and cytokine production.[4] Despite elaborate literature on leishmanial host-pathogen interaction, there is scope for an explicit revelation of compound mediated distortion of intracellular signaling cascades that can subvert parasite propagation and multiplication and can advise new therapeutic stratagems.

Botanicals or phytochemicals have proven their worth as potent therapeutics and are applied as medication against different and various types of human and animal ailments. However, unluckily VL remains an exception even after serious efforts from different groups of scientists and drug companies.[5] Reports of potent plant-derived anti-leishmanial agents are there in the literature,[6] however, mostly performed against the vector-borne promastigote form of the parasite.[7] Our previous effort demonstrated that triterpenoid saponin, Spergulin-A, isolated from a perennial shrub *Glinus oppositifolius* is a potent intracellular antileishmanial compound besides being an immune-modulator.[5] Scientific documentation in this particular area of interest also publicized that triterpenoid saponin of *Careya arborea* and some water-insoluble saponin acted as potent leishmanicidal agents.[8-10]

P2X$_7$R (a purinergic receptor) is constitutively present on macrophage surfaces with function attributes towards inflammation, apoptosis and intracellular pathogen control.[11] Modulation of macrophage biology in respect to P2X$_7$R expression alteration is monitored in

Leishmania amazonensis and other intracellular pathogens like *Toxoplasma*, *Brucella*

Cryptosporidium and *Microsporidia*.[12-16] $P2X_7R$ over-expression can be a putative

chemotherapeutic strategy in intra-macrophage pathogens. $P2X_7R$ expression is positively

correlated with inflammation, host cell apoptosis, generation of ROS and NO as well as

phagolysosomal maturation;[11] all are different strategies for intra-macrophage pathogen killing

and absence or underexpression of the $P2X_7R$ have been found with underachieved

intracellular pathogen control.[11] Existing literature suggests that activation of $P2X_7R$ can

trigger mitogen-activated protein kinases (MAPKs), including $P_{38}MAPKs$.[17]

Activation of $p_{38}MAPK$ in respect to the $P2X_7R$ over-expression leads to the generation

of ROS and NO through NADPH oxidase is also mentioned in the literature[18,19] and embrace

consequence in intra-MΦ pathogen control strategies which also comprises leishmanial

parasites. Regulation of iNOS expression in macrophages is managed by MAPK signaling

molecules with closely associated with NF- κB activation.[20] It was reported that $P_{38}MAPK$ can

phosphorylate Rab5 an important step in the endosome maturation process and critical in

intracellular leishmanial killing[21] and our previous finding denoted that $P_{38}MAPK$

phosphorylation was decisively responsible for maturation of phagolysosomal complexes and

subsequent leishmanicidal activity.[2] Importance of MAPKs in establishment and maintenance

of their infection also favoured by the fact that viscerotropic *L. donovani* parasites alleviate

MAPK phosphorylation and thus cut of the MAPK signaling cascades towards possible

leishmanicidal outcome in naive murine bone marrow-derived macrophages[22] with NF-kB

being the prime transcription factor in NO production[23] with evidence that $P_{38}MAPK$ activation

is crucial in *L. donovani* infection abolition.[24] Phagosome-lysosome fusion is also an important

innate-effector immune response of host macrophages the process of which is maintained by

sequential acquisition and removal of different proteins of hydrolases and peptidases in nature

and aided in degrading intracellular pathogens which includes leishmanial parasites.[25] The

phagolysosome fusion is also critically acquainted by enhanced $P2X_7R$ expression and

managed through bivalent calcium ion efflux and RhoA-Phospholipase D controlled pathway.[11]

Screening compounds toward puringeric receptor modifications have identified

structurally diversified compounds over two thousand and almost one-third of them are being

natural products having an affinity towards these receptors. As for example teniposide, a semi-

synthetic podophyllotoxin derivative can inhibit $P2X_7R$, whereas, agelasine and garcinolic

acid,[26] potentially activate the same. In recent years efforts are made towards the identification

of compounds that can modulate P2X receptors aiming towards novel chemotherapeutic

measurements towards diseases associated with inflammation.[27,28] Extracts of the traditional

Chinese medicinal plant, *Panax ginseng* can also activate $P2X_7$ [29] which further identified

ginsenosides, triterpenoid saponins found in the root extract of plants for $P2X_7R$ activation

properties.[30] Conversely, the *R. longifolia* extract and fractions found to have an inhibitory

effect on the $P2X_7R$ leading to their anti-inflammatory function.[31] Additionally, there are

ample reports of triterpenoid saponins interacting with different groups of MAPKs including

$P_{38}MAPK$ in conferring their cytotoxic or protective roles.[32]

To explore the molecular insights in depicting the intracellular leishmanicidal function

of the triterpenoid saponin Spergulin-A, we have emphasized on the involvement of $P2X_7R$

and downstream pathways involved. We checked the *in silico* binding affinity of Spergulin-A

with $P2X_7R$. $P2X_7R$ blockage having a hindrance towards the reduction of Spergulin-A

mediated leishmanial killing strongly advocates for its association and significance.

Involvement of $P_{38}MAPK$ shown to have specific and stern importance in leishmanial killing

as with increased NF-κBp65 mediated ROS and NO production reciprocal with parasite

reduction and this phenomenon also secondarily established by simultaneous expression

analysis of $P_{38}MAPK$ and Gp91-phox, NADPH oxidase family member. Phago-lysosomal

maturation promotion in mediating intracellular leishmanial killing by Spergulin-A also

campaigns for another contribution of P2X$_7$R as well as multipotent effectiveness of Spergulin-

A of being a natural product supposed to act like. Finally, the in vitro observation was clarified

in vivo for Spergulin-A of being a natural compound of leishmanicidal functions and acted

through the P2X$_7$R- P$_{38}$MAPK axis.

Results

Involvement of the P2X$_7$ receptor in mediating intracellular anti-leishmanial activity of

Spergulin-A. At the commencement of the evaluation of antileishmanial affectivity of

Spergulin-A against RAW 264.7 MΦ internalized parasites, increasing doses of Spergulin-A

(Fig. 1a) was applied for 24 h. Spergulin-A was found to be efficient against the intercellular

parasites in a dose-dependent manner; however above 30μg/mL of doses, the antileishmanial

activity reached a plateau. Therefore, we have selected the dose of 30μg/mL for a detailed

assessment of the intracellular anti-leishmanial activity of Spergulin-A. In our previous report,

we had also elaborated that this dose is also perfectly safe against the host RAW 264.7 MΦs

and that Spergulin-A was not effective against the acellular promastigote or intermediate

axenic amastigote forms of *L. Donovani* at diverse dose applied dose range.[5]

Use of Brilliant Blue G (BB-G) as a selective P2X$_7$R antagonist[33] was adequate in

literature against both in vivo[34] and in vitro systems[35]. Different IC50 values for different cell

lines had been reported in the literature and also against a related mouse macrophage cell line

(J774.G8) with applied dose range well within 10 μM,[35,36] however not conclusive against

RAW 264.7 MΦs. We evaluated the dose range for RAW 264.7 MΦ survival in accordance

with the view of the dose range application for its inhibitory purposes against P2X$_7$R (Fig. 1b).

The dose 10μM of BB-G was selected as P2X$_7$R antagonist in this study to be applied in vitro

against RAW 264.7 MΦs as about 92% of the exposed cells were alive after 24 h (Fig. 1b) and

the dose is well within the accepted P2X$_7$R antagonist range in literature.

Thereafter, the contribution of P2X$_7$R in intracellular *L. donovani* infection and

Spergulin-A mediated leishmanicidal function was thoroughly examined (Fig. 1c, d, e). Co-

expression of CFSE-tagged *L. donovani* and subsequent P2X$_7$R expression was monitored by

FACS during infection and Spergulin-A (30µg/mL) treatment, with or without BB-G, the

P2X$_7$R antagonist (Fig. 1c i-vi). P2X$_7$R expression was highest in the treated panel with

expected lowest CFSE-tagged *L. Donovani* existence and application of BB-G in treatment

panel lowered both the P2X$_7$R expression and *L. Donovani* removal. Spergulin-A treatment

exhibited a significant rise of P2X$_7$R expressing MΦs of about 44.3% of the parent population

in connection with intracellular parasite removal. Whereas, all other panels P2X$_7$R expressing

MΦs were about or below 5% of the parent population with or without BB-G application (Fig.

1c i-vi). And it was also noticed that P2X$_7$R antagonist BB-G application significantly lowered

Spergulin-A mediated intracellular *L. Donovani* removal (Fig. 1c iv-vi) and thus avow

important role of P2X$_7$R in Spergulin-A activated the intracellular leishmanicidal function.

The dose-dependent implication of Spergulin-A in intracellular leishmanial removal

and subsequent aversion by BB-G application which resulted in P2X$_7$R blockage was

monitored (Fig. 1d). In all the applied Spergulin-A doses, it was observed that BB-G

application lowered the leishmanicidal property of Spergulin-A significantly. Confocal

micrographs of CFSE-tagged intracellular *L. donovani* with subsequent P2X$_7$R expression were

monitored for Spergulin-A treatment as well as BB-G application (Fig. 1e). It was obvious

from the micrographs that Spergulin-A treatment was able to increase P2X$_7$R expression and

intracellular *L. Donovani* removal whereas the application of the P2X$_7$R antagonist resulted in

profuse existence of the parasite despite the Spergulin-A treatment (Fig. 1e).

This experimental evidences suggested that Spergulin-A mediated its intracellular

leishmanicidal action with close assistance with P2X$_7$R and subsequent signaling pathways

linked with over-expression of P2X$_7$R. Therefore, downstream signaling intermediates

associated with $P2X_7R$ should be closely monitored for accessing the precise involvement of

$P2X_7R$ in Spergulin-A mediated leishmanial killing.

Spergulin-A treatment has effects on the $P2X_7$ receptor-dependent intracellular Ca^{++}

release. It is documented that $P2X_7R$ activation can lead to Ca^{++} influx and resulted in

mycobacterial removal.[37] This Ca^{++} influx is definitely of great consequence in case of

intracellular leishmanial killing as this increased Ca^{++} titer is linked with phagolysosomal

fusion, $P_{38}MAPK$ activation and increased release of NO and ROS along with the altered

release of pro and anti-inflammatory cytokines; all of them have leishmanicidal function

actively or passively. Thus it is important to monitor the Ca^{++} influx during Spergulin-A

treatment and also link the event with $P2X_7R$ activation. A set of FACS analyses with $P2X_7R$

and Ca^{++} corresponding to Fura2M was expressed both graphically and through histograms of

fluorescence index and percentage of cells expressing the $P2X_7R$ and Ca^{++} for normal, *L.*

donovani parasitized, different doses of Spergulin-A treated MΦs both BB-G exposed and un-

exposed (Fig. 2a-c). Mean fluorescence index equivalent of the expressions of both $P2X_7R$ and

Ca^{++} was significantly higher in Spergulin-A treated panels and that too in a dose-dependent

manner compared to either normal or infected MΦs, and with the application of BB-G, both

these expressions got lowered compared to the corresponding BB-G unexposed panels (Fig.

2b). Subsequently, the percentage of $P2X_7R$ and Ca^{++} expressing cells against the parent

population also higher for the treated panels and the application of $P2X_7R$ agonist BB-G not

only lowered the $P2X_7R$ expressing population but also have diminishing the effect on Ca^{++}

expressing cells (Fig. 2c). These results strongly advocated the involvement and dependence

between $P2X_7R$ expression and Ca^{++} influx in Spergulin-A mediated anti-leishmanial function.

$P_{38}MAPK$ expression alteration affects Spergulin-A mediated anti-leishmanial activity

with $P2X_7$ receptor involvement. The involvement of $P_{38}MAPK$ was evaluated in *L.*

donovani infection headway and also the effect of Spergulin-A with the involvement of

P_{38}MAPK towards infection riddance as analyzed by flow cytometry (Fig. 3a). Apart from the

panels where P_{38}MAPK inhibitor (SB203580) was used (Fig. 3a iv-vi) almost all the cells were

expressing P_{38}MAPK either infected or treated with Spergulin-A (Fig. 3a i-iii). The difference

lies in the expression level of P_{38}MAPK. Compared to the control MΦ panel, infection with *L.*

donovani comprehensively reduced P_{38}MAPK (Fig. 3a ii and vii). However, Spergulin-A

treatment increased the P_{38}MAPK intensity (Fig. 3a iii and vii) and thus possibly has a very

precise correlation with the reduced intracellular parasitic count as when SB203580 was used

parasitic count also decreased along with the P_{38}MAPK expression (Fig. 3a v-vii).

207   Co-expression analyses of $P2X_7R$ and P_{38}MAPK were evaluated by confocal

microscopy (Fig. 3b). For the control MΦs, both the expressions of $P2X_7R$ and P_{38}MAPK were

at a basal level and for the infected and infected with BB-G their expression were down-

regulated more precisely P_{38}MAPK expression. Whereas, in the Spergulin-A treated panel

both $P2X_7R$ and P_{38}MAPK were visually overexpressed several folds (Fig. 3b) in accordance

with the anti-leishmanial effect demonstrated previously. However, the application of BB-G,

the $P2X_7R$ antagonist lowered the expression of P_{38}MAPK. Inhibitor mediated underexpression

of P_{38}MAPK in the treated panels demonstrated compromised intracellular parasite removal

(Fig. 3a) and confocal micrographs (Fig. 3b) suggested that $P2X_7R$ activation can positively

regulate P_{38}MAPK expression and most importantly both of them are essential in intracellular

Spergulin-A mediated the anti-leishmanial effect of which P_{38}MAPK act downstream of

$P2X_7R$.

NF-κBp65 expression alteration in Spergulin-A mediated leishmanicidal activity.

Likewise, the connection of NF-κB in Spergulin-A mediated infection control was also

monitored (Fig. 3c). The trend was almost alike of P_{38}MAPK expression, parasitic infection

down-regulated the expression altitude of NF-κB and Spergulin-A treatment was noticed to up-

regulate their expression when applied in the infection panel and in this case also the inhibitor

(Curcumin) were noticed to down-regulate NF-κB expressions with moderate increase of the

parasitic load even when co-applied with Spergulin-A (Fig. 3c vi, vii).

ROS and NO participate in anti-leishmanial effects of Spergulin-A. Changes in the

production of ROS as well as NO in respect to *L. donovani* infection and Spergulin-A

treatment was measured by flow cytometry (Fig. 4a, b) and confocal micrographs (Fig. 4c)

with the aid of fluorescent dyes DCFDA and DAF-2 DA respectively. It was observed that

ROS production was downregulated with respect to *L. donovani* infection and got up-regulated

in MΦs with Spergulin-A (30 μg/mL). In comparison to infected MΦs, Spergulin-A (30

μg/mL) exposure elevated ROS production in the infected MΦs (Fig. 4a iv, vi) which is in the

alignment of the lessened parasite load in them (Fig. 4c). However, when NAC has applied the

ROS abrogation (Fig. 4a v, vi) also leads to substantial loss of the parasiticidal activity of

Spergulin-A (Fig. 4c).

Spergulin-A (30 μg/ml) treatment was found to encourage the NO production in the

infected MΦs (Fig. 4b v, vii). It was found that the NO production has a positive effect in

checking the parasite load inside the MΦs as NMMA application in the Spergulin-A treated

infected MΦs reduced the effectivity of the compound (Fig. 4b vi, Fig. 4c) and strongly

support the anti-parasitic activity is regulated by NO.

Expression alteration of p-P$_{38}$MAPK, p-NF-κBp65, and Gp91-phox in intervening ROS

and NO involvement of Spergulin-A mediated anti-leishmanial effect. Gp91-phox the

glycosylated subunit of the heterodimer phagocytic NADPH oxidase flavocytochrome b558,

plays a crucial role in regulating the redox status of MΦs and thus can serve as a marker of

the alteration of oxidative stress due to Spergulin-A treatment in the infected MΦs, aimed

towards an explanation of the parasitic burden alteration treated with it at different doses.

Spergulin-A at different doses (10, 20 and 30μg/mL) applied in the infected MΦs and levels of

Gp91-phox was evaluated for both p-P$_{38}$MAPK and NF-κBp65 (Rel A) positive MΦs as

evaluated by both FACS (Fig. 5a) and confocal microscopy (Fig. 5b) (for only p-P_{38}MAPK).

Gp91-phox expression was simultaneously up-regulated with increasing doses of Spergulin-A

along with the increment of p-P_{38}MAPK expression (Fig. 5a i-v (upper panels) and Fig. 5a vi).

Likewise, Gp91-phox expression was also found to be elevated with respect to NF-κBp65

expression along with Spergulin-A in a dose-dependent manner (Fig. 5a i-v (lower panels) and

Fig. 5a vii).

Confocal images provided substantial evidence that with the increasing doses

Spergulin-A boosted p-P_{38}MAPK expression and also Gp91-phox expression simultaneously

(Fig. 5b). Infection with *L. donovani* lowered both p-P_{38}MAPK and Gp91-phox expression in

them compared to the control (Fig. 7b).

Phago-lysosomal maturation promotion by Spergulin-A in respect to its anti-leishmanial

activity. Phagolysosomal maturation considered a significant process to chuck out the

intracellular pathogens of phagocytic cells and the pathogens promoted multiple ways to evade

this sequential process. Spergulin-A mediated dynamics of Rab 5 an early phagosomal marker

and CathepsinD, a late phagosomal marker was observed by confocal microscopy (Fig. 6a). *L.*

donovani infection was found to suppress these markers compared to the control MΦs.

Whereas, the dose-dependent application of Spergulin-A found to amplified both Rab5 and

CathepsinD expression and signified the occurrence of phagolysosomal fusion and

demonstrated a key role of this event during leishmanicidal property arbitrated by Spergulin-A

(Fig. 6a).

In Vivo Evaluation of anti-leishmanial activity of Spergulin-A in lights of P_{38}-MAPK and

$P2X_7$ receptor involvement. Evaluation of the intracellular anti-leishmanial activity of

Spergulin-A in vitro against RAW 264.7 MΦs strongly favored the involvement of $P2X_7$R

activation and subsequent participation of P_{38}MAPK and NF-κBp65 downstream. Therefore an

in vivo evaluation of the anti-leishmanial activity of Spergulin-A with a demonstration of the

expression alteration analyses of the aforementioned key mediators could have positively

linked ours in vitro observations with its leishmanicidal affectivity in vivo. After treatment

schedule (Control and Treated group received 30mg/kg b.wt. of Spergulin-A for every alternate

day for 10 doses) was over mice were sacrificed and spleen and bone marrow MΦs were

harvested for each group and evaluated by flow cytometry for the linked expressions of

$P2X_7R$-$P_{38}MAPK$ and $P2X_7R$-NF-κBp65 against spleen MΦs (Fig. 6b) and for $P2X_7R$-

$P_{38}MAPK$ against bone marrow MΦs (Fig. 6c).

From the observations made against spleen MΦs the mean fluorescence index values

for $P2X_7R$ against both the experiments showed that the treated panels showed highest $P2X_7R$

activation and which was found to be least for the infected panels (Fig. 6b vii, viii) and

subsequent expressions of $P_{38}MAPK$ (Fig. 6b vii) and NF-κBp65 (Fig. 6b vii) showed the same

pattern; highest for the treated panels. However, for the Bone marrow MΦs although $P2X_7R$

expression was highest in the treated panel (Fig. 6c iii, iv), $P_{38}MAPK$ expression in the treated

panel was less than the infected panel (Fig. 6c ii, iv) which slightly differs from our all

previous observations both in vitro and in vivo combined.

The in vivo findings confirmed the in vitro findings and were perfectly in accordance of

the pathway of action speculated from the in vitro findings combined for the intracellular

activity evaluation of Spergulin-A; precisely with the involvement of $P2X_7R$, Ca^{++} influx,

$P_{38}MAPK$, NF-κBp65, ROS, NO and phagolysosomal maturation being the key mediators

sequentially. A graphical presentation of the step-wise anti-leishmanial activity propagation by

Spergulin-A has been demonstrated in Fig. 7d.

In silico binding assessment of Spergulin A with $P2X_7$ Receptor. The possible binding

mode of Spergulin A with its putative target $P2X_7$ was analyed by *in silico* methods. Molecular

docking study was employed to check the binding mode of investigated compound i.e.,

Spergulin A toward target $P2X_7$ (PDB: 5U1X). The molecular docking result illustrates few

hydrogen bonding interactions between the Spergulin A and P2X$_7$ receptor as shown in Fig. 7a.

The docked conformation of Spergulin A with P2X$_7$ indicates that the sulfonyl function interact

with Gln116, Trp167 (Fig. 7b). The amino acid residue Arg295 is found to form another

hydrogen bonding interaction with the oxygen atom of pyran nucleus (Fig. 7c). Moreover, it is

also noticed that the non-sugar part of Spergulin A is exposed to solvent accessible surface

area of the receptor.

Discussion

Leishmaniasis is a poorly investigated disease mainly affecting people in countries that are

developing and with challenged health practices. *Leishmania* living within the harsh

environment of phagocytes has developed strategies for prompt physiological adaptation,

escape from first-line defense systems and ability to inhibit a number of host cell functions.[38]

To endure and blossom inside macrophages *Leishmania* adopt various strategies to modulate

its physiology and signal kinetics. The direct influence of the parasite to the macrophages

linked to the suppressed generation of nitric oxide and oxidative burst and also their inhibitory

biological.[39,40] Indirectly, *Leishmania* influences the antigen presentation and T cell tasks in a

way that macrophage deactivation guides towards parasite endurance mainly by amendment of

host cell signaling of different kinases and phosphatases.[41] Being the host cell for the parasite,

any modulation of the macrophages also has an effect on the immune system and thus helping

their survival by overall immunomodulation of the host. Our previous effort[5] demonstrated that

G. oppositifolius and an isolated triterpenoid saponin, Spergulin-A was inconsistent of direct

leishmanicidal effect of against promastigotes and axenic amastigotes however proved to be a

potent intracellular antileishmanial mediator besides being an immune-modulator measured in

vitro in RAW 264.7 MΦs those infected with an *L. donovani* virulent strain. This observation

opens for the scope for precise demonstration of the pathways involved in its intracellular anti-

leishmanial effect and involvement and possible activation of P2X$_7$R has been considered.

P2X$_7$R reported being constitutively expressed on MΦ surfaces and also reported to be

associated with signaling cascades of inflammation, apoptosis and subsequent intracellular

pathogen control.[11] Additively, intracellular anti-leishmanial attributes of Spergulin-A

demonstrating the involvement of the purinergic receptor P2X$_7$ also hypothesized on the

reports of compounds screened towards the purinergic receptor amendments that about one-

third of the structurally diversified compounds are being natural products with a proven affinity

towards these purinergic receptors.[42]

Reduction of Spergulin-A mediated intra-MΦ leishmanial count was found to be linked

with P2X$_7$R activation and this reduction was hindered with P2X$_7$R antagonist BB-G

application and thus confirmed P2X$_7$R involvement and importance (Fig. 1a-e). Existing

literature suggested that P2X$_7$R possibly contributes to the removal of intracellular parasites

namely *Chlamydia*, *Leishmania*, *Toxoplasma*,[43,44] and *Mycobacterium tuberculosis* by

mediating strong pro-inflammatory and cytotoxic activity.[45] P2X$_7$R receptor activation was

found to be helpful in reducing *L. amazonensis* parasitic load in macrophages via leukotriene

B4 (LTB4) production.[46] Interestingly enough, P2X$_7$ knock out mice were observed to possess

increased lesion size and the parasitic load of *L. amazonensis* with an unwarranted CD4$^+$ T cell

proliferation.[47]

P2X$_7$R activation with extracellular stimuli like ATP can initiate opening of cation-

specific ion channel and result in the influx of Ca^{++} and the efflux of K$^+$ and with extended

exposure able to create pores in the plasma membrane resulted in the passage of larger

molecules.[11] Our experimental verification confirmed Spergulin-A activated P2X$_7$R mediated

Ca^{++} influx (Fig. 2) and unleashed a key mediator in intracellular leishmanial control.

Changes in the concentration of intracellular Ca^{2+} and PI3K activity are essential for

proper phagosomal maturation.[48] Ca^{++} is a key second messenger and cytosolic build-up of

Ca^{++} can lead to synaptic transmission, macrophage activation and apoptosis.[49] More

interestingly, the cytosolic Ca^{++} concentration affects the phagosomal maturation process by

modulating membrane fusion between phagosomes and lysosomal vesicles by activating

phospholipase D via RhoA, which can ultimately perish intracellular pathogens like

Leishmania[50] this phagolysosomal fusion can also mediated by regulating the activities of

two Ca^{2+}-dependent effector proteins, calmodulin and the multifunctional serine/threonine-

protein kinase CaMKII.[51] Spergulin-A mediated phagolysosomal maturation was confirmed

in the treated parasitized MΦs (Fig. 6a) by examining co-expression of early Rab5 and late

CathepsinD phagolysosomal markers in the treated cells and provided confirmation of the

event in intracellular parasite control.

An influx of Ca^{++} can activate the $P_{38}MAPK$, which in turn stimulates a number of

downstream effector signaling cascades.[52] The connection of phosphorylated $P_{38}MAPK$ and

leishmanial removal was closely monitored with Spergulin-A treatment (Fig. 3a, b). With

infection p-$P_{38}MAPK$ level lowered and treatment with Spergulin-A resulted in elevated p-

$P_{38}MAPK$ and diminished intracellular parasite count which again reversed with $P_{38}MAPK$

inhibitor application and thus one can conclude from the results the close association of

$P_{38}MAPK$ activation and leishmanial control. The same observation pattern was elaborated

for NF-κBp65 (Fig. 3c) wherewith the application of inhibitor compromised Spergulin-A

mediated intracellular parasite control with associated NF-κBp65 down-expression. Existing

reports also suggested that impaired $P_{38}MAPK$ expression during host-parasite interaction at

the primary hours proven beneficial during subsequent *L. donovani* infection establishment[24]

also $P_{38}MAPK$ inactivation by *Leishmania* parasites can lead to the iNOS inhibition and

inferior NO production.[39] Host invasion by intra-macrophage pathogens is frequently

associated with the restraining of NF-κBp65 total levels in infected macrophages, and this

reduction, in turn, can modulate the gene expressions necessary during the normal

immunological clearance of the parasites from the host cells.[53] Phosphorylated P_{38}MAPK

also promotes the association of NADPH oxidase at the cell membrane and resulted in

production and release of superoxide (O_2^-),[54] enhanced NF-κBp65 activation lead to the

ensuing transcription of inducible nitric oxide synthase (iNOS) and production of nitric

oxide (NO),[55] as well as production of tumour necrosis factor (TNF) and IL-6.[2] Spergulin-A

was monitored to activate Gp91-phox, a glycosylated subunit of NADPH oxidase

flavocytochrome b558 and also a marker of the NOX activity[56] with subsequent upregulated

p-P_{38}MAPK and p-NF-κBp65 expression (Fig. 5).

Enhanced production of ROS after $P2X_7R$ activation is well recognized in MΦs,[57]

mediated by NOX2 additionally also augment mitochondrial ROS.[58] *Leishmania* parasites

inside macrophages acclimatize various intimidating conditions during blood transmission,

and ROS and RNS by host macrophages.[59] At the onset of leishmanial infection, superoxide

production resulted from the oxidative burst due to phagocytosis[60] and later NO is produced

after activation of macrophages by IFN-y and TNF [61] resulting in abolition of intracellular

parasites.[62] The active subversion of ROS and NO production due to leishmanial infection

was noticed to be minimized during Spergulin-A treatment and application of their

abrogators minimized the anti-leishmanial effect of the compound (Fig. 4) and thus signifies

the importance of ROS and NO in intracellular parasite removal.

Meanwhile, a molecular docking study was performed to get a primary idea about the

binding interaction between Spergulin A and its putative target $P2X_7$. Few hydrozen bonding

interactions are observed between Spergulin A and amino acids at allosteric site binding of

$P2X_7$ (PDB: 5U1X). The docked conformation of Spergulin A with $P2X_7$ indicates the non-

sugar part of Spergulin A is exposed to solvent accessible surface area.

Finally, in vitro results from activity evaluation of Spergulin-A mediated intracellular

anti-leishmanial effect was directed to the inference that $P2X_7R$-P_{38}MAPK activation is the

key mediator of the said action with the downstream involvement of Ca^{++} regulated ROS, NO and phagolysosomal fusion as the effecter cascade. A flow diagram of which was depicted in Fig. 7d. Additionally, in vivo evaluation in determining the effect of Spergulin-A also established the fact that $P2X_7R$ and $P_{38}MAPK$ expression was evidently upregulated (Fig. 7a-c) and confirmed the in vitro findings.

Methods

Materials. Cell culture medium, Roswell Park Memorial Institute-1640 (RPMI-1640) and M-199, fetal bovine serum (FBS), penicillin-streptomycin (PS), and HEPES were acquired from Gibco BRL, Grand Island, NY, USA. Tissue culture plastic wares were acquired from NUNC (Roskilde, Denmark) and DAPI (4′,6-diamidino-2-phenylindole dihydrochloride), Lysotracker Red, CFSE and FURA 2-AM from Invitrogen, CA. Primary antibodies, secondary antibodies, and inhibitors were obtained from Santa Cruz Biotechnology, Santa Cruz, CA and Cell signaling technologies, USA.

Macrophages. Murine macrophage (MΦ) cell line RAW 264.7 (American Type Culture Collection, Rockville, MD) was cultured in Roswell Park Memorial Institute medium 1640 (RPMI-1640), supplemented with 10% heat-inactivated FBS and 100 μg/ml of antibiotic (penicillin-streptomycin)*.*

Parasites. Pathogenic *Leishmania donovani* [MHOM/IN/1983/AG83] infection was maintained by the passage in BALB/c mice. *L. donovani* promastigotes, attained from infected BALB/c mice spleen were cultured in complete M199 media (10% FBS, pH 7.4, 100U/ml penicillin G sodium, 100 μg/ml streptomycin sulfate, 25 mM HEPES, sodium bicarbonate (2.2g/L), 10% FBS and 1% Pen Strep (pH - 7.2-7.4) at 22°C. All pathogen and animal-related works were performed according to prior approval from Institutional Biosafety Committee, CSIR-Indian Institute of Chemical Biology (CSIR IICB/IBSC/CERT-32/18-19) and CSIR-

IICB-Animal Ethics Committee (IICB/AEC/Meeting/Oct/ 2017), respectively. The care and

maintenance of animals were done in agreement with the Committee for the Purpose of Control

And Supervision of Experiments on Animals (CPCSEA), guidelines.

Parasite infection and treatment. A ratio of 1:10 (MΦs: parasite) was used to infect

macrophages with *L. donovani* promastigotes. MΦs were incubated with parasites for 4 h,

washed with PBS, and either fresh media was added or treated with inhibitors for the required

time. To study the anti-leishmanial activity, normal and infected macrophages were treated

with Spergulin-A at effective concentrations up to 72 h. Miltefosine used as a positive anti-

leishmanial reference.

FACS and Confocal microscopy with CFSE tagged *L. donovani*. MΦs were parasitized with

CFSE-tagged promastigotes (25 μM/1×10^6 promastigotes in 1 ml media for 30 min) and

subjected to treatment with Spergulin-A (10, 20, 30 μg/ml)for 24 h.

For confocal microscopy, MΦs (1×10^5/glass coverslips) were washed twice with PBS

after infection and treatment, then fixed in chilled 70% ethanol. The nuclei were stained with

DAPI and observed under Olympus Fluoview FV10i confocal microscope with a 60X

objective lens. For FACS, MΦs (1×10^5/well in a six-well plate) were rinsed twice with PBS

after infection and treatment, harvested by scraping, resuspended in 400 μl PBS, and analyzed

by BD FACS LSR Fortessa with excitation at 494 nm and emission at 518 nm.

Measurement of Phospho-P$_{38}$MAPK, Gp91-phox, Phospho-NF-κBp65. For phospho-

P$_{38}$MAPK and Phospho-NF-κBp65 determination control, infected, and treated cells were

washed with PBS, then fixed with 70% ethanol, after perforation with Triton-x 100 solution

(0.5%) labeled with primary and respective fluorochrome conjugated antibodies. 20μM of

SB203580 was used to inhibit p-P$_{38}$MAPK and 25 μM of curcumin for was used as A p-NF-

κBp65 inhibitor with the necessary time of incubation periods. For the measurement of Gp91-

phox, the above procedures were followed with specific primary and secondary antibodies.

Cells were then analyzed in BD FACS LSR Fortessa.

Measurement of P2x7 with BBG. To determine $P2X_7$ level control, infected, and treated cells

were washed with PBS, then fixed with 70% ethanol, after perforation with Triton-x 100

solution (0.5%) labeled with primary and respective fluorochrome conjugated antibodies.

Before treatment with Spergulin-A, cells were treated with 10μM of Brilliant Blue G (BBG) to

inhibit $P2X_7$ in macrophage cells for 24 h of the incubation period.

Confocal Microscopy. After fixation with 70% ethanol and perforation, MΦs were stained

with primary and fluorochrome conjugated secondary antibody. Then slides were mounted and

nuclei were stained with DAPI in case of translocation of $p-P_{38}MAPK$, $P2X_7$, and Gp91-phox.

To study the leishmanicidal activity, CFSE was used to stain *L. donovani* and Lysotracker was

used to monitor acidic organelles in cells. Rab5 and CathepsinD were stained using primary

and respective fluorochrome conjugated secondary antibodies. All specimens were observed

under an Olympus Fluoview FV10i confocal microscope with a 60X objective lens.

Measurement of calcium release. Treated and untreated MΦs were collected, after washing

twice with PBS, treated with FURA 2 AM (Molecular Probes) at 37ºC. After 45 min of

incubation, the cells were washed with PBS, resuspended in fresh PBS, and examined by BD

FACS LSR Fortessa with an excitation wavelength of 380 nm and an emission wavelength of

510 nm.

FACS analyses of intracellular ROS and NO. Intracellular ROS was monitored based on the

transform in fluorescence of H2DCFDA. DAF-2 DA was used to perceive intracellular NO.[5]

Briefly, after infection and treatment, MΦs were scrapped and incubated in PBS containing 7.0

μM DAF-2DA (at 37°C for 30 min) or 20 μM DCF-DA (at 37°C for 15 min) and evaluated by

BD FACS LSR Fortessa with excitation at 480 nm and emission at 515 nm for both.

***In vivo* leishmanicidal activity.** To study leishmanicidal activity, parasite burdens in the

spleen and bone marrow of control and Spergulin-A treated mice were determined. After

confirmation of the *L. donovani* infection in the BALB/c and given an adequate time of the

infection propagation in vital organs like liver, spleen and bone marrow, infected mice were

divided into two groups arbitrarily with one control group. For phospho-P38 MAPK, phospho-

NF-κBp65 and P2X$_7$ protein determination, control and treated primary cells were stained

using primary and fluorochrome conjugated secondary antibodies. Then Cells were then

analyzed by BD FACS LSR Fortessa with respective excitation and emission wavelength.

In silico analysis. Molecular docking study[63,65] was employed to check the binding mode of

the investigated compound toward its putative target P2X$_7$R. Chem 3D ultra 8.0.3 (Chem 3D

Pro Version 8.0 And Chem Draw Ultra Version 8.0) software was used to construct the three-

dimensional structure of the investigated compound.[66] The energy minimization of the

compound was done by molecular orbital package (MOPAC) module with 100 iterations and a

minimum RMS gradient of 0.10.[64] The preparation of the ligand structure prior to docking is

performed by incorporating polar hydrogen atoms along with Gasteiger partial charges. The

torsions are allowed to rotate during docking.[67]

The crystallographic structures of P2X7 (PDB: 5U1X) were used for this analysis.[68]

The macromolecules were prepared by using Discovery Studio 3.0.[70] Polar hydrogen atoms

were added to the protein by using the AutoDock Tools (ADT).[71] The grid box was selected

5U1X centered at the corresponding ligand crystal structure by maintaining a spacing of 1 Å.[64]

Finally, the docking was performed by using the AutoDock Vina.[72] The docked poses were

visualized in Discovery Studio 3.0.[70]

Statistical analysis. All experimental values were denoted as mean ± SEM acquired from at

least three replicate experiments. Statistical significance and variations among groups were

evaluated with One-Way analysis of variance (ANOVA) followed by Dunnett's test. P values ≤

0.05 (*) or ≤ 0.01 (**) were considered as indicative of significance.

References

1. Tiuman, T.S., Santos, A.O., Ueda-Nakamura, T., Filho, B.P. & Nakamura, C.V. Recent advances in leishmaniasis treatment. *Int. J. Infect. Dis.* **15,** e525–e532 (2011).

2. Banerjee, S. et al. Attenuated Leishmania induce pro-inflammatory mediators and influence leishmanicidal activity by p38 MAPK dependent phagosome maturation in *Leishmania donovani* co-infected macrophages. *Sci. Rep.* **6,** 22335 (2016).

3. Alvar, J. et al. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One* **7,** e35671 (2012).

4. Freitas, E.O., Leoratti, F.M.S., Freire-de-Lima, C.G., Morrot, A. & Feijó, D.F. The contribution of immune evasive mechanisms to parasite persistence in visceral leishmaniasis. *Front. Immunol.* **7,** 153 (2016).

5. Banerjee, S. et al. Intracellular anti-leishmanial effect of Spergulin-A, a triterpenoid saponin of *Glinus oppositifolius*. *Infect. Drug Resist.* **12**, 2933–2942 (2019).

6. Oryan, A. Plant-derived compounds in treatment of leishmaniasis. *Iran. J Vet. Res.* **16,** 1–19 (2015).

7. Kaur, H., Thakur, A. & Kaur, S. Immunoprophylactic potential of a cocktail of three low molecular weight antigens of *Leishmania donovani* along with various adjuvants against experimental visceral leishmaniasis. *Iran. J. Parasitol.* **13,** 11–23 (2018).

8. Mandal, N. et al. A triterpenoid saponin possessing antileishmanial activity from the leaves of *Careya arborea*. *Phytochemistry* **67,** 183–190 (2006).

9. Delmas, F. et al. Antileishmanial activity of three saponins isolated from ivy, a-hederin, b-hederin and hederacolchiside A1, as compared to their action on mammalian cells cultured in vitro. *Planta Med.* **66**, 343–347 (2000).

10. Maes, L. et al. In vitro and in vivo activities of a triterpenoid saponin extract (PX-6518)

from the plant *Maesa balansae* against visceral leishmania species. *Antimicrob. Agents*

Chemother. **48**, 130–136 (2004).

11. Miller, C.M. et al. The Role of the P2X7 Receptor in Infectious Diseases. *PLoS Pathog.* **7,**

e1002212 (2011).

12. Aga, E. et al. Inhibition of the spontaneous apoptosis of neutrophil granulocytes by the

intracellular parasite *Leishmania major*. *J. Immunol.* **169**, 898–905 (2002).

13. Lees, M.P. et al. P2X7 receptor-mediated killing of an intracellular parasite, *Toxoplasma*

gondii by human and mouse macrophages. *J. Immunol.* **184**, 7040–7046 (2010).

14. Eskra, L., Mathison, A. & Splitter, G. Microarray analysis of mRNA levels from RAW

264.7 macrophages infected with *Brucella abortus*. *Infect. Immun.* **71**, 1125–1133 (2003).

15. Liu, J., Enomoto, S., Lancto, C.A., Abrahamsen, M.S. & Rutherford, M.S. Inhibition of

apoptosis in *Cryptosporidium parvum*-infected intestinal epithelial cells is dependent on

surviving. *Infect. Immun.* **76**, 3784–3792 (2008).

16. del Aguila, C. et al. *Encephalitozoon microsporidia* modulates p53-mediated apoptosis in

infected cells. *Int. J. Parasitol.* **36**, 869–876 (2006).

17. Chen, S. et al. P2X7 receptor antagonism inhibits p38 MAPK activation and ameliorates

neuronal apoptosis after subarachnoid hemorrhage in the rat. *Crit. Care Med.* **41**, e466-74.

(2013).

18. Pfeiffer, Z.A. et al. Nucleotide receptor signalling in murine macrophages is linked to

reactive oxygen species generation. *Free Radic. Biol. Med.* **42,** 1506–1516 (2007).

19. Gavala, M.L., Pfeiffer, Z.A. & Bertics, P.J. The nucleotide receptor P2RX7 mediates ATP-

induced CREB activation in human and murine monocytic cells. *J. Leukoc. Biol.* **84,** 1159–

1171 (2008).

20. Lee S.H. et al. Fucofuroeckol-A from *Eisenia bicyclis* inhibits inflammation in

lipopolysaccharide-induced mouse macrophages via downregulation of the MAPK/NF-κB

Signaling Pathway. *J. Chem.* **2016**, 6509212 (2016).

21. Cavalli, V. et al. The stress induced MAP kinase p38 regulates endocytic trafficking via

GDI:Rab5 complex. *Mol. Cell.* **7**, 421–432 (2001).

22. Prive, C. & Descoteaux, A. *Leishmania donovani* promastigotes evade the activation of

mitogen-activated protein kinases p38, c-Jun N-terminal kinase, and extracellular signal-

regulated kinase-1/2 during infection of naive macrophages. *Eur. J. Immunol.* **30,** 2235–

2244 (2000).

23. Jeong, H.J. et al. Inhibition of TNF-alpha and IL-6 production by Aucubin through

blockade of NF-kappaB activation RBL-2H3 mast cells. Cytokine **18,** 252–259 (2002).

24. Junghae, M. & Raynes, J. G. Activation of p38 mitogen-activated protein kinase attenuates

Leishmania donovani infection in macrophages. *Infect. Immun.* **70,** 5026–5035 (2002).

25. Gutierrez, M.G. Functional role(s) of phagosomal Rab GTPases. *Small GTPases* **4**, 148–

158 (2013).

26. Fischer, W., Urban, N., Immig, K., Franke, H. & Schaefer, M. Natural compounds with

P2X7 receptor-modulating properties. *Purinergic Signal.* **10,** 313–326 (2014).

27. Bartlett, R., Stokes, L. & Sluyter, R. The P2X7 receptor channel: recent developments and

the use of P2X7 antagonists in models of disease. *Pharmacol. Rev.* **66**, 638–675 (2014).

28. Stokes, L., Layhadi, J.A., Bibic, L., Dhuna, K. & Fountain, S.J. P2X4 Receptor Function

in the Nervous System and Current Breakthroughs in Pharmacology. *Front Pharmacol.* **8**,

291 (2017).

29. Helliwell, R.M. et al. Selected ginsenosides of the protopanaxdiol series are novel positive

allosteric modulators of P2X7 receptors. *Br. J. Pharmacol.* **172**, 3326–3340 (2015).

30. Orioli, E., De Marchi, E., Giuliani, A.L. & Adinolfi, E. P2X7 Receptor Orchestrates

Multiple Signalling Pathways Triggering Inflammation, Autophagy and Metabolic/Trophic

Responses. *Curr Med Chem.* 24, 2261-2275 (2017).

31. Santos, J.A.A. et al. Effect of *Rheedia longifolia* Leaf Extract and Fractions on the

P2X$_7$ Receptor *In Vitro*: Novel Antagonists? *J. Med. Food* **14**, 920-929 (2011).

32. Podolak, I., Galanty, A. & Sobolewska, D. Saponins as cytotoxic agents: a review.

Phytochem. Rev. **9**, 425–474 (2010).

33. Peng, W. et al. Systemic administration of an antagonist of the ATP-sensitive receptor

P2X7 improves recovery after spinal cord injury. *Proc. Natl. Acad. Sci. USA* **106**, 12489–

12493 (2009).

34. Wang, X.H., Xie, X., Luo, X.G., Shang, H. & He, Z.Y. Inhibiting purinergic P2X7

receptors with the antagonist brilliant blue G is neuroprotective in an intranigral

lipopolysaccharide animal model of Parkinson's disease. *Mol. Med. Rep.* **15**, 768–776

(2017).

35. Soares-Bezerra, R.J. et al. An Improved Method for P2X7R Antagonist Screening. *PLoS*

ONE **10**, e0123089 (2015).

36. Namovic, M.T., Jarvis, M.F. & Donnelly-Roberts, D. High throughput functional assays

for P2X receptors. *Curr. Protoc. Pharmacol.* Chapter 9:Unit 9 15 (2012).

37. Biswas, D. et al. ATP induced autophagy is associated with rapid killing of intracellular

mycobacteria within human monocytes/macrophages. *BMC Immunol.* **9,** 35 (2008).

38. Freitas, E.O., Leoratti, F.M.S., Freire-de-Lima, C.G., Morrot, A. & Feijó, D.F. The

contribution of immune evasive mechanisms to parasite persistence in visceral

leishmaniasis. *Front. Immunol.* **7**, 153 (2016).

39. Olivier, M., Gregory, D.J. & Forget, G. Subversion mechanisms by which *Leishmania*

parasites can escape the host immune response: a signaling point of view. *Clin. Microbiol.*

Rev. **18,** 293–305 (2005).

40. Van Assche, T., Deschacht, M., da Luz, R.A., Maes. L. & Cos, P. Leishmania-macrophage

interactions: insights into the redox biology. *Free Radic. Biol. Med.* **51**, 337–351(2011).

41. Soulat, D. & Bogdan, C. Function of macrophage and parasite phosphatases in

Leishmaniasis, *Front Immunol.* **8**, 1838 (2017).

42. Fischer, W., Urban, N., Immig, K., Franke, H. & Schaefer, M. Natural compounds with

P2X7 receptor-modulating properties. *Purinergic Signal.* **10,** 313–326 (2014).

43. Coutinho-Silva, R. et al. Inhibition of chlamydial infectious activity due to P2X7R-

dependent phospholipase D activation. *Immunity* **19**, 403–412 (2003).

44. Coutinho-Silva, R. & Ojcius, D.M. Role of extracellular nucleotides in the immune

response against intracellular bacteria and protozoan parasites. *Microbes Infect.* **14,** 1271–

1277 (2012).

45. Amaral, E.P. et al. Pulmonary infection with hypervirulent Mycobacteria reveals a crucial

role for the P2X7 receptor in aggressive forms of tuberculosis. *PLoS Pathog.* **10**, e1004188

(2014).

46. Chaves, S.P. et al. Modulation of P2X7 purinergic receptor in macrophages by *Leishmania*

amazonensis and its role in parasite elimination. *Microbes Infect.* **11,** 842–849 (2009).

47. Figliuolo, V.R. et al. The role of the P2X7 receptor in murine cutaneous leishmaniasis:

aspects of inflammation and parasite control. *Purinergic Signal.* **13,** 143–152 (2017).

48. Nunes, P. & Demaurex, N. The role of calcium signaling in phagocytosis. *J. Leukoc. Biol.*

88, 57–68 (2010).

49. Pinton, P., Giorgi, C., Siviero, R., Zecchini, E. & Rizzuto, R. Calcium and apoptosis: ER-

mitochondria Ca^{2+} transfer in the control of apoptosis. *Oncogene* **27,** 6407–6418 (2008).

50. Moradin, N. & Descoteaux, A. *Leishmania* promastigotes: building a safe niche within
macrophages. *Front. Cell Infect. Microbiol.* **2**, 121 (2012).

51. Swulius, M.T. & Waxham M.N. Ca(2+)/calmodulin-dependent protein kinases. *Cell Mol.*
Life Sci. **65**, 2637–2657 (2008).

52. Corre, I., Paris, F. & Huot, J. The p38 pathway, a major pleiotropic cascade that transduces
stress and metastatic signals in endothelial cells. Oncotarget. **8**, 55684–55714 (2017).

53. Calegari-Silva, T.C. et al. NF-κB-mediated repression of iNOS expression in *Leishmania*
amazonensis macrophage infection. *Immunol. Lett.* **127**, 19–26 (2009).

54. Tarafdar, A. & Pula G. The role of nadph oxidases and oxidative stress in
neurodegenerative disorders. *Int. J. Mol. Sci.* **19**, 3824 (2018).

55. McAdam, E. et al. Inducible nitric oxide synthase (iNOS) and nitric oxide (NO) are
important mediators of reflux-induced cell signalling in esophageal cells. **33**, 2035–43
(2012).

56. Yu, L., Quinn, M.T., Cross, A.R. & Dinauer, M.C. Gp91(phox) is the heme binding
subunit of the superoxide-generating NADPH oxidase. *Proc. Natl. Acad. Sci. USA*. **95**,
7993–7998 (1998).

57. Lenertz, L.Y., Gavala, M.L., Hill, L.M. & Bertics, P.J. Cell signaling via the P2X(7)
nucleotide receptor: linkage to ROS production, gene transcription, and receptor trafficking.
Purinergic Signal. **5**, 175–187 (2009).

58. Nakahira, K. et al. Autophagy proteins regulate innate immune responses by inhibiting the
release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat. Immunol.* **12**,
222–230 (2011).

59. Carneiro, P.P. et al. The role of nitric oxide and reactive oxygen species in the killing
of *Leishmania braziliensis* by monocytes from patients with cutaneous leishmaniasis. *PLoS*
ONE **11**, e0148084 (2016).

60. Almeida, T.F., Palma, L.C., Mendez, L.C., Noronha-Dutra, A.A. & Veras, P.S. *Leishmania amazonensis* fails to induce the release of reactive oxygen intermediates by CBA macrophages. *Parasite Immunol.* **34**, 492–498 (2012).

61. Gantt, K.R. et al. Oxidative responses of human and murine macrophages during phagocytosis of *Leishmania chagasi*. *J. Immunol.* **167,** 893–901 (2001).

62. Bogdan, C., Rollinghoff, M. & Diefenbach A. The role of nitric oxide in innate immunity. *Immunol. Rev.* **173**, 17–26 (2000).

63. Amin, S.A. et al. Pharmacoinformatics study of piperolactam a from Piper beetle root as new lead for nonsteroidal anti fertility drug development. *Comput. Biol. Chem.* **67,** 213–224 (2017a).

64. Amin, S.A., Adhikari, N., Agrawal, R.K., Jha, T. & Gayen, S. Possible Binding Mode Analysis of Pyrazolo-triazole Hybrids as Potential Anticancer Agents through Validated Molecular Docking and 3D-QSAR Modeling Approaches. *Lett. Drug Des. Discov.* **14,** 515–527 (2017b).

65. Bhardwaj, B. et al. Insight into structural features of phenyltetrazole derivatives as ABCG2 inhibitors for the treatment of multidrug resistance in cancer. *SAR QSAR Environ. Res.* **30,** 457–475 (2019).

66. Chem 3D Pro Version 8.0 And Chem Draw Ultra Version 8.0 Are Software Programs Developed by Cambridge Soft Corporation. USA.

67. Lv, P.C. Li, H.Q., Sun, J., Zhou, Y. & Zhu, H.L. Synthesis and biological evaluation of pyrazole derivatives containing thiourea skeleton as anticancer agents. *Bioorg. Med. Chem.* **18,** 4606–4614 (2010).

68. https://www.rcsb.org/structure/5U1X: https://www.rcsb.org/structure/5U1X (as accessed on 14th August 2019)

69. Karasawa, A. & Kawate, T. Structural basis for subtype-specific inhibition of the P2X7 receptor. *Elife*. **5,** e22153 (2016).

70. Discovery Studio 3.0 (DS 3.0), Accelrys Inc., CA, USA, 2015.

71. Autodock Tools 1.5.6 (ADT)/MGL Tools 1.5.6, The Scripps Research Institute, CA, USA, 2012. Available at http://mgltools.scripps.edu

72. Trott, O. & Olson, A.J. AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J. Comput. Chem.* 31, 455–461 (2010).

Acknowledgments

Sincere thanks are given to the Science and Engineering Research Board (SERB), Govt. of India (Grant No. PDF/2016/001437 and EMR/2015/001674) for financial support. The manuscript has been critically checked by the full version of the Grammarly software and Urkund for language corrections and plagiarism.

Author contributions

N.M., K.D.S., and S.B. conceptualize the work, N.M., and S.B. acquired, analyzed and written the biological data. S.A.A. and T.J. acquired and analyzed the in silico binding assay data. N.M., S.B. and SAA wrote the manuscript. N.M., T.J, S.D., and K.D.S. supervised the work.

Figure Captions

Fig. 1 Evaluation of $P2X_7$ receptor involvement in the intracellular anti-leishmanial activity of Spergulin-A. **a** Spergulin-A reduces intracellular *L. donovani* expressed as the number of parasites/20 MΦs dose-dependently after 24 h of treatment with miltefosine as positive control expressing its LC_{50} and LC_{90} values. **b** RAW 264.7 MΦ survivability in accordance with the $P2X_7$ receptor antagonist BB-G application after 24 h. **c** FACS monitored co-expression

analyses of CFSE-tagged *L. donovani* and subsequent $P2X_7R$ expression during infection and

Spergulin-A (30μg/mL) treatment. **d** Bar diagram of Spergulin-A mediated intracellular

leishmanial parasite removal and subsequent aversion by BB-G (10μM) application expressed

as the number of parasites/20 MΦs. **e** Confocal micrographs of consequent expression of

CFSE-tagged intracellular *L. donovani* with $P2X_7R$ expression during Spergulin-A (30μg/mL)

and BB-G (10μM) application. Spergulin-A enhanced $P2X_7R$ expression and intracellular *L.*

Donovani removal, whereas, BB-G caused reversal of the Spergulin-A mediated intracellular

parasite removal and $P2X_7R$ over-expression. Scale bars, 10 μm. Images are representative of

three separate experiments. All values are expressed as mean ± SEM from triplicate assays

from three independent experiments (*P*-values ≤0.05 (*) or ≤0.01 (**) vs infected panel except

for **b**, where the reference is the RAW 264.7 MΦs; ns denotes not significant).

Fig. 2 FACS monitored Ca^{++} influx evaluation with $P2X_7R$ expression analyses of *L. donovani*

parasitized, Spergulin-A and BB-G exposed MΦs by Fura2M. **a** FACS analyses with $P2X_7R$

expression and Ca^{++} corresponding to Fura2M. The four quadrants represent different stages of

$P2X_7R$ and Ca^{++} expressing populations, i.e. Q1 (only $P2X_7R$ expressing), Q2 (both $P2X_7R$ and

Ca^{++} expressing), Q3 (double negative) and Q4 (only Ca^{++} expressing). **b** Mean fluorescence

index values equivalent of the expressions of both $P2X_7R$ and Ca^{++} during *L. donovani*

infection, Spergulin-A treatment, and BB-G application. **c** Percentage of $P2X_7R$ and Ca^{++}

expressing RAW 264.7 MΦ cells against the parent population during *L. donovani* infection,

Spergulin-A treatment and BB-G application. Images are representative of three separate

experiments. All values are expressed as mean ± SEM from triplicate assays from three

independent experiments (*P*-values ≤0.05 (*) or ≤0.01 (**) vs infected panel).

Fig. 3 $P_{38}MAPK$ expression alteration with $P2X_7$ receptor involvement during Spergulin-A

treatment. **a** Flow cytometric analyses of $p\text{-}P_{38}MAPK$ expression alteration towards *L.*

donovani infection, Spergulin-A treatment and $P_{38}MAPK$-inhibitor (SB203580) application

and their co-dependence in Spergulin-A mediated anti-leishmanial effect with the histogram of

mean fluorescence values of p-P_{38}MAPK and CFSE-tagged intracellular *L. donovani* in

respective populations. **b** Confocal micrographs of co-expression analyses of $P2X_7R$ and

P_{38}MAPK in *L. donovani* parasitized, Spergulin-A and BB-G exposed MΦs. Scale bars, 10

μm. **c** NF-κBp65 expression alteration with $P2X_7$ receptor involvement during Spergulin-A

treatment. Flow cytometric analyses of NF-κBp65 expression alteration towards *L. donovani*

infection, Spergulin-A treatment and NF-κB-inhibitor (Curcumin) application and their co-

dependence in Spergulin-A mediated anti-leishmanial effect with the histogram of mean

fluorescence values of NF-κBp65 and CFSE-tagged intracellular *L. donovani* in respective

populations. Images are representative of three separate experiments. All values are expressed

as mean ± SEM from triplicate assays from three independent experiments (*P*-values ≤0.05 (*)

or ≤0.01 (**) vs infected panel).

Fig. 4 a Evaluation of changes in ROS production in control, infected, and Spergulin-A (30

μg/mL) treated MΦs with H2DCFDA by FACS with or without NAC. **b** Intracellular NO

production alterations were also examined with fluorescent probe DAF-2 DA in control,

infected, and Spergulin-A (30 μg/mL) treated MΦs by FACS with or without NMMA. **c**

Changes in CFSE-tagged intracellular *L. donovani* parasites were micrographed with respect to

Spergulin-A treatment and co-application of NAC and NMMA for evaluation of ROS and NO

involvement in Spergulin-A mediated anti-leishmanial effect by confocal microscopy, scale

bars, 10 μm. Images are representative of three separate experiments. All values are expressed

as mean ± SEM from triplicate assays from three independent experiments (*P*-values ≤0.05 (*)

or ≤0.01 (**) vs infected panel).

Fig. 5 Co-expression alteration of p-P_{38}MAPK and Gp91-phox, p-NF-κBp65 and Gp91-phox

in Spergulin-A mediated anti-leishmanial effect. **a** Spergulin-A at different doses (10, 20 and

30μg/mL) applied in the infected MΦs found to elevate levels of Gp91-phox for p-P_{38}MAPK

positive MΦs as evaluated by FACS. Histograms of percent parent populations for p-P$_{38}$MAPK positive MΦs and both Gp91-phox and p-P$_{38}$MAPK for different infected and Spergulin-A treated populations denoted their inter-connections. Spergulin-A at different doses (10, 20 and 30μg/mL) applied in the infected MΦs found to elevate levels of Gp91-phox for NF-κBp65 (Rel A) positive MΦs as evaluated by FACS. Histograms of percent parent populations for NF-κBp65 positive MΦs and both Gp91-phox and NF-κBp65 for different infected and Spergulin-A treated populations denoted their inter-connections. **b** Confocal microscopy elaborated that Gp91-phox expression was simultaneously up-regulated with increasing doses of Spergulin-A along with the increment of p-P$_{38}$MAPK expression. Images are representative of three separate experiments, scale bars, 10 μm. All values are expressed as mean ± SEM from triplicate assays from three independent experiments (*P*-values ≤0.05 (*) or ≤0.01 (**) vs infected panel).

Fig. 6 a Dynamics of phagolysosomal maturation in demonstrating the dose-dependent anti-leishmanial effect of Spergulin-A. Early phagosome marker Rab-5 expression got upregulated in the treated panel with respect to the control and infected panels. Cathepsin-D (Late phagolysosomal marker) expression also got upregulated with the exposure of Spergulin-A and signifies the phagolysosomal maturation process. Images are representative of three separate experiments, scale bars, 10 μm. **b** In vivo evaluation of the anti-leishmanial effect of Spergulin-A with P$_{38}$-MAPK and P2X$_7$ receptor involvement. Graphical presentation of the in-vivo treatment schedule and observation parameters. After confirmation of the *L. donovani* infection in the BALB/C mice, infected mice were divided into two groups arbitrarily with one control group. Control and treated groups received 30mg/kg b.wt. of Spergulin-A for every alternate day for 10 doses then sacrificed and spleen and bone marrow MΦs were harvested after treatment schedule and examined for P2X$_7$R, p-P$_{38}$MAPK and NF-κBp65 expression. Mice from each group were sacrificed and spleen MΦs were harvested for each group and evaluated

by flow cytometry for the linked expressions of $P2X_7R$-p-P_{38}MAPK and $P2X_7R$-NFκBp65.

Mean fluorescence index values equivalent of the expressions of both $P2X_7R$- p-P_{38}MAPK and

$P2X_7R$-NFκBp65 from spleen MΦs of different experimental mice groups. **c** Mice from each

group were sacrificed and bone marrow MΦs were harvested from each group and evaluated

by flow cytometry for the linked expressions of $P2X_7R$-p-P_{38}MAPK. Mean fluorescence index

values equivalent of the expressions of both $P2X_7R$- p-P_{38}MAPK from bone marrow MΦs of

different experimental mice groups. Images are representative of three separate experiments.

All values are expressed as mean ± SEM from triplicate assays from three independent

experiments (*P*-values ≤0.05 (*) or ≤0.01 (**) vs infected panel).

Fig. 7 In-silico docking study of the interdependence of Spergulin-A with P2X7 receptor. **a**

The proposed interaction investigated compound Spergulin-A with the P2X7 receptor. The

green-colored stick is the inbound crystal compound. **b** The proposed 3D interaction

investigated compound Spergulin-A with the amino acid residues of the $P2X_7$ receptor. **c** The

proposed 2D interaction investigated compound Spergulin-A with the amino acid residues of

the $P2X_7$ receptor. **d** A graphical presentation of the step-wise anti-leishmanial activity

propagation by Spergulin-A.

Spergulin-A

P2X₇ Receptor

$[Ca^{++}]$

P₃₈MAPK

IκB   NF-κB   IκB

P₃₈MAPK

NF-κB

Phago-Lysosomal Fusion

ROS
NO
Cytokines

Minimized MΦ

Spergulin-A

P2X₇ Receptor

Inside RAW 264.7 MΦ (In vitro Study)

Groups
  I.   Control
  II. Infected
  III. Sp-A Treated

- 30 mg/kg b.wt.
- Every alternate day, 10 doses
- Intraperitoneal

Spleen MΦ
Bone Marrow MΦ

L. Donovani promastigote

60 Days
Infection confirmed

Sacrificed on 20th day

Expression of
- P2X7
- p-P38 MAPK
- p-P65 Nf-κβ

FACS

Inside Mice Spleen and Bone Marroe MΦ (In vivo confirmation)

a — Survivality % of intracellular leishmanial forms vs Parasites (Amastigotes)/20 MΦ (RAW 264.7) 24 h post-infection

Legend:
- Infected MΦ
- Sp A 5µg/ml
- Sp A 10µg/ml
- Sp A 20µg/ml
- Sp A 30µg/ml
- Sp A 40µg/ml
- Miltefosine 25 µM
- Miltefosine 42 µM

b — MΦ (264.7) Survivality

Selected Dose: 91.92% viability

X-axis: Control Mφ, 1 µM, 10 µM, 20 µM, 30 µM, 40 µM, 50 µM, 60 µM, 70 µM, 80 µM, 90 µM, 100 µM

c

(i) Infected MΦ — *L. Donovani* (CFSE) P3 | P2X7 R+ P4
(ii) Treated (Sp A 30µg/ml) — *L. Donovani* (CFSE) P3 | P2X7 R+ P4
(iii) Infected MΦ + BB-G — *L. Donovani* (CFSE) P3 | P2X7 R+ P4
(iv) Treated (Sp A 30µg/ml) + BB-G — *L. Donovani* (CFSE) P3 | P2X7 R+ P4

(v) Mean Fluorescence Index Values vs Mean CFSE (*L. donovani*)
- Infected MΦ
- Treated (Sp A 30µg/ml)
- Infected MΦ + BB-G
- Treated (Sp A 30µg/ml) + BB-G

(vi) % of P Parent Populations expressing P2X7 R
- Infected MΦ
- Treated (Sp A 30µg/ml)
- Infected MΦ + BB-G
- Treated (Sp A 30µg/ml) + BB-G

d — Parasites/20 MΦ (RAW 264.7) 24 h post-infection vs Compounds Applied

Legend:
- Infected MΦ
- BB-G 10µM
- Sp A 30µg/ml
- Sp A 20µg/ml
- Sp A 10µg/ml
- Sp A 30µg/ml +BB-G
- Sp A 20µg/ml +BB-G
- Sp A 10µg/ml +BB-G

e

Columns: DAPI | *L. Donovani* Ag-83 (CFSE) | P2X7 (PE) | Merged

Rows:
- MΦ
- MΦ+ *L. donovani*
- MΦ+ *L. donovani* + BB-G
- MΦ+ *L. donovani* + Sp A 30µg/ml
- MΦ+ *L. donovani* + Sp A 30µg/ml+ BB-G

a

	DAPI	Rab 5 (PE)	Cathepsin D (FITC)	Merged
MΦ				
Infected MΦ				
Infected MΦ+ Sp A 10µg/ml				
Infected MΦ+ Sp A 20µg/ml				
Infected MΦ+ Sp A 30µg/ml				

b

Groups
I. Control
II. Infected
III. Sp-A Treated

- 30 mg/kg b.wt.
- Every alternate day, 10 doses
- Intraperitoneal

Spleen MΦ
Bone Marrow MΦ

Expression of
- P2X7
- p-P38 MAPK
- p-P65 Nf-κβ

FACS

L. Donovani promastigotes

60 Days
Infection confirmed

Sacrificed on 20th day

(i) MΦ — P2X7R+ / p-P38MAPK+
(ii) Infected MΦ — P2X7R+ / p-P38MAPK+
(iii) Treated MΦ — P2X7R+ / p-P38MAPK+
(iv) MΦ — P2X7R+ / NF-κBp65+
(v) Infected MΦ — P2X7R+ / NF-κBp65+
(vi) Treated MΦ — P2X7R+ / NF-κBp65+

(vii) / (viii) Cont. MΦ / Infected MΦ / Treated MΦ (30µg/ml)

Isolated Spleen MΦ after treatment schedule

c

(i) MΦ — P2X7R+ / p-P38MAPK+
(ii) Infected MΦ — P2X7R+ / p-P38MAPK+
(iii) Treated MΦ — P2X7R+ / p-P38MAPK+

(iv) Cont. MΦ / Infected MΦ / Treated MΦ (30µg/ml)

Isolated Bone marrow MΦ after treatment schedule