Inventive nesting behaviour in the keyhole wasp *Pachodynerus nasidens*

Latreille (Hymenoptera: Vespidae) in Australia, and the risk to aviation safety

Alan P.N. House[1]*, Jackson G. Ring[2&] and Philip P. Shaw[3&]

[1] Eco Logical Australia Pty Ltd, Brisbane, Queensland, Australia

[2] Operations Section, Brisbane Airport Corporation, Brisbane Airport, Queensland,

Australia

[3] Ecosure Pty Ltd, Burleigh Heads, Queensland, Australia

* Corresponding author

Email: alan.house@ecoaus.com.au

[&] These authors contributed equally to this work.

15
16

Abstract

The keyhole wasp (*Pachodynerus nasidens* Latreille 1812), a mud-nesting wasp native to South and Central America and the Caribbean, is a relatively recent (2010) arrival in Australia. In its native range it is known to use man-made cavities to construct nests. A series of serious safety incidents Brisbane Airport related to the obstruction of vital airspeed measuring pitot probes on aircraft possibly caused by mud-nesting wasps at prompted an assessment of risk. An experiment was designed to determine the species responsible, the types of aircraft most affected, the seasonal pattern of potential risk and the spatial distribution of risk on the airport. A series of replica pitot probes were constructed using 3D-printing technology, representing aircraft with high numbers of movements (landings and take-offs), and mounted at four locations at the airport. Probes were monitored for 39 months. Probes blocked by mud nesting wasps were retrieved and incubated in mesh bags. Emerging wasps were identified to species. Results show that all nests in probes were made by *P. nasidens*, and peak nesting occurs in the summer months. Nesting success (as proportion of nests with live adult emergents) was optimal between 24 and 31°C and that probes with apertures of more than 3 mm diameter are preferred. Not all areas on the airport are affected equally, with the majority of nests constructed in one area. The proportion of grassed areas within 1000 m of probes was a significant predictor of nesting, and probe volume may determine the sex of emerging wasps.

Introduction

Interactions between aircraft and wildlife are frequent and can have serious financial and safety consequences. Birds are the most common threat to aircraft [1, 2], with a host of terrestrial animals also implicated [3]. There were over 16,500 reported incidents involving birds and a further 397 involving other vertebrates at Australian airfields between 2008 and 2017 [4]. The majority of these incidents occurred during take-off (23.45%) or landing (34.24%), the two most vulnerable phases of flight. However, the risk posed by wildlife when aircraft are on the ground is much less understood and despite widespread anecdotal knowledge amongst the tropical and sub-tropical aviation community about insects and flight safety, specific threats posed by insects have not been quantified before. Here we report on an emerging insect-aircraft interaction that is a potentially lethal threat to flight safety.

In 2012, an exotic mud-nesting wasp (the keyhole wasp, *Pachodynerus nasidens* Latreille (Vespidae: Eumeninae)) was detected at Brisbane Airport (caught in an office off the international arrivals hall: Ross Wylie pers. comm.). This was the second record of the species in Australia after one was detected in northern Brisbane during a routine quarantine inspection of cargo initially received at the Port of Brisbane in 2010. *Pachodynerus nasidens* is native to tropical Central and South America and the Caribbean, and is also found in the southern USA (Florida, Texas and Arizona where it is possibly adventive [5, 6]. It has also been recorded from a number of Pacific islands including Hawaii, Polynesia, Micronesia and Japan [7, 8].

Pachodynerus nasidens is well known as an inquiline species, using abandoned or empty nests of other wasps, and for using man-made cavities (e.g. window crevices, keyholes, electrical sockets), to the extent that wholly constructed nests by the species are rare [9]. Indeed, it is known in USA as the "keyhole wasp". It will also

nest in the ground and construct mud nests attached to plants: this plasticity in

nesting behaviour is remarkable [5] and allows the species to adapt to new

opportunities for nesting in novel environments. It is a relatively small wasp,

distinguished from other species in the genus by the relatively short thorax and dull

yellow-brown markings (as opposed to bright yellow) on distal abdominal tergites

[10].

In its native range it is found in grasslands and swamps dominated by grasses and

sedges [11]. It is found at similar latitudes in South America (e.g. Guarapuava, Brazil)

as in Brisbane, from high altitudes (1120 m) in climates with cool wet seasons and

frosts to sea level [11].

The risk to flight safety

Airspeed is a critically important measure in aviation. Without an accurate measure of

airspeed pilots cannot easily judge take-off and landing and cannot ensure that the

aircraft flies in a safe speed range; fast enough to generate lift and remain airborne

but also slow enough to avoid structural failure under excess aerodynamic loads.

Pitot probes are hollow tube-like instruments that are commonly mounted on the

fuselage behind the nose cone and below to the cockpit. The probes measure

airspeed by subtracting mechanically the ambient "static pressure", sampled by a

separate sheltered hollow static port, from the "total pressure" of air entering the

exposed pitot probe to give a "dynamic pressure". Dynamic pressure directly equates

to speed through the air.

Aircraft manufacturers and aviation authorities recognise that amongst the issues

that might cause the pitot static system to fail to indicate correct airspeed, a blockage

(or even partial blockage) of a pitot probe or static port by insects (or other agents

such as ice) is a significant hazard [12, 13, 14, 15, 16, 17,18]. Anomalies between

separate airspeed indicators can lead to costly and hazardous rejected take-offs or

turn backs [e.g. 15, 16] or even catastrophic consequences [12, 13, 17, 18]. In

February 1996 a Boeing 757 crashed shortly after take-off from the Dominican

Republic, killing all 189 passengers and crew. Anomalous airspeed readings from the

pitot probes were responsible for the pilots misjudging the aircraft's speed. A sphecid

(mud-dauber) wasp was believed to have made a nest in one of the pitot probes,

although none were recovered [19]. The plane had been standing at Gregorio

Luperón International Airport in Puerto Plata, and for the two days prior to the fateful

flight the pitot probes were not covered as recommended by the manufacturer [13].

At Brisbane Airport, five incidents were reported between January and March 2006 in

which pitot probes gave inconsistent readings and flights were either rejected on

take-off or proceeded to their destination. All aircraft were Airbus A330s. An incident

on 19 March 2006 led to dangerous brake heating and tyre deflation when take-off

was rejected, resulting in the deployment of airport rescue and fire-fighting services

[15]. On inspection, it was found that the pitot probe on the pilot-in-command's side

had "wasp-related debris". Material retrieved from this probe by the Australian

Museum included fragments of an insect body including the head of a wasp.

In November 2013, an A330 prepared to take off but returned to the bay after

airspeed discrepancies between the captain's and first officer's readings. After

checking electronic instrumentation, the plane was cleared for a second take-off, but

another airspeed discrepancy occurred, and the captain returned under emergency

procedures. Sand and mud consistent with a mud dauber wasp nest was found to be

blocking the captain's probe [16]. Since this incident, more detailed records of wasp-

related issues have been recorded at Brisbane and a total of 26 were reported

between November 2013 and April 2019.

Partial completion of a nest in a pitot probe can take place very quickly. An A320 was

found to have a blocked total air temperature probe on arrival in Newcastle from

Brisbane in August 2015. The aircraft had been on the ground in Brisbane for only 30

minutes.

This paper reports on an experimental approach at Brisbane Airport to determine

which species are implicated in pitot probe blockages, which aircraft are being

targeted, and where on the airport the greatest risk of blockage occurs. Data on

nesting preferences, seasonality, reproductive success will be essential in directing

management to reduce the risks posed by this issue to airlines and the flying public.

124   Materials and methods

Study site

Brisbane is located on the eastern seaboard of Australia (27.3911°S, 153.1197°E),

on the shores of Moreton Bay, Queensland at approximately 4 m above sea level. It

is flanked on the east by the Brisbane River, and on the west by ecologically

significant coastal wetlands (mangroves and saltmarshes) and to the south by the

city of Brisbane. The climate is sub-tropical, with a long-term mean annual rainfall of

1190 mm, 50% of which falls between December and March. The annual mean

maximum temperature is 25.4°C, and annual mean minimum is 15.7°C, with fewer

than 2 frost days per year [20].

Probe deployment

Replica pitot probes were installed at Brisbane Airport between February 2016 and

April 2019. Pitot probes vary in design between each airframe type, so aircraft

movement data (frequency of arrivals and departures) from 2015 at Brisbane Airport

and aircraft location (which terminals and gates aircraft used) were used to determine

which airframes should be included in the study. Airframes were also filtered for pitot

probe dimensions, so that a range of aperture diameters and chamber depths (i.e.

distance to first baffle) were used. Six airframes/pitot probe types were selected

(Table 1).

Table 1. Dimensions of pitot probes for six airframes types and deployment locations
at Brisbane Airport. QLINK, QantasLink apron; DTBN, Domestic Terminal North; DTBS,
Domestic Terminal South; ITB, International Terminal (see Fig. 1).

Airframe	Aperture diameter (mm)	Distance to first baffle (mm)	Volume (mm^3)	Tip form	Locations
Embraer ERJ90	2.5	29.0	570	Concave	QLINK, DTBN, DTBS
De Havilland Dash 8	4.0	19.0	239	Concave	QLINK, DTBN, DTBS
Boeing 737-800	5.0	62.0	1218	Concave	QLINK, DTBN, DTBS, ITB
Airbus A330	5.5	33.0	784	Concave	QLINK, DTBN, DTBS, ITB
Boeing 737-400	7.2 (9/32")	33.0	1344	Convex	QLINK, ITB
Boeing 747-400	8 (5/16")	36.0	1810	Convex	QLINK, DTBN, DTBS, ITB

As sufficient decommissioned pitot probes were difficult to source, we manufactured

probes from UV-resistant ABS plastic using 3-D printing technology, based on

detailed engineering designs for accuracy. Probes were mounted on 3 mm x 900 mm

x 1200 mm white-painted steel sheets (to simulate an aircraft fuselage) which were

themselves attached to a weldmesh panel for rigidity (Fig. 2). Panels were secured to

structures as close as possible to aircraft parking locations (e.g. gate light poles at

the domestic terminal, aerobridges at the international terminal; Fig. 2) with

packaging straps and heavy duty stretch straps and flagged for visibility. They were

erected at heights similar to the aircraft-mounted probes (2-2.5 m at the domestic

terminal for e.g. A320, 4 m at the international terminal for e.g. 747-400 aircraft) (Fig.

2).

Three panels were established at each of four locations at Brisbane Airport (Fig. 2)

where incidents involving potential mud wasp activity was suspected or where wasps

had been recorded as potentially threatening aircraft: the northern domestic terminal

(QLINK and DTBN), southern domestic terminal (DTBS) and the international

terminal (ITB). The minimum distance between groups of three panels was 233 m.

Intercept traps

In addition to the mounted replica probes at the gates, a series of intercept traps

were established in November 2018 in both airside (16 traps) and landside (11 traps)

locations. These were included to determine if nesting in pitot probes could be

reduced by attracting females to alternative nest sites away from stationary aircraft.

Each trap consisted of 68 cardboard tubes 8 mm in diameter and 153 mm long, with

one open end. Blocked tubes were removed and replaced by clean ones at weekly

intervals.

Figure 1. Location of pitot probe panels at Brisbane Airport.

Figure 2. Pitot panels installed on gate light pole at Domestic Terminal (left) and on
aerobridge at International Terminal (right).

Monitoring

Probes were inspected weekly to monthly, depending on season and wasp activity.

Blocked probes were removed from the panels and placed in fine mesh bags with

labels recording date, probe type and location. Bags were kept in a mildly air-

conditioned room (25-27°C) and monitored for emergence. Emerging invertebrates

were placed in vials with methanol, and the date, species, number of individuals and

sex were recorded. Species that could not be identified were sent to the Queensland

Museum for determination. Retrieved probes that remained blocked after 70 days

were opened to determine if there were unhatched wasps in front and/or rear cells.

Probes that did not result in successful emergence were examined for remains of

undeveloped wasps and unconsumed prey.

Data analysis

We analysed the wasps' probe choice, blockage rates, locations and the relationship

between nesting and environment. As we did not have a perfectly balanced design

(in respect of numbers of probes of each airframe), probe blockage rates were

expressed as blockages per 12 probes for analyses of probe choice. Chi-squared

and Fisher exact tests (for small samples) were performed on blockage rates from

pitot probe types and airport locations, and ANOVA on blockage rates and pitot

probe diameter and volume.

Broad habitat types that wasps might respond to and be influenced by in terms of

resource availability (i.e. water and sediment for nest building, vegetation for

Lepidopteran prey) were mapped around each pitot panel location using Nearmaps®

satellite imagery from May 2016. Habitats were mapped into the classes in Table 2,

and the areas of each habitat class calculated for 20 m, 50 m, 100 m, 200 m, 500 m

and 1000 m radii around each panel in ArcGIS. Multiple regression analyses were

used to investigate relationships between nesting and environment.

Table 2. Broad habitat types mapped at Brisbane Airport.

Habitat	Description	Value to wasps	Location on airport
Natural horizontal	Predominantly grassed areas e.g. around runways, taxiways, aprons	Source of prey items; water and sediment for nesting	Airside, landside
Natural vertical	Treed areas e.g. *Casuarina* plantations	Source of prey items; water and sediment for nesting	Landside
Mixed natural	Tree/shrub/ground cover mixtures e.g. garden areas around landside infrastructure	Source of prey items; water and sediment for nesting	Landside
Built horizontal	Bitumen or concreted areas e.g. runways, taxiways, major roads (includes construction sites)	Sediment and water for nests	Airside, landside

Results

Seasonality of pitot blockages

The first pitot probe blockages occurred two weeks after the probe panels were

installed. In total, 93 instances of fully blocked probes were recorded during the 39-

month study. There is a seasonal pattern, with 96% of nesting occurring between

November and May (Fig. 3). Data from 2017 indicate that nesting of *P. nasidens*

extended into July, with hatching as late as August. Nesting has been recorded in

every month of the year except July and September. There is no relationship

between nesting and rainfall ($F_{1.37}$ = 0.0001, p = 0.99), or with rainfall in the previous

3 months ($F_{1,37}$ = 0.837, p = 0.37). However, nesting was significantly correlated with

the average monthly temperature ($F_{1,37}$ = 7.33, p = 0.01).

Figure 3. Pitot probe blockages by mud-nesting wasps and rainfall at Brisbane Airport,

February 2016 – April 2019.

Probe types blocked

The blockage rate by aircraft type indicate that probes with apertures ≥2.5 mm are

preferred (Table 3). Proportionally, probes from 737 airframes (737-400 and 737-800,

diameters 5-7.2 mm) are most likely to be blocked (56.3% of all blockages),

especially the 400 series, followed by the A330 (19.3%, diameter 5.5 mm). The

smallest probe (Embraer ERJ90, diameter 2.5 mm) had the lowest rate of blockage.

Although the standardised rates of pitot probe blockage were significantly different

from an even distribution (χ^2 = 28.08, P < 0.01), neither probe aperture diameter nor

distance to first baffle were significant predictors of nest choice ($F_{2,57}$ = 0.909, P >

0.05).

Table 3. Pitot blockages by aircraft type, February 2016-April 2019 at Brisbane Airport.

Airframe type	Standardised rate of blockages (per 12 probes)	Probe aperture diameter (mm)
ERJ90	1.3	2.5
DHC-8	14.6	4.0
B737-800	23	5.0
A330	23	5.5

B737-400	44	7.2
B747-400	13	8

An ANOVA on blockages showed that there were no significant differences in rates

of nesting between probe types ($F_{5,54}$ = 1.411, p > 0.05), even though 86% occurred

in probes with a diameter range of 5-8 mm.

Location on airport

Of the 93 blockages, 93.5% were detected at the QLINK apron: this results in a highly

significant deviation from expected ratios (χ^2 = 91.59, p = <<0.01), with no significant

differences between other locations in pair-wise tests.

In addition to this data on nesting in pitot probes, *P. nasidens* was recorded nesting

in a number of other locations around the airport including on-ground servicing

equipment, and in timber trap nests that were separately deployed to trap wasps.

Habitat distributions

Probes closer to natural habitats are more likely to be blocked than those further

away (Fig. 4). Linear regression between the number of blockages and the minimum

distance of panels to each habitat type shows that the proximity to natural horizontal

habitat is significant (R = -0.73, P < 0.01) but proximity to natural vertical habitat is

not (R = -0.025, p > 0.05) (Fig. 4). A multiple regression of the number of probe

blockages and the proportion of habitat types at increasing distance shows that the

amount of natural horizontal habitat at 1000 m is also significant (Table 4).

Figure 4. Relationship between pitot probe blockages and minimum distance to (a)
natural horizontal habitat and (b) natural vertical habitat.

Table 4. Multiple regression analysis of proportions of each broad habitat type at increasing distance from probe panels and rate of probe blockage. *P* values in bold indicate positive significant relationship; P values in italics indicate significant negative relationship.

				Habitat										
	ANOVA		natural horizontal		natural vertical		mixed natural		built horizontal		built vertical		water	
Distance (m)	F	*p*	t	*p*	t	*p*	t	*p*	t	*p*	t	*p*	t	*p*
20	2.343	0.157	-	-	-	-	-	-	-1.531	0.157	-	-	-	-
50	4.849	**0.037**	-	-	-	-	-	-	-1.345	0.211	-2.809	*0.020*	-	-
100	2.913	0.101	0.122	0.913	-	-	-	-	0.091	0.930	0.074	0.943	-	-
200	7.782	**0.010**	-2.166	0.067	-	-	-3.340	*0.012*	-2.610	*0.035*	-2.836	*0.025*	-	-
500	2.402	0.177	1.425	0.214	1.429	0.212	1.399	0.221	1.423	0.214	1.401	0.220	0.921	0.399
1000	5.399	**0.032**	2.461	**0.049**	2.305	0.061	-1.637	0.153	2.229	0.067	2.191	0.071	-0.887	0.395

Nesting success

There were 93 occurrences of blocked pitot probes. Of these, 37 (39.8%) produced live adult wasps, 18 (19.4%) had developed but unhatched wasp imagos, and 38 (40.9%) had contents that were either undeveloped or parasitised. All adult mud-nesting wasps that emerged from pitot probes were *P. nasidens*.

There is no consistent trend in successful nesting (i.e. completed nests and live hatching) with time of year (Fig. 5), and no significant relationship between hatching and rainfall during nesting ($F_{1,37} = 0.010$, P >> 0.05): however, rainfall during the previous 3 months and nesting success was positively related ($F_{1,37} = 7.998$, p < 0.01). Similarly, mean maximum temperature during nest development and ultimate nesting success were not related ($F_{1,37} = 0.409$, p > 0.05). Probes blocked during the peak of summer temperatures and rainfall (January-March: man max. temp. over 3 years 29.6°C) had lower success rates, nests completed late in the nesting season (April-June: mean max. temp. 24.1°C) had a greater likelihood of successful emergence and probes blocked after mid-May-October (mean max. temp. 23.4°C) developed to the adult stage but did not hatch.

Figure 5. Mean number of days (from collection of probes) to emergence of live wasps, February 2016 – April 2019.

Incubation times varied greatly, from 16 to 138 days with an average of 45. In the 2016-17 and 2017-18 summer nesting periods, nests that were completed at the start and the end of these periods appear to mature more quickly (Fig. 5).

Blocked probes have had 1-3 cells built, but only two (both A330), have produced two adults, and one produced three (B737-800). There was no significant difference

in probe types in the number of cells yielding more than one adult wasp (χ^2 = 5.64, p

= 0.130).

Probes that did not result in successful hatching were examined for remains of

undeveloped wasps and unconsumed prey. All had either pre-adult *P. nasidens*

pupae or had evidence of caterpillar and/or beetle larvae: none had evidence of other

native wasps or spiders (which would indicate other genera of Vespidae).

Only one probe produced live adults of a potential parasitoid of *P. nasidens*: five

Chrysis lincea (Chrysididae) adults hatched from a B737-400 probe retrieved in May

2017.

Sex ratios

A male-biased sex ratio of 1.53:1 was recorded from the pitot probes: this does not

significantly deviate from a 1:1 relationship (χ^2 = 0.748, p = 0.945) overall, but when

data is grouped by probe type there is a significant difference between males and

females (χ^2 = 2.233, p = 0.693, Fisher's exact test). Males and females were equally

likely to hatch from A330 and 747-800 probes (observed ratio 1:1), but males were 3

and 1.75 times more likely to hatch from 737-800 and 737-400 probes respectively.

There is also a close curvilinear relationship (r^2 = 0.942) between the proportion of

hatching males and pitot probe volume with the greatest proportion hatching from

mid-sized probes, but this relationship is not as close for females (r^2 = 0.517) (Fig. 6).

Figure 6. Relationship between pitot probe volume and sex of emerging live adult
wasps. Second order polynomial line of best fit fitted. r^2_{male} = 0.942 (p < 0.05), r^2_{female} =
0.517 (ns).

Discussion

Nesting by *P. nasidens* at Brisbane Airport follows general seasonal patterns found within its native range [21], with peak nesting occurring in warmer, wetter months although the only significant relationship was to temperature. Nesting success (as expressed by the proportion of nests producing live adults) is very consistent with other studies of this species (39.8% compared with 39.7% in [21], and incubation times are also generally within ranges published for native populations (16-138 days compared with 23-41 days [22], 5-59 days [21], 20-24 days [23].

Nesting activity is mostly confined to the summer months, which broadly agrees with other studies [21, 23], where egg development was optimal between 26-31°C. At Brisbane, nests completed after the peak of summer (when mean maximum temperatures fell below 26°C) still developed and hatched adults, suggesting some local climatic adaptation to slightly lower temperatures.

Sex ratios of *P nasidens* emerging from trap nests within its native range are variable, from 2:1 male to female (from a very small sample size) [11], 1:1 [24], and 0.56:1 [21]. Evidence suggests that local conditions of weather, prey availability and nest opportunities can affect sex ratios in cavity-nesting species [25, 26]. Data from this study suggests that there may be an effect of pitot probe nest volume on the proportion of males successfully hatching, but not the number of females.

The emergence of *Chrysis lincea* from blocked pitot probes indicates that some parasitism of *P. nasidens* is occurring, and the presence of *Melittobia* sp., *Cotesia* sp. and *Perilampus* sp. from mud nests on adjacent terminal buildings suggests that these parasitoids may also contribute to control of *P. nasidens*. A *Melittobia* species parasitised *P. nasidens* in Cuba, affecting between 41.6 and 75% of all cells [27].

The occurrence of *P. nasidens* at Brisbane Airport and the high number of pitot

incidents there due to mud wasps indicates that this species is primarily responsible,

but other species cannot be excluded. The earliest recorded aircraft incident in 2006,

if attributable to this species, indicates that it may have arrived some time before the

first official record in north Brisbane 2010 and at Brisbane Airport in 2012. Other mud

nesting wasp species known to be present at the airport include (from observation

and live captures) include *Abispa ephippium, Anterhynchium nigricinctum, Delta*

campaniforme, Delta sp. aff. *nigricornis*, *Delta* sp. (Vespidae: Eumeninae), *Chalybion*

sp. aff. *bengalense, Sceliphron formosum* (Sphecidae: Sceliphrinae), *Pison*

pyrrhicum, Pison sp. (Crabronidae: Crabroninae).

The choice of pitot probe type for nesting in *P. nasidens* appears to be based on a

minimum aperture of 2.5-3 mm. Adult females are 10-12 mm long, and observations

of active nesting in probes showed that wasps leave head-first, indicating that they

turn around inside the probe. A smaller aperture may be enough to allow a pupa to

develop and hatch, but adult females cannot physically negotiate the narrow bore. In

Brazil, *P. nasidens* was found to prefer apertures of 6-9 mm and did not use traps

with apertures of 4 mm which were also available [11]. [21] found *P. nasidens* and its

congeners did not nest in cavities with apertures above 7 mm in diameter. However,

P. nasidens has been found nesting in bamboo canes between 17 and 23 mm in

diameter [28]. Of the other mud wasps at Brisbane Airport (above), only *Pison*

pyrrhicum is likely to be small enough (7.4-7.7 mm long) to negotiate the pitot probes.

Females of the other species are 18 mm (*Chalybion bengalense*) to 30 mm (*Abispa*

ephippium) in length.

The environment available at Brisbane Airport to *P. nasidens* for securing nesting

resources is entirely modified. All grassed areas are managed (i.e. mown) to reduce

their attraction to birds; plantings of native species in plantations and gardens are

similarly semi-natural at best. The closest environments to the pitot probe panels are

highly modified structures that provide few nesting resources (prey, sediment and

water), except for pitot probes. There is evidence that the extent of grassy habitats at

1000 m from the panels may influence nesting success: this shows that wasps are

prepared to use the pitot probes despite making longer flights to gather nest building

resources.

Nesting in this species is extremely efficient at Brisbane Airport. Time to complete a

two-cell nest may be as short as a few hours: anecdotal evidence suggests that a

complete nest can be constructed in a pitot probe within 5 hours – this occurred at

night between arrival at 20:41 and 05:49 on 1 March 2014 (Boeing 737). In its native

range (i.e. Jamaica) the total time to complete a cell is between 2.5 and 4.75 hrs,

with a 3-cell nest in a 6 mm diameter cavity completed in 3.5 hrs [23]. As most nests

in pitot probes at Brisbane are only single-celled, nest-building is considerably more

rapid and may be limited to provisioning a single cell and applying a closing plug of

mud. In respect of dangers to aircraft however, the nest does not need to be

complete: the first addition of mud for the rearmost cell wall (if required) or

introduction of the first prey item is enough to cause anomalous airspeed readings as

air flow in the pitot probe is impeded. Wasps have been observed inspecting aircraft

noses within a few minutes of arrival at the gate, suggesting some experiential

learning and memory of the nesting resource: the mobile and transient nature of the

probes on aircraft (i.e. planes come and go) makes this choice of real pitot probes for

nesting even more remarkable.

P. nasidens is native to tropical South America, extending to the southern USA

(Florida, Texas and Arizona), including islands in the Caribbean region, although it is

possibly adventive in the USA [5]. It has been recorded from a number of Pacific

islands including Hawaii and Micronesia. The spread of *P. nasidens* across the

Pacific region does not follow a precise chronology [7] (recorded dates mark when

the species was first observed) but does indicate that the species has been well-

established outside its natural range since at least 1912. This dispersal is likely to

have been through shipping, although the possibility that wasps are carried on

aircraft cannot be discounted, especially as much of the spread is post WW2 when

air traffic began to increase in the region: aircraft luggage bay temperatures may be

sufficiently high (7°C to over 25°C: [29]) to allow adult wasps or pupae in nests to

survive, and even wheel-wells may be suitable for shorter and lower altitude flights

[30]. The appearance of *P. nasidens* in Hawaii as early as 1912 confirms that it

arrived there by boat, as the first crossing of the Pacific to Hawaii (O'ahu) by

aeroplane from the west coast of USA did not take place until 1927. The progression

of the species across the Pacific is not a neat progression over time from east to

west: this may reflect the opportunistic nature of dispersal, or simply the pattern of

investigation of island invertebrate faunas by entomologists, or both.

As an adaptable, inventive and highly mobile species outside of its natural range, *P.*

nasidens has the potential to spread from Brisbane to other locations in Australia where

climates are suitable. Having arrived in Australia, the species has established in a

challenging environment but one that provides all the basic requirements for population

persistence and has identified a potential nesting opportunity that is both transient and

mobile. In doing so, *P. nasidens* poses a significant risk to aviation safety, and further

work is warranted to determine the prospects for its control or, preferably, eradication.

Acknowledgments

We would like to thank Qantas (Matt Hill and Brian Ford) for their support of this project

and their assistance with manufacturing the probes and panels. Matt has also been an

extremely enthusiastic champion of the work and provided critical information on pitot

probes and aviation operations. Dave Selby and Peter Dunlop (Brisbane Airport

Corporation) were early supporters of this work, and Ross Wylie (Biosecurity

Queensland) gave useful advice on wasp ecology. Chris Burwell (Queensland

Museum) provided wasp identifications, and Jeff McKee, Susan House and Carissa

Free provided helpful comments on an earlier manuscript.

Financial disclosure statement

This work was fully funded by Brisbane Airport Corporation (https://www.bne.com.au/)

through contracts to Ecosure Pty Ltd and Eco Logical Australia Pty Ltd.

Competing interests

There are no competing interests.

Author contributions

APNH and PPS conceived the study; APNH, PPS and JGR collected the data; APNH

analysed the data; APNH wrote the original draft with inputs from PPS and JGR.

Data accessibility

Data and figures for this study are available at DOI 10.17605/OSF.IO/T87ZY

References

1. DeVault T, Blackwell BF, Seamans TW, Begier MJ, Kougher JD, Washburn JE, Miller PR, Dolbeer RA. Estimating interspecific economic risk of bird strikes with aircraft. Wildl Soc Bull. 2018; 42: 94-101.

2. Marra PP, Dove CJ, Dolbeer R, Dahlan NF, Heacker M, Whatton JF, et al. Migratory Canada geese cause crash of US Airways Flight 1549. Front Ecol Environ. 2009; 7: 297-301.

3. Schwarz KB, Belant JL, Martin JA, DeVault TL, Wang G. Behavioral traits and airport type affect mammal incidents with U.S. civil aircraft. Environ Manage. 2014; 54: 908-18.

4. Australian Transport Safety Bureau. Airspeed indication failure on take-off involving Airbus A330, 9M-MTK, Brisbane Airport, Queensland, on 18 July 2018. Australian Transport Safety Bureau, Canberra; 2018.

5. Carpenter JM. The genus *Pachodynerus* in North America (Hymenoptera: Vespidae: Eumeninae). Proc Entomol Soc Wash. 1986; 88: 572-77.

6. Global Biodiversity Information Facility. Global Biodiversity Information Facility search for *Pachodynerus nasidens*; 2019. [cited 27 May 2019]. GBIF Occurrence Download. Available from: https://doi.org/10.15468/dl.f65ys6.

7. Yamane S, Gusenleitner J, Menke AS. *Pachodynerus nasidens* (Latreille) (Hymenoptera, Vespidae), an adventive potter wasp new to Japan. Spec Divers. 1996; 1: 93-7.

8. Barrera-Medina R. Notas sobre el género *Pachodynerus* Saussure, 1870 (Hymenoptera: Vespidae: Eumeninae) en Chile. Bol Biodivers Chile 2010; 4: 94-8.

9. Freeman B, Jayasingh D. Population dynamics of *Pachodynerus nasidens* (Hymenoptera) in Jamaica. Oikos. 1975; 26: 86-91.

10. Bequaert JC. The genus *Pachodynerus* (Hymenoptera, Vespidae) in the Antilles. Psyche. 1948; 55, 105-12.

11. Buschini M, Buss C. Biologic aspects of different species of *Pachodynerus* (Hymenoptera; Vespidae; Eumeninae). Braz J Biol. 2010; 70: 623-29.

12. Bureau d'Enquêtes et d'Analyses pour la sécurité de l'aviation civile (BEA). Final Report On the accident on 1st June 2009 to the Airbus A330-203 registered F-GZCP operated by Air France flight AF 447 Rio de Janeiro – Paris. Bureau d'Enquêtes et d'Analyses pour la sécurité de l'aviation civile. Le Bourget, France; 2012.

13. Boeing. Erroneous: flight instrument information – safety and guidance. Aero (Boeing). 2003; 29.

14. Civil Aviation Safety Authority. Wasp infestation – Alert: Civil Aviation Safety Authority Canberra; 2018. Airworthiness Bulletin AWB 02-052 Issue 4-3 May 2018.

15. Australian Transport Safety Bureau. Rejected takeoff – Brisbane Airport, Qld – 19 March 2006 – VN-QPB, Airbus A330-303: Australian Transport Safety Bureau, Canberra; 2008. Report no. 200601453.

16. Australian Transport Safety Bureau 2016. Air data system failure involving Airbus A330-243, A6-EYJ, near Brisbane Airport, Qld on 21 November 2013: Australian Transport Safety Bureau, Canberra; 2016. Aviation Occurrence Investigation AO-2013-212.

17. Australian Transport Safety Bureau. Australian aviation wildlife strike statistics 2008–2017. Australian Transport Safety Bureau, Canberra; 2018.

18. National Transportation Safety Bureau. Aircraft Accident Report. Florida Commuter Airlines Inc. Douglas DC-3, N75KW Grand Bahama Island, Bahamas September 12, 1980: National Transportation Safety Board, Washington DC; 1981.

19. Junta Investigadora de Accidentes Aéreos. Final Aviation Accident report: Birgenair Flight ALW-301. Junta Investigadora de Accidentes Aéreos; 1996. Director General of Civil Aeronautics, Dominican Republic.

20. Bureau of Meteorology. Climate data: Bureau of Meteorology, Canberra. Available from http://www.bom.gov.au/climate/data/.

21. Oliveira Nascimento AL, Garófalo CA. Trap-nesting solitary wasps (Hymenoptera: Aculeata) in an insular landscape: Mortality rates for immature wasps, parasitism, and sex ratios. Sociobiology. 2014; 61: 207-17.

22. Wilke R., Bazilio S. & Homczinski I. 2013. Diversidade, sazonalidade e aspectos biológicos de vespas solitárias em ninhos armadilhas na região de São Miguel da Serra, Porto União − SC. (Diversity, seasonality and biological aspects of solitary wasps in nest traps in the region of Sao Miguel da Serra, Porto Uniao). In: Krupek, RA (ed.) Teaching & Research No. 11, Vol. 1, University of Paraná, Brazil.

23. Jayasingh DB, Taffe CA. The biology of the eumenid mud-wasp *Pachodynerus nasidens* in trapnests. Ecol Entomol. 1982; 27: 283-89.

24. Silva JF. Vespas solitárias (Hymenoptera: Aculeata) nidificando em ninhos-armadilha na Estação Ecológica de Ribeirão Preto, SP (Solitary wasps (Hymenoptera: Aculeata) nesting in trap nests at the Ecological Station of

Ribeirão Preto, SP) [MSc thesis]. Faculty of Philosophy, Sciences and Letters of Ribeirão Preto: University of São Paulo; 2008.

25. Buschini MLT. Life-history and sex allocation in *Trypoxylon* (syn. *Trypargilum*) *lactitarse* (Hymenoptera; Crabronidae). J Zool Syst Evol Res. 2007; 45: 206-13.

26. Polidori C, Boesi R, Borsato W. Few, small, and male: Multiple effects of reduced nest space on the offspring of the solitary wasp, *Euodynerus* (*Pareuodynerus*) *posticus* (Hymenoptera: Vespidae). C R Biol. 2011; 334: 50-60.

27. Fernandez-Triana JL, Gonzalez GG, Ferrer EP, Ruiz, AS. Ecological relationships in nests of six Cuban aculeate wasps (Hymenoptera: Vespidae, Sphecidae, Apidae). Bol SEA. 2005; 36 115-18.

28. Assis JMF, Camillo E. Diversidade, sazonalidade e aspectos biológicos de vespas solitárias (Hymenoptera: Sphecidae: Vespidae) em ninhos armadilhas na região de Ituiutaba, MG (Diversity, seasonality and nesting biology of solitary wasps (Hymenoptera: Sphecidae: Vespidae) in trap-nests in the Ituiutaba region, MG). An Soc Entomol Bras. 1997; 26: 335-47.

29. Emond J-P, Mercier F, Nunes MCN. In-flight temperature conditions in the holds of a widebody aircraft. Paper 281, 20th International Congress of Refrigeration; 1999. International Institute of Refrigeration, Sydney.

30. Perry G. Wheel-well and cargo compartment temperatures of large aircraft in flight: implications for stowaways. Aviat Space Environ Med. 2002; 73: 673-76.

Figure Legends

Figure 1. Location of pitot probe panels at Brisbane Airport.

Figure 2. Pitot panels installed on gate light pole at Domestic Terminal (left) and on

aerobridge at International Terminal (right).

Figure 3. Pitot probe blockages by mud-nesting wasps and rainfall at Brisbane Airport,

February 2016 – April 2019.

Figure 4. Relationship between pitot probe blockages and minimum distance to (a)

natural horizontal habitat and (b) natural vertical habitat.

Figure 5. Mean number of days (from collection of probes) to emergence of live wasps,

February 2016 – April 2019.

Figure 6. Relationship between pitot probe volume and sex of emerging live adult

wasps. Second order polynomial line of best fit fitted. $r^2_{male} = 0.942$ ($p < 0.05$), r^2_{female}

$= 0.517$ (ns).

Figures

Figure 1

Figure 2

Figure 3

551   Figure 4

554   Figure 5

557   Figure 6