In vitro pancreatic islet cluster expansion facilitated by hormones and chemicals

Jing-Yu Lin[#1], Jie Cheng[#1], Ya-Qin Du[#2], Wei Pan[#1], Zhong Zhang[3], Jin Wang[4], Jie An[4], Fan

Yang[4], Yun-Fei Xu[6], Hui Lin[1], Wen-Tao An[1], Jia Wang[5], Zhao Yang[5], Ren-Jie Chai[3], Hui-Li

Hu*[7], Jin-Peng Sun*[2,5] and Xiao Yu*[1]

Author details

[1]Key Laboratory Experimental Teratology of the Ministry of Education and Department of

Physiology, School of Basic Medical Sciences, Shandong University, 250012 Jinan, China

[2]Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Peking

University, Key Laboratory of Molecular Cardiovascular Science, Ministry of Education,

100191 Beijing, China

[3]Key Laboratory for Developmental Genes and Human Disease of the Ministry of Education,

School of Life Science and Technology, Southeast University, 210096 Nanjing, China

[4]Department of Pharmacology, School of Basic Medical Sciences, Shandong University,

250012 Jinan, China

[5]Key Laboratory Experimental Teratology of the Ministry of Education and Department of

Biochemistry and Molecular Biology, School of Basic Medical Sciences, Shandong University,

250012 Jinan, China

[6]Key Laboratory Experimental Teratology of the Ministry of Education and Department of

Biochemistry and Molecular Biology, Shandong University School of Medicine, 250012 Jinan,

China

[7]The Key Laboratory of Experimental Teratology, Ministry of Education and Department of

Genetics, School of Basic Medical Sciences, Shandong University, 250012 Jinan, China

[#]These authors contributed equally to this work

*Corresponding author: Xiao Yu (corresponding author)

E-mail: yuxiao@sdu.edu.cn

Jin-Peng Sun (corresponding author)

E-mail: sunjinpeng@bjmu.edu.cn

Hui-Li Hu (corresponding author)

E-mail: huhuili@sdu.edu.cn

Abstract

Tissue regeneration, such as pancreatic islet tissue propagation in vitro, could serve as a

promising strategy for diabetes therapy and personalized drug testing. However, such a protocol has not been realized yet. Propagation could be divided by expansion in vitro and passaging. Even the in vitro expansion of the islet has not been achieved to date. Here, we describe a method to enable the expansion of islet clusters isolated from pregnant mice or wild-type rats by employing a combination of specific regeneration factors and chemical compounds in vitro. The expanded islet clusters expressed insulin, glucagon and somatostatin, which are markers corresponding to pancreatic β cells, α cells and δ cells, respectively. These different types of cells grouped together, were spatially organized and functioned similarly to primary islets. Further mechanistic analysis revealed that forskolin in our recipe contributed to renewal and regeneration, whereas exendin4 was essential for preserving islet cell identity. Our results provide a novel method for the in vitro expansion of islet clusters, which is an important step forward in developing future protocols and medium used for islet tissue propagation in vitro. Such method is important for future regenerative diabetes therapies and personalized medicine using large amounts of pancreatic islets derived from the same person.

Introduction

Diabetes is characterized by insulin resistance and dysfunctional insulin secretion. Recent progress in the generation of functional insulin-producing pancreatic β cells from stem cells differentiation or transdifferentiation in vitro has shed light on cell-replacement therapies for the treatment of diabetes[1-4]. However, compared to tissue/organ replacement therapy derived from somatic islets of the same person for curing diabetes[5,6], these methods still have the following deficits: (1) The function of a tissue, such as pancreatic islets, cannot be fully substituted by a single β cell type. Islets contain at least three main cell types, and together, they contribute the functional integrity adaptations to responses to different physiological changes[7-10]. In fact, recent reports indicate the essential roles of the islet circuit in maintaining glucose metabolism homeostasis[7,10]. (2) The identity of the β cells generated from transdifferentiation or stem cells is not defined, and their functions are not guaranteed. These cells normally have different gene expression patterns or certain mutations compared to somatic β cells. (3) Some of the β-like cells generated from stem cells may still have pluripotency potential, which may correlate to a risk of tumour development. (4) Finally, these cells may elicit unwanted immune responses.

Ideally, propagating islets isolated from the same patient to the desired number and then transplanting them into the patient is a promising regenerative strategy for diabetes therapy with

benefits that stem cell or transdifferentiation based cell-replacement treatment cannot match.

This method will minimize the immune response, decrease the identity difference and result in

a better ability to adapt to different growth niche after transplantation. In addition, the organoids

generated from pancreatic islets propagated from patients can be used for drug testing because

they mostly modelling composition, architecture and function of primary tissue. Recently,

significant progress has been made in which somatic intestinal, gastric, colonal, hepatic and

pancreatic ducts were cultured and propagated to organoids in vitro[11-19], which lays the

foundation for future regenerative therapy. However, these organoids mostly exhibit epithelial

properties and are difficult to be induced into endocrine cells. Thus, the in vitro propagation of

functional endocrine organoids especially islets, is still required to be established.

Propagation of islets could be divided by two processes: 1) expansion of the islet cells in

vitro. 2) Propagating the expanded islet cells by multiple passages. Here, we reported a method

for 3D culture with a novel special medium recipe that enabled the in vitro expansion of

pancreatic islet clusters isolated from pregnant mice or wild-type rats. Further mechanistic

analysis suggested that forskolin (FSK) in the recipe is required for maintaining cell renewal

and regeneration, whereas exendin4 is essential for preserving islet identity. Therefore, our

results identified a method allowed an in vitro expansion of the pancreatic islet clusters, which

could be a key step for propagating patient-specific pancreatic islets for future regenerative

diabetes therapy.

Results

Recipe for the in vitro pancreatic islet expansion medium (PIEM)

To identify the optimal conditions for pancreatic islet expansion in vitro, we reviewed the

chemicals, proliferation and regeneration factors used in the culture of other organoids, such as

those in pancreatic ducts, cholangiocytes and hepatocytes [14,16,20,21], and pancreatic islet

organoids derived from stem cells or fibroblasts (Figure 1) [1-4]. According to their reported

functions, these chemicals and factors are classified into the following groups: islet identity,

cell division and proliferation, and cell renewal and regeneration (Figure 1). By screening

different combinations of compounds and factors, a medium composed of 12 chemicals and

hormones, as well as the general nutrient nicotinamide, B27 and GlutaMAX, was created and

was found to robustly support the in vitro expansion of dispersed islet clusters isolated from

pregnant mice or wild-type rats. We named this in vitro pancreatic islet expansion medium

PIEM.

Expansion of mouse islet clusters in vitro

Previous studies have shown that pregnant mouse islets maintain expansion ability in vivo,

whereas wild-type mouse islets undergo little expansion[22,23]. We therefore used islets from both

pregnant and wild-type mice, as pregnant mice could serve as an easier starting point for

investigating the regeneration process. We isolated pancreatic islets from both wild-type and

pregnant C57BL/6 mice by collagenase P perfusion, sedimentation and handpicking. These

pancreatic islets were separated into single cells or cell clusters by dispase II digestion and

mechanical dissociation. The single cells or cell clusters were cultured in 3-dimensional (3D)

Matrigel. Only dispersed islet clusters isolated from pregnant mice showed significant

expansion during nine-day culture (Figure 2A). Approximately 5~15% of the pancreatic islet

clusters isolated from pregnant mice were able to expand from a surface area of 6000-9000 μm^2

to a surface area of 20000-30000 μm^2 (Figure 2A and 2C). In contrast, the dispersed single islet

cells from pregnant or wild-type mice or islet clusters from wild-type mice showed no

significant expansion (Figure 2B and 2D).

We next used qRT-PCR to examine the expression of key islet markers in the expanded islet

clusters. The expanded islet clusters maintain expression of the crucial pancreatic β cell markers

insulin-1 (Ins1), insulin-2 (Ins2) and *Mafa*; the α cell markers *glucagon (Gcg)* and *Mafb*; and

the δ cell marker *somatostatin (Sst)* (Figure 2E).

To confirm that specific cell types coexisted in the in vitro expanded islet clusters, which is

a hallmark of expanded islet clusters, we performed immunostaining for insulin, glucagon and

somatostatin, which are markers for pancreatic α cells, β cells and δ cells, respectively. The

expanded islet clusters in Matrigel were too tiny, which caused great difficulties in slice

preparation for immunostaining. We therefore immunostained the whole clusters as described

in the material and methods section. The result unambiguously identified that glucagon, insulin,

and somatostatin staining was present in different cells grouped together in the same expanded

islet clusters (Figure 3A and 3B).

We next examined the functional integrity of the expanded islet clusters. In response to

combined stimulation with 20 mM glucose and 100 nM GLP-1, the expanded islet clusters

demonstrated a significant increase in insulin secretion (Figure 3C). In response to high glucose

stimulation, both other labs and our lab have shown that UCN3 serves an endogenous paracrine

factor secreted by pancreatic β cells to stimulate endogenous somatostatin secretion from

pancreatic δ cells[7,10,24]. We therefore stimulated the expanded islet clusters with both glucose

and UCN3 to minimize the amount required for islet cluster usage. The in vitro expanded islet

clusters displayed significantly more somatostatin release in response to glucose and UCN3

stimulation (Figure 3D). Moreover, the expanded islet clusters secreted more glucagon in

response to 25mM Arginine stimulation, hallmarking the normal function of pancreatic islet α

cells (Figure 3E). These results confirmed the functional integrity of the in vitro expanded islet

clusters isolated from pregnant mice.

Although the expanded pancreatic islet clusters showed similar SST and Glucagon secretion

compared to the primary isolated islets in response to specific physiological stimulations, it is

worth to note that their insulin secretion in response to the combined stimulation of glucose and

GLP-1 is significantly weakened. Some extent degeneration during the islet expansion may

occur during current in vitro expansion condition, which awaits for further investigation and

methods optimization in our future work.

Expansion of rat islet clusters in vitro

We then examined the expansion of the dispersed rat islet single cells and clusters in PIEM.

Similar to those from pregnant mice, approximately 5~15% of the rat pancreatic islet clusters

were able to expand from a surface area of 5000-8000 μm^2 to a surface area of 20000-25000

μm^2, whereas dispersed rat islet single cells showed no such expansion ability (Figure 4A and

4B). Furthermore, qRT-PCR revealed that the expanded rat islet clusters maintained the same

expression of the α cell markers *Gcg* and *Mafb* as the primary rat islets but showed significantly

higher expression of the β cell markers *Ins-1, Ins-2* and *Mafa* and the δ cell marker *Sst* (Figure

4C).

Increased gene expression related to dedifferentiation, pluripotency and proliferation in

expanded islet clusters

We next examined whether the expanded islet clusters increased the expression level of genes

functionally associated with cell proliferation, renewal and regeneration, which are important

factors for in vitro regeneration. Importantly, significantly higher expression of *Ki67*, *Ccnd1*

and *Pcna* was found in expanded islet clusters than in islets derived from pregnant mice or

wild-type rats, highlighting the increased proliferation ability of the expanded islet clusters

(Figure 5A and 5D). Moreover, significantly higher *Nanog* and *Sox9* expression were found in

expanded islet clusters derived from both pregnant mice and rats (Figure 5B and 5E). These

results provide putative explanations accounting for the better proliferation and renewal

abilities of isolated islet clusters than integral islets.

We suspected that these islets gained proliferation and pluripotency due to dedifferentiation.

Thus, we examined dedifferentiation markers in expanded islet clusters and compared them

with those in isolated pancreatic islets. Interestingly, the expanded islet clusters showed

significantly higher expression of *Ngn3* and *Hlxb9*, markers that characterize pancreatic islet

progenitor cells (Figure 5C and 5F). Although the expanded islet clusters derived from pregnant

mice showed higher expression of *Gata6*, they showed decreased expression of *Gata4*, both of

which are pancreatic progenitor markers (Figure 5C and 5F). Similarly, the expanded islet

clusters derived from rats did not display significant pancreatic progenitor characteristics, as

they showed decreased expression of *Gata6* (Figure 5C and 5F). The data suggested that these

expanded islet clusters underwent one-step dedifferentiation towards islet progenitor cells, but

they did not ultimately reach the pancreatic progenitor cell stage. Taken together, this one-step

dedifferentiation of isolated islet clusters cultured in PIEM towards pancreatic islet progenitor

cells contributed to the gain of cell proliferation, renewal and regeneration functions of

expanded islet clusters.

Essential role of FSK and exendin4 in PIEM

During our formula component screening and recipe formation for PIEM, we identified that

FSK and exendin4 are both required for robust islet cluster expansion in vitro. It is known that

exendin4 induced cAMP accumulation after it activates GLP-1R and downstream Gs proteins[25].

FSK is known as a cAMP agonist through direct binding to adenyl cyclase[26]. Moreover, recent

reports indicate that FSK is required for proliferation of liver organoids in addition to A83-01[27].

Interestingly, removing FSK or exendin4 from PIEM has different effects on the gene

expression profiles of islet identity, proliferation, renewal and regeneration markers. Removing

FSK from PIEM had partial effects on islet identity and cell proliferation markers, including

decreased expression of *Mafa*, *Sst* and *Ccnb1* (Figure 6A and 6B). In particular, FSK was

required for the expression of cell renewal and regeneration markers, including *Nanog* and

Sox17 (Figure 6C). In contrast, exendin4 was essential for islet identity and proliferation

because it maintained the expression of the pancreatic β cell markers *Ins-1*, *Pdx1* and *Mafa* and

the proliferation markers *Ki67*, *Ccnb1* and *Cdk4* (Figure 6A and 6B). In particular, exendin4

was required for the expression of *Ngn3* and *Hxlb9*, two islet progenitor markers (Figure 6D).

These results indicated that FSK in PIEM renders cell renewal and regeneration ability to

isolated islet clusters, whereas exendin4 is essential for maintaining islet β cell identity,

proliferation and dedifferentiation.

Discussion

The in vitro expansion of primary pancreatic β cells or pancreatic islet tissues has not been

reported before. Here, we developed a medium recipe, which we named PIEM, that enabled the

proliferation of dispersed islet clusters isolated from pregnant mice or wild-type rats in vitro

(Figure 7). Isolated islet clusters normally grow from an initial surface area of 5000-9000 μm^2

to a surface area of 20000-30000 μm^2, with an estimated approximately 5-fold increase in cell

number. Compared to primary pancreatic islets, expanded cell clusters have similar or higher

mRNA expression levels of insulin, glucagon and somatostatin, which are markers of

pancreatic β cells, α cells and δ cells, respectively. Expanded islet clusters demonstrated normal insulin and somatostatin secretion in response to physiological stimulation and normal glucagon secretion in response to amino acids, suggesting that the clusters recapitulate specific functions of pancreatic islets. Immunostaining further confirmed that the cells expressing insulin, glucagon and somatostatin were grouped together and spatially organized in expanded islet clusters similar to primary islets. These data indicated that the expanded islet clusters behaves similar to an in vitro organoid, thus serving as an initial step for future islet propagation methods developing in vitro. The ultimate goal of developing efficient methods for propagation islets in vitro is to fulfil the demand for regenerative diabetes therapy and personalized drug tests.

It is worth noting that we observed the expansion of islet clusters isolated from only pregnant mice and wild-type rats in PIEM, not those isolated from wild-type mice. Pregnant mice are well known for their increased pancreatic β cell proliferation[28-31], which provides a useful starting point for testing medium recipes. Although we did not observe the proliferation of islet clusters isolated from wild-type mice, we did observe the expansion of islet clusters isolated from wild-type rats, which share more properties identical to human islets than mouse islets. An urgent need is to test whether PIEM enables the expansion of primary islet clusters isolated from human patients.

Another notable observation is that only dispersed islet clusters were able to proliferate in vitro, whereas dispersed single primary islet cells could not. Two important factors may contribute to this discrepancy. First, the islet clusters contain multiple cell types, which are not only the origins of the different cell types in the finally expanded islet clusters but also may form certain cell circuits and gradient hormone concentrations to support in vitro expansion. Second, primary single cells must undergo harsher digestion than islet clusters, which means that the extracellular parts of the membrane or matrix proteins, including receptors, ion channels and transporters, may be digested and their function impaired. These membrane or matrix proteins may be key for in vitro tissue expansion. Therefore, future analysis of the different expression patterns and functions of membrane proteins and signalling circuit differences between isolated islet clusters and single cells may provide more clues for unravelling the secrets of the in vitro expansion of islets and provide guidance for a better strategy for in vitro islet cluster expansion.

We noticed that expanded islet clusters have higher expression of *Ngn3* and *Hlxb9*, which are markers of pancreatic islet progenitor cells. This observation indicated that in vitro expanded islets underwent certain dedifferentiation. However, we doubt that the islet clusters gained proliferation ability through this dedifferentiation. Further experiments with withdrawal of key chemicals in the medium will test this hypothesis. For clinical usage, the in vitro expansion of 100 patient islets to 1 million islets without significant gene mutations, while

preserving pancreatic islet function, and transferring them back to the patient to recover glucose homeostasis is ideal. However, the current approach of PIEM allows only approximately 5-fold growth of approximately 5%-15% of isolated islet clusters. The expansion of the islet clusters was only observed in the first passage. Therefore, it is unlikely to cause significant genomic instability. However, the current acquired in vitro expanded islet cluster cells do not contain sufficient amounts for RNA-seq to verify the gene expression profile or transplantation to confirm their in vivo functions. More important, the final usage of in vitro propagated islet clusters for regenerative therapy to treat diabetes in general require millions of pancreatic islets. Therefore, a future revision of the current PIEM composition and method for in vitro islet propagation is in an urgent demand.

One of the future directions for the optimization of in vitro islet propagation methods could be as follows: (1) initial expansion and dedifferentiation of islet clusters to islet progenitor-like cell clusters, (2) rapid proliferation of islet progenitor-like cell clusters to a sufficient amount, (3) differentiation of progenitor-like cell clusters to functional integral islets, (4) transplantation or (5) exploitation of these expanded islet tissues for drug tests. Therefore, manifesting the markers for each stage and searching chemicals that are useful for each procedure in the above protocol is our next goal to fulfil expanded islet clusters regeneration in vitro in the future.

Taken together, we have developed a recipe for PIEM, and our 3D culture method enabled the initial generation of expanded islet clusters via the in vitro expansion of isolated islet clusters from pregnant mice or wild-type rats in the first passage. Our study is one important step forward for the realization of a method to fulfil in vitro propagation of pancreatic islet in vitro, which may facilitate new therapies development to treat diabetes.

Materials and methods

Key resources table

Reagent	Source	Identifier
Antibodies		
Anti-Insulin	Santa	sc-9168
Anti-Glucagon	Sigma	G2654
Anti-Somatostatin	Abcom	ab30788
Donkey anti-Mouse IgG (H+L) (secondary antibody)	Invitrogen	A10037
Donkey anti- Rabbit IgG (H+L) (secondary antibody)	Invitrogen	A21206

Donkey anti-Rat IgG(H + L) (secondary antibody)	Abcam	ab150154

Chemicals, Peptides, and Recombinant Proteins

TRIzol	Thermo	15596-026
Cell Recovery Solution	Corning	354253
Matrigel	Corning	356231
GlutaMax	Thermo	35050-061
Penicillin-Streptomycin	Biological Industries	03-031-1B
recombinant human EGF	Peprotech	AF-100-15
recombinant human FGF10	Peprotech	100-26
CHIR-99021	BioGems	2520691-1MG
N-Acetylcysteine	Sigma-Aldrich	A9165-25G
Nicotinamide	Sigma-Aldrich	N0636
B27 Supplement (minus Vitamin A)	Thermo	12587-010
gastrin-1, human	MedChemExpress	HY-P1097
A83-01	Adooq Bioscience	A12358
Y-27632	Adooq Bioscience	A11001
Forskolin	TargetMol	T2939
Exendin4	ChinaPeptides	
5-Iodotubercidin	Adooq Bioscience	A13948
L-Ascorbic acid	Sigma-Aldrich	A4544
Triton X-100	Solarbio	T8200
Donkey Serum	Solarbio	SL050
DAPI	Beyotime	C1002

Critical Commercial Assays

Glucagon ELISA Kit	Cloud-Clone Corp	CEB266Mu
Insulin ELISA Kit	merck-millipore	EZRMI-13K
Somatostatin EIA Kit	Phoenix Pharmaceuticals	EK-060-03

Software and Algorithms

Prism Graphpad		https://www.graphpad.com
Adobe illustator	Adobe	http://www.adobe.com/products/illustrator.html
ImageJ		https://imagej.nih.gov/ij/

Experimental Models: Animals/Cell Lines

C57B6	Shandong University animals care facility	
Wistar	Shandong University animals care facility	

Oligonucleotides

Primer: Gcg Forward for mouse cDNA: GGCACATTCACCAGCGACTA	This paper	N/A
Primer: Gcg Reverse for mouse cDNA: GAGAAGGAGCCATCAGCGTG	This paper	N/A
Primer: Ins1 Forward for mouse cDNA: TGGACTATAAAGCTGGTGGGC	This paper	N/A
Primer: Ins1 Reverse for mouse cDNA: TGTGTAGAAGAAGCCACGCT	This paper	N/A
Primer: Ins2 Forward for mouse cDNA: GCTATCCTCAACCCAGCCTAT	This paper	N/A
Primer: Ins2 Reverse for mouse cDNA: CTCCAGTGCCAAGGTCTGAA	This paper	N/A
Primer: Mafa Forward for mouse cDNA: GGCACATTCTGGAGAGCGAG	This paper	N/A
Primer: Mafa Reverse for mouse cDNA: TCACAGAAAGAAGTCGGGTGC	This paper	N/A
Primer: Mafb Forward for mouse cDNA: TACCAGCAGATGAACCCCGA	This paper	N/A
Primer: Mafb Reverse for mouse cDNA: GCTAGTGGGTAGCTGTTGCG	This paper	N/A
Primer: Sst Forward for mouse cDNA: CCACCGGGAAACAGGAACTG	This paper	N/A
Primer: Sst Reverse for mouse cDNA: TTGCTGGGTTCGAGTTGGC	This paper	N/A
Primer: Ngn3 Forward for mouse cDNA: ACGCAATTTACTCCAGGCGA	This paper	N/A
Primer: Ngn3 Reverse for mouse cDNA: GAGGCGCCATCCTAGTTCTC	This paper	N/A
Primer: Hlxb9 Forward for mouse cDNA: GAACCTCTTGGGGAAGTGCC	This paper	N/A
Primer: Hlxb9 Reverse for mouse cDNA:	This paper	N/A

TCTTTGGCCTTTTTGCTGCG		
Primer: Gata4 Forward for mouse cDNA: AGATCGCGCCGGTTTTCTG	This paper	N/A
Primer: Gata4 Reverse for mouse cDNA: GATCACCCACCGGCTAAAGA	This paper	N/A
Primer: Gata6 Forward for mouse cDNA: TTTCCGGCAGAGCAGTAAGAG	This paper	N/A
Primer: Gata6 Reverse for mouse cDNA: GAAACGCTTTGGCAGGCAC	This paper	N/A
Primer: Sox9 Forward for mouse cDNA: CGGAACAGACTCACATCTCTCC	This paper	N/A
Primer: Sox9 Reverse for mouse cDNA: GCTTGCACGTCGGTTTTGG	This paper	N/A
Primer: Ki67 Forward for mouse cDNA: CAAGGCGAGCCTCAAGAGATA	This paper	N/A
Primer: Ki67 Reverse for mouse cDNA: TGTGCTGTTCTACATGCCCTG	This paper	N/A
Primer: Nanog Forward for mouse cDNA: AGGACAGGTTTCAGAAGCAGA	This paper	N/A
Primer: Nanog Reverse for mouse cDNA: CCATTGCTAGTCTTCAACCACTG	This paper	N/A
Primer: Ccnb1 Forward for mouse cDNA: GCGTGTGCCTGTGACAGTTA	This paper	N/A
Primer: Ccnb1 Reverse for mouse cDNA: CCTAGCGTTTTTGCTTCCCTT	This paper	N/A
Primer: Ccnd1 Forward for mouse cDNA: TGACTGCCGAGAAGTTGTGC	This paper	N/A
Primer: Ccnd1 Reverse for mouse cDNA: CTCATCCGCCTCTGGCATT	This paper	N/A
Primer: Pcna Forward for mouse cDNA: TTGCACGTATATGCCGAGACC	This paper	N/A
Primer: Pcna Reverse for mouse cDNA: GGTGAACAGGCTCATTCATCTCT	This paper	N/A
Primer: Pdx1 Forward for mouse cDNA: CCTTTCCCGAATGGAACCGA	This paper	N/A
Primer: Pdx1 Reverse for mouse cDNA: GGGCCGGGAGATGTATTTGT	This paper	N/A
Primer: Sox17 Forward for mouse cDNA: CCAAAGCGGAGTCTCGCAT	This paper	N/A
Primer: Sox17 Reverse for mouse cDNA: GCCTAGCATCTTGCTTAGCTC	This paper	N/A
Primer: Cdk4 Forward for mouse cDNA: GCCACTCGATATGAACCCGT	This paper	N/A

Primer: Cdk4 Reverse for mouse cDNA: CACAGACATCCATCAGCCGT	This paper	N/A
Primer: Gcg Forward for rat cDNA: TTCACAGGGCACATTCACCA	This paper	N/A
Primer: Gcg Reverse for rat cDNA: CTATGGCGACTTCTTCCGGG	This paper	N/A
Primer: Insulin1 Forward for rat cDNA: ACCCTAAGTGACCAGCTACAATC	This paper	N/A
Primer: Insulin1 Reverse for rat cDNA: CGGGTCCTCCACTTCACGAC	This paper	N/A
Primer: Insulin2 Forward for rat cDNA: ACCTTTGTGGTTCTCACTTGGT	This paper	N/A
Primer: Insulin2 Reverse for rat cDNA: CAGTGCCAAGGTCTGAAGGTCA	This paper	N/A
Primer: Mafa Forward for rat cDNA: GACAAGTTTGCGCAGGCCG	This paper	N/A
Primer: Mafa Reverse for rat cDNA: GTATTCACCGTTCTCGGGGC	This paper	N/A
Primer: Mafb Forward for rat cDNA: GCAACGGTAGTGTGGAGGAC	This paper	N/A
Primer: Mafb Reverse for rat cDNA: GAGCTGCGTCTTCTCGTTCT	This paper	N/A
Primer: Sst Forward for rat cDNA: GCTACTGGAGTCGTCTCTGC	This paper	N/A
Primer: Sst Reverse for rat cDNA: GGCATCGTTCTCTGTCTGGT	This paper	N/A
Primer: Ngn3 Forward for rat cDNA: GTTCCAATTCCACCCCACCT	This paper	N/A
Primer: Ngn3 Reverse for rat cDNA: CGCAGGGTCTCGATCTTTGT	This paper	N/A
Primer: Hlxb9 Forward for rat cDNA: CGGCGCTTTCCTACTCGTAT	This paper	N/A
Primer: Hlxb9 Reverse for rat cDNA: TCCCCAAGAGGTTCGATTGC	This paper	N/A
Primer: Gata4 Forward for rat cDNA: TGAATGGTATCAACCGGCCC	This paper	N/A
Primer: Gata4 Reverse for rat cDNA: TTTGAATCCCCTCCTTCCGC	This paper	N/A
Primer: Gata6 Forward for rat cDNA: TCATCACCACCCGACCTACT	This paper	N/A
Primer: Gata6 Reverse for rat cDNA: GCATGCGTTGCACAGGTAAT	This paper	N/A
Primer: Sox9 Forward for rat cDNA: CACAAGAAAGACCACCCCGA	This paper	N/A

Primer: Sox9 Reverse for rat cDNA: TGCACGTCTGTTTTGGGAGT	This paper	N/A
Primer: Ki67 Forward for rat cDNA: ACAGGGCTTAGGAAACAGTCC	This paper	N/A
Primer: Ki67 Reverse for rat cDNA: GGTTCTAACTGGTCTTCCTGGT	This paper	N/A
Primer: Nanog Forward for rat cDNA: AAGTCCCTTCCCTTGCCGT	This paper	N/A
Primer: Nanog Reverse for rat cDNA: CTCGGGACCAGACAGCTTTAG	This paper	N/A
Primer: Ccnb1 Forward for rat cDNA: GGGTGTCTTCTCAGATCGGC	This paper	N/A
Primer: Ccnb1 Reverse for rat cDNA: TCCACAGGTTTTGGTAGGGC	This paper	N/A
Primer: Ccnd1 Forward for rat cDNA: TCAAGTGTGACCCGGACTG	This paper	N/A
Primer: Ccnd1 Reverse for rat cDNA: CTACTTGGTGACTCCCGCCT	This paper	N/A
Primer: Pcna Forward for rat cDNA: GACGGGGTGAAGTTTTCTGC	This paper	N/A
Primer: Pcna Reverse for rat cDNA: GACAGTGGAGTGGCTTTTGTG	This paper	N/A

Animals

C57BL/6 mice and rats of Wistar strain were housed and bred under specific-pathogen-free conditions at Shandong University animals care facility. Female animals were used. The numbers of animals are indicated in the figure legends within each experiment. All animal care and experiments were reviewed and approved by the Animal Use Committee of Shandong University School of Medicine.

Pancreatic islet isolation and digestion into single cells or cell clusters.

We isolated pancreatic islets from C57BL/6 mice, pregnant C57BL/6 mice and Wistar rat. Briefly, mice and rat were killed by cervical dislocation and then pancreas were isolated from them individually. Adult pancreas were digested by Collagenase P (Roche, 11213873001) at 37℃ for 18-25 minutes. Digestion was stopped by cold Hank's balanced salt solution (136.9 mM NaCl; 5.4 mM KCl; 1.3 mM $CaCl_2$; 0.8 mM $MgSO_4$; 0.44 mM KH_2PO_4; 0.34 mM Na_2HPO_4; 5.55 mM D-glucose; 4.4 mM $NaHCO_3$, pH=7.4) followed by sedimentation for three

or five times at 4°C. The islets were collected by hand picking using a stereomicroscope and were cultured overnight in islet complete media containing 5.6 mM glucose (Biological Industries), 10% FBS (Gibco),0.1% penicillin/streptomycin.Next we obtained single cells and cell clusters of different diameters (20-150μm)under the microscope by controlling enzyme concentration, digestion and settling time. Islets briefly after naturally settling for 1 minute in KRBB buffer (135 mM NaCl; 4.7 mM KCl; 1.2 mM KH_2PO_4; 10 mM Hepes; 3 mM D-glucose; 5 mM $NaHCO_3$; 0.1% BSA; 1% penicillin/streptomycin, pH=7.4), preheated EDTA-KRBB solution was added, gently piped for 2 min, then placed in a 33°C water bath for 10 min; after clearing the supernatant, digested with dispase II(0.1-0.5 U/ml, Roche, 04942078001) in a 33°C water bath for 3~8 min to obtain single cells or islet cell clusters. Mechanical fragmentation was achieved by gentle pipetting (5–10 times) with a glass dropper in a 15ml glass centrifuge tube. Using DMEM/F-12 (Biological Industries) medium with 10% FBS (Gibco) and 1% penicillin/streptomycinfor termination of digestion.Collecting single cells or cell clusters by centrifugation (300×g, 5 min, Room temperature).

Production of Rspo1-conditioned medium

The RSPO1 conditioned medium is home-made and will be described in other papers.

Expanded islet clusters culture

Isolated islet cell clusters were washed twice with DMEM/F-12 (Biological Industries), counted and mixed with Matrigel in glass centrifuge tubes. 30,000–50,000 cells or 10-20 cell clusters were used per well of a pre-warmed 24-well plate (the volume of 50 μl). After Matrigel was solidified for 30min at 37°C, pancreatic islet expansion medium (PIEM) was added. PIEM consists of DMEM/F-12 (1% GlutaMax,1% Penicillin-Streptomycin) plus 15% RSPO1 conditioned medium (home-made), 3 μM CHIR-99021 (BioGems), 1 μM A83-01 (Adooq Bioscience), 10 nM gastrin-1 (MedChemExpress), 1.25 mM N-acetylcysteine (Sigma), 10 μM Y-27632 (Adooq Bioscience), 10 μM forskolin (TargetMol), 50 ng/ml Exendin-4 (ChinaPeptides), 50 μg/ml L-Ascorbic acid (Sigma), 250 nM 5-Iodotubercidin (Adooq Bioscience), 50 ng/ml FGF10 (PeproTech), 50 ng/ml EGF (PeproTech), 10 mM Nicotinamide (Sigma), B27 Supplement (minus Vitamin A) (Thermo). Cultures were kept at 37°C, 5% CO_2 in a humidified incubator. During culturing, medium was refreshed every three days.

RNA extraction and qRT-PCR.

RNA from islets and islet single cells as well as islet cell clusters derived from C57BL/6 mice, pregnant C57BL/6 mice and Wistar rat, Including the corresponding islet clusters were

extracted with TRIzol reagent (Thermo). We used the tip to pick up the expanded islet clusters

from the Matrigel using a stereomicroscope. Cell Recovery Solution(Corning) is used to

recover cells from Matrigel. Add 0.2-0.5 ml per 24-well plate cold Cell Recovery Solution.

Scrape the gel layer into an cold 1.5 ml Microcentrifuge Tube and leave it on ice for 0.5-1 hour

until the gel complete dissolved. Collecting cells by centrifugation (200×g, 5 min, 4°C). And

we used TURBO DNA-free TMKit (Invitrogen) to remove DNA contamination from the

extracted RNA. cDNA synthesis were used the qRT-PCR Kit (Toyobo, FSQ-101). We

conducted quantitative reverse-transcriptase PCR (qRT-PCR) in the LightCyclerqPCR

apparatus (Bio-Rad) with the FastStart SYBR Green Master (Roche). All primers used for qRT-

PCR assay are listed in Key Resources Table.

Immunofluorescence staining

Expanded islet clusters in martrigel were fixed in 4% paraformaldehyde for 30 min at room

temperature, followed with blocking and permeabilizing in PBS with 0.5% Triton X-100

(Solarbio, T8200) and 5% donkey serum (Solarbio, SL050) for 30min at room temperature.

Then, samples were incubated with primary antibody at 4℃ overnight, followed by incubation

with secondary antibody for 2h at room temperature. DAPI (Beyotime, C1002) was used to

stain the nucleus and find islets. The following antibodies were used for immunofluorescence:

anti-INSULIN (1:200, sc-9168; Santa), anti-SOMATOSTATIN (1:600, ab30788; Abcam),

anti-GLUCAGON (1:200, G2654; Sigma). Expanded islet clusters imaging was performed on

Zeiss LSM 780 and processed using ImageJ or Adobe illustator software.

Insulin secretion measurement

The insulin secretions were preformed similar to previously described in our group[7,9,24]. Fifty

day 7 expanded islet clusters (cell number is comparable to 1/10 of primarily isolated islets,

1000~2000 cells) or ten islets for each group were starved in 2.8 mM glucose MKRBB buffer

(5 mM KCl, 120 mM NaCl, 15 mM Hepes, 24 mM NaHCO$_3$, 1 mM MgCl$_2$, 2 mM CaCl$_2$,

pH=7.4) for 1 hour at 37℃, and then treated with 20 mM glucose and 100 nM GLP1, whereas

the control groups were treated with 2.8 mM glucose for 30 min. The supernatant fractions were

collected to measure the secreted insulin. The insulin levels were measured with the Millipore

358 Rat / Mouse Insulin (Cat. # EZRMI-13K), as what indicated by the manufacturer's instructions.

360 **Somatostatin measurement**

361  The somatostatin secretions were preformed similar to previously described in our group[7,9,24].

362 Fifty day 7 expanded islet clusters or ten primary islets for each group were starved in 1 mM

363 glucose MKRBB buffer for 1 hour at 37℃, then treated with 20 mM glucose and 100 nM UCN3,

364 whereas the control groups were treated with 1 mM glucose for 1 hour. The supernatant

365 fractions were collected for somatostatin measurement using the Phoenix Pharmaceuticals

366 ELISA kit (EK-060-03), as what indicated by the manufacturer's instructions.

368 **Glucagon measurement**

369  The glucagon secretions were preformed similar to previously described in our group[7,9,24].

370 Fifty day 7 expanded islet clusters or ten primary islets for each group were starved in 12 mM

371 glucose MKRBB buffer for 1 hour at 37℃, and then treated with 1 mM glucose and 25 mM

372 Arginine, whereas the control groups were treated with 5 mM Arginine for 1 hour. The

373 supernatant fractions were collected to measure the secreted glucagon. The glucagon levels

374 were measured with the Cloud-Clone Corp Mouse Glucagon (Cat. CEB266Mu), as what

375 indicated by the manufacturer's instructions.

377 **Statistical analysis**

378  All the qRT-PCR data were performed independently for at least three times. Statistical

379 analyses were performed using GraphPad Prism 7 software. Experimental data were performed

380 using unpaired two-tailed Student's t-test. Results are presented as mean ± SEM; $P < 0.05$ was

381 considered statistically significant.

383 **Acknowledgements**

384  This work was supported by funding from the National Science Fund for Excellent Young

385 Scholars Grant (81822008 to X.Y.,), the National Natural Science Foundation of China

386 (31671197 to X.Y., 81773704 to J.-P.S., 31970779 to H-L.H., and 81173086 to J.A.,), National

387 Key Research and Development Program of China (2019YFA0111400 to H-L.H.), and the

388 Shandong Key Research and Development Program (2017GSF18138 to J.A.,).

Competing interests

391   The authors declare no competing interests.

Author contributions

X.Y. started the idea for in vitro expansion of islet tissues since 2010. X.Y. J.-P.S supervised the overall project design and execution. H.-L.H provided key experience and protocols for in vitro 3D tissue culture and initial components advices. X.Y. J.-P.S. and H.-L.H. initiated the project. J.P.-S organized the screening idea. X.Y. J.-P.S and J.-Y.L designed the screening details for chemicals and other factors. X.Y., J.-P.S., J.-Y.L. and J.C. designed the cell culture and all other experiments. J.-Y.L., J.C., Y.-Q.D., W.P., X.Y. and J.-P.S. participated in data analysis and interpretation. Z.Z., Y.-Q.D. and R.-J.C. provided the new material preparation methods. J.W., Z.Y., F.Y. and J.A. provided insightful idea. Y.-F.X., H.L., W.-T.A. and J.W. participated in molecular biology and animal experiments. X.Y. and J.-P.S wrote the manuscript. H.-L.H, R.-J.C., J.-Y.L., J.C., Y.-Q.D. and W.P. revised manuscript and provided key insights for discussion.

Contact for reagent and resource sharing

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Professor Xiao Yu. (yuxiao@sdu.edu.cn), Jin-Peng Sun (sunjinpeng@bjmu.edu.cn) or Hui-Li Hu. (huhuili@sdu.edu.cn).

References

1.   Zhu, S., et al. Human pancreatic beta-like cells converted from fibroblasts. *Nature communications.* **7**, 10080 (2016).

2.   Li, K., et al. Small molecules facilitate the reprogramming of mouse fibroblasts into pancreatic lineages. *Cell Stem Cell.* **14**, 228-236 (2014).

3.   D'Amour, K.A., et al. Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells. *Nature biotechnology.* **24**, 1392-1401 (2006).

4.   Jiang, W., et al. In vitro derivation of functional insulin-producing cells from human embryonic stem cells. *Cell Res.* **17**, 333-344 (2007).

5.   Lancaster, M.A. & Knoblich, J.A. Organogenesis in a dish: modeling development and disease using organoid technologies. *Science.* **345**, 1247125 (2014).

6.   Clevers, H. Modeling Development and Disease with Organoids. *Cell.* **165**, 1586-1597 (2016).

7.   Li, Q., et al. A cullin 4B-RING E3 ligase complex fine-tunes pancreatic delta cell paracrine interactions. *J Clin Invest.* **127**, 2631-2646 (2017).

8.   Dou, H., et al. Calcium influx activates adenylyl cyclase 8 for sustained insulin secretion in rat pancreatic beta cells. *Diabetologia.* **58**, 324-333 (2015).

9.   Wang, H.M., et al. A stress response pathway in mice upregulates somatostatin level and transcription in pancreatic delta cells through Gs and beta-arrestin 1. *Diabetologia.* **57**, 1899-1910 (2014).

10.   van der Meulen, T., et al. Urocortin3 mediates somatostatin-dependent negative feedback

431   control of insulin secretion. *Nat Med.* **21**, 769-776 (2015).

432 11. Sato, T., et al. Long-term expansion of epithelial organoids from human colon, adenoma,
433   adenocarcinoma, and Barrett's epithelium. *Gastroenterology.* **141**, 1762-1772 (2011).

434 12. Boj, S.F., et al. Organoid models of human and mouse ductal pancreatic cancer. *Cell.* **160**, 324-
435   338 (2015).

436 13. Bartfeld, S., et al. In vitro expansion of human gastric epithelial stem cells and their responses
437   to bacterial infection. *Gastroenterology.* **148**, 126-136.e126 (2015).

438 14. Broutier, L., et al. Culture and establishment of self-renewing human and mouse adult liver and
439   pancreas 3D organoids and their genetic manipulation. *Nat Protoc.* **11**, 1724-1743 (2016).

440 15. Sampaziotis, F., et al. Reconstruction of the mouse extrahepatic biliary tree using primary
441   human extrahepatic cholangiocyte organoids. *Nature medicine.* **23**, 954-963 (2017).

442 16. Hu, H., et al. Long-Term Expansion of Functional Mouse and Human Hepatocytes as 3D
443   Organoids. *Cell.* **175**, 1591-1606.e1519 (2018).

444 17. Sachs, N., et al. Long-term expanding human airway organoids for disease modeling. *Embo j.*
445   **38**(2019).

446 18. Schutgens, F., et al. Tubuloids derived from human adult kidney and urine for personalized
447   disease modeling. *Nat Biotechnol.* **37**, 303-313 (2019).

448 19. Xiang, C., et al. Long-term functional maintenance of primary human hepatocytes in vitro.
449   *Science (New York, N.Y.).* **364**, 399-402 (2019).

450 20. Huch, M., et al. Unlimited in vitro expansion of adult bi-potent pancreas progenitors through
451   the Lgr5/R-spondin axis. *EMBO J.* **32**, 2708-2721 (2013).

452 21. Hou, P., et al. Pluripotent stem cells induced from mouse somatic cells by small-molecule
453   compounds. *Science (New York, N.Y.).* **341**, 651-654 (2013).

454 22. Sorenson, R.L. & Brelje, T.C. Adaptation of islets of Langerhans to pregnancy: beta-cell growth,
455   enhanced insulin secretion and the role of lactogenic hormones. *Hormone and metabolic*
456   *research = Hormon- und Stoffwechselforschung = Hormones et metabolisme.* **29**, 301-307
457   (1997).

458 23. Schraenen, A., et al. mRNA expression analysis of cell cycle genes in islets of pregnant mice.
459   *Diabetologia.* **53**, 2579-2588 (2010).

460 24. Li, N., et al. Ablation of somatostatin cells leads to impaired pancreatic islet function and
461   neonatal death in rodents. *Cell Death Dis.* **9**, 682 (2018).

462 25. Liang, Y.L., et al. Phase-plate cryo-EM structure of a biased agonist-bound human GLP-1
463   receptor-Gs complex. *Nature.* **555**, 121-125 (2018).

464 26. Hu, Q.X., et al. Constitutive Galphai coupling activity of very large G protein-coupled receptor
465   1 (VLGR1) and its regulation by PDZD7 protein. *The Journal of biological chemistry.* **289**, 24215-
466   24225 (2014).

467 27. Huch, M., et al. Long-term culture of genome-stable bipotent stem cells from adult human liver.
468   *Cell.* **160**, 299-312 (2015).

469 28. Staels, W., et al. Conditional islet hypovascularisation does not preclude beta cell expansion
470   during pregnancy in mice. *Diabetologia.* **60**, 1051-1056 (2017).

471 29. Kim, H., et al. Serotonin regulates pancreatic beta cell mass during pregnancy. *Nat Med.* **16**,
472   804-808 (2010).

473 30. Karnik, S.K., et al. Menin controls growth of pancreatic beta-cells in pregnant mice and
474   promotes gestational diabetes mellitus. *Science (New York, N.Y.).* **318**, 806-809 (2007).

31.   Zhao, X., et al. Involvement of the STAT5-cyclin D/CDK4-pRb pathway in beta-cell proliferation stimulated by prolactin during pregnancy. *American journal of physiology. Endocrinology and metabolism.* **316**, E135-e144 (2019).

Figure legends

Figure 1. Flowchart for the development of the medium recipe used for pancreatic islet expansion.

The chemicals and hormones used in previous studies for stem cell expansion, differentiation and pancreatic islet cell conversion (upper panel) are classified by their reported function related to cell division and proliferation, islet identity, and cell renewal and regeneration (middle panel). The common factors are summarized and combined for the formulation of the medium recipe used for in vitro pancreatic islet cluster expansion.

Figure 2. In vitro expansion of pregnant mouse islet clusters.

(A) Representative images of single islet cells or islet clusters (approximately 6000-9000 μm^2 original surface area) derived from pregnant mice after 3D culture in pancreatic islet cluster expansion medium (PIEM) at the indicated time points. Experiments were performed on pregnant mice. The scale bar in the single-cell image represents 50 μm, and one in the cluster image represents 100 μm.

(B) Representative images of dispersed wild-type mouse islet single cells or clusters at various times after 3D culture with PIEM. The scale bar in the single-cell image represents 50 μm, and one in the cluster image represents 100 μm.

(C) Numbers of expanded clusters per 100 islet cell clusters. The experiments were performed in triplicate and with pregnant mice. The data are represented as the mean ± SEM.

(D) Numbers of expanded clusters per 100 islet cell clusters. The experiments were performed in triplicate on wild-type mice. The data are represented as the mean ± SEM.

(E) Comparison of islet cell identity marker expression in pregnant mouse pancreatic islet clusters vs isolated primary mouse islets. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; clusters were compared with primary mouse islets. The data are shown as the mean ± SEM of at least three independent experiments. The data statistics were analysed using an unpaired two-tailed Student's t-test.

Figure 3. Immunostaining for pancreatic islet cell markers and functional examination of in vitro expanded pancreatic islet clusters.

(A) Immunofluorescence staining for insulin and glucagon in expanded pancreatic islet clusters derived from pregnant mice; the scale bar represents 100 μm.

(B) Immunofluorescence staining for insulin and somatostatin in expanded pancreatic islet clusters derived from pregnant mice; the scale bar represents 100 μm.

(C) Insulin secretion of primary islets or expanded islet clusters treated with 20 mM glucose and 100 nM glucagon peptide 1 (GLP-1) were measured for 30 minutes. The control groups were treated with 2.8 mM glucose. ***, $p < 0.001$, stimulation groups were compared with control groups. ###, $p < 0.001$, expanded islet clusters were compared with primary islets.

(D) Somatostatin secretion of primary islets and expanded islet clusters treated with 20 mM glucose and 100 nM UCN3 were measured for 1 hour. The control groups were treated with 1 mM glucose. **, $p < 0.01$, stimulation groups were compared with control groups. NS, no significance, expanded islet clusters were compared with primary islets.

(E) Glucagon secretion of primary islets and expanded islet clusters treated with 25 mM Arginine were measured for 1 hour. The control groups were treated with 5mM Arginine. ***, $p < 0.001$, stimulation groups were compared with control groups. NS, no significance, expanded islet clusters were compared with primary islets.

(C-E) The data are shown as the mean ± SEM of at least three independent experiments. The data statistics were analysed using an unpaired two-tailed Student's t-test.

Figure 4. In vitro expansion of rat islet clusters.

(A) Representative images of rat islet single cells or clusters (approximately 5000-8000 μm² original surface area) at various times after 3D culture with PIEM. The scale bar in the single-cell image represents 50 μm, and one in the cluster image represents 82.5 μm.

(B) Numbers of expanded clusters per 100 islet cell clusters. The experiments were performed on rats in triplicate. The data are represented as the mean ± SEM.

(C) Comparison of the expression of islet cell identity markers in rat expanded clusters compared with primary islets. *, $p < 0.05$; **, $p < 0.01$; clusters were compared with primary rat islets. The data are shown as the mean ± SEM of at least three independent experiments. The data statistics were analysed using an unpaired two-tailed Student's t-test.

Figure 5. qRT-PCR analysis of gene expression in expanded clusters **compared to primary islets.**

(A) Comparison of the expression of proliferation and cell cycle markers in pregnant mouse expanded clusters vs. isolated primary islets.

(B) Comparison of the expression of pluripotency and regeneration markers in pregnant mouse expanded clusters vs. isolated primary islets.

(C) Comparison of the expression of islet and pancreatic progenitor markers in pregnant mouse expanded clusters vs. isolated primary islets.

(D) Comparison of the expression of proliferation and cell cycle markers in rat expanded

clusters vs. isolated primary islets.

(E) Comparison of the expression of pluripotency and regeneration markers in rat expanded

clusters vs. isolated primary islets.

(F) Comparison of the expression of islet and pancreatic progenitor markers in rat expanded

clusters vs. isolated primary islets.

(A-F) *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; clusters were compared with primary islets. The

data are shown as the mean ± SEM of at least three independent experiments. The data statistics

were analysed using an unpaired two-tailed Student's t-test.

Figure 6. Essential roles of FSK and exendin4 in in vitro pancreatic islet expansion.

(A-D) qRT-PCR analysis of the expression of islet cell identity markers (A), proliferation and

cell cycle markers (B), pluripotency and regeneration markers (C) and islet and pancreatic

progenitor markers (D) in pregnant mouse clusters after 3 days of culture with PIEM or PIEM

without forskolin or exendin4.

(A-D) *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; cell clusters cultured in PIEM without FSK or

exendin4 were compared with those cultured in complete PIEM. The data are shown as the

mean ± SEM of at least three independent experiments. The data statistics were analysed using

an unpaired two-tailed Student's t-test.

Figure 7. Scheme of the current approach and future direction.

Schematic depicting the isolation, seeding and the expansion of primary islet cell clusters. After

the islets were isolated from pregnant mice or wild-type rats, the islets were digested into

appropriate cell clusters by incubation with the dispase II. The digestion time were optimized

and the mechanical blow force was used for clusters generation. These clusters were cultured

3D in PIEM to achieve the in vitro expansion of the islet cell clusters.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7