**Randomly primed, strand-switching MinION-based sequencing for the detection and**

**characterization of cultured RNA viruses**

Kelsey T. Young, Kevin K. Lahmers, Holly S. Sellers, David E. Stallknecht, Rebecca L.

Poulson, Jerry T. Saliki, S. Mark Tompkins, Ian Padykula, Chris Siepker, Elizabeth W. Howerth,

Michelle Todd, and James B. Stanton[1]

Department of Pathology (Young, Siepker, Howerth, Stanton), Poultry Diagnostic and Research

Center, Department of Population Health (Sellers), Southeastern Cooperative Wildlife Disease

Study Department of Population Health (Stallknecht, Poulson), Athens Veterinary Diagnostic

Laboratory (Saliki), Center for Vaccines and Immunology, Department of Infectious Diseases

(Tompkins, Padykula) College of Veterinary Medicine, University of Georgia, Athens, GA,

USA; Department of Biomedical Sciences & Pathobiology (Lahmers, Todd), VA-MD College of

Veterinary Medicine, Virginia Tech University, Blacksburg, VA, USA.

[1]Corresponding author: James B. Stanton, Department of Pathology, College of Veterinary

Medicine, University of Georgia, 501 DW Brooks Drive, Athens, GA 30602. jbs@uga.edu

Running title: Random strand-switching sequencing of cultured RNA viruses

**Abstract.** RNA viruses rapidly mutate, which can result in increased virulence, increased escape

from vaccine protection, and false negative detection results. Targeted detection methods have a

limited ability to detect unknown viruses and often provide insufficient data to detect

coinfections or identify antigenic variants. Random, deep sequencing is a method that can more

fully detect and characterize RNA viruses and is often coupled with molecular techniques or

culture methods for viral enrichment. Viral culture coupled with third-generation sequencing

were tested for the ability to detect and characterize RNA viruses. Cultures of bovine viral

diarrhea virus, canine distemper virus, epizootic hemorrhagic disease virus, infectious bronchitis

virus, two influenza A viruses, and porcine respiratory and reproductive syndrome virus were

sequenced on the MinION platform using a random, reverse primer in a strand-switching

reaction, coupled with PCR-based barcoding. Reads were taxonomically classified and used for

reference-based sequence building using a stock personal computer. This method accurately

detected and identified complete coding sequence genomes with a minimum of 20× coverage

depth for all seven viruses, including a sample containing two viruses. Each lineage-typing

region had at least 26× coverage depth for all viruses. Furthermore, analyzing the canine

distemper virus sample through a pipeline devoid of canine distemper virus reference sequences

modeled the ability of this protocol to detect unknown viruses. These results show the ability of

this technique to detect and characterize dsRNA, negative- and positive-sense ssRNA,

nonsegmented, and segmented RNA viruses.

**Key Words: metagenomic; MinION; RNA viruses; sequencing; strand-switching**

**Introduction**

RNA viruses are common etiological agents of animal diseases. Many of these diseases, such as

influenza,[1] Newcastle disease (ND),[2] epizootic hemorrhagic disease (EHD),[3] infectious

bronchitis (IB),[4] and porcine respiratory and reproductive syndrome (PRRS)[5] are global

economic and/or health burdens for domestic and wild animal populations. Moreover, RNA

viruses account for most emerging diseases due to the swift production of genetic variants,

enabling rapid evolution for adapting to environments and hosts.[6] The low fidelity of the viral-

encoded RNA-dependent RNA polymerases significantly contributes to the genetic diversity,

causing mutations up to a million times higher compared to cells.[7] Additionally, genetically

related viral species with segmented genomes (e.g., reoviruses and orthomyxoviruses) can

reassort genomic segments, resulting in increased genetic variation. Enhanced virulence,

resistance to vaccines, and ability for novel tissue tropism can also occur due to nucleotide

mutations, and reassortment events.[8-10] Therefore, there is a need for rapid detection of variants

through whole genome sequencing for proper diagnosis, treatment, control, and prevention.[11]

The molecular detection of RNA viruses is traditionally done using methods such as

polymerase chain reaction (PCR), real-time PCR (rtPCR), cloning, and *in situ* hybridization

(ISH).[12,13] However, these methods use targeted approaches that require prior knowledge of the

viral genome for detection and are inefficient for the discovery of novel viruses, mixed

infections, and identifying whole genomes. The advent of next generation sequencing (NGS) has

permitted new techniques that circumvent some issues with targeted RNA sequencing. Platforms

like Illumina allow untargeted deep sequencing for novel virus detection; but can be expensive

and labor intensive, particularly for whole genome sequencing.[14] Major drawbacks of these

platforms is the generation of large volumes of raw data and short reads which require high

performance computers and extensive computational analysis.[15] Another limitation of untargeted

RNA sequencing is the low relative abundance of viral RNA compared to host cellular RNA,

which may require depletion of ribosomal RNA (rRNA) and/or enrichment of viral RNA to

obtain an ample number of reads for viral strain detection.[16,17] For these reasons, using NGS for

accurate, whole viral genome sequencing remains challenging.

The long read sequencing technology provided by MinION sequencing (Oxford

Nanopore Technologies (ONT), Oxford, UK) has enabled rapid, inexpensive, high-throughput,

whole-genome sequencing of viruses.[18] MinION-based viral metagenomics studies have

accurately sequenced and identified whole genomes of chikungunya virus, Zaire ebolavirus,

hepatitis C virus, Venezuelan equine encephalitis virus, and Zika virus using untargeted

approaches by targeting poly-A sites and directly sequencing RNA (dRNA-seq) or using primer-

extension pre-amplification method (Round A/B); however, both of these methods resulted in

low quality reads and poor depth of coverage across the viral genome.[19,20] Other methods such as

sequence independent primer amplification (SISPA) have been used with MinION to obtain

whole genome sequences for bovine enterovirus from culture and canine distemper virus from

the brain of an affected dog.[21,22] While virus targeting can be accomplished with the previously

mentioned rRNA depletion or sequence targeting, it is also possible to couple this newest

sequencing technology with classical virus culture for viral enrichment. The aim of this study

was to develop a method to simultaneously detect and characterize various RNA viruses from

culture by using a randomly primed, strand-switching approach and sequencing on MinION.

Viruses were selected to include a double-stranded RNA virus (epizootic hemorrhagic disease

virus 2 [EHDV-2; family *Reoviridae*, genus *Orbivirus*]), positive-stranded viruses (infectious

bronchitis virus [IBV; family *Coronaviridae*, genus *Gammacoronavirus*, species *Avian*

*coronavirus*], porcine reproductive and respiratory syndrome virus [PRRSV; family

*Arteriviridae*, genus *Betaarterivirus*, species *Betaartervirus suid 2*], and bovine viral diarrhea

virus [BVDV; family *Flaviviridae*, genus *Pestivirus*, species *Pestivirus B*]), and negative-

stranded viruses (canine distemper virus [CDV; family *Paramyxoviridae*, genus *Morbillivirus*,

species *Canine morbillivirus*] and influenza A viruses [IAV; family *Orthomyxoviridae*, genus

*Alphainfluenzavirus*, species *Influenza A virus*] isolated from a dog and from a pig). Segmented

(EHDV-2 and IAV) and unsegmented genomes (BVDV, CDV, and PRRSV)  were also

represented in this study. This approach provides rapid, complete genome coding sequences

(CDS) for unknown RNA viruses in culture fluids and demonstrates the utility of the random

hexamer-based strand-switching primer for MinION library synthesis.

98                                    **Materials and methods**

**Samples**

The EHDV-2 isolate was propagated on cattle pulmonary artery endothelial cells (CPAE) from

spleen/lung tissue from a white-tailed deer (*Odocoileus virginianus*) collected in Georgia, USA

in 2016 at the Southeastern Cooperative Wildlife Disease Study at the University of Georgia

(UGA); CPAE cells are also persistently infected with BVDV. The CDV sample was isolated

from the brain of an infant, female raccoon (*Procyon lotor*) from Kentucky in 2018 using

African green monkey kidney cells expressing canine signaling lymphocytic activation molecule

(Vero-Dog SLAM cell line) in the Athens Veterinary Diagnostic Laboratory (AVDL). The

canine-origin IAV sample was collected from a nasal swab of a 7-year-old, male, boxer dog in

2015 from Georgia and was cultured in embryonated chicken eggs at the Poultry Diagnostic and

Research Center (PDRC), UGA. The Center for Vaccines and Immunology, UGA provided the

swine-origin influenza sample, which was isolated in 2019 from a 6-month-old, female,

Hampshire-cross pig from Georgia after testing positive for IAV by immunohistochemistry and

PCR. Swine IAV from lung homogenate was propagated on Madin-Darby canine kidney

(MDCK) cells. The IBV sample (Mass vaccine) was cultured in embryonated chicken eggs and

supplied by the PDRC. Isolate VR2385 of PRRSV was cultured on MARC-145 cells at the

Veterinary Diagnostic Laboratory, Iowa State University. Reverse-transcription real-time PCR

(RT-rtPCR) was conducted for BVDV, EHDV-2,[3] CDV, canine-origin influenza, and IBV

(Table 1). The CDV (AVDL), canine IAV (PDRC), and IBV (PDRC) RT-rtPCR assays were

done using in-house diagnostic methods. For BVDV, (RT-rtPCR) was performed using a

modified protocol[23] with SuperScript® III First-Strand Synthesis SuperMix for qRT-PCR

(Invitrogen, Carlsbad, CA) and amplified in a CFX96™ Touch Real-Time PCR Detection

System (Bio-Rad Laboratories, Inc, Hercules, CA) thermal cycler.

**Total RNA extraction**

All RNA extraction to sequencing steps for all viruses, except for PRRSV and a CDV replicate,

were conducted at UGA. Library preparation and sequencing of PRRSV and a CDV replicate

were conducted at Virginia Tech University. Total RNA for CDV, EHDV-2, BVDV, swine IAV,

and canine IAV was extracted using 1 ml of culture supernatant using Trizol® LS Reagent

(ThermoFisher Scientific, Waltham, MA) following manufacturer's protocol. Total RNA was

eluted in 88.5 µl of nuclease-free water (Qiagen, Hilden, Germany). DNase treatment was

performed using RNase-Free DNase Set (Qiagen) and then purified using RNeasy® MinElute®

Cleanup Kit (Qiagen) per manufacturer's instructions. Total RNA for IBV was extracted using

the QIAamp® Viral RNA Mini Kit (Qiagen) following manufacturer's protocol. At Virginia

Tech, RNA was extracted from PRRSV and one replicate of CDV using the QIAamp Viral RNA

Mini Kit (Qiagen) according to the manufacturer's instructions. Concentrations were measured

using the Qubit® RNA HS Assay Kit (ThermoFisher Scientific) on a Qubit® 3.0 Fluorometer.

**Strand-switching cDNA synthesis**

Strand-Switching cDNA synthesis for MinION sequencing was completed by modifying the 1D

PCR barcoding cDNA (SQK-LSK108) protocol from ONT. Reverse transcription was performed

by combining 8 µl of total RNA, 2 µl of 1 µM PCR-RH-RT primer (5' -

/5Phos/ACTTGCCTGTCGCTCTATCTTCNNNNNN- 3'; synthesized by Integrated DNA

Technologies [IDT], Coralville, IA, with standard desalting, ONT adapter sequence is

underlined), and 1 µl of 10 mM dNTPs. The reaction mixture was incubated at 65°C for 5

minutes, then snap cooled in an ice-water slurry for 1 minute. Then 4 µl of 5× RT buffer, 1 µl of

100mM DTT, 1 µl of 40 U/µl RNaseOUT™ (Invitrogen), and 2 µl of 10 µM strand-switching

oligo (PCR_Sw_mod_3G) (5' -

TTTCTGTTGGTGCTGATATTGCTGCCATTACGGCCmGmGmG- 3'; ONT provided

sequence, synthesized by IDT with HPLC purification) were added and incubated at 42°C for 2

minutes. SuperScript™ IV Reverse Transcriptase (Invitrogen) was added at a volume of 1 µl and

the reaction was incubated at the following conditions: 30 minutes at 50°C, 10 minutes at 42°C,

and 10 minutes at 80°C. The cDNA was purified by KAPA Pure Beads (Kapa Biosystems,

Wilmington, MA) or with AMPure XP beads (Beckman Coulter, Indianapolis, IN) at 0.7× beads:

solution ratio.

**Barcoding PCR**

The reverse-transcribed cDNA was amplified following ONT's 1D PCR barcoding cDNA

(SQK-LSK108) protocol using the PCR Barcoding Expansion 1-12 (EXP-PBC001) kit (ONT)

and LongAmp Taq 2× Master Mix (New England Biolabs, MA) with the following

thermocycling conditions: 95°C for 3 minutes; 18 cycles of 95°C for 15 seconds, 62°C for 15

seconds, 65°C for 22 minutes; 65°C for 23 minutes. The barcoded, amplified DNA was bead

purified at a 0.8× ratio.

**MinION library preparation and sequencing**

After the barcoding PCR, two to three samples were pooled by equal volume to make a total

volume of 47 µl. MinION libraries were prepared from the pooled barcoded amplicons using the

Ligation Sequencing Kit (SQK-LSK109) (ONT) following the 1D amplicon/cDNA by Ligation

per ONT's instructions. Briefly, the pooled samples were end prepped with NEBNext FFPE

Repair Mix (New England Biolabs) and NEBNext End repair/dA-tailing Module (New England

Biolabs) and bead purified at 1.0× bead volume. Then, sequencing adapters (ONT) were ligated

onto the end-prepped library using NEBNext Quick T4 DNA Ligase (New England Biolabs) and

bead purified (0.4× beads ratio) with the Long Fragment Buffer (LFB) (ONT). The final libraries

were combined with Sequencing Buffer (SQB) and Loading Beads (LB) per ONT's instructions

and were sequenced on used or new FLO-MIN106 R9.4 flow cells (ONT), with the exception of

the PRRSV and replicated CDV library, which were sequenced on a FLO-MIN107 R9.5, with

the MinION Mk1b sequencer. Per the pre-run quality control check for previously used flow

cells, a minimum of 1,000 active pores were required for sequencing. To maximize data

generation from single flow cells, used flow cells were nuclease flushed, before or after

sequencing, by adding a mixture of 10 µl DNase I and 190 µl DNase I Reaction Buffer (New

England Biolabs) to the priming port and incubated for 30 minutes at room temperature.

Sequencing was initiated by MinKNOW v.18.12.4–19.05.0 (ONT) using the 48-hour sequencing

protocol with the live basecalling option turned off. During the run, available FAST5 files were

processed as below to estimate number and length of viral reads to estimate sequencing time

required to obtain sufficient genome coverage.

**Pre-processing of raw MinION sequence data**

The FAST5 files produced from sequencing for IBV, CDV, swine-origin and canine-origin

influenza, and EHDV-2 and BVDV libraries were processed using an in-house script that

sequentially basecalled, demultiplexed, and trimmed adapters on a Macbook Pro (3.1 GHz Intel

Core i7, 8GB) using OS X El Capitan v.10.11.6. In summary, reads were basecalled using the

CPU version of Guppy v.2.1.3 (ONT) by defining the appropriate configurations for the flowcell

(FLO-MIN106) and kit (SQK-LSK109) used for sequencing and library preparation. Calibration

strand detection and filtering was enabled with the --calib_detect parameter. The basecalled fastq

files with a qscore ≥7 were sorted into a "pass" folder using the --qscore-filtering parameter and

used for further analysis. Then, reads were demultiplexed using Porechop v.0.2.4

(https://github.com/rrwick/Porechop) based on barcodes (-b output_dir) with --

require_two_barcodes setting enabled, 1,000,000 reads were aligned to all known adapter sets (--

check_reads 1000000), adapters with 99% identity were trimmed from read ends (--

adapter_threshold 99), and chimeric reads with middle adapters were removed.

The raw FAST5 files produced for PRRSV and the CDV replicate from Virginia Tech

were basecalled using the GPU version of Guppy v.3.1.5 on a Dell Precision 3060 Tower (Intel

Core i7, 31.3GB, NVIDIA GeForce GTX 1080) using Ubuntu 16.04 LTS. Guppy was initiated

using the same parameters as above with the flowcell configuration defined as "FLO-MIN107".

The basecalled reads were then trimmed and demultiplexed using Porechop with the same

parameters described previously.

**Virus classification and lineage typing**

After basecalling, demultiplexing, and adapter trimming, reads were classified with Centrifuge

v.1.0.4[24] by the lowest taxonomical rank with default parameters with the addition of allowing

50 assignments (or hits) for each read (-k 50) using a custom index built for each library based

on the propagation system. Custom indices were constructed using an exhaustive search for

complete genome sequences of all possible viruses infecting vertebrates downloaded from

NCBI's nucleotide database, as of 15 March 2019, that included 80,550 viral reference genomes

(Supplementary Table 1). Low-complexity regions in sequences were masked using dustmasker

(NCBI C++ Toolkit, https://ncbi.github.io/cxx-toolkit). Dustmasked whole genomes of the

species of cell line (see below), and/or the bovine genome (GCF_002263795.1_ARS-UCD1.2),

to account for fetal bovine serum used in cell culture, were obtained using `centrifuge-download`

and concatenated with the dustmasked vertebrate virus sequences to classify reads from host(s)

cellular RNA. Using the concatenated files, indices were built by `centrifuge-build` with

default parameters. The indices were then used to cluster reads based on short alignments to

respective host and viral sequences to determine the presence and abundance of viral species in

each sample using the standard out file. Reads to unexpected viral species were clustered and

compared to the Genbank database using web-based nucleotide Basic Local Alignment Tool

(BLASTn) (https://blast.ncbi.nlm.nih.gov) with default settings. Percentage of viral reads and

host reads were determined by removing duplicate reads and then dividing the number of reads

clustered by the number of total reads after demultiplexing.

Once the viral species was identified, custom lineage-typing Centrifuge indices were also

built to further classify beyond viral species and assess for the possibility of mixed infections of

the same virus species, as previously described for IBV.[25] Briefly, for each main lineage per

virus, one complete sequence of the lineage-typing region was selected and then combined with

the respective genome of the cell line. Indices were constructed using the following: 20 N-

terminal protease fragment ($N^{pro}$) sequences for BVDV[26] with the bovine genome

(GCF_002263795.1_ARS-UCD1.2), 13 hemagglutinin (*H*) gene sequences for CDV[27] with the

African green monkey genome (GCF_000409795_Chlorocebus_sabeus_1.1), nine *VP2*

sequences for EHDV[28] with the bovine genome (GCF_002263795.1_ARS-UCD1.2), 32 spike 1

(*S1*) sequences for IBV[29] with the chicken genome (GCF_000002315.4_Gallus_gallus-5.0), 18

hemagglutinin (*HA*) sequences and 11 neuraminidase (*NA*) sequences for IAV[30] with the canine

genome (GCF_000002285.3_CanFam3.1) or chicken genome (GCF_000002315.6_GRCg6a),

and 18 *ORF5* sequences for PRRSV[31] with the swine genome (GCF_000003025.6_Sscrofa11.1)

(Supplementary Table 2). All sequences were dustmasked, assigned a unique taxonomy

identification number, with the exception of the IAV sequences, and indices were built with

default settings using `centrifuge-build`. Viral lineage typing was evaluated by aligning the

individual trimmed and barcoded FASTQ files produced from Porechop to the respective custom

lineage-typing indexes with Centrifuge, and reads were clustered by taxonomy identifications

that represent potential lineage types in the sample. Each cluster of reads, representing

potentially different lineages, were then used for reference based consensus building using

Geneious v.11.1.3 (Biomatters, Aukland, New Zealand) to build a consensus for each lineage

using the "Map to Reference" tool with medium sensitivity, re-iterating up to five times, and

using the same reference sequences chosen for generating the custom Centrifuge lineage-typing

databases. Consensus sequences derived for each potential lineage were analyzed using BLASTn

with default settings. Results from BLASTn were sorted by query coverage and subject

coverage. Subjects with the highest query, subject coverages, and identical bit-scores were

considered the "top hit(s)" to identify the virus and lineage in each cluster. A coinfection was

defined as different clusters of reads creating consensus sequences that matched to different

lineages and reads could be parsed accordingly for final consensus building. If no coinfection

was detected, all viral reads were used for consensus building.

**Viral genome consensus sequence generation**

After viruses were lineage typed for each sample, all reads that aligned to respective viral species

from Centrifuge output were imported into Geneious for whole-genome, reference-based

consensus building with the "Map to Reference" tool, after removing duplicate reads, using the

same methods from above. Consensus sequence building was done with the requirement of at

least 20× coverage at each base. References used for mapping were selected from using the "top

hit" result in BLASTn from the lineage-typing analysis. Due to the known errors within long

homopolymer regions during MinION sequencing,[32] consensus sequences were manually

inspected and homopolymer areas that resulted in frameshifts were manually edited. The CDS

regions were extracted from each edited consensus sequence and analyzed using BLASTn. For

viruses with gapped genomes (e.g., CDV, IBV, and PRRSV), complete CDS and non-coding

intergenic regions were extracted and analyzed with BLASTn.  The "top hit" for each consensus

sequence was determined by using the same criteria from above. The CDS regions for lineage-

typing regions for each virus were also extracted and analyzed with BLASTn. The best "top hit",

percent pairwise identity, mean, minimum, maximum, and standard deviation of base coverage,

and percentage of genome coverage were evaluated for each viral genome and lineage-typing

region consensus sequences with the statistics tool in Geneious.

**Simulating novel virus identification**

In addition to the above method, sequencing data from the CDV sample were analyzed in a

manner that simulated novel virus identification by constructing an all vertebrate virus, custom

Centrifuge index with the omission of all canine morbillivirus sequences downloaded from

NCBI as of 12 April 2019 (Supplementary Table 1). The index was assembled as described

previously with African green monkey genome. Trimmed and demultiplexed reads from the

CDV sample were classified with Centrifuge using parameters described above. Reads were

grouped by taxonomy ID and exported into Geneious to build consensus sequences as described

previously. The consensus was analyzed with BLASTn by excluding canine morbillivirus (taxid:

11232) in the search set. For comparison, Kraken-style reports were created using `centrifuge-`

`kreport` from the Centrifuge output files from the CDV-absent pipeline and the CDV-present

pipeline from above. Output files were visualized using Pavian v.1.0.[33] To discern the phylogeny

of the "novel" virus among other pathogens in the *Morbillivirus* genus, the final genome CDS

and phosphoprotein (*P*) gene CDS consensus sequences were aligned with 18 complete genome

CDS and *P* gene CDS acquired from GenBank using the neighbor-joining algorithm in

*ClustalW*[34] with default settings. Phylogenetic trees for complete genome CDS and *P* gene CDS

were inferred using the Maximum Likelihood statistical method based on the Tamura 3-

parameter substitution model[35] with bootstrap values calculated at 1000 replicates in MEGA X.[36]

**Sanger sequencing for validation**

Sanger sequencing was performed for the 5' UTR region of BVDV, *H* gene of CDV, and *VP2*

segment of EHDV-2 to confirm the identity of isolates and to compare MinION consensus

sequences for the lineage-typing regions. Briefly, using 8 µl of total RNA from each sample,

cDNA was synthesized using SuperScript® III First-Strand Synthesis System for RT-PCR

(Invitrogen) with random hexamers (50 ng/µl) following manufacturer's instructions. The cDNA

from each sample was amplified using DreamTaq Green PCR Master Mix (2×) (ThermoFisher

Scientific) following manufacturer's protocol with 10 µM of each respective primer (Table 2)

targeting partial sequences of the lineage-typing regions and 1 µl of cDNA. Thermocycling

conditions for PCR amplification of the 5' UTR region for BVDV are as follows: 95°C for 5

minutes; 34 cycles of 95°C for 30 seconds, 58°C for 30 seconds, 72°C for 30 seconds; 72°C for 5

minutes. Thermocycling conditions for PCR amplification of the *H* gene for CDV are as follows:

95°C for 5 minutes; 40 cycles of 95°C for 30 seconds, 50°C for 30 seconds, 72°C for 1.5

minutes; 72°C for 10 minutes. Thermocycling conditions for PCR amplification of the *VP2*

segment for EHDV are as follows: 95°C for 3 minutes; 40 cycles of 95°C for 30 seconds, 57°C

for 30 seconds, 72°C for 45 seconds; 72°C for 5 minutes. Electrophoresis with a 1.0% agarose

gel was performed to confirm PCR products. Amplicons were then purified using the QIAquick

PCR Purification Kit (Qiagen) following manufacturer's protocol and eluted in 30 µl nuclease-

free water (Qiagen). Final concentration and purity were measured using NanoDrop™ 2000

Spectrophotometer (ThermoFisher Scientific). The purified PCR products and 5 µM of each

primer (Table 2) were submitted to GENEWIZ (South Plainfield, NJ) for bidirectional Sanger

sequencing.

**Sanger sequencing analysis and pairwise identity with MinION consensus sequences**

Results from Sanger sequencing were analyzed with Geneious. Using the chromatograms, low

quality regions (≥Q40) from the 5□ and 3□ ends were trimmed and ambiguous bases were

manually edited. Primers were trimmed before aligning forward and reverse sequences from

each sample using Geneious Alignment with default settings. The consensus sequence was

compared using BLASTn and selecting the best "top hit" with the same criteria described above

for MinION data.

Partial lineage-typing sequences from Sanger sequencing for BVDV, CDV, and EHDV

were compared with the full-length lineage-typing consensus sequences from MinION using

Geneious Alignment with default settings. Pairwise identity of the alignment was calculated with

Geneious.

318                                     **Results**

**MinION sequencing, viral classification, and lineage typing**

Libraries were prepared for seven cultured samples for seven different viruses by using random

hexamer primed, strand-switching for reverse transcription, PCR-based barcoding, and pooled

sequencing using MinION. Sequencing occurred for a minimum of about 2 hrs and 45 min and a

maximum of about 47 hrs and 37 min to obtain 567,780–6,984,000 raw reads. Total sequencing

time varied between runs based on the time of day the run was started, the number of viral reads

detected early in the sequencing run, and the ability to re-use a flow cell (i.e., if the flow cell was

not to be reused, then it was often sequenced to near exhaustion). After the raw reads were

basecalled, demultiplexed, and trimmed, 7,523–1,173,058 reads were assigned to barcodes of

interest and used for viral classification (Table 3).

Barcoded reads for each of the seven libraries were aligned to custom-built indices with

Centrifuge that detected seven different viruses belonging to three main types of RNA viral

genomes: BVDV (positive-sense ssRNA), CDV (negative-sense ssRNA), EHDV-2 (dsRNA),

IBV (positive-sense ssRNA), canine-origin IAV (negative-sense ssRNA), swine-origin IAV

(negative-sense ssRNA), and PRRSV (positive-sense ssRNA). Additionally, three of the viruses,

EHDV-2, canine-origin influenza, and swine-origin influenza, are comprised of segmented

genomes. The randomly primed strand-switching method was also able to classify a sample

containing EHDV-2 and BVDV. All viruses detected were categorized with viral reads

representing 0.97–63.3% of all reads sequenced for each sample with percentage of host reads

ranging from 2.5 to 80.4% (Table 3). Reads to unexpected viral species were analyzed with

BLASTn and were determined to be short alignments to various host, often ribosomal,

sequences.

Viral sequences were parsed and further classified into lineage types using custom built

lineage-typing indices for Centrifuge. Reads were clustered based on the lineage-typed alignment

and consensus sequences were built using Geneious. While the Centrifuge alignments suggested

the possibility of 2–19 lineages per virus isolate, only one lineage per species was detected in

each isolate after consensus building of each potential lineage (Table 4).

The complete CDS for each virus's lineage-typing region had at least 33× depth of

coverage with a mean range of 105.1–22,968.0 (Table 4). The BLASTn comparison for the

genotyping regions resulted in 96.49–100.0% pairwise identity with each "top hit" (Table 4).

Using the $N^{pro}$ region, the lineage of BVDV was determined as genotype BVDV-2a with 96.23%

identity with McCart_C strain. Sequences from both libraries of CDV were genotyped as

America II with 96.49% identity of the *H* gene to A75/17. The genotype for EHDV-2 was

determined as EHDV-2w with 99.90% identity to the OV215 strain using the *VP2* segment. IBV

was determined to be genotype 1-lineage 4, with 100.0% identity of the *S1* gene to the Ma5

strain. The *HA* and *NA* segments of the canine-origin IAV showed 100.0% and 99.79% identity,

respectively, to an H3N2 subtype circulating in dogs in North America. The porcine-origin

influenza was determined to be an H1N2 subtype similar to other North American porcine-origin

IAVs, with 99.65% identity to the *HA* segment and 99.79% identity to the *NA* segment. The

*ORF5* for PRRSV had 99.67% identity to VR2385 and was determined as a Type 2, lineage 5.

**Viral genome consensus sequence evaluation**

For all seven viruses, complete genome CDS were acquired with a minimum of 26× depth of

each base, with the exception of the replicated CDV with 90.2% genome coverage at 20× depth

(Table 5). Additionally, complete CDS for all segments of three segmented viruses, EHDV-2

(dsRNA), swine influenza, and canine influenza (negative-sense ssRNA), were obtained with a

minimum of 43× depth of coverage and an average of 99.89% identity across all segments. The

mean depth for each virus was 72.1–28,014.8. The complete genome CDS consensus sequences

had a high pairwise identity of 97.37–100.0% to their respective "top hit" when using BLASTn

to compare sequences with GenBank databases (Table 5). The complete CDS consensus

sequences for BVDV, EHDV, and canine IAV were deposited in GenBank under the following

accession numbers: BVDV = MN824468; EHDV= MN824457–MN824466; canine IAV =

MN812282–MN812289. The complete CDS with intergenic regions for CDV was also deposited

in Genbank as accession number MN824467.

**Simulating novel virus identification**

Under simulated conditions in which CDV would be an unknown virus, the Centrifuge standard

out file for CDV analyzed showed that approximately 500 reads aligned to a morbillivirus, with

239 reads aligned to phocine distemper virus (PDV; family *Paramyxoviridae*, genus

*Morbillivirus*, species *Phocine morbillivirus*) and fewer reads (<70) aligning to other more

distantly related morbilliviruses: feline morbilliviruses (species *Feline morbillivirus*), rinderpest

virus (species *Rinderpest morbillivirus*), peste-des-petits-ruminants virus (species *Small

ruminant morbillivirus*), cetacean morbillivirus (species *Cetacean morbillivirus*), and measles

virus (species *Measles morbillivirus*) (Figure 1a). In contrast, when CDV was included in the

Centrifuge database, approximately 4,000 reads aligned to CDV, which was the only

morbillivirus detected (Figure 1b). In the CDV-absent analysis, reads scattered across different

morbilliviruses suggesting that the actual species was absent from the database but is most

similar to PDV. A total of 251 (12 reads aligned to 3 or more sequences, resulting in these reads

not being counted in the Centrifuge standard out file) reads were classified as PDV and were

exported to Geneious for reference-based consensus building by aligning to

PDV/Wadden_Sea.NLD/1988 (accession: NC_028249). The consensus sequence was analyzed

with BLASTn by excluding canine morbillivirus in the search set which resulted in a 78.71%

identity with PDV/Wadden_Sea.NLD/1988 (accession: KC802221.1) with 78.0% query

coverage. Furthermore, phylogenetic analysis of the whole genome CDS and *P* gene CDS

consensus of the dubbed "novel" sequence clustered with PDV but shows sequence divergence

suggestive of a species similar to, but different from, PDV (Figure 2).

**Sanger Sequencing and pairwise identity with MinION**

Sanger sequencing targeting partial sequences of the lineage-typing regions for BVDV, CDV,

and EHDV was used to confirm lineage types in the samples (Table 6). For BVDV, primers

targeting a partial sequence of the 5☐ UTR were used and resulted in a "top hit" to BVDV-2

isolate 95-1501. Sequencing of the *H* gene for CDV resulted in a "top hit" to CDV isolate

THA/VG. The *VP2* segment was targeted for EHDV and resulted in a "top hit" to EHDV-2

isolate OV617. The consensus sequences from Sanger and MinION sequencing had 100.0%

pairwise identities for all three viruses (Table 6). MinION consensus sequences compared to

Sanger for BVDV and EHDV had identical "top hits". For CDV, MinION sequencing had a "top

hit" to CDV A75/15 with 96.49% identity but the shorter 923 bp fragment from Sanger had a top

alignment with CDV THA/VG with 96.53% identity (Table 6). The Sanger sequence is based on

smaller fragments of the lineage-typing region and, therefore, the BLASTn-based pairwise

identities cannot be directly compared between MinION and Sanger sequences.

406                                   **Discussion**

A large proportion of emerging diseases are caused by RNA viruses[40] and, due to their

mutability, rapid detection is needed; however, this can be hindered due to the series of PCR

panels used for identification of unknown viruses and inefficient for the discovery of

coinfections.[41] Deep sequencing-based approaches using viral nucleic acid enrichment methods

have been described to address this issue, including targeted and untargeted library preparations,

like SISPA.[21,22] The methodology in this study demonstrates the application of culture-based

viral enrichment followed by random, strand-switching MinION sequencing for accurately

detecting and characterizing RNA viruses. RNA viruses with varying genome compositions

(single stranded [positive- and negative- sense], double stranded, and segmented) were used to

demonstrate the ability of untargeted strand-switching to obtain complete CDS of genotyping

regions and whole genomes with viral culture enrichment methods. Moreover, two viruses

(EHDV-2 and BVDV) were detected from a single sample, illustrating the utility of the random

sequencing approach. Lastly, data analysis for one sample (CDV) was treated as a novel virus,

highlighting the feasibility of this method to identify a new or poorly characterized virus.

This random sequencing approach proved to be robust across various RNA viruses in

obtaining full length CDS of the complete genome after using an unbiased, fast aligner to

identify the likely virus, followed by reference-based consensus building to identify the lineage

type. With at least $26\times$ depth of coverage across the genome for all viruses, whole genome CDS

had 96.99–100.0% identity to the top BLASTn hits. Complete CDS for all segments of three

segmented viruses, EHDV-2 (dsRNA), swine influenza, and canine influenza (negative-sense

ssRNA), were obtained with a minimum of $43\times$ depth of coverage and an average of 99.89%

identity across all segments. The best "top hit" for each segment does not match across all

segments to the same isolate, which may be due to reassortment events for segmented genomes[8,9]

and the unavailable sequence data for all segments for some isolates in NCBI.

The alignments of the whole genome CDS and the lineage-typing regions were similar

and consistent with the origin of the samples in this study. The identification of the EHDV as an

EHDV-2w with highest similarity to EHDVs circulating in white-tailed deer in the southeastern

United States in 2017 is consistent with the collection of this sample from a white-tailed deer in

Georgia in 2016. The influenza sample of canine origin was determined as a canine H3N2 strain,

matching the typing completed as part of the original diagnostic case workup. The full

sequencing provided by this method was able to confirm that the isolate in this study was most

similar to canine H3N2 viruses circulating in the southeastern United States in 2015, consistent

with the time and geographic location in which this sample was collected. Lineage typing of the

swine-origin influenza virus categorized it as an H1N2 most similar to an H1N2 IAV isolated

from a swine in 2018 from Ohio. It is possible that the slightly lower percent identity for the

swine-origin influenza is due to a relative paucity of sequence data available for 2019 swine-

origin IAV at this time. The CDV America II lineage, the classification of CDV in this study, is a

common lineage found in North American wildlife[42] and is consistent with the collection of this

sample from a raccoon in Kentucky. The sequences for IBV and PRRSV were highly similar to

the known sequences of those isolates.

This protocol was also tested on two scenarios. The first was the detection of two viruses

in a single culture system, demonstrating its ability to simultaneously obtain accurate, complete

CDS for rapid detection and characterization of viruses in coinfected samples, comparable to

other studies using advanced sequencing for analysis of cultured viruses.[43] The second was the

data analysis under the simulating conditions of CDV being a novel virus. This resulted in read

alignments that spanned across the *Morbillivirus* genus. Excluding the background hits, the

largest proportion of reads hit to PDV and consensus building with these reads had a low

pairwise identity (78.71%), suggesting it was not a PDV. The phylogenetic divergence gives

evidence that the virus in the sample belongs to the *Morbillivirus* genus and is phylogenetically

related to PDV. If CDV was truly an unknown virus, the Centrifuge output would have

suggested the identification of a novel or divergent virus that is most similar to PDV sequences,

consistent with the known close genetic relationship between CDV and PDV.[44] Furthermore, in

the event of an unknown viral etiology and due to the multiplexing capabilities of MinION

sequencing, DNA library approaches can be applied and sequenced concurrently with the

random strand-switching library to identify RNA or DNA virus in the sample.[45,46]

Sanger sequencing was performed on partial lineage-typing sequences for BVDV, CDV,

and EHDV and compared with the full-length lineage-typing regions obtained from MinION

sequencing, resulting in 100.0% pairwise identity. The 100% pairwise identity between the

Sanger and MinION sequences is consistent with previous results using MinION sequencing,[2,25]

that demonstrated the ability to obtain accurate MinION sequences by increasing depth of

coverage.[47,48] Sanger sequencing and BLASTn results for the 5$\square$ UTR for BVDV and *VP2* for

EHDV matched the MinION lineage-typing region BLASTn results. For the *H* gene for CDV,

Sanger sequencing had a top BLASTn alignment with isolate THA/VG with 96.53%; however,

the results did show 96.0% identity with isolate A75/15, comparable to MinION sequencing

result of 96.49% identity to isolate A75/15. It is noteworthy that aligning the Sanger and

MinION results for the *H* gene in CDV showed 100.0% pairwise identity. The differences in

BLASTn top alignments are attributed to the shorter sequence from Sanger sequencing, causing

a slightly lower sequence specificity. Longer fragments, particularly whole genome sequencing,

have shown improved resolution for genotyping analyses.[49,50] Additionally, the limitations of

partial sequences used in Sanger sequencing are illustrated by the inability of Sanger sequencing

to identify a single best hit for EHDV-2 and CDV, which resulted in multiple hits with identical

BLASTn scores; whereas, the MinION-based method obtained complete CDS of the *VP2*

segment and *H* gene allowing for a single best hit for these viruses. In addition to more accurate

classification, whole genome sequencing of viruses is advantageous in providing more data for

detection, epidemiology, genotyping and phylogenetic analysis compared to partial and/or

targeted sequences obtained from other classical sequencing methods. The relatively limited data

provided by targeted sequencing may require further sequencing and expenses to obtain

additional biological information. In particular, different genetic sequences are used for

phylogenetic analysis and genotyping. Some viruses also have multiple genotyping regions, such

as BVDV, EHDV, and influenza.[26,28,30] Furthermore, recombination events are often difficult to

identify with partial sequences and are important in investigating host range, virulence, and

vaccine evasion.[51-53] Thus, routine, whole genome sequencing of viruses gives quick turnaround

data for various analyses and could provide more comprehensive databases needed to understand

viral evolution.

Random, deep sequencing for many high-throughput sequencing platforms has some

caveats such as false hits. As one example, illustrated in Figure 1, some reads were classified as

lassa mammarenavirus, various herpesviruses, and others. Endogenous retroviruses and some

DNA viruses were also frequently found to be present after annotating sequences. After

clustering and aligning these reads to GenBank using BLASTn, it was determined that the reads

were short sequences that matched to various host sequences. Reference sequences of the host

and propagation systems of the cultured viruses were included in the Centrifuge indices to reduce

these misalignments; however, these genomes are not as well described as other commonly

studied organisms and may not represent the full sequence diversity of the host genome. Users

can adjust alignment settings to help reduce the number of false hits, but as with any test,

increasing the specificity of the test will negatively impact sensitivity. Similar to other fields

(e.g., background lesions in pathology, growth of nonpathogenic bacteria in bacterial cultures),

confirmatory tests may be required and analysis of deep sequencing results requires evaluation

by a trained user, one experienced with bioinformatic methodology and with knowledge of

veterinary infectious diseases.

While rapid, cost-effective whole-genome sequencing may be useful in a diagnostic

setting, ease of use and robustness across laboratories is key for deployment. For this reason,

libraries for CDV and PRRSV were prepared at different laboratories. Both libraries for CDV

had similar results; however, the differences can possibly be attributed to the use of a sequencing

kit intended for 1D sequencing with a FLO-MIN107 (R9.5) flowcell that is typically used for

$1D^2$ sequencing. The repercussions of the 1D kit with a $1D^2$ flowcell combination are not well

known but it is of interest to note that trimming the basecalled files resulted in 1,691,792 reads

removed due to middle adapters, significantly decreasing the total number of reads assigned to

each barcode.

The present study provides promising results for quick identification of unknown

cultured RNA viruses by using MinION sequencing with a previously untested random

approach. Future studies to compare the varying random methods of deep sequencing are

required to determine the most efficient methods. As with other deep sequencing methods,[50]

MinION-based sequencing will also likely be used to metagenomically detect viruses directly

from clinical samples (e.g., serum, swabs, tissues), and work is ongoing to investigate this usage.

MinION sequencing is a cost-effective way to multiplex samples and achieve long reads for

more accurate genome consensus building compared to other short read sequencing technologies.

Overall, the addition of full-genome sequencing to more routine diagnostic use will increase the

available knowledge regarding sequence diversity and allow for improved tracking of viruses

and a better understanding of the genetic determinants of viral pathogenesis.

**Acknowledgments**

The authors acknowledge Vanessa Gauthiersloan, Samantha Day and Erich Linnemann from the

Poultry Diagnostic & Research Center, University of Georgia, Clara Kienzle-Dean from the

Southeastern Cooperative Wildlife Disease Study, University of Georgia, and Pablo Pinyero

from the Veterinary Diagnostic Lab, Iowa State University for their technical help.

**Declaration of conflicting interests**

The authors declared no potential conflicts of interest with respect to the research, authorship,

and/or publication of this article.

**Funding**

The project described was supported by Agriculture and Food Research Initiative Competitive

Grant no. 2018-67015-28306 from the USDA National Institute of Food and Agriculture. This

project has been funded in part with funds from the National Institute of Allergy and Infectious

Diseases, National Institutes of Health, Department of Health and Human Services, under Center

of Excellence of Influenza Research and Surveillance (CEIRS) under contract number

HHSN272201400004C (SMT).

**Supplementary material**

Supplementary material for this article is available online.

**References**

1. Sartore S, et al. The effects of control measures on the economic burden associated with
epidemics of avian influenza in Italy. Poult Sci 2010;89:1115–1121.
546
547 2. Butt SL, et al. Rapid virulence prediction and identification of Newcastle disease virus
548     genotypes using third-generation sequencing. Virol J 2018;15:179.
549
550 3. Clavijo A, et al. An improved Real-Time Polymerase Chain Reaction for the Simultaneous
551     Detection of All Serotypes of Epizootic Hemorrhagic Disease Virus. J Vet Diagn Invest
552     2010;22:588–593.
553
554 4. Cavanagh, D and Naqi SA. Infectious bronchitis. In: Diseases of Poultry. 11th ed. Ames,
555     Iowa: Iowa State Univer Pr, 2003:101–119.
556
557 5. Zhang J, et al. High-throughput whole genome sequencing of Porcine reproductive and
558     respiratory syndrome virus from cell culture materials and clinical specimens using next-
559     generation sequencing technology. J Vet Diagn Invest 2016;29:41–50.
560
561 6. Woolhouse MEJ, et al. Assessing the Epidemic Potential of RNA and DNA Viruses. Emerg
562     Inf Dis 2016;22:2037–2044.
563
564 7. Rubio L, et al. Genetic variability and evolutionary dynamics of viruses of the family
565     Closteroviridae. Front. Microbiol. 2013;4:151–151.
566
567 8. Read SJ, et al. Molecular techniques for clinical diagnostic virology. J Clin Pathol
568     2000;53:502–506.
569
570 9. Chan RWY, et al. Tissue tropism of swine influenza viruses and reassortants in ex vivo
571     cultures of the human respiratory tract and conjunctiva. J Virol 2011;85:11581–11587.
572
573 10. Flint, J. Synthesis of RNA from RNA templates. In: Principles of Virology. 4th ed.
574     Washington, DC: ASM Press, 2015: 157–184.
575
576 11. Gilchrist CA, et al. Whole-genome sequencing in outbreak analysis. Clin Microbiol Rev
577     2015;28:541–563.
578
579 12. Cobo F. Application of molecular diagnostic techniques for viral testing. Open Virol J
580     2012;6:104–114.
581
582 13. Gullett JC, et al. Quantitative Nucleic Acid Amplification Methods for Viral Infections. Clin
583     Chem 2015;61:72–78.
584
585 14. Ranjan R, et al. Analysis of the microbiome: Advantages of whole genome shotgun versus
586     16S amplicon sequencing. Biochem Biophys Res Commun 2016;469:967–977.
587
588 15. Kasibhatla, SM, et al. Analysis of next-generation sequencing data in virology- Opportunities
589     and challenges. IntechOpen 2016; DOI: 10.5772/61610.

590
591  16. Head SR, et al. Library construction for next-generation sequencing: overviews and
592      challenges. BioTechniques 2014;56:61–64
593
594  17. Marston DA, et al. Next generation sequencing of viral RNA genomes. BMC Genomics
595      2013;14:444.
596
597  18. Quick J, et al. Real-time, portable genome sequencing for Ebola surveillance. Nature
598      2016;530:228–232.
599
600  19. Greninger AL, et al. Rapid metagenomic identification of viral pathogens in clinical samples
601      by real-time nanopore sequencing analysis. Genome Med 2015;7:99.
602
603  20. Kilianski A, et al. Use of Unamplified RNA/cDNA-Hybrid Nanopore Sequencing for Rapid
604      Detection and Characterization of RNA Viruses. Emerg Infect Dis 2016;22:1448–1451.
605
606  21. Beato MS, et al. Identification and genetic characterization of bovine enterovirus by
607      combination of two next generation sequencing platforms. J Virol Methods 2018;260:21–25.
608
609  22. Peserico A, et al. Diagnosis and characterization of canine distemper virus through
610      sequencing by MinION nanopore technology. Sci Rep 2019;9:1714–1714.
611
612  23. Mahlum CE, et al. Detection of bovine viral diarrhea virus by TaqMan® reverse transcription
613      polymerase chain reaction. J Vet Diagn Invest 2002;14:120–125.
614
615  24. Kim D, et al. Centrifuge: rapid and sensitive classification of metagenomic sequences.
616      Genome Res 2016;26:1721–1729.
617
618  25. Butt SL, et al. Real-time, MinION-based, amplicon sequencing for lineage typing of
619      infectious bronchitis virus from upper respiratory samples. J Vet Diagn Invest 2019.
620      Accepted.
621
622  26. Yeşilbağ K, et al. Variability and Global Distribution of Subgenotypes of Bovine Viral
623      Diarrhea Virus. Viruses 2017;9:128.
624
625  27. Piewbang C, et al. Genetic and evolutionary analysis of a new Asia-4 lineage and naturally
626      recombinant canine distemper virus strains from Thailand. Sci Rep 2019;9:3198.
627
628  28. Brown-Joseph T, et al. Identification and characterization of epizootic hemorrhagic disease
629      virus serotype 6 in cattle co-infected with bluetongue virus in Trinidad, West Indies. Vet
630      Microbiol 2019;229:1–6.
631
632  29. Valastro V, et al. S1 gene-based phylogeny of infectious bronchitis virus: An attempt to
633      harmonize virus classification. Infect Genet and Evol 2016;39:349–364.
634

30. Shi W, et al. A complete analysis of HA and NA genes of influenza A viruses. PloS One
2010;5:e14454–e14454.

31. Kang H, et al. Geographic distribution and molecular analysis of porcine reproductive and
respiratory syndrome viruses circulating in swine farms in the Republic of Korea between
2013 and 2016. BMC Vet Res 2018;14:160.

32. O'Donnell CR, et al. Error analysis of idealized nanopore sequencing. Electrophoresis
2013;34:2137–2144.

33. Breitwieser, FP, et al. Pavian: interactive analysis of metagenomics data or microbiome
studies and pathogen identification. Bioinformatics 2019.

34. Thompson JD, et al. CLUSTAL W: improving the sensitivity of progressive multiple
sequence alignment through sequence weighting, position-specific gap penalties and weight
matrix choice. Nucleic Acids Res 1994;22:4673–4680.

35. Tamura K. Estimation of the number of nucleotide substitutions when there are strong
transition-transversion and G+C-content biases. Mol Biol Evol 1992;9:678–687.

36. Kumar S, et al. MEGA X: Molecular Evolutionary Genetics Analysis across Computing
Platforms. Mol Biol Evol 2018;35:1547–1549.

37. Vilček Š, et al. Pestiviruses isolated from pigs, cattle and sheep can be allocated into at least
three genogroups using polymerase chain reaction and restriction endonuclease analysis.
Arch of Virol 1994;136:309–323.

38. Martella V, et al. Genotyping canine distemper virus (CDV) by a hemi-nested multiplex PCR
provides a rapid approach for investigation of CDV outbreaks. Vet Microbiol 2007;122:32–
42.

39. Sun F, et al. Molecular Typing of Epizootic Hemorrhagic Disease Virus Serotypes by One-
Step Multiplex RT-PCR. J Wildl Dis 2014;50:639–644.

40. J Woolhouse ME, et al. RNA Viruses: A case study of the biology of emerging infectious
diseases. Microbiol Spectr 2013;1:10.1128/microbiolspec.OH-0001–2012.

41. Ginocchio CC, et al. Current best practices for respiratory virus testing. J Clin Microbiol
2011;49:S44–S48.

42. Wostenberg DJ, et al. Evidence of two cocirculating canine distemper virus strains in
mesocarnivores from northern Colorado, USA. J. of Wildlife Diseases 2018;54:534-543.

43. Ahasan MS, et al. Complete genome sequence of Mobuck Virus isolated from a Florida
white-tailed deer (*Odocoileus virginianus*). Microbiol Resour Announc 2019;8:e01324–
01318.

44. Pfeffermann K, et al. Chapter Four - Morbillivirus Pathogenesis and Virus–Host Interactions.
In: Advances in Virus Research. eds. Kielian M, Mettenleiter TC and Roossinck MJ, Vol.
100: Academic Press, 2018:75–98.

45. Jain M, et al. Improved data analysis for the MinION nanopore sequencer. Nat Methods
2015;12:351–356.

46. McNaughton AL, et al. Illumina and Nanopore methods for whole genome sequencing of
hepatitis B virus (HBV). Sci Rep 2019;9:7081.

47. Jain M, et al. MinION Analysis and Reference Consortium: Phase 2 data release and analysis
of R9.0 chemistry. F1000Res 2017;6:760–760.

48. Magi A, et al. Characterization of MinION nanopore data for resequencing analyses. Brief
Bioinform 2016;18:940–953.

49. Goldstein EJ, et al. Integrating patient and whole-genome sequencing data to provide insights
into the epidemiology of seasonal influenza A (H3N2) viruses. Microb Genom
2018;4:e000137.

50. Pallen MJ, et al. High-throughput sequencing and clinical microbiology: progress,
opportunities and challenges. Curr Opin in Microbiol 2010;13:625–631.

51. Runckel C, et al. Identification and manipulation of the molecular determinants influencing
poliovirus recombination. PLOS Pathog 2013;9:e1003164.

52. Su S, et al. Epidemiology, genetic recombination, and pathogenesis of coronaviruses. Trends
in Microbiol 2016;24:490–502.

53. Park E-S, et al. Identification of a natural recombination in the F and H genes of feline
morbillivirus. Virology 2014;468–470:524–531.

**Tables**

714 **Table 1**. Sample background information and reverse-transcription real-time PCR (RT-rtPCR) results.

| Virus | Original Host | Propagation System | RT-rtPCR (Ct) | Collection | |
|---|---|---|---|---|---|
| | | | | State (USA) | Year |
| BVDV | BVDV positive CPAE cells | CPAE | 24.90 | N/A | N/A |
| CDV | North American raccoon | Vero-Dog SLAM | 20.35 | Kentucky | 2018 |
| EHDV | White-tailed deer | CPAE | 15.15 | Georgia | 2016 |
| IBV | N/A | ECE | 12.48 | N/A | N/A |
| Canine-origin IAV | Dog | ECE | 16.94 | Georgia | 2015 |
| Swine-origin IAV | Hampshire-cross pig | MDCK | ND | Georgia | 2019 |
| PRRSV | N/A | MARC-145 | ND | N/A | N/A |

715 BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; Ct = cycle threshold; CPAE = cattle pulmonary artery
716 endothelial; ECE = embryonated chicken eggs; EHDV = epizootic hemorrhagic disease virus; IAV = influenza A virus; IBV =
717 infectious bronchitis virus; MDCK = Madin-Darby Canine Kidney; N/A = not applicable; ND = not determined; PRRSV = porcine
718 reproductive and respiratory syndrome virus.
719
720
721 Table 2. Reverse-transcription PCR (RT-PCR) primer information for Sanger sequencing.

| Virus | Target Region | Primer Name | Primer Sequence (5′ → 3′) | Amplicon Size (bp) | Reference |
|---|---|---|---|---|---|
| BVDV | 5′ UTR | 324 | ATGCCCT/ATAGTAGGACTAGCA | 288 | [37] |
| | | 326 | TCAACTCCATGTGCCATGTAC | | |
| CDV | *H* | 204 (+) | GAATTCGACTTCCGCGATCTCC | 1,160 | [38] |
| | | 232 (-) | TAGGCAACACCACTAATTTRGACTC | | |
| EHDV | *VP2* | EHDV-2F | TGGTGAAAATACGGTATATAACC | 246 | [39] |
| | | EHDV-2R | GTTCAAATTCATCTGGGCTCATACT | | |

722 BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; EHDV = epizootic hemorrhagic disease virus; *H* =
723 hemagglutinin; UTR = untranslated region.
724

725    **Table 3**. Sequencing data from Porechop and Centrifuge.

| Virus | Estimated Total Run Time (HH:MM) | Porechop | | Viral Reads (%) | Host Reads (%) |
|---|---|---|---|---|---|
| | | Total Reads | Total Bases | | |
| BVDV & EHDV | 8:20 | 756,373 | 299,505,703 | 12.7 | 75.2 |
| CDV | 3:20 | 183,206 | 63,116,488 | 2.10 | 80.4 |
| CDV* | 47:37 | 75,492 | 33,681,975 | 0.97 | 45.8 |
| IBV | 7:40 | 7,523 | 11,361,873 | 63.3 | 2.50 |
| Canine IAV | 2:45 | 745,382 | 341,878,915 | 36.6 | 53.2 |
| Swine IAV | 5:20 | 1,173,058 | 450,544,038 | 31.1 | 63.4 |
| PRRSV* | 47:37 | 28,418 | 18,889,591 | 57.1 | 14.3 |

726    BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; EHDV = epizootic
727    hemorrhagic disease virus; IAV = influenza A virus; IBV = infectious bronchitis virus; PRRSV
728    = porcine reproductive and respiratory syndrome virus.
729    * Libraries were prepared and sequenced at Virginia Tech University.
730

**Table 4**. Consensus results for coding sequences (CDS) of lineage-typing regions for each virus.

| Virus | Lineage-typing Region | Consensus Length (bp) | BLASTn Top Hit | Accession | Lineage | Pairwise Identity (%) | Mean Depth of Coverage | Minimum Depth of Coverage | Maximum Depth of Coverage |
|---|---|---|---|---|---|---|---|---|---|
| BVDV | 5□ UTR | 381 | BVDV-2 95-1501 | MH231130 | ND | 97.91 | 12,395.6 | 10,115 | 13,980 |
|  | $N^{pro}$ | 504 | BVDV-2 McCart_c | MH806438 | 2a | 96.23 | 14,783.5 | 13,286 | 15,908 |
| CDV | $H$ | 1,824 | CDV A75/17† | AF164967 | America II | 96.49 | 817.3 | 151 | 1,739 |
| CDV* | $H$ | 1,824 | CDV A75/17† | AF164967 | America II | 96.49 | 105.1 | 33 | 176 |
| EHDV | $VP2$ | 2,949 | EHDV-2 OV617 | MK958997 | 2w | 99.90 | 523.7 | 43 | 896 |
| IBV | $S1$ | 3,489 | Avian coronavirus strain Ma5 | KY626045 | GI-L4 | 100.0 | 287.5 | 219 | 350 |
| Canine IAV | $HA$ | 1,701 | Influenza A virus (A/canine/Georgia/101875/ 2015(H3N2))† | MF173286 | H3g1 | 100.0 | 13,981.0 | 960 | 21,463 |
|  | $NA$ | 1,410 | Influenza A virus (A/canine/Georgia/95391/2015 (H3N2))† | KX570998 | N2g2 | 99.79 | 22,968.0 | 674 | 37,816 |
| Swine IAV | $HA$ | 1,698 | Influenza A virus (A/swine/Ohio/18TOSU1194/2018(H1N2))† | MN198216 | H1g4 | 99.65 | 2,360.7 | 424 | 3,148 |
|  | $NA$ | 1,410 | Influenza A virus (A/swine/Ohio/18TOSU1194/2018(H1N2))† | MN198218 | N2g2 | 99.79 | 4,752.8 | 517 | 6,928 |
| PRRSV* | $ORF5$ | 603 | PRRSV VR2385 | JX044140 | Type 2, Lineage 5 | 99.67 | 546.2 | 470 | 588 |

BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; EHDV = epizootic hemorrhagic disease virus; Gx-Lx= genotype-lineage; $H$ = hemagglutinin; $HA$ = hemagglutinin; IAV = influenza A virus; IBV = infectious bronchitis virus; $NA$ = neuraminidase ND = not determined; $N^{pro}$ = N-terminal protease; $ORF$ = open reading frame; PRRSV = porcine reproductive and respiratory syndrome virus; $S1$ = Spike 1.

* Libraries were prepared and sequenced at Virginia Tech University.

† BLASTn results showed 2-29 sequence alignments with identical Bit-Scores, query coverage, and pairwise identities to other isolates. The provided "top hit" was included in those alignments.

741 **Table 5**. Consensus results of whole genome coding sequences (CDS) for each virus.

| Virus | Segment | Reads Per Consensus | Consensus Length (bp) | BLASTn Top Hit | Accession | Pairwise Identity (%) | Mean Depth of Coverage | Minimum Depth of Coverage | Maximum Depth Coverage |
|---|---|---|---|---|---|---|---|---|---|
| BVDV | | 54,561 | 11,689 | BVDV-2 95-1501 | MH231130 | 97.37 | 4,277.3 | 518 | 15,908 |
| CDV | | 3,428 | 15,477 | CDV A75/17 | AF164967 | 97.01 | 299.6 | 26 | 1,739 |
| CDV* | | 717 | 13,957 | CDV A75/17 | AF164967 | 96.99 | 72.1 | 20 | 176 |
| EHDV | *VP1* | 2,091 | 3,909 | EHDV-2 OV617 | MK958997 | 99.97 | 576.4 | 231 | 1,333 |
| | *VP2* | 1,423 | 2,949 | EHDV-2 OV617 | MK958997 | 99.90 | 523.7 | 43 | 896 |
| | *VP3* | 1,807 | 2,700 | EHDV-2 OV215 | MF688818 | 99.93 | 606.3 | 100 | 1,346 |
| | *VP4* | 1,602 | 1,935 | EHDV-2 OV215 | MF688819 | 99.95 | 625.3 | 218 | 1,205 |
| | *NS1* | 3,598 | 1,656 | EHDV-2 OV215 | MF688823 | 99.88 | 2,295.5 | 185 | 2,711 |
| | *VP5* | 3,249 | 1,584 | EHDV-2 OV617 | MK959001 | 100.0 | 1,473.6 | 168 | 2,693 |
| | *VP7* | 3,415 | 1,050 | EHDV-2 OV215 | MF688822 | 100.0 | 1,751.5 | 117 | 2,804 |
| | *NS2* | 6,485 | 903 | EHDV-2 OV617† | MK959005 | 99.91 | 2,884.1 | 392 | 3,930 |
| | *VP6* | 15,516 | 1,080 | EHDV-2 OV617 | MK959002 | 99.91 | 6,335.3 | 196 | 13,629 |
| | *NS3* | 3,512 | 687 | EHDV-2 OV617 | MK959006 | 100.0 | 2,879.7 | 181 | 3,409 |
| IBV | | 4,695 | 26,574 | Avian coronavirus strain Ma5 | KY626045 | 100.0 | 416.3 | 158 | 1,011 |
| Canine IAV | *PB2* | 50,753 | 2,280 | Influenza A virus (A/canine/Florida/269770/2015(H3N2)) | MF173191 | 100.0 | 19,415.2 | 1,136 | 30,101 |
| | *PB1* | 38,930 | 2,274 | Influenza A virus (A/canine/Florida/269770/2015(H3N2))† | MF173194 | 100.0 | 13,406.6 | 2,339 | 21,726 |
| | *PA* | 28,755 | 2,151 | Influenza A virus (A/canine/Florida/269770/2015(H3N2)) | MF173220 | 100.0 | 1,1662.4 | 923 | 17,815 |
| | *HA‡* | 33,293 | 1,701 | Influenza A virus (A/canine/Georgia/101875/ 2015(H3N2))† | MF173286 | 100.0 | 13,981.0 | 960 | 21,463 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| Swine IAV | NP | 32,490 | 1,497 | Influenza A virus (A/canine/Georgia/95391/ 2015 (H3N2))† | KX571004 | 100.0 | 1,3119.1 | 2,544 | 19,511 |
| | NA‡ | 41,130 | 1,410 | Influenza A virus (A/canine/Georgia/95391/ 2015 (H3N2))† | KX570998 | 99.79 | 22,968.0 | 674 | 37,816 |
| | M1 & M2 | 39,862 | 982 | Influenza A virus (A/canine/Texas/2100186/ 2015(H3N2))† | MF173280 | 100.0 | 23,887.5 | 962 | 34,210 |
| | NEP & NS1 | 3,937 | 838 | Influenza A virus (A/canine/Georgia/101875 / 2015(H3N2))† | MF173122 | 100.0 | 2,291.5 | 524 | 2,971 |
| | PB2 | 41,421 | 2,280 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198223 | 99.74 | 9,835.2 | 223 | 33,925 |
| | PB1 | 14,908 | 2,274 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198222 | 99.74 | 2,303.0 | 352 | 3,438 |
| | PA | 26,431 | 2,151 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198221 | 99.72 | 7,941.8 | 417 | 20,275 |
| | HA‡ | 6,673 | 1,698 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198216 | 99.65 | 2,360.7 | 424 | 3,148 |
| | NP§ | 64,370 | 1,497 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198219 | 99.40 | 28,014.8 | 305 | 61,449 |
| | NA‡ | 8,519 | 1,410 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198218 | 99.79 | 4,752.8 | 517 | 6,928 |
| | M1 & M2 | 9,655 | 982 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198217 | 100.0 | 4,855.6 | 178 | 7,536 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | *NEP & NS1* | 3,276 | 838 | Influenza A virus (A/swine/Ohio/18TOSU1194/ 2018(H1N2))† | MN198220 | 99.88 | 1,637.4 | 283 | 2,076 |
| PRRSV | | 16,216 | 14,636 | PRRSV VR2385 | JX044140 | 99.84 | 2,045.4 | 27 | 13,535 |

BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; EHDV = epizootic hemorrhagic disease virus; *HA* =
hemagglutinin; IAV = influenza A virus; IBV = infectious bronchitis virus; *M* = matrix protein; *NA* = neuraminidase; *NEP* = nuclear
export protein; *NP* = nucleocapsid; *NS* = nonstructural protein; *PA* = polymerase; *PB* = polymerase; PRRSV = porcine reproductive and
respiratory syndrome virus.
* Libraries prepared and sequenced at Virginia Tech University.
† BLASTn results showed 2-77 sequence alignments with identical Bit-Scores, query coverage, and pairwise identities to other isolates.
The provided "top hit" was included in those alignments.
‡ Repeated from Table 4 for completeness of genome assessment.
§ Due to a large number of reads, only half the reads classified as the NP gene were used for reference-based consensus building.
751

752 **Table 6**. Sanger and MinION sequencing consensus pairwise identity results.

| Virus | Target Region | MinION | | | Sanger | | | Sanger vs. MinION Pairwise Identity (%) |
|---|---|---|---|---|---|---|---|---|
| | | Consensus Length (bp) | BLASTn Top Hit | Pairwise Identity (%) | Consensus Length (bp) | BLASTn Top Hit | Pairwise Identity (%) | |
| BVDV | 5□ UTR | 381 | BVDV-2 95-1501 (MH231130) | 97.91 | 268 | BVDV-2 95-1501 (MH231130) | 99.22 | 100.0 |
| CDV | *H* | 1,824 | CDV A75/17 (AF164967) | 96.49 | 923 | CDV THA/VG (JX886780)* | 96.53 | 100.0 |
| EHDV | *VP2* | 2,949 | EHDV-2 OV617 (MK958998) | 99.90 | 158 | EHDV-2 OV617 (MK958998)* | 100.0 | 100.0 |

753 BVDV= bovine viral diarrhea virus; CDV = canine distemper virus; EHDV = epizootic hemorrhagic disease virus; *H* =
754 hemagglutinin; UTR = untranslated region.
755 * BLASTn results showed 5-12 sequence alignments with identical Bit-Scores, query coverage, and pairwise identities to other
756 isolates. The provided "top hit" was included in those alignments.
757

**Figure Legends**

**Figure 1.** Sankey diagram displaying the top ten nodes per taxonomic classification level. Centrifuge v.1.0.4[24] Kraken-style reports

from analyzing sequences from canine distemper virus (CDV) with (a) pipeline containing CDV sequences and (b) novel virus

simulation using a pipeline devoid of CDV sequences in custom Centrifuge indices were visualized with Pavian v.1.0.[33] Number of

reads assigned to taxonomic classification correspond with the width of the flow and are listed above each node. Reads hitting to the

bovine and African green monkey genomes were excluded. Abbreviations are: D = domain; F = family; G = genus; P=phylum; S =

species.

**Figure 2.** Phylogenetic analysis of morbilliviruses of (a), whole genome CDS and (b), *P* gene CDS after a data analysis of CDV

sequences using a pipeline devoid of CDV reads to simulate novel virus detection and identification. Each sequence is designated by

species abbreviation and GenBank accession number. The sequence denoted with an asterisk was generated from the data analysis

pipeline with MinION sequencing. The evolutionary history was inferred by using the Maximum Likelihood method and Tamura 3-

parameter model.[35] These analyses involved 19 nucleotide sequences and a total of 14,578 positions (a) or 1,460 positions (b) in the

final dataset. All positions containing gaps and missing data were eliminated. Evolutionary analyses were conducted in MEGA X.[36]

Abbreviations are: CeMV = cetacean morbillivirus; FeMV = feline morbillivirus; MeV = measles virus; PDV = phocine distemper

virus; PPRV = peste-des-pestits-ruminants virus; RPV = rinderpest virus.