Mutation of a PER2 phosphodegron perturbs the circadian phosphoswitch

Shusaku Masuda[1], Rajesh Narasimamurthy[2], Hikari Yoshitane[1],
Jae Kyoung Kim[3], Yoshitaka Fukada[1,*], and David M. Virshup[2,4,*]

[1] Department of Biological Sciences, School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan.

[2] Programme in Cancer and Stem Cell Biology, Duke-NUS Medical School, 169857 Singapore

[3] Department of Mathematical Sciences, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Korea

[4] Department of Pediatrics, Duke University Medical Center, Durham, NC 27710

* Correspondence should be addressed to:

Dr. David M. Virshup
Programme in Cancer and Stem Cell Biology,
Duke-NUS Medical School,
169857 Singapore
E-mail: david.virshup@duke-nus.edu.sg

Dr. Yoshitaka Fukada
Department of Biological Sciences, School of Science,
The University of Tokyo,
Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan
Tel: +81-3-5841-4381 Fax: +81-3-5841-0763
E-mail: sfukada@mail.ecc.u-tokyo.ac.jp

Abstract

Casein kinase 1 (CK1) plays a central role in regulating the period of the circadian clock. In mammals, PER2 protein abundance is regulated by CK1-mediated phosphorylation and proteasomal degradation. On the other hand, recent studies have questioned whether the degradation of the core circadian machinery is a critical step in clock regulation. Prior cell-based studies found that CK1 phosphorylation of PER2 at Ser478 recruits the ubiquitin E3 ligase β-TrCP, leading to PER2 degradation. Creation of this phosphodegron is regulated by a phosphoswitch that is also implicated in temperature compensation. However, *in vivo* evidence that this phosphodegron influences circadian period is lacking. Here, we generated and analyzed PER2-Ser478Ala knock-in mice. The mice showed longer circadian period in behavioral analysis. Molecularly, mutant PER2 protein accumulated in both the nucleus and cytoplasm of the mouse liver, while *Per2* mRNA levels were minimally affected. Nuclear PER1, CRY1 and CRY2 proteins also increased, probably due to stabilization of PER2-containing complexes. In mouse embryonic fibroblasts derived from PER2-Ser478Ala::LUC mice, three-phase decay and temperature compensation of the circadian period was perturbed. These data provide direct *in vivo* evidence for the importance of phosphorylation-regulated PER2 stability in the circadian clock and validate the phosphoswitch in a mouse model. (200 words)

Introduction

The cell-autonomous clock systems oscillating with a period of ~24 h have evolved to coordinate physiological functions with daily environmental changes [1]. The molecular mechanism at the heart of the circadian clock is based on transcriptional/translational feedback loops that consist of clock genes and the encoded clock proteins. In the mammalian core clockwork, two proteins, circadian locomotor output cycles kaput (CLOCK) and brain muscle arnt-like 1 (BMAL1), form a heterodimer that binds to the cis-regulatory DNA element E-boxes to activate transcription of *Period* (*Per*) and *Cryptochrome* (*Cry*) genes. Translated PER and CRY proteins form a multimeric complex with casein kinase 1 (CK1δ/ε) in the cytoplasm, and then translocate to the nucleus, where they inhibit the transcriptional activity of CLOCK-BMAL1 complex [2–4]. The key molecular components of the clock in all organisms are subject to various posttranslational modifications, most notably phosphorylation and ubiquitination, and disruption of these posttranslational modifications can cause circadian dysfunction [5, 6]

Casein kinase 1 (CK1) is an evolutionarily conserved core clock component [6]. The role of CK1 in the clock is complex. Mutations in CK1, first identified in hamsters [7, 8] and Drosophila [9, 10], can cause both period shortening and lengthening, while pharmacologic inhibition of CK1 consistently causes period lengthening in animals [11–13] and in plants [14]. Intriguingly, the kinase activity of CK1 on synthetic peptides is temperature-insensitive [12, 15]. This is consistent with a central role for CK1 in temperature compensation of the circadian period, the mechanism that maintains a stable circadian period despite changes in ambient temperature [16–18]. In mammals, CK1 delta and epsilon (CK1δ/ε) are essential kinases for the circadian clock [19]. The first identified mammalian circadian clock mutant, the *tau* hamster, harbors a semidominant mutation in CK1ε, and *tau* homozygous animals exhibit markedly shortened locomotor rhythms with a period of 20 h [7, 8]. In humans, mutations in CK1δ also cause Familial Advanced Sleep Phase (FASP) syndrome and shorten the circadian period [20, 21].

The biochemical function of CK1δ/ε in the clock arises from its stoichiometric tight interaction with and phosphorylation of PER proteins [4, 8, 22, 23]. One major consequence of PER2 phosphorylation by CK1δ/ε is the regulation of PER2 stability via the phosphoswitch [11, 17, 18, 24–27]. The PER2 phosphoswitch requires at least two phosphorylation domains. Cell-based and

biochemical studies have established that Ser478 of PER2 protein is phosphorylated by CK1δ/ε, and Ser478 phosphorylation creates a phosphodegron that recruits an E3 ubiquitin ligase, beta-transducin repeat-containing homologue protein (β-TrCP) [11, 28]. β-TrCP is a substrate recognition subunit of the Skp1-Cullin-F-box (SCF) complex that catalyzes formation of a polyubiquitin chain on a substrate targeted for proteasomal degradation. The activity of CK1δ/ε on the S478 phosphodegron is controlled by a second PER2 phosphorylation domain called the FASP region. In this region, CK1δ/ε phosphorylates Ser659, followed by rapid phosphorylation of additional downstream serines in the motif pSxxSxxS. Cell based assays have demonstrated that this multi-phosphorylation of the FASP domain stabilizes PER2 by inhibiting phosphorylation of the S478 phosphodegron [17, 18, 24, 26]. Indeed, the FASP region is so named because a mutation of human PER2 Ser662 (corresponding to Ser659 in mouse) causes FASP syndrome [25, 29]. In the phosphoswitch model, the short-period phenotype common to *tau* and FASP mutations has been attributed to enhanced phosphorylation of the β-TrCP site leading to more rapid PER2 degradation. To date, however, there is no *in vivo* evidence that the β-TrCP site plays a role in determining the circadian period. To test this, we generated two knock-in mouse lines carrying Ser-to-Ala mutation at position 478 of PER2 protein. We find that PER2-Ser478Ala protein is stabilized in the mouse liver, and the mutation caused lengthening of the circadian period of mouse behavioral rhythms. These data provide the first genetic evidence for the importance of the β-TrCP site in the time-keeping mechanism of the circadian clock *in vivo*.

Result

The PER2-S478A mutation lengthens the period of behavioral rhythms

The phosphoswitch model holds that PER2 stability is bidirectionally regulated by two key phosphorylation sites, the β-TrCP site and the FASP site (Figure 1A). The model also predicts that a mutation of PER2 at the β-TrCP site should result in a more stable protein and a longer circadian period length *in vivo*. To test this hypothesis, we used CRISPR-Cas9 to generate a mouse with PER2 Ser478Ala (PER2-S478A)(Figure 1A). PER2-S478A homozygous mice were morphologically normal and fertile and were born in the expected Mendelian ratio. The mutant mice showed robust wheel-running activity rhythms under a 12 h light:

12 h dark (LD) cycle, and no apparent abnormality was observed in the light
entrainment (Figure 1B). On the other hand, when transferred to constant dark
(DD) condition, the mutant mice showed ~1 h longer period of the behavioral
rhythms (24.58 ± 0.08 h) than wild-type (23.67 ± 0.06 h) (Figure 1C). This is the
first *in vivo* demonstration that the Ser478 phosphodegron of PER2 determines
the circadian period.

Excessive accumulation of PER2-S478A mutant protein in the mouse liver

When over-expressed in cultured cells, PER2-S478A mutant was more stable
than wild-type PER2 protein [17]. Here we examined the abundance of
endogenous *Per2* mRNA and PER2 protein in the livers of PER2-S478A mice.
Livers were harvested at six time points during the second day in DD condition.
Per2 transcripts showed the expected circadian oscillation in both wild-type and
PER2-S478A mice, with no significant difference observed between the
genotypes (Figure 2A). In contrast, peak levels of PER2 protein in both the
nucleus and cytoplasm were ~2-fold higher at CT22 in PER2-S478A mice (Figure
2B and 2C). Thus, the S478A mutation stabilizes PER2 protein *in vivo* similar to
what was seen in *in vitro* over-expression assays [17]. It should be noted that,
although the peak levels of PER2-S478A protein were higher, PER2-S478A
protein was still almost undetectable at CT10 and CT14, suggesting that
phosphorylation of Ser478 is required for temporal degradation of PER2, and that
other degradation mechanism(s) may be contributing as well [30].

The PER2-S478A mutation stabilizes CRY proteins

PER2 has been proposed to be rate-limiting for *in vivo* formation of a stable
multi-protein complex containing PER1, CRY1/2 and CK1δ/ε [2, 4] that
assembles in the cytoplasm and then translocates to the nucleus. To assess the
consequence of stabilization of PER2-S478A protein on these core circadian
proteins, we examined the expression of a series of clock proteins in the livers of
wild-type and mutant mice. At the time of peak PER2 abundance (CT22, Figure
2B, C), the cytosolic protein levels of PER1 and CRY2 were also increased
(Figure 3A). Lagging by several hours, the nuclear abundance of PER1, CRY1
and CRY2 was also increased in the livers of PER2-S478A mice (Figure 3B).
Thus, stabilized PER2 increases the abundance of the full circadian repressor
complex.

Several studies proposed that PER2 stabilizes CRY proteins by interfering
with the interaction between CRYs and FBXL3, an E3 ubiquitin ligase that directs

degradation of CRYs [31–33]. Up-regulation of CRY protein levels in the PER2-
S478A mutant liver (Figure 3A, B) suggests that CRYs are stabilized in response
to elevation of PER2 protein levels. To test this, we measured the half-lives of
CRY1::LUC and CRY2::LUC fusion proteins in HEK 293T cells by monitoring the
bioluminescence decay after treatment of cycloheximide (CHX) inhibiting new
protein translation. When PER2-S478A was co-expressed, CRY1::LUC and
CRY2::LUC were stabilized in a dose-dependent manner and to the same extent
as wild-type PER2 protein (Figure S1). Thus, stabilization of PER2 appears to
stabilize CRY proteins in both cultured cells and in mouse liver.

PER abundance has been proposed to be the rate-limiting step in the
assembly and nuclear entry of the repressor complex [2, 4, 34]. Notably, the
nuclear protein levels of CRY1 and CRY2 were still high at CT10 (Figure 3B)
when a large proportion of PER2-S478A protein was degraded (Figure 2B). This
suggests that CRY proteins can repress CLOCK-BMAL1 transcriptional activity
even in the absence of PER proteins, consistent with a previous study [35]. In
fact, the mRNA expression profiles of typical E-box-regulated genes, such as
albumin D-site binding protein (*Dbp*), *Nr1d1* (also known as *Rev-erbA*) and *Nr1d2*
(also known as *Rev-erbB*), were phase-delayed in the PER2-S478A mutant liver
(Figure 3C). The delayed expression profiles were also observed in RRE-
regulated genes *Bmal1, Clock,* and *Cry1* (Figure 3C). These profiles are
consistent with the longer behavioral rhythms of the PER2-S478A mice.
Concomitantly, nuclear expression profiles of DBP and NR1D1 proteins were
delayed in the mutant mouse liver (Figure 3B). Overall, our data indicate that the
PER2-S478A mutation results in accumulation of circadian repressor complex in
the nucleus, leading to delayed de-repression of the CLOCK-BMAL1 activator.

The S478A mutation of PER2 compromises temperature compensation
The phosphoswitch model arose from the observation that PER2::LUC
degradation occurred in three phases [17]: an initial rapid decay (the first phase),
a plateau-like slow decay (the second phase), and finally, a more rapid terminal
decay (the third phase). This was visualized by continuous monitoring of
bioluminescence signals from PER2::LUC knock-in MEFs. To investigate how the
S478A mutation affects the three-phase degradation *in vivo*, we in parallel
introduced the S478A mutation into the PER2::LUC allele (PER2-S478A::LUC)
encoding a PER2 fusion protein with firefly luciferase [36]. In the analysis of the
wheel running activity rhythms, PER2-S478A::LUC mice exhibited a significantly

longer circadian period than PER2::LUC control mice (Figure 4A, B), similar to

that seen in the PER2-S478A mice. We established mouse embryonic fibroblasts

(MEFs) from both PER2-S478A::LUC and control PER2::LUC mice.

We then tested if disruption of the phosphoswitch by the S478A mutation

affects the temperature compensation, which is a key feature of the circadian

clock [37]. While the rate of most enzymatic reactions doubles or triples with every

10°C increase in temperature (*i.e.*, temperature coefficient, Q_{10} = 2-3), circadian

rhythms are relatively stable in the face of environmental temperature alterations,

and in fact are often over-compensated, with a $Q_{10} < 1.0$. Indeed, in the

PER2::LUC MEFs, as the temperature decreased from 37°C to 30°C the clock

sped up ~1.5 hours, giving a $Q_{10} = 0.80$ (Figure 4C, D). However, in the PER2-

S478A::LUC MEFs, the clock only sped up ~0.5 hours, for a Q_{10} of 0.92 (Figure

4C, D). Thus, temperature compensation was compromised by the S478A

mutation, underscoring the importance of this site in the phosphoswitch-regulated

temperature compensation.

Finally, we examined whether mutation of the phosphodegron can disrupt the

previously observed three-phase decay of PER2::LUC. When protein translation

was inhibited by addition of cycloheximide (CHX) at two distinct time points

(CT19.8 and CT21.4) in the PER2 accumulation phase, we again observed three-

phase degradation of endogenous PER2::LUC (Figure 4E). The duration of the

second phase was dependent on the time when CHX was added, as previously

reported [17]. At both time points, the three-phase degradation of PER2 protein

was markedly attenuated by the S478A mutation (Figure 4E), resembling what

was seen when CK1 was inhibited [17].

Collectively, these data demonstrate that the phosphodegron in PER2 plays

a key role in the phosphoswitch, temperature compensation and circadian period

in vivo.

Discussion

Robust genetic data indicate that CK1δ/ε controls the speed of the

circadian clock oscillation, for which multiple potential molecular mechanisms

have been proposed. These include nuclear-cytoplasmic shuttling of PER

proteins, phosphorylation of PER-associated proteins in the repression

complex, phosphorylation of CLOCK to decrease its DNA binding, and a

phosphoswitch to regulate PER protein stability [4, 17, 38]. We previously

demonstrated that phosphorylation of the FASP region inhibited phosphorylation
of the β-TrCP phosphodegron, but it has not been experimentally addressed
whether the β-TrCP degron actually regulates the circadian period. In the
present study, we find that phosphorylation of the β-TrCP site controlling PER2
stability is an essential step in determining the period length *in vivo*. Mice with
the PER2-S478A mutation showed significantly longer behavioral rhythms than
control mice (Figure 1). The period lengthening effect (by ~1 h) is similar to
those observed in mice injected with a CK1 inhibitor [13] and in transgenic mice
carrying human PER2 S662D, a phospho-mimic mutant at the FASP site [25].
These data show that the β-TrCP site is important for the regulation of
behavioral rhythm by the phosphoswitch. Unlike what was observed in hPER2-
S662D transgenic mice [25], *Per2* mRNA levels were unaltered in the PER2-
S478A mutant. The peak levels of PER2 protein were increased in cytosol and
nucleus (Figure 2), consistent with disruption of the Ser478 phosphodegron.
Thus, this study provides a strong genetic evidence that phosphorylation-
regulated degradation of PER2 is indeed a key regulator of the clock speed.
CK1δ and CK1ε may not be the only kinases regulating PER2 stability. For
example, a chemical screening identified CK1α as a regulator of PER protein
stability [39]. This effect of CK1α may be indirect because CK1α binds to PER
proteins with a much lower affinity than CK1δ/ε does [38–40]. In *Drosophila*,
CK1α enhances PER degradation elicited by DBT (the CK1δ/ε homolog in
Drosophila) [41]. CK2 phosphorylates PER2 at Ser53 and promotes PER2
proteasomal degradation in both CK1δ/ε-dependent and -independent pathways
[42]. PKCα [43] and GRK2 [44] regulate the nuclear entry of PER2 and might
affect the stability of PER2. In this study, we showed that PER2-S478A mutation
attenuated CK1δ/ε-mediated degradation of PER2 protein and lengthened the
circadian period. Loss of function of CK1 in *Neurospora* also lengthened the
circadian period [45]. The clock genes are not orthologous across species in
eukaryotes, but architectures of the molecular clockworks are analogous.
FREQUENCY (FRQ), a negative component of the circadian clock in *Neurospora*,
is analogous to PER. CK1 phosphorylates FRQ and decreases its protein stability
as well [46]. In contrast, recent studies proposed that the binding affinity of CK1
for FRQ, instead of FRQ stability, might be a critical factor for the period
determination through phosphorylation of other clock proteins [47, 48]. Our data
establish a role for the phosphodegron in regulating circadian rhythms but do not

rule out an alternative model in which CK1-mediated phosphorylation of other
substrates also plays a role for regulation of clock timing.

In this study, we propose a model (Figure 5), in which PER2 Ser478
phosphorylation governs decay of PER/CRY-containing repressor complex that
determines restart timing of the E-box-dependent transcriptional activation. The
peak level of PER2 protein is regulated by the balance between the competing
phosphorylations of stabilizing (FASP) and destabilizing (β-TrCP) sites. The
PER2-S478A mutation disrupts this balance and results in excessive PER2
accumulation (Figure 2) that interferes with the degradation of PER1 and CRY1/2
proteins (Figure 3). The mature repressive complex enters the nucleus to inhibit
CLOCK-BMAL1. The nuclear PER proteins are then degraded, presumably in a
β-TrCP site-independent fashion. The remaining CRY proteins stay on CLOCK-
BMAL1 activator complex to keep repressing the transcriptional activity in the
absence of PERs [35, 50, 51], also contributing to the lengthened period
phenotype (Figure 1).

Temperature compensation (or over-compensation) of the circadian period
has been described for more than 60 years [37]. Mathematical models predicted
that temperature-dependent opposing reactions contribute to the temperature-
compensated clockwork [49], although the processes which virtually act in this
mechanism have not been understood. We previously demonstrated that the
PER2 phosphoswitch may act as the compensating process in mammalian
clockwork [17]. The phosphoswitch leads to two pools of PER2, one less stable
due to degron phosphorylation, and the other more stable due to FASP
phosphorylation inhibiting degron phosphorylation. Therefore, three-phase
degradation can be observed following CHX treatment in PER2::LUC MEFs [17].
Here, the three-phase degradation was compromised in PER2-S478A::LUC
MEFs, consistent with this model. The mutant MEFs showed also cellular rhythms
with a longer period at lower temperature. These results are consistent with the
idea that the β-TrCP site is more susceptible to phosphorylation at lower
temperature to make the circadian period compensated for temperature changes
[17].

Although mutation of the Ser478 phosphodegron increased PER2 stability,
PER2 was still degraded between CT22 and CT2 (Figure 2). Furthermore, the
three-phase degradation was indeed perturbed in the PER2-S478A mutant but
was not completely abolished (Figure 4E, left panels). These data suggest
alternative pathway(s) for PER2 degradation. One possible pathway is via an

additional β-TrCP site. A potential phosphodegron motif TpSGCSpS is conserved

in PER1 (aa121-126) and PER2 (aa92-97), whereas the primary β-TrCP degron

SpSGYGpS (aa477-482 in PER2) is not present in PER1 [52]. In PER2, the

phosphorylation of the primary degron plays a dominant role for β-TrCP-mediated

degradation, but a role of the second degron has been reported when the primary

degron is mutated [28]. In PER2-S478A mutant, this additional degron may

therefore contribute to the alternative degradation pathway. Second, it was

recently reported that oncoprotein MDM2 targets PER2 for proteasomal

degradation, independent of the phosphorylation status of PER2 protein [30].

Third, it is possible that β-TrCP recruited to phosphorylated PER1 might

contribute to PER2 degradation in the repressor complex. Future studies on

detailed mechanisms underlying the PER2 stability control will help to understand

the temperature compensation of the circadian clock.

Acknowledgements

We thank Dale Cowley of TransViragen/UNC Chapel Hill Animal Models

Core for design and production of the Ser478Ala mutant mice, and Dr. Atsu Aiba

in the University of Tokyo for their help with mouse embryo freezing and the

embryo transfer. This work was partially supported by Singapore's

NMRC/CIRG/1465/2017 to D.V., by Grants-in-Aid for Specially Promoted

Research to Y.F. (17H06096), and for Scientific Research (B) to H.Y.

(25440041) from MEXT of Japan, and by the PRIME from Japan Agency for

Medical Research and Development to H.Y. (17937210).

Author contributions statement

295   S.M., R.N., H.Y., Y.F., and D.V. designed the study. S.M. performed the

296 almost all experiments. R.N. performed the three-phase degradation assay and

297 J.K.K analyzed it. H.Y., Y.F., and D.V. supervised the whole project. S.M. and

298 H.Y. wrote the first draft of the manuscript. Y.F. and D.V. edited the manuscript.

299 All authors reviewed and approved the final manuscript.

Competing interests

302   The authors declare no competing interests.

Materials and methods

Mice

Mice used in this study (C57BL/6 background) were handled in accordance with the Guidelines for the Care and Use of Laboratory Animals at The University of Tokyo. Mice were housed in cages with free access to commercial chow (CLEA Japan) and tap water. The animals were maintained in a light-tight chamber at a constant temperature (23°C ± 1°C) and humidity (65% ± 10%).

Generation of PER2-S478A mice and PER2-S478A::LUC mice

Four Cas9 guide RNAs targeting the mouse *Per2* Ser478 codon were selected for activity testing. Guide RNAs were cloned into a T7 promoter vector followed by *in vitro* transcription (HiScribe T7 High Yield RNA Synthesis Kit, New England BioLabs) and spin column purification (RNeasy, Qiagen). Functional testing was performed by incubating Cas9 protein and guide RNA with PCR-amplified target site followed by gel electrophoresis to detect target site cleavage. The guide RNA selected for genome editing in embryos was Per2-sg72T (protospacer sequence 5'- GCGGCTCCAGTGGCTAT-3'). Donor oligonucleotide Per2-S478A-T was used as a donor for point mutation insertion in the *Per2* gene (5'- CTTCTACACCTTTCTATGTGCTTTCCCCCCAGCCTGTCCCCCACAGCGGCT CAGCTGGCTATGGGAGCCTGGGCAGTAACGGATCCCACGAACACCTCATG AGCCAGA -3'). SpCas9 protein was expressed and purified by the UNC Chapel Hill Protein Expression and Purification Core Facility.

Embryos were produced by in vitro fertilization with C57BL/6J females (Jackson Laboratory) as egg donors and $Per2^{luc/luc};CK1\varepsilon^{+/+};CK1\delta^{flox/flox}$ males as sperm donors. Single-cell fertilized embryos were microinjected with 400 nM Cas9 protein, 50 ng/ul guide RNA and 50 ng/ul donor oligonucleotide in microinjection buffer (5 mM Tris-HCl pH 7.5, 0.1 mM EDTA). Injected embryos were implanted in recipient pseudopregnant females and resulting pups were screened by PCR and sequencing for the presence of the $Per2^{S478A}$ allele. Founders harboring the $Per2^{S478A}$ allele were mated to wild-type C57BL/6J females for germline transmission of the targeted allele and to determine if the targeted allele was in cis with the luciferase knock-in. Note that because the $Per2^{luc}$ allele was produced by gene targeting in ES cells from strain 129, the *Per2* locus in the embryos was heterozygous with 129-derived and C57BL/6J-

derived alleles. PCR amplicon sequencing documented polymorphisms in the
vicinity of the Ser478 codon that could be used to distinguish the 129 and
C57BL/6J alleles. Founders with the S478A mutation in the C57BL/6J allele
were used to generate animals with the S478A mutation without the luciferase.
Founders with the S478A allele inserted in cis with luciferase in the 129 allele
were used to establish the combined S478A-Luc strain.
For the establishment of the Ser478Ala mutant lines, biopsy DNA was
genotyped by PCR amplification of a 1,042 bp region encompassing the Ser478
codon with primers Per2-ScF2 (5'- CGGGTCTCTCTGTGCACTCTTG -3') and
Per2-ScR2 (5'- GATGCTTCCTTCTGTCCTCCA -3'). The PCR product was
sequenced with primer Per2-SqR2 (5'- GTGACTTTTGGTTTGAATCTTGC -3').
For genotyping, DNA was extracted from the tail at and analyzed by the
polymerase chain reaction with the primers Per2-ScF2, Per2-ScFwt (5'-
CACAGCGGCTCCAGTGG-3'), Per2-ScRmut (5'-
AGGCTCCCATAGCCAGCTG-3') and Per2-ScR3 (5'-
GCTTCTCAGGGAGAGGAACAG-3').
355

Behavioral experiments
Six- to ten-week-old male mice were individually housed in cages equipped
with running wheels and were entrained to the 12-h /12-h LD cycle. After more
than two weeks, wheel revolutions were recorded under the DD condition for 24
days or longer. Their spontaneous locomotor activities were recorded as the
number of wheel revolutions in 5-min bins and were analyzed with ClockLab
software (Actimetrics). The circadian period of the activity rhythms under the DD
condition was determined via a chi-square periodogram procedure from animals
that showed rhythmicities with $P < 0.001$, based on locomotor activity in days
11–24 after the start of DD condition.
366

Cell culture and real-time monitoring of rhythmic gene expression
MEFs from PER2::LUC mice and PER2-S478A::LUC mice were maintained
at 37°C under 5% CO_2, 95% air in Dulbecco's modified Eagle's medium
(SIGMA) supplemented with 100 units/ml penicillin, 100 µg/ml streptomycin, and
10% fetal bovine serum. We carried out real-time monitoring of luciferase
expression as previously described [53] with minor modifications. Briefly, cells

were treated with 0.1 µM (final) dexamethasone for 30 min, and then the media
were replaced by recording media [phenol-red free Dulbecco's modified Eagle's
medium (SIGMA) supplemented with 10% fetal bovine serum, 3.5 g/l glucose,
25 U/ml penicillin, 25 µg/ml streptomycin, 0.1 mM luciferin, and 10 mM HEPES-
NaOH; pH 7.0]. The bioluminescence signals were continuously recorded for 5-
10 days at 37ºC in air with Dish Type Luminescencer, LumiCycle (Actimetrics).
379

RNA preparation
Tissues from 7–8-week-old male mice were collected every 4 h from 38 h
after the beginning of the DD condition (projected circadian time (CT)). Livers
with three biological replicates were lysed with TRIzol reagent (Invitrogen), and
total RNA was prepared with the RNeasy Mini Kit (Qiagen) according to the
manufacturer's protocol.
386

RT-qPCR
For quantification of gene expression, RNA was reverse-transcribed by Go
Script Reverse Transcriptase (Promega) with both anchored (dT) 15 primers
and random oligo primers. We subjected the cDNA to real-time PCR
(StepOnePlus Real-Time PCR Systems, Applied Biosystems) using GoTaq
Master Mix (Promega) with the gene specific primers (Table 1).
393

Preparation of nuclear and cytosolic fractions of mice liver
The nuclear proteins and cytosolic proteins were isolated as previously
described [54]. Briefly, the mouse tissue (1 g wet weight) was washed with ice-
cold PBS and homogenized on ice with 11 ml of ice-cold buffer A [10 mM
HEPES-NaOH (pH 7.8), 10 mM KCl, 0.1 mM EDTA, 1 mM dithiothreitol (DTT), 1
mM phenylmethylsulfonyl fluoride (PMSF), 4 µg/ml aprotinin, 4 µg/ml leupeptin,
50 mM NaF, and 1 mM Na_3VO_4]. The homogenate was centrifuged (5 min each,
700 xg), and the supernatant was used as "cytosolic fraction". the resultant
precipitate was rinsed and centrifuged (5 min each, 700 xg). The precipitate
was resuspended in 2 ml of ice-cold buffer C [20 mM HEPES-NaOH (pH 7.8),
400 mM NaCl, 1 mM EDTA, 5 mM $MgCl_2$, 2% (v/v) glycerol, 1 mM DTT, 1 mM
PMSF, 4 µg/ml aprotinin, 4 µg/ml leupeptin, 50 mM NaF, and 1 mM Na_3VO_4].

After being gently mixed at 4°C for 1 h, the suspension was centrifuged (30 min
each, 21,600 x g) twice, and the final supernatant was used as "nuclear
fraction".
409

Antibodies and immunoblot analysis
410
Liver nuclear and cytosolic proteins separated by SDS-PAGE were
transferred to polyvinylidene difluoride membrane (Millipore). The blots were
blocked in a blocking solution (1%[w/v] skim milk in TBS [50 mM Tris-HCl, 140
mM NaCl, 1 mM $MgCl_2$ (pH 7.4)]) for 1 h at 37°C and then incubated overnight
at 4°C with a primary antibody in the blocking solution. The signals were
visualized by an enhanced chemiluminescence detection system (PerkinElmer
Life Science). The blot membrane was subjected to densitometric scanning and
the band intensities were quantified using ImageQuant TL (GE Healthcare). The
primary antibodies used in this study are as follows; anti-β-actin (Sigma,
A2228), anti-TBP (Santa Cruz Biotechnology, sc-421), CLSP3 anti-CLOCK [54]
(CLSP3, Medical & Biological Laboratories, D333-3), B1BH2 anti-ARNTL [54]
(B1BH2, Medical & Biological Laboratories, D335-3), anti-PER1 (Medical &
Biological Laboratories, PM091), anti-PER2 (Medical & Biological Laboratories,
PM083), anti-CRY1 (Medical & Biological Laboratories, PM081), anti-CRY2
(Medical & Biological Laboratories, PM082), anti-DBP (Medical & Biological
Laboratories, PM079) and anti-NR1D1 (Medical & Biological Laboratories,
PM092).
428

Degradation assay
429
Degradation assays were performed as previously described [55] with minor
modifications. Plasmids driving expression of CRY1::LUC or CRY2::LUC were
co-transfected with myc-PER2 or myc-PER2-S478A plasmids in HEK 293T
cells. After 48 h, the transfected cells were treated with 100 µg/ml cycloheximide
(Nakalai tesque) before monitoring the bioluminescence in LumiCEC (Churitsu),
or were harvested for immunoblotting. Degradation assay in cultured MEFs was
performed in essentially the same way except that DNA was not transfected.
To calculate half-lives of CRY1::LUC and CRY2::LUC, each bioluminescence
data was fitted by two-term exponential model;

$$f(x) = a_1(1/2)^{(x/\tau_1)} + a_2(1/2)^{(x/\tau_2)} + b,$$

where x represent time, a_i, b and τ_i are constant value such as initial quantities, baseline and half-lives, respectively. These values were computed by linear least square fittings. The fact that the data were fitted well by two-term exponential model indicates CRY1::LUC and CRY2::LUC can exist at least in two states with different stability. The half-life of unstable form (τ_i) was likely to be independent of PER2 abundance, whereas that of stable form (τ_2) was significantly increased in a PER2 dependent manner. The accumulation of CRY proteins in PER2-S478A mice liver can be explained by stabilization of the stable form of CRY proteins, thus, we described τ_2 as the half-lives in Figure S1.

To calculate the instantaneous half-life of PER2, we divide the PER2 decay curve into segments of 1h with a sliding window, which is moved at increments of 0.1 h as done in Zhou et al [17]. Then, each segment of decay curve is fitted to the exponential decay curve.

Reference

1. Reppert SM, Weaver DR (2002) Coordination of circadian timing in mammals. Nature 418:935–941. https://doi.org/10.1038/nature00965
2. Lee C, Etchegaray J-P, Cagampang FRA, et al (2001) Posttranslational Mechanisms Regulate the Mammalian Circadian Clock. Cell 107:855–867. https://doi.org/10.1016/S0092-8674(01)00610-9
3. Takahashi JS (2017) Transcriptional architecture of the mammalian circadian clock. Nat Rev Genet 18:164–179. https://doi.org/10.1038/nrg.2016.150
4. Aryal RP, Kwak PB, Tamayo AG, et al (2017) Macromolecular Assemblies of the Mammalian Circadian Clock. Mol Cell 67:770–782.e6. https://doi.org/10.1016/j.molcel.2017.07.017
5. Hirano A, Fu YH, Ptáek LJ (2016) The intricate dance of post-translational modifications in the rhythm of life. Nat Struct Mol Biol 23:1053–1060. https://doi.org/10.1038/nsmb.3326
6. Gallego M, Virshup DM (2007) Post-translational modifications regulate the ticking of the circadian clock. Nat Rev Mol Cell Biol 8:139–148. https://doi.org/10.1038/nrm2106
7. Ralph M, Menaker M (1988) A mutation of the circadian system in golden hamsters. Science (80-) 241:1225–1227. https://doi.org/10.1126/science.3413487
8. Lowrey PL, Shimomura K, Antoch MP, et al (2000) Positional Syntenic Cloning and Functional Characterization of the Mammalian Circadian Mutation. Science (80-) 288:483–491. https://doi.org/10.1126/science.288.5465.483
9. Kloss B, Price JL, Saez L, et al (1998) The Drosophila clock gene double-time encodes a protein closely related to human casein kinase Iε. Cell 94:97–107. https://doi.org/10.1016/S0092-8674(00)81225-8

10.   Price JL, Blau J, Rothenfluh A, et al (1998) double-time Is a Novel
Drosophila Clock Gene that Regulates PERIOD Protein Accumulation.
Cell 94:83–95. https://doi.org/10.1016/S0092-8674(00)81224-6

11.   Eide EJ, Woolf MF, Kang H, et al (2005) Control of Mammalian Circadian
Rhythm by CKI ε -Regulated Proteasome-Mediated PER2 Degradation
Control of Mammalian Circadian Rhythm by CKI ε -Regulated
Proteasome-Mediated PER2 Degradation. Mol Cell Biol 25:2795–2807.
https://doi.org/10.1128/MCB.25.7.2795

12.   Isojima Y, Nakajima M, Ukai H, et al (2009) CKIε/δ-dependent
phosphorylation is a temperature-insensitive, period-determining process
in the mammalian circadian clock. Proc Natl Acad Sci 106:15744–15749.
https://doi.org/10.1073/pnas.0908733106

13.   Meng Q-J, Maywood ES, Bechtold DA, et al (2010) Entrainment of
disrupted circadian behavior through inhibition of casein kinase 1 (CK1)
enzymes. Proc Natl Acad Sci 107:15240–15245.
https://doi.org/10.1073/pnas.1005101107

14.   Uehara TN, Mizutani Y, Kuwata K, et al (2019) Casein kinase 1 family
regulates PRR5 and TOC1 in the Arabidopsis circadian clock. Proc Natl
Acad Sci 201903357. https://doi.org/10.1073/pnas.1903357116

15.   Shinohara Y, Koyama YM, Ukai-Tadenuma M, et al (2017) Temperature-
Sensitive Substrate and Product Binding Underlie Temperature-
Compensated Phosphorylation in the Clock. Mol Cell 67:783–798.e20.
https://doi.org/10.1016/j.molcel.2017.08.009

16.   Tosini G, Menaker M (1998) The tau mutation affects temperature
compensation of hamster retinal circadian oscillators. Neuroreport
9:1001–5

17.   Zhou M, Kim JK, Eng GWL, et al (2015) A Period2 Phosphoswitch
Regulates and Temperature Compensates Circadian Period. Mol Cell
60:77–88. https://doi.org/10.1016/j.molcel.2015.08.022

18.   Narasimamurthy R, Hunt SR, Lu Y, et al (2018) CK1δ/ε protein kinase
primes the PER2 circadian phosphoswitch. Proc Natl Acad Sci 115:5986–
5991. https://doi.org/10.1073/pnas.1721076115

19.   Lee H, Chen R, Lee Y, et al (2009) Essential roles of CKIdelta and
CKIepsilon in the mammalian circadian clock. Proc Natl Acad Sci U S A
106:21359–64. https://doi.org/10.1073/pnas.0906651106

20.   Xu Y, Padiath QS, Shapiro RE, et al (2005) Functional consequences of a
CKIδ mutation causing familial advanced sleep phase syndrome. Nature
434:640–644. https://doi.org/10.1038/nature03453

21.   Brennan KC, Bates EA, Shapiro RE, et al (2013) Casein kinase Iδ
mutations in familial migraine and advanced sleep phase. Sci Transl Med
5:. https://doi.org/10.1126/scitranslmed.3005784

22.   Keesler GA, Camacho F, Guo Y, et al (2000) Phosphorylation and
destabilization of human period I clock protein by human casein kinase Iε.
Neuroreport 11:951–955. https://doi.org/10.1097/00001756-200004070-
00011

23.   Maywood ES, Chesham JE, Smyllie NJ, Hastings MH (2014) The Tau
Mutation of Casein Kinase ε Sets the Period of the Mammalian
Pacemaker via Regulation of Period1 or Period2 Clock Proteins. J Biol
Rhythms 29:110–118. https://doi.org/10.1177/0748730414520663

24.   Vanselow K, Vanselow JT, Westermark PO, et al (2006) Differential
effects of PER2 phosphorylation: Molecular basis for the human familial
advanced sleep phase syndrome (FASPS). Genes Dev 20:2660–2672.
https://doi.org/10.1101/gad.397006

25.   Xu Y, Toh KL, Jones CR, et al (2007) Modeling of a Human Circadian

Mutation Yields Insights into Clock Regulation by PER2. Cell 128:59–70. https://doi.org/10.1016/j.cell.2006.11.043

26. Shanware NP, Hutchinson JA, Kim SH, et al (2011) Casein kinase 1-dependent phosphorylation of familial advanced sleep phase syndrome-associated residues controls PERIOD 2 stability. J Biol Chem 286:12766–12774. https://doi.org/10.1074/jbc.M111.224014

27. Fustin J, Kojima R, Itoh K, et al (2018) Two Ck1δ transcripts regulated by m6A methylation code for two antagonistic kinases in the control of the circadian clock. Proc Natl Acad Sci 115:5980–5985. https://doi.org/10.1073/pnas.1721371115

28. Ohsaki K, Oishi K, Kozono Y, et al (2008) The role of β-TrCP1 and β-TrCP2 in circadian rhythm generation by mediating degradation of clock protein PER2. J Biochem 144:609–618. https://doi.org/10.1093/jb/mvn112

29. Toh KL, Jones CR, He Y, et al (2001) An hPer2 phosphorylation site mutation in familiar advanced sleep phase syndrome. Science (80-) 291:1040–1043. https://doi.org/10.1126/science.1057499

30. Liu J, Zou X, Gotoh T, et al (2018) Distinct control of PERIOD2 degradation and circadian rhythms by the oncoprotein MDM2. 715:1–18. https://doi.org/10.1126/scisignal.aau0715

31. Chen R, Schirmer A, Lee Y, et al (2009) Rhythmic PER Abundance Defines a Critical Nodal Point for Negative Feedback within the Circadian Clock Mechanism. Mol Cell 36:417–430. https://doi.org/10.1016/j.molcel.2009.10.012

32. Nangle SN, Rosensweig C, Koike N, et al (2014) Molecular assembly of the period-cryptochrome circadian transcriptional repressor complex. Elife 3:1–14. https://doi.org/10.7554/eLife.03674

33. Xing W, Busino L, Hinds TR, et al (2013) SCF FBXL3 ubiquitin ligase targets cryptochromes at their cofactor pocket. Nature 496:64–68. https://doi.org/10.1038/nature11964

34. Ye R, Selby CP, Chiou Y, et al (2014) Dual modes of CLOCK:BMAL1 inhibition mediated by Cryptochrome and Period proteins in the mammalian circadian clock. Genes Dev 28:1989–1998. https://doi.org/10.1101/gad.249417.114

35. Koike N, Yoo S-H, Huang H-C, et al (2012) Transcriptional Architecture and Chromatin Landscape of the Core Circadian Clock in Mammals. Science (80-) 338:349–354. https://doi.org/10.1126/science.1226339

36. Yoo SH, Yamazaki S, Lowrey PL, et al (2004) PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. Proc Natl Acad Sci U S A 101:5339–5346. https://doi.org/10.1073/pnas.0308709101

37. Hastings JW, Sweeney BM (1957) on the Mechanism of Temperature Independence in a Biological Clock. Proc Natl Acad Sci 43:804–811. https://doi.org/10.1073/pnas.43.9.804

38. Vielhaber E, Eide E, Rivers A, et al (2000) Nuclear Entry of the Circadian Regulator mPER1 Is Controlled by Mammalian Casein Kinase I varepsilon. Mol Cell Biol 20:4888–4899. https://doi.org/10.1128/mcb.20.13.4888-4899.2000

39. Hirota T, Lee JW, Lewis WG, et al (2010) High-throughput chemical screen identifies a novel potent modulator of cellular circadian rhythms and reveals CKIα as a clock regulatory kinase. PLoS Biol 8:. https://doi.org/10.1371/journal.pbio.1000559

40. Dahlberg CL, Nguyen EZ, Goodlett D, Kimelman D (2009) Interactions between casein kinase Iε (CKIε) and two substrates from disparate signaling pathways reveal mechanisms for substrate-kinase specificity.

PLoS One 4:. https://doi.org/10.1371/journal.pone.0004766

41. Lam VH, Li YH, Liu X, et al (2018) CK1α collaborates with DOUBLETIME
to regulate PERIOD function in the Drosophila circadian clock. J Neurosci
38:10631–10643. https://doi.org/10.1523/JNEUROSCI.0871-18.2018

42. Tsuchiya Y, Akashi M, Matsuda M, et al (2009) Involvement of the Protein
Kinase CK2 in the Regulation of Mammalian Circadian Rhythms. Sci
Signal 2:ra26-ra26. https://doi.org/10.1126/scisignal.2000305

43. Jakubcakova V, Oster H, Tamanini F, et al (2007) Light Entrainment of
the Mammalian Circadian Clock by a PRKCA-Dependent
Posttranslational Mechanism. Neuron 54:831–843.
https://doi.org/10.1016/j.neuron.2007.04.031

44. Mehta N, Cheng AH, Chiang CK, et al (2015) GRK2 Fine-Tunes
Circadian Clock Speed and Entrainment via Transcriptional and Post-
translational Control of PERIOD Proteins. Cell Rep 12:1272–1288.
https://doi.org/10.1016/j.celrep.2015.07.037

45. He Q, Cha J, He Q, et al (2006) CKI and CKII mediate the FREQUENCY-
dependent phosphorylation of the WHITE COLLAR complex to close the
Neurospora circadian negative feedback loop. Genes Dev 20:2552–2565.
https://doi.org/10.1101/gad.1463506

46. Querfurth C, Diernfellner ACR, Gin E, et al (2011) Circadian
Conformational Change of the Neurospora Clock Protein FREQUENCY
Triggered by Clustered Hyperphosphorylation of a Basic Domain. Mol Cell
43:713–722. https://doi.org/10.1016/j.molcel.2011.06.033

47. Larrondo LF, Olivares-Yañez C, Baker CL, et al (2015) Decoupling
circadian clock protein turnover from circadian period determination.
Science (80-) 347:. https://doi.org/10.1126/science.1257277

48. Liu X, Chen A, Caicedo-Casso A, et al (2019) FRQ-CK1 interaction
determines the period of circadian rhythms in Neurospora. Nat Commun
10:4352. https://doi.org/10.1038/s41467-019-12239-w

49. Ruoff P (1992) Introducing temperature‐compensation in any reaction
kinetic oscillator model. J Interdisiplinary Cycle Res 23:92–99.
https://doi.org/10.1080/09291019209360133

50. Shearman LP, Sriram S, Weaver DR, et al (2000) Interacting molecular
loops in the mammalian circadian clock. Science (80-) 288:1013–1019.
https://doi.org/10.1126/science.288.5468.1013

51. Ye R, Selby CP, Ozturk N, et al (2011) Biochemical analysis of the
canonical model for the mammalian circadian clock. J Biol Chem
286:25891–25902. https://doi.org/10.1074/jbc.M111.254680

52. Shirogane T, Jin J, Ang XL, Harper JW (2005) SCFβ-TRCP controls
Clock-dependent transcription via casein kinase 1-dependent degradation
of the mammalian period-1 (Per1) protein. J Biol Chem 280:26863–
26872. https://doi.org/10.1074/jbc.M502862200

53. Imamura K, Yoshitane H, Hattori K, et al (2018) ASK family kinases
mediate cellular stress and redox signaling to circadian clock. Proc Natl
Acad Sci 201719298. https://doi.org/10.1073/pnas.1719298115

54. Yoshitane H, Takao T, Satomi Y, et al (2009) Roles of CLOCK
Phosphorylation in Suppression of E-Box-Dependent Transcription. Mol
Cell Biol 29:3675–3686. https://doi.org/10.1128/MCB.01864-08

55. Hirano A, Yumimoto K, Tsunematsu R, et al (2013) FBXL21 regulates
oscillation of the circadian clock through ubiquitination and stabilization of
cryptochromes. Cell 152:1106–1118.
https://doi.org/10.1016/j.cell.2013.01.054

Figure legends

Figure 1. The PER2 Ser478Ala mutation lengthens circadian rhythms
(A) The domain structure of PER2 protein is depicted (upper). The genome
sequences in wild-type (WT) and PER2-S478A (SA) knock-in mouse and
corresponding amino acid sequences are shown (lower). PAS, Per-Arnt-Sim
domain; CKBD, CK1-binding domain; CBD, CRY-binding domain.
(B) Wheel-running activities of representative PER2-S478A homozygous mice
and littermate wild-type mice are shown. Mice were entrained to LD for two
weeks or longer and then exposed to DD.
(C) The circadian period of the activity rhythms under the DD condition was
determined via a chi-square periodogram procedure based on locomotor activity
in days 11–24 after the start of DD condition. Mean values ± SEM obtained from
ten wild-type (WT) mice and twelve homozygous PER2-S478A (SA) mice are
given. *** $p = 4.2$ x E-8, two-sided Student's *t*-test.

**Figure 2. PER2 expression levels are post-transcriptionally upregulated in
PER2-S478A mouse liver**
(A) Temporal expression of *Per2* mRNA normalized by *Rps29*. Livers were
harvested at 4-h intervals followed by real-time RT-qPCR. Mean values ± SEM
obtained from three animals of each genotype are given.
(B, C) Temporal expression of PER2 proteins in liver. Liver nuclear extracts (B)
and cytoplasmic lysates (C) were prepared at 4-h intervals following SDS-PAGE
and immunoblotting. Data are represented as dots for individual experiments
and as lines for means.

**Figure 3. The PER2-S478A mutation alters the expression profiles of clock
genes and proteins in the liver**
(A, B) Temporal expression of clock proteins in liver cytosol. Liver cytoplasmic
lysates (A) and nuclear extracts (B) were prepared at 4-h intervals and analyzed
by SDS-PAGE and immunoblotting with indicated antibodies. Data are
represented as dots for individual experiments and as lines for means.

674 (C) Temporal expression of clock genes. Livers were harvested at 4-h intervals
675 followed by real-time RT-qPCR for the indicated mRNAs. Mean values ± SEM
676 obtained from three animals of each genotype are given.
677

Figure 4. The mutation of PER2 at Ser478 perturbs the phosphoswitch
(A) Wheel-running activities of representative PER2::LUC and PER2-
S478A::LUC homozygous mice are shown. Mice were entrained to LD for two
weeks or longer and then exposed to DD.
(B) The circadian period of the activity rhythms under the DD condition was
determined via a chi-square periodogram procedure based on locomotor activity
in days 11–24 after the start of DD condition. Mean values ± SEM obtained from
ten PER2::LUC mice (WT) and ten PER2-S478A::LUC (SA) mice are given. ***
p = 5.6 x E-7, two-sided Student's t-test.
(C) Representative recordings of bioluminescence from PER2::LUC or PER2-
S478A::LUC MEFs are shown. MEFs were synchronized with dexamethasone
and the bioluminescence was then continually measured in the Lumicycle.
(D) The circadian period of cellular rhythms was calculated. Mean values ±
SEM obtained from three PER2::LUC MEFs (WT) and three PER2-S478A::LUC
(SA) MEFs are given. *** p = 0.001, two-sided Student's t-test. n.s. not
significant (p = 0.17, two-sided Student's t-test).
(E) Bioluminescence from PER2::LUC (WT) and PER2-S478A::LUC (SA) MEFs
are shown. MEFs were synchronized with dexamethasone and the
bioluminescence was continually measured in the Lumicycle. CHX was added
at CT 19.8 (left) or at CT 21.4 (right). By fitting the exponential curve to the
small segment of the decay curves, the instantaneous half-life is calculated (see
Materials and methods for details).

Figure 5. PER2-S478A mutation lengthens the circadian period
PER2 and CRY proteins accumulate during night and disappear before CT2 in
wild-type. In contrast, PER2 and CRY proteins are stabilized and excessively
accumulate around CT 22 in PER2-S478A mutants. Thereafter, PER2 proteins
are degraded by CT2 whereas CRY proteins keep repressing the CLOCK-
BMAL complexes.
706

Supplemental Figure S1. PER2 stabilizes CRY1::LUC and CRY2::LUC

(A, B) Half-lives of the bioluminescence derived from CRY1::LUC (A, upper panel) or CRY2::LUC (B, upper panel) after CHX treatment. The detailed method for calculation of half-lives is described in Materials and methods. Myc-PER2 (WT) or Myc-PER2-S478A (SA) were co-transfected with CRY1::LUC or CRY2::LUC. The abundance of PER2 proteins were analyzed by immunoblotting with anti-PER2 antibody (A, B, lower panel). Mean values ± SEM obtained from three biological replicates are given.

(C) Representative western blot for PER2, corresponding to (A, lower panel).

(D) Representative western blot for PER2, corresponding to (B, lower panel).

A

B Wild-type   PER2-S478A   C

A

B   CT 2   6   10 14 18 22

C   CT 2   6   10 14 18 22

Table 1. Primer sequences for RT-qPCR

Primer	Sequence (5'-3')
mRps29-Fw	TGAAGGCAAGATGGGTCAC
mRps29-Rv	GCACATGTTCAGCCCGTATT
mPer1-Fw	CAGGCTAACCAGGAATATTACCAGC
mPer1-Rv	CACAGCCACAGAGAAGGTGTCCTGG
mPer2-Fw	GGCTTCACCATGCCTGTTGT
mPer2-Rv	GGAGTTATTTCGGAGGCAAGTGT
mCry1-Fw	CCCAGGCTTTTCAAGGAATGGAACA
mCry1-Rv	TCTCATCATGGTCATCAGACAGAGG
mCry2-Fw	GGGACTCTGTCTATTGGCATCTG
mCry2-Rv	GTCACTCTAGCCCGCTTGGT
mClock-Fw	CCTATCCTACCTTGGCCACACA
mClock-Rv	TCCCGTGGAGCAACCTAGAT
mArntl-Fw	GCAGTGCCACTGACTACCAAGA
mArntl-Rv	TCCTGGACATTGCATTGCAT
mDbp-Fw	AATGACCTTTGAACCTGATCCCGCT
mDbp-Rv	GCTCCAGTACTTCTCATCCTTCTGT
mNr1d1-Fw	CGTTCGCATCAATCGCAACC
mNr1d1-Rv	GATGTGGAGTAGGTGAGGTC
mNr1d2-Fw	ACGGATTCCCAGGAACATGG
mNr1d2-Rv	CCTCCAGTGTTGCACAGGTA