L-type voltage-gated calcium channel agonists improve hearing loss and modify ribbon synapse

morphology in the zebrafish model of Usher Syndrome Type 1

Alaa Koleilat[1,2,3], Joseph A. Dugdale[4], Trace A. Christenson[5], Jeffrey L. Bellah[3,6], Aaron M.

Lambert[7,8], Mark A. Masino[7], Stephen C. Ekker[3,9], Lisa A. Schimmenti[4,9,10,11,12,13]

[1]College of Continuing and Professional Studies, University of Minnesota, Minneapolis, MN,

USA;

[2]Mayo Clinic Graduate School of Biomedical Sciences, Clinical and Translational Science Track,

Rochester, MN, USA (current position);

[3]Center for Clinical and Translational Science, Mayo Clinic, Rochester, MN, USA

[4]Department of Ophthalmology and Visual Neuroscience, University of Minnesota, Minneapolis,

MN, USA

[5]Mayo Clinic Microscopy and Cell Analysis Core, Rochester, MN, USA

[6]Department of Genetics and Development, Columbia University, New York City, NY, USA

[7]Department of Neuroscience, University of Minnesota, Minneapolis, MN, USA

[8]Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA, USA

[9]Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, MN, USA

[10]Department of Pediatrics, University of Minnesota, Minneapolis, MN, USA

[11]Department of Genetics, Cell Biology and Development, University of Minnesota, Minneapolis,

MN, USA

[12]Department of Otorhinolaryngology, Mayo Clinic, Rochester, MN, USA (current position)

[13]Department of Clinical Genomics, Mayo Clinic, Rochester, MN, USA (current position)

Key words: zebrafish, hair cell, ribbon synapse, myo7aa, hearing loss

Summary Statement

We identified that the hair cell biology and behavior of the *myo7aa*[-/-] mutant differs from wildtype

and this difference can be rescued using L-type voltage-gated calcium channel agonists.

Abstract

Usher Syndrome (USH) is the most common cause of human deaf/blindness. The zebrafish *myo7aa*[-/-] mutant, faithfully models USH1; homozygous zebrafish are deaf and exhibit circular swimming. We hypothesized that hair cell morphology would differ in *myo7aa*[-/-] mutants compared to wild type. We also tested the hypothesis that agonists of L-type voltage-gated calcium channels would alter ribbon synapse morphology and behavior of zebrafish *myo7aa*[-/-] mutants. We discovered that *myo7aa*[-/-] zebrafish have fewer glutamatergic vesicles tethered to hair cell ribbon synapses, yet maintain a comparable ribbon area. We identified that *myo7aa*[-/-] mutants have fewer total active hair cells, fewer total CTBP2 expressing puncta, and an altered distribution of CTBP2 puncta compared to wildtype. We also identified that *myo7aa*[-/-] mutants have fewer active post-synaptic cells and fewer total MAGUK puncta, compared to wildtype. Behaviorally, *myo7aa*[-/-] mutant fish have abnormal swimming as measured by larger absolute smooth orientations and have little to no acoustic startle. Treatment with L-type voltage-gated calcium channel agonists altered the abnormal cell and behavioral phenotypes toward wildtype. Our data supports that L-type voltage-gated calcium channel agonists induce morphological changes at the ribbon synapse—in both the number of tethered vesicles and the distribution of CTBP2 puncta, shifts swimming behavior towards wildtype swimming and improves acoustic startle response.

INTRODUCTION

A reduction or the complete absence of the ability to detect sounds is the most common impairment of the human senses (Ahmed et al., 2013). About 20 percent of Americans report some degree of hearing loss, making this a major public health concern (Communication) (Prevention, 2016, Prevention, April 1996). Hearing loss is caused by genetic or environmental/infectious factors, with more than fifty percent of hearing loss is attributed to genetics (Morton and Nance, 2006).

Up to 11% of deaf/hard of hearing children and 1 in 6000 people in the United States have Usher Syndrome (USH) (Kimberling et al., 2010). USH is characterized by congenital onset of sensorineural deafness/hard of hearing with later onset of retinitis pigmentosa resulting in significant vision impairment (Usher, 1914). There are three types of USH characterized by degree of hearing, vestibular abnormalities and onset of vision loss (Davenport SLH, 1977).

Usher Syndrome Type I (USH1) is the most severe form of Usher Syndrome with profound bilateral congenital hearing loss and onset of retinitis pigmentosa within the first decade of life. Ten unique loci have been associated with USH1 with variants in *MYO7A* as the most common cause, accounting for 53-63% of affected individuals (Lentz J, 1999). Additionally, pathogenic variants in CDH23, PCDH15, USH1C and USH1G are responsible for 19-35%, 11-19%, 6-7% and 7% respectively (hereditaryhearingloss.org). Each gene encodes structural and motor proteins important for mechanotransduction in the inner ear hair cells (Beurg et al., 2009, Grati and Kachar, 2011, Grillet et al., 2009a, Kazmierczak et al., 2007, Marcotti, 2012, Pepermans and Petit, 2015, Siemens et al., 2004).

Gibson et al. in 1995 identified the first USH locus in the *shaker-1* (*sh1*) mouse. The *shaker-1* mouse presented with hearing loss, head tossing and circling behaviors due to vestibular dysfunction, and upon examination of inner ear hair cells was found to have disorganized stereocilia. Through positional cloning techniques homozygous mutations at the *sh1* locus were identified in *Myo7a*. Two years later, Weil et al. identified that USH1B in humans was also due to pathogenic variants in *MYO7A* (Weil et al., 1997). Ernest et al. in 2000 described a zebrafish model of USH1B caused by a premature stop codon in *myo7aa*, the *mariner* mutant, in which the phenotype of the homozygous recessive larval fish consisted of a circular swimming pattern, defective balance, morphological and functional defects in the inner ear hair cells and most notably the lack of a startle response (Ernest et al., 2000).

MYO7A encodes an unconventional actin-binding motor protein, which is important for development and function of the inner ear hair cells. It is specifically involved in upholding the structural integrity of the hair bundle allowing for a mechanical stimulus to be converted into a

chemical stimulus. The myosin VIIa protein is localized at the upper tip link density of stereocilia

in sensory hair cells (Hasson et al., 1995). Myosin VIIa along with Harmonin and Sans interact

with one another to connect the upper end of the tip link (composed of cadherin 23 and

protocadherin 15) to the actin cytoskeleton of the stereocilium (Ahmed et al., 2006, Caberlotto et

al., 2011, Grati and Kachar, 2011, Grillet et al., 2009b, Siemens et al., 2004). Myosin VIIa is

involved in maintaining the tension of the tiplink structure upon positive hair cell deflection.

When sound is administered the stereocilia of hair cells are deflected towards the tallest

stereocilia allowing for the mechanotransduction channel (MET) located at the apical region of

the stereocilia to open (Fig. 1A). The opening of the MET channel causes positively charged

cations such as potassium and calcium to flow into the cell and effect depolarization.

Once the proper membrane potential is reached, L-type voltage-gated calcium channels

(Ca_v1.3)-localized to the active zones in hair cells adjacent to the ribbon synapse-open, further

increasing intracellular calcium concentrations (Brandt et al., 2005, Moser and Vogl, 2016, Sidi

et al., 2004). Although calcium has many roles in sensory hair cells, entry of calcium through

Ca_v1.3 is necessary to mediate the release of synaptic vesicles tethered to the ribbon synapse

structure (Fuchs, 2005, Moser et al., 2006, Nicolson, 2015, Wong et al., 2014). The ribbon

synapse is composed primarily of a central protein, Ribeye encoded by CTBP2, and surrounded

by a halo of glutamatergic vesicles (Schmitz et al., 2000). When glutamine is released into the

synaptic cleft, it binds onto post-synaptic cell receptors to stimulate cochlear afferents, thus

transducing the sound signal (Fig.1A).

In the absence of myo7a expression, the MET channel is unable to gate properly in

response to stereocilia deflections, therefore cell depolarization is not reached and L-type

voltage-gated calcium channels do not open, limiting synaptic transmission (Fig. 1B). In this

study, we investigated the hypothesis that increasing intracellular calcium in the inner ear hair

cell of a *myo7aa*[-/-] mutant would restore hearing and swimming abnormalities as well as change

hair cell biology. In order to address this hypothesis, we treated *myo7aa*[-/-] mutants with L-type

voltage-gated calcium channel agonists. This class of drugs was previously used in a variety of

studies to treat other neurologic conditions, but not hearing loss (Hiramatsu et al., 1997, Kitano

et al., 1994, Starkstein et al., 2016).

Here we report that treatment of *myo7aa*[-/-] zebrafish with three separate drugs with L-

type voltage-gated calcium channel agonistic activity, specifically (±)-Bay K 8644, Nefiracetam

and (R)-Baclofen, can restore hair cell biology and behavior abnormalities observed in untreated

mutant zebrafish toward wildtype.

RESULTS

Hair cell ribbon synapse ultrastructure, CTPB2 *expression,* MAGUK *expression, and behavior of the myo7aa$^{-/-}$ mutant is different from wildtype*

The *myo7aa$^{-/-}$* mutant was first described in 2000 to have absent acoustic startle, balance abnormalities, a deflated swim bladder, and disorganized stereocilia (Ernest et al., 2000); Fig. 2A-D) It was previously reported that the mechanotransduction channel did not uptake the vital dye, FM 1-43, and is therefore inactive in this mutant model (Seiler and Nicolson, 1999); Fig. 2A, B). To elucidate the ultrastructural state of the ribbon synapse structure in hair cells we used transmission electron microscopy (TEM) (Fig. 2E, F). The mean ± standard error of the mean of wildtype ribbon area is 37866.7 ± 3166.2 nm^2 (n=35 ribbons; 6 zebrafish) and the mean of the *myo7aa$^{-/-}$* mutants is 30992.6 ± 3347.2 nm^2 (n=35 ribbons; 5 zebrafish); therefore the ribbon area between *myo7aa$^{-/-}$* mutants and wildtype larvae is comparable (Fig. 2G, t-test, p=0.14). The mean number of tethered vesicles for wildtype is 19 ± 1 vesicles (n=32 ribbons, 6 zebrafish) which is statistically different from the mean for the *myo7aa$^{-/-}$* mutants 15 ± 1 vesicles (n=34 ribbons, 5 zebrafish) (Fig. 2H, t-test, p<0.01).

We further explored the ribbon synapse in the *myo7aa$^{-/-}$* mutants and performed immunohistochemistry to assess differences in pre- and post-synaptic markers of the hair cells. The ribeye protein is the result of alternative splicing of the transcription factor CTBP2. Ribeye has two main roles: clustering of the Ca$_v$1.3 channels at the presynapse and stabilizing contacts with the afferent neurons (Fig. 3A) (Lv et al., 2016, Sheets et al., 2011). CTBP2 puncta can be observed in hair cells and we define an active hair cell as any hair cell with at least one CTBP2 puncta. We assessed CTBP2 expression in zebrafish lateral line hair cells and quantified the number of active hair cells, total puncta and the distribution of puncta. The wildtype neuromasts had a mean of 19 ± 1 active hair cells (n=22 neuromasts, n=18 zebrafish) and a mean of 62 ± 4 total CTBP2 puncta per neuromast (n=22 neuromasts, n=17 zebrafish). These two parameters are statistically significant compared to the *myo7aa$^{-/-}$* mutants with a mean of 12 ± 1 active hair cells (n=18 neuromasts, n=17 zebrafish) and a mean of 34 ± 3 total CTBP2 puncta per neuromast (n=18 neuromasts, n=17 zebrafish) (t-test, p<0.0001, p<0.0001) indicating that there are fewer active hair cells and fewer CTBP2 puncta per neuromast in the *myo7aa$^{-/-}$* mutants (Fig. 3D, E). We also investigated the distribution of CTBP2 puncta and identified that most hair cells from the *myo7aa$^{-/-}$* mutant neuromasts had two puncta per cell whereas most wildtype hair cells had three puncta per cell (Fig. 3F). Additionally, the proportion of hair cells with two CTBP2 puncta and three CTBP2 puncta are statistically significant between wildtype and *myo7aa$^{-/-}$*

mutants accompanied by no differences in the proportion of 1, 4, 5 and 6+ CTBP2 puncta

between the two groups (Table S2).

We also explored any differences in the post-synaptic hair cell marker, MAGUK.

Membrane associated guanylate kinase, MAGUK, was used to assess afferent post-synaptic

structures in the afferent fiber terminal (Anderson 1996; Lv et al. 2016; Sheets et al. 2011). This

kinase family targets and anchors glutamate receptors to the synaptic terminal on the post-

synaptic cell as part of the postsynaptic density (PSD) (Fig. 4A). Wildtype neuromasts had a

mean of 22 ± 1 active post-synaptic cells (n=29 neuromasts, n=29 zebrafish) and a mean of 59

± 3 total MAGUK puncta per neuromast (n=29 neuromasts, n=29 zebrafish). These two

parameters are statistically significant compared to the *myo7aa$^{-/-}$* mutants which have a mean of

16 ± 1 active post-synaptic cells (n=23 neuromasts, n=23 zebrafish) and a mean of 46 ± 2 total

MAGUK puncta per neuromast (n=23 neuromasts, n=23 zebrafish) (t-test, p<0.0001, p=0.001).

This indicates that there are fewer active post-synaptic cells and fewer MAGUK puncta per

neuromast in the *myo7aa$^{-/-}$* mutants (Fig. 4D, E). We also investigated the distribution of

MAGUK puncta and identified that most post-synaptic cells from the *myo7aa$^{-/-}$* mutant

neuromasts had 3 puncta per cell whereas most wildtype hair cells had either 2 or 3 puncta per

cell; however, this difference is not statistically significant (Fig. 4F, Table S2)

The abnormal circular and loop-like swimming behavior of the *myo7aa$^{-/-}$* mutant is

apparent by observation using a tactile stimulus to the tail or through spontaneous swimming.

We were able to quantify this abnormal swimming behavior by using a previously described

behavior assessment (Branson et al. 2009). We quantified the absolute smooth orientation or

global change in body orientation as a function of time. The absolute smooth orientation of

wildtype fish was 295 ± 5° (n=46 zebrafish) compared to the *myo7aa$^{-/-}$* mutant larvae which had

a mean absolute smooth orientation of 486 ± 5° (n=61 zebrafish)(t-test; p<0.0001) (Fig. 5).

L-type voltage gated calcium channel agonists alter myo7aa$^{-/-}$ ribbon synapse

ultrastructure

Through transmission electron microscopy we established that ribbon synapse

structures differ between 5 days post fertilization (dpf) wildtype and *myo7aa$^{-/-}$* mutant fish,

specifically the number of vesicles tethered to the electron-dense structure, ribeye. To explore

whether L-type voltage-gated calcium channel agonists can modify hair cell ribbon synapse

components we incubated 4 dpf *myo7aa$^{-/-}$* mutant embryos overnight in either 5 µM (±)-Bay K

8644, 250 µM Nefiracetam or 125 µM (R)-Baclofen and performed transmission electron

microscopy on 5 dpf. We observed that the mean area of the electron dense structure, ribeye,

for *myo7aa*$^{-/-}$ mutant fish incubated with either 5 µM (±)-Bay K 8644 or 250 µM Nefiracetam was

27274.97 ± 2231.61 nm^2 (n=35 ribbons, n=5 zebrafish)(t-test, p=0.36) and 24024.14 ± 2335.09

nm^2 (n=29 ribbons, n=4 zebrafish)(t-test, p=0.09), respectively, and thus did not alter the area of

the ribbon synapse. Additionally, neither drug altered the number of vesicles tethered to ribeye

with a mean of 17 ± 1 (n=35 ribbons, n=5 zebrafish) (t-test, p=0.07) and a mean of 14 ± 1 (n=29

ribbons, n=4 zebrafish) (t-test, p=0.54), respectively. However, treatment with 125 µM (R)-

Baclofen did modify both parameters with a mean area of 44071.6 ± 3008 nm^2 (n=50 ribbons,

n=5 zebrafish) (t-test, p=0.02) and 17 ± 1 tethered vesicles (n=46 ribbons, n=5 zebrafish) (t-test,

p=0.03) (Fig. 6). None of the three drugs showed any adverse effect on wildtype ribbon synapse

ultrastructure compared to untreated wildtype (Fig. S1).

L-type voltage-gated calcium channel agonists alter CTBP2 and MAGUK expression

We previously identified that the *myo7aa*$^{-/-}$ mutant has different CTBP2 and MAGUK gene

expression compared to wildtype. We discovered that *myo7aa*$^{-/-}$ mutants have fewer active hair

cells and post-synaptic cells, fewer total CTBP2 and MAGUK puncta and a different distribution

of CTBP2 puncta compared to wildtype. Here we explored the effect of L-type voltage-gated

calcium channel agonists on CTBP2 and MAGUK expression. We identified that wildtype larvae

treated with either 250 µM Nefiracetam or 125 µM (R)-Baclofen did not alter the overall

distribution of CTBP2 puncta, however incubation with 5 µM (±)-Bay K 8644 did shift the curve

so that most hair cells had 2 puncta per cell compared to 3 puncta per cell in the untreated

groups (Fig. 7 A-D, I; Table S3). There was generally no change in the distribution of MAGUK

puncta with incubation of either 5 µM (±)-Bay K 8644, 250 µM Nefiracetam or 125 µM (R)-

Baclofen (Fig. 7E-H, J; Table S4). Additionally, there was an increase in the total number of

active hair cells per neuromast and the total number of CTBP2 puncta per neuromast for

wildtype fish incubated in 5 µM (±)-Bay K 8644, but no change in both parameters upon

incubation with either 250 µM Nefiracetam or 125 µM (R)-Baclofen (Fig. S2A). There was,

however, an increase in total number of active post-synaptic cells per neuromast and the total

number of MAGUK puncta per neuromast for wildtype fish incubated in 5 µM (±)-Bay K 8644 or

125 µM (R)-Baclofen, but no change with 250 µM Nefiracetam (Fig. S3A). We also identified

that *myo7aa*$^{-/-}$ mutant larvae treated with either 5 µM (±)-Bay K 8644, 250 µM Nefiracetam or

125 µM (R)-Baclofen did alter the distribution of CTBP2 puncta so that most hair cells had 3

puncta per cell (Fig. 7K-N, S; Table S3). There was also generally no statistically significant

change in the distribution of MAGUK puncta with incubation of either 5 µM (±)-Bay K 8644, 250

µM Nefiracetam or 125 µM (R)-Baclofen, (Fig. 7O-R, T; Table S4). Moreover, there was an

increase in the total number of active hair cells per neuromast, total number of CTBP2 puncta

per neuromast, total number of post-synaptic cells per neuromast and total number of MAGUK

puncta per neuromast for *myo7aa*$^{-/-}$ fish incubated in 5 µM (±)-Bay K 8644, but no change in any

of the parameters upon incubation with either 250 µM Nefiracetam or 125 µM (R)-Baclofen (Fig.

S2B, Fig. S3B).

L-type voltage-gated calcium channel agonists restore myo7aa*$^{-/-}$ *abnormal swimming

245   To determine the effect of L-type voltage-gated calcium channel agonists on swimming

behavior in the *myo7aa*$^{-/-}$ mutants, we used a behavior assessment that quantifies turning

angles as a function of time (absolute smooth orientation). Wildtype fish incubated in 5 µM (±)-

Bay K 8644 had smaller absolute smooth orientations with a mean of 218 ± 6° (n=15 zebrafish)

compared to untreated wildtype larvae with a mean of 295 ± 5° (n=46 zebrafish), however

swimming traces were similar (t-test; p<0.0001) (Fig. 8A, B, C). Incubation with 250 µM

Nefiracetam or 125 µM (R)-Baclofen had similar absolute smooth orientations compared to

untreated wildtype fish 309 ± 10° (n=17 zebrafish) and 313 ± 13° (n=11) and swimming

trajectories, respectively (t-test, p=0.22, p=0.23) (Fig. 8A, B, C). *Myo7aa*$^{-/-}$ mutant fish incubated

in 5 µM (±)-Bay K 8644, 250 µM Nefiracetam or 125 µM (R)-Baclofen showed improved

swimming behavior with decreased absolute smooth orientation 252 ± 16° (n=11), 450 ± 13°

(n=22) , 451 ± 11° (n=20) compared to untreated controls, respectively (t-test, p<0.0001,

p=0.03, p=0.01) (Fig.8A, C). *Myo7aa*$^{-/-}$ fish incubated with any of the three drugs displayed

smoother swimming trajectories with less circular swimming and loop-like swimming and more

zigzag swimming (Fig. 8B). Dose response and time response experiments were performed for

Nefiracetam and (R)-Baclofen on wildtype and *myo7aa*$^{-/-}$ mutants (Fig. S4-7). Three doses of

Nefiracetam were tested, 85 µM, 125 µM and 250 µM. Wildtype absolute smooth orientations

did not differ from untreated fish except at 125 µM and *myo7aa*$^{-/-}$ mutant absolute smooth

orientations did not differ from untreated fish except at 250 µM (Fig. S4). We also tested if 1

hour incubation would suffice to see any effect and we concluded that there was no difference in

either group after 1 hour incubation (Fig. S5). Five doses of (R)-Baclofen were tested, 75 µM,

125 µM, 250 µM, 500 µM and 1 mM. Wildtype absolute smooth orientations did not differ from

untreated fish except at 75 µM, 250 µM, 500 µM and *myo7aa*$^{-/-}$ mutant absolute smooth

orientations did differ from untreated fish at all doses (Fig. S6). We also tested a shorter

incubation time at the 125uM dose and identified that a 1 hour incubation of (R)-Baclofen is

sufficient to see an effect in the *myo7aa*$^{-/-}$ mutants (Fig. S7). There was no difference between

wildtype untreated and treated with 125 µM (R)-Baclofen after the 1 hour incubation (Fig. S7).

Lastly, we assessed if there was a change in swim bladder inflation upon incubation in
the *myo7aa*[-/-] mutants with 250 µM Nefiracetam and 125 µM (R)-Baclofen and did not see any
change (Fig. S8).
275
L-type voltage-gated calcium channel agonists increase *myo7aa*[-/-] acoustic startle
response
The *myo7aa*[-/-] mutant is profoundly deaf and has little to no acoustic startle to sounds at
200, 300, 400, 500 or 600 Hz (Fig.8E), however a profound deafness does not equate to an
abolished ability to detect sounds if in the right environment given the correct cellular machinery.
We assessed whether L-type voltage-gated calcium channel agonists can improve the ability of
myo7aa[-/-] mutants to startle to an acoustic stimulus. Wildtype fish untreated responded most
frequently at 200 Hz with a mean response rate of 0.70 and the lowest mean response rate at
600 Hz. Wildtype fish treated with 5 µM (±)-Bay K 8644 had decreased responses at all
frequencies, except 200 Hz (Table S5). Incubation with 125 µM (R)-Baclofen also decreased
acoustic startle responses at all frequencies, however this may be due to the sedation effect of
this drug (Finnimore et al., 1995, Renier et al., 2007) (Fig. 8D, Table S5). Interestingly, both 5
µM (±)-Bay K 8644 and 250 µM Nefiracetam increased acoustic startle response in the *myo7aa*[-]
[/-] mutant fish, however, (±)-Bay K 8644 had a more robust effect and increased responses at all
frequencies. We did not see any improvement of *myo7aa*[-/-] mutant acoustic startle response
upon incubation with 125 µM (R)-Baclofen (Fig. 8E, Table S5).
292
DISCUSSION
The zebrafish model of USH1 was first characterized in 2000 and shows phenotypes
similar to human patients (Ernest et al., 2000). The *myo7aa*[-/-] mutants are deaf at the onset of
hearing (5 dpf), have vestibular abnormalities manifested in abnormal circular and loop like
swimming, have irregular stereocilia and do not uptake the FM1-43 vital dye, indicating an
inactive MET channel. In the present study we identified that 1) the *myo7aa*[-/-] mutants have
comparable areas of the ribeye protein in the ribbon synapse and significantly fewer tethered
vesicles; 2) the *myo7aa*[-/-] mutants have significantly fewer numbers of active hair cells, fewer
total CTBP2 puncta and a different distribution of CTBP2 puncta compared to wildtype; 3) the
myo7aa[-/-] mutants have significantly fewer numbers of active post-synaptic cells and fewer total
MAGUK puncta; 4) the turning angles of the *myo7aa*[-/-] mutants are larger compared to wildtype
turning angles; and 5) the *myo7aa*[-/-] mutants lack startle to a sound stimulus. We hypothesized
that using L-type voltage gated calcium channel agonists will restore the abnormal phenotypes.

We discovered that in the *myo7aa*[-/-] mutants 1) (R)-Baclofen increases ribbon area and number

of tethered vesicles 2) All three L-type voltage-gated calcium channel agonists change the

distribution of CTBP2 puncta to resemble wildtype distribution. 3) L-type voltage-gated calcium

channel agonists improve swimming behavior and decrease turning angle 4) (±)-Bay K 8644

and Nefiracetam improve acoustic startle response in the *myo7aa*[-/-] mutants.

Ribbon synapses are only found in sensory cells of vertebrate animals. They are

specialized presynaptic synapses that contain an electron dense structure that tethers synaptic

vesicles near voltage-gated calcium channels at the active zone of the hair cell. The main

component of a ribbon synapse is Ribeye with a monolayer of tethered glutamatergic synaptic

vesicles (Schmitz et al., 2000). It has previously been reported that there are 3-5 single ribbons

per hair cell in zebrafish (Nicolson, 2015). In this study we investigated if there is a difference in

the ribbon synapse structure between the *myo7aa*[-/-] mutants and wildtype in ultrastructure and

expression of genes involved in ribbon synapse structure and function. Through TEM we

observed that synaptic ribbons were present at the synapse in the *myo7aa*[-/-] mutants indicating

that the machinery to generate and localize the ribbons is functional in this model. We also

observed that the ribbons in the *myo7aa*[-/-] mutants possess an electron dense structure that has

a halo of tethered glutamatergic vesicles. Current understanding of this model's hair cell is that it

is not functional and is unable to transmit signal, however we challenge this view with our data

that indicates the ribbons are generated with the correct components, are localizing to the

synapse and the vesicles are fusing with the membrane to release neurotransmitter into the

synaptic cleft. It has also been reported that hair cell innervation is important for the correct

localization of the ribbons at the plasma membrane and for the regulation of ribbon size and

number (Suli et al., 2016). They also conclude that the physical presence of afferent fibers is

enough to regulate ribbon formation. Therefore, our observation of the presence of ribbon

structures at the synapse in the *myo7aa*[-/-] mutants indicates that the hair cell is capable of

stimulating afferent fibers. We argue that this signal in the *myo7aa*[-/-] mutants may not be

sufficient to detect meaningful sound; however the ability of the cell to perform is intact. These

observations support that cells are present and could respond to pharmacologic therapy.

Additionally, Nicolson et al. conducted whole-cell patch-clamp recordings of wildtype and

myo7aa[-/-] mutants and showed that the potential of the *myo7aa*[-/-] mutant is within the range of

wildtype indicating that defects in this mutant hair cell do not cause an abnormal resting

membrane potential and that the hair cell is still capable of transmitting a signal (Nicolson et al.,

1998).

We identified that the area of the ribbon synapse does not differ between wildtype and

myo7aa[-/-] mutants. It was also observed that ribbon size in another mechanotransduction

mutant, *sputnik*-which harbors a mutation in *cadherin 23*- did not differ from wildtype (Suli et al.,

2016). However, the *myo7aa*[-/-] mutants did have significantly fewer tethered vesicles. This

further prompts the questions as to why there are fewer tethered vesicles. We hypothesize that

this could be due to a decreased number of free cytosolic synaptic vesicles that would then be

recruited at the opposite pole of the ribbon to replace docked vesicles that exocytose. This

decrease in cytosolic synaptic vesicles would then lead to a decreased number of docked

vesicles and therefore less neurotransmitter released into the synaptic cleft. In order to test this

hypothesis cytosolic synaptic vesicles and docked vesicles would need to be quantified. Another

possibility that could explain why there are fewer tethered vesicles in the *myo7aa*[-/-] mutants is

that there is a disruption in the checkpoint for the attachment of synaptic vesicles to the dense

body.

CTBP2 is alternatively spliced to produce CTBP2 and Ribeye, which is the central

component of a ribbon synapse (Sheets et al., 2011, Wan et al., 2005). Sheets et al. identified

that in zebrafish Ribeye is essential to cluster L-type voltage-gated calcium channels ($Ca_v1.3a$)

to the active zone at the basal end of the hair cell (2011). This was also described in cochlear

inner ear hair cells (Frank et al., 2010). We explored the expression profile of CTBP2 in the

myo7aa[-/-] mutants and discovered that *myo7aa*[-/-] mutants have significantly fewer active hair

cells –defined as a hair cell containing at least one CTBP2 puncta-, fewer total CTBP2 puncta

and a different distribution of CTBP2 puncta compared to wildtype. It was previously reported

that hair cells of zebrafish larvae typically contain 3-5 ribbons (Nicolson, 2015, Obholzer et al.,

2008). We identified that most wildtype hair cells contain three CTBP2 puncta and that most

myo7aa[-/-] mutant hair cells have two CTBP2 puncta. Regus-Leidig et al. identified that

regulation of RIBEYE A and B-domains controls the assembly and plasticity of synaptic ribbons

and interactions between these two domains are inhibited by low concentrations of $NAD(H^+)$

(Magupalli et al., 2008, Regus-Leidig et al., 2010). Therefore, we suspect that the decreased

expression of CTBP2 in the *myo7aa*[-/-] mutants could be explained by possible mitochondrial

defects, however further analysis of mitochondria structure, function and by-products are

necessary.

It is well established that pre-synaptic cell activity affects post-synaptic cell activity and

vice versa (Luscher et al., 2000, Matus, 1999, Muller, 1997). Therefore, we used the post-

synaptic cell marker, MAGUK to assess the post-synaptic cell and how it relates to what was

observed in the ribbon synapse in the pre-synaptic cell. MAGUK is a kinase family that targets

and anchors glutamate receptors to the synaptic terminal (Oliva et al., 2012). We discovered

that *myo7aa*$^{-/-}$ mutants have significantly fewer number of active post-synaptic cells-defined as a

cell with at least one MAGUK puncta- and fewer total MAGUK puncta per neuromast. We

identified that most wildtype hair cells contain two to three MAGUK puncta and that most

myo7aa$^{-/-}$ mutant hair cells have three MAGUK puncta. This is the first report of the distribution

of CTBP2 and MAGUK puncta in wildtype and in any hearing defective mutant. By

understanding how hair cell biology affects the post-synaptic cell we are able to speculate as to

why the hair cell functions in a specific way. We hypothesize that the smaller number of CTBP2

puncta in the *myo7aa*$^{-/-}$ mutant is because the hair cell is not reaching the threshold required for

normal expression of CTBP2. This may be due to a decrease in intracellular calcium

concentrations or a decrease in expression of calcium binding proteins required for ribbon

synapse formation. The decreased CTBP2 expression coupled with the data that the ribbon is

localized to the synapse supports the hypothesis that the hair cell is releasing neurotransmitter

into the synaptic cleft. It is interesting to note that although the total number of active post-

synaptic cells and total number of MAGUK puncta is decreased, the distribution of the puncta is

not different from wildtype. We know that MAGUKs are scaffolding proteins that maintain and

modulate synaptic strength; therefore we believe that the fewer total number of MAGUK puncta

in *myo7aa*$^{-/-}$ mutant post-synaptic cells is a compensatory mechanism in which the post-synaptic

cells are responding to the decreased glutamate release from the hair cell. We can conclude

that there is some regulation of AMPA and NMDA receptors leading to synaptic plasticity. This

would need to be confirmed by analyzing the number of intact synapses by co-labeling with both

antibodies and then determining if there are orphan post-synaptic densities.

A large-scale mutagenesis screen conducted in 1996 resulted in a large number of

homozygous recessive mutants that exhibit behavioral defects (Granato et al., 1996). This study

also included 15 morphologically normal mutants with balance abnormalities including the

mariner mutant which displays looping and circular swimming starting at 4 dpf, develops a

curved spine (that augments the abnormal swimming) at 5 dpf, and fails to develop a properly

inflated swim bladder. We utilized a previously published behavior assessment that includes

tracking zebrafish swimming in response to a static stimulus and were able to quantify this

abnormal swimming behavior. We quantified the absolute smooth orientation, which is the sum

of all turning angles a fish swam in one swimming episode as a function of time. The absolute

smooth orientation of *myo7aa*$^{-/-}$ mutants is much larger compared to wildtype. This is also

observed in the swimming trajectories where the *myo7aa*$^{-/-}$ mutants display circular and looping

swimming and the wildtype animals have smooth trajectories and swim close to the edge of the
well.
The objective of this study was to test the hypothesis that L-type voltage-gated calcium
channel agonists can improve the abnormal cellular and behavioral phenotypes observed in the
myo7aa[-/-] mutant. The rationale for this hypothesis stems from understanding that the *myo7aa*[-/-]
mutant does not have a functional myo7aa protein therefore is unable to properly gate the MET
channel, the membrane potential required for depolarization is not achieved, the L-type voltage-
gated calcium channel does not open and synaptic transmission does not occur. We
hypothesize that increasing the downstream signal in *myo7aa*[-/-] mutant hair cells by increasing
intracellular calcium can reconstitute a new functional response to sound by increasing the
sensitivity of the calcium channel through treatment with L-type voltage-gated calcium channel
agonists. The three drugs we tested are (±)-Bay K 8644, Nefiracetam, and (R)-Baclofen. (±)-Bay
K 8644 is a potent L-type calcium channel activator (Nowycky et al., 1985, Schramm et al.,
1983). It has been used extensively to study regulation of calcium influx and the role of L-type
calcium channels in synaptic transmission in neuronal cells (Bonci et al., 1998, Monjaraz et al.,
2000, Watanabe et al., 1998). Nefiracetam is a nootropic or cognitive enhancer that activates
both L and N-type calcium channels (Yoshii and Watabe, 1994, Yoshii et al., 2000) and may
modulate the GABA receptor currents by interacting with a PKA pathway (Huang et al., 1996)
and PKC pathway (Nishizaki et al., 1998). Nootropic agents can influence various
neurotransmissions including dopaminergic, cholinergic, glutamatergic and GABAergic
pathways (Funk and Schmidt, 1984, Giovannini et al., 1993, Huang et al., 1996, Marchi et al.,
1990, Nabeshima et al., 1990, Nishizaki et al., 1998, Oyaizu and Narahashi, 1999, Watabe et
al., 1993). Additionally, it has been demonstrated that Nefiracetam affects acetylcholine induced
currents and NMDA receptors (Moriguchi et al., 2003, Narahashi et al., 2003, Zhao et al., 2001).
NMDA and AMPA receptors are ionotropic glutamate receptors that are nonselective cation
channels (K^+ and Na^+). Activation of these channels produces excitatory post-synaptic
responses. NMDA receptors allow not only the entry of potassium and sodium ions, but also
calcium; this increase in intracellular calcium acts as a secondary messenger to augment
intracellular signaling cascades leading to the enrichment of synaptic activity. Previous studies
have used Nefiracetam hypothesizing that this mechanism of action could improve post-stroke
depression, post-stroke apathy and Alzheimer's disease (Robert G. Robinson et al., 2009,
Starkstein et al., 2016). Lastly, (R)-Baclofen was previously shown to act as $GABA_B$ agonist
(Bowery et al., 1983). $GABA_B$ receptors are coupled through G-proteins to potassium channels,
calcium channels, and adenylate-cyclase. However, more recently $GABA_B$ receptors have been

shown to modulate various voltage-gated calcium channels (Carter and Sabatini, 2004, Shen

and Slaughter, 1999), specifically a direct interaction was identified between the L-type voltage-

gated calcium channel ($Ca_v1.3$) and $GABA_B$ receptor subunit 2 (Park et al., 2010). They

discovered that activation of the $GABA_B$ receptor increases L-type calcium currents. (R)-

Baclofen has been used in multiple recent clinical trials to treat Fragile X (Berry-Kravis et al.,

2017) and autism (Veenstra-VanderWeele et al., 2017); and racemic Baclofen is an FDA

approved drug currently used to treat muscle spasticity (Ertzgaard et al., 2017, Rizzo et al.,

2004).

Modulating the L-type voltage-gated calcium channel activity was the central aim for

using these three compounds. Sheets et al. reports that $Ca_v1.3a$ conductance affects the

morphology of ribbon synapses; specifically identifying that 3 dpf zebrafish embryos had a

decrease in $Ca_v1.3a$ conductance causing a larger ribbon area, but when they blocked $Ca_v1.3a$

with Isradipine at 5 dpf there was no change indicating that there is not much of an effect at

more mature ribbon synapses (Sheets et al., 2012). They suggested that there is a period of

synaptic maturation that is highly influenced by calcium flux through the $Ca_v1.3$ that affects

synaptic ribbons. We argue that this point is at 4 dpf in the development of inner ear hair cells

and is the reason we chose to treat zebrafish at this developmental state for an overnight

incubation.

Using TEM, we demonstrate that (R)-Baclofen increases ribbon area and number of

tethered vesicles in the *myo7aa*$^{-/-}$ mutant ribbon synapse which is indistinguishable from

wildtype. Incubation with (±)-Bay K 8644 and Nefiracetam did not have an effect on ribbon area

and number of tethered vesicles on the *myo7aa*$^{-/-}$ mutants. Therefore, the mechanism in which

(R)-Baclofen acts on the *myo7aa*$^{-/-}$ mutant ribbon synapses to increase ribbon area and number

of tethered vesicles may not be directly and exclusively attributed to agonistic action of $Ca_v1.3a$,

however our study is the first to demonstrate that ribbon synapses of the *myo7aa*$^{-/-}$ mutant can

be altered with drug incubation to more closely resemble wildtype ribbon synapses. We did not

see any adverse effects in the ribbon area or number of tethered vesicles on wildtype larvae

upon incubation with the three drugs.

In addition to demonstrating that (R)-Baclofen can alter ribbon synapse area and number

of tethered vesicles, we demonstrated that all three compounds change the distribution of

CTBP2 puncta. Our finding that most wildtype hair cells have two CTBP2 puncta and most

myo7aa$^{-/-}$ mutant hair cells have three CTBP2 puncta establishes that this difference may be

contributing to the phenotype of the *myo7aa*$^{-/-}$ mutant. Our data shows that incubation with all

three L-type voltage-gated calcium channel agonists shifts the distribution of CTBP2 puncta and

restores the distribution to resemble wildtype. This is an important discovery, because it

indicates that all three compounds are likely acting in the same manner in their effect on CTBP2

distribution. Since (±)-Bay K 8644 only has one mechanism of action, L-type voltage-gated

calcium channel agonist-we can presume that Nefiracetam and (R)-Baclofen effects may be

primarily through the same pathway. Increasing intracellular calcium through this mechanism

may have an effect on the regulation of CTBP2. We also identified that (±)-Bay K 8644

significantly increased the total number of active hair cells and post-synaptic cells per

neuromast and the total number of CTBP2 and MAGUK puncta per neuromast. It is likely that

only (±)-Bay K 8644 had this effect because it is the most potent of the three and its only

mechanism of action is to act on the $Ca_v1.3a$ channel. We also observed that upon incubation

with $5~\mu M$ (±)-Bay K 8644 the distribution of CTBP2 puncta did shift in wildtype larvae so that

most hair cells have two CTBP2 puncta rather than three which is observed in the untreated

wildtype groups. One possible explanation could be that we are observing a dose response that

takes a bell-shaped curve meaning that some doses (lower or higher) do not give a specific

outcome whereas those in the middle do. In this case, we suspect that $5~\mu M$ is a dose that falls

in the middle of a bell shaped curve and would elicit a response. This hypothesis could be

tested by measuring CTBP2 expression with higher and lower doses of (±)-Bay K 8644. Recent

work shows that evoked presynaptic calcium influx through L-type voltage-gated calcium

channels stimulates calcium uptake in the mitochondria causing decreased cellular NAD(H) thus

decreasing ribbon formation (Wong et al., 2019), which may explain why $5~\mu M$ (±)-Bay K 8644

shifts the distribution of CTBP2 puncta in wildtype hair cells, however this is unlikely in this

situation because we see that (±)-Bay K 8644 actually increases the total number of active hair

cells and post-synaptic cells per neuromast and the total number of CTBP2 and MAGUK puncta

per neuromast. Lastly, (R)-Baclofen also increased the number of active post-synaptic cells and

total number of MAGUK puncta in wildtype larvae.

It is well characterized that zebrafish demonstrate a short-latency startle response, the

C-start, in response to tactile and auditory stimuli that consists of a C-bend of the body followed

by a counter bend and then a swimming burst (Eaton et al., 1991, Kimmel et al., 1974). This

response is a result of activation of the Mauthner cells (M-cells). Zottoli et al. linked Mauthner

cell activity to escape behavior (Zottoli, 1977). We utilized a behavior assessment that tracks

zebrafish swimming behavior and quantifies the absolute smooth orientation or turning angle as

a function of time in response to a static stimulus. We identified that the absolute smooth

orientation of *myo7aa*$^{-/-}$ mutants is much larger compared to wildtype. We showed improved

swimming behavior in the *myo7aa*$^{-/-}$ mutants meaning a decrease in turning angle using all three

L-type voltage-gated calcium channel agonists; however the most robust response was with (±)-
Bay K 8644, which resulted in turning angles indistinguishablefrom wildtype. We tested all three
drugs in wildtype animals as well and observed no change in the absolute smooth orientation at
these specific doses with the exception of (±)-Bay K 8644. This could be explained by the fact
that the doses chosen for Nefiracetam and (R)-Baclofen were chosen based on dose-response
studies (Figs. S4 & S6). These data show that changes in the absolute smooth orientation are
observed in wildtype larvae depending on the dose. We did not conduct a dose response study
for (±)-Bay K 8644, therefore we would expect that other doses would result in no significant
changes in swimming behavior.
Not only did our quantitative analysis of the swimming behavior indicate improvement,
swimming trajectories of individual fish from each group also demonstrate less circular and
looping swimming and more smooth and zig-zag swimming. It is interesting to note that all
wildtype fish (untreated or treated) had a preference for swimming toward the edge of the well, a
process called thigmotaxis. Peitsaro et al. report that thigmotaxis in zebrafish is an indicator of
anxiety (Peitsaro et al., 2003). This has also been shown in mice (Simon et al., 1994). It is
unclear if this process is disrupted in the *myo7aa$^{-/-}$* mutants or if the abnormal swimming
behavior is what prevents the fish from remaining at the edge of the well.
The acoustic startle response can be elicited starting at 5 dpf, which is when the ear is
completely developed (Eaton and Didomenico, 1986, Zeddies and Fay, 2005). The zebrafish
has two sensory organs: the inner ear and lateral line system. The inner ear contains three
otolithic organs: the utricle, the saccule and at 11-12 dpf the lagena develops (Popper and Fay,
1993). Saccular hair cells respond to relatively high frequencies (200 Hz-1200 Hz) and fibers
that innervate the saccule synapse on M-cells (Bang et al., 2002, Fay, 1995, Lin and Faber,
1988, Zottoli, 1977). The lateral line responds to much lower frequencies (50-100 Hz) (Coombs,
1994, Coombs and Janssen, 1990). There are multiple hearing assessments previously
reported including acoustically evoked behavioral responses, tone burst, pre-pulse inhibition
assays, microphonic potentials and auditory brainstem responses (Bang et al., 2002,
Bhandiwad et al., 2013, Burgess and Granato, 2007, Higgs et al., 2002, Yang et al., 2017, Yao
et al., 2016, Zeddies and Fay, 2005). However, all of these assessments require specialized
equipment and analysis tools that would provide more information than what we required to
asses presence or absence of an acoustic startle response. We, therefore, sought to develop a
cost-effective and accurate hearing assessment tool that can be accessible to others. In our
hearing assessment we acquired videos of 5 dpf larval fish placed on a speaker and
administered frequencies ranging from 200 to 600 Hz. We used *myo7aa$^{-/-}$* mutants as a control

for the experimental design and utilized an intensity level that stimulated the wildtype fish but did

not cause a startle in the *myo7aa*[-/-] mutants. Using this paradigm, we identified that untreated

wildtype fish do not respond with a 100% response rate at any frequency and we see the most

robust responses at 200 and 400 Hz. Incubation with (±)-Bay K 8644 and (R)-Baclofen

decreased mean response rate in the wildtype fish, however incubation with Nefiracetam

improved the mean response rate. A potential explanation as to why we are observing adverse

effects in wildtype fish upon incubation with (±)-Bay K 8644 in not only the acoustic startle but

also in the distribution of CTBP2 puncta and the absolute smooth orientation is that excessive

cytosolic calcium (altering calcium homeostasis) in the hair cell can be detrimental (Esterberg et

al., 2013). Another explanation could be that the elevated cytosolic calcium in the wildtype hair

cells due to (±)-Bay K 8644 could be causing excessive activation of the glutamate receptors

that has been linked to apoptotic hair-cell death (Sheets, 2017). We may be observing this

affect across various assessments with only (±)-Bay K 8644, in the wildtype animals because it

is the more potent drug. In other words, in a normal hair cell if the calcium concentration falls

above the threshold of homeostasis then it can have adverse effects on the cell. However, in a

hair cell that already has a starting calcium concentration below the threshold (*myo7aa*[-/-]

mutant) than pharmacologic strategies to increase intracellular calcium will bring the calcium

concentration to the level of homeostasis.

It has previously been reported that incubation with (R)-Baclofen can cause a sedative

effect in zebrafish that decreases locomotor activity that is dose dependent (Renier et al., 2007).

This study reports that doses as low as 10^{-4} M dropped the locomotor activity to about 25-30%

of what was observed in the controls. We are seeing that wildtype fish treated with 125 µM (R)-

Baclofen exhibit about 20% of the mean response rate of control fish at each individual

frequency.

Lastly, we tested the hypothesis that L-type voltage gated calcium channel agonists can

restore some hearing by increasing mean acoustic startle response rate in the *myo7aa*[-/-]

mutants. Our data suggests that (±)-Bay K 8644 and Nefiracetam improve acoustic startle

response in the *myo7aa*[-/-] mutants. This is an important discovery because it is the first report of

a pharmacotherapy restoring hearing in a zebrafish model of USH1. Additionally, like other

otophysan fishes, the zebrafish has a swimbladder that is coupled to the saccule through

Weberian ossicles that facilitates responses to sound pressure (Higgs et al., 2003). The

myo7aa[-/-] mutant does not develop a swimbladder. We tested if L-type voltage gated calcium

channels affect inflation of the swim bladder in the *myo7aa*[-/-] mutant and identified that

Nefiracetam and (R)-Baclofen did not result in an inflated swim bladder and did not negatively

affect wildtype swim bladders. This supports our hearing assessment data suggesting that the

myo7aa$^{-/-}$ mutant fish incubated in L-type voltage-gated calcium channel agonists that are

responding to acoustic stimuli are responding to direct acceleration of the cells within the otolith

organs, specifically the saccule and not to changes in sound pressure.

Conclusion

Here we report the first study using L-type voltage-gated calcium channel agonists to

treat the zebrafish model of USH1, the *myo7aa*$^{-/-}$ mutant. We have shown through transmission

electron microscopy and immunohistochemistry that the hair cell biology of the *myo7aa*$^{-/-}$ mutant

differs from wildtype in that there are fewer glutamatergic vesicles tethered to the ribbon

synapse, fewer total active hair cells and post-synaptic cells, fewer total CTBP2 and MAGUK

puncta, and a different distribution of CTBP2 puncta. We quantified the previously observed

behavioral phenotypes and identified that *myo7aa*$^{-/-}$ fish have larger turning angles and little to

no acoustic startle response. Upon treatment with L-type voltage-gated calcium channel

agonists, we discovered changes in hair cell morphology/ultrastructure, gene expression and

behavior through swimming and hearing assessments. Our data support that treatment with

these compounds restores the abnormalities in the *myo7aa*$^{-/-}$ mutants to more closely resemble

wildtype by 1) increasing the number of tethered vesicles to the ribbon 2) shifting the distribution

of CTBP2 puncta 3) decreasing turning angle and improving swimming behavior and 4)

improving acoustic startle. However, there are still many unanswered questions in regards to

localization and density of L-type voltage-gated calcium channels, ratio of ribbons at the

synapse versus those that are not, mitochondrial contribution to the phenotype or the rescue

and direct quantification of calcium concentrations with and without treatment. We are confident

that the results we report in this study represent a significant step towards understanding the

mechanism of disease and discovering compounds to treat hearing loss caused by pathogenic

variants in *MYO7A.*

MATERIALS AND METHODS

Zebrafish maintenance and husbandry

All animals used in this study were zebrafish (*Danio rerio).* The zebrafish strains used for all

experiments were the tc320b allele (c.2699T>A; p.Tyr846Stop) in exon 21 of *myo7aa* (Ernest et

al., 2000) and wildtype Segrest strains were obtained from the laboratory stock at Mayo Clinic

(Leveque et al. 2016). Heterozygous adults for the *myo7aa* mutation were used to generate

homozygous larval fish (*myo7aa*$^{-/-}$). Animals were raised in a 10-hour dark and 14-hour light

cycle with embryos and adults kept at 28.5°C (Westerfield, 1995). Embryos were kept in Petri

dishes and fresh embryo water was replaced daily. Larvae were kept at room temperature for all

experiments and were maintained in the incubator at all other times. Wildtype Segrest strains

and heterozygous adult *myo7aa* fish were also maintained at the University of Minnesota, Twin

Cities Zebrafish Core Facility. The Institutional Animal Care and Use Committee approved all

experimental procedures both at Mayo Clinic and the University of Minnesota.

Phenotyping and genotyping

The *myo7aa*[-/-] tc320b allele results from a single nucleotide polymorphism A-T, which results in

an early stop codon p.Tyr846Stop (Ernest et al., 2000). Homozygous recessive mutant larval

fish were identified at 4 dpf (days post fertilization) based on various phenotypic traits, including

defective balance, absent startle to sound, and circular or looping swimming patterns. DNA from

whole embryos or fin biopsies was extracted for allele-specific quantitative PCR for genotyping

as previously described (Lee et al., 2016).

Hair cell staining

In order to stain the stereocilia of the saccule hair cells, embryos were initially fixed overnight in

freshly thawed 4% paraformaldehyde at 4°C. Two washes for 5 minutes each in 0.2% PBSTx

were conducted at room temperature. Embryos were then incubated for 4 days in 2% PBSTx at

4°C to completely dissolve the otoliths. F-actin staining occurred for up to 2 hours at room

temperature using Alexa Fluor 488 Phalloidin (Molecular Probes, Eugene, OR), which was

diluted 1:20 in PBSTw. Embryos were then washed six times for 10 minutes at room

temperature in 0.2% PBSTx. Hair cell stereocilia were observed and imaged using the LSM 780

inverted confocal microscope run on Zen software package (Zeiss).

MET channel assessment

Mechanotransduction channel activity was assessed through FM 1-43 dye uptake (Molecular

Probes, Eugene, OR). A working stock of 3 µM FM 1-43 was used for a 30 second incubation

and rinsed three times immediately with embryo water. Lateral line neuromasts were imaged for

FM 1-43 uptake using a Lightsheet Z.1 microscope (Zeiss). Lateral-orientated z-stacks at 5X/1.0

NA water-dipping objective (Zeiss) and an RFP (emission filter BP 505-545 LP 585) optical filter

(Zeiss). Each corresponding dorsal image is a maximal image projection generated from z-

dimension stacks.

Drug incubation

The maximum tolerated dose was determined for all drugs used in this study. This was

performed by incubating 4 dpf wildtype embryos individually in 24-well plates in 1 mL total

volume of embryo water including various doses of dissolved drug. Control fish were incubated

in the same concentration of embryo water with only drug vehicle (e.g. DMSO, H_2O). The 24-

well plates were placed in a 29°C embryo incubator for the duration of the drug treatment. Heart

rates, startle responses, swimming behavior and overall health were observed after one hour,

four hours and overnight incubation to determine any adverse effects or possible toxicity. Drug

doses started at 0.5 µM and increased based on overall health and survival of the embryos. The

drugs used in this study are as follows: (±)-Bay K 8644 (catalog # 1544, Batch No. 4; Tocris),

Nefiracetam (catalog # 2851, Batch No. 1; Tocris) and (R)-Baclofen (catalog # 0796, Batch No.

3; Tocris). (±)-Bay K 8644 and Nefiracetam were dissolved in DMSO to a 100 mM stock solution

and (R)-Baclofen was dissolved in water (with gentle warming) to a 20 mM stock solution. The

final doses that would be used in the drug treatments were determined by identifying the highest

tolerated dose and using the dose one step lower. A control experiment was conducted to

determine the effect of DMSO on behavior and at the concentrations used for dissolving drugs,

and no adverse effects were identified.

Video acquisition of swimming behavior

Videos were made according to the video acquisition protocol previously described (Lambert et

al. 2012). High-speed videos of 5dpf larvae were acquired at either 120 or 180 frames per

second (fps) using FlyCap software and the Point Grey Firefly. A LED light stage was used

underneath a 20 mm arena in order to increase illumination for a higher quality video. A 50 V

electric shock (5 ms duration) was used to startle the animal every 20 seconds. An LED light

was used to confirm the administration of the electric shock.

Video tracking and analysis

Videos acquired from the FlyCap software were processed as fmf files through Ctrax: The

Caltech Multiple Fly Tracker, free source software (http://ctrax.sourceforge.net/) (Branson et al.,

2009). The tracking settings for motion, observation and hindsight are listed in Table S1. Videos

were exported as mat files, fixed for errors using the Fix Errors Matlab Toolbox (FEMT),

provided by (Branson et al., 2009), and analyzed in Matlab version 2012b using in-house scripts

from the Masino Laboratory (available upon request). Swimming trajectories were generated

using the Ctrax Behavioral Microarray Matlab toolbox. A circle indicating the perimeter of the

testing well was added to each image as a reference point. Each electric shock was confirmed

by determining the frame in which the LED light appears using Image J. The frame range

corresponding to that swimming episode was found in the mat files. The values for the absolute

smooth orientation of the fish during one swimming episode were totaled and then divided by

the episode duration. This was performed for all swimming episodes.

Hearing assessment

Up to ten 4 dpf embryos were placed in 60 X 15 mm Petri dishes and placed in a 29°C embryo

incubator in an isolated behavior room for overnight incubation. At 5dpf embryos were placed on

top of a Bose Soundlink Color Speaker and allowed up to 5 min to acclimate. The speaker was

placed on top of an illuminator for a better quality video. A white piece of paper was placed

beneath the Petri dish to lighten the background in order to create contrast between the larval

fish and the speaker. A script written in Matlab (available upon request) was used to generate 3

one second stimuli every 20 seconds for the following frequencies: 200, 300, 400, 500 and 600

Hz and administered via Bluetooth connection between the laptop and the speaker. The

intensity of the sounds administered was determined using an Extech sound level meter, around

77 dB. Videos were taken using a Sony Handycam HDR cx560 and were de-identified and

blindly scored by a trained laboratory member for startle responses. Responses were scored as

either 0 (no response) or 1 (response) for each fish at each stimulus. Mean values of responses

and no responses were calculated and the mean variance was determined.

Processing and imaging of zebrafish for TEM

5dpf embryos were fixed in Trump's fixative (4 % paraformaldehyde with 1% glutaraldehyde) for

a minimum of 24 hr. Following fixation, embryos were secondarily fixed with 1% osmium

tetroxide and 2% uranyl acetate, dehydrated through an ethanol series and embedded into

Embed 812/Araldite resin. Processing was facilitated with the use of a BioWave laboratory

microwave oven (Pelco Biowave 3450, Ted Pella, Inc., Redding, CA). After a 24 hr

polymerization at 60°C, ultrathin sections (0.1 mm) were stained with 2% uranyl acetate and

lead citrate. Micrographs were acquired in the Mayo Clinic Microscopy and Cell Analysis Core

using a JEOL1400 or JEOL1400Plus transmission electron microscope operating at 80 kV

(Peabody, MA) equipped with a Gatan Orius camera (Gatan, Inc., Warrendale, PA).

Analysis of ribbon synapse TEM images

Transmission electron microscopy images of ribbon synapse structures were de-identified and
analyzed for ribbon area and number of tethered vesicles. ImageJ software was used to
determine the area of the electron dense structure and the number of vesicles was quantified
manually.
715

Immunohistochemistry
The preparation and the immunohistochemistry (IHC) experiments were conducted across five
days using a nutator unless otherwise specified. The first day included fixing 5dpf embryos in
4% paraformaldehyde at 4°C for overnight incubation. Samples were then rinsed twice and
washed three times for 10 minutes each with PBTr (0.02% Triton-X 100 in 1X PBS), washed for
5 minutes in a 1:1 mixture (PBTr and 100% MetOH) and then washed three times for 5 minutes
in 100% MetOH. They were stored at -20°C if not used immediately. The third day included
rehydrating the embryos through a 5-minute wash of the 1:1 mixture followed by three 5-minute
washes with PBTr. Samples were permeabilized for 50 minutes with proteinase K (diluted
1:2000 in PBTr) and then washed five times for 5 minutes with PBTr. Block buffer included 10%
Normal Goat Serum, 0.2% Triton-x 100, 2% DMSO in 1X PBS. Samples were incubated in
block buffer for at least 3 hours. The primary antibodies used in this study were mouse
monoclonal against membrane associated guanylate kinase, a post-synaptic membrane marker
(MAGUK; 1:250 NeuroMab Clone K28/86, Catalog # 75-029) (Anderson 1996; Lv et al. 2016;
Sheets et al. 2011)) and CTBP2 (1:500 Santa Cruz Catalog # Sc-55502)(Lv et al. 2016) diluted
in prepared block buffer . Samples were incubated in diluted primary antibody at 4°C overnight.
The fourth day included washing samples six times for 30 minutes with PBTr. The secondary
antibodies used were Alexa Fluor 647 goat anti-mouse (1:250 Thermo-Fisher/Invitrogen Catalog
A21235) and Alexa Fluor 488 goat anti-mouse (1:500 Thermo-Fisher/Invitrogen Catalog
A21121) respectively and were diluted in PBTr. Samples were incubated in diluted secondary
antibody at 4°C overnight. Lastly, samples were washed again six times for 30 minutes each
wash with PBTr and washed three times for 5 minutes with 1X PBS. Samples were stored in 1X
PBS in 4°C. Images were acquired using the LSM 780 inverted confocal microscope run on Zen
software package (Zeiss). Samples were mounted with Vectashield Hardset DAPI Mounting
Medium, and covered with a cover slip for approximately 10 minutes before imaging. All images
were captured with the 20x objective and a 5.0 magnitude zoom. All images were de-identified
prior to analysis using ImageJ software. The proportion of active cells with 1,2,3,4,5,6+ puncta
was determined by identifying the number of cells with n puncta divided by the total number of
active cells in that specific neuromast.

Statistical analysis

Statistical analyses were performed using JMP software (JMP® 1989-2019). Each treatment

group was individually compared to the corresponding control group (wildtype or *myo7aa*[-/-]

untreated) through a student's t-test. Acoustic startle response and CTBP2/MAGUK puncta

statistical analysis was performed using a two-tailed Fischer's exact test

(https://www.graphpad.com/quickcalcs/contingency1/). All statistical analyses are presented

with appropriate significant digits.

Acknowledgements

We thank Dr. Teresa Nicolson for providing us with the tc320b allele of *mariner*. We are grateful

to Mark Curry and Patrick Jochim in the Mayo Clinic Media Services. We also thank Gabriel

Martinez and Dr. Timothy Wiggins for help with Adobe Illustrator, Matlab and the behavior

assessment. We are also grateful to Dr. Lavinia Sheets for her helpful discussions. Lastly, we

would like to thank the staff members in the zebrafish core facilities at Mayo Clinic and the

University of Minnesota, Twin Cities.

Competing Interests

No competing interests declared**.**

Funding

A.K was supported by the CTSA Grant Number UL1 TR000135 from the National Center for

Advancing Translational Sciences (NCATS), a component of the National Institutes of Health

(NIH). L.A.S. was supported by funding from the Mayo Clinic Center for Individualized Medicine,

Philanthropic support from the Nelson Family Fund and from the Mayo Clinic Department of

Otorhinolaryngology. This paper's contents are solely the responsibility of the authors and do

not necessarily represent the official view of the National Institutes of Health.

Data Availability

All data required for analysis is included within the main text.

Author Contributions

Conceptualization-A.K, S.C.E, L.A.S

Methodology- A.K, A.M.L, M.A.M

Software- A.K, A.M.L, M.A.M

Validation- A.K, J.D, M.A.M

Formal Analysis- A.K

Investigation- A.K, J.D, J.B, T.C

Resources- A.K, J.D, M.A.M, S.C.E, L.A.S

Data Curation- A.K., A.M.L, M.A.M

Writing-original draft preparation- A.K.

Writing-review and editing-A.K, J.D., T.C., J.B., A.M.L, M.A.M, S.C.E, L.A.S

Visualization- A.K.

Supervision- S.C.E, L.A.S

Project administration-A.K, S.C.E, L.A.S

Funding acquisition-L.A.S

References

AHMED, Z. M., FROLENKOV, G. I. & RIAZUDDIN, S. 2013. Usher proteins in inner ear
structure and function. *Physiol Genomics,* 45**,** 987-9.
AHMED, Z. M., GOODYEAR, R., RIAZUDDIN, S., LAGZIEL, A., LEGAN, P. K., BEHRA, M.,
BURGESS, S. M., LILLEY, K. S., WILCOX, E. R., RIAZUDDIN, S., GRIFFITH, A. J.,
FROLENKOV, G. I., BELYANTSEVA, I. A., RICHARDSON, G. P. & FRIEDMAN, T. B.
2006. The tip-link antigen, a protein associated with the transduction complex of sensory
hair cells, is protocadherin-15. *J Neurosci,* 26**,** 7022-34.
BANG, P. I., YELICK, P. C., MALICKI, J. J. & SEWELL, W. F. 2002. High-throughput behavioral
screening method for detecting auditory response defects in zebrafish. *J Neurosci*
Methods, 118**,** 177-87.
BERRY-KRAVIS, E., HAGERMAN, R., VISOOTSAK, J., BUDIMIROVIC, D., KAUFMANN, W.
E., CHERUBINI, M., ZAREVICS, P., WALTON-BOWEN, K., WANG, P., BEAR, M. F. &
CARPENTER, R. L. 2017. Arbaclofen in fragile X syndrome: results of phase 3 trials. *J*
Neurodev Disord, 9**,** 3.
BEURG, M., FETTIPLACE, R., NAM, J. H. & RICCI, A. J. 2009. Localization of inner hair cell
mechanotransducer channels using high-speed calcium imaging. *Nat Neurosci,* 12**,** 553-
8.
BHANDIWAD, A. A., ZEDDIES, D. G., RAIBLE, D. W., RUBEL, E. W. & SISNEROS, J. A. 2013.
Auditory sensitivity of larval zebrafish (Danio rerio) measured using a behavioral
prepulse inhibition assay. *J Exp Biol,* 216**,** 3504-13.
BONCI, A., GRILLNER, P., MERCURI, N. B. & BERNARDI, G. 1998. L-Type Calcium Channels
Mediate a Slow Excitatory Synaptic Transmission in Rat Midbrain Dopaminergic
Neurons. *The Journal of Neuroscience,* 18**,** 6693-6703.
BOWERY, N. G., HILL, D. R. & HUDSON, A. L. 1983. Characteristics of GABAB receptor
binding sites on rat whole brain synaptic membranes. *Br J Pharmacol,* 78**,** 191-206.
BRANDT, A., KHIMICH, D. & MOSER, T. 2005. Few CaV1.3 channels regulate the exocytosis
of a synaptic vesicle at the hair cell ribbon synapse. *J Neurosci,* 25**,** 11577-85.
BRANSON, K., ROBIE, A. A., BENDER, J., PERONA, P. & DICKINSON, M. H. 2009. High-
throughput ethomics in large groups of Drosophila. *Nat Methods,* 6**,** 451-7.

BURGESS, H. A. & GRANATO, M. 2007. Sensorimotor gating in larval zebrafish. *J Neurosci,*
27**,** 4984-94.
CABERLOTTO, E., MICHEL, V., FOUCHER, I., BAHLOUL, A., GOODYEAR, R. J.,
PEPERMANS, E., MICHALSKI, N., PERFETTINI, I., ALEGRIA-PREVOT, O.,
CHARDENOUX, S., DO CRUZEIRO, M., HARDELIN, J. P., RICHARDSON, G. P.,
AVAN, P., WEIL, D. & PETIT, C. 2011. Usher type 1G protein sans is a critical
component of the tip-link complex, a structure controlling actin polymerization in
stereocilia. *Proc Natl Acad Sci U S A,* 108**,** 5825-30.
CARTER, A. G. & SABATINI, B. L. 2004. State-dependent calcium signaling in dendritic spines
of striatal medium spiny neurons. *Neuron,* 44**,** 483-93.
COMMUNICATION, C. F. H. A. *Statistics and Facts About Hearing Loss* [Online]. Available:
http://chchearing.org/facts-about-hearing-loss/ [Accessed].
COOMBS, S. 1994. Nearfield detection of dipole sources by the goldfish (Carassius auratus)
and the mottled sculpin (Cottus bairdi). *J Exp Biol,* 190**,** 109-29.
COOMBS, S. & JANSSEN, J. 1990. Behavioral and neurophysiological assessment of lateral
line sensitivity in the mottled sculpin, Cottus bairdi. *J Comp Physiol A,* 167**,** 557-67.
DAVENPORT SLH, O. G. 1977. The heterogeneity of Usher's syndrome. *In: Proceedings of the*
5th International Conference of Birth Defects, abstract 215.
EATON, R. C. & DIDOMENICO, R. 1986. Role of the Teleost Escape Response during
Development. *Transactions of the American Fisheries Society,* 115**,** 128-142.
EATON, R. C., DIDOMENICO, R. & NISSANOV, J. 1991. Role of the Mauthner cell in
sensorimotor integration by the brain stem escape network. *Brain Behav Evol,* 37**,** 272-
85.
ERNEST, S., RAUCH, G. J., HAFFTER, P., GEISLER, R., PETIT, C. & NICOLSON, T. 2000.
Mariner is defective in myosin VIIA: a zebrafish model for human hereditary deafness.
Hum Mol Genet, 9**,** 2189-96.
ERTZGAARD, P., CAMPO, C. & CALABRESE, A. 2017. Efficacy and safety of oral baclofen in
the management of spasticity: A rationale for intrathecal baclofen. *J Rehabil Med,* 49**,**
193-203.
ESTERBERG, R., HAILEY, D. W., COFFIN, A. B., RAIBLE, D. W. & RUBEL, E. W. 2013.
Disruption of intracellular calcium regulation is integral to aminoglycoside-induced hair
cell death. *J Neurosci,* 33**,** 7513-25.
FAY, R. R. 1995. Physiology of primary saccular afferents of goldfish: implications for Mauthner
cell response. *Brain Behav Evol,* 46**,** 141-50.
FINNIMORE, A. J., ROEBUCK, M., SAJKOV, D. & MCEVOY, R. D. 1995. The effects of the
GABA agonist, baclofen, on sleep and breathing. *Eur Respir J,* 8**,** 230-4.
FRANK, T., RUTHERFORD, M. A., STRENZKE, N., NEEF, A., PANGRSIC, T., KHIMICH, D.,
FEJTOVA, A., GUNDELFINGER, E. D., LIBERMAN, M. C., HARKE, B., BRYAN, K. E.,
LEE, A., EGNER, A., RIEDEL, D. & MOSER, T. 2010. Bassoon and the synaptic ribbon
organize Ca(2)+ channels and vesicles to add release sites and promote refilling.
Neuron, 68**,** 724-38.
FUCHS, P. A. 2005. Time and intensity coding at the hair cell's ribbon synapse. *J Physiol,* 566**,**
7-12.
FUNK, K. F. & SCHMIDT, J. 1984. Changes of dopamine metabolism by hypoxia and effect of
nootropic drugs. *Biomed Biochim Acta,* 43**,** 1301-4.
GIOVANNINI, M. G., RODINÒ, P., MUTOLO, D. & PEPEU, G. 1993. Oxiracetam and
aniracetam increase acetylcholine release from the rat hippocampus in vivo. *Drug*
Development Research, 28**,** 503-509.
GRANATO, M., VAN EEDEN, F. J., SCHACH, U., TROWE, T., BRAND, M., FURUTANI-SEIKI,
M., HAFFTER, P., HAMMERSCHMIDT, M., HEISENBERG, C. P., JIANG, Y. J., KANE,
D. A., KELSH, R. N., MULLINS, M. C., ODENTHAL, J. & NUSSLEIN-VOLHARD, C.

1996. Genes controlling and mediating locomotion behavior of the zebrafish embryo and
larva. *Development,* 123**,** 399-413.
GRATI, M. & KACHAR, B. 2011. Myosin VIIa and sans localization at stereocilia upper tip-link
density implicates these Usher syndrome proteins in mechanotransduction. *Proc Natl*
Acad Sci U S A, 108**,** 11476-81.
GRILLET, N., KAZMIERCZAK, P., XIONG, W., SCHWANDER, M., REYNOLDS, A.,
SAKAGUCHI, H., TOKITA, J., KACHAR, B. & MULLER, U. 2009a. The
mechanotransduction machinery of hair cells. *Sci Signal,* 2**,** pt5.
GRILLET, N., XIONG, W., REYNOLDS, A., KAZMIERCZAK, P., SATO, T., LILLO, C.,
DUMONT, R. A., HINTERMANN, E., SCZANIECKA, A., SCHWANDER, M., WILLIAMS,
D., KACHAR, B., GILLESPIE, P. G. & MULLER, U. 2009b. Harmonin mutations cause
mechanotransduction defects in cochlear hair cells. *Neuron,* 62**,** 375-87.
HASSON, T., HEINTZELMAN, M. B., SANTOS-SACCHI, J., COREY, D. P. & MOOSEKER, M.
S. 1995. Expression in cochlea and retina of myosin VIIa, the gene product defective in
Usher syndrome type 1B. *Proc Natl Acad Sci U S A,* 92**,** 9815-9.
HIGGS, D. M., ROLLO, A. K., SOUZA, M. J. & POPPER, A. N. 2003. Development of form and
function in peripheral auditory structures of the zebrafish (Danio rerio). *J Acoust Soc Am,*
113**,** 1145-54.
HIGGS, D. M., SOUZA, M. J., WILKINS, H. R., PRESSON, J. C. & POPPER, A. N. 2002. Age-
and size-related changes in the inner ear and hearing ability of the adult zebrafish
(Danio rerio). *J Assoc Res Otolaryngol,* 3**,** 174-84.
HIRAMATSU, M., SHIOTANI, T., KAMEYAMA, T. & NABESHIMA, T. 1997. Effects of
nefiracetam on amnesia animal models with neuronal dysfunctions. *Behav Brain Res,*
83**,** 107-15.
HUANG, C. S., MA, J. Y., MARSZALEC, W. & NARAHASHI, T. 1996. Effects of the nootropic
drug nefiracetam on the GABAA receptor-channel complex in dorsal root ganglion
neurons. *Neuropharmacology,* 35**,** 1251-61.
KAZMIERCZAK, P., SAKAGUCHI, H., TOKITA, J., WILSON-KUBALEK, E. M., MILLIGAN, R.
A., MULLER, U. & KACHAR, B. 2007. Cadherin 23 and protocadherin 15 interact to form
tip-link filaments in sensory hair cells. *Nature,* 449**,** 87-91.
KIMBERLING, W. J., HILDEBRAND, M. S., SHEARER, A. E., JENSEN, M. L., HALDER, J. A.,
TRZUPEK, K., COHN, E. S., WELEBER, R. G., STONE, E. M. & SMITH, R. J. 2010.
Frequency of Usher syndrome in two pediatric populations: Implications for genetic
screening of deaf and hard of hearing children. *Genet Med,* 12**,** 512-6.
KIMMEL, C. B., PATTERSON, J. & KIMMEL, R. O. 1974. The development and behavioral
characteristics of the startle response in the zebra fish. *Dev Psychobiol,* 7**,** 47-60.
KITANO, Y., MAKINO, M., USUI, C., TAKASUNA, K., KASAI, Y., HIROHASHI, M., TAMURA,
K., TAKAYAMA, S., KOJIMA, H., IIZUKA, H. & ET AL. 1994. General pharmacological
profile of the new cognition-enhancing agent nefiracetam. *Arzneimittelforschung,* 44**,**
199-210.
LEE, H. B., SCHWAB, T. L., KOLEILAT, A., ATA, H., DABY, C. L., CERVERA, R. L.,
MCNULTY, M. S., BOSTWICK, H. S. & CLARK, K. J. 2016. Allele-Specific Quantitative
PCR for Accurate, Rapid, and Cost-Effective Genotyping. *Hum Gene Ther,* 27**,** 425-35.
LENTZ J, K. B. 1999. Usher Syndrome Type I. *GeneReviews®.*
LIN, J. W. & FABER, D. S. 1988. Synaptic transmission mediated by single club endings on the
goldfish Mauthner cell. I. Characteristics of electrotonic and chemical postsynaptic
potentials. *J Neurosci,* 8**,** 1302-12.
LUSCHER, C., NICOLL, R. A., MALENKA, R. C. & MULLER, D. 2000. Synaptic plasticity and
dynamic modulation of the postsynaptic membrane. *Nat Neurosci,* 3**,** 545-50.
LV, C., STEWART, W. J., AKANYETI, O., FREDERICK, C., ZHU, J., SANTOS-SACCHI, J.,
SHEETS, L., LIAO, J. C. & ZENISEK, D. 2016. Synaptic Ribbons Require Ribeye for

Electron Density, Proper Synaptic Localization, and Recruitment of Calcium Channels.
Cell Rep, 15**,** 2784-95.
MAGUPALLI, V. G., SCHWARZ, K., ALPADI, K., NATARAJAN, S., SEIGEL, G. M. & SCHMITZ,
F. 2008. Multiple RIBEYE–RIBEYE Interactions Create a Dynamic Scaffold for the
Formation of Synaptic Ribbons. *The Journal of Neuroscience,* 28**,** 7954-7967.
MARCHI, M., BESANA, E. & RAITERI, M. 1990. Oxiracetam increases the release of
endogenous glutamate from depolarized rat hippocampal slices. *Eur J Pharmacol,* 185**,**
247-9.
MARCOTTI, W. 2012. Functional assembly of mammalian cochlear hair cells. *Exp Physiol,* 97**,**
438-51.
MATUS, A. 1999. Postsynaptic actin and neuronal plasticity. *Curr Opin Neurobiol,* 9**,** 561-5.
MONJARAZ, E., NAVARRETE, A., LÓPEZ-SANTIAGO, L. F., VEGA, A. V. & COTA, G. 2000.
L-type calcium channel activity regulates sodium channel levels in rat pituitary GH3 cells.
The Journal of Physiology, 523**,** 45-55.
MORIGUCHI, S., MARSZALEC, W., ZHAO, X., YEH, J. Z. & NARAHASHI, T. 2003.
Potentiation of N-methyl-D-aspartate-induced currents by the nootropic drug nefiracetam
in rat cortical neurons. *J Pharmacol Exp Ther,* 307**,** 160-7.
MORTON, C. C. & NANCE, W. E. 2006. Newborn hearing screening--a silent revolution. *N Engl*
J Med, 354**,** 2151-64.
MOSER, T., BRANDT, A. & LYSAKOWSKI, A. 2006. Hair cell ribbon synapses. *Cell Tissue*
Res, 326**,** 347-59.
MOSER, T. & VOGL, C. 2016. New insights into cochlear sound encoding. *F1000Res,* 5.
MULLER, D. 1997. Ultrastructural plasticity of excitatory synapses. *Rev Neurosci,* 8**,** 77-93.
NABESHIMA, T., NODA, Y., TOHYAMA, K., ITOH, J. & KAMEYAMA, T. 1990. Effects of DM-
9384 in a model of amnesia based on animals with GABAergic neuronal dysfunctions.
Eur J Pharmacol, 178**,** 143-9.
NARAHASHI, T., MARSZALEC, W., MORIGUCHI, S., YEH, J. Z. & ZHAO, X. 2003. Unique
mechanism of action of Alzheimer's drugs on brain nicotinic acetylcholine receptors and
NMDA receptors. *Life Sci,* 74**,** 281-91.
NICOLSON, T. 2015. Ribbon synapses in zebrafish hair cells. *Hear Res,* 330**,** 170-7.
NICOLSON, T., RUSCH, A., FRIEDRICH, R. W., GRANATO, M., RUPPERSBERG, J. P. &
NUSSLEIN-VOLHARD, C. 1998. Genetic analysis of vertebrate sensory hair cell
mechanosensation: the zebrafish circler mutants. *Neuron,* 20**,** 271-83.
NISHIZAKI, T., MATSUOKA, T., NOMURA, T., SUMIKAWA, K., SHIOTANI, T., WATABE, S. &
YOSHII, M. 1998. Nefiracetam modulates acetylcholine receptor currents via two
different signal transduction pathways. *Mol Pharmacol,* 53**,** 1-5.
NOWYCKY, M. C., FOX, A. P. & TSIEN, R. W. 1985. Three types of neuronal calcium channel
with different calcium agonist sensitivity. *Nature,* 316**,** 440-443.
OBHOLZER, N., WOLFSON, S., TRAPANI, J. G., MO, W., NECHIPORUK, A., BUSCH-
NENTWICH, E., SEILER, C., SIDI, S., SOLLNER, C., DUNCAN, R. N., BOEHLAND, A.
& NICOLSON, T. 2008. Vesicular glutamate transporter 3 is required for synaptic
transmission in zebrafish hair cells. *J Neurosci,* 28**,** 2110-8.
OLIVA, C., ESCOBEDO, P., ASTORGA, C., MOLINA, C. & SIERRALTA, J. 2012. Role of the
MAGUK protein family in synapse formation and function. *Dev Neurobiol,* 72**,** 57-72.
OYAIZU, M. & NARAHASHI, T. 1999. Modulation of the neuronal nicotinic acetylcholine
receptor-channel by the nootropic drug nefiracetam. *Brain Res,* 822**,** 72-9.
PARK, H. W., JUNG, H., CHOI, K. H., BAIK, J. H. & RHIM, H. 2010. Direct interaction and
functional coupling between voltage-gated CaV1.3 Ca2+ channel and GABAB receptor
subunit 2. *FEBS Lett,* 584**,** 3317-22.

PEITSARO, N., KASLIN, J., ANICHTCHIK, O. V. & PANULA, P. 2003. Modulation of the
histaminergic system and behaviour by alpha-fluoromethylhistidine in zebrafish. *J*
Neurochem, 86**,** 432-41.
PEPERMANS, E. & PETIT, C. 2015. The tip-link molecular complex of the auditory mechano-
electrical transduction machinery. *Hear Res,* 330**,** 10-7.
POPPER, A. N. & FAY, R. R. 1993. Sound detection and processing by fish: critical review and
major research questions. *Brain Behav Evol,* 41**,** 14-38.
PREVENTION, C. F. D. C. A. 2016. Summary of 2016 national CDC EHDI data.
PREVENTION, C. F. D. C. A. April 1996. National Occupational Research Agenda.
REGUS-LEIDIG, H., SPECHT, D., TOM DIECK, S. & BRANDSTATTER, J. H. 2010. Stability of
active zone components at the photoreceptor ribbon complex. *Mol Vis,* 16**,** 2690-700.
RENIER, C., FARACO, J. H., BOURGIN, P., MOTLEY, T., BONAVENTURE, P., ROSA, F. &
MIGNOT, E. 2007. Genomic and functional conservation of sedative-hypnotic targets in
the zebrafish. *Pharmacogenet Genomics,* 17**,** 237-53.
RIZZO, M. A., HADJIMICHAEL, O. C., PREININGEROVA, J. & VOLLMER, T. L. 2004.
Prevalence and treatment of spasticity reported by multiple sclerosis patients. *Mult Scler,*
10**,** 589-95.
ROBERT G. ROBINSON , M. D., RICARDO E. JORGE , M. D., KATHLEEN CLARENCE-
SMITH , M. D., PH.D. , & SERGIO STARKSTEIN , M. D. 2009. Double-Blind Treatment
of Apathy in Patients with Poststroke Depression Using Nefiracetam. *The Journal of*
Neuropsychiatry and Clinical Neurosciences, 21**,** 144-151.
SCHMITZ, F., KONIGSTORFER, A. & SUDHOF, T. C. 2000. RIBEYE, a component of synaptic
ribbons: a protein's journey through evolution provides insight into synaptic ribbon
function. *Neuron,* 28**,** 857-72.
SCHRAMM, M., THOMAS, G., TOWART, R. & FRANCKOWIAK, G. 1983. Novel
dihydropyridines with positive inotropic action through activation of Ca2+ channels.
Nature, 303**,** 535-7.
SEILER, C. & NICOLSON, T. 1999. Defective calmodulin-dependent rapid apical endocytosis in
zebrafish sensory hair cell mutants. *J Neurobiol,* 41**,** 424-34.
SHEETS, L. 2017. Excessive activation of ionotropic glutamate receptors induces apoptotic
hair-cell death independent of afferent and efferent innervation. *Scientific Reports,* 7**,**
41102.
SHEETS, L., KINDT, K. S. & NICOLSON, T. 2012. Presynaptic CaV1.3 channels regulate
synaptic ribbon size and are required for synaptic maintenance in sensory hair cells. *J*
Neurosci, 32**,** 17273-86.
SHEETS, L., TRAPANI, J. G., MO, W., OBHOLZER, N. & NICOLSON, T. 2011. Ribeye is
required for presynaptic Ca(V)1.3a channel localization and afferent innervation of
sensory hair cells. *Development,* 138**,** 1309-19.
SHEN, W. & SLAUGHTER, M. M. 1999. Metabotropic GABA receptors facilitate L-type and
inhibit N-type calcium channels in single salamander retinal neurons. *J Physiol,* 516 (Pt
3)**,** 711-8.
SIDI, S., BUSCH-NENTWICH, E., FRIEDRICH, R., SCHOENBERGER, U. & NICOLSON, T.
2004. gemini encodes a zebrafish L-type calcium channel that localizes at sensory hair
cell ribbon synapses. *J Neurosci,* 24**,** 4213-23.
SIEMENS, J., LILLO, C., DUMONT, R. A., REYNOLDS, A., WILLIAMS, D. S., GILLESPIE, P.
G. & MULLER, U. 2004. Cadherin 23 is a component of the tip link in hair-cell stereocilia.
Nature, 428**,** 950-5.
SIMON, P., DUPUIS, R. & COSTENTIN, J. 1994. Thigmotaxis as an index of anxiety in mice.
Influence of dopaminergic transmissions. *Behav Brain Res,* 61**,** 59-64.
STARKSTEIN, S. E., BROCKMAN, S., HATCH, K. K., BRUCE, D. G., ALMEIDA, O. P., DAVIS,
W. A. & ROBINSON, R. G. 2016. A Randomized, Placebo-Controlled, Double-Blind

Efficacy Study of Nefiracetam to Treat Poststroke Apathy. *J Stroke Cerebrovasc Dis,* 25**,**
1119-1127.
SULI, A., PUJOL, R., CUNNINGHAM, D. E., HAILEY, D. W., PRENDERGAST, A., RUBEL, E.
W. & RAIBLE, D. W. 2016. Innervation regulates synaptic ribbons in lateral line
mechanosensory hair cells. *J Cell Sci,* 129**,** 2250-60.
USHER, C. H. 1914. On the inheritance of retinitis pigmentosa, with notes of cases. *Roy. Lond.*
*Ophthal. Hosp. Rep***,** 130-236.
VAN CAMP G, S. R. *Hereditary Hearing Loss Homepage* [Online]. Available:
https://hereditaryhearingloss.org [Accessed].
VEENSTRA-VANDERWEELE, J., COOK, E. H., KING, B. H., ZAREVICS, P., CHERUBINI, M.,
WALTON-BOWEN, K., BEAR, M. F., WANG, P. P. & CARPENTER, R. L. 2017.
Arbaclofen in Children and Adolescents with Autism Spectrum Disorder: A Randomized,
Controlled, Phase 2 Trial. *Neuropsychopharmacology,* 42**,** 1390-1398.
WAN, L., ALMERS, W. & CHEN, W. 2005. Two ribeye genes in teleosts: the role of Ribeye in
ribbon formation and bipolar cell development. *J Neurosci,* 25**,** 941-9.
WATABE, S., YAMAGUCHI, H. & ASHIDA, S. 1993. DM-9384, a new cognition-enhancing
agent, increases the turnover of components of the GABAergic system in the rat
cerebral cortex. *Eur J Pharmacol,* 238**,** 303-9.
WATANABE, Y., LAWLOR, G. F. & FUJIWARA, M. 1998. Role of nerve terminal L-type Ca2+
channel in the brain. *Life Sci,* 62**,** 1671-5.
WEIL, D., KUSSEL, P., BLANCHARD, S., LEVY, G., LEVI-ACOBAS, F., DRIRA, M., AYADI, H.
& PETIT, C. 1997. The autosomal recessive isolated deafness, DFNB2, and the Usher
1B syndrome are allelic defects of the myosin-VIIA gene. *Nat Genet,* 16**,** 191-3.
WESTERFIELD, M. 1995. *The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish*
(Danio rerio), Eugene, OR, University of Oregon Press.
WONG, A. B., RUTHERFORD, M. A., GABRIELAITIS, M., PANGRSIC, T., GOTTFERT, F.,
FRANK, T., MICHANSKI, S., HELL, S., WOLF, F., WICHMANN, C. & MOSER, T. 2014.
Developmental refinement of hair cell synapses tightens the coupling of Ca2+ influx to
exocytosis. *EMBO J,* 33**,** 247-64.
WONG, H.-T. C., ZHANG, Q., BEIRL, A. J., PETRALIA, R. S., WANG, Y.-X. & KINDT, K. S.
2019. Synaptic mitochondria are critical for hair-cell synapse formation and function.
*bioRxiv***,** 671701.
YANG, Q., SUN, P., CHEN, S., LI, H. & CHEN, F. 2017. Behavioral methods for the functional
assessment of hair cells in zebrafish. *Front Med,* 11**,** 178-190.
YAO, Q., DESMIDT, A. A., TEKIN, M., LIU, X. & LU, Z. 2016. Hearing Assessment in Zebrafish
During the First Week Postfertilization. *Zebrafish,* 13**,** 79-86.
YOSHII, M. & WATABE, S. 1994. Enhancement of neuronal calcium channel currents by the
nootropic agent, nefiracetam (DM-9384), in NG108-15 cells. *Brain Res,* 642**,** 123-31.
YOSHII, M., WATABE, S., MURASHIMA, Y. L., NUKADA, T. & SHIOTANI, T. 2000. Cellular
mechanism of action of cognitive enhancers: effects of nefiracetam on neuronal Ca2+
channels. *Alzheimer Dis Assoc Disord,* 14 Suppl 1**,** S95-102.
ZEDDIES, D. G. & FAY, R. R. 2005. Development of the acoustically evoked behavioral
response in zebrafish to pure tones. *J Exp Biol,* 208**,** 1363-72.
ZHAO, X., KURYATOV, A., LINDSTROM, J. M., YEH, J. Z. & NARAHASHI, T. 2001. Nootropic
drug modulation of neuronal nicotinic acetylcholine receptors in rat cortical neurons. *Mol*
Pharmacol, 59**,** 674-83.
ZOTTOLI, S. J. 1977. Correlation of the startle reflex and Mauthner cell auditory responses in
unrestrained goldfish. *J Exp Biol,* 66**,** 243-54.
1072
1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

Fig 1. L-type voltage-gated calcium channel agonists may restore function in hair cell.

(A) In a normal hair cell, sound causes stereocilia to deflect towards the tallest stereocilia and

the mechanotransduction channels (MET) at the top of the stereocilia open in response allowing

cations such as calcium and potassium to flow into the cell. This causes a change in

membrane potential which leads to L-type voltage gated calcium channels opening at

the basolateral sides of the cell. Calcium enters the cell thus increasing intracellular calcium

concentrations. Calcium mediates neurotransmitter release from the synaptic vesicles within the

ribbon synapse into the synaptic cleft thus stimulating afferent neurons. (B) In cells lacking

myo7a expression, proper MET channel gating does not occur and thus the appropriate

membrane potential is not reached to allow L-type voltage gated calcium channel to open and

thus there is insufficient synaptic transmission to the auditory nerve to create meaningful

interactions. We hypothesize that by augmenting the downstream signal in *myo7aa*[-/-] mutant

hair cells, we can reconstitute a new functional response to sound by increasing the

sensitivity of the calcium channel through treatment with L-type voltage-gated calcium channel

agonists (C).

1102

Fig. 2 *myo7aa*^{-/-} larvae at 5 dpf have altered mechanotransduction activity, stereocilia

structure and ribbon synapse structure. (A, B) Representative lateral view of 5 dpf wildtype

larvae (A) and *myo7aa*^{-/-} larvae (B) briefly exposed to FM1-43 obtained using Light Sheet

Fluorescence microscopy. Labeling with FM1-43 indicates functional mechanotransduction.

White arrowhead points to the swimbladder; *myo7aa*^{-/-} larvae do not have an inflated

swimbladder organ. (C, D) Representative images of 5 dpf wildtype stereocilia (C) and *myo7aa*^{-/-}

(D) stained with Alexa Fluor 488 Phalloidin, a high-affinity filamentous actin probe, obtained

using confocal microscopy. White arrow highlights organized and smooth stereocilia in wildtype

hair cells and disorganized and splayed stereocilia in myo7aa^{-/-} hair cells. Note: not all

stereocilia are splayed in *myo7aa*^{-/-} hair cells. (E, F) Representative images of 5dpf wildtype

ribbon synapse structure (E) and myo7aa^{-/-} ribbon synapse (F) obtained with transmission

1114   electron microscopy. White arrow points to ribeye protein and white star indicates a halo of

1115   tethered vesicles. Scale bars: 200 nm. (G) Wildtype ribbon synapses have a comparable ribbon

1116   area to *myo7aa*^{-/-} ribbon synapses (t-test). (H) Wildtype ribbon synapses have an increased

1117   number of tethered vesicles compared to *myo7aa*^{-/-} ribbon synapses (t-test). Black bold line

1118   represents the mean of the data set and error bars are 95% confidence intervals.

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

Fig. 3 *myo7aa*^{-/-} CTBP2 expression in 5 dpf larvae is different from 5 dpf wildtype larvae.

(A) Cartoon depiction of the basal end of a hair cell innervated by the auditory nerve. CTBP2 is

alternatively spliced to produce ribeye protein, which is a component of the synaptic ribbon.

Ribeye has a halo of tethered vesicles which forms the synaptic ribbon. (B, C) Representative

maximum intensity projection (z-stack top-down image) of neuromast from 5dpf wildtype larvae

(B) and *myo7aa*^{-/-} larvae (C) labeled with DAPI labeled nuclei (blue) and Ribeye (red). The

dotted circle outlines one hair cell. Scale bars: 10 µm. (D) 5dpf wildtype neuromasts have a

greater number of active hair cells compared to *myo7aa*^{-/-} neuromasts. An active hair cell is

defined as a hair cell with at least one CTBP2 puncta (t-test). (E) 5dpf wildtype neuromasts

have a greater number of total CTBP2 puncta to *myo7aa*^{-/-} neuromasts (t-test). Black bold line

represents the mean of the data set and error bars are 95% confidence intervals. (F) Further

1147   analysis of the distribution of CTBP2 puncta across a collection of active hair cells reveals that

1148   most 5dpf wildtype active hair cells contain 2 CTBP2 puncta compared to 3 CTBP2 puncta in

1149   *myo7aa*$^{-/-}$ hair cells. All error bars are 95% confidence intervals.

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

Fig. 4 _myo7aa⁻ᐟ⁻_ MAGUK expression in 5 dpf larvae is different from 5 dpf wildtype larvae.

(A) Cartoon depiction of the basal end of a hair cell innervated by the auditory nerve. MAGUK

(membrane associated guanylate kinases) are a kinase family that target and anchor glutamate

receptors to the synaptic terminals on the post-synaptic cell as part of the postsynaptic density

(PSD). (B, C) Representative maximum intensity projection (z-stack top-down image) of

neuromast from 5 dpf wildtype larvae (B) and _myo7aa⁻ᐟ⁻_ larvae (C) labeled with DAPI labeled

nuclei (blue) and MAGUK (green). The dotted circle outlines one hair cell. Scale bars: 10 μm.

(D) 5dpf wildtype neuromasts have a greater number of active post-synaptic cells compared to

myo7aa⁻ᐟ⁻ neuromasts. An active post-synaptic cell is defined as a cell with at least one MAGUK

puncta (t-test). (E) 5dpf wildtype neuromasts have a greater number of total MAGUK puncta to

myo7aa⁻ᐟ⁻ neuromasts (t-test). Black bold line represents the mean of the data set and error bars

1180   are 95% confidence intervals. (F) Further analysis of the distribution of MAGUK puncta across

1181   a collection of active hair cells reveals that most 5 dpf wildtype active hair cells contain either 2

1182   or 3 MAGUK puncta compared to 3 MAGUK puncta in *myo7aa*[-/-] hair cells. All error bars are

1183   95% confidence intervals.

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

Fig. 5 5 dpf *myo7aa*^{-/-} larvae exhibit larger turning angles. Movement tracking of 5dpf

wildtype and *myo7aa*^{-/-} larvae over a 2.5 minute interval with a 5 ms electric stimulus (50 mV)

administered every 20 s. Ctrax software was used for video processing and Matlab 2012b for

video analysis. (A, B) *myo7aa*^{-/-} larvae have larger absolute smooth orientations (turning angles)

compared to wildtype larvae. Individual turning angles from a population of wildtype and

myo7aa^{-/-} larvae were used to construct the lines (t-test). Black bold line represents the mean of

the data set and error bars are 95% confidence intervals.

1201

1202

1203

1204

Fig. 6 *myo7aa*[-/-] **mutant ribbon synapse abnormalities diminished with exposure to (R)-**
Baclofen. (A-E) Representative images of 5dpf wildtype untreated ribbon synapse structure (A)
myo7aa[-/-] untreated ribbon synapse (B) and *myo7aa*[-/-] treated with 5 µM (±)-Bay K 8644 (C),
250 µM Nefiracetam (D) and 125 µM (R)-Baclofen (E) obtained with transmission electron
microscopy. Scale bars: 1000 nm. (F) *myo7aa*[-/-] untreated ribbon synapse area is smaller
compared to myo7aa[-/-] treated with 125 µM (R)-Baclofen. Additionally, wildtype untreated ribbon
synapses have a comparable ribbon area to myo7aa[-/-] treated with 125 µM (R)-Baclofen ribbon
synapses (t-test). (G) *myo7aa*[-/-] untreated ribbons have fewer tethered vesicles compared to
myo7aa[-/-] treated with 125 µM (R)-Baclofen. Wildtype untreated ribbons have a comparable
number of tethered vesicles to *myo7aa*[-/-] treated with 125 µM (R)-Baclofen ribbon synapses (t-
test). This indicates that there is a rescue in the number of tethered vesicles upon treatment

1217   with (R)-Baclofen. Black bold line represents the mean of the data set and error bars are 95%

1218   confidence intervals.

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

Fig 7. Distribution of CTBP2 and MAGUK puncta more closely resembles wildtype

distribution with exposure to L-type voltage-gated calcium channel agonists in 5dpf

***myo7aa*^{-/-} neuromasts.** (A-D, J-N) Representative maximum intensity projection (z-stack top-

down image) of neuromast from 5 dpf wildtype larvae untreated and treated (A-D) and *myo7aa*^{-/-}

larvae untreated and treated (K-N) labeled with DAPI labeled nuclei (blue) and Ribeye (red).

Scale bars: 10 µm. (I) A majority of 5 dpf untreated wildtype active hair cells have 3 CTBP2

puncta. Upon incubation with 5 µM (±)-Bay K 8644 a majority of active hair cells shifted to 2

CTBP2 puncta. There was no change with incubation of either 250 µM Nefiracetam or 125 µM

(R)-Baclofen. (S) A majority of 5dpf untreated *myo7aa*^{-/-} active hair cells have 2 CTBP2 puncta.

Upon incubation with 5 µM (±)-Bay K 8644, 250 µM Nefiracetam or 125 µM (R)-Baclofen a

majority of hair cells have 3 CTBP2 puncta. (E-H, O-R) Representative maximum intensity

projection (z-stack top-down image) of neuromast from 5 dpf wildtype larvae untreated and

treated (E-H) and *myo7aa*^{-/-} larvae untreated and treated (O-R) labeled with DAPI labeled nuclei

(blue) and Ribeye a (red). Scale bars: 10 μm. (J) A majority of 5 dpf untreated wildtype active

hair cells have 2 or 3 MAGUK puncta. Upon incubation with 5 μM (±)-Bay K 8644, 250 μM

Nefiracetam or 125 μM (R)-Baclofen a majority of active hair cells have 2 MAGUK puncta. (T) A

majority of 5 dpf untreated *myo7aa*^{-/-} active hair cells have 3 MAGUK puncta. Upon incubation

with 5 μM (±)-Bay K 8644, 250 μM Nefiracetam or 125 μM (R)-Baclofen a majority of hair cells

shift to 2 MAGUK puncta. All error bars are 95% confidence intervals.

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

Fig. 8 *myo7aa*[-/-] **swimming behavior and acoustic startle response improved with L-type**
voltage-gated calcium channel agonists. Movement tracking of 5 dpf wildtype and *myo7aa*[-/-]
larvae over a 2.5 minute interval with a 5 ms electric stimulus (50 mV) administered every 20 s.
Ctrax software was used for video processing and Matlab 2012b for video analysis. (A, B) 250
µM Nefiracetam and 125 µM (R)-Baclofen did not affect the absolute smooth orientation of
wildtype larvae, however, 5 µM (±)-Bay K 8644 resulted in a decreased absolute smooth
orientation compared to untreated wildtype larvae. *Myo7aa*[-/-] larvae treated with 5 µM (±)-Bay K
8644, 250 µM Nefiracetam and 125 µM (R)-Baclofen individually all gave decreased absolute
smooth orientations, with the most robust response observed in 5 µM (±)-Bay K 8644
incubation. Individual turning angles from a population of wildtype and *myo7aa*[-/-] larvae were
used to construct the lines (t-test). Black bold line represents the mean of the data set and error

bars are 95% confidence intervals. (C) Sample movement traces of wildtype and *myo7aa*[-/-]

larvae in individual treatment groups indicates that in the presence of 5 µM (±)-Bay K 8644, 250

µM Nefiracetam and 125 µM (R)-Baclofen *myo7aa*[-/-] larvae swimming changes to include

smoother trajectories with more zig-zag like swimming and fewer circling episodes and overall

movement traces of wildtype in the 4 treatments is unaffected. Diameter of the wells is 20 mm.

(D, E) Acoustic startle response was captured by administering 3 stimuli at each frequency per

experiment. Videos were scored blindly and the mean number of responses per total stimuli

(mean response rate) was determined. 5dpf untreated wildtype larvae respond most at 200 Hz

and *myo7aa*[-/-] larvae have little to no response at all frequencies. Upon incubation with 5 µM (±)-

Bay K 8644 wildtype acoustic startle decreased at all frequencies, however *myo7aa*[-/-] larvae had

a significant increase with over 20% response rate at 200 and 600 Hz. Incubation in 250 µM

Nefiracetam increased acoustic startle in wildtype and *myo7aa*[-/-] larvae at all frequencies,

although the increase in the *myo7aa*[-/-] larvae is modest. 125 µM (R)-Baclofen decreased

acoustic startle response significantly in wildtype larvae and had little to no effect on acoustic

startle in the *myo7aa*[-/-] larvae. Black bold line represents the mean of the data set and error bars

are mean variance.

1297