Mountain stoneflies may tolerate warming streams: evidence from organismal

physiology and gene expression

Scott Hotaling[1,*], Alisha A. Shah[2,*], Kerry L. McGowan[1], Lusha M. Tronstad[3], J. Joseph

Giersch[4], Debra S. Finn[5], H. Arthur Woods[2], Michael E. Dillon[6,a], and Joanna L. Kelley[1,a]

Affiliations:

[1] School of Biological Sciences, Washington State University, Pullman, WA, USA

[2] Division of Biological Sciences, University of Montana, Missoula, MT, USA

[3] Wyoming Natural Diversity Database, University of Wyoming, Laramie, WY, USA

[4] U.S. Geological Survey, Northern Rocky Mountain Science Center, West Glacier, MT, USA

[5] Department of Biology, Missouri State University, Springfield, MO, USA

[6] Department of Zoology and Physiology, University of Wyoming, Laramie, WY, USA

* *Contributed equally*

[a] *Co-supervised research*

Correspondence:

Scott Hotaling, School of Biological Sciences, Washington State University, Pullman, WA,

99164, USA; Email: scott.hotaling@wsu.edu; Phone: (828) 507-9950; ORCID: 0000-0002-5965-

0986

Alisha A. Shah, Division of Biological Sciences, University of Montana, Missoula, MT, 59812,

USA, Email: alisha.shah@mso.umt.edu; Phone: (512) 694-7532; ORCID: 0000-0002-8454-

7905

Running head: Thermal tolerance of mountain stoneflies

Keywords: critical thermal maximum; RNAseq; Plecoptera; alpine streams; thermal tolerance;

climate change; *Lednia tumana*; endangered species; glacier biology

Abstract:

Rapid glacier recession is altering the physical conditions of headwater streams. Stream

temperatures are predicted to rise and become increasingly variable, potentially putting entire

meltwater-associated biological communities at risk of extinction. Thus, there is a pressing need

to understand how thermal stress affects mountain stream insects, particularly where glaciers

are likely to vanish on contemporary timescales. In this study, we tested the critical thermal

maximum (CT_{MAX}) of stonefly nymphs representing multiple species and a range of thermal

regimes in the high Rocky Mountains, USA. We then collected RNA-sequencing data to assess

how organismal thermal stress translated to the cellular level. Our focal species included the

meltwater stonefly, *Lednia tumana*, which was recently listed under the U.S. Endangered

Species Act due to climate-induced habitat loss. For all study species, critical thermal maxima

($CT_{MAX} > 20°C$) far exceeded the stream temperatures mountain stoneflies experience (< 10°C).

Moreover, while evidence for a cellular stress response was present, we also observed

constitutive expression of genes encoding proteins known to underlie thermal stress (i.e., heat

shock proteins) even at low temperatures that reflected natural conditions. Our results challenge

the prevailing notion that high-elevation aquatic insects are physiologically threatened by

warming temperatures. Rather, we posit that other factors (e.g., competition) may better explain

their extreme distributions.

Introduction:

Predicting how species will respond to climate change is a central goal of contemporary

ecology (Araújo & New, 2007, Urban *et al.*, 2016). This goal is difficult, however, because at a

minimum it requires knowledge of extant distributions, physiological limits, and future conditions

in relevant habitats. Mountain streams around the world are being transformed by climate

change, primarily through rapid recession of glaciers and perennial snowfields (Hotaling *et al.*,

2017). Warmer air temperatures are predicted to cause rapid loss of permanent snow and ice,

drive generally higher, more variable stream temperatures, and eventually lower flows in

meltwater-dominated catchments (Huss & Hock, 2018, Jones *et al.*, 2014). Expected ecological

responses include a reduction of biodiversity in headwater streams across multiple levels of

biological organization and taxonomic groups (Bálint *et al.*, 2011, Finn *et al.*, 2013, Giersch *et*

al., 2017, Hotaling *et al.*, 2019a, Jordan *et al.*, 2016). Considerable attention has been devoted

to potential losses of aquatic insect diversity (e.g., Jacobsen *et al.*, 2012). However, the specific

mechanisms underlying physiological limits in alpine stream insects remain unknown. This

knowledge gap is particularly important in light of the widely held assumption that aquatic

insects living at high-elevations are cold-adapted stenotherms that will not tolerate warming

streams (Giersch *et al.*, 2015, Jacobsen *et al.*, 2012). Recent evidence showing that the thermal

maxima of high-elevation stream taxa can greatly exceed maximum water temperatures (e.g.,

Shah *et al.*, 2017b) and that spring-dwelling cold stenotherms exhibit little variability in heat

shock protein (HSP) expression across temperatures (e.g., Ebner *et al.*, 2019) contradicts this

assumption, raising new questions about whether climate warming directly threatens headwater

biodiversity. To better understand the degree to which headwater species can tolerate warming,

links between relevant traits at the organismal (thermal stress) and cellular (e.g., gene

expression) level are needed.

As ectotherms, insect body temperatures depend strongly on their external environment.

Insects are therefore threatened by rising global temperatures, and recent studies have

documented declines in their diversity (Lister & Garcia, 2018, Sánchez-Bayo & Wyckhuys,

2019). The effects of temperature on ectotherm performance and survival, however, are

complex. Ectotherms may respond to stressful temperatures through plasticity or acclimatization

(Seebacher *et al.*, 2015), the evolution of higher thermal limits (Angilletta Jr *et al.*, 2007), or

behavioral thermoregulation (Kearney *et al.*, 2009). Temperature can also affect organismal

distributions indirectly. For instance, changing temperatures can alter ratios of oxygen supply

and demand (Pörtner *et al.*, 2007, Verberk *et al.*, 2016). Cold habitats can also provide natural

buffering against invasions by competitors or predators (Isaak *et al.*, 2015). Thus, temperature

likely shapes both the evolution of aquatic insect physiology as well as local networks of biotic

interactions. To understand the relationship between temperature and ectotherm tolerance, trait-

based approaches (e.g., testing upper thermal tolerance) can be effective. However, a focus on

physiological traits at the whole-organism level may overlook other key aspects of a species'

potential for response, perhaps limiting predictions of whether species can evolve in response to

changing thermal regimes (Chown *et al.*, 2010) or tolerate them *in situ* via plasticity. Thus, there

is a need to connect traits from cellular to organismal levels and consider findings holistically.

Most aquatic insect life histories begin with an egg state followed by an extended period

of development as nymphs before winged adult emergence. Due to the high heat capacity of

water, stream temperatures are less thermally variable than air. However, a surprising amount

of variation still exists in streams due to many factors, including latitude, elevation, flow, and

canopy cover (Shah *et al.*, 2017b). At high-elevations, an additional factor—the primary source

of water input—plays an outsized role in dictating thermal variation downstream (Hotaling *et al.*,

2017). High-elevation freshwaters are fed by four major hydrological sources: glaciers,

snowfields, groundwater aquifers, and subterranean ice (Hotaling *et al.*, 2019a, Tronstad *et al.*,

2019, Ward, 1994). Glaciers and subterranean ice (e.g., rock glaciers) promote near constant,

extremely cold conditions (i.e., less than 3°C year-round) whereas snowmelt- and groundwater-

fed streams are warmer and often more thermally variable (Hotaling *et al.*, 2019a, Tronstad *et

al.*, 2019). However, these general thermal "rules" only apply in close proximity to a primary

source. Patterns can change dramatically downstream as flows are altered (e.g., pooling into a

high-elevation pond) and sources mix (e.g., a warmer groundwater-fed stream flows into a

glacier-fed stream). With extensive thermal variation over small geographic scales and

abundant, putatively cold-adapted resident invertebrates, high-elevation waters provide an ideal,

natural model for testing hypotheses of physiological limits in a framework relevant to global

change predictions.

In this study, we investigated gene expression as a function of tolerance to heat stress

for stonefly nymphs collected from high-elevation streams in the northern Rocky Mountains. We

focused on three taxa—*Lednia tetonica*, *Lednia tumana*, and *Zapada* sp.—all of which have

habitat distributions closely aligned with cold, meltwater stream conditions. *Lednia tumana* was

recently listed under the U.S. Endangered Species Act due to climate-induced habitat loss (U.S.

Fish & Wildlife Service, 2019). To test thermal tolerance at the organismal level, we measured

the critical thermal maximum (CT_{MAX}), a widely used metric for comparing tolerance to heat

stress among animals (Healy *et al.*, 2018). We specifically addressed three overarching

questions: (1) Does natural thermal variation in stream temperature predict mountain stonefly

CT_{MAX}? (2) Do high-elevation stoneflies mount cellular stress responses when subjected to heat

stress? And, if so, which genes are involved? (3) Is there a link between habitat conditions,

organismal limits, and underlying gene expression? Following Shah *et al.* (2017b), we expected

nymphs from streams with higher maximum temperatures to have correspondingly higher

values of CT_{MAX}. We also expected to observe a signal of cellular stress with genes typical of

heat stress responses (e.g., HSPs) upregulated. Finally, we expected nymphs that experience

higher temperatures in nature to exhibit a correspondingly muted cellular stress response.

Collectively, our study sheds new light on thermal stress in high-elevation stream insects and

contributes new perspective to a pressing challenge for the field: clarifying if species living in

cold headwaters are as sensitive to warming temperatures as their extreme distributions

suggest.

Table 1. Environmental variation, mountain range, and habitat types included in this study.
GNP: Glacier National Park, Montana. GRTE: Teton Range, Wyoming. T_{MAX}: the maximum
temperature observed, T_{RANGE}: the difference between the maximum and minimum
temperatures observed, and T_{MEAN}: the mean temperature observed. All temperature data are in
degrees Celsius. SPC: specific conductivity (μS cm^{-1}), PI: Pfankuch Index, a measure of stream
channel stability (higher values correspond to a less stable streambed). Temperatures were
measured on a representative day in late July 2019 for all sites except Lunch Creek (data from
late July 2014). See Table S1 for specific dates of temperature data collection.
137

Population	Range	Taxa	Type	T_{MAX}	T_{RANGE}	T_{MEAN}	SPC	PI
Lunch Creek	GNP	*L. tumana*	Snowmelt	9.9	5.7	6.2	40.7	25
Wind Cave	GRTE	*Zapada* sp.	Icy seep	3.2	0.5	2.8	101.1	18
Mt. St. John	GRTE	*L. tetonica*	Icy seep	4.6	2.2	3.0	25.0	34
Cloudveil Dome	GRTE	*L. tetonica*	Glacier-fed	2.1	0.3	2.0	4.1	32
Skillet Glacier	GRTE	*L. tetonica*	Glacier-fed	7.1	4.4	4.4	3.1	34
Tetonica Pond[a]	GRTE	*L. tetonica*	Pond	4.9	2.3	3.1	29.3	n/a

138　[a] Named by the authors. Does not reflect official conventions.
139

Materials and Methods:

Specimen collection

During the summer of 2018 (29 July-6 August), we collected late-instar stonefly nymphs

representing at least three species (*Lednia tetonica, Lednia tumana*, and *Zapada* sp.; Family

Nemouridae) from six streams in Glacier National Park (GNP), Montana, and Grand Teton

National Park and the surrounding region (GRTE), Wyoming, USA (Figure 1; Tables 1, S1). We

selected a later summer timepoint because it represents the warmest stream temperatures

nymphs experience before emerging in August. Also, given the acclimation capacity of CT_{MAX} in

temperate aquatic insects (Shah *et al.*, 2017a), we measured CT_{MAX} during this period because

it is also when we expected CT_{MAX} to be greatest. Specimens were collected by turning over

rocks and gently transferring nymphs to a small tray filled with streamwater. Nymphs were

brought to the laboratory in 1 L Whirl-Pak bags (Nasco) filled with streamwater surrounded by

snow or ice. Species were identified based on morphological variation following previous studies

(e.g., Giersch *et al.*, 2017). Unlike *Lednia*, multiple *Zapada* species can be present in the same

stream and previous genetic data has indicated the potential for cryptic diversity in the group

(Hotaling *et al.*, 2019b). Therefore, we cannot exclude the possibility of more than one species

of *Zapada* in the Wind Cave population and thus only identified *Zapada* to genus (Table 1).

157
158
Figure 1. (A) The region of the Rocky Mountains where this study was conducted including (B)
Glacier National Park, MT and (C) Grand Teton National Park, WY and the surrounding region.
(D) A thermograph of hourly temperatures for each study site in late July. Site acronyms (top to
bottom): Lunch Creek (LC), Skillet Glacier (SG), Tetonica Pond (TP), Mt. St. John (MSJ), Wind
Cave (WC), and Cloudveil Dome (CD). A complete 24-hour thermograph is not shown for MSJ
and TP because only 21 and 19 hours of continuous data were recorded for those sites,
respectively. More extensive thermal data are provided in Figure S1.

Environmental data and aquatic habitat classifications

For each study stream, we measured temperature by placing *in situ* HOBO loggers
(Temperature Pro v2, Onset Computer Corporation) that recorded temperature hourly. Lengths
of logger deployments ranged from less than 24 hours (Mt. St. John, Tetonica Pond) to several
days (Cloudveil Dome) or a full year (Lunch Creek, Skillet Glacier, Wind Cave). Using these
data, we constructed a one-day thermograph for each site based on a representative day in late
July (exact dates provided in Table S1) and estimated the highest (T_{MAX}), range (T_{RANGE}), and
mean (T_{MEAN}) temperatures for that day. For two sites with more than one year of temperature
data (Wind Cave: 2016, 2019; Lunch Creek: 2012, 2013, 2014), we compared multiple complete
thermographs for July to ensure that our results were not biased by an unusual year- or day-
specific pattern (Figure S1). We also collected two additional environmental variables to inform
our habitat classifications (see below): specific conductivity (SPC), measured with a YSI
Professional ProPlus multiparameter probe which was calibrated at the trailhead before each
sampling trip, and stream channel stability, calculated via a modified version of the Pfankuch
Index (PI), a standard metric for assessing channel stability in mountain systems that integrates
five key physical characteristics of the stream into a single value (Peckarsky *et al.*, 2014).

We classified sites into habitat types following previous studies (Giersch *et al.*, 2017,
Hotaling *et al.*, 2019a, Tronstad *et al.*, 2019). Briefly, we incorporated a site's primary
hydrological source, environmental variation, and geomorphology, to group them into one of
four habitat types: streams fed by a surface glacier ("glacier-fed"), a perennial snowfield
("snowmelt-fed"), emanating from subterranean ice (e.g., rock glaciers, "icy seep"), or slow-
flowing, alpine ponds ("pond"). We categorized a stream as glacier-fed if it had a named glacier
upstream and an extremely unstable streambed (PI > 30). Any other streams fed by perennial
surface ice and snow were categorized as snowmelt-fed. We classified streams as icy seeps if
we observed evidence of a subterranean ice source (e.g., lobes of a rock glacier), they were
extremely cold (e.g., $T_{MAX} < 5°C$), and had high conductivity (SPC > 50; Hotaling *et al.*, 2019a).
Ponds were identified by their low-angle profile and the presence of standing water.

Table 2. Morphological and physiological data included in this study. Holding: time (hours) that specimens were held at 3°C with no access to food before testing. *N*: sample size for each population. Mean body lengths were used as a proxy for mass and are reported in millimeters with standard errors. RNAseq: sample sizes for RNA sequencing for treatment (T; CT_{MAX}) and control (C; held at 3°C) specimens. Mean CT_{MAX} is given in degrees Celsius.

Population	Taxon	Length	Holding	*N*	Mean CT_{MAX}	RNAseq (*N*)
Lunch Creek	*L. tumana*	4.9 ± 0.5	72	24	28.7	3T / 3C
Wind Cave	*Zapada* sp.	4.4 ± 0.6	48	23	25.9	--
Mt. St. John	*L. tetonica*	5.6 ± 0.7	12	24	26.6	3T / 3C
Cloudveil Dome	*L. tetonica*	4.5 ± 0.5	12	23	26.1	--
Skillet Glacier	*L. tetonica*	5.6 ± 0.4	12	17	28.6	--
Tetonica Pond	*L. tetonica*	4.6 ± 0.6	12	23	28.6	3T / 3C

Measuring critical thermal maxima (CT_{MAX})

Nymphs were brought into the laboratory as quickly as possible (typically less than 12 hours after collection) and transferred to holding chambers in 150-quart coolers filled with water from a nearby stream (Pacific Creek: 43.9036°, -110.5892°). We used aquarium chilling units (1/10 HP, Coralife) to maintain the holding baths at ~3°C (Figure S2). Each holding chamber contained 12 nymphs in a ~2 L plastic container immersed in the bath such that both water and nymphs were isolated from the rest of the system. We included plastic mesh in each chamber to give nymphs substrate to cling to. We maintained high levels of water flow and dissolved oxygen by air stone bubbling in each chamber. Nymphs had no access to food during the holding period to ensure they were tested in a fasting state (i.e., after available food had been digested, absorbed, and cleared from the digestive tract). All nymphs were held in these conditions for at least ~12 hours before testing (Table 2).

We measured CT_{MAX}, a survivable temperature at which nymph locomotor function becomes disorganized. We placed up to 12 nymphs in individual mesh chambers in a water bath held at 3°C. Because warming is the most obvious effect of climate change in high-mountain streams, we chose to only vary temperature, while maintaining natural flow and oxygenation with air pumps. Four thermo-electric cooling (TEC) plates attached to a temperature controller were used to increase temperature at ~0.25°C per minute. We recorded CT_{MAX} when an individual nymph could no longer right itself after being turned onto its back (Videos S1-S2). After a nymph reached its CT_{MAX}, we immediately transferred it to an 8°C bath for recovery and assessed survival by monitoring nymphs until they resumed normal movement. Nymphs were later preserved in ~95% ethanol. We measured body length to the nearest ¼ mm using a dissecting microscope and a millimeter grid attached to the base of the microscope. A

subset of nymphs were flash frozen at either their CT_{MAX} or holding temperature for RNA

sequencing (RNAseq).

For CT_{MAX}, all statistical analyses were conducted in R v3.4.0 (R Core Team, 2013). We

focused our main analysis on *L. tetonica* because we had data for multiple populations. We first

analyzed the effect of body size (length) on CT_{MAX} with a linear model. Because body size was

not a significant predictor of CT_{MAX} (see Results), our final linear models assessing the effect of

maximum stream temperature (T_{MAX}) on CT_{MAX} included only T_{MAX} as the predictor variable. We

also assessed the effect of T_{MAX} on CT_{MAX} in a second, broader analysis which included the two

populations of *L. tumana* and *Zapada* sp. While valuable, this 'all-species' analysis cannot be

used to determine if CT_{MAX} varies among species or if the variation we observed is due to

population-level differences because we lacked replicates for *L. tumana* and *Zapada* sp.

RNA sequencing

During the thermal tolerance experiment, a subset of individuals from three populations

and both *Lednia* species (Lunch Creek, *L. tumana*; Mt. St. John and Tetonica Pond, *L. tetonica*;

Figure 1A, Table 2) were sampled for RNAseq. Nymphs at their CT_{MAX} (treatment) and others

that remained at the holding temperature (control) were flash frozen in liquid nitrogen. We

sampled three treatment and three control nymphs for each population ($N = 18$ total; Table 2).

Samples were stored in liquid nitrogen until they were transferred to a -80° freezer. We

extracted total RNA from entire nymphs following the NucleoSpin RNA (Macherey-Nagel Inc.)

protocol. For extraction, specimens were re-flash frozen with liquid nitrogen in a 1.5 mL

microcentrifuge tube and ground into a fine powder with a sterilized pestle. We quantified RNA

with a Qubit 2.0 fluorometer (Thermo Fisher Scientific) and assessed RNA extraction quality via

fragment analysis with an ABI 3730 DNA Analyzer (Thermo Fisher Scientific).

We prepared RNAseq libraries from 1 μg of total RNA with the NEBNext Poly(A) mRNA

Magnetic Isolation Module (NEB) according to the manufacturer protocol. We targeted a 300-

450 basepair (bp) fragment size distribution. For cDNA amplification, fifteen PCR cycles were

used for all libraries. Presence of a PCR product was visually assessed using an eGel (Thermo

Fisher Scientific). Final libraries were quantified with a Qubit 2.0 fluorometer and further

assessed for quality, amount of cDNA, and fragment size distribution using a 2100 BioAnalyzer

with the High Sensitivity DNA Analysis kit (Agilent). Libraries were then pooled in equal

nanomolar concentrations and sequenced on one lane of an Illumina HiSeq4000 with 100 bp

paired-end chemistry by the Roy J. Carver Biotechnology Center at the University of Illinois

Urbana-Champaign.

Gene expression analyses and protein annotation

We assessed raw sequence data quality with fastQC v0.11.4 (Andrews, 2010) and

visualized a combined output for all libraries with MultiQC v1.5 (Ewels *et al.*, 2016). Next, we

trimmed reads in three successive rounds, all with Trim Galore! v0.4.1 (Krueger, 2015) and

default settings except as noted. First, we removed adapter sequences (--illumina --stringency

6). Next, we trimmed for quality and poly-A tails (--quality 20 --stringency 6 --adapter A{30} --

adapter2 A{30}). We then trimmed for poly-T tails and discarded reads that had become too

short (--stringency 6 --length 50 --adapter T{30} --adapter2 T{30}). We then assessed the quality

of the trimmed reads with fastQC v0.11.4. We randomly subsampled one library (Library 3;

Control, Mt. St. John) to 80% of its original amount because its sequencing depth was much

higher than the rest of the data set. For this, we used the reformat function of BBTools v37.80

(Bushnell, 2014). We removed one library (Library 9; Control, Mt. St. John) from all downstream

analyses as it had just 2.6 million reads, far fewer than any other library (see Results).

We mapped reads to the *L. tumana* reference genome (GenBank #QKMV00000000.1)

with the mitochondrial genome (GenBank #MH374046; Hotaling *et al.*, 2019c) appended to it.

We used HiSat2 v2.1.0 (Pertea *et al.*, 2015) with default settings, first building an index of the

reference with the hisat2-build command. To ensure no bias was introduced by differential

mapping rates between *L. tumana* and *L. tetonica* samples to the *L. tumana* reference genome,

we compared the mean mapping rates for both species with an unpaired *t*-test. Because HiSat2

outputs unsorted SAM files, we converted the output to sorted BAM files with samtools v1.7 (Li

et al., 2009).

We generated a gene count matrix for each library with StringTie v1.3.5 (Pertea *et al.*,

2015). We first ran StringTie with the default settings to assemble alignments into potential

transcripts without a reference annotation (-G) because none is available for *L. tumana.* Next,

we used the --merge utility to combine library-specific sets of transcripts into a merged,

putatively non-redundant set of isoforms. This tool outputs a merged Gene Transfer Format

(GTF) file. We then re-ran StringTie using the merged GTF (-G) and the flags -B and -e to

enable the output of Ballgown GTF files for the global set of transcripts shared by all samples.

Next, we ran the prepDE.py script, also part of the StringTie package, to generate counts

matrices for all genes and transcripts identified in the previous steps.

We performed differential expression analyses using edgeR v3.26.8 (Robinson *et al.*,

2010) in R version 3.5.2 (R Core Team, 2013). We filtered our data set by requiring transcripts

to have more than five total reads and to be present in at least two samples. To visually

compare expression variation across groups of interest (i.e., treatments, species, and

populations), we used the plotPCA function. After filtering, we identified structure in global gene

expression that could not be explained by sample preparation, library size, species, population,

or treatment (Figure S3). We removed this unwanted variation with RUVseq v1.18.0 (Risso *et*

al., 2014). Specifically, we used the "*in silico* empirical" functionality of RUVg where a set of the

least differentially expressed genes (DEGs) are identified and used as controls to globally

normalize variation in the data set. We used the default trimmed mean of M-values (TMM)

method to normalize the data and calculate effective library sizes (Figure S4). Dispersions were

estimated using a generalized linear model and a Cox-Reid profile-adjusted likelihood

(McCarthy *et al.*, 2012). We identified DEGs with quasi-likelihood F-tests (Lun *et al.*, 2016)

which were run using contrasts. We performed DEG identification across three levels of

comparison: (1) Within-populations between treatment (collected at their CT_{MAX}) and control

(held at 3°C) specimens. (2) Between treatment and control for *L. tetonica* specimens only (Mt.

St. John and Tetonica Pond). (3) Between treatment and control for all specimens. A false

discovery rate (FDR) ≤ 0.05 was used to identify DEGs.

To annotate our data set, we extracted the longest isoform for each gene using the

CGAT toolkit and the 'gtf2gtf' function (Sims *et al.*, 2014). We then extracted genes from the file

containing the longest isoforms with gffread v.0.9.9 (Trapnell *et al.*, 2012). We performed a

blastx search for each gene (E-value: 0.001) against the manually curated and reviewed Swiss-

Prot database (Boeckmann *et al.*, 2003; accessed 1 March 2019). Using the results of our

blastx search, we annotated genes, retrieved gene ontology (GO) terms, and mapped GO terms

using Blast2GO v5.2 (Conesa *et al.*, 2005). We annotated DEGs with the top BLAST hit per

transcript. For DEGs without a match in the Swiss-Prot database, we performed two additional

searches using online tools: (1) a batch search against the RFAM database (Kalvari *et al.*, 2017;

http://rfam.org/search) and (2) a manual blast search (E-value: 0.001) against the automatically

annotated and unreviewed TrEMBL database (Boeckmann *et al.*, 2003; http://uniprot.org/blast/).

In Blast2GO v5.2, we performed GO term enrichment analyses using the results of our

blastx/Swiss-Prot annotations on two test sets with one-tailed Fisher's Exact Tests and FDR ≤

0.05 after correcting for multiple tests: (1) upregulated genes for *L. tetonica* only and (2)

downregulated genes for *L. tetonica* only. We did not perform GO term enrichment analysis for

L. tumana because no DEGs were identified for the representative population we examined

(Lunch Creek; see Results). We also did not perform GO term enrichment on the overall *Lednia*

data set because of redundancy with the *L. tetonica* analysis (i.e., roughly two-thirds of the

same individuals would be included). For enrichment analyses, the complete set of transcripts
with BLAST hits were used as the reference set.

To test if stoneflies from more thermally variable environments have muted cellular
responses to stress, we identified all genes annotated as heat shock proteins based on BLAST
hit descriptions. Next, we sorted these genes by their overall expression [$\log_2$ counts per million
(logCPM)] and filtered them to a final set using two criteria: (1) We only included genes
expressed at moderate to high levels (≥ 4 logCPM) and (2) only retained the most expressed hit
(highest mean logCPM) for each unique gene. We did this to prevent any potential bias due to
one gene being represented by multiple hits (see Results). Next, we calculated the mean
difference in logCPM between treatment and control nymphs for each gene and population.
Because the data were not normally distributed (P, Shapiro-Wilk < 0.001), we compared the
distributions of mean differences for each population using a Kruskal-Wallis rank sum test
followed by a Dunn test for multiple comparisons. All scripts and commands used in this study
are available on GitHub (https://github.com/scotthotaling/Lednia_RNAseq).

Results:
Environmental data and species collection
According to the environment criteria described above, we identified one snowmelt-fed
stream (Lunch Creek: GNP), two icy seeps (Wind Cave, Mt. St. John; GRTE), two glacier-fed
streams (Cloudveil Dome, Skillet Glacier; GRTE), and one alpine pond (Tetonica Pond; GRTE;
Table 1). We collected *L. tumana* from Lunch Creek, *Zapada* sp. from Wind Cave, and *L.*
tetonica from the other four sites (Figure 1, Table 1). Lunch Creek was the warmest (T_{MEAN} =
6.2°C; T_{MAX} = 9.9°C) and most thermally variable site (T_{RANGE} = 5.7°C; Table 1). Cloudveil Dome
(T_{MAX} = 2.1°C) and Wind Cave (T_{MAX} = 3.2°C) were the coldest and least variable sites (T_{RANGE} ≤
0.5°C; Table 1). Icy seeps were the coldest and least thermally variable habitat type overall
(T_{MAX}, icy seeps = 3.9°; T_{RANGE}, icy seeps = 1.4°C). For the two sites with two or more years of
available temperature data (2 years, Wind Cave; 3 years, Lunch Creek), thermal differences
across years were negligible (Figure S1).

353
354
Figure 2: (A) The effect of maximum stream temperature (T_{MAX}) on critical thermal maximum
(CT_{MAX}) for each nymph (smaller, lighter circles) with means for each population also shown
(darker, outlined circles). Asterisks mark the species with only a single population sample
(*Zapada* sp. and *L. tumana*). Trendlines indicate significant relationships between T_{MAX} and
CT_{MAX} for two separate linear models for *L. tetonica* only (blue line) and for all species (black
line). In both analyses, stoneflies from colder streams had lower CT_{MAX} values than those from
warmer streams. (B) Box plots showing variation in CT_{MAX} across species. Black horizontal lines
in each box indicate the median with lower and upper bounds of the box representing the lower
and upper quartiles of the data, respectively. Whiskers show the maximum and minimum
values.
365

Thermal physiology

We confirmed that all nymphs survived the CT_{MAX} treatment (except for those that were

immediately flash frozen for RNAseq and could not be assessed). There was no effect of body

size on CT_{MAX} in the *L. tetonica* ($P = 0.58$) nor all-species analysis ($P = 0.28$; Figure S5). We

therefore did not include body size as a covariate in our statistical models. We observed

differences in CT_{MAX} among populations of *L. tetonica* (Figure 2A). Stoneflies inhabiting colder

sites exhibited lower CT_{MAX} values compared to those from warmer sites ($F_{1,85} = 26.108$, $P <$

0.001). We observed the lowest CT_{MAX} for *L. tetonica* in Cloudveil Dome ($T_{MAX} = 2.1°C$; $CT_{MAX} =$

26.1°C), and the highest for *L. tetonica* in Tetonica Pond ($T_{MAX} = 5°C$; $CT_{MAX} = 29.2°C$). We also

found a significant positive relationship between T_{MAX} and CT_{MAX} in the all-species analysis

which included *L. tumana* and *Zapada* sp. ($F_{1,132} = 39.054$, $P < 0.001$). Although we could not

statistically test differences in CT$_{MAX}$ among species due to a lack of replicate *L. tumana* and

Zapada sp. populations, our results indicate that CT$_{MAX}$ may be highest for *L. tumana* (Figure

2B). However, this finding may simply be reflective of the only *L. tumana* population sampled

also being from Lunch Creek, the warmest site included in this study.

RNA sequencing and annotation

We generated 368.8 million read pairs for 18 libraries with a mean per sample of 20.6

million $\pm$ 1.9 million (min. = 2.6 million, max. = 39.2 million). After filtering, subsampling of the

library with the most reads, and dropping the library with the fewest reads, we retained 354.1

million read pairs. On average, 85.2% of reads mapped to the *L. tumana* reference genome with

L. tumana libraries mapping at a slightly higher rate (mean 89.0% $\pm$ 0.5%; min. = 87.8%, max. =

91.0%) than *L. tetonica* (mean = 83.2% $\pm$ 0.6%; min. = 81.0%, max. = 84.5%; *P, t*-test <

0.0001). However, this difference in mapping rate did not extend to a difference in total reads

mapped (mean, *L. tumana* = 19.2 million, mean *L. tetonica* = 21.7 million; *P, t*-test = 0.42). Raw

reads for this study are deposited on the NCBI SRA under BioProject PRJNA587097.

393
394

Figure 3. Global differences in gene expression for stonefly nymphs color-coded by treatment (red, CT$_{MAX}$) or control (blue, held at 3°C) and grouped by species (colored polygons) and populations (shapes).

Differential expression

After filtering and processing of the data set, our gene counts matrix contained 52,954 unique entries. We observed global differences in gene expression between *L. tumana* and *L. tetonica* (Figure 3). When *L. tumana* and *L. tetonica* were combined ("*Lednia*"), 80 genes were differentially expressed: 65 upregulated and 15 downregulated in the treatment (CT$_{MAX}$) versus control group (FDR ≤ 0.05). When only *L. tetonica* populations were considered ("*Tetonica*"), 71 genes were differentially expressed: 60 upregulated, 11 downregulated. Thirty-four DEGs were shared between groups (32 upregulated, two downregulated). When each population was considered alone, no DEGs were identified (including for Lunch Creek, the only *L. tumana* population). While we report results for the *Lednia* and *Tetonica* data sets above, we focus hereafter on *Tetonica* because it contains the most statistical power (two populations) with no potential for species-specific bias. Furthermore, due to the fragmented nature of the *L. tumana* genome (contig N50: 4.7 kilobases (kb); 74,445 contigs > 1 kb; Hotaling *et al.*, 2019c), portions of the same gene were likely present on different contigs in the reference. When we assembled transcripts, this manifested as unique transcripts annotated to the same gene. Thus, in many instances (e.g., hexamerins, *HEXA*; Figures 4, S6), we recovered multiple independent hits to the same gene. While multiple hits may reflect biological reality (e.g., more than one copy of a gene in the genome perhaps reflecting a gene family expansion) we cannot draw such a conclusion. We specify how multiple hits to the same gene were handled where appropriate.

For *Tetonica*, 46 DEGs (64.8%) had BLAST hits to the high-quality, manually curated Swiss-Prot database, 32 of which were unique (Table S2). Of the remainder, three (4.2%) had hits to the RFAM database and nine (12.7%) had hits to the TrEMBL database. Because the TrEMBL protein database is unreviewed, we only refer to Swiss-Prot/RFAM annotations unless specifically noted. The most upregulated gene [MSTRG.32248; log$_2$ fold change (logFC) = 15.6; FDR = 0.015] had no annotation to any database (Figure 4). However, the next four most-upregulated genes (logFC = 7-9.1; Figure 4) included *ABCA3*, which binds ATP, a nucleolysin (*TIAL1*)*,* and two heat shock proteins *HSP70B2* and *HSP70A1*. The two heat shock proteins were also the most expressed DEGs (logCPM = 8.9 and 9.3, respectively) after three genes which were all annotated as hexamerins (*HEXA*; logCPM = 9.3-10). Fourteen DEGs had hits to the same apolipophorin gene, *APLP*, with relatively similar changes in expression (logFC, *APLP* = 2.1-3.8; Figure S6) and overall expression levels (logCPM, *APLP* = 2.2-6.9). The three most

downregulated genes did not have BLAST hits to the Swiss-Prot or RFAM databases [logFC = -

6.6 to -13.7; Figure 4], however two of them (MSTRG.1867 and MSTRG.3534) had TrEMBL hits

though they were not informative in terms of predicted function (Table S2).

Forty-one GO terms were enriched in the upregulated *Tetonica* data set (Figure S7): 26

were classified as being part of a biological process ontology, three were cellular component

related, and 11 were linked to molecular function. The top four most significantly enriched GO

terms were all lipid-related, including their transport, binding, and localization. Eight of the

enriched GO terms (19.5% overall) were associated with protein folding, and three were linked

to chaperone proteins which are commonly associated with physiological stress (Beissinger &

Buchner, 1998)(Beissinger & Buchner, 1997). In the same vein, one enriched GO term – "heat

shock protein binding" (GO:0031072; FDR = 0.015) – clearly reflected a link to heat stress at the

cellular level. No GO terms were enriched for downregulated *Tetonica* DEGs.

Environmental variability and gene expression

Across all populations and species, 38 genes were annotated as heat shock proteins

(HSPs). Of these, 12 unique genes were expressed at moderate to high levels (logCPM ≥ 4;

Figure S8). We found no support for our hypothesis that stoneflies naturally experiencing higher

(and more variable) temperatures exhibit muted cellular stress responses versus those

inhabiting colder (and more thermally stable) streams (Figure 5; *P*, Dunn's ≥ 0.66).

449
450
451 **Figure 4.** Log fold change of *Lednia tetonica* DEGs (white = BLAST annotated; black = no hit;
452 FDR ≤ 0.05). For annotated genes, only the hits to the Swiss-Prot or RFAM databases with the
453 lowest FDRs are included. The full version of this figure, including any instances of multiple hits
454 to the same protein, is provided in Figure S6. Complete information for each annotation is
455 provided in Table S2.
456

457
458
459   **Figure 5.** (A) Mean population-level differences in expression between treatment and control
460   specimens for the 12 most highly expressed, unique HSPs annotated in this study. (B)
461   Distributions of the values in (A) grouped by population. No significant differences were present
462   (P, Dunn < 0.05).
463

Discussion:

As climate change proceeds, headwaters will be dramatically altered by the reduction or

loss of meltwater from glaciers and perennial snowfields (Hotaling *et al.*, 2017). However, the

physiological limits of aquatic, high-elevation species, a group of organisms presumed to be

acutely imperiled by climate change, remain largely unknown (but see Shah *et al.*, 2017b). In

this study, we explored the thermal physiology of high-elevation stoneflies inhabiting the

meltwater of rapidly fading glaciers and snowfields in the Rocky Mountains. Our focal species

are representative of an entire community that may be at risk of climate-induced extirpation

(Giersch *et al.*, 2017, Hotaling *et al.*, 2019a, Tronstad *et al.*, 2019), and included *L. tumana*, a

species listed under the U.S. Endangered Species Act due to climate-induced habitat loss (U.S.

Fish & Wildlife Service, 2019). We show that habitat conditions, specifically maximum

temperatures, predict upper thermal limits and that nymphs mount a cellular stress response

when faced with heat stress. Contrary to our expectations, however, we saw no link between

the scale of the stress response and natural conditions nymphs experience. That is, stoneflies

from warmer environments did not exhibit a muted cellular stress response across HSPs versus

those from cooler streams. Our results shed new light on thermal tolerance of mountain

stoneflies and complement recent cellular perspectives on aquatic insect thermal biology (Ebner

et al., 2019, Gamboa *et al.*, 2017). Broadly, our findings and those of others (e.g., Ebner *et al.*,

2019, Shah *et al.*, 2017b, Treanor *et al.*, 2013), challenge the prevailing notion that aquatic

insect larvae living in extremely cold mountain streams cannot survive warming. For *Lednia,*

with the ability to tolerate short-term temperatures that likely exceed anything they naturally

experience by more than ~10°C, we hypothesize that their headwater distributions are actually a

product of other mechanisms (e.g., species interactions at lower elevation) paired with a unique

capacity to develop at near freezing temperatures.

Thermal tolerance

In mountain systems, thermal tolerance is important to organismal distributions.

Previously, thermal tolerance has been shown to explain the elevational limits of many

terrestrial taxa (Andrews, 1998, Brattstrom, 1968, Feder & Lynch, 1982, Huey & Webster,

1976), but whether it also explains limits of aquatic taxa is unknown. Our study shows that

species of high-elevation stoneflies in the Rocky Mountains, often described as cold

stenotherms that are highly susceptible to warming (e.g., Giersch *et al.*, 2017), can withstand

short-term temperatures much higher than those they experience in nature (see also Shah *et

al.*, 2017b). While the utility of CT_{MAX} has been challenged due to its sensitivity to ramping rates,

as well as acclimation and starting temperatures (Rezende *et al.*, 2011, Terblanche *et al.*,

2011), recent arguments in favor of its ecological relevance have also been made (Jørgensen *et

al.*, 2019). We contend that CT_{MAX} may be uniquely appropriate for mountain stream taxa. Many

high-elevation streams naturally experience rapid increases in temperature throughout the day

(e.g., Lunch Creek, Figure 1D) and with summer streamflows predicted to be reduced under

climate change (Huss & Hock, 2018), baseline temperatures and intraday temperature spikes

should both increase as meltwater volume declines and its buffering capacity is lost.

Simultaneous increases in temperature and reductions in flow may also result in reduced heat

tolerance in stoneflies because less oxygen can dissolve in warmer, slow-moving water, which

will in turn not meet the rising oxygen demands induced by elevated metabolic rates (Pörtner &

Knust, 2007). Evidence for this phenomenon, however, is mixed (Verberk *et al.*, 2016). It is

therefore imperative for future research to address effects of oxygen limitation on heat

tolerance, especially in high-elevation aquatic insects that will experience a potentially lethal

combination of increased temperatures and decreased oxygen (Jacobsen, 2020).

We also observed population-level CT_{MAX} variation within *L. tetonica*, suggesting that

local thermal regime is more important to thermal tolerance than regional thermal regime, and

echoing the findings of other recent studies (Gutiérrez‑Pesquera *et al.*, 2016, Shah *et al.*,

2017b). This effect of local conditions on thermal tolerance might outweigh differences that arise

from evolutionary history because all species (e.g., *Lednia tetonica* and *Zapada* sp.) from cooler

streams had lower CT_{MAX} than those from warmer streams (Figure 2). While we cannot

determine if thermal variation among populations represents evolved differences, all specimens

were held in a common thermal regime for at least 12 hours to limit the effects of previous

thermal conditions on CT_{MAX} estimates. Regardless of the mechanism, the high-elevation

stoneflies included in this study appear poised to cope with short-term warming in streams,

although some populations are likely to be more resilient than others (e.g., those experiencing

higher present-day maximum temperatures) and some life stages may be more sensitive to

warming.

Indeed, we only focused on larvae in our study, and therefore cannot discern whether

other life stages (e.g., eggs or adults) differ in their thermal tolerance. However, we chose to

focus on nymphs for three reasons. First, the larval stage is the key developmental period for

aquatic insects when the vast majority of growth and other fitness-related processes (e.g., egg

production) are occurring, and recent modeling has shown that predicted impacts of climate

change impacts on species can be greatly *underestimated* when the larval stage is overlooked

(Levy *et al.*, 2015). Second, aquatic insect larvae typically do most of their growing during

summer months, when the threat of heat stress is greatest. And, third, egg hatching success in

the laboratory was recently shown to be consistently high for mountain stream insects,

regardless of rearing temperature, including one treatment (12°C) that exceeded the highest

temperatures the focal midges naturally experienced (Schütz & Füreder, 2019). Still, future

experiments that link traits (e.g., thermal tolerance) to cellular processes across the aquatic

insect life cycle will greatly improve our understanding of how sensitivity at key life stages may

influence long-term viability of populations.

Gene expression

High-elevation stoneflies residing in cold meltwater-fed streams exhibited a cellular

stress response when faced with temperatures at their CT_{MAX}. The bulk of this response was

comprised of upregulated genes and included well-known stress response genes (e.g., HSPs;

Lindquist & Craig, 1988), lesser known but potentially stress-related genes in insects (e.g.,

APLP, Dassati *et al.*, 2014), and many DEGs that could not be annotated (Figure 4). Three

HSPs (*HSP70B2*, *HSP70A1*, *HSC70-5*) were upregulated in nymphs experiencing thermal

stress. With well-established roles as cellular protectants, preventing protein denaturation,

binding aberrant proteins, and many other stress-induced measures, the upregulation of HSPs

was unsurprising (King & MacRae, 2015). However, given the seemingly psychrophilic lifestyle

of *Lednia*, where individuals develop at temperatures near 0°C, we expected to see widespread

upregulation of HSPs in treatment nymphs. This was not the case. Rather, *Lednia* appeared to

constitutively express many HSPs even at low temperature (Figure S8). This suggests that,

contrary to the prevailing view, exposure to low temperatures may actually stress *Lednia* (see

additional discussion below). Similar patterns of constitutive HSP expression have been

observed in other cold-tolerant species. For instance, larval caddisflies (Ebner *et al.*, 2019),

polar fish (Buckley *et al.*, 2004), and Antarctic grass (Reyes *et al.*, 2003) constitutively express

many HSPs, presumably to chaperone proteins at low temperature. The potential for *Lednia* to

be stressed by cold temperatures is further supported by the inability of *L. tumana* nymphs to

tolerate being enclosed in ice (Hotaling *et al.*, In review).

We also observed upregulation of genes with lipid-related functions. Lipids, particularly

those in membranes, are extremely sensitive to changes in temperature (Hazel, 1995), and

because of their role in key biological processes (e.g., solute diffusion), are important for

thermal acclimation and adaptation (Muir *et al.*, 2016, Pernet *et al.*, 2007). Many ectotherms

remodel their membrane lipids under thermal stress to maintain fluidity, typically through

increases in saturated fatty acids at higher temperatures (Muir *et al.*, 2016). The process of

maintaining membrane fluidity in the face of changing temperatures has been termed

homeoviscous adaptation (HVA; Sinensky, 1974). The degree of saturation in cuticular lipids of

stoneflies varies across life stages, and was speculated to be related to thermal tolerance,

particularly as it relates to the aquatic versus terrestrial environment (Armold *et al.*, 1969). To

our knowledge, the presence of HVA has not been explicitly tested for in any aquatic insect.

However, before upregulation of lipid-related genes can be presumed to underlie HVA or similar

functional changes in high-elevation stoneflies, alternate explanations about the role of lipids in

development must be considered (see below).

While heat stress is presumed to drive the expression patterns we observed, aquatic

insects accelerate their development and emerge earlier at warmer temperatures (Nebeker,

1971, Rempel & Carter, 1987), sometimes even during CT_{MAX} experiments (A.A.S., personal

observation). Thus, some expression changes may be the result of developmental shifts rather

than thermal stress directly. Indeed, when exposed to long-term temperatures above those they

naturally experience (e.g., $\geq$ 15°C for ~1 month), *L. tumana* nymphs rapidly develop compared

to those held at colder temperatures. However, rapidly emerging adults can get stuck while

shedding their cuticle and die in the process (S.H. and A.A.S., unpublished data). Some of our

results appear more reflective of this developmental shift than heat stress directly. For instance,

it has been suggested that *ABCA3* is upregulated during insect wing development (Broehan *et*

al., 2013). In our study, high temperatures induced upregulation of *ABCA3*, perhaps indicating

accelerated wing development in preparation for emergence as winged adults. Lipid content in

aquatic insects also varies seasonally and tends to peak before metamorphosis, an

energetically demanding activity (Cavaletto & Gardner, 1999). In the caddisfly, *Clisoronia*

magnifica, roughly 80% of the lipid reserves needed for metamorphosis occurred during the last

larval instar (Cargill *et al.*, 1985), which was the same developmental timepoint of the stoneflies

included in this study.

The upregulation of *HEXA* raises similar, albeit more complex, questions. Stoneflies

possess two types of hexameric proteins in their hemolymph: hemocyanin (*HCD*), an oxygen-

carrying protein, and hexamerins, multi-functional proteins that likely evolved from hemocyanin

(Amore *et al.*, 2011, Hagner-Holler *et al.*, 2007). We saw some evidence for the upregulation of

HCD in heat-stressed stoneflies (Figure S9), perhaps reflecting the physiological challenge of

extracting the necessary oxygen from warmer water. However, while hexamerins likely evolved

from *HCD*, their function shifted to storage proteins after they lost the ability to bind oxygen

(Burmester, 2015, Markl & Winter, 1989). Hexamerins primarily act as sources of amino acids

during non-feeding periods (e.g., emergence, Haunerland, 1996) but may also play a role in

cuticle formation (Burmester, 2015, Hagner-Holler *et al.*, 2007), a key stage in aquatic insect

emergence. Thus, the upregulation of *HEXA* may be another cellular indicator of accelerated

emergence to escape injurious conditions.

Mountain stream insects as cold stenotherms: reconsidering a historical paradigm

Aquatic insects living in chronically cold habitats have long been assumed to be cold-

adapted and therefore intolerant of warming (e.g., Giersch *et al.*, 2017, Jacobsen *et al.*, 2012).

This assumption has rarely, if ever, been supported by direct measurements. A potential

mismatch between theory and data is particularly important for imperiled species of

conservation concern. *Lednia tumana* is federally endangered under the U.S. Endangered

Species Act due to loss of cold, meltwater habitat (U.S. Fish & Wildlife Service, 2019). As

glaciers disappear around the world (Huss & Hock, 2018), the demise of *Lednia* and similar

species (e.g., *Zapada* sp.) is presumed to be merely a matter of time (Giersch *et al.*, 2017).

While this may be true, alternative hypotheses or threats beyond temperature should be

considered. Chief among these is the question of niche breadth. Factors limiting niche breadth

are diverse and may not be directly linked to temperature (e.g., interspecific competition or food

availability, Connell, 1961, Roughgarden, 1974), although thermal sensitivity can certainly play a

major, interactive role (Gilchrist, 1995). While terrestrial habitats exhibit a wide array of thermal

variation, potentially allowing more thermal space for species with similar ecologies to exist in

sympatry, the buffering capacity of flowing water may reduce the diversity of thermal niches in

streams across similar spatial extents. Thus, with relatively high short-term thermal tolerance,

far exceeding temperatures experienced in nature, and cellular signatures of stress even at low

temperatures (e.g., constitutive expression of HSPs at 3°C), we hypothesize that the distribution

of *Lednia* and similar species reflects not a requirement for cold conditions but simply a greater

tolerance for them versus other species. Rather than being an extreme thermal specialist,

Lednia may have evolved a wide thermal niche allowing it to colonize environments free of

limiting biological factors. Our hypothesis aligns with previous experimental evidence

highlighting the potential for biotic factors beyond temperature to alter alpine stream ecosystems

(Khamis *et al.*, 2015).

When considering climate change impacts on mountain stream biodiversity, it is

important to distinguish between a species being imperiled by rising temperatures or biotic

factors. At present, the prevailing theory is that a warmer water community will shift uphill and

displace coldwater taxa as glaciers and perennial snowfields are lost (Hotaling *et al.*, 2017).

This theory assumes that coldwater species (e.g., *Lednia*) will not be able to tolerate warmer

conditions and will be extirpated while lower elevation species simultaneously track their

preferred thermal conditions upstream. However, if existing headwater communities can tolerate

warmer conditions and their lower limits are set by other factors (e.g., predation), then climate

change risks for mountain stream communities may be far less generalizable than currently

assumed. For instance, if competition at lower elevations is limiting *Lednia* distributions then

warming temperatures do not guarantee simplistic, binary outcomes of predicted presence or

absence. Rather, the future of *Lednia* and similar taxa may depend upon additional factors

including whether or not their competitors shift uphill (rather than tolerate warmer conditions *in*

situ), how resources may change, and others that are difficult to predict.

Conclusion:

High-elevation stoneflies in the Rocky Mountains can tolerate short-term temperatures

well beyond those they experience in the wild. When challenged with high temperatures,

nymphs mount a cellular stress response that includes upregulation of classic stress response

genes (e.g., HSPs) as well as genes that may be involved in developmental transitions from

aquatic to terrestrial life stages. Aquatic insects are known to develop more rapidly at warmer

temperatures (Nebeker, 1971). Our own laboratory tests of *L. tumana*, however, support a

thermal limit of ~15°C where mortality during emergence greatly increases (S.H. and A.A.S.,

unpublished data). Thus, the potential for warming, even in the short-term, to accelerate

development to the point of lethality warrants further investigation. However, in light of our

results and similar studies (Ebner *et al.*, 2019, Shah *et al.*, 2017b), we challenge the premise

that the distribution of mountain stream insects in cold, thermally stable habitats indicates

specialized preferences for cold, or evolved physiologies that are only viable in the cold. Rather,

the appearance of constitutive expression of many HSPs in *Lednia* as well as the inability of *L.*

tumana nymphs to survive ice enclosure (Hotaling *et al.*, In review) suggest that the

contemporary thermal regimes they experience may actually be injurious. Ultimately, if imperiled

species like *Lednia* are not directly threatened by warming temperatures in the near-term, then

there is clear reason for greater optimism about their future. However, explicit investigations of

their development under warmer regimes, rather than simplistic, short-term exposures, are

needed as well as an understanding of how species interactions and resource availability shape

their distributions.

Acknowledgements:

We thank the University of Wyoming-National Park Service (UW-NPS) Research Station for

funding. S.H. was supported by NSF awards #OPP-1906015 and #IOS-1557795. A.A.S. was

supported by an NSF Postdoctoral Research Fellowship in Biology (DBI-1807694). Harold

Bergman, Winsor Lowe, Taylor Price, and Lydia Zeglin provided valuable logistic, laboratory, or

field assistance. The UW-NPS Research Station provided laboratory space to perform the

experiments and the Ghalambor Lab provided insect holding equipment. We performed

computational analyses on the Kamiak High Performance Computing Cluster at Washington

State University. We thank Christopher Kozakiewicz and Mark Smithson for comments that

improved the manuscript. Any use of trade, firm, or product names is for descriptive purposes

only and does not imply endorsement by the U.S. Government.

Author contributions:

S.H. and A.A.S. conceived of the study. S.H., A.A.S., K.L.M., L.M.T., J.J.G., D.S.F., M.E.D., and

J.L.K. collected the data. S.H. and A.A.S. analyzed the data and wrote the manuscript with input

from K.L.M., H.A.W., M.E.D., and J.L.K. All authors read and approved the final version.

References:

Amore V, Gaetani B, Puig MA, Fochetti R (2011) New data on the presence of hemocyanin in
Plecoptera: Recomposing a puzzle. J Insect Sci*, **11**, 153.

Andrews RM (1998) Geographic variation in field body temperature of Sceloporus lizards.
Journal of Thermal Biology*, **23**, 329-334.

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. pp Page,
Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom.

Angilletta Jr MJ, Wilson RS, Niehaus AC, Sears MW, Navas CA, Ribeiro PL (2007) Urban
physiology: city ants possess high heat tolerance. PLoS One*, **2**, e258.

Araújo MB, New M (2007) Ensemble forecasting of species distributions. Trends in Ecology &
Evolution*, **22**, 42-47.

Armold M, Blomquist G, Jackson L (1969) Cuticular lipids of insects—III. The surface lipids of
the aquatic and terrestrial life forms of the big stonefly, Pteronarcys californica Newport.
Comparative Biochemistry

Physiology*, **31**, 685-692.

Bálint M, Domisch S, Engelhardt C *et al.* (2011) Cryptic biodiversity loss linked to global climate
change. Nature Climate Change*, **1**, 313.

Beissinger M, Buchner J (1998) How chaperones fold proteins. Journal of Biological Chemistry*,
379, 245-259.

Boeckmann B, Bairoch A, Apweiler R *et al.* (2003) The SWISS-PROT protein knowledgebase
and its supplement TrEMBL in 2003. Nucleic Acids Res*, **31**, 365-370.

Brattstrom BH (1968) Thermal acclimation in anuran amphibians as a function of latitude and
altitude. J Comparative Biochemistry & Physiology*, **24**, 93-111.

Broehan G, Kroeger T, Lorenzen M, Merzendorfer H (2013) Functional analysis of the ATP-
binding cassette (ABC) transporter gene family of Tribolium castaneum. BMC Genomics*,
14, 6.

Buckley BA, Place SP, Hofmann GE (2004) Regulation of heat shock genes in isolated
hepatocytes from an Antarctic fish, Trematomus bernacchii. Journal of Experimental
Biology*, **207**, 3649-3656.

Burmester T (2015) Expression and evolution of hexamerins from the tobacco hornworm,
Manduca sexta, and other Lepidoptera. Insect Biochemistry and Molecular Biology*, **62**,
226-234.

Bushnell B (2014) BBTools software package.

Cargill AS, Cummins KW, Hanson BJ, Lowry RR (1985) The role of lipids, fungi, and
temperature in the nutrition of a shredder caddisfly, Clistoronia magnifica. Freshwater
Invertebrate Biology, **4**, 64-78.
Cavaletto JF, Gardner WS (1999) Seasonal dynamics of lipids in freshwater benthic
invertebrates. In: *Lipids in freshwater ecosystems.* pp Page., Springer.
Chown SL, Hoffmann AA, Kristensen TN, Angilletta Jr MJ, Stenseth NC, Pertoldi C (2010)
Adapting to climate change: a perspective from evolutionary physiology. Climate
Research, **43**, 3-15.
Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M (2005) Blast2GO: a universal
tool for annotation, visualization and analysis in functional genomics research.
Bioinformatics, **21**, 3674-3676.
Connell JH (1961) The influence of interspecific competition and other factors on the distribution
of the barnacle Chthamalus stellatus. Ecology, **42**, 710-723.
Dassati S, Waldner A, Schweigreiter R (2014) Apolipoprotein D takes center stage in the stress
response of the aging and degenerative brain. Neurobiol Aging, **35**, 1632-1642.
Ebner JN, Ritz D, Von Fumetti S (2019) Comparative proteomics of stenotopic caddisfly
Crunoecia irrorata identifies acclimation strategies to warming. Mol Ecol, **28**, 4453-4469.
Ewels P, Magnusson M, Lundin S, Käller M (2016) MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics, **32**, 3047-3048.
Feder ME, Lynch JF (1982) Effects of latitude, season, elevation, and microhabitat on field body
temperatures of neotropical and temperate zone salamanders. Ecology, **63**, 1657-1664.
Finn DS, Khamis K, Milner AM (2013) Loss of small glaciers will diminish beta diversity in
Pyrenean streams at two levels of biological organization. Global Ecology and
Biogeography, **22**, 40-51.
Gamboa M, Tsuchiya MC, Matsumoto S, Iwata H, Watanabe K (2017) Differences in protein
expression among five species of stream stonefly (Plecoptera) along a latitudinal
gradient in Japan. Archives of Insect Biochemistry & Physiology, **96**, e21422.
Giersch JJ, Hotaling S, Kovach RP, Jones LA, Muhlfeld CC (2017) Climate-induced glacier and
snow loss imperils alpine stream insects. Global Change Biology, **23**, 2577-2589.
Giersch JJ, Jordan S, Luikart G, Jones LA, Hauer FR, Muhlfeld CC (2015) Climate-induced
range contraction of a rare alpine aquatic invertebrate. Freshwater Science, **34**, 53-65.
Gilchrist GW (1995) Specialists and generalists in changing environments. I. Fitness landscapes
of thermal sensitivity. Am Nat, **146**, 252-270.

Gutiérrez-Pesquera LM, Tejedo M, Olalla-Tárraga M, Duarte H, Nicieza A, Solé M (2016)
Testing the climate variability hypothesis in thermal tolerance limits of tropical and
temperate tadpoles. Journal of Biogeography, **43**, 1166-1178.

Hagner-Holler S, Pick C, Girgenrath S, Marden JH, Burmester T (2007) Diversity of stonefly
hexamerins and implication for the evolution of insect storage proteins. Insect
Biochemistry and Molecular Biology, **37**, 1064-1074.

Haunerland N (1996) Insect storage proteins: gene families and receptors. Insect Biochemistry
& Molecular Biology, **26**, 755-765.

Hazel JR (1995) Thermal adaptation in biological membranes: is homeoviscous adaptation the
explanation? Annual Review of Physiology, **57**, 19-42.

Healy TM, Brennan RS, Whitehead A, Schulte PM (2018) Tolerance traits related to climate
change resilience are independent and polygenic. Global Change Biology, **24**, 5348-
5360.

Hotaling S, Finn DS, Joseph Giersch J, Weisrock DW, Jacobsen D (2017) Climate change and
alpine stream biology: progress, challenges, and opportunities for the future. Biological
Reviews, **92**, 2024-2045.

Hotaling S, Foley ME, Zeglin LH *et al.* (2019a) Microbial assemblages reflect environmental
heterogeneity in alpine streams. Global Change Biology, **25**, 2576-2590.

Hotaling S, Giersch JJ, Finn DS *et al.* (2019b) Congruent population genetic structure but
differing depths of divergence for three alpine stoneflies with similar ecology, geographic
distributions, and climate change threats. Freshwater Biology, **64**, 335-347.

Hotaling S, Kelley JL, Weisrock DW (2019c) Nuclear and mitochondrial genomic resources for
the meltwater stonefly (Plecoptera: Nemouridae), Lednia tumana (Ricker, 1952). Aquatic
Insects, 1-8.

Hotaling S, Shah AA, Dillon ME, Giersch JJ, Tronstad LM, Finn DS, Kelley JL (In review) Cold
physiology of mountain stoneflies (Plecoptera: Nemouridae): Insights from the high
Rocky Mountains. Western North American Naturalist.

Huey RB, Webster TP (1976) Thermal biology of Anolis lizards in a complex fauna: the
Christatellus group on Puerto Rico. Ecology, **57**, 985-994.

Huss M, Hock R (2018) Global-scale hydrological response to future glacier mass loss. Nature
Climate Change, **8**, 135.

Isaak DJ, Young MK, Nagel DE, Horan DL, Groce MC (2015) The cold-water climate shield:
delineating refugia for preserving salmonid fishes through the 21st century. Global
Change Biology, **21**, 2540-2553.

Jacobsen D (2020) The dilemma of altitudinal shifts: caught between high temperature and low
oxygen. Frontiers in Ecology and the Environment.
Jacobsen D, Milner AM, Brown LE, Dangles O (2012) Biodiversity under threat in glacier-fed
river systems. Nature Climate Change, **2**, 361-364.
Jones LA, Muhlfeld CC, Marshall LA, Mcglynn BL, Kershner JL (2014) Estimating Thermal
Regimes of Bull Trout and Assessing the Potential Effects of Climate Warming on
Critical Habitats. River Research and Applications, **30**, 204-216.
Jordan S, Giersch JJ, Muhlfeld CC, Hotaling S, Fanning L, Tappenbeck TH, Luikart G (2016)
Loss of genetic diversity and increased subdivision in an endemic alpine stonefly
threatened by climate change. PLoS One, **11**, e0157386.
Jørgensen LB, Malte H, Overgaard JJFE (2019) How to assess Drosophila heat tolerance:
unifying static and dynamic tolerance assays to predict heat distribution limits. **33**, 629-
642.
Kalvari I, Argasinska J, Quinones-Olvera N *et al.* (2017) Rfam 13.0: shifting to a genome-centric
resource for non-coding RNA families. Nucleic Acids Res, **46**, D335-D342.
Kearney M, Shine R, Porter WP (2009) The potential for behavioral thermoregulation to buffer
"cold-blooded" animals against climate warming. Proceedings of the National Academy
of Sciences, **106**, 3835-3840.
Khamis K, Brown LE, Hannah DM, Milner AM (2015) Experimental evidence that predator range
expansion modifies alpine stream community structure. Freshwater Science, **34**, 66-80.
King AM, Macrae TH (2015) Insect heat shock proteins during stress and diapause. Annual
Review of Entomology, **60**, 59-75.
Krueger F (2015) Trim Galore!: a wrapper tool around Cutadapt and FastQC to consistently
apply quality and adapter trimming to FastQ files. Babraham Bioinformatics, Cambridge,
United Kingdom. pp Page.
Levy O, Buckley LB, Keitt TH, Smith CD, Boateng KO, Kumar DS, Angilletta Jr MJ (2015)
Resolving the life cycle alters expected impacts of climate change. Proceedings of the
Royal Society B: Biological Sciences, **282**, 20150837.
Li H, Handsaker B, Wysoker A *et al.* (2009) The sequence alignment/map format and SAMtools.
Bioinformatics, **25**, 2078-2079.
Lindquist S, Craig E (1988) The heat-shock proteins. Annual Review of Genetics, **22**, 631-677.
Lister BC, Garcia A (2018) Climate-driven declines in arthropod abundance restructure a
rainforest food web. Proceedings of the National Academy of Sciences, **115**, E10397-
E10406.

Lun AT, Chen Y, Smyth GK (2016) It's DE-licious: a recipe for differential expression analyses
of RNA-seq experiments using quasi-likelihood methods in edgeR. In: *Statistical*
Genomics. pp Page., Springer.

Markl J, Winter S (1989) Subunit-specific monoclonal antibodies to tarantula hemocyanin, and a
common epitope shared with calliphorin. Journal of Comparative Physiology B*,* **159**,
139-151.

Mccarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor RNA-Seq
experiments with respect to biological variation. Nucleic Acids Res*,* **40**, 4288-4297.

Muir AP, Nunes FL, Dubois SF, Pernet F (2016) Lipid remodelling in the reef-building
honeycomb worm, Sabellaria alveolata, reflects acclimation and local adaptation to
temperature. Sci Rep*,* **6**, 35669.

Nebeker AV (1971) Effect of water temperature on nymphal feeding rate, emergence, and adult
longevity of the stonefly Pteronarcys dorsata. Journal of the Kansas Entomological
Society, 21-26.

Peckarsky BL, Mcintosh AR, Horn SC *et al.* (2014) Characterizing disturbance regimes of
mountain streams. Freshwater Science*,* **33**, 716-730.

Pernet F, Tremblay R, Comeau L, Guderley H (2007) Temperature adaptation in two bivalve
species from different thermal habitats: energetics and remodelling of membrane lipids.
Journal of Experimental Biology*,* **210**, 2999-3014.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL (2015) StringTie
enables improved reconstruction of a transcriptome from RNA-seq reads. Nature
Biotechnology*,* **33**, 290.

Pörtner HO, Knust R (2007) Climate change affects marine fishes through the oxygen limitation
of thermal tolerance. Science*,* **315**, 95-97.

Pörtner HO, Peck L, Somero G (2007) Thermal limits and adaptation in marine Antarctic
ectotherms: an integrative view. Philos Trans R Soc Lond B Biol Sci*,* **362**, 2233-2258.

R Core Team (2013) R: A language and environment for statistical computing.

Rempel RS, Carter JC (1987) Temperature influences on adult size, development, and
reproductive potential of aquatic Diptera. Canadian Journal of Fisheries & Aquatic
Sciences*,* **44**, 1743-1752.

Reyes MA, Corcuera LJ, Cardemil L (2003) Accumulation of HSP70 in Deschampsia antarctica
Desv. leaves under thermal stress. Antarctic Science*,* **15**, 345-352.

Rezende EL, Tejedo M, Santos M (2011) Estimating the adaptive potential of critical thermal
limits: methodological problems and evolutionary implications. Functional Ecology, **25**,
111-121.

Risso D, Ngai J, Speed TP, Dudoit S (2014) Normalization of RNA-seq data using factor
analysis of control genes or samples. Nature Biotechnology, **32**, 896.

Robinson MD, Mccarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics, **26**, 139-140.

Roughgarden J (1974) Niche width: biogeographic patterns among Anolis lizard populations.
Am Nat, **108**, 429-442.

Sánchez-Bayo F, Wyckhuys KA (2019) Worldwide decline of the entomofauna: A review of its
drivers. Biological Conservation, **232**, 8-27.

Schütz SA, Füreder L (2019) Egg development and hatching success in alpine chironomids.
Freshwater Biology, **64**, 685-696.

Seebacher F, White CR, Franklin CE (2015) Physiological plasticity increases resilience of
ectothermic animals to climate change. Nature Climate Change, **5**, 61.

Shah AA, Funk WC, Ghalambor CK (2017a) Thermal acclimation ability varies in temperate and
tropical aquatic insects from different elevations. J Integrative Comparative Biology, **57**,
977-987.

Shah AA, Gill BA, Encalada AC et al. (2017b) Climate variability predicts thermal limits of
aquatic insects across elevation and latitude. Functional Ecology, **31**, 2118-2127.

Sims D, Ilott NE, Sansom SN et al. (2014) CGAT: computational genomics analysis toolkit.
Bioinformatics, **30**, 1290-1291.

Sinensky M (1974) Homeoviscous adaptation—a homeostatic process that regulates the
viscosity of membrane lipids in Escherichia coli. Proceedings of the National Academy of
Sciences, **71**, 522-525.

Terblanche JS, Hoffmann AA, Mitchell KA, Rako L, Le Roux PC, Chown SL (2011) Ecologically
relevant measures of tolerance to potentially lethal temperatures. J Exp Biol, **214**, 3713-
3725.

Trapnell C, Roberts A, Goff L et al. (2012) Differential gene and transcript expression analysis of
RNA-seq experiments with TopHat and Cufflinks. Nature Protocols, **7**, 562.

Treanor HB, Giersch JJ, Kappenman KM, Muhlfeld CC, Webb MaH (2013) Thermal tolerance of
meltwater stoneflyLednia tumananymphs from an alpine stream in Waterton–Glacier
International Peace Park, Montana, USA. Freshwater Science, **32**, 597-605.

Tronstad LM, Hotaling S, Giersch JJ, Wilmot OJ, Finn DS (2019) Headwater streams fed by

subterranean ice: potential climate refugia for mountain communities? bioRxiv.

U.S. Fish & Wildlife Service (2019) Endangered and Threatened Wildlife and Plants:

Threatened Species Status for Meltwater Lednian Stonefly and Western Glacier Stonefly

with a Section 4(d) Rule. Federal Register*,* **84**, 64210-64227.

Urban MC, Bocedi G, Hendry AP *et al.* (2016) Improving the forecast for biodiversity under

climate change. Science*,* **353**, aad8466.

Verberk WC, Overgaard J, Ern R, Bayley M, Wang T, Boardman L, Terblanche JS (2016) Does

oxygen limit thermal tolerance in arthropods? A critical review of current evidence.

Comparative Biochemistry & Physiology Part A: Molecular Integrative Physiology*,* **192**,

64-78.

Ward J (1994) Ecology of alpine streams. Freshwater Biology*,* **32**, 277-294.