Running head: Prezygotic isolation in Arabidopsis lyrata

Limited phenological and pollinator-mediated isolation among selfing and outcrossing

Arabidopsis lyrata populations

Authors: Courtney E. Gorman*[1], Lindsay Bond[2], Mark van Kleunen[1,3], Marcel E. Dorken[2], and

Marc Stift[1]

[1] University of Konstanz, Department of Biology, Universitaetsstrasse 10, Konstanz, BW, DE

78464

[2] Trent University, Department of Biology, DNA B108.10, Peterborough, ON, CAN K9J 7B8

[3] Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou

University, Taizhou 318000, China

* Corresponding author (email: courtney.gorman@uni-konstanz.de)

Abstract

Mating system transitions from outcrossing to selfing have been a frequent evolutionary shift in angiosperms and clearly play a role in species divergence. However, many questions remain about the initial mechanistic basis of reproductive isolation during the evolution of selfing. For instance, how important are prezygotic pre-pollination mechanisms (e.g., changes in phenology and pollinator visitation) in maintaining reproductive isolation between newly arisen selfing populations and their outcrossing ancestors? To test whether changes in phenology and pollinator visitation reproductively isolate selfing populations of Arabidopsis lyrata from conspecific outcrossing populations, we conducted a common garden experiment in the native range with plants from selfing and outcrossing populations as well as their F1 hybrids. Hybrid plants resulting from crosses between self-incompatible populations had the highest probability of flowering. Time of peak flowering differed among mating systems although the magnitude of the difference was small. Flowering duration and maximum flower number did not strongly differ between mating systems and hybrids, and these phenological traits had little influence on the frequency of pollinator visits of either solitary bees or hoverflies. Additionally, pollinators preferred to visit flowers on the same plant (~50% of the time) rather than exploring nearby plants, regardless of whether they were of an alternative mating system type or F1 hybrids, creating a large opportunity for geitonogamy. Finally, by modelling pollen-transfer probabilities, we found no significant differences in the opportunity for pollinations between selfing and outcrossing plants. Overall, we found that selfing and outcrossing *A. lyrata* populations show some phenological divergence but are likely only weakly prezygotically isolated via pre-pollination mechanisms.

Key words: *Arabidopsis lyrata*, mating system, phenology, pre-pollination, prezygotic isolation,

pollinators, reproductive isolation, selfing

Mating-system transitions from obligate outcrossing to predominantly selfing have arisen

repeatedly across almost all major plant lineages (Barrett 2002). Up to 15% of seed plants are

predominantly selfing and many share a relatively recent common ancestor with outcrossing

species (Goodwillie et al. 2005). The transition from outcrossing to selfing is usually associated

with convergent evolution of a flower morphology that optimizes self-pollination and resource

use, termed the "selfing syndrome" (Darwin 1876, Jain 1976, Sicard & Lenhard 2011). These

types of changes in flowering likely contribute to the reproductive isolation of selfing lineages

(Rieseberg & Willis 2007), but it is unclear if they or their subsequent effects on pollinators are

the main drivers of reproductive isolation in incipient selfing species.

Reproductive barriers are essential to the maintenance of evolutionary independence of

parapatric populations (i.e. populations with slightly overlapping ranges). Reproductive isolation

can be partial or complete, and most plant species are isolated by a combination of pre- and

postzygotic barriers (Coyne & Orr 2004, Rieseberg & Willis 2007, Baack et al. 2015). In plants,

prezygotic mechanisms are generally more important than postzygotic mechanisms in

contributing to the total reproductive isolation of species (Rieseberg & Willis 2007, Lowry et al.

2008, Widmer et al. 2009, Baack et al. 2015). Post-pollination mechanisms such as genetic

incompatibilities can cause differences in seed number and/or seed viability, but pre-pollination

mechanisms generally contribute more to the total reproductive isolation of plant species

(Rieseberg & Willis 2007, Lowry et al. 2008, Widmer et al. 2009). Although rarely addressed,

this principle appears to hold for cases where a shift to self-fertilization has played a part in

speciation. For example, in two closely related *Mimulus* species with a contrasting mating

system, differences in mean flowering date and floral display contributed the most to

reproductive isolation compared to other pre- and post-pollination mechanisms (Martin & Willis

2007). However, whether the transition to self-fertilization could also promote rapid prezygotic

reproductive isolation via changes in floral morphology and associated shifts in pollinator

preferences (Coyne & Orr 2004) has rarely been studied.

Plant phenological traits and the behaviour of pollinators could readily interact. For

example, pollinator behaviour and the number of flowers should both play a large role in

determining the opportunity for the flowers of self-compatible plants to be fertilized by a

different flower on the same plant (i.e. geitonogamy). Furthermore, geitonogamy could help to

reproductively isolate self-compatible individuals. For instance, if pollinators commonly visit

multiple flowers on the same individual, it could facilitate higher selfing rates of self-compatible

individuals (Karron et al. 2009). Different types of pollinators, such as flies vs. bees, could also

differ greatly in their pollination strategies (Gervasi & Schiestl 2017). Additionally, if pollinators

more commonly visit plants in very close proximity, this could contribute to greater population

viscosity and result in more matings among closely related individuals that share the same

mating system (Harder & Barrett 1996, Barrett 2003, Mitchell et al. 2009). Furthermore, due to

flower attractiveness, pollinators might preferentially and repeatedly visit plants exhibiting a

particular mating system type. Alternatively, at the earliest stages of divergence, pollinators

might show limited or no ability to differentiate between plants with alternative mating types.

Here, we use *Arabidopsis lyrata* ssp. *lyrata* (L.) to examine the role of differences in

phenology and pollinator attraction as mechanisms of reproductive isolation in a recently

diverged selfing lineage. In several populations across the range of this normally outcrossing

species (multi-locus outcrossing rate: $0.83 < T_m < 0.99$), all plants are self-compatible, have low

outcrossing rates, and therefore reproduce primarily through selfing (multi-locus outcrossing

rate: $0.09 < T_m < 0.41$) (Mable et al. 2005; Mable & Adam 2007; Foxe et al. 2010). The selfing

and outcrossing populations are geographically interspersed, therefore secondary contact

following evolutionary divergence in parapatry is likely. Also, the transition to selfing in these

populations is thought to have happened < 10,000 years ago because 1) the range now occupied

by outcrossing and selfing populations was mostly covered by ice during the last glacial

maximum (Foxe et al. 2010), and 2) the selfing populations have not developed a selfing

syndrome (e.g. smaller flower size and/or a reduction in pollen, nectar, and scent production)

(Carleial et al. 2017b). This raises the question of whether selfing populations have diverged

from their outcrossing ancestors in traits conferring reproductive isolation. Similarly, given that

outcrossing and selfing populations are at least partly interfertile and can regularly produce

healthy offspring (Willi 2013a, 2013b), F1 hybrids may be a critical factor in determining

whether secondary contact would lead to coalescence of the diverged populations or alternatively

reinforce their evolved differences.

In a common-garden experiment set within the native range of *A. lyrata*, we simulated

two phases of secondary contact between selfing and outcrossing populations. The first phase

corresponds to initial contact between parental plants from selfing and outcrossing populations.

The second phase corresponds to secondary contact between admixed plants (hybrids between

populations) and parental plants. This allowed us to test whether the evolution of selfing has led

to pre-pollination isolation through divergence in phenology and/or insect pollinator attraction.

Specifically, we asked whether there was reproductive isolation between outcrossing and selfing

plants and their F1 hybrids through differences in 1) the timing or intensity of flowering; and 2)

pollinator visitation rates and paths. Based on this, we tested whether phenological differences

and pollinator behaviour reduced the opportunities for pollen exchange between mating systems.

Moreover, as geitonogamy can also contribute to reproductive isolation, we quantified the

opportunities for geitonogamy.

Materials and Methods

Study system

Arabidopsis lyrata spp. *lyrata* (L.) is a small, short-lived perennial that is native to North

America. It occurs in dry-mesic habitats with shallow soils, such as rock outcrops and sand

dunes. Individual plants can produce several stems that terminate in racemes of numerous (>20)

small white flowers. The primary pollinators of *A. lyrata* are small solitary bees and hoverflies,

which are attracted to the nectar and pollen of the flowers. The ancestral condition in

Arabidopsis lyrata is self-incompatibility, however the barrier to self-fertilization has broken

down in several North American populations (Mable et al. 2005). Additionally, many of these

newly self-compatible populations have evolved high selfing rates (Foxe et al. 2010). There is

some variation in floral traits, such as flower size and pollen:ovule ratios, among selfing and

outcrossing populations, which is primarily explained by population genetic background and not

mating system (Carleial et al. 2017b).

Crossing designs

To generate the material needed to simulate secondary contact between diverged selfing

and outcrossing populations, we sowed field-collected seeds from 12 North

American *A. lyrata* populations with known breeding and mating systems (Foxe et al. 2010;

seeds were kindly provided by Barbara Mable, University of Glasgow). These included six

populations characterized as outcrossing (high outcrossing rates, high frequency of self-

incompatible individuals, hereafter referred to as SI populations) and six populations

characterized as selfing (low outcrossing rates, high frequency of self-compatible individuals,

hereafter referred to as SC populations) (Table S1).

In 2012 and 2013, we then produced seeds by manually cross- and self-pollinating up to

eight plants per population. To perform the pollinations, we emasculated a flower prior to anther

dehiscence, or the same individual in 'selfed' crosses, and rubbed a freshly dehisced anther from

a haphazardly chosen plant from the same population over its stigma. Progeny were produced

with the following cross-types: within SI population (SI-within), and within SC population both

by crossing (SC-within) and by selfing (SC-self).

Then to generate the material needed to simulate admixture between the parental

populations and their F1 hybrids, we performed a full diallel cross in 2014 and 2015 with six

plants of each of the six SI and six SC populations. This cross produced additional progeny of

the SI-within and SC-within cross types, as well as the following cross types: between SI

population (SIxSI), between SC population (SCxSC), between SC and SI population reciprocally

(SIxSC or SCxSI). All crosses were reciprocal and yielded a total of 1032 seed families (Table

S2).

<u>Experimental design of common garden experiment</u>

To test whether differences in phenology and flower-visitor attraction can reproductively

isolate plants from selfing populations, we set up a common garden experiment at Trent

University in Peterborough, Ontario, Canada. This location is at an intermediate latitude within

the geographic range of the source populations (Fig. 1). From March 20 to 22, 2018, for each

seed family, up to 50 seeds were sown on a moistened peat-based substrate in one pot. Plants

were grown in climate chambers with 11-hour days and a 21°C/18°C day/night cycle at 95%

humidity. Between April 18 and May 1, when seedlings had developed at least two true leaves,

we transplanted three haphazardly chosen seedlings from each seed family to individual Stuewe

and Sons Ray Leach "Cone-tainers" ™ [Tangent, Oregon, USA] with the same peat-based

substrate. On May 10, plants were moved to the common garden, prior to any flowering.

Within the common garden, plants were organized in a randomized block design. There

were three replicates of three 3x6m blocks. Each of the nine resulting blocks contained between

150-180 individual plants distributed randomly over 180 positions within 9 cone-tainer trays

with plants from each seed family and cross type evenly distributed among the blocks. In total,

1512 plants were raised in the common garden.

Phenological data

To test for potential reproductive isolation between the cross types due to differences in

phenology, we recorded daily for each plant whether it flowered and how many open flowers it

had. Open flowers were defined as flowers with visible reproductive organs (stigma and anthers)

and that still had petals attached to the flower. Besides calculating opportunities for pollen-

transfer between outcrossing and selfing populations, this allowed us to compare the time to

onset of flowering, flowering duration, time of peak flowering, and the maximum flower number

(i.e., the number of flowers on the day of peak flowering) for each individual for each of the

cross-types.

Flower visitor observations

To test for differences in insect attraction and flower visitor movements within and

between plants, we recorded flowers with GoPro Hero Session® [San Mateo, California, USA]

cameras. Specifically, we tested whether there were differences in the potential for

geitonogamous selfing (visitor movement within the same plant), and for outcrossing (i.e., visitor

movement between plants) within and between mating system. To standardize the recording

procedure, 4-6 flowering plants (depending on their size) were taken from their blocks and

placed in a tray located at the front of their respective blocks. This method ensured clear video

footage of multiple focal plants simultaneously, while minimizing interfering with the visual

context of the pollinators provided by the configuration of plants in the block design. To make

sure that focal plants had a different set of neighbors for each set of observations, we combined

flowering individuals systematically according to their position in the block, going through the

block in three different ways: 1) taking consecutive plants in a vertical direction, 2) taking

consecutive plants in a horizontal direction, and 3) taking plants from the same position but in

different trays. Due to this approach, the cross type of the focal plants combined in the video-

frames was random.

We recorded 12-15-minute-long videos that were later trimmed to the central 10 minutes

to exclude potential effects of disturbance during starting and stopping the cameras. In total, 500

videos were taken throughout the flowering period, of which a random subset of 140 videos

(23.3 hours of video) were analysed by the first author in a random order. In total, these videos

included 379 unique individuals (41% of all flowering individuals in the common garden), and

123 plants were observed in multiple videos. For each visitor, we recorded whether it was a

solitary bee or hoverfly, the duration of the visit and the path it took (see below). Finer

taxonomic identification was not possible due to the video resolution, but we took high quality

photographs to identify the most common visitors: hoverflies (Syrphidae) *Eristalis arbustorum*,

Syritta pipiens, *Sphaerophoria* sp. and *Toxomerus marginatus*, and solitary bees from the family

Halictidae (kindly identified by Bill Crins, Toronto, Canada).

The path that each visitor took after its initial visit to a flower was recorded to test

whether plants from selfing populations received fewer visits than those from outcrossing

populations as is expected in selfing plants (Fausto Jr. et al. 2001). Moreover, because pollinators

will often focus on exploiting one type of flower and/or floral scent, we tested whether visitors

were more likely to move to neighbouring plants with the same mating system than to plants

with a different mating system, and whether progeny from crosses between mating system

received fewer visits than progeny from crosses within the same mating system. We classified

visitor paths as: "away" – the visitor left the video frame after an initial visit; "same" – the visitor

visited a second flower on the same individual; or to one of the cross types as defined above

("SI-within", "SC-within", "SC-self", "SIxSI", "SCxSC", "SIxSC", "SCxSI") – the visitor went

to a flower on a different individual. This also allowed us to quantify the general behaviour of

pollinators.

<u>Statistical analyses</u>

All statistical analyses were done in R 3.5.1 (R Core Team, 2019). To test if there were

differences in the time of peak flowering and duration of flowering between SI and SC cross

types (SI-within, SC-within, SC-self) and between within population cross types and between

population cross types (SI-within, SC-within, SC-self vs SIxSI, SCxSC, SIxSC, SCxSI), we used

Gaussian linear mixed-effects models implemented in lme4 (Bates et al. 2015) using *cross type*

as a fixed effect, and *maternal population* and *paternal population* as random effects. We used

"Improper" prior distributions, i.e. distributions with density functions that do not integrate to 1

and are therefore not "proper" probability distributions (Korner-Nievergelt et al. 2016).

Specifically, $p(\beta) \propto 1$ was implemented for the model coefficients and $p(\sigma^2) \propto 1/\sigma^2$ for the

variance parameters. To obtain the posterior distribution, 5000 values were directly simulated

from the joint posterior distribution of the model parameters using the function sim of the R

package 'arm' (Gelman & Hill 2007). The means of the simulated values from the joint posterior

distributions of the model parameters were then used as estimates, and the 2.5% and 97.5%

quantiles were used as the lower and upper limits of the 95% credible intervals to make

comparisons among cross types.

To test if there were differences in the mean maximum flower number among the cross

types, a hurdle model (hurdle function, package 'pscl'; Zeileis et al. 2008, Jackman 2017) with a

negative binomial distribution that included *maximum flower number* as the response variable

and *cross type* as a fixed effect was performed. The hurdle model accounts for the excess number

of zero counts in the maximum flower number data. This model specifies one process for zero

counts and a separate process for positive counts. The zero counts (flower number as either 0 or

1) were then modelled with a binomial logit model and the positive counts (plants that flowered)

with a truncated negative binomial model. The hurdle model also allowed us to calculate the

probability that individuals from a cross type would flower.

Pollinator visitation rate (per plant) was analysed separately for the two main visitor

classes hoverflies and solitary bees. The cross type 'SC-self' was excluded from the analyses of

pollinator visitation due to low sample size. To test if there were differences in the frequency of

pollinator visits among the cross types, two identical generalized linear mixed-effects models

with negative binomial distributions with *number of visits* as the response variable (one model

for visits made by hoverflies and another one for solitary bees). The explanatory variables were

cross type and *flower number* as fixed effects, and *maternal population* and *paternal population*

as random effects. In these models, the number of adaptive Gauss-Hermite quadrature points

(nAGQ) was set to zero, which optimizes the random effects and the fixed-effects coefficients in

the penalized iteratively reweighted least squares step (Bates et al. 2015). This results in a faster

but less precise parameter estimation for generalized mixed effect models (Bates et al. 2015).

These models used a log-link function. Improper prior distributions were used, as in the analyses

of time of peak flowering and flowering duration.

Pollinator visitation paths were analysed in two ways. The probability that a pollinator

would make a certain choice after landing on a flower was analysed with a multinomial logistic

regression as implemented in the function multinom in the package 'nnet' (Venables & Ripley

2002). *Path* in the multinomial model included all cross types and the same plant (opportunity

for geitonogamy) as path options, along with the option of leaving the observation frame. This

model included both *cross type* and *flower number* as fixed effects and *path* as the response

variable using a logit link function. To further parse the pollinator preference and the effect of

flower number and distance between plants in the frame, a conditional logistic regression

(function clogit, package 'survival'; Therneau & Grambsch 2000) was performed. The

conditional logistic regression was performed separately for hoverflies and solitary bees and

included the insect's selection for any of the cross types in the same video frame as the response

variable, as well as *relative flower number*, *relative distance*, and *cross type* as fixed effects, and

finally *switch ID* as the strata. The strata command specifies the group of observations inherent

to our video recordings. The strata in this case specifies the group of choice options for each

pollinator in each video. Switch ID was defined as: what the insect selected (1) and everything

the insect did not select (0) and incorporated information about the distance to the other

individuals and the flower number relative to the other individuals. The cross type 'SI-within'

was used as the baseline as this cross type represents the ancestral condition in *A. lyrata*. *Relative*

flower number and *relative ranked distance* were obtained by dividing by the maximum value

within the same video-frame.

<u>Pollen-transfer probabilities</u>

273   To examine whether there were differences in the opportunities for self- or outcross

pollination between selfing and outcrossing plants, we used the empirical information on

phenology and pollinator behaviour to model opportunities for outcrossing between 'SI-within'

and 'SC-within' plants and for geitonogamous pollen movement within plants. Between

population cross types were excluded from these analyses. In terms of outcrossing, we were

interested in opportunities for pollen exchange within versus between mating types (e.g., whether

plants from self-compatible populations of *A. lyrata* had more opportunity to mate with each

other than with plants from self-incompatible populations) as a potential mechanism of

reproductive isolation. To do this, we used calculations of K_{ij} - the "pollen transfer

probabilities" outlined in Brunet and Charlesworth (1995). Here, we use the calculation of K to

refer to the opportunity for mating between plants from different populations in the common

garden, based on the overlap in the number of flowers of each mating type per day.

285   In Brunet and Charlesworth (1995), K_{ij} refers to the probability that flowers at the ith

position on an inflorescence are pollinated by flowers on other plants at the jth position. Here, we

are not interested in the effects of floral position on pollen transfer probabilities, but in the

possible effect of mating type. Accordingly, we estimated opportunities for pollen transfer within

versus between self-incompatible and self-compatible mating types by calculating the following

values of K:

1. K_{ss}

2. K_{so}

3. K_{os}

4. K_{oo}

where, K_{ss} refers to the opportunity for plants from self-compatible populations (SC-within

cross type) to fertilize flowers on other plants from self-compatible populations, K_{so} refers to the

opportunity for plants from self-compatible populations to fertilize flowers on plants from self-

incompatible populations (SI-within cross type), and so on.

Brunet and Charlesworth define the probability of pollen transfer between flowers of type i and

j on day c, K_{ij} as: $\frac{\sum_c f_{cj}^m N_{ci}^f}{\sum_c N_{cj}^m}$ where, $f_{cj}^m = \frac{N_{cj}^m}{\sum_j N_{cj}^m}$.

301   In the above expressions, the superscripts m and f refer to plants in the male and female

phases. For *A. lyrata*, which is a simultaneous hermaphrodite, m $= f$, but because we are

interested in pollen movement between plants, $i \neq j$. To calculate mating-type specific values of,

for example, K_{ss}, we calculated f^m as the proportion of all flowers open per day in the common

garden that were from individual plants from self-compatible populations (the SC-within cross

type). Therefore, for this calculation, the value of the numerator, N_{cj}^m, was calculated only for

those plants. Plants from self-incompatible populations were included in the calculation

of N^m for K_{os} and K_{oo}. For all values of K, all plants were included in the calculation of the

denominator of f^m, $\sum_j N_{cj}^m$.

The calculated values of K are frequency dependent - a small group of plants of one

mating type surrounded by plants of the opposite mating type would have more opportunities for

between, rather than within mating-type pollen transfer. Because we were specifically interested

in opportunities for pollen transfer driven by phenology, not frequency, we used bootstrapping to

generate 200 randomly sampled, equal-sized populations of plants of each mating type for the

calculation of K. For each mating type, we sampled 200 plants (with replacement) of each cross

type for inclusion in each calculation of K.

Two values of K refer to within mating-type fertilization opportunities and two of them

to between mating-type fertilization opportunities. To evaluate whether plants from populations

of the two different mating types (self-compatible versus self-incompatible) differed in the

proportion of within- versus between mating-type pollen transfer opportunities, we used linear,

mixed models for each set of bootstrapped values of K. Population was included as random

effect in these models. In the absence of phenological differences between plants from self-

compatible and self-incompatible populations, the expected value of the parameter estimate for

the mating-type effect is 0. Accordingly, to test whether plants representing the two mating types

differed in their opportunities for within- versus between mating-type pollen transfer, we tested

whether the distribution of parameter estimates from each set of bootstrapped values of

K differed from 0 using a two-tailed *t*-test.

The opportunity for geitonogamous self-pollination is determined by the number of

simultaneously open flowers per plant and the likelihood that pollinators will move from one

flower to another on the same plant. Videos of pollinator movements provide per-population

estimates of that likelihood. We calculated the opportunity for geitonogamous pollen

transfer, G_c, for plants with n open flowers as a geometric series of the likelihood of within-plant

pollinator movement x. That is, for each day c, $G_c = x + x^2 + x^3 + \cdots + x^{n-1}$. The total

opportunity for geitonogamous pollen transfer over the flowering season was calculated as $\sum_c G_c$.

Results

<u>Phenology</u>

Of the 1512 plants in the common garden, 938 flowered (62%). The main flowering

period lasted six weeks from June 1 to July 14, although a few individuals flowered later (nine

individuals flowered a second time and 10 individuals flowered for the first time as late as

September) (Fig. S1). The probability of an individual flowering varied by cross type. SI-within

and SC-within cross types did not strongly differ from each other in the probability of flowering

(55% and 61%, respectively; CrI overlapping; Fig. 2a), however the probability of flowering of

the SC-self cross type (29%) was substantially lower (CrI not overlapping with any of the other

cross-types; Fig. 2a). So, while progeny formed by selfing flowered less, merely having the

ability to self did not substantially decrease the probability of flowering when compared to

individuals from outcrossing populations. The F1 hybrid cross types did not differ from the

within-population cross types in the probability of flowering (54%-64%; CrIs overlapping; Fig.

2a), with the exception of the SIxSI cross type being more likely (83%) to flower than the other

F1 or within-population cross types (CrI not overlapping with any of the other cross-types; Fig.

2a). Additionally, the direction of the cross for SIxSC and SCxSI hybrid F1 crosses did not have

an obvious effect on flowering probability, as both cross types had similar probabilities (63%

and 64% respectively) for flowering (CrIs overlapping; Fig. 2a).

The time of peak flowering differed among the cross types. While there were no strong

differences in the time of peak flowering between the SI-within and SC-within cross types, the

SC-self cross type peaked in flowering about one to two days earlier (average = 27.6 days) than

the other within or F1 cross types (however, CrIs overlapped with all cross types except SIxSI;

Fig. 2b). The day of peak flowering did not differ between SIxSC and SCxSI plants, indicating

that cross direction did not influence the time of peak flowering (CrIs overlapping; Fig. 2b).

There were also no strong differences in peak flowering between the within-population and F1

cross types (CrIs overlapping), with the exception that the SIxSI cross type tended to reach peak

flowering one to two days later (Fig. 2b). Similarly, flowering duration did not strongly differ

among the cross types (Fig. 2c), but the SC-self cross type tended to have a shorter duration (6-

15 days) than the other within population cross types (11-19 days) or F1 cross types (15-21 days)

(Fig. 2c).

For flowering plants of all cross types, the mean maximum number of open flowers on a

single day ranged from 10.8 to 13.6 flowers. After correcting for zero-inflation in the maximum

flower number model, there were no strong differences in maximum flower number among the

cross types.

<u>Pollinator visitation</u>

There were no differences in flower visitation between the within population cross types

or between the within population cross types and the F1 hybrids (all CrIs overlapping; Fig. 3a,

Fig. 3b). Solitary bees and hoverflies were the predominant visitors, and they had similar

visitation frequencies and no clear pattern of preference for any of the cross types (compare Fig.

3a and 3b). The behaviour of both types of pollinators appeared to facilitate geitonogamous self-

pollination, as ~50% of the movements between flowers were to a different flower on the same

plant (Fig. 4). When cases of a pollinator visiting another flower on the same plant were not

considered, the odds of an initial visitor moving to a plant in the frame decreased by 89%

(solitary bees) and 94% (hoverflies) for each unit increasing relative distance (significantly

negative odds-ratios for relative distance; Table 1). In other words, pollinators were more prone

to visit the nearest plant, regardless of the cross type or the number of flowers on the

neighbouring plant.

Opportunities for pollen-transfer

Based on a bootstrapping approach that integrated the timing and intensity of flowering

throughout the entire flowering period, the opportunity for between vs. within cross type pollen-

transfer was nearly equal both for the SI-within and SC-within cross type (Fig. 5). In other

words, slight shifts in phenology and flowering intensity (Fig. S1) are unlikely to lead to

reproductive isolation. Additionally, a similar approach that also took into account the pollinator

paths showed that the opportunity for geitonogamy arising from plants having multiple flowers

open simultaneously and the high frequency of within plant movement of pollinators (Fig. 4) did

not differ between SI-within and SC-within plants.

Discussion

Our common garden experiment simulating secondary contact between outcrossing and

selfing populations showed that phenology largely overlapped between plants from outcrossing

and selfing populations. There were also no differences in pollinator preference related to mating

system. Regardless of mating system, pollinators tended to move between flowers on the same

plant, thus facilitating opportunities for geitonogamy. Our models of pollen-transfer

probabilities, which integrated differences in phenology (timing and intensity of flowering),

revealed equal opportunities for pollen-transfer within- and among mating systems. Together,

this suggests that prezygotic pre-pollination mechanisms do not strongly reproductively isolate

plants from selfing and outcrossing populations of *Arabidopsis lyrata*. However, because plants

simultaneously open multiple flowers, and pollinators predominantly move from flower to

flower on the same plant, our data suggest that there is a large opportunity for geitonogamy in

this system. Such geitonogamy could isolate selfers to some extent.

Pollinator visitation

Our results show that differences in pollinator preference do not play a large role in the

reproductive isolation of the newly diverged selfing populations. We found no differences in the

preferences of the two main pollinator types, hoverflies and solitary bees, among the cross types.

Additionally, when given a choice pollinators preferred to stay on the same individual versus

exploring nearby plants regardless of cross type. When they did choose to visit another plant,

pollinators preferred to go to the closest plant regardless of how many flowers it had or what

cross type it was. This lack of pollinator preference could be due to the pollinators simply being

generalists, as previous studies have shown that hoverflies, for instance, are not very choosy with

respect to the plants they visit (Gervasi & Schiestl 2017). Additionally, there could be a lack of

floral differences because selfing individuals still need pollinators to transfer their pollen as they

are not completely autonomous (Carleial et al. 2017b). Overall, pollinator preference seems to be

playing little role in differentiating the selfing from outcrossing populations of *A. lyrata*.

Nevertheless, the behaviour of the pollinators could favour selfing for several other

reasons. For instance, we found that pollinators often visit different flowers on the same

individual, irrespective of mating system. This should provide ample opportunity for within-

individual pollen transfer (Darwin 1876, Harder & Barrett 1995), and thus for self-compatible

individuals to self-fertilize through geitonogamy. Moreover, when pollinators moved between

plants, they mainly moved between nearby individuals. Given that *A. lyrata* seeds have no

mechanisms to promote dispersal, and plants can produce over 1000 seeds per season,

neighbouring plants could be highly related to each other (Willi & Määttänen 2011). As a

consequence, the observed behaviour of pollinators could cause mate limitation in self-

incompatible plants, making the transfer of cross-pollen rare and/or mainly from incompatible

partners (e.g., from relatives that share S-alleles). In self-compatible plants, on the other hand,

geitonogamy may help overcome this mate limitation and provide reproductive assurance (Baker

1955). Even without mate limitation, where geitonogamy does not provide reproductive

assurance but rather results in seed discounting, selfing could still be favoured due to the

associated inherent transmission advantage (Fisher 1941). The transmission advantage would

favour selfers if inbreeding depression is low. Indeed, the relatively low estimates of inbreeding

depression for our study populations (Carleial et al. 2017a) imply that such transmission

advantage alone could be sufficient to drive the evolution of selfing. There is usually a low

frequency of self-compatible individuals in outcrossing populations (Foxe et al. 2010; Mable et

al. 2017). Since the observed pollinator behaviour should promote geitonogamous selfing, it

remains enigmatic why selfing has not evolved in all North American populations of *A. lyrata*.

Potential consequences of admixture

We found that F1 hybrid cross types had a similar phenology and pollinator visitation as

the parental cross types. Earlier studies have shown that hybrids between outcrossing and selfing

plants can be intermediate for phenological traits. For instance, in the genus *Clarkia,*

hybridization between selfing and outcrossing populations resulted in floral characteristics and

flowering times that were intermediate between the parental populations (Moore & Lewis 1965).

Our results show similar relationships between F1 hybrids and the parental populations in regard

to flowering probability and time of peak flowering. This suggests that in a scenario of secondary

contact, F1 hybrids would likely function as a bridge to further gene exchange between selfing

and outcrossing plants, which could potentially lead to the parental populations merging

(Rhymer & Simberloff 1996). Whether the resulting admixed populations will maintain a mixed

mating system (Goodwillie et al. 2005), or evolve to become predominantly selfing or

outcrossing remains to be tested. Initially, as inbreeding depression tends to be low (Carleial et al

2017a; Li et al, in press), selfing may be favoured due to the associated inherent transmission

advantage. However, on longer timescales, expression of drift load may select against selfing as

shown in selfing populations of *A. lyrata* (Willi et al. 2013b; but see Joschinski et al. 2015). It

would therefore be of interest to monitor the performance and mating system of admixed

populations over multiple years.

<u>Conclusion</u>

Our common garden experiment showed that, although pollinator behaviour may isolate

selfers by promoting geitonogamy, outcrossing and selfing *A. lyrata* populations are only weakly

reproductively isolated via pre-pollination mechanisms. These findings differ from findings in

other systems with a recent transition to selfing (e.g., Ramsey et al. 2003, Briscoe Runquist et al.

2014). The weak isolation between selfing and outcrossing populations of *A. lyrata* is likely

because its transition to selfing is even more recent, and has not led to evolution of a selfing

syndrome (Carleial et al. 2017b). Future studies could investigate if reproductive isolation due to

prezygotic pre-pollination mechanisms are larger in natural populations, giving specific attention

to parapatric selfing and outcrossing populations. Moreover, to what extent other mechanisms

such as niche differentiation and genetic incompatibilities contribute to reproductive isolation

remains to be investigated.

Acknowledgements

Christina Steinecke and Ryan Holt helped with data collection. Colleen White helped with

transplanting. Bill Crins taxonomically identified pollinators. Fränzi Korner-Nievergelt and Pius

Korner-Nievergelt (oikostat GmbH) helped with parts of the statistical analyses. Barbara Mable

provided the source material (seeds) from wild populations. Yan Li, Ekaterina Mamonova, and

Sina Konitzer-Glöckner helped with the crosses to produce the seeds used in the experiment.

This research was supported by the German Research Foundation (Deutsche

Forschungsgemeinschaft - Project number 388824194 to MS).

Authors Contributions

MS, MVK, MD, and CG conceived the ideas and designed methodology; CG and LB collected

the data; CG and MD analysed the data; CG led the writing of the manuscript. All authors

contributed critically to the drafts and gave final approval for publication.

Data Accessibility

Data will be made available from the Dryad Digital Repository upon acceptance.

References

Baack, E., Melo, M. C., Rieseberg, L. H., & Ortiz-Barrientos, D. (2015). The origins of

reproductive isolation in plants. *New Phytologist*, *207*(4), 968–984. doi: 10.1111/nph.13424

Baker, H. G. (1955). Self-compatibility and establishment after "long-distance" dispersal.

Evolution, *9*(3), 347–349. doi: 10.1111/j.1558-5646.1955.tb01544.x

Barrett, S. C. H. (2002). The evolution of plant sexual diversity. *Nature Reviews Genetics*, *3*(4),

274–284. doi: 10.1038/nrg776

Barrett, S. C. H. (2003). Mating strategies in flowering plants: The outcrossing-selfing paradigm

and beyond. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

358(1434), 991–1004. doi: 10.1098/rstb.2003.1301

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects

models using lme4. *Journal of Statistical Software*, *67*(1). doi: 10.18637/jss.v067.i01

Briscoe Runquist, R. D., Chu, E., Iverson, J. L., Kopp, J. C., & Moeller, D. A. (2014). Rapid

evolution of reproductive isolation between incipient outcrossing and selfing *Clarkia*

species. *Evolution*, *68*(10), 2885–2900. doi: 10.1111/evo.12488

Brunet, J., & Charlesworth, D. (1995). Floral sex allocation in sequentially blooming plants.

Evolution, *49*(1), 70–79. doi: 10.1111/j.1558-5646.1995.tb05959.x

Carleial, S., van Kleunen, M., & Stift, M. (2017a). Relatively weak inbreeding depression in

selfing but also in outcrossing populations of North American *Arabidopsis lyrata*. *Journal*

of Evolutionary Biology, *30*(11), 1994–2004. doi: 10.1111/jeb.13169

Carleial, S., van Kleunen, M., & Stift, M. (2017b). Small reductions in corolla size and pollen:

ovule ratio, but no changes in flower shape in selfing populations of the North American

Arabidopsis lyrata. *Oecologia*, *183*(2), 401–413. doi: 10.1007/s00442-016-3773-4

Coyne, J.A. & Orr, H.A. (2004) Speciation. Sinauer Associates, Inc., Sunderland, MA.

Darwin, C. (1876). The effects of cross- and self-fertilization in the vegetable kingdom. John

Murray, London, UK.

Fausto, J. A., Eckhart, V. M., & Geber, M. A. (2001). Reproductive assurance and the

evolutionary ecology of self-pollination in *Clarkia xantiana* (Onagraceae). *American*

Journal of Botany, *88*(1–12), 1794–1800. doi: 10.2307/3558355

Fisher, R. A. (1941). Average excess and average effect of an allelic substitution. *Annals of*

Eugenics, *11*(1), 53–63. doi: 10.1111/j.1469-1809.1941.tb02272.x

Foxe, J. P., Stift, M., Tedder, A., Haudry, A., Wright, S. I., & Mable, B. K. (2010).

Reconstructing origins of loss of self-incompatibility and selfing in North American

Arabidopsis lyrata: A population genetic context. *Evolution*, *64*(12), 3495–3510. doi:

10.1111/j.1558-5646.2010.01094.x

Gelman A., and J. Hill. (2007). Data analysis using regression and hierarchical/multilevel

models. Cambridge University Press, Cambridge.

Gervasi, D. D. L., & Schiestl, F. P. (2017). Real-time divergent evolution in plants driven by

pollinators. *Nature Communications*, *8*, 14691. doi: 10.1038/ncomms14691

Goodwillie, C., Kalisz, S., & Eckert, C. G. (2005). The evolutionary enigma of mixed mating

systems in plants: Occurrence, theoretical explanations, and empirical evidence. *Annual*

Review of Ecology, Evolution, and Systematics, *36*(1), 47–79. doi:

10.1146/annurev.ecolsys.36.091704.175539

Harder, L. D., & Barrett, S. C. H. (1995). Mating cost of large floral displays in hermaphrodite

plants. *Nature*, *373*(6514), 512–515. doi: 10.1038/373512a0

Harder, L. D., & Barrett, S. C. H. (1996). Pollen dispersal and mating patterns in animal-

pollinated plants. In *Floral Biology* (pp. 140–190). doi: 10.1007/978-1-4613-1165-2_6

Jackman S. (2017). pscl: Classes and Methods for R Developed in the Political Science

Computational Laboratory. United States Studies Centre, University of Sydney. Sydney,

New South Wales, Australia. R package version 1.5.2.

Jain, S. K. (1976). The Evolution of Inbreeding in Plants. *Annual Review of Ecology and*

Systematics, *7*(1), 469–495. doi: 10.1146/annurev.es.07.110176.002345

Joschinski, J., van Kleunen, M., & Stift, M. (2015). Costs associated with the evolution of selfing

in North American populations of *Arabidopsis lyrata*? *Evolutionary Ecology*, *29*(5), 749–

764. doi: 10.1007/s10682-015-9786-3

Karron, J. D., Holmquist, K. G., Flanagan, R. J., & Mitchell, R. J. (2009). Pollinator visitation

patterns strongly influence among-flower variation in selfing rate. *Annals of Botany*, *103*(9),

1379–1383. doi: 10.1093/aob/mcp030

Korner-Nievergelt, F., Roth, T., von Felten, S., Guélat, J., Almasi, B., & Korner-Nievergelt, P.

(2015). Bayesian Data Analysis in Ecology Using Linear Models with R, BUGS, and Stan.

London: Academic Press.

Li, Y., van Kleunen, M., & Stift, M. (2019). Sibling competition does not magnify inbreeding

depression in North American *Arabidopsis lyrata*. *Heredity*. doi: 10.1038/s41437-019-

0268-1

Lowry, D. B., Rockwood, R. C., & Willis, J. H. (2008). Ecological reproductive isolation of

coast and inland races of *Mimulus guttatus*. *Evolution*, *62*(9), 2196–2214. doi:

10.1111/j.1558-5646.2008.00457.x

Mable, B. K., Hagmann, J., Kim, S.-T., Adam, A., Kilbride, E., Weigel, D., & Stift, M. (2017).

What causes mating system shifts in plants? *Arabidopsis lyrata* as a case study. *Heredity*,

118(1), 52–63. doi: 10.1038/hdy.2016.99

Mable, B. K., & Adam, A. (2007). Patterns of genetic diversity in outcrossing and selfing

populations of *Arabidopsis lyrata*. *Molecular Ecology*, *16*(17), 3565–3580. doi:

10.1111/j.1365-294X.2007.03416.x

Mable, B. K., Robertson, A. V., Dart, S., Berardo, C. Di, & Witham, L. (2005). Breakdown of

self-incompatibility in the perennial *Arabidopsis lyrata* (Brassicaceae) and its genetic

consequences. *Evolution*, *59*(7), 1437–1448. doi: 10.1111/j.0014-3820.2005.tb01794.x

Martin, N. H., & Willis, J. H. (2007). Ecological divergence associated with mating system

causes nearly complete reproductive isolation between sympatric *Mimulus* species.

Evolution, *61*(1), 68–82. doi: 10.1111/j.1558-5646.2007.00006.x

Mitchell, R. J., Irwin, R. E., Flanagan, R. J., & Karron, J. D. (2009). Ecology and evolution of

plant–pollinator interactions. *Annals of Botany*, *103*(9), 1355–1363. doi:

10.1093/aob/mcp122

Moore, D. M., & Lewis, H. (1965). The evolution of self-pollination in *Clarkia xantiana*.

Evolution, *19*(1), 104. doi: 10.2307/2406299

R Core Team. (2019). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ramsey, J., Bradshaw, H. D., & Schemske, D. W. (2003). Components of reproductive isolation

between the monkeyflowers *Mimulus lewisii* and *M. cardinalis* (Phrymaceae). *Evolution*,

57(7), 1520–1534. doi: 10.1111/j.0014-3820.2003.tb00360.x

Rhymer, J. M., & Simberloff, D. (2002). Extinction by hybridization and introgression. *Annual*

Review of Ecology and Systematics, *27*(1), 83–109. doi: 10.1146/annurev.ecolsys.27.1.83

Rieseberg, L. H., & Willis, J. H. (2007). Plant Speciation 10.1126/science.1137729. *Science*,

317, 910–914.

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: A model for studying the genetic and

evolutionary basis of morphological adaptation in plants. *Annals of Botany*, Vol. 107, pp.

1433–1443. doi: 10.1093/aob/mcr023

Therneau, T. M., & Grambsch, P. M. (2000). Modeling survival data: extending the Cox model.

Springer, New York.

Venables, W. N., & Ripley, B. D. (2002). Modern Applied Statistics with S. Fourth Edition.

Springer, New York.

Widmer, A., Lexer, C., & Cozzolino, S. (2009). Evolution of reproductive isolation in plants.

Heredity, *102*(1), 31–38. doi: 10.1038/hdy.2008.69

Willi, Y., & Maattanen, K. (2011). The relative importance of factors determining genetic drift:

Mating system, spatial genetic structure, habitat and census size in Arabidopsis lyrata. *New*

Phytologist, *189*(4), 1200–1209. doi: 10.1111/j.1469-8137.2010.03569.x

Willi, Y. (2013a). The battle of the sexes over seed size: Support for both kinship genomic

imprinting and interlocus contest evolution. *The American Naturalist*, *181*(6), 787–798. doi:

10.1086/670196

Willi, Y. (2013b). Mutational meltdown in selfing Arabidopsis lyrata. *Evolution*, *67*(3), 806–815.

doi: 10.1111/j.1558-5646.2012.01818.x

Zeileis, A., Kleiber, C., & Jackman, S. (2008). Regression models for count data in R. *Journal of*

Statistical Software, *27*(8), 1–25. doi: 10.18637/jss.v027.i08

Tables

Table 1. Summary of results from the conditional logistic regression model that analyzed

whether *relative flower number*, *relative distance*, or *cross type* influenced the pollination path.

Cross type 'SI- within' was used as the baseline. Hoverflies: n=546, number of events = 142,

Likelihood ratio test= 62.16 on 7 df, p=<.001*. Solitary Bees: n=541, number of events=140,

Likelihood ratio test= 45.24 on 7 df, p=<.001*. Symbols and abbreviations used in the column

headings: SE= standard error; z= Wald statistic. For both hoverflies and solitary bees, relative

plant distance had the greatest influence on pollinator choice. There was no strong preference for

any of the cross types or individuals with more flowers. Significant effects are highlighted in

bold.

Visitor	Fixed effect	Odds Ratio	SE	z	Pr(>|z|)	95% confidence interval	
Solitary bees	Relative Flower Number	1.10	0.32	0.29	0.77	0.59	2.04
	Relative Distance	**0.11**	**0.38**	**-5.73**	**<.001***	**0.05**	**0.24**
	SC-within	2.01	0.39	1.80	0.07	0.94	4.30
	SIxSI	1.32	0.39	0.71	0.48	0.62	2.82
	SCxSC	0.96	0.39	-0.12	0.91	0.44	2.07
	SIxSC	0.80	0.36	-0.64	0.52	0.40	1.60
	SCxSI	0.91	0.38	-0.25	0.81	0.43	1.91
Hoverflies	Relative Flower Number	0.90	0.34	-0.32	0.75	0.46	1.75
	Relative Distance	**0.06**	**0.43**	**-6.56**	**<.001***	**0.03**	**0.14**
	SC-within	1.06	0.45	0.14	0.89	0.44	2.55
	SIxSI	1.09	0.41	0.21	0.83	0.49	2.44
	SCxSC	1.70	0.40	1.34	0.18	0.78	3.71
	SIxSC	1.03	0.40	0.07	0.95	0.47	2.27
	SCxSI	0.83	0.41	-0.44	0.66	0.37	1.86

609 * indicate p-values <.05. The 95% confidence interval is the confidence interval of the odds

610 ratio. *Relative flower number* and *relative distance* were estimated relative to the other

611 individuals in the observation. These variables were transformed to range between 0 and 1.

Figure captions

Figure 1: Map of the location of the common garden experiment in relation to the source populations. Gray circles represent selfing populations and black diamonds represent outcrossing populations. The black X represents the location of the common garden at Trent University, Peterborough, Ontario.

Figure 2: Panel of effect plots for the differences in phenological traits among the cross types (n=1512). a) The probability of flowering per cross type. Fitted values are obtained from the hurdle model. b) The day of peak flowering per cross type. This was calculated as the experimental day where each individual had the highest number of flowers during the total flowering period. Fitted values obtained are from the Gaussian linear model that included *day of peak flowering* as the response variable. c) The flowering duration (days) per cross type. Fitted values obtained are from the Gaussian linear model that included *flowering duration* as the response variable. Vertical bars in all figures represent the 95% credible intervals. SI-within= within outcrossing population cross type, SC-within= within selfing population cross type, SC-self= selfed, SI= self-incompatible, SC= self-compatible. The probability of flowering and time of peak flowering varied among the cross types. Flowering duration did not strongly differ among the cross types.

Figure 3: Differences in pollinator visitation (number of visits) by a) solitary bees and b) hoverflies among the cross types (n=502). The fitted values obtained are from the generalized linear models with negative binomial distributions described in the methods. Vertical bars represent the 95% credible intervals. SI-within= within outcrossing population cross type, SC-within= within selfing population cross type, SI= self-incompatible, SC= self-compatible. The

number of visits did not strongly differ among the cross types and pollinator identity had a minor

influence on the number of visits. Solitary bees made slightly more visits overall.

Figure 4: a) Stacked bar plot representing the probability of pollinators making a choice to visit

an individual of any cross type after visiting an individual of a certain cross type. Probabilities

were obtained from the multinomial model (and thus corrected for the number of available

flowers on each plant in the array). Pollinators could also choose to visit a flower on the same

plant ("same") or to leave the experimental setup ("away"). The cross types on the x-axis

represent the cross type of the initial visit. The stacked blue bars represent the probability of that

cross type being selected after a visit to the cross type on the x-axis. SI-within= within

outcrossing population cross type, SC-within= within selfing population cross type, SI= self-

incompatible, SC= self-compatible. b) Graphical representation of the probabilities of pollinator

choice. Ranges represent the range of probabilities for that specific choice. Arrow thickness

corresponds to probability. Base picture used for figure modified from Wiki Commons: DBCLS

TV [CC BY 4.0 (https://creativecommons.org/licenses/by/4.0)]

Figure 5: Representative bootstrapped run of the distribution of the self- and outcross pollen

transfer opportunities for self-compatible and self-incompatible individuals. The parameter

estimate for the difference in transfer-probability for the shown run was 0.024, and thus close to

the mean value of 0.020 across all the bootstrapped samples. There were no significant

differences in the opportunities for pollen transfer.

Figure 1

 Figure 2

a)

b)

c)

 Figure 3

671 **Figure 4**

672 **a)**

674 **b)**

676 **Figure 5**

677