Dendritic Morphology and Inhibitory Regulation Distinguish Dentate Semilunar Granule Cells from Granule Cells through Distinct Stages of Postnatal Development

Akshay Gupta[1,3], Deepak Subramanian[1,3], Archana Proddutur[1,3], Yun-Juan Chang[2], Vidhatri Raturi[1], Jenieve Guevarra[1], Yash Shah[1], Fatima S. Elgammal[1], and Vijayalakshmi Santhakumar[1,3]

[1]Department of Pharmacology, Physiology and Neuroscience, [2]Office of Advance Research Computing, Rutgers New Jersey Medical School, Newark, New Jersey 07103, [3]Department of Molecular, Cell and Systems Biology, University of California Riverside, California 92521.

ORCHID ID: Gupta A (0000-0002-3788-6277); Subramanian D (0000-0001-7109-6714); Proddutur A (0000-0003-2333-4188); Chang Y-J (0000-0003-2688-6391); Santhakumar V (0000-0001-6278-4187)

Short Title SGCs during postnatal development

Word count for Abstract: 249   Number of pages: 41   Number of figures: 7

Declarations:

Funding: The project was supported by National Institutes of Health/National Institute of Neurological Diseases and Stroke R01 NS069861, R01NS097750 and New Jersey Commission on Brain Injury Research CBIR16IRG017 to V.S. and, CBIR11FEL003 to A.G

Conflicts of interest/Competing interests: No conflicts or competing interests to discuss

Availability of data and material: All reconstructions will be uploaded to Neuromorpho.org. Physiology data will be shared upon request.

Code availability: Standard software packages, Neurolucida 360, pClamp10 and R were used in analysis. Custom IGOR-Pro code for IPSC analysis will be made available upon request.

Authors' contributions: AG, DS, JG, and AP performed experiments; AG, D.S, Y-JC, YS, and VR analyzed data; AG, DS, FSE, and VS interpreted results of experiments; AG, AP, and DS prepared figures; AG and VS conceived of and designed research; AG, DS, and VS drafted manuscript.

Acknowledgements: We thank Dr. Luke Fritzky at the Rutgers Imaging Core for help with imaging and Dipika Sekhar for data entry. We thank Dr. Kelly A. Hamilton for thoughtful discussions and comments.

Correspondence: **Vijayalakshmi Santhakumar**, PhD
                     Department of Molecular, Cell and Systems Biology
                     University of California, Riverside

E-mail: vijayas@ucr.edu

Abstract

Semilunar granule cells (SGCs) have been proposed as a morpho-functionally distinct class of

hippocampal dentate projection neurons contributing to feedback inhibition and memory

processing in juvenile rats. However, the structural and physiological features that can reliably

classify granule cells (GCs) from SGCs through postnatal development remain unresolved.

Focusing on postnatal days 11-13, 28-42, and >120, corresponding with human infancy,

adolescence, and adulthood, we examined the somatodendritic morphology and inhibitory

regulation in SGCs and GCs to determine the cell-type specific features. Unsupervised cluster

analysis confirmed that morphological features reliably distinguish SGCs from GCs irrespective

of animal age. SGCs maintain higher spontaneous inhibitory postsynaptic current (sIPSC)

frequency than GCs from infancy through adulthood. Although sIPSC frequency in SGCs was

particularly enhanced during adolescence, sIPSC amplitude and cumulative charge transfer

declined from infancy to adulthood and were not different between GCs and SGCs.

Extrasynaptic GABA current amplitude peaked in adolescence in both cell types and was

significantly greater in SGCs than in GCs only during adolescence. Perforant-path evoked

dentate population responses *in vivo,* conducted to assess the circuit level changes in dentate

inhibition during development, revealed greater paired-pulse depression during adolescence. The

data delineate the structural features that can reliably distinguish GCs from SGCs through

development. The results reveal developmental differences in passive membrane properties and

steady state inhibition between GCs and SGCs which could confound their use in classifying the

cell types and identify parallel developmental time courses for regulation of GC and SGC tonic

GABA currents and dentate feedback inhibition.

Keywords: inhibition, GABA, dentate gyrus, extrasynaptic, development, granule cell,
semilunar granule cell

The dentate gyrus, the primary gateway for cortical inputs to the hippocampus, plays a unique

role in memory processing as a center for sparse coding, mediated in large part, by strong

inhibitory filtering of activity (Dengler and Coulter 2016). The classical dentate projection

neurons, granule cells (GCs), which number in over a million in the rat brain, rest at more

hyperpolarized potentials than most hippocampal neurons, receive powerful feedback inhibition,

and project strong "detonator" synapses to CA3 enabling sparse yet reliable transmission

(Dengler and Coulter 2016; Engel and Jonas 2005). Recent characterization of semilunar granule

cells (SGCs), a dentate cell-type originally identified by Ramón y Cajal (1953), has revealed a

second class of dentate projection neurons that are located in the inner molecular layer (IML),

has distinctive somato-dendritic structure, persistent firing, and robust inhibition (Gupta et al.

2012; Larimer and Strowbridge 2010; Williams et al. 2007). While SGCs have been

physiologically defined in rat (Gupta et al. 2012; Williams et al. 2007), cells with dendritic

morphology consistent with SGCs have been observed in primates (Duffy and Rakic 1983;

Seress and Frotscher 1990). Additionally, physiological recordings from cells with

morphological features of SGCs have been performed in mouse and rabbit (Sancho-Bielsa et al.

2012; Save et al. 2018), indicating their presence across mammalian species. Due to their unique

persistent firing in response to inputs, SGCs have been proposed to play a crucial role in

regulating GC feedback inhibition, sparse coding of inputs, and pattern separation (Larimer and

Strowbridge 2010; Walker et al. 2010). Recent studies suggest that SGCs are preferentially

involved in memory engrams (Erwin et al. 2020). Since SGCs lack a cell-specific neurochemical

marker, morphological characteristics, and lower input resistance (R_{in}) are currently the primary

approach to identify SGCs. However, the limited literature on SGC morphology and synaptic

inputs during development have impeded further analysis of the role of SGCs in the dentate

circuit.

Developmentally, both SGCs and GCs express the homeodomain transcription factor Prox1

(Gupta et al. 2012) and derive from the same neural precursor pool, although SGCs differentiate

in a more restricted embryonic phase, demonstrating a common lineage between SGCs and GCs

(Kerloch et al. 2018; Save et al. 2018). Based on the embryonic developmental analysis, SGCs

are estimated at 3% of the neurons generated from the dentate neurogenic pool from which GCs

derive (Save et al. 2018). Like GCs, SGCs have dendrites in the dentate molecular layer, and

axons projecting to CA3. However, they can be distinguished from GCs based on their expansive

dendritic span, large semi-lunar somata in the inner molecular layer (IML), and frequent

presence of IML axon collaterals (Gupta et al. 2012; Williams et al. 2007). In earlier studies in

rats, we defined SGCs on the basis of their wider dendritic angle (Gupta et al. 2012), which was

subsequently confirmed in mice (Save et al. 2018). To date, morphometric analysis of SGCs has

been limited to a narrow window of postnatal day (PD) 14-42 in rats, which is consistent with the

neurological developmental state of human adolescence (Semple et al. 2013; Sengupta 2013), or

in an embryonically labeled subgroup of neurons in young adult mice (Save et al. 2018).

However, whether SGCs retain their distinct structural characteristics through postnatal

development has not been examined. Recent studies have suggested layer specific differences in

GC morphology (Kerloch et al. 2018) and disease-related changes in GC dendritic features

(Freiman et al. 2011; Llorens-Martin et al. 2015; von Campe et al. 1997). However, without

knowing the specific structural features that distinguish GCs from SGCs, it is difficult to

interpret whether reports on changes in GC morphology reflect inclusion of structurally distinct

SGCs in the datasets. Moreover, the ability to reliably distinguish SGCs from GCs across animal

age is a prerequisite to elucidating the unique role that SGCs play in dentate processing.

Unsupervised analysis of dendritic morphometric features is ideally suited to examine whether

SGCs remain distinct from GCs through development and to elucidate the somato-dendritic

features that specify the cell types.

Physiologically, SGCs show prolonged firing, reduced spike frequency adaptation, and lower R_{in}

than GCs (Gupta et al. 2012; Save et al. 2018; Williams et al. 2007). Unlike GCs, SGCs respond

to perforant-path stimulation with persistent firing lasting several seconds, which correlates with

periods of increased hilar activity termed "up-states" (Larimer and Strowbridge 2010). SGCs

have larger NMDA currents (Williams et al. 2007) than GCs, make synaptic contacts with hilar

mossy cells and interneurons and have been proposed to drive granule cell feedback inhibition

(Larimer and Strowbridge 2010). Apart from excitation, we previously demonstrated that SGCs

have greater synaptic and extrasynaptic GABA currents (Gupta et al. 2012). Additionally, we

identified that GCs and SGCs have diametrically opposite changes in synaptic and tonic GABA

currents after brain injury suggesting that differences in inhibition which could contribute to

distinct roles for SGCs and GCs in the dentate circuit. GCs undergo changes in both tonic and

synaptic GABA currents during postnatal development (Hollrigel et al. 1998; Holter et al. 2010;

Lee and Liou 2013). In parallel, studies on adult-born GCs labeled at specific time points show

cell-age dependent maturation of GABA currents, ultimately reaching the levels similar to

mature GCs in age-matched animals (Dieni et al. 2012; Li et al. 2012). Yet, whether inhibition

in SGCs remains distinct from mature GCs through postnatal development is not known. While

GCs show prolonged functional maturation (Yu et al. 2013a), age-dependent changes in

inhibitory regulation of SGCs remains to be tested. This study was conducted to explicitly to

determine the characteristic structural features which can be used to distinguish SGCs from GCs

through postnatal developmental stages representing infancy (PD 11-13, prior to rodent eye

opening), adolescence (PD 28-42 days, period of cortical maturation), and adult (> PD 120,

cortical and sexual maturity) stages of development (Semple et al. 2013) and to identify the

differences in steady-state inhibitory regulation of the two cell types through maturation of the

dentate circuit.

Materials and Methods

Materials and Methods

Animals. All experiments were performed in accordance with IACUC protocols approved by

Rutgers-NJMS, Newark, NJ, and the University of California at Riverside, CA and in keeping

with the ARRIVE guidelines. The study included Wistar rats ranging in age from 11-13 days,

28-42 days, and 120-180 days designated as infancy, adolescence, and adulthood, respectively

(Semple et al. 2013; Sengupta 2013). Due to the potential effects of hormonal variation on

GABA currents, slice recordings in rats from >28 days were restricted to males. A subset of

recordings was derived from surgical or saline-injected controls for an independent study and

was pooled with data from naïve rats from which they showed no statistical difference.

Slice Physiology. Rats were anesthetized with isoflurane and decapitated. Horizontal brain

slices (300μm) were prepared in ice-cold sucrose artificial cerebrospinal fluid (sucrose-aCSF)

containing the following (in mM): 85 NaCl, 75 sucrose, 24 $NaHCO_3$, 25 glucose, 4 $MgCl_2$, 2.5

KCl, 1.25 NaH_2PO_4, and 0.5 $CaCl_2$ using a Leica VT1200S Vibratome. The slices were sagittally

bisected and incubated at 32°C for 30 min in a submerged holding chamber containing an equal

volume of sucrose-aCSF and recording aCSF, and subsequently were held at room temperature.

The recording aCSF contained the following (in mM): 126 NaCl, 2.5 KCl, 2 $CaCl_2$, 2 $MgCl_2$,

1.25 NaH_2PO_4, 26 $NaHCO_3$, and 10 D-glucose. All solutions were saturated with 95% O_2 and

5% CO_2 and maintained at a pH of 7.4 for 1–6 h. Slices were transferred to a submerged

recording chamber and perfused with oxygenated aCSF at 33°C. Whole-cell voltage-clamp

recordings from neurons in the dentate granule cell layer (GCL) and inner molecular layer (IML)

were performed using infrared differential interference contrast visualization techniques (Gupta

et al. 2012; Yu et al. 2016) with a Nikon Eclipse FN-1 microscope, using a 40X, 0.80 NA water-

immersion objective. Cells in the hilar-GCL border were not included in the study. Recordings

were obtained using MultiClamp 700B (Molecular Devices). Data were low pass filtered at 3

kHz, digitized using Digidata 1440A, and acquired using pClamp10 at 10 kHz sampling

frequency. Tonic and synaptic GABA currents were recorded in aCSF with no added GABA or

GABA transporter antagonists in the recording solution (Gupta et al. 2012; Yu et al. 2013b).

Voltage-clamp recordings of inward GABA currents were obtained from a holding potential of -

70 mV using microelectrodes (5–7 MΩ) containing (in mM): 125 $CsCl_2$, 5 NaCl, 10 HEPES, 2

$MgCl_2$, 0.1 EGTA, 2 Na-ATP, and 0.5 Na-GTP, titrated to a pH 7.25 with CsOH. Biocytin

(0.2%) was included in the internal solution for post hoc cell identification, and the glutamate

receptor antagonist kynurenic acid (3 mM KynA, Tocris Bioscience) was included in the external

solution to isolate GABA currents. Passive membrane properties including resting membrane

potential and R_{in} were recorded in current clamp using an internal containing (in mM) 126 K-

gluconate, 4 KCl, 10 HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 Phosphocreatine and 0.2% biocytin

(Gupta et al. 2012). Among neurons recorded in the IML, only cells showing obtuse dendritic

angles, presence of dendritic spines and axon projecting to the hilus (Gupta et al. 2012)

(Supplementary Figure 1) were included in the physiological analysis as putative SGCs. The

somata of all GCs included in the analysis were located in the GCL. Recordings were

discontinued if series resistance increased by >20%. Access resistance was not different between

cell-types or between developmental groups. Following establishment of whole cell mode,

baseline recordings were obtained for a minimum of five minutes in the presence of KynA prior

to addition of GABA blockers. Recordings with excessive baseline fluctuations during the 3-5

minutes of recordings in KynA were discarded. All salts were purchased from Sigma-Aldrich

(St. Louis, MO). Tonic GABA current, the steady-state current blocked by the $GABA_AR$

antagonist bicuculline methiodide (BMI, 100µM, Sigma-Aldrich) or gabazine (SR95531, 10 µM,

Sigma-Aldrich), was measured as described previously (Gupta et al. 2012; Yu et al. 2013b) using

custom macros in IgorPro7.0 software (WaveMetrics). Following physiological recordings,

slices were fixed in 0.2mM phosphate buffer containing 4% paraformaldehyde at 4°C for 2 days.

Biocytin staining was revealed using Alexa Fluor 594-conjugated streptavidin (Gupta et al. 2012;

Swietek et al. 2016).

Morphometry and hierarchical cluster analysis. Sections were visualized and imaged using a

Zeiss LSM 510 confocal microscope with a 20X, 0.5 NA objective. Cell reconstructions from

confocal image stacks were performed using the directional kernels user-guided reconstruction

algorithm in Neurolucida 360 (MBF Bioscience) followed by manual correction and validation

in 3D. About 10-20 percent of each dendritic arbor was reconstructed manually. Neurolucida

Explorer (MBF Biosciences) was used to extract non-nominal or non-ordinal somatodendritic

morphological quantitative parameters (defined in Supplementary Tables 1 & 2) for use in

statistical comparisons and hierarchical cluster analysis. A total of 42 projection neurons in

which the dendritic arbors were fully reconstructed were analyzed. The software failed to

quantify a subset of somatic parameters in 6 cells, and these cells were excluded from cluster

analysis.

Data were tested for uniform distribution and each quantified variable was fit to the sum of two

or more Gaussian functions and quality of fit determined using maximum likelihood analysis

(MLA; v2test) to assess normal distribution of parameters within each cell type. Variables with a

nonuniform distribution were used for subsequent cluster analysis. A total of 42 somato-dendritic

parameters (Supplementary Tables 1 & 2) from 36 morphologically reconstructed neurons,

including both features measured in Neurolucida Explorer from the 3D reconstructions and

parameters measured manually in 2D rendering (Neurolucida 360, MicroBrightfield) were

analyzed. Unsupervised clustering and principal component analyses of morphological properties

were conducted within R version 3.5.0, using R package Cluster, FactoMineR, and factoextra by

an investigator (Y-J C) blinded to cell types and age groups. Hierarchical clustering on the

selected principal components (PCs) was performed using Ward's Criterion with Euclidean

distance to generate the dendrogram. The clustering partition was obtained from hierarchical

clustering and improved with K-means method (Husson et al. 2010). Summary of morphological

data and statistical analysis of morphometric parameters are presented in Supplementary Tables

3 and 4, respectively.

In Vivo Recordings: Animals were anesthetized using urethane (1.5g/kg, i.p.) and all recordings

commenced 60-90 minutes post induction following confirmation of surgical anesthesia. Rats

were secured in a stereotaxic frame and placed on a warming pad (Stoelting) to maintain optimal

body temperature (37ºC) throughout the experiments. Platinum-Iridium Concentric bipolar

stimulating electrodes (Microprobes) were used to stimulate medial perforant path (mPP) (AP: -4

mm, ML: +3mm, DV: -1.5mm from bregma for infant, ML: +4-4.2 mm, DV: -2mm from lambda

for adolescent and adult). A single tungsten wire electrode (50µm diameter, California Wire

Company) was positioned in the GCL guided by responses evoked by mPP stimulation (0.6mA,

150µs, 0.1Hz monophasic pulses; ISOflex, AMPI Israel). Coordinates used for dentate gyrus

were, from bregma, AP: -2 mm, ML: +1.5 mm, DV: -2.2-2.5mm for infant, AP: -3.5 mm, ML:

+2 mm, DV: -3mm for adolescent, and AP: -4 mm, ML: +2mm, DV: -3.5mm for adult animals.

Final electrode positions were optimized for maximal evoked responses. Paired-pulse responses

were evoked with a 20ms interval between pulses, averaged over 5 sweeps each (0.8-1mA, 150

µs, 0.1Hz, monophasic pulse). Current intensity was adjusted to levels that reliably produced

population spike amplitude of 0.5mV or more. Adolescent and adult animals were transcardially

perfused with 0.9% saline followed by 4% PFA and brains were extracted for histological

confirmation of electrode track. Population spike amplitude was measured as detailed previously

(Neuberger et al. 2017). Paired pulse ratio was calculated as the amplitude of population spike 2

/population spike1. Evoked responses were recorded and analyzed using LabChart 8.0 (AD

Instruments). Data from animals in which the paired pulse ratio was >1, indicative of stimulating

electrode in the lateral perforant path, were excluded.

Analysis and Statistics: Individual spontaneous inhibitory postsynaptic currents (sIPSCs) were

detected using custom software in Igor-Pro7.0 (Gupta et al. 2012). The investigator (AG) was

blinded to cell type during analysis. Events were visualized, and any "noise" that spuriously met

trigger specifications was rejected. Cumulative probability plots of sIPSC amplitude and

frequency were constructed using IgorPro by pooling an equal number of sIPSCs from each cell.

Kinetics and charge transfer were calculated from the averaged trace of all accepted sIPSC

events. Rise time was measured as the time for amplitude to change from 20 to 80 % of peak.

Amplitude weighted τ_{decay} was calculated from a two-exponential fit to the IPSC decay. sIPSC

charge transfer was calculated as the area under the curve of the baseline adjusted average sIPSC

trace. The summed sIPSC charge transfer over one second was calculated as the product of the

sIPSC charge transfer and sIPSC frequency for each cell. R_{in} was calculated as the slope of the

current voltage plots with voltage response (averaged over the last 400 ms) to one second current

injections from -200pA to -40 pA (in 40 pA steps) in each cell. Membrane time constant

($\tau_{membrane}$) was obtained from a single exponential fit to the voltage response to a -200pA current

injection. Data are shown as mean±SEM (standard error of the mean) or median and interquartile

range (IQR) where appropriate and presented in Supplementary Table 5. Kruskal-Wallis One

Way Analysis of Variance on ranks was conducted on data which failed Shapiro Wilk test for

normalcy or equal variance test.   Two-way ANOVA (TW-ANOVA, SigmaPlot 12.3) and

Bonferroni correction was used to test for statistical differences in tonic GABA currents.

Summary data and details of the statistical tests used are included in Supplementary Tables 5-8.

Sample sizes were not predetermined and conformed with those employed in the field.

Significance was set to $p < 0.05$, subject to appropriate Bonferroni correction. All custom macros

for analysis and data sets will be made available on request upon publication of the study.

Results

Somato-dendritic characteristics distinguish SGCs from GCs throughout postnatal

development.

To examine whether SGCs are structurally distinct from GCs throughout postnatal development,

we undertook electrophysiological recordings from putative GCs and SGCs in the GCL and IML

respectively, and filled cells with biocytin to recover their morphology. As illustrated by

morphological reconstructions obtained from confocal images of biocytin filled neurons (Fig. 1),

putative GCs exhibit compact molecular layer dendritic arbors while putative SGCs recorded in

the IML consistently exhibited wider dendritic spread. These differences in dendritic spread are

maintained throughout the three developmental stages tested. Additionally, the X-Y plane

projection of the reconstructed neurons revealed the characteristic crescent shaped somata of

SGCs. Figure 1 illustrates pseudo color rendering of dendritic arbors with segments assigned on

the basis of branch order in 3D reconstructions in Neurolucida 360. Color coding of the

dendritic segments revealed that, in addition to differences in dendritic span, SGCs differed from

GCs in the extent to which individual dendritic trees branched and the number of segments in

different dendritic orders. Moreover, reconstructions suggest that GCs have a dense distribution

of dendritic arbors in a compact volume while SGCs appear to have sparsely distributed

dendrites in a larger volume. Thus, while 3D reconstructions demonstrate that SGCs maintain a

qualitative pattern of dendritic arborization distinct from GCs at all postnatal ages examined,

they reveal additional morphometric features which differ between cell types indicating a need

for comprehensive unsupervised *quantitative* analysis of dendritic structure.

To determine if somato-dendritic parameters distinguish GCs from SGCs through development,

we undertook the first unsupervised classification of dentate projection neurons on the basis of

morphometry. Projection neurons were identified based on the presence of somata in the GCL or

IML, axons with boutons entering the hilus and targeting CA3 (Supplementary Figure 1).

Additionally, we find that both SGCs and GCs, have a high density of dendritic spines

(Supplementary Figure 1), which we used as a criterion to distinguish projection neurons from

local interneurons when the axon was not fully recovered. Morphometric parameters of the cells

reconstructed in 3D were obtained from automated algorithms in Neurolucida 360 (definitions

are included in Supplementary Tables 1 and 2). Principal Component Analysis (PCA) of 42

distinct morphometric parameters from 36 cells revealed a relatively high dimensional structure.

The first three principal components (PCs) explained about 46% of the total variance in the data,

while the first seven components retained over 85% of the variance (Supplementary Figure 2).

PCA analysis of individual cells was projected on to the first three principal components and

visualized in 3D representation with confidence interval (CI) ellipsoid set to 0.95, which

suggested that the cells likely segregate by cell type (Fig. 2A) rather than developmental age of

the rat (Fig. 2B).

Interestingly, clustering by PCA matched that of the investigator (Fig. 2). A total of 36 dentate

projection neurons (16 cells in infant, 10 cells in adolescent and 10 cells in adult), recovered

based on the quantitative morphological features, were included in the clustering analysis. A

hybrid approach of hierarchical clustering on principal components (HCPC) which helps in de-

noising multidimensional dataset was adopted (Husson et al. 2010). HCPC on the first seven

components suggested two cluster partitioning groups (Fig 2C). Opening of the blinding revealed

that the two putative clusters demonstrated a tendency for cells to cluster by putative cell-type.

Cluster 1 included 16 cells classified as GCs and five classified as an SGC by the investigator.

Clusters 2 consisted entirely of cell classified as SGCs by the investigator (A.G).

Identity of principal morphometric parameters that best represent the PCs

The PCs were examined further to identify the top five morphological variables which best

represent the PC ($cos2 \geq 0.7$) and the morphological features that contributed to PC variability

(Supplementary Figure 2). The number of first, second, and third order dendritic segments (note:

number of first order dendritic segments equals number of primary dendrites), number of second

order nodes and soma width where the top five morphological features that best represented the

PCs while dendritic area and volume contributed most to variability (Supplementary Figure 2).

Although the dendritic span and angle, used by the experimenter for classification (Gupta et al.

2012; Williams et al. 2007), contributed to the PC, morphology-based clustering revealed

additional salient parameters including number of primary dendrites and dendritic segmentation

which differed between clusters. Interestingly, one of the cells classified as SGCs by the

investigator which clustered with putative GCs by unsupervised clustering was located in the

IML (a feature that was not included in the cluster analysis), had a wide dendritic angle, but had

more than one primary dendrite, typically seen in GCs, with complex pattern of branching which

could have driven the clustering with GCs (Supplementary Figure 2D). Importantly, comparison

of cell classification based on the unsupervised approach to that of the investigator (A.G.)

confirmed that the investigator and PCA-based classifier agreed on the "grouping" of >85% (31

of 36) of cells examined. These findings confirm that GCs and SGCs are structurally distinct and

demonstrate that the investigator can reliably discriminate the cell-types.

Developmental changes in somato-dendritic morphology of dentate projection neurons

Using the investigator-assigned classifications, we next examined which specific morphological

parameters showed cell-type and developmental differences. First, we focused on the parameters

that contributed to PCs underlying cell classification. As predicted based on the PCA, the

number of primary dendrites (first order segments), second order segments and nodes were

different between SGCs and GCs, with SGCs having significantly more dendritic segments in the

first three orders of dendrites (Fig. 3A, Supplementary Figure 3 & Supplementary Tables 3 &4).

However, the effect of *age* on the number of segments in each order was not statistically

significant (Supplementary Figure 3A&B, Supplementary Tables 3 & 4). Similarly, soma width

and dendritic angle were greater in SGCs and failed to show age related changes (by TW-

ANOVA; Fig. 3B&C, and Supplementary Tables 3 & 4). Thus, these parameters are ideally

suited to distinguish cell-types regardless of age. In contrast, dendritic length showed a

significant effect of age yet was not different between cell types (Fig. 3D, and Supplementary

Tables 3 & 4). Of note, SGCs had significantly lower dendritic complexity than GCs and failed

to show the age-dependent increase in complexity observed in GCs (Fig. 3E, and Supplementary

Tables 3 & 4). Additional parameters that reflected 3D dendritic structure including convex hull

surface area showed significant differences between cell-type and with age (Fig. 3F,

Supplementary Tables 3 & 4). The convex hull surface area increased with development from

infancy to adolescence and remained at adolescent levels in the adult (Fig. 3F, and

Supplementary Tables 3 & 4) and appeared to contribute substantially to variability in the PCA

analysis of pooled morphometric dataset (Supplementary Figure 2B). Finally, certain summed

dendritic parameters including total numbers of dendritic terminals (ends), nodes, and segments

showed neither cell-type nor age related differences (Supplementary Figure 3D-F,

Supplementary Tables 3 & 4).

Developmental changes in spontaneous synaptic inhibition to GCs and SGCs

We previously demonstrated that SGCs from adolescent rats receive a greater frequency of action-potential driven sIPSCs than GCs from age-matched rats (Gupta et al. 2012). To determine if SGC inhibitory drive changes through postnatal development, we recorded sIPSCs from SGCs and GCs at three developmental stages. As illustrated in Fig. 4, SGCs consistently showed a higher frequency of sIPSCs compared to GCs from age matched rats. Both cell-types showed changes in sIPSC frequency with age. The frequency of sIPSCs in GCs increased from infancy through adolescence, peaked at adolescence and showed a slight, yet, significant decrease in adults. (Fig. 4D, Supplementary Figure 4 and Supplementary Tables 5 & 6). Despite the decrease from adolescence to adulthood, the sIPSC frequency in GCs from adults was higher than that in infancy. SGCs, on the other hand, showed a distinct peak in sIPSC frequency during adolescence with a significant reduction in frequency in adults, back to the levels observed in infancy (Fig. 4D, Supplementary Figure 4, and Supplementary Tables 5 & 6). Thus, despite being consistently higher than in GCs, sIPSC frequency in SGCs appears to show a specific and transient enhancement during adolescence.

Unlike sIPSC frequency, SGC sIPSC amplitude did not differ from age-matched GCs in infant and adolescent rats (Fig. 5C and Supplementary Tables 5 & 6), consistent with our earlier reports in adolescent rats (Gupta et al. 2012). However, in adult rats, sIPSC amplitude in GCs was larger than in SGCs (Fig. 5C). In GCs, sIPSC amplitude decreased significantly from infancy to adolescence and remained constant thereafter with differences in sIPSC between adolescence and adulthood not reaching statistical significance (Fig. 5C, Supplementary Tables 5 & 6). sIPSC

amplitude in SGCs decreased through postnatal development with a significant reduction from

infancy to adolescence, and a further decline from adolescence to adulthood (Fig. 5C,

Supplementary Tables 5 & 6).

Since distal inhibitory inputs can attenuate to a smaller amplitude at the soma (Soltesz et al.

1995), we sought to assess whether changes in the proportion of proximal versus distal dendritic

synaptic inputs could contribute to developmental change in GC and SGC sIPSC amplitude. To

determine if there is a systematic change in the amplitude of sIPSCs with age, we assigned

sIPSCs to two groups based on their amplitudes: large and putative proximal and perisomatic

(>50pA), and small (<50pA), potentially dendritic and calculated the proportion of these events

classes in GCs and SGCs during development. Consistent with the developmental increase in

dendritic length (Fig.3D), the proportion of large amplitude, presumed perisomatic events

(>50pA) were highest in infants and reduced progressively with age while the smaller amplitude

events <50pA events increased with age in both the cell types (Supplementary Figure 5). These

findings suggest that developmental increase in dendritic length may contribute to decline in

sIPSC amplitude over age in both cell types.

Systematic analysis of sIPSC kinetics identified an overall decline in $\tau_{\text{decay-WT}}$ in GCs with a

significant reduction from infant to adolescent rats (Fig. 5D, Supplementary Table 5 & 7).

Similarly, 20-80 rise time in GCs showed an overall decline with age with a significant reduction

from infancy to adulthood (Fig. 5E, Supplementary Table 5 & 7). In contrast, both sIPSC 20-80

rise time and $\tau_{\text{decay-WT}}$ increased from infancy through adulthood in SGCs demonstrating a

divergence in the effect of development on sIPSC kinetics between GCs and SGCs. Since the

developmental changes in both sIPSC rise and $\tau_{\text{decay-WT}}$ showed parallel trends within a given cell

type while differing between cell types, we evaluated whether developmental changes in cellular

passive membrane parameters may underlie these changes. Consistent with the changes observed

in sIPSC kinetics in GCs, both R_{in} and membrane time constant ($\tau_{membrane}$) trended to decline

from infancy through adulthood with the decrease in R_{in} reaching statistical significance

(Supplementary Fig. 6, Supplementary Tables 5 & 8). As with sIPSC kinetics in SGCs, both

R_{in} and $\tau_{membrane}$ increased significantly from infancy through adulthood (Supplementary Fig. 6,

Supplementary Tables 8). While direct correlation of the sIPSC kinetics with $\tau_{membrane}$ was not

feasible due to use of different cohorts of recordings for sIPSC and passive parameter data, the

results are consistent with a role for developmental changes in membrane passive properties

contributing to the shift in sIPSC kinetics in cell types with postnatal development. Moreover,

the data demonstrate that postnatal development has an opposite effect on R_{in} in GCs and SGCs.

R_{in} in SGCs was lower than in GCs in infant and adolescent rats, a feature that has been used to

distinguish the cell types (Erwin et al. 2020). However, in adult rats, R_{in} in SGCs was

significantly higher than age matched GCs (Supplementary Fig. 6, Supplementary Tables 5 & 8)

revealing that a lower R_{in} cannot consistently distinguish SGCs from GCs across all

developmental time points.

Given the developmental and cell-specific changes in sIPSC peak amplitude and kinetics, we

examined whether GCs and SGCs showed changes in sIPSC charge transfer during postnatal

development. Consistent with changes in peak amplitude, sIPSC charge transfer declined with

development in both cell types (Fig. 5F, Supplementary Tables 5 & 7). Despite the changes in

kinetics, the net charge transfer was not different between GCs and SGCs from age-matched rats

(Fig. 5F, Supplementary Tables 5 & 6). To comprehensively assess spontaneous synaptic

inhibition in GCs and SGCs during development in a manner that would include divergent

changes in frequency amplitude and kinetics between cell-types and during development, we

estimated the cumulative sIPSC charge trasnfer over one second as a product of the sIPSC

frequency and charge transfer for each cell. Interestingly, the cumulative sIPSC charge transfer

over one second was maximum during infancy and declined with development in both GCs and

SGCs (Supplementary Fig. 4C, Supplementary Table 5 & 7). Moreover, the cumulative sIPSC

charge transfer over one second was not different between cell-types at any developmental stage

examined (Supplementary Fig. 4C, Supplementary Table 5 & 7).

Extrasynaptic GABA currents in SGCs peak during adolescence

Apart from GABAergic synaptic inputs, dentate GCs are known to express extra- and peri-

synaptic $GABA_A$ receptors that contribute to steady-state tonic GABA currents (Stell et al.

2003). We previously demonstrated the presence of tonic GABA currents in SGCs and identified

that the amplitude of tonic GABA current in SGCs was greater than in age-matched adolescent

GCs (Gupta et al. 2012). Here we find that although tonic GABA current amplitude in SGCs,

measured as the baseline currents blocked by a saturating concentration of $GABA_A$ receptor

antagonists, was significantly greater than in GCs during adolescence, tonic $GABA_A$ current

amplitude was not different between GCs and SGCs during infancy or adulthood (Fig. 6). Both

GCs and SGCs showed a significant increase in tonic GABA currents from infancy to

adolescence which returned back to pre-adolescent levels in adults (Fig. 6 and Supplementary

Tables 5 & 6).

Enhanced inhibitory regulation of dentate activity *in vivo* during adolescence

Developmental shifts in GABAergic inhibition would be expected to alter dentate output in

response to input activation. Having assessed spontaneous and tonic steady state synaptic

inhibition in dentate projection neuronal classes during development we sought to assess whether

dentate inhibition at the level of the circuit tracked the developmental profiles of synaptic

inhibitory charge transfer which declined with development or extrasynaptic inhibition which

was selectively enhanced during adolescence. Paired stimuli to the angular bundle evoked with

an interval of 20 ms have been reliably used to evaluate hippocampal network inhibition by local

neurons (Jedlicka et al. 2010; Jedlicka et al. 2011). *In vivo* examination of GCL population

response evoked by a pair of stimuli to the angular bundle at 20 ms interval in infant, adolescent,

and adult rats identified a significant decrease in paired pulse ratios (PPR) in adolescent group

compared to infant and adult groups (Fig. 7; PPR: 0.28 ± 0.03 in n=7 adolescents vs. 0.79 ± 0.06 in

n=8 infants and 0.61 ± 0.09 in n=6 adults; $p<0.05$, Kruskal-Wallis one way ANOVA on ranks).

In line with previous developmental studies evaluating PPR in rats (DiScenna and Teyler 1994),

we did not see a significant difference in PPR between the infants and adult groups. Although the

magnitude of steady-state inhibition may not reflect recruitment of inhibition during afferent

activation, results from anesthetized rats in vivo demonstrating heightened feedback inhibitory

regulation of dentate activity during adolescence parallel the developmental peak in tonic

inhibition in GCs and SGCs in adolescent rats.

Discussion:

Contemporary literature on the dentate gyrus describes the projection neurons as a largely homogeneous population of GCs with limited diversity (Kesner 2018). Since GCs are a unique subset of neurons that undergo neurogenesis and maturation through adulthood, structural and functional diversity in GCs is largely attributed to the maturation state of neurons within the circuit (Toda and Gage 2018). At any given postnatal developmental stage of the animal, immature GCs tend to be located closer the hilar border of the cell layer, have less elaborate dendritic structures and are functionally more excitable than their mature counterparts (Kerloch et al. 2018; Overstreet-Wadiche and Westbrook 2006). However, emerging recognition of a structurally and functionally distinct subset of dentate projection neurons challenges the prevailing view that dentate projection neurons are a homogenous class of cells distinguished by developmental stages (Gupta et al. 2012; Save et al. 2018; Williams et al. 2007). SGCs, which have been characterized in the IML of the dentate gyrus, differ from GCs in their expansive dendritic arbors and show more sustained firing activity (Larimer and Strowbridge 2010; Williams et al. 2007). These and other features, including their distinct window of embryonic day 12-15 for SGC development (Save et al. 2018), enhanced excitatory drive (Larimer and Strowbridge 2010), and distinct synaptic and extrasynaptic inhibitory currents (Gupta et al. 2012) indicate that SGCs are distinct from GCs. Although SGCs have been proposed to sculpt feedback inhibition of GCs, gate dentate activity, and contribute to memory processing (Larimer and Strowbridge 2010; Walker et al. 2010), there is limited information on the structural and physiological differences between GCs and SGCs across postnatal development of the animal. This information is needed to determine which features consistently and reliably distinguish the cell types at all age groups. Our detailed and objective morphometric analysis conducted at three

distinct developmental stages, namely, infant, adolescent and adult age groups demonstrates that

clustering dentate projection neurons based on somato-dendritic morphology distinguishes them

into different "subtypes" corresponding to SGCs and GCs. Additionally, SGCs consistently

maintained higher frequency of inhibitory synaptic inputs than GCs at all ages. Our results

demonstrate that although both GCs and SGCs exhibit peaks in frequency of synaptic inhibitory

inputs and magnitude of tonic GABA currents during adolescence, these parameters were

significantly higher in SGCs than in GCs at this time point. However, the peak sIPSC amplitude

and cumulative charge transfer were highest in infants, decreased with development and were not

different between cell types. These findings demonstrate that SGCs are a structurally and

functionally distinct subtype of dentate projection neurons which is in keeping with the emerging

recognition of subpopulations among hippocampal and entorhinal projection neurons (Pilli et al.

2012; Soltesz and Losonczy 2018). Moreover, the significantly heightened tonic inhibition in

SGCs during adolescence suggests that SGC activity could be more strongly modulated than

GCs by a variety of neuroactive compounds including alcohol and neurosteroids, which

selectively augment extrasynaptic GABA currents (Maguire and Mody 2009; Mody et al. 2007).

Interestingly, *in vivo* analysis of paired pulse inhibition, demonstrated that feedback inhibition at

the level of the network was maximal during adolescence. While the parallel peak in feedback

inhibition *in vivo* and tonic GABA currents during adolescence are intriguing, it is paradoxical

that dentate feedback inhibition is enhanced when SGCs, which have been proposed to drive

feedback inhibition, are under relatively higher tonic inhibitory regulation. These findings raise

the possibility that higher extrasynaptic inhibitory tone in SGCs may be a compensatory

mechanism to reduce feedback inhibition during adolescence. Future studies examining this

possibility coupled with developmental changes in excitatory drive and active properties of GCs

and SGCs can provide a more comprehensive understanding of the network role of SGCs.

In addition to using unbiased approaches to classify SGCs as a distinct neuronal class, our data

identify key age-invariant features to distinguish the SGCs from GCs. We find that the number

of primary dendrites, dendritic angle, and soma ratio are significantly higher in SGCs and can be

used to categorize SGCs and GCs. In particular, the multiple primary dendrites observed in

SGCs stands in stark contrast to the typical one to two apical dendrites observed in granule cell

reconstructions (Thind et al. 2008). Indeed, the striking >85% correspondence between the

unsupervised clustering and investigator assigned clustering likely stems from the investigator's

use of dendritic angle, soma width, which are significant contributors to the first PC, in addition

to soma location to classify cell types. Dendritic length, on the other hand, increased during

development but was not different between cell types indicating that processes reflecting

developmental maturation are common to the cell types. Consistent with the presence of larger

dendritic angle, the convex hull 3D volume and 3D surface area of SGCs were greater than that

of GCs. However, there was also a developmental increase in these parameters from infancy to

adolescence and a further stabilization into adulthood in both cell types, likely reflecting the

developmental increase in hippocampal volume. Interestingly, although the number of terminal

nodes was not different, the dendritic complexity was significantly lower in SGCs than in GCs

demonstrating differences in branching patterns. Difference in branching patterns including

dendritic complexity can impact neuronal firing and intracellular signaling, as has been

demonstrated in modeling studies (Li et al. 2015; van der Velden et al. 2012; van Elburg and van

Ooyen 2010). Whether dendritic morphology can account for differences in intrinsic physiology

between the dentate projection neuron types (Gupta et al. 2012; Save et al. 2018; Williams et al.

2007) remains to be examined. The other aspect where distinguishing cell type based on

morphology becomes critical is in disease. Dentate granule cells are known to undergo changes

in dendritic structure including alterations in complexity under physiological conditions and in

trauma and neurodegenerative diseases (Llorens-Martin et al. 2015; Redila and Christie 2006;

Villasana et al. 2015). The ability to distinguish GCs and SGCs across multiple developmental

stages would be crucial to quantifying and interpreting changes in morphology in trauma,

epilepsy, and neurodegenerative disease.

While there are clear differences in the structure, SGCs and GCs share several characteristics.

Both SGCs and GCs are projection neurons, with dendrites in the dentate molecular layer and

axons projecting to hippocampal CA3 (Gupta et al. 2012; Save et al. 2018; Williams et al. 2007).

Several dendritic parameters including total numbers of dendritic segments, nodes, terminals,

and dendritic tortuosity (Supplementary Table 3) showed neither cell type nor age-related

differences. Similarly, SGCs, like GCs, have dendritic spines which can aid in distinguishing

them from inhibitory neurons. Additionally, hilar axon collaterals of SGCs have "mossy fiber

boutons" typically attributed to GCs (Supplementary Fig. 1 and Save et al. 2018). Notably, in

pilot clustering analysis which included a few molecular layer inhibitory neurons, the

interneurons clustered on a different branch of the dendrogram than SGCs and GCs (data not

shown). The similarities between GCs and SGCs are not surprising, since we previously

identified that SGCs express the Prospero homeobox protein 1 (Prox1) present in GCs indicating

a shared lineage (Gupta et al. 2012). This shared lineage was confirmed by a recent study which

identified a narrow developmental window of embryonic day 12-15 during which SGCs are

produced from the same precursor niche as GCs (Save et al. 2018). Yet, the morphology of

embryonically labeled SGCs were distinct from that of GCs labeled on the same day (Save et al.

2018), demonstrating that they are a distinct population of cells rather than a cohort of GCs with

a different maturation state. In this context, studies examining dendritic properties of GCs with

somata located in the outer third of the molecular layer have consistently reported wider

dendritic fields and distinct dendritic arbors consistent with the possibility that the outer third of

the granule cell layer may consist of a mixed population of GCs and SGCs (Green and Juraska

1985; Kerloch et al. 2018; Sun et al. 2013).   Whether specific genetic, molecular or

developmental cues guide the development of SGCs during neurogenesis or whether local

molecular and spatial factors in the dentate GCL-molecular layer border contribute to elaboration

of distinct dendritic arbors remains to be determined (Hatami et al. 2018; Lefebvre et al. 2015).

Although the mechanisms underlying specification of SGCs as a distinct population needs

further investigation, our objective unsupervised morphometric analysis identifies the key

somato-dendritic structural features that distinguish SGCs from GCs through postnatal

development.

A defining feature of the dentate gyrus is the presence of heavy inhibitory regulation (Coulter

and Carlson 2007). Dentate GCs receive synaptic inhibition from a diverse population of neurons

and are also under steady-state, tonic extrasynaptic inhibition (Coulter and Carlson 2007; Ewell

and Jones 2010; Harney and Jones 2002; Stell et al. 2003). In an earlier study, we identified that

SGCs are under stronger inhibitory regulation than GCs with higher frequency of inhibitory

synaptic currents and greater amplitude of tonic GABA currents (Gupta et al. 2012). Here, we

find that SGCs continue to receive greater spontaneous synaptic inhibitory events than GCs

through postnatal development. This contrasts with the lack of difference in inhibitory current

between embryonically-born and adult-born mature granule cells (Laplagne et al. 2007).

Interestingly, while the frequency of sIPSCs peaked during adolescence in both cell types, the

amplitude decreases progressively with development. Additionally, while synaptic membrane

kinetics decreased with development in GCs, they increased with development of SGCs. The

combined effect of cell specific and developmental changes resulted in an overall decrease in

cumulative synaptic inhibitory charge transfer from infancy to adulthood while maintaining

similar charge transfer between age-matched GCs and SGCs. In addition to developmental

changes in $\tau_{membrane}$ which could contribute to cell-specific regulation of synaptic kinetics, the

roles of changes in GABA receptor subunits, dendritic pruning and synaptic distribution need to

be considered in future works. Although lower R_{in} in SGCs has been considered a defining

distinction from GCs (Erwin et al. 2020; Gupta et al. 2012; Williams et al. 2007), our

demonstration that SGCs in adult rats have higher R_{in} than in GCs suggests that R_{in} may not be

an age-invariant feature for cell classification.

Can dendritic structural features explain the cell-type specific and developmental differences in

sIPSC frequency in dentate projection neurons? The consistently higher sIPSC frequency in

SGCs is surprising as the dendritic lengths and location are not different between GCs and SGCs

(Fig. 3). The wider dendritic distribution of SGCs, also reflected in the greater convex hull 3D

volume, could allow for inputs from a larger group of inhibitory neurons to impinge on SGC

dendrites, while GCs with their compact dendritic distribution may receive fewer inputs. It is

also possible that the relatively early embryonic development of SGCs increases the chance of

SGCs to receive more synaptic inputs compared to GCs which develop later into adulthood.

However, since both dendritic length and convex hull 3D volume increase with postnatal

development of both cells, changes in dendritic morphology and embryonic development are

unlikely to account for the peak in sIPSC frequency in adolescence followed by decline in adults.

The possibility of developmental increase in synapses from infancy to adolescence followed by

pruning or synapse elimination into adulthood (Riccomagno and Kolodkin 2015; Tran et al.

2009) may be considered in future works. The progressive decrease in sIPSC amplitude with

development in both SGCs and GCs could reflect the increase in dendritic length with

development. Consistent with this proposal, the proportion of large amplitude events decrease

progressively with development in both cell types (Supplementary Figure 5).

In parallel, the amplitude of extrasynaptic GABA currents in SGCs peaked during adolescence.

Similarly, tonic GABA currents in GCs showed a peak during adolescence confirming the

developmental increase reported in previous studies (Holter et al. 2010). Increase in expression

of the extrasynaptic $GABA_AR$ δ subunit in both GCs and SGCs (Gupta et al. 2012; Maguire and

Mody 2009) is likely to mediate the increase in tonic GABA currents from infancy to

adolescence. Additionally, changes in synaptically released GABA accompanying changes in

sIPSC frequency, reported here, could contribute to the peak in extrasynaptic GABA exhibited in

adolescence. An interesting feature of tonic GABA currents mediated by $GABA_AR$ δ subunit is

their robust enhancement by neurosteroids, raising the possibility that increases in ambient

neurosteroids during adolescence (Harden and MacLusky 2004; Maguire and Mody 2009) could

augment tonic GABA currents. Since tonic GABA currents mediated by $GABA_AR$ δ subunits are

exquisitely sensitive to ethanol (Mody et al. 2007), the greater magnitude of tonic GABA

currents in SGCs during adolescence is likely to render SGCs vulnerable to ethanol modulation

and impact their role in dentate processing. Indeed, behavioral deficits following alcohol

administration have been shown to be particularly accentuated during adolescence (Lacaille et al.

2015; Spanos et al. 2012). It would be important for future studies to ascertain how the

differences in basal synaptic inhibition impact the recruitment and circuit function of SGCs and

GCs during network activity and with development. Since basal inhibition in the dentate gyrus

regulates synaptic plasticity and pattern separation at the level of the circuit (Dengler and Coulter

2016; Madar et al. 2019), and SGCs have been proposed to contribute to dentate pattern

separation (Larimer and Strowbridge 2010), developmental changes in GC and SGC inhibition

are likely to influence memory and cognitive performance.

Together, the structural and functional data identify SGCs as a cell type which differs from GCs

in somato-dendritic structure and developmental plasticity of inhibitory inputs, extrasynaptic

inhibition and membrane kinetics. Our data demonstrate that SGCs have heightened

extrasynaptic inhibition during adolescence which would make them susceptible to endogenous

and exogenous modulation of activity levels during adolescence. Notably, our results delineate

salient structural features that enable anatomical identification of this subpopulation of dentate

projection neurons. The novel data defining the structural features of SGCs will allow for future

targeted analysis of their molecular profile and microcircuit connectivity to better understand

their role in circuit function and behaviors. In conclusion, the fundamental characterization of

SGCs presented here will support incorporation of SGCs into current models of the dentate gyrus

and consideration of their role in dentate microcircuit processing in health and disease.

Reference List

Coulter DA, Carlson GC (2007) Functional regulation of the dentate gyrus by GABA-mediated

inhibition ProgBrain Res 163:235-243

Dengler CG, Coulter DA (2016) Normal and epilepsy-associated pathologic function of the

dentate gyrus Prog Brain Res 226:155-178 doi:10.1016/bs.pbr.2016.04.005

Dieni CV, Chancey JH, Overstreet-Wadiche LS (2012) Dynamic functions of GABA signaling

during granule cell maturation Front Neural Circuits 6:113 doi:10.3389/fncir.2012.00113

DiScenna PG, Teyler TJ (1994) Development of inhibitory and excitatory synaptic transmission

in the rat dentate gyrus Hippocampus 4:569-576 doi:10.1002/hipo.450040506

Duffy CJ, Rakic P (1983) Differentiation of granule cell dendrites in the dentate gyrus of the

rhesus monkey: a quantitative Golgi study J Comp Neurol 214:224-237

doi:10.1002/cne.902140210

Engel D, Jonas P (2005) Presynaptic action potential amplification by voltage-gated Na+

channels in hippocampal mossy fiber boutons Neuron 45:405-417

doi:10.1016/j.neuron.2004.12.048

Erwin SR, Sun W, Copeland M, Lindo S, Spruston N, Cembrowski MS (2020) A Sparse,

Spatially Biased Subtype of Mature Granule Cell Dominates Recruitment in Hippocampal-

Associated Behaviors Cell Rep 31:107551 doi:10.1016/j.celrep.2020.107551

Ewell LA, Jones MV (2010) Frequency-tuned distribution of inhibition in the dentate gyrus The

Journal of neuroscience : the official journal of the Society for Neuroscience 30:12597-12607

doi:10.1523/JNEUROSCI.1854-10.2010

Freiman TM, Eismann-Schweimler J, Frotscher M (2011) Granule cell dispersion in temporal

lobe epilepsy is associated with changes in dendritic orientation and spine distribution Exp

Neurol 229:332-338 doi:10.1016/j.expneurol.2011.02.017

Green EJ, Juraska JM (1985) The dendritic morphology of hippocampal dentate granule cells

varies with their position in the granule cell layer: a quantitative Golgi study Exp Brain Res

59:582-586

Gupta A, Elgammal FS, Proddutur A, Shah S, Santhakumar V (2012) Decrease in tonic

inhibition contributes to increase in dentate semilunar granule cell excitability after brain injury J

Neurosci 32:2523-2537 doi:10.1523/JNEUROSCI.4141-11.2012

Harden C, MacLusky NJ (2004) Aromatase inhibition, testosterone, and seizures Epilepsy Behav

5:260-263 doi:10.1016/j.yebeh.2003.12.001

Harney SC, Jones MV (2002) Pre- and postsynaptic properties of somatic and dendritic

inhibition in dentate gyrus Neuropharmacology 43:584-594

Hatami M, Conrad S, Naghsh P, Alvarez-Bolado G, Skutella T (2018) Cell-Biological

Requirements for the Generation of Dentate Gyrus Granule Neurons Front Cell Neurosci 12:402

doi:10.3389/fncel.2018.00402

Hollrigel GS, Ross ST, Soltesz I (1998) Temporal patterns and depolarizing actions of

spontaneous GABAA receptor activation in granule cells of the early postnatal dentate gyrus J

Neurophysiol 80:2340-2351 doi:10.1152/jn.1998.80.5.2340

Holter NI, Zylla MM, Zuber N, Bruehl C, Draguhn A (2010) Tonic GABAergic control of

mouse dentate granule cells during postnatal development Eur J Neurosci 32:1300-1309

doi:10.1111/j.1460-9568.2010.07331.x

Husson F, Josse J, Pages J (2010) Principal component methods - hierarchical clustering -

partitional clustering: why would we need to choose for visualizing data?

Jedlicka P, Deller T, Schwarzacher SW (2010) Computational modeling of GABAA receptor-

mediated paired-pulse inhibition in the dentate gyrus J Comput Neurosci 29:509-519

doi:10.1007/s10827-010-0214-y

Jedlicka P et al. (2011) Increased dentate gyrus excitability in neuroligin-2-deficient mice in vivo

Cereb Cortex 21:357-367 doi:10.1093/cercor/bhq100

Kerloch T, Clavreul S, Goron A, Abrous DN, Pacary E (2018) Dentate Granule Neurons

Generated During Perinatal Life Display Distinct Morphological Features Compared With Later-

Born Neurons in the Mouse Hippocampus Cereb Cortex doi:10.1093/cercor/bhy224

Kesner RP (2018) An analysis of dentate gyrus function (an update) Behav Brain Res 354:84-91

doi:10.1016/j.bbr.2017.07.033

Lacaille H, Duterte-Boucher D, Liot D, Vaudry H, Naassila M, Vaudry D (2015) Comparison of

the deleterious effects of binge drinking-like alcohol exposure in adolescent and adult mice J

Neurochem 132:629-641 doi:10.1111/jnc.13020

Laplagne DA, Kamienkowski JE, Esposito MS, Piatti VC, Zhao C, Gage FH, Schinder AF

(2007) Similar GABAergic inputs in dentate granule cells born during embryonic and adult

neurogenesis Eur J Neurosci 25:2973-2981 doi:10.1111/j.1460-9568.2007.05549.x

Larimer P, Strowbridge BW (2010) Representing information in cell assemblies: persistent

activity mediated by semilunar granule cells Nat Neurosci 13:213-222 doi:10.1038/nn.2458

Lee CY, Liou HH (2013) GABAergic tonic inhibition is regulated by developmental age and

epilepsy in the dentate gyrus Neuroreport 24:515-519 doi:10.1097/WNR.0b013e32836205bc

Lefebvre JL, Sanes JR, Kay JN (2015) Development of dendritic form and function Annu Rev

Cell Dev Biol 31:741-777 doi:10.1146/annurev-cellbio-100913-013020

Li L, Gervasi N, Girault JA (2015) Dendritic geometry shapes neuronal cAMP signalling to the

nucleus Nat Commun 6:6319 doi:10.1038/ncomms7319

Li Y, Aimone JB, Xu X, Callaway EM, Gage FH (2012) Development of GABAergic inputs

controls the contribution of maturing neurons to the adult hippocampal network Proc Natl Acad

Sci U S A 109:4290-4295 doi:10.1073/pnas.1120754109

Llorens-Martin M, Rabano A, Avila J (2015) The Ever-Changing Morphology of Hippocampal

Granule Neurons in Physiology and Pathology Front Neurosci 9:526

doi:10.3389/fnins.2015.00526

Madar AD, Ewell LA, Jones MV (2019) Temporal pattern separation in hippocampal neurons

through multiplexed neural codes PLoS Comput Biol 15:e1006932

doi:10.1371/journal.pcbi.1006932

Maguire J, Mody I (2009) Steroid hormone fluctuations and GABA(A)R plasticity

Psychoneuroendocrinology 34 Suppl 1:S84-90 doi:10.1016/j.psyneuen.2009.06.019

Mody I, Glykys J, Wei W (2007) A new meaning for "Gin & Tonic": tonic inhibition as the

target for ethanol action in the brain Alcohol 41:145-153 doi:10.1016/j.alcohol.2007.03.009

Neuberger EJ, Swietek B, Corrubia L, Prasanna A, Santhakumar V (2017) Enhanced Dentate

Neurogenesis after Brain Injury Undermines Long-Term Neurogenic Potential and Promotes

Seizure Susceptibility Stem Cell Reports 9:972-984 doi:10.1016/j.stemcr.2017.07.015

Overstreet-Wadiche LS, Westbrook GL (2006) Functional maturation of adult-generated granule

cells Hippocampus 16:208-215 doi:10.1002/hipo.20152

Pilli J, Abbasi S, Richardson M, Kumar SS (2012) Diversity and excitability of deep-layer

entorhinal cortical neurons in a model of temporal lobe epilepsy J Neurophysiol 108:1724-1738

doi:10.1152/jn.00364.2012

Ramón y Cajal S (1953) Histologie du système nerveux de l'homme & des vertébrés. Éd.

franc☐aise rev. & mise à jour edn. Consejo Superior de Investigaciones Científicas, Instituto

Ramón y Cajal, Madrid,

Redila VA, Christie BR (2006) Exercise-induced changes in dendritic structure and complexity

in the adult hippocampal dentate gyrus Neuroscience 137:1299-1307

doi:10.1016/j.neuroscience.2005.10.050

Riccomagno MM, Kolodkin AL (2015) Sculpting neural circuits by axon and dendrite pruning

Annu Rev Cell Dev Biol 31:779-805 doi:10.1146/annurev-cellbio-100913-013038

Sancho-Bielsa FJ, Navarro-Lopez JD, Alonso-Llosa G, Molowny A, Ponsoda X, Yajeya J,

Lopez-Garcia C (2012) Neurons of the dentate molecular layer in the rabbit hippocampus PLoS

One 7:e48470 doi:10.1371/journal.pone.0048470

Save L, Baude A, Cossart R (2018) Temporal Embryonic Origin Critically Determines Cellular

Physiology in the Dentate Gyrus Cereb Cortex doi:10.1093/cercor/bhy132

Semple BD, Blomgren K, Gimlin K, Ferriero DM, Noble-Haeusslein LJ (2013) Brain

development in rodents and humans: Identifying benchmarks of maturation and vulnerability to

injury across species Prog Neurobiol 106-107:1-16 doi:10.1016/j.pneurobio.2013.04.001

Sengupta P (2013) The Laboratory Rat: Relating Its Age With Human's Int J Prev Med 4:624-

630

Seress L, Frotscher M (1990) Morphological variability is a characteristic feature of granule cells

in the primate fascia dentata: a combined Golgi/electron microscope study J Comp Neurol

293:253-267

Soltesz I, Losonczy A (2018) CA1 pyramidal cell diversity enabling parallel information

processing in the hippocampus Nat Neurosci 21:484-493 doi:10.1038/s41593-018-0118-0

Soltesz I, Smetters DK, Mody I (1995) Tonic inhibition originates from synapses close to the

soma Neuron 14:1273-1283

Spanos M, Besheer J, Hodge CW (2012) Increased sensitivity to alcohol induced changes in

ERK Map kinase phosphorylation and memory disruption in adolescent as compared to adult

C57BL/6J mice Behav Brain Res 230:158-166 doi:10.1016/j.bbr.2012.02.010

Stell BM, Brickley SG, Tang CY, Farrant M, Mody I (2003) Neuroactive steroids reduce

neuronal excitability by selectively enhancing tonic inhibition mediated by delta subunit-

containing GABAA receptors Proc Natl Acad Sci U S A 100:14439-14444

doi:10.1073/pnas.2435457100

Sun GJ, Sailor KA, Mahmood QA, Chavali N, Christian KM, Song H, Ming GL (2013)

Seamless reconstruction of intact adult-born neurons by serial end-block imaging reveals

complex axonal guidance and development in the adult hippocampus J Neurosci 33:11400-

11411 doi:10.1523/JNEUROSCI.1374-13.2013

Swietek B, Gupta A, Proddutur A, Santhakumar V (2016) Immunostaining of Biocytin-filled and

Processed Sections for Neurochemical Markers J Vis Exp doi:10.3791/54880

Thind KK, Ribak CE, Buckmaster PS (2008) Synaptic input to dentate granule cell basal

dendrites in a rat model of temporal lobe epilepsy J Comp Neurol 509:190-202

doi:10.1002/cne.21745

Toda T, Gage FH (2018) Review: adult neurogenesis contributes to hippocampal plasticity Cell

Tissue Res 373:693-709 doi:10.1007/s00441-017-2735-4

Tran TS et al. (2009) Secreted semaphorins control spine distribution and morphogenesis in the

postnatal CNS Nature 462:1065-1069 doi:10.1038/nature08628

van der Velden L, van Hooft JA, Chameau P (2012) Altered dendritic complexity affects firing

properties of cortical layer 2/3 pyramidal neurons in mice lacking the 5-HT3A receptor J

Neurophysiol 108:1521-1528 doi:10.1152/jn.00829.2011

van Elburg RA, van Ooyen A (2010) Impact of dendritic size and dendritic topology on burst

firing in pyramidal cells PLoS Comput Biol 6:e1000781 doi:10.1371/journal.pcbi.1000781

Villasana LE, Kim KN, Westbrook GL, Schnell E (2015) Functional Integration of Adult-Born

Hippocampal Neurons after Traumatic Brain Injury(1,2,3) eNeuro 2

doi:10.1523/ENEURO.0056-15.2015

von Campe G, Spencer DD, de Lanerolle NC (1997) Morphology of dentate granule cells in the

human epileptogenic hippocampus Hippocampus 7:472-488 doi:10.1002/(SICI)1098-

1063(1997)7:5<472::AID-HIPO4>3.0.CO;2-J

Walker MC, Pavlov I, Kullmann DM (2010) A 'sustain pedal' in the hippocampus? Nature

neuroscience 13:146-148 doi:10.1038/nn0210-146

Williams PA, Larimer P, Gao Y, Strowbridge BW (2007) Semilunar granule cells: glutamatergic

neurons in the rat dentate gyrus with axon collaterals in the inner molecular layer J Neurosci

27:13756-13761 doi:10.1523/JNEUROSCI.4053-07.2007

Yu EP, Dengler CG, Frausto SF, Putt ME, Yue C, Takano H, Coulter DA (2013a) Protracted

postnatal development of sparse, specific dentate granule cell activation in the mouse

hippocampus J Neurosci 33:2947-2960 doi:10.1523/JNEUROSCI.1868-12.2013

Yu J, Proddutur A, Elgammal FS, Ito T, Santhakumar V (2013b) Status epilepticus enhances

tonic GABA currents and depolarizes GABA reversal potential in dentate fast-spiking basket

cells J Neurophysiol 109:1746-1763 doi:10.1152/jn.00891.2012

Yu J, Proddutur A, Swietek B, Elgammal FS, Santhakumar V (2016) Functional Reduction in

Cannabinoid-Sensitive Heterotypic Inhibition of Dentate Basket Cells in Epilepsy: Impact on

Network Rhythms Cereb Cortex 26:4229-4314 doi:10.1093/cercor/bhv199

 Figure Legends

Figure 1. Somato-dendritic differences in dentate projection neurons - GCs and SGCs.

Representative neuronal reconstructions of GC (1A-F) and SGC (1G-L) showing distinct

dendritic arbors from infant (top), adolescent (middle) and adult (bottom) age groups. Note that

images are shown in different planes for each age group. The complexity of dendrites is shown

in XY planes for GCs and SGCs 'A, B, C' and 'G, H, I' respectively and perpendicular view is

shown in 'D, E, F' for GCs and 'J, K, L' for SGCs. Inset images in the center represent the color

coding for dendritic segments from proximal to distal axes. Scale bar: 100µm.

Figure 2. Hierarchical clustering on principal component reveals two major clusters of

dentate projection neurons. (A-B) 3D representation of the principal component analysis of

individuals resolved by first three principal components, with confidence interval (CI) ellipsoid

set to 0.95. The plot suggests grouping by cell type (A) but not by age (B). (C) Hierarchical

clustering on Principal Components based on 42 morphometric parameters (Supplementary

Tables 1 & 2) was performed using Ward's method with Euclidean distance to generate the

dendrogram. Dendrogram classifies neurons into two putative clusters. Cells in which the

classifier and experimenter failed to concur on classification are represented by green

arrowheads based on investigator assigned classification.

Figure 3: Comparison of morphometric parameters between GCs and SGCs at distinct

developmental stages. Summary plots of averages no. of primary dendrites (A), soma width (B),

dendritic angle (C), total dendritic length (D), dendritic complexity (E), and convex hull surface

area (F) of GCs and SGCs at three developmental time points. *, #, and $ denote $p<0.05$ for

differences between cell types, in GC across age groups and SGC across age groups, respectively

by TW-ANOVA followed by post-hoc pairwise comparison (Supplementary Tables 3 & 4).

N= GCs, 6 infant, 6 adolescent and 4 adult and SGCs, 9 infant, 5 adolescent and 6 adult.

Figure 4: Developmental differences in sIPSC frequencies of GCs and SGCs. Representative

sIPSC traces in GCs (left, A1, B1, C1) and SGCs (right, A2, B2, C2) in infant, adolescent, and

adult age groups. Summary plot of developmental differences in sIPSC frequency between and

within GCs and SGCs (D). *, #, and $ denote $p<0.05$ for differences between cell types, in GC

across age groups and SGC across age groups, respectively by TW-ANOVA followed by post-

hoc pairwise comparison (Supplementary Tables 5 & 6). Cumulative probability plots of sIPSCs

frequency show differences between cell types at infant (E1), adolescent (E2) and adult (E3) age

groups * denotes $p<0.05$ for differences between cell types by Kruskal-Wallis Test

(Supplementary Tables 5 & 6).

Figure 5: Developmental changes in average sIPSC parameters in GCs and SGCs. Overlay

of representative average sIPSC waveforms from GCs (A) and SGCs (B) in different age

groups. Summary plots of sIPSC amplitude (C), weighted τ_{decay} (D), 20-80 rise time (E), and

charge transfer (F) at three developmental stages in both cell types. *, #, and $ denote $p<0.05$ for

differences between cell types, in GC across age groups and SGC across age groups, respectively

by TW-ANOVA followed by post-hoc pairwise comparison (Supplementary Tables 5 - 8).

Figure 6: Extrasynaptic tonic GABAergic currents in SGCs peak during adolescence.

Representative baseline current recordings in GCs (left, A1-3) and SGCs (right, B1-3) in infant,

adolescent and adult age groups. Tonic GABA current is measured as the difference in baseline

current upon perfusion of the GABA receptor antagonist BMI (100µM). Gaussian fit to the

positive half of the baseline current, under basal conditions and in BMI, is illustrated on the right

of each trace and was used to quantify tonic GABA. (C) Summary plot showing average tonic

GABAergic currents at three distinct time points. *, #, and $ denote $p < 0.05$ for differences

between cell types, in GCs across age groups and in SGCs across age groups, respectively by

TW-ANOVA followed by post-hoc pairwise comparison (Supplementary Tables 5 & 6).

Figure 7: Paired pulse inhibition as a measure of local feedback inhibition. (A) Shows

representative traces showing paired pulse inhibition among infant, adolescent, and adult groups

in response to activation of medial Perforant path (mPP) in anesthetized rats. Note the difference

in y scale bars. (B) Box plot showing the distribution of paired pulse ratio's between the groups

(infant vs. adolescent p=<0.001; adolescent vs. adults p=0.007; infants vs. adults p=0.068 : *

indicates $p < 0.05$ by One-way ANOVA followed by pairwise comparison by Post-hoc Holm-

Sidak method.

Supplementary Figure legends

Supplementary Figure 1: Representative images of a GC and SGC. Images of a typical GC (A) and SGC (B) illustrate the somatic location, dendritic arbor, with high density of spines (insets, white arrows) and axons with boutons (white arrow heads) targeting CA3 used by experimenter to classify SGCs and GCs. Scale bar: 100µm; Inset scale bar: 20 µm.

Supplementary Figure 2: Analysis of Morphological Variables underlying Principal Components. A. Histogram illustrates percentage of information retained by each dimension (principal component, PC). B. Factor maps illustrate the quality of representation of the morphometric variables measured by cos2 (square cosine, squared coordinates). The darker color indicates stronger contribution to variability to PC. C. Representation of the top five variables in the first two dimensions. D. Neuronal reconstructions of SGC in first cluster (notice higher complexity than other SGCs in figure 1). Scale bar: 100µm.

Supplementary Figure 3: Comparison of morphometric parameters between GCs and SGCs at distinct developmental stages. Summary plot of number of second (A) and third (B) order dendritic segments, second order nodes (C), total number of dendritic ends (D), total dendritic nodes (E) and total dendritic segments (F) in GCs and SGCs at the three age groups examined. * denotes $p<0.05$ for differences between cell types by TW-ANOVA followed by post-hoc pairwise comparison (Supplementary Tables 3 & 4). N= GCs, 6 infant, 6 adolescent and 4 adult and SGCs, 9 infant, 5 adolescent and 6 adult.

Supplementary Figure 4: Cumulative plots of sIPSC frequency in GCs and SGCs through

development. Cumulative plots of sIPSC frequency comparing three age groups * denotes

p<0.05 for differences between cell types by Kruskal-Wallis Test (Supplementary Table 6) in

GCs (A) and SGCs (B). Summary plot of cumulative charge transfer over one second (C) at

three developmental stages in both cell types. #, and $ denote p<0.05 for differences in GC

across age groups and SGC across age groups, respectively by TW-ANOVA followed by post-

hoc pairwise comparison.

Supplementary Figure 5: Developmental differences in distribution of sIPSC amplitudes in

GCs and SGCs. Pie chart distributions illustrating subjective percentage distribution of high

peak amplitude (>50pA) and smaller peak amplitude (<50 pA) amplitude sIPSCs in GCs (top)

and SGCs (bottom) across age groups.

Supplementary Figure 6: Analysis of developmental changes in passive membrane

properties of GCs and SGCs. Representative voltage traces in response to a -200pA current

injection for one second in GCs and SGCs at the developmental stages under investigation

reveal differences in R_{in} (A). Summary of R_{in} in the cell types (B). Summary plot of membrane

time constant ($\tau_{membrane}$) obtained from single exponential fits to the voltage response to -200pA

current injection (C). *, #, and $ denote p<0.05 for differences between cell types, in GC across

age groups and SGC across age groups, respectively by TW-ANOVA followed by post-hoc

pairwise comparison (Supplementary Table 8).

Figure 3

Figure 4