1   Complete genome sequence and comparative genomics of the fish

2   pathogen *Streptococcus iniae* QMA0248

Areej S. Alsheikh-Hussain[1,2,3], Nouri L. Ben Zakour[1,2,3,5], Brian M. Forde[1,2,3],

Oleksandra Silayeva[4], Andrew C. Barnes[4], and Scott A. Beatson*[1,2,3].

[1]School of Chemistry & Molecular Biosciences, [2]Australian Infectious Diseases

Research Centre, [3]Australian Centre for Ecogenomics, and [4]School of Biological

Science, The University of Queensland, Brisbane, Queensland, Australia;

Present address: [5]The Westmead Institute for Medical Research and the University

of Sydney, Sydney, New South Wales, Australia

Key words: SMRT sequencing, Insertion Sequence, reference guided assembly,

misassembly, mobile genetic elements

* Corresponding author.

Mailing address: School of Chemistry and Molecular Biosciences, Building 76, The

University of Queensland, Brisbane QLD 4072, Australia.

Phone: +617 33654863; E-mail: s.beatson@uq.edu.au

Abstract

Fish mortality caused by *Streptococcus iniae* is a major economic problem in fish

aquaculture in warm and temperate regions globally. There is also risk of zoonotic

infection by *S. iniae* through handling of contaminated fish. In this study, we

present the complete genome sequence of *S. iniae* strain QMA0248, isolated from

farmed barramundi in South Australia. The 2.12 Mb genome of *S. iniae* QMA0248

carries a 32 Kb prophage, a 12 Kb genomic island, and 92 discrete insertion

sequence (IS) elements. These include 9 novel IS types that belong mostly to the

IS*3* family. Comparative and phylogenetic analysis between *S. iniae* QMA0248 and

publicly available complete *S. iniae* genomes revealed discrepancies that are likely

due to misassembly in the genomes of isolates ISET0901 and ISNO. We also

determined by long-range PCR that a tandem duplication of an rRNA region in the

PacBio assembly of QMA0248 was an assembly error. We suggest that a similar

rRNA duplication in the PacBio genome of *S. iniae* 89353 may also be a

misassembly. Our study not only highlights assembly problems in existing

genomes, but provides a high quality reference genome for *S. iniae* QMA0248,

including manually curated mobile genetic elements, that will assist future *S. iniae*

comparative genomic and evolutionary studies.

Introduction

Streptococcus iniae is a fish pathogen that causes mortality in a wide range of fish

species in wild and farmed, marine and freshwater environments, resulting in

large economic losses to aquaculture (1, 2). *S. iniae* is also considered an

opportunistic human pathogen, causing sporadic infections mostly in the elderly

who have more than one underlying health condition such as diabetes mellitus, or

chronic rheumatic heart disease (3, 4). *S. iniae* pathogenesis is imparted through

a repertoire of virulence factors (VF) including surface proteins, secreted toxins,

and capsular polysaccharide (CPS) (4). VFs can be acquired through lateral gene

transfer (LGT) of mobile genetic elements (MGE) such as composite transposons,

genomic islands (GI) or prophages.

MGEs are a means by which bacterial pathogens acquire traits that help adapt to

changing conditions including vaccination, antibiotics, a new host or environment

(5, 6). Indeed, they are considered the main drivers of gene flux in bacteria,

contributing to diversity within species (7). Insertion sequence (IS) elements, for

instance, are small MGEs (0.7-3.5 Kb) that have an important role in evolution and

genome plasticity. IS insertion within bacterial chromosomes or plasmids can

result in genetic modifications through insertional inactivation of genes or up-

regulation of adjacent intact genes through outward-facing promoter sequences

carried by some IS (8, 9). In some cases pairs of IS can mobilize intervening

sequence as a composite transposon (8). The mobility of IS elements leads to their

expansion or loss within bacterial lineages. Expansion is associated with

accumulation of pseudogenes, which is considered an early stage in genome

reduction as a mechanism for adaptation (9). Accordingly, to obtain the complete

evolutionary picture within bacterial species it is important to study the

distribution and abundance of IS elements.

As yet there is no study that focuses on the diversity and distribution of MGEs in

S. iniae genomes. In fact, only four complete *S. iniae* genomes were deposited in

Genbank at the commencement of the present study, none of which had

comprehensive annotations for MGEs: SF1, YSFST01-82, ISET0901, and ISNO. *S.*

iniae SF1 (accession: CP005941) was cultured from moribund flounder from a fish

farm that experienced an epidemic in North China (10). YSFST01-82 (accession:

CP010783) is an isolate from a diseased Olive flounder from South Korea (2012)

(11). ISET0901 (accession: CP007586) is reported to be a highly virulent strain,

which was isolated from diseased Nile tilapia during an outbreak in Israel (2005)

(12). The strain ISNO (accession: CP007587) was obtained through selection for

resistance of *S. iniae* ISET0901 to novobiocin (13).

There are currently 25 different IS families in the ISFinder database, but only one

IS described for *S. iniae* (IS*Stin1* of the family IS*256*). As is typical for complete

bacterial genomes, the annotation of the four complete *S. iniae* genomes available

at Genbank are limited to some of the transposase genes associated with IS

elements without definition of the IS boundaries. Indeed, the difficulty in

annotating IS elements and the lack of reliable automated annotation methods

means that only a small subset of complete genomes have accurate IS annotations.

Small partial IS elements are often disregarded although they reflect valuable

insights on ancestral recombination events. With the increasing availability of

long-read sequencing there is a need for a high quality, well-characterized *S. iniae*

reference genome that better enables the impact of IS elements on evolution and

diversity to be determined.

In this study, we have completely characterized the genome of *S. iniae* QMA0248.

Manual curation of annotations for IS, genomic islands, prophages, and CRISPR

was carried out along with a comparison with the four publicly available complete

genomes from NCBI (SF1, YSFST01-82, ISET0901, and ISNO). Comparative and

phylogenetic analyses revealed discrepancies between the MGE content

indicating likely misassembly in the genomes ISNO, and ISET0901. The complete

genome of QMA0248 will provide an important scaffold for future phylogenomic

studies of the *S. iniae* species.

Results and Discussion

Genomic features of *S. iniae* QMA0248

The genome of *S. iniae* QMA0248 consists of a single circular chromosome of

2,116,570 bp with no plasmids (Figure 1 and Table 1). The QMA0248

chromosome has an average GC content of 36.8 %, consistent with the other four

S. iniae genomes (SF1, YSFST01-82, ISET0901, and ISNO) and in common with

several other *Streptococcus* spp. such as *S. agalactiae* (35.6%) (14), *S. pneumoniae*

(39.7%) (15), and *S. pyogenes* (38.5%) (16). Of note, there is a high degree of

strand bias in the genome of *S. iniae* QMA0248 where genes are preferentially

oriented in the leading strand, which is typical for Firmicutes (Figure 1). Only

seven percent of QMA0248 protein-coding genes are annotated as hypothetical

proteins compared with a quarter in SF1 and 11-14% in the other three published

genomes: YSFST01-82, ISET0901, and ISNO. There are 68 pseudogenes identified

in QMA0248 (GenBank assembly accession: CP022392.1), most due to

interruption by IS (Figure 1, Supplementary Table S1). This is approximately five

times greater than the number of pseudogenes predicted in SF1. Collectively,

these differences between the compared genomes QMA0248, SF1, YSFST01-82,

ISET0901, and ISNO are likely to reflect different approaches to annotation. The

other 25 pseudogenes are caused by in-frame stop codons or frame-shifts, all

supported by additional mapping of Illumina reads of the strain QMA0248 against

its PacBio assembly.

QMA0248 has a single CRISPR locus (Clustered regularly interspaced short

palindromic repeats) (Figure 1), which harbors a tandem array of 15 identical 36

bp repeats, separated by 14 distinct 30 bp spacers, which is about double the size

of the CRISPR array in SF1, ISET0901, and ISNO (Supplementary Figure S1).

Upstream of QMA0248 CRISPR are four Cas genes, *csn2*, *cas2*, *cas1*, and *cas9*, which

alongside the CRISPR locus provide adaptive immunity against foreign DNA (e.g.

phage and plasmids) (17).

QMA0248 harbors 58 tRNA genes and 15 rRNA loci, consisting of 5S, 16S, and 23S

genes, arranged in five loci. In contrast, there is one rRNA operon fewer in SF1,

ISET0901, and ISNO than in QMA0248 (Table 1). Furthermore, during the

preparation of this manuscript, a PacBio complete genome was published for the

strain *S. iniae* 89353 (accession: CP017952) which has an identical rRNA

arrangement to QMA0248 except that one rRNA locus encodes a ~7 Kb tandem

duplication (i.e. six loci in total) (18). Such intra-species variation in the number

of rRNA genes (and tRNA genes) is not uncommon in bacteria (including

streptococci) (19, 20), and prompted us to investigate further.

Confirmation of rRNA assembly in *S. iniae* QMA0248 genome

PacBio reads for QMA0248 were initially assembled into a large ~2 Mb contig

representing most of the chromosome of *S. iniae* QMA0248 and three contigs less

than 10 Kb in length. The short contigs appeared to be single read chimeras that

were discarded from the final assembly. However, the identification of the tandem

rRNA region in *S. iniae* 89353 prompted us to review the assembled short contigs.

One of these ~7 Kb discarded contigs encoded an rRNA region (5S, 16S, and 23S

genes in tandem with an intervening cluster of tRNA genes). Subsequent

reassembly and visualization of mapped raw reads indicated that the additional

rRNA contig could be placed in three of the five rRNA operon locations to form a

~13 Kb putative tandem duplication of 5S, 16S and 23 rRNA genes as seen in the

89353 genome (18). Closer examination of the read pileup for the ~7 Kb rRNA

contig revealed that the tandem duplication was not well supported by

overlapping reads (Supplementary Figure S2) suggesting that it may be a chimeric

assembly of reads from more than one rRNA locus. To investigate the location of

the putative tandem rRNA duplication in *S. iniae* QMA0248 we carried out long-

range PCR across each of the five potential rRNA loci in the chromosome. PCR

revealed no tandem rRNA duplication in any locus (Supplementary Figure S3).

Never-the-less, we cannot rule out that during the culturing step, prior to DNA

extraction and genome sequencing, there existed a sub-population of QMA0248

cells with the tandem rRNA duplication. This would be consistent with finding

only a small number of reads spanning the tandem repeat. Similar doubt exists

over the tandem rRNA repeat in 89353, which should also be investigated at the

read-level.

Characterization of large mobile genetic elements in *S. iniae* QMA0248

Our investigation of the rRNA discrepancies in *S. iniae* SF1, ISET0901, and ISNO

also revealed differences in the mobile genetic element (MGE) content within the

available complete genomes. The chromosome of QMA0248 has a single ~12 Kb

genomic island (GI-Leu) inserted within the tRNA-Leu downstream of a large

number of consecutive ribosomal genes (Table 2). GI-Leu encodes an integrase at

its 5' end (QMA0248_0125), which is predicted to be responsible for the GI

insertion. A ~2.8 Kb region at the 5' end of the GI (87881–90708) is homologous

to the fish pathogen *S. parauberis* KCTC 11537, with 90% nucleotide sequence

identity, including part of the integrase, plasmid replication gene, and two

hypothetical proteins. The island encodes a collagen-binding surface protein (Cna,

B-type domain), which is a virulence factor with an LPxTG cell wall anchor motif,

a conserved Gram-positive *cocci* pentapeptide (21). An IS*30* family transposon

insertion truncates the collagen-binding domain encoded by *cna*, likely leading to

loss of functionality in QMA0248. This adhesin has been shown to play an

important pathogenic role in *Staphylococcus aureus* by facilitating bacterial cells

adherence to host collagen (22). Most of this GI appears to have been deleted in

the genomes of SF1, ISET0901, and ISNO only, along with the rRNA operon

upstream of it (Figure 3), which explains the difference with QMA0248 in total

number of rRNA and tRNA genes (Table 1).

The genome of *S. iniae* QMA0248 harbors a single ~32 Kb incomplete phage (Phi1)

(1991661–2023508), including a 5' integrase gene (QMA0248_1936), inserted

upstream of a tRNA-Cys gene (Table 2). A total of 44 genes encoding phage

proteins were identified, including genes involved in DNA replication such as DNA

polymerase III, tail morphogenesis, as well as host lysis such as holin and lysin, in

addition to 24 phage hypothetical proteins. More than half of the genes encoding

phage proteins carried by QMA0248 are homologous to proteins encoded by

temperate bacteriophage *Streptococcus* PH10 (56.8% according to PHAST) (23).

Furthermore, Phi1 in QMA0248 exhibits a remarkable nucleotide sequence

identity (99%) to a prophage encoded within the SF1 genome in the same locus,

whereas it is entirely absent in YSFST01-82, ISET0901, and ISNO (Figure 3).

Characterization of *S. iniae* QMA0248 insertion sequences

Insertion sequences (IS) were analyzed in the *S. iniae* QMA0248 genome using the

ISFinder database coupled with manual curation. The analysis revealed 92 IS

(Table 3), which is higher than the average number per bacterial genome (n=38)

but consistent with the lifestyle of *S. iniae* as a facultative pathogen (24, 25).

Furthermore, the number of IS found in *S. iniae* QMA0248 is substantially higher

than other Streptococci such as *S. mitis* strain B6 (n=63) but comparable to that of

the Gram positive fish pathogen *Lactococcus garvieae* (26, 27). The 92 IS elements

belong to 7 different IS families and 20 IS types. These include 9 novel types

belonging to IS*3*, IS*30*, IS*1182*, and IS*200*/IS*605* families, which we have submitted

to the ISFinder database (IS*Stin2*–IS*Stin10*) (Table 3 and Supplementary Tables

S2–4). Around half of all IS copies in QMA0248 belong to these 9 novel types

consistent with expansion of *S. iniae* specific IS since speciation (Table 3).

Amongst those genes disrupted by IS in QMA0248 is the restriction enzyme

component of a type II Restriction Methylation system that probably recognizes

"GCNGC" (28). This insertion renders the cognate methyltransferase (MTase)

(QMA0248_0516) an orphan and, given the high number of GCNGC sites

(3814/Mb in QMA0248) across the genome, suggests a potential role for this

MTase in global gene regulation. By comparison, there are 7762/Mb available

GATC sites in the *E. coli* K-12 MG1655 genome available for methylation by Dam,

the archetypal orphan type II MTase known to play a major role in gene regulation

(29). Further discussion of the methylome data generated by PacBio sequencing

of QMA0248 is provided as Supplementary Information. . Taken altogether, the *S.*

iniae genome harbors a large repertoire of IS elements, which may be associated

with adaptation to host or environment. Indeed, it is well accepted that IS

expansion is an early sign of genome reduction as a mechanism of adaptation to

host (25, 30, 31).

Phylogenetic and comparative analysis of *S. iniae* QMA0248, SF1, YSFST01-

82, ISET0901, and ISNO

The core genome of *S. iniae* QMA0248, SF1, YSFST01-82, ISET0901, and ISNO

accounts for ~75% of the chromosome. IS were compared between the reference

chromosome of QMA0248 and each of the *S. iniae* chromosomes (SF1, YSFST01-

82, ISET0901, and ISNO) using Artemis Comparison Tool (ACT) (32). Although IS

elements typically result in genomic rearrangements and loss of synteny, this is

not seen in *S. iniae.* This lack of rearrangement is reflected by the consistent

pattern of GC skew in the genome of *S. iniae* QMA0248 (Figure 1). Eight IS copies

out of the 92 detected in QMA0248 are absent in the genomes of SF1, ISET0901

and ISNO only (Supplementary Table S2). Other IS elements are unique to the

genomes SF1, ISET0901 and ISNO in syntenic positions. An interesting example is

an IS*981* (SF1 locus_tag: K710_0799 and K710_0800), inserted in *cas9* gene in the

CRISPR/Cas region only in those three genomes (Figure 2). Another example is

IS*Stin1*, inserted only in SF1, ISET0901, and ISNO (SF1 locus_tag: K710_0761).

Additionally, three IS copies are present in syntenic positions only in QMA0248,

ISET0901, and ISNO. Eight insertions are absent in YSFST01-82 only and another

seven IS copies are found exclusive to QMA0248 (Supplementary Table S2).

In addition to the IS differences, the five *S. iniae* genomes have major differences

in the content of other mobile genetic elements (MGE) that reflect variations in

the length of their respective chromosomes (Table 1). These MGEs include two

prophages (Phi1 and Phi2), a genomic island (GI), and two other regions of

difference (ROD) (Table 2, Figure 3). This includes a ~28 Kb region that is only

found in YSFST01-82 (ROD1), and a ~20 Kb region that is present in four genomes

but almost entirely absent from YSFST01-92 (ROD2) (Table 2, Figure 3).

Most variations in MGEs (including IS) were found to be incongruent with the core

SNP phylogeny. For instance, the deleted ~12 Kb GI-Leu and the *cas9* gene

disrupted by IS*981* exist only in SF1, ISET0901, and ISNO (Figures 2 and 3), but

these three isolates appear on divergent branches, indicating potential

independent events (Supplementary Figure S4). To investigate the discrepancies

between MGEs and phylogeny we compared multiple phylogenetic trees that were

constructed using different methods, including the core genome, core SNP and

using different software (Supplementary Figure S4). All phylogenies consistently

revealed that *S. iniae* isolates QMA0248 and SF1 cluster together in one clade,

whereas ISET0901 and ISNO cluster in another, and all four isolates cluster

separately to YSFST01-82, the latter diverging earliest from the root

(Supplementary Figure S4).

Discrepancies between the *S. iniae* genomes are likely due to misassembly

Taken together, our comparative analyses suggest that one or more of the *S. iniae*

genomes under comparison have been misassembled. The genomes ISET0901 and

ISNO were both assembled from Illumina data using BioNumerics (Applied Math)

and the genome *S. iniae* SF1 as a reference (12, 13). *S. iniae* SF1 was assembled *de*

novo from a combination of 454 GS FLX+, Illumina MiSeq and Sanger sequencing

(12). During the preparation of the present manuscript SF1 was removed from the

NCBI RefSeq database (https://www.ncbi.nlm.nih.gov/refseq/). YSFST01-92 was

also a hybrid assembly (454 GS FLX Titanium, Opgen optical mapping and Sanger

sequencing) but it remains in the RefSeq database. Reference-guided assembly

was introduced to enable comparisons between two very closely related isolates.

However, this practice can result in the erroneous inclusion of MGEs that exist in

the template genome but are absent from the comparison strain. Even with careful

curation it is impossible to avoid misplacing repetitive sequences such as IS, as

observed here in the case of *cas9* insertion and the other 8 IS copies that are absent

in SF1, ISET0901, and ISNO only. Moreover, reference-guided assemblies may

result in the loss of novel regions that are only present in the newly sequenced

strain, in which case a *de novo* approach is always required (33). Although

reference-guided assembly is no longer generally accepted for prokaryote

genomes, a number of examples remain available in public repositories such as

GenBank. For both ISET0901 and ISNO the assembly strategy is clearly outlined in

the comment field of the GenBank file, and the primary publications (12, 13).

Never-the-less, the consequences of using such genomes in downstream analyses

may not be apparent to all (for example, all three genomes are available in widely

used genome databases such as PATRIC (www.patricbrc.org version 3.5.36)(34)).

Removal of some early hybrid 454 complete genomes from public repositories like

RefSeq should help maintain the quality of available complete genomes. Long-read

sequencing data from Pacific Biosciences and Oxford Nanopore bring complete

bacterial genomes within reach of most laboratories, but here also significant care

is often required to avoid misassembly. Furthermore, as illustrated here and in

other studies (35, 36), what appear to be misassemblies may in fact be biologically

relevant. Ultimately the onus is on the user of public data to exercise caution when

validating the source, assembly strategy, and quality of available complete

genomes.

Conclusions

We assembled and annotated a high quality complete genome sequence for *S. iniae*

QMA0248, including manual curation of 92 insertion sequences. Comparative

analysis with publicly available complete genomes *S. iniae* SF1, YSFST01-82,

ISET0901, and ISNO revealed discrepancies in the MGE content consistent with

errors introduced by reference guided assembly of ISNO and ISET0901. Such

problems are not new, but many bacterial genomes assembled in this way remain

in public repositories of complete genomes. Our results emphasize the need to

critically appraise complete genome assemblies prior to comparative analysis.

Despite long-read sequencing becoming the gold standard for complete genome

assembly of bacterial isolates, caution is needed to avoid misassembly. To better

understand how insertion sequences, genomic islands, and other mobile elements

contribute to *S. iniae* diversity, there is a need for larger genomic studies using

global collections of *S. iniae* isolates from dissimilar origins. The genome of *S. iniae*

QMA0248 represents an important resource for future *S. iniae* comparative

genomic and evolutionary studies.

Materials and Methods

Bacterial strain and sequencing

Streptococcus iniae strain QMA0248 was isolated from diseased barramundi

(*Lates calcarifer*) from a farm in South Australia in 2009 (37). Genomic DNA was

prepared from several well-isolated colonies of *S. iniae* QMA0248 grown for 18 h

on Todd-Hewitt agar from a master seed stock (non-subcultured) with the

Genomic Tip 20 kit (Qiagen). Pre-incubation for 2 h at 37°C of cells suspended in

500 μL 50 mM EDTA containing 200 units of mutanolysin and 2 mg lysozyme was

found to improve cell lysis prior to following the manufacturers protocol for

purification of high molecular weight DNA. The genome of the strain QMA0248

was sequenced using 3 SMRT cells on the Pacific Biosciences (PacBio) RS II

platform and P4C2 sequencing chemistry, which generated a total of 57,083 reads

with an average length of 6,178 bp. Reads were deposited at BioProject

[PRJNA385746](#) under accession [SRP109617](#). The genome of the strain QMA0248

was also sequenced using Illumina Nextera libraries on HiSeq2000.

Genome assembly and modified bases detection

PacBio sequencing reads derived from *S. iniae* QMA0248 genomic DNA were

assembled using HGAP (hierarchical genome assembly process) using the PacBio

Single Molecule Real Time (SMRT) Portal (V2.3.0) (38), with default settings and

minimum seed read length of 4,509 bp. Contiguity was used to visualize the assembly and the overlap between contigs using BLASTn (39, 40). The resulting assembly was used as a reference in the RS Resequencing module of PacBio's SMRT Analysis v2.3.0 to map the raw reads onto the reference genome producing a highly accurate genome consensus. Illumina-sequenced reads of the strain QMA0248 were mapped to the PacBio assembly using Snippy v3.0 (https://github.com/tseemann/snippy). To analyze read pileup in the potential rRNA tandem operon, raw reads of QMA0248 were mapped onto the ~7 Kb rRNA contig using BLASR v2.2, as part of the PacBio's SMRT Analysis Suite, and visualized using Artemis (41). Methylated DNA bases were identified in the resulting genome assembly using the RS Modification and Motif Analysis protocol and Motif Finder v1 within the SMRT Analysis suite v2.3.0 using Quality Value (QV) cutoff of 30. DNA Methyltransferases (MTases) were identified using nucleotide comparisons against REBASE (42).

PCR to investigate the rRNA tandem duplication

To investigate the presence of a rRNA tandem repeat, long range PCR from unique adjacent region to unique adjacent region was performed using specific primers for each of 5 rRNA regions (Supplementary Table S1), which were designed using Primer3 (43). PCR was done using LongAmp® Taq PCR Kit (NEB) from 40 ng of QMA0248 *S. iniae* genomic DNA as follows: 5 min at 94°C; 30 cycles of (30 sec at 94°C, 30 sec at 56°C, and 15 min at 65°C); and 10 min final extension at 65°C. The gel was loaded with 5 μL QUICK-LOAD® 1 Kb Extend DNA Ladder and 1μL of PCR products and run for 90 min at 70V using 0.7% TAE buffer solution and stained with HydraGreen.

Genome and mobile genetic elements annotation

Automated genome annotation was done using Prokka v1.11 (Prokaryotic

Genome Annotation System) (44) and then manually curated using Artemis (41)

and Geneprimp (45). The start codons of known coding sequences (CDS), such as

in the capsule and streptolysin S operons, were further adjusted where

appropriate using UniProtKB (http://www.uniprot.org/) and Pfam (46). CRISPRs

(Clustered Regularly Interspaced Short Palindromic Repeats) were predicted

using CRISPR finder tool (http://crispr.u-psud.fr/crispr/) (47). Prophage

annotation was done using PHAST (Phage Search Tool)

(http://phast.wishartlab.com/) (48). Island Viewer

http://www.pathogenomics.sfu.ca/islandviewer/) was used to predict genomic

islands (GI) (49). Boundaries of phage and GI regions were manually adjusted to

their respective attachment sites. IS Saga (http://www-genome.biotoul.fr/) was

used for the initial identification of insertion sequences (IS). Additional manual

curation was carried out to confirm the boundaries of complete and partial IS

elements using the IS Finder database (http://www-is.biotoul.fr/). IS element

matches against the database that were at ≥95% nucleotide identity were

assigned the top matching IS name. IS elements of less percent identity are novel

and were assigned the names IS*Stin2*–IS*Stin10* by ISFinder. The impact of IS

elements on flanking coding sequences was analyzed using Artemis and by

searching the amino acid sequence in UniProt KB (http://www.uniprot.org/), and

Pfam databases (41, 46). The complete annotated genome sequence was

deposited at GenBank under the accession number CP022392.

Comparative genomics and phylogenetic analysis

Alignments of the whole-genome or genomic sub-regions, such as the CRISPR and

genomic islands, were done using BLASTn implemented in EasyFig v2.1 (40, 50).

Detailed analysis of regions of difference and comparison of insertion sequences

(IS) were done using Artemis Comparison Tool (ACT) (32). The core genome of

the five genomes QMA0248, SF1, YSFST01-82, ISET0901, and ISNO was defined

using Roary (51). Phylogenetic trees were constructed using the core genome and

the core SNP methods, using multiple programs (see below), using the strain

QMA0140 as the out-group (52). The strain *S. iniae* QMA0140 is a dolphin isolate

from USA in 1976, and was sequenced using Illumina HiSeq 2000 (See Bioproject

PRJNA417543). For quality control, the first 20 bp of each read derived from

QMA0140 genomic DNA were hard trimmed using Nesoni v0.132

(https://github.com/Victorian-Bioinformatics-Consortium/nesoni) with a

minimum length and quality of 70, and 20, respectively. Hard trimmed filtered

reads of QMA0140 were assembled using SPAdes v3.9.0 where contigs < 10X

coverage and smaller than 100 bp were removed (53). For core genome

phylogenies, whole-genome alignment of the five complete genomes and

QMA0140 draft assembly was done using Mauve v2.4.0 and Parsnp v1.2 with

default parameter settings (54, 55). For the alignment using Mauve, conserved

blocks in all 6 genomes longer than 500 bp were selected and concatenated using

the stripSubsetLCBs script producing the core genome alignment. For core SNP

phylogenies, error-free simulated reads were created using wgsim v0.3.2

(https://github.com/lh3/wgsim) and mapped to the reference genome QMA0248

along with QMA0140 hard trimmed and filtered raw reads using bowtie v1.0.0

(56), where variants were called using Nesoni v0.132

(https://github.com/Victorian-Bioinformatics-Consortium/nesoni) using default

parameters. Core SNPs were also identified by mapping the reads of the 6

genomes to QMA0248 using BWA-MEM v0.7.15 (r1140), implemented in Snippy

v3.0 (https://github.com/tseemann/snippy) (57). All trees were produced by

RAxML v8.2.9 (58) using the general time-reversible (GTR) and GAMMA

distribution model of among-site rate variation with bootstrapping from 1000

replicates, and viewed using FigTree v1.4.0

(http://tree.bio.ed.ac.uk/software/figtree).

1.   Agnew W, Barnes AC. Streptococcus iniae: an aquatic pathogen of global

veterinary significance and a challenging candidate for reliable vaccination.

Veterinary microbiology. 2007;122(1-2):1-15.

2.   Shoemaker CA, Klesius PH, Evans JJ. Prevalence of Streptococcus iniae in

tilapia, hybrid striped bass, and channel catfish on commercial fish farms in the

United States. American journal of veterinary research. 2001;62(2):174-7.

3.   Lau SK, Woo PC, Luk W-k, Fung AM, Hui W-t, Fong AH, et al. Clinical

isolates of Streptococcus iniae from Asia are more mucoid and β-hemolytic than

those from North America. Diagnostic microbiology and infectious disease.

2006;54(3):177-81.

4.   Baiano J, Barnes AC. Towards control of Streptococcus iniae. Emerg Infect

Dis. 2009;15(12):1891.

5.   Frost LS, Leplae R, Summers AO, Toussaint A. Mobile genetic elements:

the agents of open source evolution. Nature reviews Microbiology.

2005;3(9):722.

6.   van Elsas JD, Bailey MJ. The ecology of transfer of mobile genetic

elements. FEMS microbiology ecology. 2002;42(2):187-97.

7.   Ochman H, Lawrence JG, Groisman EA. Lateral gene transfer and the

nature of bacterial innovation. Nature. 2000;405(6784):299-304.

8.   Partridge SR. Analysis of antibiotic resistance regions in Gram-negative

bacteria. FEMS microbiology reviews. 2011;35(5):820-55.

9.   Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their

genomic impact and diversity. FEMS microbiology reviews. 2014;38(5):865-91.

10.   Cheng S, Hu YH, Jiao XD, Sun L. Identification and immunoprotective

analysis of a Streptococcus iniae subunit vaccine candidate. Vaccine.

2010;28(14):2636-41.

11.   Rajoo S, Jeon W, Park K, Yoo S, Yoon I, Lee H, et al. Complete genome

sequence of Streptococcus iniae YSFST01-82, isolated from olive flounder in Jeju,

South Korea. Genome announcements. 2015;3(2):e00319-15.

12.   Pridgeon JW, Zhang D, Zhang L. Complete genome sequence of a virulent

strain, Streptococcus iniae ISET0901, isolated from diseased tilapia. Genome

announcements. 2014;2(3):e00553-14.

13.   Pridgeon JW, Zhang D, Zhang L. Complete genome sequence of the

attenuated novobiocin-resistant Streptococcus iniae vaccine strain ISNO.

Genome announcements. 2014;2(3):e00510-14.

14.   Glaser P, Rusniok C, Buchrieser C, Chevalier F, Frangeul L, Msadek T, et al.

Genome sequence of Streptococcus agalactiae, a pathogen causing invasive

neonatal disease. Molecular microbiology. 2002;45(6):1499-513.

15.   Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, et al.

Complete genome sequence of a virulent isolate of Streptococcus pneumoniae.

Science. 2001;293(5529):498-506.

16.   Ferretti JJ, McShan WM, Ajdic D, Savic DJ, Savic G, Lyon K, et al. Complete

genome sequence of an M1 strain of Streptococcus pyogenes. Proceedings of the

National Academy of Sciences. 2001;98(8):4658-63.

17.   Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, et

al. CRISPR provides acquired resistance against viruses in prokaryotes. Science.

2007;315(5819):1709-12.

18.   Gong H-Y, Wu S-H, Chen C-Y, Huang C-W, Lu J-K, Chou H-Y. Complete

Genome Sequence of Streptococcus iniae 89353, a Virulent Strain Isolated from

Diseased Tilapia in Taiwan. Genome Announcements. 2017;5(4):e01524-16.

19.   Acinas SG, Marcelino LA, Klepac-Ceraj V, Polz MF. Divergence and

redundancy of 16S rRNA sequences in genomes with multiple rrn operons.

Journal of bacteriology. 2004;186(9):2629-35.

20.   Lim K, Furuta Y, Kobayashi I. Large variations in bacterial ribosomal RNA

genes. Molecular biology and evolution. 2012;29(10):2937-48.

21.   Fischetti V, Pancholi V, Schneewind O. Conservation of a hexapeptide

sequence in the anchor region of surface proteins from gram‐positive cocci.

Molecular microbiology. 1990;4(9):1603-5.

22.   Switalski LM, Patti JM, Butcher W, Gristina AG, Speziale P, Höök M. A

collagen receptor on Staphylococcus aureus strains isolated from patients with

septic arthritis mediates adhesion to cartilage. Molecular microbiology.

1993;7(1):99-107.

23.   van der Ploeg JR. Genome sequence of the temperate bacteriophage PH10

from Streptococcus oralis. Virus genes. 2010;41(3):450-8.

24.   Aziz RK, Breitbart M, Edwards RA. Transposases are the most abundant,

most ubiquitous genes in nature. Nucleic Acids Res. 2010;38(13):4207-17.

25.   Ochman H, Davalos LM. The nature and dynamics of bacterial genomes.

Science. 2006;311(5768):1730-3.

26.   Denapaite D, Brückner R, Nuhn M, Reichmann P, Henrich B, Maurer P, et

al. The genome of Streptococcus mitis B6-what is a commensal? PLoS One.

2010;5(2):e9426.

27.   Eraclio G, Ricci G, Fortina MG. Insertion sequence elements in Lactococcus

garvieae. Gene. 2015;555(2):291-6.

28.   Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA

restriction and modification: enzymes, genes and genomes. Nucleic acids

research. 2009;38(suppl_1):D234-D6.

29.   Lobner-Olesen A, Skovgaard O, Marinus MG. Dam methylation:

coordinating cellular processes. Curr Opin Microbiol. 2005;8(2):154-60.

30.   Moran NA, Plague GR. Genomic changes following host restriction in

bacteria. Current opinion in genetics & development. 2004;14(6):627-33.

31.   Richards VP, Lang P, Bitar PDP, Lefébure T, Schukken YH, Zadoks RN, et al.

Comparative genomics and the role of lateral gene transfer in the evolution of

bovine adapted Streptococcus agalactiae. Infection, Genetics and Evolution.

2011;11(6):1263-75.

32.   Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG,

Parkhill J. ACT: the Artemis comparison tool. Bioinformatics. 2005;21(16):3422-

3.

33.   Nijkamp J, Winterbach W, Van den Broek M, Daran J-M, Reinders M, De

Ridder D. Integrating genome assemblies with MAIA. Bioinformatics.

2010;26(18):i433-i9.

34.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, et al.

Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis

Resource Center. Nucleic Acids Res. 2017;45(D1):D535-D42.

35.   Forde BM, Ben Zakour NL, Stanton-Cook M, Phan MD, Totsika M, Peters

KM, et al. The complete genome sequence of Escherichia coli EC958: a high

quality reference sequence for the globally disseminated multidrug resistant E.

coli O25b:H4-ST131 clone. PLoS One. 2014;9(8):e104400.

36.   Draper JL, Hansen LM, Bernick DL, Abedrabbo S, Underwood JG, Kong N,

et al. Fallacy of the Unique Genome: Sequence Diversity within Single

Helicobacter pylori Strains. MBio. 2017;8(1).

37.   Millard CM, Baiano JC, Chan C, Yuen B, Aviles F, Landos M, et al. Evolution

of the capsular operon of Streptococcus iniae in response to vaccination. Applied

and environmental microbiology. 2012;78(23):8219-26.

38.   Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al.

Nonhybrid, finished microbial genome assemblies from long-read SMRT

sequencing data. Nature methods. 2013;10(6):563-9.

39.   Sullivan MJ, Zakour NLB, Forde BM, Stanton-Cook M, Beatson SA.

Contiguity: contig adjacency graph construction and visualisation. PeerJ

PrePrints; 2015. Report No.: 2167-9843.

40.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment

search tool. Journal of molecular biology. 1990;215(3):403-10.

41.   Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream M-A, et al.

Artemis: sequence visualization and annotation. Bioinformatics.

2000;16(10):944-5.

42.   Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA

restriction and modification: enzymes, genes and genomes. Nucleic acids

research. 2009:gkp874.

43.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et

al. Primer3—new capabilities and interfaces. Nucleic acids research.

2012;40(15):e115-e.

44.   Seemann T. Prokka: rapid prokaryotic genome annotation.

Bioinformatics. 2014:btu153.

45.   Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A,

et al. GenePRIMP: a gene prediction improvement pipeline for prokaryotic

genomes. Nature methods. 2010;7(6):455-7.

46.   Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths‐Jones S, et al.

The Pfam protein families database. Nucleic acids research. 2004;32(suppl

1):D138-D41.

47.   Grissa I, Vergnaud G, Pourcel C. CRISPRFinder: a web tool to identify

clustered regularly interspaced short palindromic repeats. Nucleic acids

research. 2007;35(suppl 2):W52-W7.

48.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast phage

search tool. Nucleic acids research. 2011:gkr485.

49.   Dhillon BK, Laird MR, Shay JA, Winsor GL, Lo R, Nizam F, et al.

IslandViewer 3: more flexible, interactive genomic island discovery, visualization

and analysis. Nucleic acids research. 2015:gkv401.

50.   Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison

visualizer. Bioinformatics. 2011;27(7):1009-10.

51.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al.

Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics.

2015;31(22):3691-3.

52.   Baiano JC, Tumbol RA, Umapathy A, Barnes AC. Identification and

molecular characterisation of a fibrinogen binding protein from Streptococcus

iniae. BMC microbiology. 2008;8:67.

53.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. Journal of computational biology. 2012;19(5):455-77.

54.   Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of

conserved genomic sequence with rearrangements. Genome research.

2004;14(7):1394-403.

55.   Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid

core-genome alignment and visualization of thousands of intraspecific microbial

genomes. Genome biology. 2014;15(11):524.

56.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. Genome

biology. 2009;10(3):R25.

57.   Li H. Aligning sequence reads, clone sequences and assembly contigs with

BWA-MEM. arXiv preprint arXiv:13033997. 2013.

58.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 2014;30(9):1312-3.

59.   Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter:

circular and linear interactive genome visualization. Bioinformatics.

2009;25(1):119-20.

Table 1: General features of five *S. iniae* complete genomes.

Strain	QMA0248	SF1	YSFST01-82	ISET0901	ISNO
Accession Number	CP022392	CP005941	CP010783	CP007586	CP007587
Genome Size (bp)	2,116,570	2,149,844	2,086,959	2,070,822	2,070,182
GC Content (%)	36.8	36.7	36.8	36.8	36.8
Total CDS number	1,946	2,125	1,897	1,872	1,865
Total gene number	2,196	2,196	2,029	1,997	1,996
rRNA (5S, 16S, 23S)	15	12	15	12	12
tRNA	58	45	58	45	45
Reference	This study	(12)	(11)	(12)	(13)
Assembly type	PacBio RSII P2C4	454 FLX+/ Illumina MiSeq/ Sanger	454 FLX Titanium/ Opgen/ Sanger	Illumina 1500 HiSeq/ Reference guided assembly	Illumina 1500 HiSeq/ Reference guided assembly

594 **Table 2: Large mobile genetic elements (MGE) and regions of difference (ROD) identified in the 5 *S.***
595 ***iniae* genomes analyzed (QMA0248, SF1, YSFST01-82, ISET0901, and ISNO).**

Region	Genomic Island (GI-leu)	ROD1	Prophage 2	ROD2	Prophage 1 (Phi1)
Coordinates[1]	87374-100014	177819-206359 (YSFST01-82)	848479-890501 (SF1)	1767227-1787619	1991661-2023508
Length (Kb)	12.6	28.5	42.0	20.4	31.8
GC content (%)	35.9	37.2	35.3	34.3	37.6
Features	MGE; 13 tRNA and 1 rRNA operon (5S, 16S, and 23S) upstream; integrase; IS*30* and IS*256* family IS elements	Integrase; IS*3* and IS*256* family IS elements	MGE; integrase	IS*3* family IS element	MGE; integrase; 1 tRNA-Cys
No. of CDSs	11	33	63	18	53
Major CDSs	Cro/CI family transcriptional regulator; ECF subfamily RNA polymerase sigma factor; plasmid replication protein; membrane protein	ESAT-6-like protein; two-component sensor histidine kinase; galactose mutarotase; 3 PTS galactitol transporter subunits (IIA, IIC, and IIB)	Phage DNA replication protein; prophage antirepressor, phage capsid and scaffold protein; putative tail protein; holin; endolysin; antigen C; several phage hypothetical proteins	ESAT-6-like protein; O-glycosyl hydrolase; phage infection protein; 3 lipoproteins; protein kinase	DNA helicase; Cro/CI family transcriptional regulator; tail and capsid proteins; holin; lysin; DNA N-4 cytosine methyltransferase; site specific recombinase; several phage hypothetical proteins
Best hit (% identity, % coverage)	*S. parauberis* KCTC 11537 (90, 41)	*S. dygalactiae subsp. equisimilis ATCC 12394* (70, 34)	*S. parauberis* KCTC 11537 (91, 45)	*S. thermophilus* JIM 8232 (81, 13)	Bacteriophage PH10 of *Streptococcus* (71, 34)

596 [1]Coordinates are in QMA0248 GenGenBank annotation (GCA_002220115.1) unless otherwise indicated.

598 **Table 3: Summary of all insertion sequences (IS) identified in QMA0248. Partial IS are suffixed by –p.**

IS Family in QMA0248	Number of IS copies	IS types (copy number, mean % amino acid identity)
IS*3*	32	IS*Sag2* (10, 98.6), IS*Sag2*-p (2, 98.7)
		IS*981* (8, 99.8)
		IS*Spy1* (1, 86.7), IS*Spy1*-p (1, 90.5)
		*IS*Stin6* (3, 94.5), *IS*Stin7* (4, 90.0), *IS*Stin5* (2, 69.3), unclassified most similar to *IS*Stin5-p* (1, 73.8)
IS*30*	22	IS*Sag9* (3, 99.6), IS*Sag9*-p (5, 99.6)
		*IS*Stin4* (2, 93.3), *IS*Stin4*-p (2, 89.6), *IS*Stin9* (9, 81.4), *IS*Stin2* (1, 84.7)
IS*256*	17	IS*Stin1* (16, 91.0), unclassified most similar to IS*Stin1*-p (1, 90.2)
IS*1182*	13	*IS*Stin8* (7, 86.3), *IS*Stin8*-p (2, 89.8), *IS*Stin3* (1, 87.7), *IS*Stin3*-p (2, 87.1), unclassified most similar to *IS*Stin8-p* (1, 70.5)
IS*200*/IS*605*	5	*IS*Stin10* (3, 99), *IS*Stin10*-p (1, 98.6), unclassified most similar to *IS*Stin10* (1, 86.5)
IS*L3*	1	Unclassified most similar to IS*Sth1*-p (1, 77.8)
IS*110*	2	Unclassified most similar to IS*L4* (2, 65.8)

599 ** Novel IS element.*

Forward CDSs

Reverse CDSs **Pseudogenes**

rRNA **Insertion Sequences**

tRNA **GI, CRISPR/Cas, Phage**

Figure 1: Circular map of the genome *Streptococcus iniae* QMA0248. Genomic

features from outer ring to the inner ring are described in the key to the left, where

the innermost two rings correspond to the GC skew (inner), and GC plot (outer).

CDS: coding sequence. GI: genomic island. The circular map was generated using

DNAPlotter (59).

608

Figure 2: Comparison of the CRISPR/Cas region between QMA0248, SF1,

YSFST01-82, ISET0901, and ISNO. Alignment of Cas genes where the genomes

are ordered according to their position in the phylogenetic tree (left). The

maximum Likelihood (ML) phylogeny was rooted to QMA0140 (not shown) and

built using 1,111 SNPs. Arrows correspond to Cas genes, which are labeled at the

bottom. Figure was produced using EasyFig (50) using 500 bp as minimum length,

90% as minimum identity value, and 0.001 as maximum e-value.

Figure 3: Whole-genome alignment of the 5 genomes QMA0248, SF1,
YSFST01-82, ISET0901, and ISNO. The genomes are ordered according to their
position in the core SNP-based phylogenetic tree. The maximum Likelihood (ML)
phylogeny was rooted to QMA0140 (not shown) and built using 1,111 SNPs. The
scale bar indicates the number of substitutions represented by branch lengths.
BLASTn comparison was produced using EasyFig (50) using 2000 bp as minimum
length, 50% as minimum identity value, and 1×10^{-17} as maximum e-value.

Importance

The present manuscript provides *Streptococcus iniae* PacBio sequenced genome (QMA0248) with
complete characterization of its mobile genetic elements (MGE) content including genomic islands,
prophages, and insertion sequences (IS). The genome of *S. iniae* QMA0248 harbours 92 IS elements
including 9 novel ones belonging mostly to IS*3*, which were submitted to ISFinder database. Here we
also present the first analysis for *S. iniae* methylome. We also performed comparative and phylogenetic
analysis between *S. iniae* QMA0248 and publicly available complete *S. iniae* genomes, which to our
knowledge has not been done for *S. iniae* before. This analysis revealed discrepancies that are likely due
to misassembly in the genomes of strains ISET0901 and ISNO.Our study not only highlights assembly
problems in existing *S. iniae* genomes, but provides a high quality reference genome for *S.*
iniae QMA0248, including manually curated MGEs, that will assist future *S. iniae* comparative genomic
and evolutionary studies.