1   # A high quality reference genome for the fish pathogen

2   *Streptococcus iniae*

3
Areej S. Alsheikh-Hussain[1,2], Nouri L. Ben Zakour[1,2,3], Brian M. Forde[1,2],

Oleksandra Silayeva[4], Andrew C. Barnes[4], and Scott A. Beatson*[1,2].

[1]School of Chemistry & Molecular Biosciences, [2]Australian Infectious Diseases

Research Centre and [4]School of Biological Science, The University of Queensland,

Brisbane, Queensland, Australia;

Present address: [3]The Westmead Institute for Medical Research and the University

of Sydney, Sydney, New South Wales, Australia

Key words: SMRT sequencing, Insertion Sequence, reference guided assembly,

misassembly, mobile genetic elements

* Corresponding author.

Mailing address: School of Chemistry and Molecular Biosciences, Building 76, The

University of Queensland, Brisbane QLD 4072, Australia.

Phone: +617 33654863; E-mail: s.beatson@uq.edu.au

Abstract

Fish mortality caused by *Streptococcus iniae* is a major economic problem in fish

aquaculture in warm and temperate regions globally. There is also risk of zoonotic

infection by *S. iniae* through handling of contaminated fish. In this study, we

present the complete genome sequence of *S. iniae* strain QMA0248, isolated from

farmed barramundi in South Australia. The 2.12 Mb genome of *S. iniae* QMA0248

carries a 32 Kb prophage, a 12 Kb genomic island, and 92 discrete insertion

sequence (IS) elements. These include 9 novel IS types that belong mostly to the

IS*3* family. Comparative and phylogenetic analysis between *S. iniae* QMA0248 and

publicly available complete *S. iniae* genomes revealed discrepancies that are likely

due to misassembly in the genomes of isolates ISET0901 and ISNO. We also

determined by long-range PCR that a tandem duplication of an rRNA region in the

PacBio assembly of QMA0248 was an assembly error. A similar rRNA duplication

in the PacBio genome of *S. iniae* 89353 may also be a misassembly. Our study not

only highlights assembly problems in existing genomes, but provides a high

quality reference genome for *S. iniae* QMA0248, including manually curated

mobile genetic elements, that will assist future *S. iniae* comparative genomic and

evolutionary studies.

Introduction

45 *Streptococcus iniae* is a fish pathogen that causes mortality in a wide range of fish

46 species in wild and farmed, marine and freshwater environments, resulting in

47 large economic losses to aquaculture (1, 2). *S. iniae* is also considered an

48 opportunistic human pathogen, causing sporadic infections mostly in the elderly

49 who have more than one underlying health condition such as diabetes mellitus, or

chronic rheumatic heart disease (3, 4). *S. iniae* pathogenesis is imparted through

a repertoire of virulence factors (VF) including surface proteins, secreted toxins,

and capsular polysaccharide (CPS) (4). VFs can be acquired through lateral gene

transfer (LGT) of mobile genetic elements (MGE) such as composite transposons,

genomic islands (GI) or prophages.

MGEs are a means by which bacterial pathogens acquire traits that help adapt to

changing conditions including vaccination, antibiotics, a new host or environment

(5, 6). Indeed, they are considered the main drivers of gene flux in bacteria,

contributing to diversity within species (7). Insertion sequence (IS) elements, for

instance, are small MGEs (0.7-3.5 Kb) that have an important role in evolution and

genome plasticity. IS insertion within bacterial chromosomes or plasmids can

result in genetic modifications through insertional inactivation of genes or up-

regulation of adjacent intact genes through outward-facing promoter sequences

carried by some IS (8, 9). In some cases pairs of IS can mobilize intervening

sequence as a composite transposon (8). The mobility of IS elements leads to their

expansion or loss within bacterial lineages. Expansion is associated with

accumulation of pseudogenes, which is considered an early stage in genome

reduction as a mechanism for adaptation (9). Accordingly, to obtain the complete

evolutionary picture within bacterial species it is important to study the

distribution and abundance of IS elements.

As yet there is no study that focuses on the diversity and distribution of MGEs in

S. iniae genomes. In fact, only four complete *S. iniae* genomes were deposited in

Genbank at the commencement of the present study, none of which had

comprehensive annotations for MGEs: SF1, YSFST01-82, ISET0901, and ISNO. *S.*

iniae SF1 (accession: CP005941) was cultured from moribund flounder from a fish

farm that experienced an epidemic in North China (10). YSFST01-82 (accession:

CP010783) is an isolate from a diseased Olive flounder from South Korea (2012)

(11). ISET0901 (accession: CP007586) is reported to be a highly virulent strain,

which was isolated from diseased Nile tilapia during an outbreak in Israel (2005)

(12). The strain ISNO (accession: CP007587) was obtained through selection for

resistance of *S. iniae* ISET0901 to novobiocin (13).

There are currently 25 different IS families in the ISFinder database, but only one

IS described for *S. iniae* (IS*Stin1* of the family IS*256*). As is typical for complete

bacterial genomes, the annotation of the four complete *S. iniae* genomes available

at Genbank are limited to some of the transposase genes associated with IS

elements without definition of the IS boundaries. Indeed, the difficulty in

annotating IS elements and the lack of reliable automated annotation methods

means that only a small subset of complete genomes have accurate IS annotations.

Small partial IS elements are often disregarded although they reflect valuable

insights on ancestral recombination events. With the increasing availability of

long-read sequencing there is a need for a high quality, well-characterized *S. iniae*

reference genome that better enables the impact of IS elements on evolution and

diversity to be determined.

In this study, we have completely characterized the genome of *S. iniae* QMA0248.

Manual curation of annotations for IS, genomic islands, prophages, and CRISPR

was carried out along with a comparison with the four publicly available complete

genomes from NCBI (SF1, YSFST01-82, ISET0901, and ISNO). Comparative and

phylogenetic analyses revealed discrepancies between the MGE content

indicating likely misassembly in the genomes ISNO, and ISET0901. The complete

genome of QMA0248 will provide an important scaffold for future phylogenomic

studies of the *S. iniae* species.

Results and Discussion

Genomic features of *S. iniae* QMA0248

The genome of *S. iniae* QMA0248 consists of a single circular chromosome of

2,116,570 bp with no plasmids (Figure 1 and Table 1). The QMA0248

chromosome has an average GC content of 36.8 %, consistent with the other four

S. iniae genomes (SF1, YSFST01-82, ISET0901, and ISNO) and in common with

several other *Streptococcus* spp. such as *S. agalactiae* (35.6%) (14), *S. pneumoniae*

(39.7%) (15), and *S. pyogenes* (38.5%) (16). Of note, there is a high degree of

strand bias in the genome of *S. iniae* QMA0248 where genes are preferentially

oriented in the leading strand, which is typical for Firmicutes (Figure 1). Only

seven percent of QMA0248 protein-coding genes are annotated as hypothetical

proteins compared with a quarter in SF1 and 11-14% in the other three published

genomes: YSFST01-82, ISET0901, and ISNO. There are 68 pseudogenes identified

in QMA0248 (GenBank assembly accession: CP022392.1), most due to

interruption by IS (Figure 1, Supplementary Table S1). This is approximately five

times greater than the number of pseudogenes predicted in SF1. Collectively,

these differences between the compared genomes QMA0248, SF1, YSFST01-82,

ISET0901, and ISNO are likely to reflect different approaches to annotation. The

other 25 pseudogenes are caused by in-frame stop codons or frame-shifts, all supported by additional mapping of Illumina reads of the strain QMA0248 against its PacBio assembly.

QMA0248 has a single CRISPR locus (Clustered regularly interspaced short palindromic repeats) (Figure 1), which harbors a tandem array of 15 identical 36 bp repeats, separated by 14 distinct 30 bp spacers, which is about double the size of the CRISPR array in SF1, ISET0901, and ISNO (Supplementary Figure S1). Upstream of QMA0248 CRISPR are four Cas genes, *csn2*, *cas2*, *cas1*, and *cas9*, which alongside the CRISPR locus provide adaptive immunity against foreign DNA (e.g. phage and plasmids) (17).

QMA0248 harbors 58 tRNA genes and 15 rRNA loci, consisting of 5S, 16S, and 23S genes, arranged in five loci. In contrast, there is one rRNA operon fewer in SF1, ISET0901, and ISNO than in QMA0248 (Table 1). Furthermore, during the preparation of this manuscript, a PacBio complete genome was published for the strain *S. iniae* 89353 (accession: CP017952) which has an identical rRNA arrangement to QMA0248 except that one rRNA locus encodes a ~7 Kb tandem duplication (i.e. six loci in total) (18). Such intra-species variation in the number of rRNA genes (and tRNA genes) is not uncommon in bacteria (including streptococci) (19, 20), and prompted us to investigate further.

Confirmation of rRNA assembly in *S. iniae* QMA0248 genome

PacBio reads for QMA0248 were initially assembled into a large ~2 Mb contig

representing most of the chromosome of *S. iniae* QMA0248 and three contigs less

than 10 Kb in length. The short contigs appeared to be single read chimeras that

were discarded from the final assembly. However, the identification of the tandem

rRNA region in *S. iniae* 89353 prompted us to review the assembled short contigs.

One of these ~7 Kb discarded contigs encoded an rRNA region (5S, 16S, and 23S

genes in tandem with an intervening cluster of tRNA genes). Subsequent

reassembly and visualization of mapped raw reads indicated that the additional

rRNA contig could be placed in three of the five rRNA operon locations to form a

~13 Kb putative tandem duplication of 5S, 16S and 23 rRNA genes as seen in the

89353 genome (18). Closer examination of the read pileup for the ~7 Kb rRNA

contig revealed that the tandem duplication was not well supported by

overlapping reads (Supplementary Figure S2) suggesting that it may be a chimeric

assembly of reads from more than one rRNA locus. To investigate the location of

the putative tandem rRNA duplication in *S. iniae* QMA0248 we carried out long-

range PCR across each of the five potential rRNA loci in the chromosome. PCR

revealed no tandem rRNA duplication in any locus (Supplementary Figure S3).

Never-the-less, we cannot rule out that during the culturing step, prior to DNA

extraction and genome sequencing, there existed a sub-population of QMA0248

cells with the tandem rRNA duplication. This would be consistent with finding

only a small number of reads spanning the tandem repeat. Similar doubt exists

over the tandem rRNA repeat in 89353, which should also be investigated at the

read-level.

Characterization of large mobile genetic elements in *S. iniae* QMA0248

Our investigation of the rRNA discrepancies in *S. iniae* SF1, ISET0901, and ISNO

also revealed differences in the mobile genetic element (MGE) content within the

available complete genomes. The chromosome of QMA0248 has a single ~12 Kb

genomic island (GI-Leu) inserted within the tRNA-Leu downstream of a large

number of consecutive ribosomal genes (Table 2). GI-Leu encodes an integrase at

its 5' end (QMA0248_0125), which is predicted to be responsible for the GI

insertion. A ~2.8 Kb region at the 5' end of the GI (87881–90708) is homologous

to the fish pathogen *S. parauberis* KCTC 11537, with 90% nucleotide sequence

identity, including part of the integrase, plasmid replication gene, and two

hypothetical proteins. The island encodes a collagen-binding surface protein (Cna,

B-type domain), which is a virulence factor with an LPxTG cell wall anchor motif,

a conserved Gram-positive *cocci* pentapeptide (21). An IS*30* family transposon

insertion truncates the collagen-binding domain encoded by *cna*, likely leading to

loss of functionality in QMA0248. This adhesin has been shown to play an

important pathogenic role in *Staphylococcus aureus* by facilitating bacterial cells

adherence to host collagen (22). Most of this GI appears to have been deleted in

the genomes of SF1, ISET0901, and ISNO only, along with the rRNA operon

upstream of it (Figure 3), which explains the difference with QMA0248 in total

number of rRNA and tRNA genes (Table 1).

The genome of *S. iniae* QMA0248 harbors a single ~32 Kb incomplete phage (Phi1)

(1991661–2023508), including a 5' integrase gene (QMA0248_1936), inserted

upstream of a tRNA-Cys gene (Table 2). A total of 44 genes encoding phage

proteins were identified, including genes involved in DNA replication such as DNA

polymerase III, tail morphogenesis, as well as host lysis such as holin and lysin, in

addition to 24 phage hypothetical proteins. More than half of the genes encoding

phage proteins carried by QMA0248 are homologous to proteins encoded by

temperate bacteriophage *Streptococcus* PH10 (56.8% according to PHAST) (23).

Furthermore, Phi1 in QMA0248 exhibits a remarkable nucleotide sequence

identity (99%) to a prophage encoded within the SF1 genome in the same locus,

whereas it is entirely absent in YSFST01-82, ISET0901, and ISNO (Figure 3).

Characterization of *S. iniae* QMA0248 insertion sequences

Insertion sequences (IS) were analyzed in the *S. iniae* QMA0248 genome using the

ISFinder database coupled with manual curation. The analysis revealed 92 IS

(Table 3), which is higher than the average number per bacterial genome (n=38)

but consistent with the lifestyle of *S. iniae* as a facultative pathogen (24, 25).

Furthermore, the number of IS found in *S. iniae* QMA0248 is substantially higher

than other Streptococci such as *S. mitis* strain B6 (n=63) but comparable to that of

the Gram positive fish pathogen *Lactococcus garvieae* (26, 27). The 92 IS elements

belong to 7 different IS families and 20 IS types. These include 9 novel types

belonging to IS*3*, IS*30*, IS*1182*, and IS*200*/IS*605* families, which we have submitted

to the ISFinder database (IS*Stin2*–IS*Stin10*) (Table 3 and Supplementary Tables

S2–4). Around half of all IS copies in QMA0248 belong to these 9 novel types

consistent with expansion of *S. iniae* specific IS since speciation (Table 3).

Amongst those genes disrupted by IS in QMA0248 is the restriction enzyme

component of a type II Restriction Methylation system that probably recognizes

"GCNGC" (28). This insertion renders the cognate methyltransferase (MTase)

(QMA0248_0516) an orphan and, given the high number of GCNGC sites

(3814/Mb in QMA0248) across the genome, suggests a potential role for this

MTase in global gene regulation. By comparison, there are 7762/Mb available

GATC sites in the *E. coli* K-12 MG1655 genome available for methylation by Dam,

the archetypal orphan type II MTase known to play a major role in gene regulation

(29). Further discussion of the methylome data generated by PacBio sequencing

of QMA0248 is provided as Supplementary Information. . Taken altogether, the *S.*

iniae genome harbors a large repertoire of IS elements, which may be associated

with adaptation to host or environment. Indeed, it is well accepted that IS

expansion is an early sign of genome reduction as a mechanism of adaptation to

host (25, 30, 31).

Phylogenetic and comparative analysis of *S. iniae* QMA0248, SF1, YSFST01-

82, ISET0901, and ISNO

The core genome of *S. iniae* QMA0248, SF1, YSFST01-82, ISET0901, and ISNO

accounts for ~75% of the chromosome. IS were compared between the reference

chromosome of QMA0248 and each of the *S. iniae* chromosomes (SF1, YSFST01-

82, ISET0901, and ISNO) using Artemis Comparison Tool (ACT) (32). Although IS

elements typically result in genomic rearrangements and loss of synteny, this is

not seen in *S. iniae.* This lack of rearrangement is reflected by the consistent

pattern of GC skew in the genome of *S. iniae* QMA0248 (Figure 1). Eight IS copies

out of the 92 detected in QMA0248 are absent in the genomes of SF1, ISET0901

and ISNO only (Supplementary Table S2). Other IS elements are unique to the

genomes SF1, ISET0901 and ISNO in syntenic positions. An interesting example is

an IS*981* (SF1 locus_tag: K710_0799 and K710_0800), inserted in *cas9* gene in the

CRISPR/Cas region only in those three genomes (Figure 2). Another example is

IS*Stin1*, inserted only in SF1, ISET0901, and ISNO (SF1 locus_tag: K710_0761).

Additionally, three IS copies are present in syntenic positions only in QMA0248,

ISET0901, and ISNO. Eight insertions are absent in YSFST01-82 only and another

seven IS copies are found exclusive to QMA0248 (Supplementary Table S2).

In addition to the IS differences, the five *S. iniae* genomes have major differences

in the content of other mobile genetic elements (MGE) that reflect variations in

the length of their respective chromosomes (Table 1). These MGEs include two

prophages (Phi1 and Phi2), a genomic island (GI), and two other regions of

difference (ROD) (Table 2, Figure 3). This includes a ~28 Kb region that is only

found in YSFST01-82 (ROD1), and a ~20 Kb region that is present in four genomes

but almost entirely absent from YSFST01-82 (ROD2) (Table 2, Figure 3).

Most variations in MGEs (including IS) were found to be incongruent with the core

SNP phylogeny. For instance, the deleted ~12 Kb GI-Leu and the *cas9* gene

disrupted by IS*981* exist only in SF1, ISET0901, and ISNO (Figures 2 and 3), but

these three isolates appear on divergent branches, indicating potential

independent events (Supplementary Figure S4). To investigate the discrepancies

between MGEs and phylogeny we compared multiple phylogenetic trees that were

constructed using different methods, including the core genome, core SNP and

using different software (Supplementary Figure S4). All phylogenies consistently

revealed that *S. iniae* isolates QMA0248 and SF1 cluster together in one clade,

whereas ISET0901 and ISNO cluster in another, and all four isolates cluster

separately to YSFST01-82, the latter diverging earliest from the root

(Supplementary Figure S4).

Discrepancies between the *S. iniae* genomes are likely due to misassembly

Taken together, our comparative analyses suggest that one or more of the *S. iniae*

genomes under comparison have been misassembled. The genomes ISET0901 and

ISNO were both assembled from Illumina data using BioNumerics (Applied Math)

and the genome *S. iniae* SF1 as a reference (12, 13). *S. iniae* SF1 was assembled *de*

novo from a combination of 454 GS FLX+, Illumina MiSeq and Sanger sequencing

(12). During the preparation of the present manuscript SF1 was removed from the

NCBI RefSeq database. YSFST01-82 was also a hybrid assembly (454 GS FLX

Titanium, Opgen optical mapping and Sanger sequencing) but it remains in the

RefSeq database and is the designated genome for *S. iniae* at NCBI

(https://www.ncbi.nlm.nih.gov/genome/?term=streptococcus+iniae; accessed

23rd Jan 2020).

We have no reason to suspect that the YSFST01-82 genome assembly is inferior to

that of QMA0248, but adopting the latter as an alternative representative *S. iniae*

genome is justified for investigators wishing to take advantage of a manually

curated set of MGEs. In contrast we strongly recommend not using ISET0901 or

ISNO in future comparative studies of *S. iniae* genomes. Reference-guided

assembly was introduced to enable comparisons between two very closely related

isolates. However, this practice can result in the erroneous inclusion of MGEs that

exist in the template genome but are absent from the comparison strain. Even with

careful curation it is impossible to avoid misplacing repetitive sequences such as

IS, as observed here in the case of *cas9* insertion and the other 8 IS copies that are

absent in SF1, ISET0901, and ISNO only. Moreover, reference-guided assemblies

may result in the loss of novel regions that are only present in the newly

sequenced strain, in which case a *de novo* approach is always required (33).

Although reference-guided assembly is no longer generally accepted for

prokaryote genomes, a number of examples remain available in public

repositories such as GenBank. For both ISET0901 and ISNO the assembly strategy

is clearly outlined in the comment field of the GenBank file, and the primary

publications (12, 13). Never-the-less, the consequences of using such genomes in

downstream analyses may not be apparent to all (for example, all three genomes

are available in widely used genome databases such as PATRIC

(www.patricbrc.org version 3.5.36)(34)).

Removal of some early hybrid 454 complete genomes from public repositories like

RefSeq should help maintain the quality of available complete genomes. Long-read

sequencing data from Pacific Biosciences and Oxford Nanopore bring complete

bacterial genomes within reach of most laboratories, but here also significant care

is often required to avoid misassembly. Furthermore, as illustrated here and in

other studies (35, 36), what appear to be misassemblies may in fact be biologically

relevant. Ultimately the onus is on the user of public data to exercise caution when

validating the source, assembly strategy, and quality of available complete

genomes.

Conclusions

We assembled and annotated a high quality complete genome sequence for *S. iniae* QMA0248, including manual curation of 92 insertion sequences. Comparative analysis with publicly available complete genomes *S. iniae* SF1, YSFST01-82, ISET0901, and ISNO revealed discrepancies in the MGE content consistent with errors introduced by reference guided assembly of ISNO and ISET0901. Such problems are not new, but many bacterial genomes assembled in this way remain in public repositories of complete genomes. Our results emphasize the need to critically appraise complete genome assemblies prior to comparative analysis. Despite long-read sequencing becoming the gold standard for complete genome assembly of bacterial isolates, caution is needed to avoid misassembly. To better understand how insertion sequences, genomic islands, and other mobile elements contribute to *S. iniae* diversity, there is a need for larger genomic studies using global collections of *S. iniae* isolates from dissimilar origins. The genome of *S. iniae* QMA0248 represents an important resource for future *S. iniae* comparative genomic and evolutionary studies.

Materials and Methods

Bacterial strain and sequencing

Streptococcus iniae strain QMA0248 was isolated from diseased barramundi (*Lates calcarifer*) from a farm in South Australia in 2009 (37). Genomic DNA was prepared from several well-isolated colonies of *S. iniae* QMA0248 grown for 18 h on Todd-Hewitt agar from a master seed stock (non-subcultured) with the Genomic Tip 20 kit (Qiagen). Pre-incubation for 2 h at 37°C of cells suspended in

500 μL 50 mM EDTA containing 200 units of mutanolysin and 2 mg lysozyme was

found to improve cell lysis prior to following the manufacturers protocol for

purification of high molecular weight DNA. The genome of the strain QMA0248

was sequenced using 3 SMRT cells on the Pacific Biosciences (PacBio) RS II

platform and P4C2 sequencing chemistry, which generated a total of 57,083 reads

with an average length of 6,178 bp. Reads were deposited at BioProject

PRJNA385746 under accession SRP109617. The genome of the strain QMA0248

was also sequenced using Illumina Nextera libraries on HiSeq2000.

Genome assembly and modified bases detection

PacBio sequencing reads derived from *S. iniae* QMA0248 genomic DNA were

assembled using HGAP (hierarchical genome assembly process) using the PacBio

Single Molecule Real Time (SMRT) Portal (V2.3.0) (38), with default settings and

minimum seed read length of 4,509 bp. Contiguity was used to visualize the

assembly and the overlap between contigs using BLASTn (39, 40). The resulting

assembly was used as a reference in the RS Resequencing module of PacBio's

SMRT Analysis v2.3.0 to map the raw reads onto the reference genome producing

a highly accurate genome consensus. Illumina-sequenced reads of the strain

QMA0248 were mapped to the PacBio assembly using Snippy v3.0

(https://github.com/tseemann/snippy). To analyze read pileup in the potential

rRNA tandem operon, raw reads of QMA0248 were mapped onto the ~7 Kb rRNA

contig using BLASR v2.2, as part of the PacBio's SMRT Analysis Suite, and

visualized using Artemis (41). Methylated DNA bases were identified in the

resulting genome assembly using the RS Modification and Motif Analysis protocol

and Motif Finder v1 within the SMRT Analysis suite v2.3.0 using Quality Value

(QV) cutoff of 30. DNA Methyltransferases (MTases) were identified using

nucleotide comparisons against REBASE (42).

PCR to investigate the rRNA tandem duplication

To investigate the presence of a rRNA tandem repeat, long range PCR from unique

adjacent region to unique adjacent region was performed using specific primers

for each of 5 rRNA regions (Supplementary Table S6), which were designed using

Primer3 (43). PCR was done using LongAmp® Taq PCR Kit (NEB) from 40 ng of

QMA0248 *S. iniae* genomic DNA as follows: 5 min at 94°C; 30 cycles of (30 sec at

94°C, 30 sec at 56°C, and 15 min at 65°C); and 10 min final extension at 65°C. The

gel was loaded with 5 μL QUICK-LOAD® 1 Kb Extend DNA Ladder and 1μL of PCR

products and run for 90 min at 70V using 0.7% TAE buffer solution and stained

with HydraGreen.

Genome and mobile genetic elements annotation

Automated genome annotation was done using Prokka v1.11 (Prokaryotic

Genome Annotation System) (44) and then manually curated using Artemis (41)

and Geneprimp (45). The start codons of known coding sequences (CDS), such as

in the capsule and streptolysin S operons, were further adjusted where

appropriate using UniProtKB (http://www.uniprot.org/) and Pfam (46). CRISPRs

(Clustered Regularly Interspaced Short Palindromic Repeats) were predicted

using CRISPR finder tool (http://crispr.u-psud.fr/crispr/) (47). Prophage

annotation　　was　　done　　using　　PHAST　　(Phage　　Search　　Tool)

(http://phast.wishartlab.com/)　　　　　(48).　　　　　Island　　　　　Viewer

http://www.pathogenomics.sfu.ca/islandviewer/) was used to predict genomic

islands (GI) (49). Boundaries of phage and GI regions were manually adjusted to their respective attachment sites. IS Saga (http://www-genome.biotoul.fr/) was used for the initial identification of insertion sequences (IS). Additional manual curation was carried out to confirm the boundaries of complete and partial IS elements using the IS Finder database (http://www-is.biotoul.fr/). IS element matches against the database that were at ≥95% nucleotide identity were assigned the top matching IS name. IS elements of less percent identity are novel and were assigned the names IS*Stin2*–IS*Stin10* by ISFinder. The impact of IS elements on flanking coding sequences was analyzed using Artemis and by searching the amino acid sequence in UniProt KB (http://www.uniprot.org/), and Pfam databases (41, 46). The complete annotated genome sequence was deposited at GenBank under the accession number CP022392.

Comparative genomics and phylogenetic analysis

Alignments of the whole-genome or genomic sub-regions, such as the CRISPR and genomic islands, were done using BLASTn implemented in EasyFig v2.1 (40, 50). Detailed analysis of regions of difference and comparison of insertion sequences (IS) were done using Artemis Comparison Tool (ACT) (32). The core genome of the five genomes QMA0248, SF1, YSFST01-82, ISET0901, and ISNO was defined using Roary (51). Phylogenetic trees were constructed using the core genome and the core SNP methods, using multiple programs (see below), using the strain QMA0140 as the out-group (52). The strain *S. iniae* QMA0140 is a dolphin isolate from USA in 1976, and was sequenced using Illumina HiSeq 2000 (See Bioproject PRJNA417543). For quality control, the first 20 bp of each read derived from QMA0140 genomic DNA were hard trimmed using Nesoni v0.132

(https://github.com/Victorian-Bioinformatics-Consortium/nesoni)   with   a

minimum length and quality of 70, and 20, respectively. Hard trimmed filtered

reads of QMA0140 were assembled using SPAdes v3.9.0 where contigs < 10X

coverage and smaller than 100 bp were removed (53). For core genome

phylogenies, whole-genome alignment of the five complete genomes and

QMA0140 draft assembly was done using Mauve v2.4.0 and Parsnp v1.2 with

default parameter settings (54, 55). For the alignment using Mauve, conserved

blocks in all 6 genomes longer than 500 bp were selected and concatenated using

the stripSubsetLCBs script producing the core genome alignment. For core SNP

phylogenies, error-free simulated reads were created using wgsim v0.3.2

(https://github.com/lh3/wgsim) and mapped to the reference genome QMA0248

along with QMA0140 hard trimmed and filtered raw reads using bowtie v1.0.0

(56),   where   variants   were   called   using   Nesoni   v0.132

(https://github.com/Victorian-Bioinformatics-Consortium/nesoni) using default

parameters. Core SNPs were also identified by mapping the reads of the 6

genomes to QMA0248 using BWA-MEM v0.7.15 (r1140), implemented in Snippy

v3.0 (https://github.com/tseemann/snippy) (57). All trees were produced by

RAxML v8.2.9 (58) using the general time-reversible (GTR) and GAMMA

distribution model of among-site rate variation with bootstrapping from 1000

replicates,   and   viewed   using   FigTree   v1.4.0

(http://tree.bio.ed.ac.uk/software/figtree).

1.   Agnew W, Barnes AC. Streptococcus iniae: an aquatic pathogen of global

veterinary significance and a challenging candidate for reliable vaccination.

Veterinary microbiology. 2007;122(1-2):1-15.

2.   Shoemaker CA, Klesius PH, Evans JJ. Prevalence of Streptococcus iniae in

tilapia, hybrid striped bass, and channel catfish on commercial fish farms in the

United States. American journal of veterinary research. 2001;62(2):174-7.

3.   Lau SK, Woo PC, Luk W-k, Fung AM, Hui W-t, Fong AH, et al. Clinical

isolates of Streptococcus iniae from Asia are more mucoid and β-hemolytic than

those from North America. Diagnostic microbiology and infectious disease.

2006;54(3):177-81.

4.   Baiano J, Barnes AC. Towards control of Streptococcus iniae. Emerg Infect

Dis. 2009;15(12):1891.

5.   Frost LS, Leplae R, Summers AO, Toussaint A. Mobile genetic elements:

the agents of open source evolution. Nature reviews Microbiology.

2005;3(9):722.

6.   van Elsas JD, Bailey MJ. The ecology of transfer of mobile genetic

elements. FEMS microbiology ecology. 2002;42(2):187-97.

7.   Ochman H, Lawrence JG, Groisman EA. Lateral gene transfer and the

nature of bacterial innovation. Nature. 2000;405(6784):299-304.

8.   Partridge SR. Analysis of antibiotic resistance regions in Gram-negative

bacteria. FEMS microbiology reviews. 2011;35(5):820-55.

9.   Siguier P, Gourbeyre E, Chandler M. Bacterial insertion sequences: their

genomic impact and diversity. FEMS microbiology reviews. 2014;38(5):865-91.

10.   Cheng S, Hu YH, Jiao XD, Sun L. Identification and immunoprotective

analysis of a Streptococcus iniae subunit vaccine candidate. Vaccine.

2010;28(14):2636-41.

11.   Rajoo S, Jeon W, Park K, Yoo S, Yoon I, Lee H, et al. Complete genome

sequence of Streptococcus iniae YSFST01-82, isolated from olive flounder in Jeju,

South Korea. Genome announcements. 2015;3(2):e00319-15.

12.   Pridgeon JW, Zhang D, Zhang L. Complete genome sequence of a virulent

strain, Streptococcus iniae ISET0901, isolated from diseased tilapia. Genome

announcements. 2014;2(3):e00553-14.

13.   Pridgeon JW, Zhang D, Zhang L. Complete genome sequence of the

attenuated novobiocin-resistant Streptococcus iniae vaccine strain ISNO.

Genome announcements. 2014;2(3):e00510-14.

14.   Glaser P, Rusniok C, Buchrieser C, Chevalier F, Frangeul L, Msadek T, et al.

Genome sequence of Streptococcus agalactiae, a pathogen causing invasive

neonatal disease. Molecular microbiology. 2002;45(6):1499-513.

15.   Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, et al.

Complete genome sequence of a virulent isolate of Streptococcus pneumoniae.

Science. 2001;293(5529):498-506.

16.   Ferretti JJ, McShan WM, Ajdic D, Savic DJ, Savic G, Lyon K, et al. Complete

genome sequence of an M1 strain of Streptococcus pyogenes. Proceedings of the

National Academy of Sciences. 2001;98(8):4658-63.

17.   Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, et

al. CRISPR provides acquired resistance against viruses in prokaryotes. Science.

2007;315(5819):1709-12.

18.   Gong H-Y, Wu S-H, Chen C-Y, Huang C-W, Lu J-K, Chou H-Y. Complete

Genome Sequence of Streptococcus iniae 89353, a Virulent Strain Isolated from

Diseased Tilapia in Taiwan. Genome Announcements. 2017;5(4):e01524-16.

19.   Acinas SG, Marcelino LA, Klepac-Ceraj V, Polz MF. Divergence and

redundancy of 16S rRNA sequences in genomes with multiple rrn operons.

Journal of bacteriology. 2004;186(9):2629-35.

20.   Lim K, Furuta Y, Kobayashi I. Large variations in bacterial ribosomal RNA

genes. Molecular biology and evolution. 2012;29(10):2937-48.

21.   Fischetti V, Pancholi V, Schneewind O. Conservation of a hexapeptide

sequence in the anchor region of surface proteins from gram‐positive cocci.

Molecular microbiology. 1990;4(9):1603-5.

22.   Switalski LM, Patti JM, Butcher W, Gristina AG, Speziale P, Höök M. A

collagen receptor on Staphylococcus aureus strains isolated from patients with

septic arthritis mediates adhesion to cartilage. Molecular microbiology.

1993;7(1):99-107.

23.   van der Ploeg JR. Genome sequence of the temperate bacteriophage PH10

from Streptococcus oralis. Virus genes. 2010;41(3):450-8.

24.   Aziz RK, Breitbart M, Edwards RA. Transposases are the most abundant,

most ubiquitous genes in nature. Nucleic Acids Res. 2010;38(13):4207-17.

25.   Ochman H, Davalos LM. The nature and dynamics of bacterial genomes.

Science. 2006;311(5768):1730-3.

26.   Denapaite D, Brückner R, Nuhn M, Reichmann P, Henrich B, Maurer P, et

al. The genome of Streptococcus mitis B6-what is a commensal? PLoS One.

2010;5(2):e9426.

27.   Eraclio G, Ricci G, Fortina MG. Insertion sequence elements in Lactococcus

garvieae. Gene. 2015;555(2):291-6.

28.   Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA

restriction and modification: enzymes, genes and genomes. Nucleic acids

research. 2009;38(suppl_1):D234-D6.

29.   Lobner-Olesen A, Skovgaard O, Marinus MG. Dam methylation:

coordinating cellular processes. Curr Opin Microbiol. 2005;8(2):154-60.

30.   Moran NA, Plague GR. Genomic changes following host restriction in

bacteria. Current opinion in genetics & development. 2004;14(6):627-33.

31.   Richards VP, Lang P, Bitar PDP, Lefébure T, Schukken YH, Zadoks RN, et al.

Comparative genomics and the role of lateral gene transfer in the evolution of

bovine adapted Streptococcus agalactiae. Infection, Genetics and Evolution.

2011;11(6):1263-75.

32.   Carver TJ, Rutherford KM, Berriman M, Rajandream M-A, Barrell BG,

Parkhill J. ACT: the Artemis comparison tool. Bioinformatics. 2005;21(16):3422-

3.

33.   Nijkamp J, Winterbach W, Van den Broek M, Daran J-M, Reinders M, De

Ridder D. Integrating genome assemblies with MAIA. Bioinformatics.

2010;26(18):i433-i9.

34.   Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, et al.

Improvements to PATRIC, the all-bacterial Bioinformatics Database and Analysis

Resource Center. Nucleic Acids Res. 2017;45(D1):D535-D42.

35.   Forde BM, Ben Zakour NL, Stanton-Cook M, Phan MD, Totsika M, Peters

KM, et al. The complete genome sequence of Escherichia coli EC958: a high

quality reference sequence for the globally disseminated multidrug resistant E.

coli O25b:H4-ST131 clone. PLoS One. 2014;9(8):e104400.

36.   Draper JL, Hansen LM, Bernick DL, Abedrabbo S, Underwood JG, Kong N,

et al. Fallacy of the Unique Genome: Sequence Diversity within Single

Helicobacter pylori Strains. MBio. 2017;8(1).

37.   Millard CM, Baiano JC, Chan C, Yuen B, Aviles F, Landos M, et al. Evolution

of the capsular operon of Streptococcus iniae in response to vaccination. Applied

and environmental microbiology. 2012;78(23):8219-26.

38.   Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al.

Nonhybrid, finished microbial genome assemblies from long-read SMRT

sequencing data. Nature methods. 2013;10(6):563-9.

39.   Sullivan MJ, Zakour NLB, Forde BM, Stanton-Cook M, Beatson SA.

Contiguity: contig adjacency graph construction and visualisation. PeerJ

PrePrints; 2015. Report No.: 2167-9843.

40.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment

search tool. Journal of molecular biology. 1990;215(3):403-10.

41.   Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream M-A, et al.

Artemis: sequence visualization and annotation. Bioinformatics.

2000;16(10):944-5.

42.   Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA

restriction and modification: enzymes, genes and genomes. Nucleic acids

research. 2009:gkp874.

43.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et

al. Primer3—new capabilities and interfaces. Nucleic acids research.

2012;40(15):e115-e.

44.   Seemann T. Prokka: rapid prokaryotic genome annotation.

Bioinformatics. 2014:btu153.

45.   Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A,

et al. GenePRIMP: a gene prediction improvement pipeline for prokaryotic

genomes. Nature methods. 2010;7(6):455-7.

46.   Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths‐Jones S, et al.

The Pfam protein families database. Nucleic acids research. 2004;32(suppl

1):D138-D41.

47.   Grissa I, Vergnaud G, Pourcel C. CRISPRFinder: a web tool to identify

clustered regularly interspaced short palindromic repeats. Nucleic acids

research. 2007;35(suppl 2):W52-W7.

48.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast phage

search tool. Nucleic acids research. 2011:gkr485.

49.   Dhillon BK, Laird MR, Shay JA, Winsor GL, Lo R, Nizam F, et al.

IslandViewer 3: more flexible, interactive genomic island discovery, visualization

and analysis. Nucleic acids research. 2015:gkv401.

50.   Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison

visualizer. Bioinformatics. 2011;27(7):1009-10.

51.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al.

Roary: rapid large-scale prokaryote pan genome analysis. Bioinformatics.

2015;31(22):3691-3.

52.   Baiano JC, Tumbol RA, Umapathy A, Barnes AC. Identification and

molecular characterisation of a fibrinogen binding protein from Streptococcus

iniae. BMC microbiology. 2008;8:67.

53. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al.

SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. Journal of computational biology. 2012;19(5):455-77.

54. Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of

conserved genomic sequence with rearrangements. Genome research.

2004;14(7):1394-403.

55. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid

core-genome alignment and visualization of thousands of intraspecific microbial

genomes. Genome biology. 2014;15(11):524.

56. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. Genome

biology. 2009;10(3):R25.

57. Li H. Aligning sequence reads, clone sequences and assembly contigs with

BWA-MEM. arXiv preprint arXiv:13033997. 2013.

58. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-

analysis of large phylogenies. Bioinformatics. 2014;30(9):1312-3.

59. Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter:

circular and linear interactive genome visualization. Bioinformatics.

2009;25(1):119-20.

Table 1: General features of five *S. iniae* complete genomes.

Strain	QMA0248	SF1	YSFST01-82	ISET0901	ISNO
Accession Number	CP022392	CP005941	CP010783	CP007586	CP007587
Genome Size (bp)	2,116,570	2,149,844	2,086,959	2,070,822	2,070,182
GC Content (%)	36.8	36.7	36.8	36.8	36.8
Total CDS number	1,946	2,125	1,897	1,872	1,865
Total gene number	2,196	2,196	2,029	1,997	1,996
rRNA (5S, 16S, 23S)	15	12	15	12	12
tRNA	58	45	58	45	45
Reference	This study	(12)	(11)	(12)	(13)
Assembly type	PacBio RSII P2C4	454 FLX+/ Illumina MiSeq/ Sanger	454 FLX Titanium/ Opgen/ Sanger	Illumina 1500 HiSeq/ Reference guided assembly	Illumina 1500 HiSeq/ Reference guided assembly

607 **Table 2: Large mobile genetic elements (MGE) and regions of difference (ROD) identified in the 5 *S.***
608 ***iniae* genomes analyzed (QMA0248, SF1, YSFST01-82, ISET0901, and ISNO).**

Region	Genomic Island (GI-leu)	ROD1	Prophage 2	ROD2	Prophage 1 (Phi1)
Coordinates[1]	87374-100014	177819-206359 (YSFST01-82)	848479-890501 (SF1)	1767227-1787619	1991661-2023508
Length (Kb)	12.6	28.5	42.0	20.4	31.8
GC content (%)	35.9	37.2	35.3	34.3	37.6
Features	MGE; 13 tRNA and 1 rRNA operon (5S, 16S, and 23S) upstream; integrase; IS*30* and IS*256* family IS elements	Integrase; IS*3* and IS*256* family IS elements	MGE; integrase	IS*3* family IS element	MGE; integrase; 1 tRNA-Cys
No. of CDSs	11	33	63	18	53
Major CDSs	Cro/CI family transcriptional regulator; ECF subfamily RNA polymerase sigma factor; plasmid replication protein; membrane protein	ESAT-6-like protein; two-component sensor histidine kinase; galactose mutarotase; 3 PTS galactitol transporter subunits (IIA, IIC, and IIB)	Phage DNA replication protein; prophage antirepressor, phage capsid and scaffold protein; putative tail protein; holin; endolysin; antigen C; several phage hypothetical proteins	ESAT-6-like protein; O-glycosyl hydrolase; phage infection protein; 3 lipoproteins; protein kinase	DNA helicase; Cro/CI family transcriptional regulator; tail and capsid proteins; holin; lysin; DNA N-4 cytosine methyltransferase; site specific recombinase; several phage hypothetical proteins
Best hit (% identity, % coverage)	*S. parauberis* KCTC 11537 (90, 41)	*S. dygalactiae subsp. equisimilis ATCC 12394* (70, 34)	*S. parauberis* KCTC 11537 (91, 45)	*S. thermophilus* JIM 8232 (81, 13)	Bacteriophage PH10 of *Streptococcus* (71, 34)

609 [1]Coordinates are in QMA0248 GenGenBank annotation (GCA_002220115.1) unless otherwise indicated.
610

611 **Table 3: Summary of all insertion sequences (IS) identified in QMA0248. Partial IS are suffixed by –p.**

IS Family in QMA0248	Number of IS copies	IS types (copy number, mean % amino acid identity)
IS3	32	IS*Sag2* (10, 98.6), IS*Sag2*-p (2, 98.7)
		IS*981* (8, 99.8)
		IS*Spy1* (1, 86.7), IS*Spy1*-p (1, 90.5)
		*IS*Stin6* (3, 94.5), *IS*Stin7* (4, 90.0), *IS*Stin5* (2, 69.3), unclassified most similar to *IS*Stin5-p* (1, 73.8)
IS30	22	IS*Sag9* (3, 99.6), IS*Sag9*-p (5, 99.6)
		*IS*Stin4* (2, 93.3), *IS*Stin4*-p (2, 89.6), *IS*Stin9* (9, 81.4), *IS*Stin2* (1, 84.7)
IS256	17	IS*Stin1* (16, 91.0), unclassified most similar to IS*Stin1*-p (1, 90.2)
IS1182	13	*IS*Stin8* (7, 86.3), *IS*Stin8*-p (2, 89.8), *IS*Stin3* (1, 87.7), *IS*Stin3*-p (2, 87.1), unclassified most similar to *IS*Stin8-p* (1, 70.5)
IS200/IS605	5	*IS*Stin10* (3, 99), *IS*Stin10*-p (1, 98.6), unclassified most similar to *IS*Stin10* (1, 86.5)
ISL3	1	Unclassified most similar to IS*Sth1*-p (1, 77.8)
IS110	2	Unclassified most similar to IS*L4* (2, 65.8)

612 *Novel IS element.*

Forward CDSs

Reverse CDSs | Pseudogenes

rRNA | Insertion Sequences

tRNA | GI, CRISPR/Cas, Phage

613
614
615 **Figure 1: Circular map of the genome *Streptococcus iniae* QMA0248.** Genomic

616 features from outer ring to the inner ring are described in the key to the left, where

617 the innermost two rings correspond to the GC skew (inner), and GC plot (outer).

618 CDS: coding sequence. GI: genomic island. The circular map was generated using

619 DNAPlotter (59).

621

Figure 2: Comparison of the CRISPR/Cas region between QMA0248, SF1,

YSFST01-82, ISET0901, and ISNO. Alignment of Cas genes where the genomes

are ordered according to their position in the phylogenetic tree (left). The

maximum Likelihood (ML) phylogeny was rooted to QMA0140 (not shown) and

built using 1,111 SNPs. Arrows correspond to Cas genes, which are labeled at the

bottom. Figure was produced using EasyFig (50) using 500 bp as minimum length,

90% as minimum identity value, and 0.001 as maximum e-value.

631

Figure 3: Whole-genome alignment of the 5 genomes QMA0248, SF1, YSFST01-82, ISET0901, and ISNO. The genomes are ordered according to their position in the core SNP-based phylogenetic tree. The maximum Likelihood (ML) phylogeny was rooted to QMA0140 (not shown) and built using 1,111 SNPs. The scale bar indicates the number of substitutions represented by branch lengths. BLASTn comparison was produced using EasyFig (50) using 2000 bp as minimum length, 50% as minimum identity value, and 1 x 10^{-17} as maximum e-value.

Supplementary Information

A high quality reference genome for the fish pathogen

Streptococcus iniae

Areej S. Alsheikh-Hussain[1,2], Nouri L. Ben Zakour[1,2,3], Brian M. Forde[1,2],

Oleksandra Silayeva[4], Andrew C. Barnes[4], and Scott A. Beatson*[1,2].

[1]School of Chemistry & Molecular Biosciences, [2]Australian Infectious Diseases Research Centre and [4]School of Biological Science, The University of Queensland, Brisbane, Queensland, Australia;

Present address: [3]The Westmead Institute for Medical Research and the University of Sydney, Sydney, New South Wales, Australia

* Corresponding author.

Mailing address: School of Chemistry and Molecular Biosciences, Building 76, The University of Queensland, Brisbane QLD 4072, Australia.

Phone: +617 33654863; E-mail: s.beatson@uq.edu.au

S. iniae QMA0248 methylome

DNA methylation guides numerous critical processes including defence against foreign DNA, DNA replication and repair, gene expression and virulence. Analysis of PacBio sequence data enabled the detection of genome-wide DNA methylation. Three DNA methyltransferases (MTases) were encoded in the QMA0248 genome. On the basis of homology to known MTases, QMA0248_0514 (annotated as M.Sin248ORF514P in REBASE) likely targets the GCNGC motif and QMA0248_1949 (annotated as M.Sin248ORF1949P in REBASE) likely targets GCCHR (1). QMA0248_0505 (annotated as M.Sin248ORF0505P in REBASE) is encoded ~5kb upstream of QMA0248_0514 but has no close functional homologs and thus has an unknown recognition sequence (1). A small fraction of GCNGC and GCCHR motifs were methylated in QMA0248 (<3%). Collectively these two motifs account for 21,421 sites across the chromosome (or 1 every 100 bp) suggesting a potential role for methylation in the regulation of gene expression in QMA0248. This figure is roughly equivalent to the frequency of Dam GATC sites in *Escherichia coli*. Further work is required to determine if the activity detected here is biologically meaningful.

5-methylcytosine DNA methyltransferase QMA0248_0514

A putative Type II 5-methylcytosine (m5C) DNA MTase (QMA0248_0514, annotated as M.Sin248ORF514P in REBASE) likely targets the GCNGC motif (1). Uncharacterised homologs with 99% amino acid identity are found in 8 other available *S. iniae* complete or draft genomes including SF1, YSFST01-82, ISET0901, ISNO and 89353 (1). In most cases restriction enzymes predicted to recognise GCNGC are predicted to be encoded nearby, or immediately adjacent to

the respective MTase gene. Notably, in QMA0248 the adjacent restriction enzyme (QMA0248_0515) is a pseudogene that has been truncated by an IS*981*. In *S. iniae* KCTC 11634BP, the orthologous gene in its draft quality 454 genome was truncated the same point by a contig break (2), suggesting that in both QMA0248 and KCTC 11634BP the MTase does not function as part of a restriction-modification system. The GCNGC motif is found 8074 times in the QMA0248 genome suggesting that methylation activity could have wide-ranging regulatory consequences.

The closest homologs to QMA0248_0514 for which MTase activity has been determined is the M.CmaLM2II enzyme from *Clostridium mangenotii* LM2 and the M.LmoJ3I enzyme from *Listeria monocytogenes* J3115. Despite sharing modest overall amino acid similarity to QMA0248_0514 (59% and 45%, respectively), regions of high amino acid identity within their predicted target recognition domains (34/34 for M.CmaLM2II and 32/24 for M.LmoJ3I) support the prediction that QMA0248_0514 would also methylate the 2nd cytosine of the GCNGC motif. Detection of m5C using PacBio data is normally unreliable (3). As expected, methylation was detected at only a small fraction of GCNGC sites in the QMA0248 genome and the consensus motif determined by the PacBioSMRT-Portal software includes additional bases that are probably artefactual (e.g. GCNGCAGC) (Supplementary Table S5). Further experimental work (such as using Tet1 pre-treatment to enhance detection of m5C with PacBio sequencing, or Oxford Nanopore sequencing) is needed to determine the true extent of cytosine methylation in the *S. iniae* genome and its role in gene regulation*.*

N4-methylcytosine DNA methyltransferase QMA0248_1949

The second *S. iniae* QMA0248 MTase (QMA0248_1949, known as M.Sin248ORF1949P in REBASE) shares 95% amino acid identity with the Orphan (gamma) N4-methylcytosine (m4C) DNA MTase M.NgoDCXV (gb|AJ004687.2) from *Neisseria gonorrhoeae*, which specifically targets the GCCHR motif. M.NgoDCXV homologs are remarkably rare and confined to a few streptococcal species (including *S. iniaie* SF1). No specific methylation of GCCHR was detected in the QMA0248 genome but 126 motifs that partially overlapped with GCCHR showed evidence of methylation (Supplementary Table S5). The GCCHR motif is present in 13,347 locations in the QMA0248 genome so this represents only a fraction of available sites. The m4C modification is normally detectable from PacBio sequence data, therefore further work is required to determine if QMA0248_1949 is expressed and functional.

Putative Type II N4-methylcytosine or N6-methyladenine DNA methyltransferase QMA0248_0505

The third MTase in *S. iniae* QMA0248 (QMA0248_0505, known as M.Sin248ORF0505P in REBASE) is encoded ~5kb upstream of QMA0248_0514 and shares a similar strain distribution. There are no close homologs of QMA0248_0505 for which a recognition site has been determined. Accordingly it has been annotated by REBASE as a putative Type II N4-cytosine or N6-adenine DNA methyltransferase of unknown recognition sequence (1).

Supplementary Figures

Figure S1: Graphical representation of spacers in CRISPR between YSFST01-82, ISET0901, ISNO, SF1, and QMA0248. Each box represents a spacer, where the same colour and number indicate identical spacers. *: A spacer with an additional direct repeat sequence.

Figure S2: PacBio raw reads pileup of *Streptococcus iniae* QMA0248 on the spurious 7.2 Kb rRNA contig. Alignment of reads is visualised using BamView implemented in Artemis. The lower panel corresponds to the forward and reverse strands of the 7.2 Kb contig, where vertical black lines indicate stop codons. Horizontal lines on the upper panel correspond to the aligned reads, where Artemis colours duplicate reads that span the same region in green black.

Figure S3: Long-range PCR across rRNA operons. Gel electrophoresis of long range PCR products across five rRNA operons of *S. iniae* QMA0248 (lanes 1-5 in operon-wise order as listed in Table 4). Primers are listed in Table 4. "L" denotes 1 Kb Extend DNA Ladder (NEB).

Figure S4: Alternative tree-building methods. Phylogenetic trees constructed for *S. iniae* based on core genome using Mauve (A) and Parsnp (B), and core SNP using Nesoni (C) and Snippy (D). Maximum likelihood (ML) phylograms were built using RAxML where bootstrap support values are shown on branches. Trees are rooted using QMA0140 (not shown). Branch lengths correspond to the number of substitutions (C and D), and the number of substitutions per site (A and B). ML phylograms in A and B were built from 1,928,667 bp and 1,856,820 bp core genome alignment, respectively, whereas C and D were built using 1,241 and 1,116 substitutions, respectively.

1.	Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. Nucleic acids research. 2009;38(suppl_1):D234-D6.

2.	Choi HS, Kwon MG, Kim MS, Park MA, Kim D-W, Park J-Y, et al. Draft genome sequence of beta-hemolytic Streptococcus iniae KCTC 11634. Genome announcements. 2013;1(6):e00897-13.

3.	Clark TA, Lu X, Luong K, Dai Q, Boitano M, Turner SW, et al. Enhanced 5-methylcytosine detection in single-molecule, real-time sequencing via Tet1 oxidation. BMC biology. 2013;11(1):4.