A novel virus alters gene expression and vacuolar morphology in *Malassezia* cells and

induces a TLR3-mediated inflammatory immune response

Minji Park[a], Yong-Joon Cho[b], Donggyu Kim[c], Chul-Su Yang[c,d], Shi Mun Lee[e], Thomas L.

Dawson Jr.[e,f], Satoshi Nakamizo[e], Kenji Kabashima[e,g], Yang Won Lee[h,i], Won Hee Jung [a,#]

[a]Department of Systems Biotechnology, Chung-Ang University, Anseong, Korea

[b]The Research Institute of Basic Sciences, Seoul National University, Seoul, Korea

[c]Department of Molecular and Life Science, Hanyang University, Ansan, Korea

[d]Department of Bionano Technology, Hanyang University, Seoul, Korea

[e]Singapore Immunology Network (SIgN) and Skin Research Institute of Singapore (SRIS),

Agency for Science, Technology and Research (A*STAR), Biopolis, Singapore

[f]Center for Cell Death, Injury & Regeneration, Departments of Drug Discovery &

Biomedical Sciences and Biochemistry & Molecular Biology, Medical University of South

Carolina, Charleston, South Carolina, United States

[g]Department of Dermatology Kyoto University Graduate School of Medicine, Kyoto, Japan

[h]Department of Dermatology, School of Medicine, Konkuk University, Seoul, 05029, Korea

[i]Research Institute of Medicine, Konkuk University, Seoul, Korea

Running Head: A novel virus in *Malassezia*

#Address correspondence to Won Hee Jung, whjung@cau.ac.kr

Word counts

Abstract: 196 words; Text: 4,987 words

Abstract

Most fungal viruses have been identified in plant pathogens, whereas the presence of viral

particles in human pathogenic fungi is less well studied. In the present study, we observed

extrachromosomal double-stranded RNA (dsRNA) segments in various clinical isolates of

Malassezia species. *Malassezia* is the most dominant fungal genus on the human skin surface,

and species in this group are considered etiological factors of various skin diseases including

dandruff, seborrheic dermatitis, and atopic dermatitis. We identified novel dsRNA segments,

and our sequencing results revealed that the virus, named MrV40, belongs to the Totiviridae

family and contains an additional satellite dsRNA segment encoding a novel protein. The

transcriptome of virus-infected *Malassezia restricta* cells was compared to that of virus-cured

cells, and the results showed that transcripts involved in ribosomal biosynthesis were

downregulated and those involved in energy production and programmed cell death were

upregulated. Moreover, transmission electron microscopy revealed significantly larger

vacuoles in virus-infected *M. restricta* cells, indicating that MrV40 infection dramatically

altered *M. restricta* physiology. Our analysis also revealed that viral nucleic acid from MrV40

induced a TLR3-mediated inflammatory immune response in bone marrow-derived dendritic

cells, suggesting that a viral element contributes to the pathogenicity of *Malassezia*.

Importance

Malassezia is the most dominant fungal genus on the human skin surface and is associated

with various skin diseases including dandruff and seborrheic dermatitis. Among *Malassezia*

species, *Malassezia restricta* is the most widely observed species on the human skin. In the

current study, we identified a novel dsRNA virus, named MrV40, in *M. restricta* and

characterized the sequence and structure of the viral genome along with an independent

satellite dsRNA viral segment. Moreover, expression of genes involved in ribosomal

synthesis and programmed cell death was altered, indicating that virus infection affected the

physiology of the fungal host cells. Our data also showed that the viral nucleic acid from

MrV40 induces a TLR3-mediated inflammatory immune response in bone marrow-derived

dendritic cells, indicating that a viral element likely contributes to the pathogenicity of

Malassezia. This is the first study to identify and characterize a novel mycovirus in

Malassezia.

Introduction

Viruses have been observed in many fungal species since their first identification in mushrooms (1). Fungal viruses, also known as mycoviruses, possess different forms of viral genomes including double-stranded RNA (dsRNA), single-stranded RNA (ssRNA), and single-stranded DNA (ssDNA). It is estimated that 30–80% of all fungal species, mainly endophytic fungi, are infected with viruses. Unlike viruses that infect other organisms, the transmission of fungal virus occurs vertically by cell division or horizontally via mating or hyphal anastomosis, with no extracellular phase of the virus life cycle. dsRNA segments have predominantly been found in fungal viruses, and, taxonomically, the fungal dsRNA viruses are classified into seven families: Chrysoviridae, Endornaviridae, Megabirnaviridae, Quadriviridae, Partitiviridae, Reoviridae, and Totiviridae (2).

The model fungus *Saccharomyces cerevisiae* also carries a dsRNA virus that belongs to the Totiviridae family and is known as the L-A virus. A unique feature of fungal viruses of the Totiviridae family is their capability to produce a killer toxin that lyses susceptible neighbor strains, whereas the virus-containing strain (also known as the killer strain) is immune to the toxin. Studies of how the virus produces killer toxins in *S. cerevisiae* showed that the killer toxins are encoded by a satellite dsRNA segment, known as the M satellite, within the L-A virus. To date, four different killer toxins, K1, K2, K28, and Klus, have been described (3-6). The *S. cerevisiae* L-A virus forms icosahedral particles with a diameter of approximately 39 nm (7). The virus possesses a non-segmented 4.6-kb dsRNA genome consisting of two open reading frames (ORFs), *gag* and *pol*, which overlap by 130 base pairs (bp) (8). A major 76-kDa capsid protein is encoded by *gag*, and a 180-kDa minor protein species is encoded as a Gag-Pol fusion protein by a −1 ribosomal frame-shift event (9, 10). The C-terminus of the Gag-Pol fusion protein possesses viral RNA-dependent RNA polymerase (RDRP) activity (8).

The ribosomal frameshift is an interesting feature in a compact viral genome and has been commonly found in various viral genomes as a mechanism to allow viruses to express overlapping ORFs. Studies of *S. cerevisiae* L-A virus revealed that the mechanism of frameshifting is based on the sequence structures including a canonical slippery heptamer, 5'-X XXY YYZ-3' (X= A, U or G; Y=A or U; Z=A, U, or C) and RNA pseudoknot (11).

Fungal viruses have also been considered biocontrol agents in the field of agriculture. For example, a virus causes hypovirulence in the chestnut blight fungus *Cryphonectria parasitica* (12, 13), and a virus mediates the biocontrol of other phytopathogenic fungi such as *Helminthosporium victoriae*, *Sclerotinia sclerotiorum*, and *Botrytis cinerea* (14).

Although viral infections in fungal cells are widespread, the interactions between the fungal virus and its host are not well understood. One of the most studied host defense mechanisms is RNA silencing. Several studies have shown that RNA silencing functions as an antiviral defense mechanism against *C. parasitica* in fungi. Disruption of the dicer pathway in *C. parasitica* increases the susceptibility to virus infections (15), and p29 was identified as a suppressor that inhibits expression of the genes required for RNA silencing-mediated viral defense in the fungus (16). Similarly, conserved RNA silencing-mediated antiviral defense systems have been identified in *Aspergillus nidulans*, *Rosellinia necatrix*, and *Fusarium graminearum* (17-19).

Malassezia is the most dominant fungal genus on the human skin surface and is considered an etiological factor in various skin diseases including dandruff, seborrheic dermatitis, and atopic dermatitis (20-23). Eighteen *Malassezia* species have been identified; among them, *Malassezia restricta* is the most abundant on the human skin (20). Recent studies showed an increased burden of *M. restricta* on the scalp of patients with dandruff, indicating an association between dandruff and the fungus, although its role as a pathogenic organism is

still unclear, and host susceptibility should be taken into consideration (21, 24-27). Most

fungal viruses are found in plant pathogenic fungi, whereas few examples of viral particles

have been identified in human pathogenic fungi such as *Candida albicans* (28). In the present

study, we observed extrachromosomal dsRNA segments in various *M. restricta* clinical

isolates which represented a novel viral genome and its satellite. Sequence analysis revealed

that the virus belongs to the Totiviridae family and that the additional satellite dsRNA

segment encodes a novel protein. The interactions between the viral elements and the fungal

host, and the impact of the virus on fungal interactions with immune cells were evaluated.

Results

Identification of extrachromosomal dsRNA segments in *Malassezia*

Extrachromosomal nucleic acid bands were observed in total nucleic acid extracts of the *M.*

restricta strains isolated in our recent study (29). Among the strains, *M. restricta* KCTC

27540 was used to identify extrachromosomal segments. Total nucleic acids were extracted

from the strain and digested with DNase I, RNase A, and RNase T1. The extrachromosomal

segments and ribosomal RNA were resistant to DNase I, indicating that they were neither

ssDNA nor dsDNA. RNase A degraded all nucleic acids except for genomic DNA, whereas

RNase T1, which catalyzes the degradation of ssRNA, removed ribosomal RNA only (30).

These results suggested that the extrachromosomal segments correspond to dsRNA and that

M. restricta KCTC 27540 possesses two separate extrachromosomal segments as estimated

by agarose gel electrophoresis to be approximately 4.5 and 1.5 kb (Fig. 1A).

To confirm whether the extrachromosomal segments observed in other *M. restricta* clinical

isolates were also dsRNA, total nucleic acid extracts from strains other than *M. restricta*

KCTC 27540 were treated with DNase I and RNase T1. The extrachromosomal segments

remained unaffected after enzyme treatment, indicating that they are also dsRNA, as in *M.*

restricta KCTC 27540 (Fig. 1B). Moreover, other *Malassezia* species including *M. globosa*,

M. pachydermatis, and *M. sympodialis* showed similar extrachromosomal dsRNA segments

suggesting that these segments are common characteristics of *Malassezia* species (Fig. 1C).

Agarose gel electrophoresis revealed extrachromosomal segments composed of at least two

separate dsRNA fragments except for *M. restricta* KCTC 27543, which showed a single

dsRNA fragment. Additionally, the large fragments of dsRNA showed similar sizes (~5.0 kb),

whereas the small dsRNA segments varied in size in different strains (Fig. 1B and 1C). We

hypothesized that the dsRNA segments from *Malassezia* strains represent the dsRNA

elements of mycoviruses, which are prevalent in all major fungal taxa (2). Indeed, a dsRNA

mycovirus infected soil fungus, *Trichoderma atroviride* NFCF028, displayed similar

extrachromosomal segments (Fig. S1) (31).

Sucrose gradient ultracentrifugation was conducted to purify virus particles to confirm that

the dsRNA segments in the *Malassezia* strains were indeed viral elements. The separated

nucleic acids and proteins in each fraction were analyzed. Two dsRNA fragments (~5.0 and

~1.7 kb) were clearly visible in fractions 1–6 following agarose gel electrophoresis (Fig. 2A).

Moreover, the results of sodium dodecyl sulfate-poly acrylamide gel electrophoresis (SDS-

PAGE) showed that fractions 3–6 contained protein bands with an estimated molecular

weight of ~77 kDa (Fig. 2B). This molecular weight is similar to that of the known capsid

protein of the *S. cerevisiae* mycovirus (9, 32, 33). Fractions 3–6 were subsequently evaluated

by microscopy to visualize mycovirus particles in *M. restricta*. Transmission electron

microscopy (TEM) images showed virus-like particles with an isometric shape and a

diameter of 43 nm (Fig. 2C). These results support the hypothesis that the extrachromosomal

dsRNA segments form the genome of mycovirus in *M. restricta* KCTC27540. We named the

viral particle as MrV40 (_M. restricta_ KCTC 275_40_ Myco_v_irus). The large and small dsRNA

viral fragments were named as MrV40L and MrV40S, respectively. In addition to evaluating

images of the purified virus particles, we examined the morphology of the virus-infected and

the virus-cured _M. restricta_ KCTC 27540 strain pairs by TEM. The virus-cured _M. restricta_

KCTC 27540 strain was prepared by sterilizing the virus-infected _M. restricta_ KCTC 27540

through numerous serial passages (Supplemental Materials and Methods, and Fig. S2). In

general, there was no morphological difference between the virus-infected and the virus-

cured KCTC 27540 strains. However, we noticed that the size of vacuoles in the virus-

infected strain was significantly larger than that in the virus-cured strain, suggesting that the

virus influences vacuole size in _M. restricta_ (Fig. 3).

Determination of dsRNA sequence of MrV40L

The complete sequence of MrV40L was determined by a combination of the Illumina MiSeq

technique and the Sanger sequencing method using purified viral dsRNA. The length of the

complete assembled sequence of MrV40L was 4,606 bp, and two overlapping open reading

frames (ORFs), designated as ORF1 and ORF2, were identified (Fig. 4A). ORF1 corresponds

to the region from nucleotides (nt) 28 to 2,097 and encodes a polypeptide of 689 amino acids

with a molecular weight of 77 kDa. ORF2 corresponds to the region from nt 1,949 to 4,538

and encodes a polypeptide of 862 amino acids with a molecular weight of 98 kDa. The results

of BLAST analysis showed that the protein sequences of ORF1 and ORF2 were highly

similar to that of the capsid protein (the Pfam families of LA-virus_coat, PF09220) and viral

RNA-directed RNA polymerase (RDRP; the Pfam family of RDRP_4, PF02123) of _S._

segobiensis virus L belonging to the genus _Totivirus_ (Totiviridae family) with 53%

(YP_009507830.1) and 52% (YP_009507831.1) identities, respectively (34). Eight conserved

motifs, which are commonly found in totiviruses, were found within ORF2, supporting the

classification of MrV40 as a totivirus (Fig. 4B) (35). Moreover, phylogenetic analysis with

known amino acid sequences of RDRPs from dsRNA mycoviruses demonstrated that the

RDRP encoded by the MrV40L genome was clustered with totiviruses (Fig. 4C). Thus,

MrV40L is a dsRNA viral genome encoding a capsid protein and a RDRP, and we propose

that MrV40 belongs to the *Totivirus* genus.

Next, we analyzed the genomic sequences of dsRNA viruses in other clinical *M. restricta*

isolates to confirm that they were similar to those of MrV40L, and thus belong to the

Totivirus genus. RT-PCR was performed using four primer sets corresponding to the sequence

of the conserved regions of the capsid protein and RDRP in the MrV40L genome (Fig. 5A).

The results showed that six of 11 viruses generated RT-PCR products, suggesting the

presence of viral genomes similar to MrV40Ls, while the remaining samples had dissimilar

viral genomes (Fig. 5B).

Furthermore, we performed phylogenetic classification of the same viruses found in other

clinical *M. restricta* isolates by multilocus sequence typing of the 1,075-bp region, 638 bp of

ORF1, and 437 bp of ORF2, corresponding to *gag* and *pol*, respectively, within the viral

genomes. The results revealed that the viruses were classified into three clades: clade □

(MrV12L, MrV16L, MrV40L, MrV79L, MrV80L, and MrV82L), clade □ (MrV18L,

MrV43L, MrV83L, and MrV50L), and clade □ (MrV24L) (Fig. 5C). Additionally, the

sequences of the 1,075-bp region of MrV40L, MrV79L, and MrV82L in the clade I were 100%

identical, suggesting that they originated from the same lineage.

Determination of dsRNA sequence of MrV40S

To determine the sequence of MrV40S, the dsRNA segments of MrV40S were extracted from

agarose gels and then subjected to cDNA cloning and sequencing (See Materials and

Methods). Using the partial sequences obtained from cDNA clones and the sequences

obtained from repeated 5′ rapid amplification of cDNA ends (RACE), we successfully

determined the complete sequence of MrV40S. The sequence of MrV40S was 1,355 bp, and

a single ORF was identified in the 3′ region (from nt 773 to 1,097), which encoded a

polypeptide of 124 amino acids with a molecular weight of 15.6 kDa. The ORF was

designated as ORF3. Although we obtained the complete sequence of MrV40S ORF3, no

homologous protein sequence was identified by BLAST analysis using all currently available

databases.

Mycoviruses belonging to *Totivirus*, particularly the *S. cerevisiae* LA-virus, often possess a

satellite dsRNA segment known as M dsRNA, which is responsible for producing a killer

toxin that excludes neighboring yeast cells. Because MrV40L resembles M dsRNA, we

predicted that MrV40S produces a protein that inhibits other *Malassezia* strains and/or other

fungal and/or bacterial cells residing on the skin. To test the toxin-like activity of the protein

produced from MrV40S, ORF3 was cloned and the protein was heterologously expressed in

Escherichia coli and purified (Fig. 6A). The activity of the purified protein was evaluated

against several pathogenic fungi and bacteria including *M. restricta*, *C. albicans*,

Cryptococcus neoformans, E. coli, and *Staphylococcus aureus*. Unexpectedly, the purified

protein displayed no growth inhibitory effect on the microbial cells tested (data not shown).

Based on these results, we concluded that the novel protein produced from ORF3 likely has

no toxin-like activity against the microorganisms tested, and further functional studies are

required to characterize its function.

M dsRNA in *S. cerevisiae* produce several variants of killer toxins, namely, M1, M2, M3, and

Mlus, and they showed limited homology in their protein sequences (4-6). Based on this

information, we compared the sequences of satellite dsRNA genomes between MrV40S and

those in other *M. restricta* strains containing the small dsRNA segments. Reverse

transcription (RT)-PCR was conducted using a series of primers specific to MrV40S. The

results suggested that among the nine strains tested, only the satellite dsRNA in *M. restricta*

KCTC 27582 possessed a similar dsRNA segment. These data indicate that the satellite

dsRNA genome sequences are highly variable (Fig. 6B).

MrV40 altered transcriptome profiles in the host *M. restricta* strain

To understand the influence of MrV40 on the host physiology, we compared the

transcriptomes of virus-infected and virus-cured *M. restricta* KCTC27540 strain pairs by

RNA sequencing. Transcriptome analysis showed that 258 and 809 genes were up- and down-

regulated by more than 2-fold, respectively, in the virus-infected strain compared to those in

the virus-cured strain (Table 1 and Table S1). These results indicate that the presence of the

virus may impact a number of physiological processes in *M. restricta*.

In *S. cerevisiae*, numerous genes are known to be involved in maintaining its dsRNA virus

(36) and are categorized into two groups: *MAK* (maintenance of killer) and *SKI* (superkiller).

At least 25 *MAK* genes have been reported (37); among them, *MAK3*, *MAK10*, and *PET18*

were shown to be required for the maintenance of both *S. cerevisiae* L-A virus and its M

satellite, whereas all other *MAK* genes are responsible only for the M satellite (38-43). We

could not find homologs of *MAK10* or *PET18* in the genome of *M. restricta*, suggesting that a

different or modified virus maintenance mechanism may be present in the fungus compared

to *S. cerevisiae* (Table 2). The results also showed that among the *MAK* homologs, *MAK1*,

MAK5, *MAK11*, *MAK16*, *MAK21*, *MAK7*, and *MAK8* were downregulated by more than 2-

fold. In *S. cerevisiae*, these genes, except for *MAK1*, were involved in 60S ribosomal subunit

biosynthesis and M satellite maintenance. Thus, we predicted that MrV40 reduced ribosomal

biogenesis within the host. Additionally, *M. restricta* contained a series of *SKI* homologs.

However, no *SKI* homologs were differentially expressed in the virus-infected *M. restricta*

KCTC27540 compared to those in the virus-cured strain, indicating that the function of *SKI*

genes was not critical for maintaining the virus. In addition to the genes involved in virus

maintenance, we observed upregulation of numerous genes required for the TCA cycle and

the electron transport chain, including MRET_1104 (NADH dehydrogenase (ubiquinone) Fe-

S protein 7), MRET_1378 (succinate dehydrogenase (ubiquinone) iron-sulfur subunit),

MRET_1953 (NADH dehydrogenase (ubiquinone) Fe-S protein 1), MRET_2042 (fumarate

hydratase, class II), MRET_2097 (succinate dehydrogenase (ubiquinone), MRET_2956 (2-

oxoglutarate dehydrogenase E1 component), MRET_3173 (dihydrolipoamide dehydrogenase)

and MRET_4117 (aconitate hydratase) (Table 3). These results suggest that virus

maintenance and propagation may require higher energy production in the host cell.

The overall dysregulation of primary metabolism may disturb the normal cell physiology in

the *M. restricta* strain infected with MrV40. Indeed, we observed up-regulation of genes

involved in programmed cell death in the fungal host cells. For example, the expression of

MRET_3200 (p38 MAP kinase), MRET_1134 (programmed cell death 6-interacting protein),

and MRET_2499 (autophagy-related protein, *ATG101*), which are associated with

programmed cell death, was upregulated by 4.43-, 3.14-, and 2.83-fold in the virus-infected

fungal cells, respectively. Moreover, MRET_0131 (FAS-associated factor), which is involved

in Fas-induced apoptosis, was found to be strongly upregulated (8.80-fold) in the presence of

the virus within the fungal host (44). It is well-known that programmed cell death is triggered

during virus infection (45, 46), and our findings agree with this observation. To confirm that

there was an increase in programmed cell death, we stained cells with Annexin V and

propidium iodide, which is commonly used to quantitate the apoptosis (47, 48). Using flow

cytometric analysis, we found higher fluorescence intensity in the virus-infected KCTC

27540 strain than in the virus-cured strain, confirming that programmed cell death is

increased in the virus-infected *M. restricta* strain (Fig. S5). We also tested whether increased

programmed cell death influence other physiological characteristics of the virus-infected cells

such as susceptibility to an antifungal drug. However, antifungal susceptibility of the virus-

infected KCTC 27540 was similar to that of the virus-cured strain (data not shown),

suggesting that the viral infection may have a limited effect on the physiological

characteristics of the host cells.

MrV40 induced the TLR3-mediated immune system

Since it was first reported that fungal viral dsRNA induces cytokine production in rabbits (49-

51), several studies have demonstrated that Toll-like receptor 3 (TLR3) plays a central role in

viral dsRNA recognition and production of inflammatory cytokines in innate immune cells

(52, 53). Additionally, a recent study showed that *S. cerevisiae* viral dsRNA stimulates the

immune system through TLR3 in a human embryonic kidney cell line (54). We therefore

investigated whether MrV40 itself or the virus-containing *Malassezia* cells alter the

expression patterns of TLRs and cytokine production in mammalian cells. To this end, bone

marrow-derived dendritic cells (BMDCs) obtained from C57BL/6 mice, which have been

used as a model to study the interactions between fungal cells including *Malassezia* and the

innate immune system in a mammalian host, were used in our study (55, 56).

Two independent strain pairs, virus-infected and virus-cured *M. restricta* KCTC 27540

strains as well as virus-infected and virus-cured *M. restricta* KCTC 27524 strains were used

for the study. Live fungal cells, fungal cell lysates, purified capsid protein, purified satellite

protein (produced from MrV40S ORF3), and purified MrV40 dsRNA were prepared and co-

incubated with BMDCs, and the expression of TLRs and cytokines were analyzed. We found

that TLR3 expression was significantly upregulated by live fungal cells and fungal cell

lysates of the virus-infected *M. restricta* strains, while live cells and their lysates of the virus-

cured strains did not enhance the expression of TLR3. Considering that the expression of

other TLRs were unchanged upon addition of live cells and cell lysates, we concluded that

host response upon treatment with virus-infected *M. restricta* cells was mainly mediated by

TLR3 in murine BMDCs. Moreover, upregulation of TLR3 expression was observed in

BMDCs treated with purified MrV40 dsRNA but not with purified capsid and satellite

proteins, suggesting that MrV40 dsRNA is the causative agent that enhances the expression

of TLR3 (Fig. 7A). Moreover, our data suggested that purified capsid and satellite proteins

from MrV40 did not influence the expression of TLRs.

The expression of several inflammatory cytokines including tumor necrosis factor (TNF)-α,

interleukin (IL)-6, IL-10, interferon (IFN)-α and IFN-γ, in response to live fungal cells,

fungal cell lysates, purified capsid protein, purified satellite protein, and purified MrV40

dsRNA, were evaluated. Particularly, IFN-α and IFN-γ were included because of their

involvement in the antiviral response in mammals (57). The results of the cytokine analysis

indicated that virus-infected live *M. restricta* cells and their lysates induced the production of

TNF-α, IL-6, IL-10, IFN-α and IFN-γ, while no change in cytokine expression was observed

in BMDCs treated with virus-cured live *M. restricta* cells and their cell lysates. Similarly,

capsid protein and MrV40 dsRNA also enhanced the expression of all cytokines tested (Fig.

7B). Cytokine profiles were also measured in homozygous TLR3 knockout mice, as our

observations suggested a TLR3-mediated host response in BMDCs treated with virus-

infected fungal cells. As shown in Fig. 7B, cytokine levels in BMDCs isolated from TLR3

knockout mice were not significantly altered except in those treated with the capsid and

satellite proteins. These data suggest that MrV40 increased the production of the cytokines

TNF-α, IL-6, IL-10, IFN-α, and IFN-γ in a TLR3-dependent manner in BMDCs. However, as

suggested above, the cytokine production induced by the capsid protein seemed to be TLR3-

independent. Moreover, the purified satellite protein also caused a TLR3-independent

increase in cytokine production except of TNF-α.   These results suggest that the capsid and

the satellite proteins also contributed to upregulation of cytokine expression in a TLR3-

independent manner. Overall, our data suggest that dsRNA in MrV40 triggers an increase in

the production of cytokines involved in inflammation and that TLR3 plays a central role in

the host response.

Discussion

In the current study, we detected dsRNA virus in several clinical isolates of *Malassezia*

species. Among them, MrV40 identified in *M. restricta* KCTC 27540 was selected, and its

genome structure and effects on host gene expression and on the mammalian immune

response were evaluated. Our data showed that MrV40 consists of two RNA segments, which

we named MrV40L and MrV40S. The results of the genome sequence analysis suggested that

these segments were 4,606 and 1,355 bp in length, respectively, and belong to the genus

Totivirus. Typically, the genomes of the viruses belonging to the genus *Totivirus* consist of

non-segmented dsRNA with sizes between 4.6 and 7.0 kb, and contain two ORFs, *gag* and

pol. Studies have specifically examined the genome structure of *Totivirus* because of the

overlapping nature of the two ORFs, where a -1 frameshift occurs, resulting translation of the

fusion protein (2). The overlapping ORFs and frameshift were frequently observed in a

compact viral genome to translate proteins and were found in several dsRNA and ssRNA

viruses; these ORFs allow ribosomes to translate CP and RDRP continuously with a missing

CP termination codon (11, 58, 59). The mechanism of frameshifting in the viral genome is

based on the pseudoknot structure of the mRNA for efficient slipping via a slippery site (60).

For example, in the genome of the *S. cerevisiae* L-A virus, the 5′ ORF (*gag*) encodes a 76-

kDa CP and the 3′ ORF (*pol*) encodes an RDRP, which is expressed as a 180-kDa CP-RDRP

fusion protein generated by a -1 ribosomal frameshift (8, 11). In our study, MrV40L (the

major dsRNA segment in MrV40) contained two overlapping ORFs, ORF1 and ORF2. We

identified a putative slippery site heptamer, 5′-GGGTTTT-3′, at the region from nt 1, 968 to

1,974 for the -1 ribosomal frameshift, which may be associated with production of the fusion

protein of 170 kDa in MrV40L. A previous study suggested that in the *S. cerevisiae* L-A virus,

the rate of ribosomal frameshifts was approximately 1.8%, giving 120 CP and 2 CP-RDRP

fusion protein molecules per virus particle (11). Considering the low efficiency of producing

the fusion proteins by ribosomal frameshifting, we expected to observe a significantly lower

translation rate of the fusion protein; indeed, the putative 170-kDa band was not detected by

SDS-PAGE. In addition to MrV40L, we determined the sequence of the satellite dsRNA

segment MrV40S and found that it consists of 1,355 nt containing a single ORF, ORF3,

producing a novel 15.6-kDa protein. As observed for other totiviruses, the possible toxin-like

activity of the protein was investigated in our study, but no growth inhibitory activity against

several bacteria and fungi was observed.

It has been estimated that 30–80% of fungal species in nature are infected with viruses, and a

fungal host normally shows no specific symptoms upon infection (14). However, several

genes were shown to be required for maintaining and propagating viruses in the host fungal

cells. In *S. cerevisiae*, numerous chromosomal genes are known to be involved in viral

propagation and expression of the viral killer toxin (36). Furthermore, several genes are

known to be responsible for maintaining the L-A virus and M dsRNA in *S. cerevisiae*. The

MAK genes are required for the propagation and maintenance of the L-A virus and M dsRNA

in *S. cerevisiae* (61). Among the *MAK* genes, *MAK3*, *MAK10*, and *PET18* are required for the

maintenance of both L-A virus and M dsRNA, whereas all other *MAK* genes are responsible

only for M dsRNA (38-43). Particularly, *MAK3* encodes an N-acetyltransferase and is

required for N-acetylation of the coat protein (38, 62). A previous study showed that the coat

proteins without acetylation failed to self-assemble, resulting in the loss of all dsRNA viruses

(39). *MAK10* and *PET18* (*MAK30*+*MAK31*) encode a non-catalytic subunit of N-terminal

acetyltransferase and a protein of unknown function, respectively. Mutant strains lacking

each gene contained unstable viral particles, indicating that the genes are involved in the

structural stability of LA-virus and M dsRNA (40). *MAK1* (*MAK17*, *TOP1*) encodes DNA

topoisomerase I, and other *MAK* genes including *MAK2*, *MAK5*, *MAK11*, *MAK16*, *MAK21*,

MAK7 (*RPL4A*), *MAK8* (*TCM1*), and *MAK18* (*RPL41B*) are related to 60S ribosomal subunit

assembly (43, 63, 64). All mutant strains lacking the above genes showed decreased levels of

free 60S ribosomal subunits and the inability to maintain M dsRNA, suggesting that stable

propagation of the satellite dsRNA depends on 60S ribosome synthesis (42, 43). In addition

to the *MAK* genes, the *SKI* gene family has been shown to be involved in the maintenance

and propagation of virus in *S. cerevisiae*. *SKI1* (*XRN1*) encoding a 5'-3' exonuclease is

involved in the degradation of uncapped mRNA including viral mRNA, and *SKI2*, *SKI3*,

SKI6, *SKI7*, and *SKI8* block the translation of viral mRNAs (65-67).

In the current study, homologs of most *MAK* and *SKI* genes were identified in *M. restricta*.

The results of the transcriptome analysis suggested that most *MAK* genes were downregulated,

which may in turn reduce ribosome synthesis in the *M. restricta* strain containing MrV40. In

contrast, no *SKI* homolog showed significantly altered transcript levels in the *M. restricta*

strain harboring MrV40. Moreover, we found enhanced expression of genes involved in

energy metabolism and programmed cell death in the *M. restricta* strain containing the virus.

S. cerevisiae also displayed relatively small changes in fungal host gene expression upon

virus infection, possibly because of co-adaptation of the virus within the fungal host (68).

Maintenance and propagation of virus within the fungal host may be involved in the post-

transcriptional mechanism and may contribute to the minimal changes in host gene

expression. Notably, the possibility that an RNA silencing pathway in *M. restricta* cells

influences virus maintenance was excluded because of the absence of homologous genes

required for the pathway in the genome of the fungus.

In addition to transcriptome analysis, we directly investigated whether the virus influences

the cellular morphology of *M. restricta* and the structures of its intracellular organelles by

TEM. We observed significantly larger vacuoles in virus-infected *M. restricta* cells. An

increased vacuole size upon virus infection has been reported previously. The

phytopathogenic fungus *Botrytis porri* infected with dsRNA virus 1 showed the formation of

abundant vacuoles (69), and turnip mosaic virus induced the formation of a large central

vacuole in *Nicotiana benthamiana* plant cells (70). Particularly, in *N. benthamiana*, turnip

mosaic virus particles were shown to accumulate in vacuoles and be protected by the

organelle membranes against the harsh host environment, particularly during xylem vessel

differentiation. Furthermore, virus accumulation in vacuoles in plant-to-plant virus

transmission has been suggested (70). Although the expression of genes involved in vacuole

biogenesis was not altered in transcriptome analysis in virus-infected *M. restricta* cells, there

may be a relationship between vacuole functions and MrV40 in *M. restricta*. Therefore,

additional physiological studies are needed.

TLR3 is well-conserved in most vertebrates, localizes on the endosomal transmembrane, and

plays a role in immune and non-immune cells. It has been suggested that TLR3 senses viral

dsRNA taken up through endocytosis and contributes to defending the host against viral

infection by regulating the expression of a range of cytokines (52, 71). A previous study

demonstrated a significant decrease in cytokines (IFN-α/β, IFN-γ, and IL-12p40) and increase

in viral PFU in the spleen of $Tlr3^{-/-}$ mice infected with mouse cytomegalovirus (72). We

found that TLR3 expression was significantly increased by the live cells of two independent

virus-infected *M. restricta* strains and their cell lysates, while no change was found by the

live cells and cell lysates of the virus-cured strains. The causative agent was very likely to be

dsRNA of MrV40 in the virus-infected *M. restricta* strains since purified dsRNA from the

MrV40 particles enhanced TLR3 expression. The virus-infected live *M. restricta* cells, their

cell lysates, and purified dsRNA from the MrV40 particles also showed increased production

of cytokines involved in inflammation. However, in $Tlr3^{-/-}$ BMDCs, the production of all

cytokines in the cells treated with the virus-infected live *M. restricta* cells, their cell lysates,

and purified dsRNA from the MrV40 particles was attenuated, suggesting that TLR3 plays a

central role in the host response against dsRNA of MrV40 and mediates the production of the

cytokines.

Similarly, the *Leishmania* parasite *L. guyanensis* infected with the dsRNA virus belonging to

Totivirus induced a TLR3-mediated inflammatory immune response within the vertebrate

host, indicating that the *Leishmania* RNA virus (LRV1) functions as an innate immunogen.

Moreover, in *Leishmania*, LRV1 was suggested to stimulate the inflammatory immune

response and increase the severity and persistence of the parasite (73). However, whether

MrV40 contributes to hyper- or hypo-pathogenicity of *M. restricta* remains unclear.

Therefore, further studies to characterize the potential function of MrV40 as an innate

immunogen are needed. In addition, since our data suggest contribution of the MrV40 capsid

and the satellite proteins to host cytokine expression in a TLR3-independent manner,

additional studies are needed to define a role of the capsid and the satellite proteins in

interactions between these viral components and the mammalian host cells.

Overall, our study demonstrated the existence of a dsRNA virus within *M. restricta*. We also

determined the sequences and structure of the viral genome along with the independent RNA

segment. Moreover, we identified viruses not only from different strains of *M. restricta* but

also from other *Malassezia* species, although variations were observed in the viral genomes.

Evidence that the viral nucleic acid from MrV40 induces a TLR3-mediated inflammatory

immune response was obtained, suggesting that a viral element contributes to the

pathogenicity of *Malassezia*.

Materials and Methods

Strains and growth conditions

M. restricta KCTC 27512, KCTC 27516, KCTC 27518, KCTC 27524, KCTC 27540, KCTC

27543, KCTC 27550, KCTC 27879, KCTC 27880, KCTC 27882, and KCTC 27883, *M.*

globosa CBS 7966, *M. pachydermatis* KCTC 27587, and *M. sympodialis* KCTC 27817 were

obtained as previously described and cultured on Leeming and Notman agar (LNA) medium

at 34°C for 3 days (29, 75-77). Among these strains, *M. restricta* KCTC 27540 was used to

identify dsRNA viruses. *E. coli* DH10B and BL21 were grown in Luria-Bertani broth at 37°C.

Purification of virus particles

The virus particles were purified as previously described with some modifications (69).

Briefly, approximately 10 g of *M. restricta* cells was harvested and suspended in 10 mL of

extraction buffer (0.1 M sodium phosphate buffer containing 3% (v/v) Triton X-100, pH 7.0).

The suspended cells were vortexed with glass beads and centrifuged at 10,000 ×*g* at 4°C for

20 min to remove cell debris. The supernatant containing crude extracts was ultracentrifuged

at 119,000 ×*g* at 4°C for 2 h to precipitate all particles. The resulting pellet was suspended in

1 mL of 0.1 M sodium phosphate buffer (pH 7.0), and the suspension was centrifuged at

16,000 ×*g* at 4°C for 30 min to remove large particles. The supernatant was overlaid on

sucrose solution with a gradient concentration ranging from 10% to 40% (w/v) and

centrifuged at 70,000 ×*g* at 4°C for 2 h. Fractions (700 µL/each fraction) were collected from

the top of the ultracentrifuged sucrose gradient solution. Each fraction was analyzed by 1%

agarose gel electrophoresis and 8% SDS-PAGE to detect dsRNA segments and protein,

respectively. Methods for cell fixation and TEM are described in the Supplementary

Materials and Methods.

Determination of MrV40L and MrV40S sequences

The reference genome sequence was required to obtain the complete sequence of MrV40L by

Illumina Miseq. We first obtained the partial fragment of MrV40L and determined its

nucleotide sequence. Briefly, 1 µg of cDNA obtained as described above was amplified using

1 µM of the tagged oligo_1 primer to bind the tag and 1.25 units of *Taq* DNA polymerase

(BIONEER, Daejeon, Korea). The amplicons were cloned into the pGEM-T vector (Promega,

Madison, WI, USA), transformed into *E. coli* DH10B, and sequenced with the universal M13

primers. The resulting partial sequences of MrV40L showed high similarity with the genome

of *Scheffersomyces segobiensis* virus L isolate NRRL Y-11571, which was used to assemble

the reference genome in our study. To sequence the entire genome of the virus from *M.*

restricta KCTC 27540, dsRNA was purified using Cellulose A (Advantec, Taipei, Taiwan)

from total cell extract as described previously (78). After purification, purified dsRNA was

treated with Ambion^TM DNase I (Thermo Fisher Scientific, Waltham, MA, USA) to remove

remaining genomic DNA remained. The sequencing library was constructed using the TruSeq

Stranded Total RNA Sample Prep Kit (Illumina, San Diego, CA, USA) following the

manufacturer's instructions and excluding poly-A selection. The resulting library was

sequenced on an Illumina MiSeq instrument according to the manufacturer's instructions.

The generated raw sequences, 250-bp paired-end reads, were assembled by CLC Genomics

Workbench v7.5 (Qiagen, Hilden, Germany), and the resulting contigs were identified by

BLASTN of the NCBI nucleotide database. Based on the reference genome, two contigs of

MrV40L were assembled. The gap between the contigs was sequenced by gap-filling RT-PCR

using primers Linkage1_V40 and Linkage2_V40) (see Table S2), and both termini were

sequenced by RACE PCR using a 5′/3′ RACE Kit, 2^nd Generation (Roche, Basel, Switzerland)

according to the 5′ RACE protocol described by the manufacturer.

To determine the MrV40S genome, the RACE PCR method was used because we failed to

find any contig of MrV40S. Like MrV40L, the partial sequence of cDNA from MrV40S was

amplified using the tagged oligo_1 primer. The amplicons were cloned into the pGEM-T

vector (Promega) and sequenced. Using the partial sequence obtained, 5′ RACE PCR was

performed and repeated until no additional sequence was found.

***In silico* analysis**

To identify the ORFs in MrV40L and MrV40S, the nucleotide sequences of the genomes

were analyzed by ORFfinder (https://www.ncbi.nlm.nih.gov/orffinder/). To predict ribosomal

frameshift signals, a putative slippery site was identified using FSFinder2 (79). The amino

acid sequences of CP and RDRP of MrV40L were analyzed using the Pfam database

(http://pfam.xfam.org/) and BLASTP (https://blast.ncbi.nlm.nih.gov/). For sequence

alignment, Jalview2.0 was used (80). The phylogenetic tree showing the relationship of

MrV40L with RDRP of other dsRNA fungal viruses was constructed using MEGA X, and

phylogenetic reconstruction analysis was performed by the neighbor-joining method with

1,000 bootstrap replications (81, 82).

Mice and cell culture

Wild-type C57BL/6 mice were purchased from Orient Bio Co. (Gyeonggi-do, Korea). *TLR3*[-/-]

(B6;129S1-Tlr3[tm1Flv]/J, 005217) mice were obtained from Jackson Laboratory (Bar Harbor,

ME, USA). All animal experimental procedures were reviewed and approved by the

Institutional Animal Care and Use Committee of Hanyang University (protocol 2018-0085)

and performed in accordance with Korean Food and Drug Administration guidelines. All

animals were maintained in a specific pathogen-free environment. Primary BMDCs were

isolated from C57BL/6 mice and cultured in Dulbecco's modified Eagle medium for 3–5

days in the presence of 20 ng/mL recombinant granulocyte-macrophage colony-stimulating

factor (R&D Systems, Minneapolis, MN, USA) as described previously (83). Cell cultures

were stained to detect dendritic cells with CD11c-FITC (eBiosciences, San Diego, CA, USA).

Cytokine measurement

Mouse cytokines in the culture supernatants were measured with a BD OptEIA ELISA kit

(BD Biosciences, Franklin Lakes, NJ, USA) as described previously (84). All assays were

performed according to the manufacturer's instructions. Two independent strain pairs, virus-

infected and virus-cured *M. restricta* KCTC 27540 and virus-infected and virus-cured *M.*

restricta KCTC 27540, were used. To treat BMDCs with live fungal cells, a 20:1 ratio

$(2.0 \times 10^6$ and 1.0×10^5 cells/mL of *M. restricta* and BMDCs, respectively) was used. A final

concentration of 100 µg/mL (total protein/volume) of fungal cell lysates was used to treat

BMDCs. A final concentration of 1 µM of purified MrV40 capsid and satellite proteins and

500 ng/mL of purified MrV40 dsRNA from virus-infected M. restricta KCTC 27540 strain

were incubated with BMDCs. Phosphate-buffered saline (PBS) and polyinosinic-

polycytidylic acid [poly (I:C), 25 µg/mL] served as negative and positive controls,

respectively. Cytokines were measured 6 h after treatment. All experiments were repeated

four times.

cDNA synthesis, RNA isolation, transcriptome analysis by RNA sequencing,

quantitative real-time PCR, and heterologous expression and purification of MrV40

proteins

See Supplementary Materials and Methods

Data availability

The complete genome sequences of MrV40L and MrV40S were deposited into GenBank

under accession numbers MN603497 and MN603498, respectively. The transcriptome data

were deposited in Gene Express Omnibus under the accession number GSE138985.

Acknowledgments

We thank James W. Kronstad for critical reading of the manuscript and Taegun Seo for

helpful discussions. There is a companion paper from the Joe Heitman group at Duke

University, and we thank Joe Heitman and Giuseppe Ianiri for communication. This work

was supported by the National Research Foundation of Korea (NRF) grant funded by the

Korea government (MSIT) 2019R1F1A1061930 (WHJ), 2019R1A4A1024764 (WHJ), and

2019R1I1A2A01064237 (CSY).

References

1.   Hollings M. 1962. Viruses associated with a die-back disease of cultivated mushroom.

Nature 196:962.

2.   Ghabrial SA, Caston JR, Jiang D, Nibert ML, Suzuki N. 2015. 50-plus years of fungal

viruses. Virology 479-480:356-68.

3.   Bussey H. 1991. K1 killer toxin, a pore-forming protein from yeast. Mol Microbiol

5:2339-43.

4.   Hannig EM, Leibowitz MJ. 1985. Structure and expression of the M2 genomic

segment of a type 2 killer virus of yeast. Nucleic Acids Res 13:4379-400.

5.   Schmitt MJ, Tipper DJ. 1990. K28, a unique double-stranded RNA killer virus of

Saccharomyces cerevisiae. Mol Cell Biol 10:4807-15.

6.   Rodriguez-Cousino N, Maqueda M, Ambrona J, Zamora E, Esteban R, Ramirez M.

2011. A new wine Saccharomyces cerevisiae killer toxin (Klus), encoded by a double-

stranded rna virus, with broad antifungal activity is evolutionarily related to a

chromosomal host gene. Appl Environ Microbiol 77:1822-32.

7.   Cheng RH, Caston JR, Wang G-j, Gu F, Smith TJ, Baker TS, Bozarth RF, Trus BL,

Cheng N, Wickner RB. 1994. Fungal virus capsids, cytoplasmic compartments for the

replication of double-stranded RNA, formed as icosahedral shells of asymmetric Gag

dimers. Elsevier.

8.   Icho T, Wickner RB. 1989. The double-stranded RNA genome of yeast virus L-A

encodes its own putative RNA polymerase by fusing two open reading frames. J Biol

Chem 264:6716-23.

9.   Hopper JE, Bostian K, Rowe L, Tipper D. 1977. Translation of the L-species dsRNA

genome of the killer-associated virus-like particles of Saccharomyces cerevisiae.

Journal of Biological Chemistry 252:9010-9017.

10.   Fujimura T, Wickner RB. 1988. Gene overlap results in a viral protein having an RNA

binding domain and a major coat protein domain. Cell 55:663-671.

11.   Dinman JD, Icho T, Wickner RB. 1991. A -1 ribosomal frameshift in a double-

stranded RNA virus of yeast forms a gag-pol fusion protein. Proc Natl Acad Sci U S A

88:174-8.

12.   Van Alfen NK, Jaynes RA, Anagnostakis SL, Day PR. 1975. Chestnut Blight:

Biological Control by Transmissible Hypovirulence in Endothia parasitica. Science

189:890-1.

13.   Choi GH, Nuss DL. 1992. Hypovirulence of chestnut blight fungus conferred by an

infectious viral cDNA. Science 257:800-3.

14.   Ghabrial SA, Suzuki N. 2009. Viruses of plant pathogenic fungi. Annu Rev

Phytopathol 47:353-84.

15.   Segers GC, Zhang X, Deng F, Sun Q, Nuss DL. 2007. Evidence that RNA silencing

functions as an antiviral defense mechanism in fungi. Proc Natl Acad Sci U S A

104:12902-6.

16.   Zhang X, Segers GC, Sun Q, Deng F, Nuss DL. 2008. Characterization of hypovirus-

derived small RNAs generated in the chestnut blight fungus by an inducible DCL-2-

dependent pathway. J Virol 82:2613-9.

17.   Hammond TM, Andrewski MD, Roossinck MJ, Keller NP. 2008. Aspergillus

mycoviruses are targets and suppressors of RNA silencing. Eukaryot Cell 7:350-7.

18.   Yaegashi H, Shimizu T, Ito T, Kanematsu S. 2016. Differential Inductions of RNA

Silencing among Encapsidated Double-Stranded RNA Mycoviruses in the White Root

Rot Fungus Rosellinia necatrix. J Virol 90:5677-92.

19.   Yu J, Lee KM, Cho WK, Park JY, Kim KH. 2018. Differential Contribution of RNA

Interference Components in Response to Distinct Fusarium graminearum Virus

Infections. J Virol 92.

20.   Findley K, Oh J, Yang J, Conlan S, Deming C, Meyer JA, Schoenfeld D, Nomicos E,

Park M, Program NIHISCCS, Kong HH, Segre JA. 2013. Topographic diversity of

fungal and bacterial communities in human skin. Nature 498:367-70.

21.   Clavaud C, Jourdain R, Bar-Hen A, Tichit M, Bouchier C, Pouradier F, El Rawadi C,

Guillot J, Ménard-Szczebara F, Breton L. 2013. Dandruff is associated with

disequilibrium in the proportion of the major bacterial and fungal populations

colonizing the scalp. PloS one 8:e58203.

22.   Patino-Uzcategui A, Amado Y, Cepero de Garcia M, Chaves D, Tabima J, Motta A,

Cardenas M, Bernal A, Restrepo S, Celis A. 2011. Virulence gene expression in

Malassezia spp from individuals with seborrheic dermatitis. J Invest Dermatol

131:2134-6.

23.   Zhang E, Tanaka T, Tajima M, Tsuboi R, Nishikawa A, Sugita T. 2011.

Characterization of the skin fungal microbiota in patients with atopic dermatitis and in

healthy subjects. Microbiol Immunol 55:625-32.

24.   Park T, Kim HJ, Myeong NR, Lee HG, Kwack I, Lee J, Kim BJ, Sul WJ, An S. 2017.

Collapse of human scalp microbiome network in dandruff and seborrhoeic dermatitis.

Experimental dermatology 26:835-838.

25.   Wang L, Clavaud C, Bar☐Hen A, Cui M, Gao J, Liu Y, Liu C, Shibagaki N, Guéniche

A, Jourdain R. 2015. Characterization of the major bacterial–fungal populations

colonizing dandruff scalps in Shanghai, China, shows microbial disequilibrium.

Experimental dermatology 24:398-400.

26.   Xu Z, Wang Z, Yuan C, Liu X, Yang F, Wang T, Wang J, Manabe K, Qin O, Wang X.

2016. Dandruff is associated with the conjoined interactions between host and

microorganisms. Scientific reports 6:srep24877.

27. Grice EA, Dawson TL, Jr. 2017. Host-microbe interactions: Malassezia and human
642 skin. Curr Opin Microbiol 40:81-87.

28. Sharma S, Gupta S, Shrivastava JN. 2011. Presence of Virus like Particles in Human
644 Pathogenic Fungi: Chrysosporium sps and Candida albicans. Indian J Virol 22:104-10.

29. Park M, Cho YJ, Lee YW, Jung WH. 2017. Whole genome sequencing analysis of the
646 cutaneous pathogenic yeast Malassezia restricta and identification of the major lipase
647 expressed on the scalp of patients with dandruff. Mycoses 60:188-197.

30. EDY VG, SZEKELY M, LOVINY T, DREYER C. 1976. Action of Nucleases on
649 Double□Stranded RNA. European journal of biochemistry 61:563-572.

31. Lee SH, Yun S-H, Chun J, Kim D-H. 2017. Characterization of a novel dsRNA
651 mycovirus of Trichoderma atroviride NFCF028. Archives of virology 162:1073-1077.

32. Sommer SS, Wickner RB. 1982. Yeast L dsRNA consists of at least three distinct
653 RNAs; evidence that the non-Mendelian genes [HOK],[NEX] and [EXL] are on one
654 of these dsRNAs. Cell 31:429-441.

33. Park CM, Lopinski JD, Masuda J, Tzeng TH, Bruenn JA. 1996. A second double-
656 stranded RNA virus from yeast. Virology 216:451-4.

34. Taylor DJ, Ballinger MJ, Bowman SM, Bruenn JA. 2013. Virus-host co-evolution
658 under a modified nuclear genetic code. PeerJ 1:e50.

35. Bruenn JA. 1993. A closely related group of RNA-dependent RNA polymerases from
660 double-stranded RNA viruses. Nucleic Acids Res 21:5667-9.

36. Schmitt MJ, Breinig F. 2002. The viral killer system in yeast: from molecular biology
662 to application. FEMS microbiology reviews 26:257-276.

37. Wickner RB, Leibowitz MJ. 1979. Mak mutants of yeast: mapping and
characterization. J Bacteriol 140:154-60.

38. Tercero J, Dinman J, Wickner R. 1993. Yeast MAK3 N-acetyltransferase recognizes

the N-terminal four amino acids of the major coat protein (gag) of the LA double-

stranded RNA virus. Journal of bacteriology 175:3192-3194.

39. Tercero J, Wickner RB. 1992. MAK3 encodes an N-acetyltransferase whose

modification of the LA gag NH2 terminus is necessary for virus particle assembly.

Journal of Biological Chemistry 267:20277-20281.

40. Fujimura T, Wickner RB. 1987. LA double-stranded RNA viruslike particle

replication cycle in Saccharomyces cerevisiae: particle maturation in vitro and effects

of mak10 and pet18 mutations. Molecular and cellular biology 7:420-426.

41. Lee Y-J, Wickner R. 1992. MAK10, a glucose-repressible gene necessary for

replication of a dsRNA virus of Saccharomyces cerevisiae, has T cell receptor alpha-

subunit motifs. Genetics 132:87-96.

42. Wickner RB, Ridley SP, Fried HM, Ball SG. 1982. Ribosomal protein L3 is involved

in replication or maintenance of the killer double-stranded RNA genome of

Saccharomyces cerevisiae. Proceedings of the National Academy of Sciences

79:4706-4708.

43. Ohtake Y, Wickner RB. 1995. Yeast virus propagation depends critically on free 60S

ribosomal subunit concentration. Molecular and Cellular Biology 15:2772-2781.

44. Ryu S-W, Lee S-J, Park M-Y, Jun J-i, Jung Y-K, Kim E. 2003. Fas-associated factor 1,

FAF1, is a member of Fas death-inducing signaling complex. Journal of Biological

Chemistry 278:24003-24010.

45. Fujikura D, Miyazaki T. 2018. Programmed Cell Death in the Pathogenesis of

Influenza. Int J Mol Sci 19.

46. Jorgensen I, Rayamajhi M, Miao EA. 2017. Programmed cell death as a defence

against infection. Nat Rev Immunol 17:151-164.

47.   Carmona-Gutierrez D, Bauer MA, Zimmermann A, Aguilera A, Austriaco N,

Ayscough K, Balzan R, Bar-Nun S, Barrientos A, Belenky P. 2018. Guidelines and

recommendations on yeast cell death nomenclature. Microbial Cell 5:4.

48.   Newbold A, Martin BP, Cullinane C, Bots M. 2014. Detection of apoptotic cells using

propidium iodide staining. Cold Spring Harb Protoc 2014:1202-6.

49.   Banks GT, Buck KW, Chain EB, Himmelweit F, Marks JE, Tyler JM, Hollings M,

Last FT, Stone OM. 1968. Viruses in fungi and interferon stimulation. Nature

218:542-5.

50.   Kleinschmidt WJ, Cline JC, Murphy EB. 1964. Interferon Production Induced by

Statolon. Proc Natl Acad Sci U S A 52:741-4.

51.   Lampson GP, Tytell AA, Field AK, Nemes MM, Hilleman MR. 1967. Inducers of

interferon and host resistance. I. Double-stranded RNA from extracts of Penicillium

funiculosum. Proc Natl Acad Sci U S A 58:782-9.

52.   Perales-Linares R, Navas-Martin S. 2013. Toll-like receptor 3 in viral pathogenesis:

friend or foe? Immunology 140:153-67.

53.   Thompson MR, Kaminski JJ, Kurt-Jones EA, Fitzgerald KA. 2011. Pattern

recognition receptors and the innate immune response to viral infection. Viruses

3:920-40.

54.   Claudepierre MC, Hortelano J, Schaedler E, Kleinpeter P, Geist M, Remy-Ziller C,

Brandely R, Tosch C, Laruelle L, Jawhari A, Menguy T, Marchand JB, Romby P,

Schultz P, Hartmann G, Rooke R, Bonnefoy JY, Preville X, Rittner K. 2014. Yeast

virus-derived stimulator of the innate immune system augments the efficacy of virus

vector-based immunotherapy. J Virol 88:5242-55.

55.   Ishikawa T, Itoh F, Yoshida S, Saijo S, Matsuzawa T, Gonoi T, Saito T, Okawa Y,

Shibata N, Miyamoto T, Yamasaki S. 2013. Identification of distinct ligands for the C-

type lectin receptors Mincle and Dectin-2 in the pathogenic fungus Malassezia. Cell

Host Microbe 13:477-88.

56. Vargas G, Rocha JD, Oliveira DL, Albuquerque PC, Frases S, Santos SS, Nosanchuk

JD, Gomes AM, Medeiros LC, Miranda K, Sobreira TJ, Nakayasu ES, Arigi EA,

Casadevall A, Guimaraes AJ, Rodrigues ML, Freire-de-Lima CG, Almeida IC,

Nimrichter L. 2015. Compositional and immunobiological analyses of extracellular

vesicles released by Candida albicans. Cell Microbiol 17:389-407.

57. Samuel CE. 2001. Antiviral actions of interferons. Clinical microbiology reviews

14:778-809.

58. Brierley I, Boursnell ME, Binns MM, Bilimoria B, Blok VC, Brown TD, Inglis SC.

1987. An efficient ribosomal frame-shifting signal in the polymerase-encoding region

of the coronavirus IBV. EMBO J 6:3779-85.

59. Jacks T, Varmus HE. 1985. Expression of the Rous sarcoma virus pol gene by

ribosomal frameshifting. Science 230:1237-42.

60. Jacks T, Madhani HD, Masiarz FR, Varmus HE. 1988. Signals for ribosomal

frameshifting in the Rous sarcoma virus gag-pol region. Cell 55:447-58.

61. Magliani W, Conti S, Gerloni M, Bertolotti D, Polonelli L. 1997. Yeast killer systems.

Clinical microbiology reviews 10:369-400.

62. Tercero JC, Riles LE, Wickner RB. 1992. Localized mutagenesis and evidence for

post-transcriptional regulation of MAK3. A putative N-acetyltransferase required for

double-stranded RNA virus propagation in Saccharomyces cerevisiae. J Biol Chem

267:20270-6.

63. Wickner RB. 1996. Double-stranded RNA viruses of Saccharomyces cerevisiae.

Microbiological reviews 60:250.

64. Thrash C, Bankier AT, Barrell BG, Sternglanz R. 1985. Cloning, characterization, and

sequence of the yeast DNA topoisomerase I gene. Proceedings of the National

Academy of Sciences 82:4374-4378.

65. Widner WR, Wickner RB. 1993. Evidence that the SKI antiviral system of

Saccharomyces cerevisiae acts by blocking expression of viral mRNA. Molecular and

cellular biology 13:4331-4341.

66. Masison DC, Blanc A, Ribas JC, Carroll K, Sonenberg N, Wickner RB. 1995.

Decoying the cap-mRNA degradation system by a double-stranded RNA virus and

poly (A)-mRNA surveillance by a yeast antiviral system. Molecular and Cellular

Biology 15:2763-2771.

67. Benard L, Carroll K, Valle RC, Masison DC, Wickner RB. 1999. The ski7 antiviral

protein is an EF1-α homolog that blocks expression of non-Poly (A) mRNA in

Saccharomyces cerevisiae. Journal of virology 73:2893-2900.

68. McBride RC, Boucher N, Park DS, Turner PE, Townsend JP. 2013. Yeast response to

LA virus indicates coadapted global gene expression during mycoviral infection.

FEMS yeast research 13:162-179.

69. Wu M, Jin F, Zhang J, Yang L, Jiang D, Li G. 2012. Characterization of a novel

bipartite double-stranded RNA mycovirus conferring hypovirulence in the

phytopathogenic fungus Botrytis porri. J Virol 86:6605-19.

70. Wan J, Basu K, Mui J, Vali H, Zheng H, Laliberte JF. 2015. Ultrastructural

Characterization of Turnip Mosaic Virus-Induced Cellular Rearrangements Reveals

Membrane-Bound Viral Particles Accumulating in Vacuoles. J Virol 89:12441-56.

71. Matsumoto M, Oshiumi H, Seya T. 2011. Antiviral responses induced by the TLR3

pathway. Reviews in Medical Virology 21:67-77.

72. Tabeta K, Georgel P, Janssen E, Du X, Hoebe K, Crozat K, Mudd S, Shamel L,

Sovath S, Goode J. 2004. Toll-like receptors 9 and 3 as essential components of innate

immune defense against mouse cytomegalovirus infection. Proceedings of the

National Academy of Sciences 101:3516-3521.

73.   Hartley MA, Ronet C, Zangger H, Beverley SM, Fasel N. 2012. Leishmania RNA

virus: when the host pays the toll. Frontiers in Cellular and Infection Microbiology 2.

74.   Shelly Applen Clancey FR, Salomé LeibundGut-Landmann, Joseph Heitman,

Giuseppe Ianiri. 2019. A novel mycovirus evokes transcriptional rewiring in

Malassezia and provokes host inflammation and an immunological response. bioRxiv

doi:https://doi.org/10.1101/2019.12.18.880518.

75.   Kim M, Cho Y-J, Park M, Choi Y, Hwang SY, Jung WH. 2018. Genomic tandem

quadruplication is associated with ketoconazole resistance in Malassezia

pachydermatis. J Microbiol Biotechnol 28:1937-1945.

76.   Park M, Jung WH, Han SH, Lee YH, Lee YW. 2015. Characterisation and Expression

Analysis of MrLip1, a Class 3 Family Lipase of Malassezia restricta. Mycoses

58:671-8.

77.   Leeming JP, Notman FH. 1987. Improved methods for isolation and enumeration of

Malassezia furfur from human skin. J Clin Microbiol 25:2017-9.

78.   Okada R, Kiyota E, Moriyama H, Fukuhara T, Natsuaki T. 2015. A simple and rapid

method to purify viral dsRNA from plant and fungal tissue. Journal of General Plant

Pathology 81:103-107.

79.   Moon S, Byun Y, Kim HJ, Jeong S, Han K. 2004. Predicting genes expressed via -1

and +1 frameshifts. Nucleic Acids Res 32:4884-92.

80.   Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. 2009. Jalview Version

2--a multiple sequence alignment editor and analysis workbench. Bioinformatics

25:1189-91.

81.   Kumar S, Nei M, Dudley J, Tamura K. 2008. MEGA: a biologist-centric software for

evolutionary analysis of DNA and protein sequences. Brief Bioinform 9:299-306.

82.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol 35:1547-

1549.

83.   Vargas G, Rocha JD, Oliveira DL, Albuquerque PC, Frases S, Santos SS, Nosanchuk

JD, Gomes AMO, Medeiros LC, Miranda K. 2015. Compositional and

immunobiological analyses of extracellular vesicles released by C andida albicans.

Cellular microbiology 17:389-407.

84.   Koh H-J, Kim Y-R, Kim J-S, Yun J-S, Kim S, Kim SY, Jang K, Yang C-S. 2018.

CD82 hypomethylation is essential for tuberculosis pathogenesis via regulation of

RUNX1-Rab5/22. Experimental & molecular medicine 50:62.

85.   Saitou N, Nei M. 1987. The neighbor-joining method: a new method for

reconstructing phylogenetic trees. Mol Biol Evol 4:406-25.

86.   Felsenstein J. 1985. Confidence Limits on Phylogenies: An Approach Using the

Bootstrap. Evolution 39:783-791.

87.   Sanderlin RS, Ghabrial SA. 1978. Physicochemical properties of two distinct types of

virus-like particles from Helminthosporium victoriae. Virology 87:142-51.

88.   Shapira R, Choi GH, Nuss DL. 1991. Virus-like genetic organization and expression

strategy for a double-stranded RNA genetic element associated with biological control

of chestnut blight. EMBO J 10:731-9.

89.   Hillman BI, Halpern BT, Brown MP. 1994. A viral dsRNA element of the chestnut

blight fungus with a distinct genetic organization. Virology 201:241-50.

90.   Van der Lende TR, Duitman EH, Gunnewijk MG, Yu L, Wessels JG. 1996. Functional

analysis of dsRNAs (L1, L3, L5, and M2) associated with isometric 34-nm virions of

Agaricus bisporus (white button mushroom). Virology 217:88-96.

91.   Nomura K, Osaki H, Iwanami T, Matsumoto N, Ohtsu Y. 2003. Cloning and

characterization of a totivirus double-stranded RNA from the plant pathogenic fungus,

Helicobasidium mompa Tanaka. Virus Genes 26:219-26.

92.   Tuomivirta TT, Hantula J. 2003. Two unrelated double-stranded RNA molecule

patterns in Gremmeniella abietina type A code for putative viruses of the families

Totiviridae and Partitiviridae. Arch Virol 148:2293-305.

93.   Jiang D, Ghabrial SA. 2004. Molecular characterization of Penicillium chrysogenum

virus: reconsideration of the taxonomy of the genus Chrysovirus. J Gen Virol

85:2111-21.

94.   Urayama S, Ohta T, Onozuka N, Sakoda H, Fukuhara T, Arie T, Teraoka T, Moriyama

H. 2012. Characterization of Magnaporthe oryzae chrysovirus 1 structural proteins

and their expression in Saccharomyces cerevisiae. J Virol 86:8287-95.

95.   Wu Q, Luo Y, Lu R, Lau N, Lai EC, Li WX, Ding SW. 2010. Virus discovery by deep

sequencing and assembly of virus-derived small silencing RNAs. Proc Natl Acad Sci

U S A 107:1606-11.

96.   Isawa H, Kuwata R, Hoshino K, Tsuda Y, Sakai K, Watanabe S, Nishimura M, Satho

T, Kataoka M, Nagata N, Hasegawa H, Bando H, Yano K, Sasaki T, Kobayashi M,

Mizutani T, Sawabe K. 2011. Identification and molecular characterization of a new

nonsegmented double-stranded RNA virus isolated from Culex mosquitoes in Japan.

Virus Res 155:147-55.

97.   Mor SK, Phelps NB. 2016. Molecular detection of a novel totivirus from golden

shiner (Notemigonus crysoleucas) baitfish in the USA. Arch Virol 161:2227-34.

98.   Yang X, Zhang Y, Ge X, Yuan J, Shi Z. 2012. A novel totivirus-like virus isolated

from bat guano. Arch Virol 157:1093-9.

99.   Zhai Y, Attoui H, Mohd Jaafar F, Wang HQ, Cao YX, Fan SP, Sun YX, Liu LD,

840   Mertens PP, Meng WS, Wang D, Liang G. 2010. Isolation and full-length sequence

841   analysis of Armigeres subalbatus totivirus, the first totivirus isolate from mosquitoes

842   representing a proposed novel genus (Artivirus) of the family Totiviridae. J Gen Virol

843   91:2836-45.

844 100. Dihanich M, Van Tuinen E, Lambris J, Marshallsay B. 1989. Accumulation of

845   viruslike particles in a yeast mutant lacking a mitochondrial pore protein. Molecular

846   and cellular biology 9:1100-1108.

847 101. Leibowitz MJ, Wickner RB. 1978. pet 18: A chromosomal gene required for cell

848   growth and for the maintenance of mitochondrial DNA and the killer plasmid of yeast.

849   Molecular and General Genetics MGG 165:115-121.

850 102. Toh□e A, Sahashi Y. 1985. The PET18 locus of Saccharomyces cerevisiae: a complex

851   locus containing multiple genes. Yeast 1:159-171.

852 103. Thrash C, Voelkel K, DiNardo S, Sternglanz R. 1984. Identification of

853   Saccharomyces cerevisiae mutants deficient in DNA topoisomerase I activity. Journal

854   of Biological Chemistry 259:1375-1377.

855 104. Icho T, Wickner RB. 1988. The MAK11 protein is essential for cell growth and

856   replication of M double-stranded RNA and is apparently a membrane-associated

857   protein. Journal of Biological Chemistry 263:1467-1475.

858 105. Wickner RB. 1988. Host function of MAK16: G1 arrest by a mak16 mutant of

859   Saccharomyces cerevisiae. Proceedings of the National Academy of Sciences

860   85:6007-6011.

861 106. Carroll K, Wickner RB. 1995. Translation and M1 double-stranded RNA propagation:

862   MAK18= RPL41B and cycloheximide curing. Journal of bacteriology 177:2887-2891.

863 107. Toh-E A, Guerry P, Wickner RB. 1978. Chromosomal superkiller mutants of

864   Saccharomyces cerevisiae. Journal of bacteriology 136:1002-1007.

865   108.   Ball SG, Tirtiaux C, Wickner RB. 1984. Genetic control of LA and L-(BC) dsRNA
866   copy number in killer systems of Saccharomyces cerevisiae. Genetics 107:199-217.
867   109.   Ridley SP, Sommer SS, Wickner RB. 1984. Superkiller mutations in Saccharomyces
868   cerevisiae suppress exclusion of M2 double-stranded RNA by LA-HN and confer cold
869   sensitivity in the presence of M and LA-HN. Molecular and Cellular Biology 4:761-
870   770.
871   110.   Dmochowska A, Dignard D, Henning D, Thomas DY, Bussey H. 1987. Yeast KEX1
872   gene encodes a putative protease with a carboxypeptidase B-like function involved in
873   killer toxin and α-factor precursor processing. Cell 50:573-584.
874   111.   Wickner RB. 1974. Chromosomal and nonchromosomal mutations affecting the"
875   killer character" of Saccharomyces cerevisiae. Genetics 76:423-432.
876   112.   Wickner RB, Leibowitz MJ. 1976. Two chromosomal genes required for killing
877   expression in killer strains of Saccharomyces cerevisiae. Genetics 82:429-442.

882 **Figure legends**

885   **Tables**

887   **Table 1. Number of differentially expressed genes**

	Number of differential expressed genes (+virus/-virus)				
	2-fold	3-fold	4-fold	>5-fold	Total
Upregulated	204	32	12	10	258
Downregulated	597	152	70	60	809

     Table 2. Differential expression of genes involved in maintaining dsRNA virus

S. cerevisiae gene	An encoded protein and/or function	Needed by *S. cerevisiae* virus	*M. restricta* gene ID	Fold change (+virus/−virus)	Reference(s)
MAK3	N-acetyltransferase modifying Gag		MRET_0174	0.62	(39, 62)
MAK10	Non-catalytic subunit of N-terminal acetyltransferase	L-A, M	No hits		(41, 100)
PET18 (MAK30+MAK31)	Unknown		No hits		(101, 102)
MAK1 (TOP1)	DNA topoisomerase I		MRET_0706	0.41	(103)
MAK2			MRET_4192	0.88	
MAK5			MRET_0265	0.44	(43)
MAK21	60S subunit biosynthesis		MRET_2922	0.24	
MAK11		M	MRET_2252	0.40	(43, 104)
MAK16			MRET_2745	0.48	(43, 105)
MAK7 (RPL4A)	60S subunit protein L4		No hits		(43)
MAK8 (TCM1)	60S subunit protein L3		MRET_1592	0.24	(42)
MAK18 (RPL41B)	60S subunit protein L41		MRET_2856	0.60	(106)
SKI1 (XRN1)	5′-3′ exonuclease	L-A, M	MRET_4129	0.73	(107)

SKI2	RNA helicase		MRET_2647	0.91	
SKI3	Tetratricopeptide repeat protein		MRET_1481	0.78	
SKI4 (*CSL4*)	Exosome non-catalytic core component		MRET_2576	0.96	(108, 109)
SKI6	Exosome non-catalytic core component		MRET_3188	0.55	
SKI7	GTP-binding protein		No hits		
SKI8	WD-repeat protein		No hits		
KEX1	Cell death protease essential for hypochlorite-induced apoptosis	M	MRET_4176	0.93	(110-112)
KEX2	Kexin, a subtilisin-like protease (proprotein convertase)		MRET_0618	1.13	(111, 112)

892 **Table 3. Differential expression of genes involved in TCA cycle and electron transport**

893 **chain**

Gene	Annotation	Fold change (+virus/−virus)
MRET_1104	NADH dehydrogenase (ubiquinone) Fe-S protein 7	2.85
MRET_1378	succinate dehydrogenase (ubiquinone) iron-sulfur subunit	3.55
MRET_1709	cytochrome c	2.56
MRET_1953	NADH dehydrogenase (ubiquinone) Fe-S protein 1	2.44
MRET_2042	fumarate hydratase, class II	2.42
MRET_2097	succinate dehydrogenase (ubiquinone) flavoprotein subunit	2.19
MRET_2956	2-oxoglutarate dehydrogenase E1 component	2.21
MRET_3173	dihydrolipoamide dehydrogenase	2.27
MRET_4117	aconitate hydratase	3.55

     Figures

Fig. 1. Extrachromosomal dsRNA segments in *Malassezia*. (A) Nucleic acids from

Malassezia restricta KCTC 27540 were separated on a 0.7% agarose gel. Lane 1, total

nucleic acids; lane 2, total nucleic acids treated with DNase I; lane 3, total nucleic acids

treated with RNase A; lane 4, total nucleic acids treated with RNase T1. (B) Presence of

dsRNA in *M. restricta* strains. Total nucleic acids were extracted and treated with DNase I

and RNase T1 and were separated on a 0.7% agarose gel. (C) Total nucleic acids were

extracted from different *Malassezia* species, treated with DNase I and RNase T1, and were

separated on a 0.7% agarose gel.

Fig. 2. Isolation of virus particles from *Malassezia restricta* KCTC 27540, and their TEM images. The collected fractions after sucrose gradient ultracentrifugation were analyzed for their nucleic acids and proteins on an agarose gel (A) and SDS-PAGE gel (B), respectively. Images of the viral particles (red arrows) were obtained using a transmission electron microscope (C).

Fig. 3. Morphology of vacuoles in virus-infected and virus-cured *Malassezia restricta*

cells. Ultrathin sections of virus-infected (A) and virus-cured (B) *M. restricta* KCTC 27540

strains were observed by TEM. Cytoplasmic vacuole formation in a cell, expressed as the

percentage of the area occupied by vacuoles (C) and the number of vacuoles (D) per cell, was

estimated (n = 100 cells for each strain). The plot was created with BioVinci software

(BioTuring Inc., San Diego, CA, USA). Statistical analysis for differences between the strains

was performed with unpaired t-tests. *p < 0.001.

Fig. 4. Genome and phylogenetic analysis of MrV40L. (A) Comparison of the genomic

organization of MrV40L with that of other totiviruses. (B) Eight conserved motifs within the

viral RDRP of MrV40L. The red boxes indicate conserved motifs. The amino acid sequences

of ScVL-A, ScVL-BC, SsVL, and MrV40L were aligned using Jalview. (D) Multiple

alignment of 18 amino acid sequences of RDRPs from dsRNA viruses was analyzed using the

neighbor-joining method with bootstrap test (1,000 replicates) using MEGA X (82, 85, 86).

The evolutionary distances are in units of the number of amino acid substitutions per site. All

ambiguous positions were removed for each sequence pair. CHV1, *Cryphonectria* hypovirus

1; CHV2, *Cryphonectria* hypovirus 2; AsV, *Armigeres subalbatus* virus; DTV, *Drosophila*

melanogaster totivirus; OMRV, *Omono river* virus; ToV, *Tianjin* totivirus; GSTV, *Golden*

shiner totivirus; ScVL-A, *S. cerevisiae* virus L-A; ScVL-BC, *S. cerevisiae* virus L-BC; SsVL,

S. segobienesis virus L; GaRV-L1, *Gremmeniella abietina* virus L1; Hv190SV,

Helminthosporium victoriae virus-190S; MoV1, *Magnaporthe oryzae* virus 1; HmV-17,

Helicobasidium mompa totivirus 1-17; PcV, *Penicillium chrysogenum* virus; Hv145SV,

Helminthosporium victoriae 145S virus; AbV1, *Agaricus bisporus* virus 1 (8, 33, 34, 87-99).

Fig. 5. Phylogenetic classification of viruses found in *Malassezia restricta*. (A, B) The

cDNAs of dsRNAs from *M. restricta* strains were amplified using primers for genes encoding

CP and RDRP (ORF1 and ORF2, respectively) homologs in the viral genome. The following

primers were used: lane 1, MrV40L_CP_F1 and MrV40L_CP_R1; lane 2, MrV40L_CP_F2

and MrV40L_CP_R2; lane 3, MrV40L_RDRP_F1 and MrV40L_RDRP_R1; lane 4,

MrV40L_RDRP_F2 and MrV40L_RDRP_R2 (see Table S2 in supplemental material). (C)

MrV-Ls were clustered into three clades (clade ▯, ▯, and ▯). Multiple alignment of

nucleotide sequences of combined parts of *gag* and *pol* from 11 MrV-Ls was analyzed by the

neighbor-joining method with bootstrap test (1,000 replicates) in MEGA X (82, 85, 86). The

evolutionary distances are in units of the number of base substitutions per site.

Fig. 6. Characterization and similarity analysis of MrV40S. (A) The protein encoded by

ORF3 was heterologously expressed in *E. coli* and purified using a His-tag column. (B) RT-

PCR results. The primers specific to MrV40S were used to amplify satellite dsRNA from

other *M. restricta* strain containing small dsRNA segments. The following primers were used:

lane 1, MrV40S_F5 and V40S_SP1; lane 2, MrV40S_ORF_F1 and MrV40S_R1 (see Table

S2 in supplemental material).

Fig. 7. Evaluation of TLR and cytokine levels upon treatment with the viral elements.

TLR expression (A) and cytokine production (B) in BMDCs co-incubated with live fungal cells, total fugal cell lysates, purified MrV40 capsid protein, purified satellite protein, and purified MrV40 dsRNA. PBS and poly (I:C) served as negative and positive controls, respectively. WT, wild-type mice (C57BL/6); *tlr3*[-/-], TLR3 knock out mice (B6;129S1-Tlr3[tm1Flv]/J, 005217); FC, live fungal cells; LY, cell lysate; 40+, virus-infected KCTC 27540; 40-, virus-cured KCTC 27540; 24+, virus-infected KCTC 27524; 24-, virus-cured KCTC 27524; CP, purified capsid protein; p15, purified MrV40S satellite protein; dsRNA, purified

983   MrV40 dsRNA. Values represent the average from quadruplicates with standard deviations.

984   Statistical analysis for differences between PBS and each sample was performed with

985   unpaired t-tests. *p < 0.001 and **p < 0.0001.