Full title:

Increased juvenile survival may not be universally linked to longevity: ecological, social and life-history
drivers of age-specific mortality in birds

Authors:

Emeline Mourocq[1]*, emeline.mourocq@laposte.net
Szymon M. Drobniak[2,3], szymek.drobniak@uj.edu.pl
Michael Griesser[1,4], michael.griesser@gmail.com
9
10 # Authors Affiliations:

11
[1]*University of Zurich, Department of Anthropology, University of Zurich, Zurich, Switzerland*
[2]*Institute of Environmental Sciences, Jagiellonian University, Kraków, Poland*
[3]*School of Biological, Environmental and Earth Sciences, University of New South Wales, Sydney,*
Australia
[4]*State Key Laboratory of Biocontrol, Department of Ecology and School of Life Sciences, Sun Yat-sen*
University, Guangzhou, 510275, China
18
*Corresponding author:

Name: Emeline Mourocq
Address: calle Vela y Mora 43 2ºIz, 11660 Prado del Rey (Cadiz, Spain)
Phone: +34692186305
e-mail: emeline.mourocq@laposte.net

Short title:

Juvenile survival-longevity relationship

Abstract

A classical prediction of the traditional evolutionary theories of ageing (tETA) is that longevity should

be positively correlated with survival early on in life. However, large and unexplained variation exists

in juvenile survival-longevity combinations. Here, we provide the first comparative study

investigating the life-history, ecological and social correlates of juvenile survival, longevity and their

combinations in 204 bird species. Overall, both measurements were positively correlated, but

multiple survivals' combinations evolved, some in accordance with tETA ("positive JS-L

combinations") while others contrasting it ("JS-L mismatches"). Positive JS-L combinations covaried

with the pace of life proxies, whereas mismatching combinations covaried with the growing season

length, where long growing seasons promoted juvenile survival, while short growing seasons

promoted longevity. Interestingly, sociality explained only positive combinations, while life-history

and ecological traits explained both positive and mismatching combinations. Overall, these findings

challenge a main prediction of the tETA, and identify key evolutionary forces driving the coevolution

between juvenile survival and longevity.

Key words:

Traditional theories of aging (tETA: "mutation accumulation"[1], "antagonistic pleiotropy"[2],

"antagonistic pleiotropy"[3]) propose that extrinsic mortality is the main driver of longevity[4,5]. They

predict that higher extrinsic mortality early on in life leads to relatively few individuals reaching old

age, and the fitness value of prolonged lifespan is therefore small in such cases. Thus, selection to

extend longevity is only strong in populations with high survival early on in life (juvenile survival

henceforth). Accordingly, these theories predict that longevity should be positively correlated with

juvenile survival[2,4,6-8].

Although this classical prediction of tETA underlies many life-history studies, and is

commonly cited as being largely corroborated by existing data[9,10], support has been mixed and

alternative theories exist[5,11-13]. Moreover, recent theoretical and empirical studies do challenge this

prediction[9,14-16], and state that juvenile survival (extrinsic mortality early on in life) is not a random

process but does depend on age, individual condition, or population density. Accordingly, species

can deviate from the expected relationship between juvenile survival and longevity, by having a low

juvenile survival but being long-lived, or by having a high juvenile survival but being short-lived[5,9,13-

17]. However, it remains unclear whether these deviations represent evolved strategies modulated by

specific life-history, ecological and/or social factors, or whether they are pieces of a continuum of

randomly varying combinations.

Longevity is a pivotal factor shaping life-histories[18,19], but survival can vary among the stages

of life and differently influence the evolution of life-history traits[20-22]. Theoretical[12,23-26] and empirical

work on birds[27], fishes[22] and mammals[28] have highlighted the importance of considering age-specific

survival to understand the evolution of life-history traits. Specifically, these studies showed that age-

specific survival patterns that deviate from the classical prediction of tETA (e.g., low chance of

survival early in life but a high longevity), are linked to unusual combinations of life-history traits that

are characteristic to both slow- and fast-living animals[19]. For instance, turtles and crocodiles suffer

from high juvenile mortality, and accordingly females lay many eggs in each reproductive event (like

fast-living animals) despite that they are exceptionally long-lived (like slow-living animals)[29]. Thus,

considering the factors affecting age-specific survivals and their combination is critical to understand

life history evolution in general.

Longevity varies considerably across species. In vertebrates it ranges from a few months to

over 100 years[30]. Comparative work did show that adaptations that reduce extrinsic mortality,

including protective shells[6] or the ability to fly[31], are linked with increased longevity. Moreover, long-

lived species tend to be active during the period of day with the lowest predation risk[31], have a low

number of co-occurring predators of adults[32] or life-history traits characteristic of a slow pace of life

(e.g., produce few offspring, which develop slowly and mature relatively late in life)[12,18]. Additionally,

larger mammals and birds live longer than smaller ones[32-34].

In many taxa, juveniles usually have lower and more variable survival than adults[35-38]. The

few studies investigating juvenile survival showed that the small body size of juveniles may explain

their low survival in lineages with slow growth (mammals, reptiles) and indeterminate growth

(fish)[39,40]. In lineages with rapid body growth (birds), low juvenile survival can reflect age-dependent

social dominance[35] or lacking skills[41]. Besides, juvenile survival tends to be high in birds with long

nestling periods[42], low reproductive allocation[43,44], prolonged post-fledging care[45], or prolonged

association with the parents beyond independence (i.e., family-living species, see[46])[47,48] (Table 1).

Although a number of studies have investigated inter-specific variation in longevity[31,32,34,49,50], it is

unknown which factors influence survival early on in life and how this relates to longevity[51].

Importantly, comparative studies are lacking.

Here, we use phylogenetic comparative analyses to understand interspecific variation in

juvenile survival (measured as post-fledglings to first-year survival rate) and maximum longevity, as

well as their relationship, in 204 bird species. Firstly, we compare the association of (i) juvenile

survival and (ii) maximum longevity with life-history, ecological and social parameters. Secondly, we

investigate how juvenile survival and maximum longevity relate to each other, and assess which life-

history, ecology and social traits better explain (i) positive associations between juvenile survival and

longevity (i.e., as expected by the classical prediction of tETA: low-low and high-high combinations,

referred to as "positive JS-L combinations" henceforth), and (ii) mismatches between juvenile

survival and longevity (i.e., deviation from the classical prediction of tETA: low-high and high-low

combinations, referred to as "JS-L mismatches" henceforth).

Table 1| Description and prediction of the parameters investigated in this study.

	parameter's name	description	prediction	source prediction
life-history	adult body mass*	mean adult body mass (g)	larger body size confers better ability to cope with temporary food shortages, climatic fluctuations and extreme weather than smaller body size; large body size may associate with higher juvenile survival and higher longevity	[39,52]
	incubation period*	number of days from laying to hatching	longer incubation period may associate with higher juvenile survival and higher longevity	[12,18]
	nestling period*	number of days from hatching to fledging	longer nestling period may associate with higher juvenile survival and higher longevity	[42]
	annual parental investment*	body-mass scaled annual reproductive investment (total mass of eggs produced annually divided by adult body mass)[a]	higher parental investment may associate with lower juvenile survival and lower longevity	[12,18,53]
	chick development mode	precocial vs. non precocial; semi-altricial or semi precocial species were categorised as non precocial	precocial species should have lower juvenile survival but higher longevity because of lower parental care after hatching while the opposite is expected for altricial species	[54]
ecological	sedentariness	resident vs. migratory; based on the species maximum movement; sedentary species or with local movement were categorised as resident and the one with regional or inter-continental movement as migratory	costs associated with migration could translate into lower juvenile survival and lower longevity in migratory than in non-migratory species	[55]
	period of activity	diurnal vs. nocturnal; crepuscular species (i.e. active at dawn and dusk) were categorised as nocturnal	species that are active at night are likely to be harder for predators to detect and predators are more scarce at night thus, nocturnal species might have higher juvenile survival and live longer than diurnal species	[31]
	nest predation risk*	based on both most commonly used nest location and nest type; ordinally ranked: 1 = inaccessible nests in cavities, 2 = open nests in cliffs or tree, 3 = open nest in shrub-layer or the ground[b]	nest predation risk may alter the developmental phase of the nestling and the reproductive effort of the parents which may affect juvenile survival and longevity; greater nest predation risk may associate with lower juvenile survival and lower longevity	[20]
	foraging exposure*	level of exposure to predators during foraging time based on most commonly used foraging area; ordinally ranked: 1 = pelagic species, 2 = aerial foragers, 3 = terrestrial foragers	pelagic or aerial forager should have lower predation risk and be more capable of escaping from predators than species that feed on the ground; juvenile survival and longevity may be reduced in the latter more than in the formers	[56,57]
	vegetation cover*	cover of woody vegetation in habitat (%)	more open habitats provide less visual cover than habitats dense in vegetation, increasing the risk of being killed; thus, low vegetation density may associate with lower juvenile survival and lower longevity. The reverse may be true if vegetation	[57-59]

		cover, by obstructing the view of the prey, affects its survival	
	caloric content of food*	energy content of the food in kcal/100g[(c)]	food calory content can influence the energy available for maintenance; high calory diet may associate with higher juvenile survival and higher longevity	60,61
	fibre content of food*	fibre food content in g/100g[(c)]	food fibre content can influence digestion efficiency and thus the level of resource acquired and health; high fibre diet may associate with higher juvenile survival and higher longevity	60,61
	foraging cost*	level of energy demand for foraging based on most commonly used foraging technics; ordinally ranked: 1 = sit and wait hunters, 2 = swimming or short perch & short flights, 3 = aerial or under water foraging, 4 = terrestrial or gleaners [(d)]	species with highly energetically demanding foraging strategies may have lower juvenile survival and lower longevity than species with less energetically demanding technics	62
	diet specialisation	specialist (only one diet class) vs. generalist (more than one diet class)	a change in the food availability can have higher costs for specialist than generalist species as the later can deviate to other food resources; specialisation may associate with lower juvenile survival and lower longevity	63
	habitat specialisation	specialist (only one habitat type) vs. generalist (more than one habitat type)[(e)]	a change in habitat availability can have higher costs for specialist than generalist species as the latter can occupy other habitat types; specialisation is predicted to associate with lower juvenile survival and lower longevity	64,65
	MGS duration*	mean duration of the growing season in months[(f)] (i.e., month(s) of the yeuyuar in which temperature and rainfall allow significant plant productivity)	a short growing season implies changes in environmental conditions over the year, thus MGS duration can be seen as a proxy of environemental variability; less variable environments (long growing season) may associate with higher juvenile survival and longevity than highly variable environment (short growing season)	66
	region	breeding distribution range: northern or southern hemisphere, both hemispheres, island	southern hemisphere and island species may have higher juvenile survival and higher longevity compared to northern hemisphere species	34,66,67
	N avian predators*	number of sympatric adult's or independent juveniles' predator species[(g)]	a higher number of predators increases the risk of being predated; higher number of predators may associate with a lower juvenile survival and lower longevity	32
Social	parental care mode	uniparental, biparental, cooperative breeding	the presence of additional carer can reduce survival risks on young and survival costs on the other carer(s); biparental and cooperative breeding species may have higher juvenile survival and higher longevity than uniparental species	68,but see[69,70]
	social system	family living (offspring remain at least 50 days beyond nutritional independence with parents) vs. non-family living[(h)]	species with prolonged post-fledging parental care or having a prolonged association with the parents beyond independence, as in family-living species, may have higher juvenile survival and lower longevity	45,71

Because experience (e.g., foraging, competition, reproductive strategies, anti-predation behaviours)
differs between young individuals (inexperienced) and adults (experienced), we also assumed each of
the abovementioned parameters to diferentially influence juvenile survival and longevity, and
potentially explain variation in juvenile survival/longevity relationships.
* Included in the PCA (Table 2). The other parameters are categorical variables. (a)[56],
(b)[72,73], (c)[60], (d)[74], (e)[75], (f)[76], (g)[32], (h)[46].

Materials and Methods

Survival data. We collected data on juvenile survival and maximum longevity for 293 bird species

covering 20 taxonomic orders and 74 families (Fig. S1 in Supporting Information), using existing

datasets[32], the Handbook of the Birds of the World[77], the Birds of North America[78], the Handbook of

Australian, New Zealand and Antarctic Birds[79], the Handbook of Southern Africa[80], the Australian

Birds and Bats Banding Scheme database[81] and Animal Ageing and Longevity database[82] (available at

http://genomics.senescence.info/species/).

Juvenile survival was assessed as the proportion of fledglings that survive their first year of

life, where many juveniles die due to extrinsic mortality[83]. For species where multiple values of

juvenile survival were available we used their mean. Maximum longevity (maximum observed

lifespan) was mostly assessed with mark-recapture of ringed wild birds, but for 19 species longevity

was of unknown origin (captivity or wild). Earlier studies showed that longevity records in captivity

and the wild are highly correlated[32,34] and thus, we also included longevity data of unknown origin.

Longevity estimates are influenced by the sampling effort because the larger the sample the higher

is the chance to sample a long-lived indvidual[32]. Therefore, to adjust for any bias associated with

maximum longevity estimates we included the independent number of Web of Science records per

species (research effort) as a covariate in our analyses (available at

http://apps.webofknowledge.com).

Life-history, ecology and social parameters. We used a published dataset[84] that was

complemented with data from the sources listed above, and compiled data on life-history, ecological

and social parameters that may influence juvenile survival and longevity (Table 1). We could find

data for the 20 parameters listed in Table 1 for 204 of the 293 species (Fig. S2). Thus, 293 species

were considered in descriptive analyses, while a subset 204 species entered detailed phylogenetic

mixed models.

Statistical analyses. *General procedures.* All statistical analyses were performed in R version

3.2.2[85]. We used phylogenetic controlled mixed models in ASReml-R 3[86] to control for the

phylogenetic dependency among species (VSN International, Hempstead, U.K.[87]). We included

phylogeny as a random effect in the model in the form of a correlation matrix of distances from the

root of the tree to the most recent common ancestor between two species. We tested the

phylogenetic effect with a likelihood ratio test where 2 times the difference in log-likelihood

between the model with and without the phylogeny is tested against a χ^2 distribution with one

degree of freedom[88]. To account for phylogenetic uncertainty, all ASReml-R models were run with

300 different phylogenetic trees obtained from www.birdtree.org[89]. We averaged the estimates

from the 300 models and present the averaged estimates and the Fs_{300} (proportion of trees for

which the p-value associated with an estimate was <0.05). Individual p-values were obtained

through a conditional Wald F-test. All continuous variables were standardised by centring (around

the mean) and scaling (by the standard deviation) them, to allow direct comparison of the model

estimates[90], but we present raw data in the figures. We checked for the assumptions of normally

distributed and homogeneous residuals by visually inspecting histograms and qq-plots of the

residuals as well as residuals plotted against fitted values.

To reduce the multidimensionality of our predictor variables and to reduce their

collinearity[91], we performed a principal component analysis (PCA) with varimax rotation including all

12 continuous predictors, and extracted 7 PC's given in Table 2. Prior to the PCA, the distribution of

these predictors was checked graphically and, if necessary, transformed to obtain a more

symmetrical distribution, and then standardised (see above).

Table 2 | Results of the Principal Component Analysis (PCA) with varimax rotation on the 12 continuous predictors.

category	transformation	variable	nº component	life-history pace	exposure to predators	food fibre	nest predation risk	N avian predators	foraging cost	MGS duration	h2
				1	7	2	6	3	5	4	h2
life-history	ln	adult body mass		**0.90**	-0.25	-0.08	0.13	-0.01	-0.17	-0.12	0.95
	none	annual parental investment		**-0.89**	0.14	-0.17	0.13	0.14	-0.05	-0.02	0.87
	sqrt	incubation period		**0.76**	-0.35	0.23	-0.03	-0.10	-0.28	-0.04	0.84
	none	nestling period		0.51	0.04	0.44	-0.58	-0.13	0.01	0.20	0.85
ecological	none	nest predation risk		0.05	-0.10	-0.12	**0.94**	-0.12	0.09	-0.07	0.93
	none	foraging cost		-0.17	0.02	-0.11	0.08	-0.07	**0.96**	-0.05	0.98
	none	calorie content of food		0.15	0.49	0.62	0.02	0.44	0.01	-0.01	0.83
	ln	fibre content of food		-0.11	0.16	**-0.89**	0.22	-0.01	0.15	-0.10	0.92
	none	foraging exposure		-0.30	**0.86**	-0.04	-0.01	0.04	-0.02	0.00	0.83
	sqrt	vegetation cover		-0.21	**0.85**	-0.01	-0.13	-0.08	0.05	0.11	0.80
	sqrt	N avian predators		-0.18	-0.03	0.08	-0.07	**0.95**	-0.05	0.00	0.95
	none	MGS duration		-0.06	0.08	0.08	-0.11	-0.01	-0.05	**0.98**	0.99
		SS loadings		2.68	1.94	1.5	1.34	1.15	1.07	1.05	
		cumulative variance explained (%)		22	39	51	62	72	81	89	

We considered coefficients of correlation greater than 0.7 or less than -0.7 to be high loadings (highlighted in bold). h2 is the communality of the 7
components. ln: natural logarithm, sqrt : Square root

Full mixed models included the 7 PC's (Table 2), the 8 categorical variables described in Table

1, and as covariates research effort (log transformed) and body mass (log transformed) to control for

allometry[32,49]. Since the life-history pace PC was loaded by adult body mass (Table 2) and therefore

partially controlling for allometry, we only included the residuals from a linear model between the

natural logarithm of adult body mass and the life-history pace component as body-mass covariate.

This way the presence in the model of both the life-history pace PC and the residual body mass

allows to fully control for allometry.

The importance of first-year survival for fitness benefits is likely to depend on the age at first

reproduction (AFR) (63.8% of the species had an AFR ≤ 1 year old, 17%]1; 2], 9.6%]2; 3] and 9.6% >

3 years old, Fig. S3). Therefore, we re-ran the PCA and all the following analyses on a subset of

species for which AFR was available (N=188, Fig. S4). PCA output remained the same, and AFR

loaded positively on the life-history pace PC (Table S1). The linear mixed-effects models gave

qualitatively similar output (Tables S2, S3, S4) suggesting that in our set of species it is unlikely that

AFR affected our analyses, and thus we present in the manuscript the analyses including all species

(N=204).

Correlates of juvenile survival and longevity. We ran two phylogenetically controlled linear mixed-

effects models including the same life-history, ecological and social predictors to assess the factors

correlating with juvenile survival and with longevity. We fitted in both cases the full models (i.e., no

model selection applied) to obtain comparable estimates of the same set of predictors in both

models. To compare the influence of each predictor on both response variables, juvenile survival and

longevity were standardised[90].

Combinations of juvenile survival and longevity. The second set of analyses assessed factors that

were associated with combinations of juvenile survival and longevity that (i) concurred with (positive

JS-L combinations) or (ii) deviated from (JS-L mismatches) the positive correlation between juvenile

survival and longevity, predicted by tETA. We captured the natural patterns of association between

juvenile survival rate and maximum longevity using a PCA approach on the two log-transformed and

standardised survival variables. The PCA resulted in two principal components (PCs, Table S5). Due

to the properties of a PC data rotation, PC1 was loaded positively by both survival estimates (Table

S5). Thus, it describes a tied link between juvenile survival and longevity, capturing patterns that

concur with the classical prediction of tETA (cases positioned on PC1 represent the most typical

cases of positive JS-L combinations). PC2 was loaded positively by juvenile survival rate and

negatively by maximum longevity (Table S5). Being perpendicular to PC1, it captures how much a

species deviates from the overall expected association, and thus, how much it deviates from the

classical prediction of tETA (JS-L mismatches).

We ran two separate phylogenetically controlled linear mixed-effects models to assess the

factor associated with absolute values of (i) PC1 (positive JS-L combinations) and (ii) PC2 (JS-L

mismatches). We included the same set of predictors and covariates as in the full models of juvenile

survival and longevity analyses, and included the sign (positive or negative) of the corresponding PC

as a factor and in interaction with each predictor. The latter allowed us to assess the correlates of

each possible combination of juvenile survival and longevity, i.e., to investigate how species

attributes associated with (i) high juvenile survival-high longevity vs. low juvenile survival-low

longevity combinations (positive JS-L combinations, analysis of PC1), and (ii) deviation towards

higher juvenile survival-lower longevity vs. lower juvenile survival-higher longevity (JS-L mismatches,

analysis of PC2). For both models, we used a backward model selection process. We successively

removed terms with p > 0.10, starting with the highest-order interactions and following with the

simple effects. We compared models including and excluding the focal predictor using *model.sel*

function from the MuMIn package[92]. The decision to exclude the predictor was based on the AICc

criterion using a ΔAICc (i.e., $AICc_{included} - AICc_{excluded}$) > 2 as threshold[93]. Results of the full models are

provided in Table S6 and S7.

		juvenile survival		longevity	
		(first-year survival rate)		(maximum longevity)	
		estimates*	F_{S300}	estimates*	F_{S300}
(intercept)		0.20	0.02	**-0.70**	**1**
residual adult body mass (covariate)		-0.02	0	0.03	0
ln (research effort) (covariate)		**-0.14**	**1**	**0.18**	**1**
life-history pace PC		**0.50**	**1**	**0.42**	**1**
nest predation risk PC		**0.19**	**0.92**	-0.10	0
exposure to predators PC		-0.17	0	**-0.22**	**0.88**
N avian predators PC		0.07	0	-0.05	0
MGS duration PC		-0.01	0	-0.08	0
foraging cost PC		-0.14	0	-0.06	0
food fibre PC		-0.13	0	0.03	0
diet specialisation	generalist	0.00	0	0.00	0
	specialist	0.04		-0.07	
habitat specialisation	generalist	0.00	0	0.00	0
	specialist	0.10		-0.11	
period of activity	diurnal	0.00	0	0.00	0
	nocturnal	0.31		-0.38	
sedentariness	resident	0.00	0	0.00	0
	migratory	0.14		0.07	
region	both	0.00	0	0.00	0
	island	-0.32		-0.12	
	northern	0.39		-0.16	
	southern	0.35		-0.21	
chick development mode	non-precocial	0.00	0	0.00	0
	precocial	-0.10		-0.36	
parental care mode	biparental	0.00	0	0.00	0
	cooperation	-0.08		-0.17	
	uniparental	0.53		0.34	
social system	family-living	0.00	0	0.00	0
	non family-living	-0.23		-0.25	

Bold estimates correspond to predictors with significant effect.
PC: principal component from Table 2.
F_{S300}: frequency of trees for which p-values < 0.05.
*: reference level of categorical variables have an estimate of 0; estimates reflect difference in slope
between the reference level and focal level.
Phylogenetic effect longevity model: likelihood ratio test: LRT = 29.52, df = 1, p < 0.001
Phylogenetic effect juvenile survival model: likelihood ratio test: LRT = 3.33, df = 1, p = 0.0

Results

Correlates of juvenile survival and longevity. Juvenile survival rate ranged from 0.08 to 0.95 (0.39

± 0.16; mean ± SD) and maximum longevity ranged from 5 to 51 (17.7 ± 9.0) years. Juvenile survival

and longevity both correlated with the life-history pace PC, where species with a slow life-history

pace (large body size, low annual reproductive investment, long incubation period; Table 1) had

significantly higher juvenile survival and greater longevity compared to species with a fast life-history

pace (small body size, high annual reproductive investment, short incubation period; Tables 1 and 3).

Moreover, juvenile survival was higher in species with a high nest predation risk (open nest close to

the ground or on the ground; Tables 1 and 3), while longevity was greater in species with a low

exposure of adults to predators (pelagic forager, living in open habitat; Tables 1 and 3). The

phylogenetic effect was only significant for longevity (Table 3).

Figure 1 | Correlation between juvenile survival (first-year survival) and maximum longevity on 293 bird species.
RMA slope = 53.15, 95% CI (34.13, 81.71); $r_{Spearman}$ = 0.28, S = 3003600, p <0.0001. 64 species (22%) are inside and 229 (78%) outside the 95%CI of the regression line (shaded area). See Fig. S6 for species identification.

Combinations of juvenile survival and longevity. Juvenile survival and longevity were positively

correlated ($r_{Spearman}$= 0.28, p <0.0001) (Figs. 1 and S5) and the slope of their linear regression was

significant (N = 293, RMA slope = 53.15, 95% CI of the slope: 34.13, 81.71, p < 0.0001). However,

there were major deviations from the regression line (R^2 = 0.07), and 229 of 293 species (78%) fell

outside the 95% confidence interval (CI) of the RMA regression (Figs. 1 and S6). We note that the

percentage of species that deviate from the overall juvenile survival-longevity relationship was only

a slightly lower (71%) when using a more conservative CI (99%CI: Fig. S7).

Positive associations between juvenile survival and longevity. In general, positive JS-L

combinations were associated with family living, or a high risk of nest predation (open nest on or

close to the ground, Table 1) but those effects were independent of the direction of the relationship

(significant simple effects: Table S8). Opposite positive combinations of juvenile survival and

longevity (low-low vs. high-high) were differently associated with specific-species attributes. This

was reflected by the significant two-way interactions between the sign of PC1 (negative: low-low vs.

positive: high-high JS-L combinations, Fig. 2) and sedentariness, exposure to predators, life-history

pace and parental care mode. Species with high juvenile survival-high longevity combinations were

migratory, had a low exposure to adult predators (pelagic forager, living in open habitat; Table 1), a

slow life-history pace or uniparental care. In contrast, species with low juvenile survival-low

longevity combinations were sedentary, had high exposure to predators, a fast life-history pace, or

had bi-parental or cooperative offspring care) (Figs. 2 and S8, Table S8).

Mismatches between juvenile survival and longevity. In general, JS-L mismatches were associated

with low exposure to adult predators (pelagic foraging, living in open habitat, Table 1) or being a

habitat generalist, but these effects were independent of the direction of the relationship between

juvenile survival and longevity (significant simple effects: Table S9). Opposite JS-L mismatches (low-

high vs. high-low) were differently associated with specific-species attributes as reflected by the

significant two-way interactions between the sign of PC2 (negative: low-high vs. positive: high-low

JS-L combinations, Fig. 2) and period of activity, MGS duration and life-history pace. Species with

stronger than expected combinations of high juvenile survival-low longevity lived in stable

environments with long growing seasons (Table 1), or were nocturnal. In contrast, species with

outstandingly low juvenile survival-high longevity combinations lived in variable environment with

short growing seasons (Table 1) or had a slow life-history pace (Figs. 2 and S9, Table S9).

Figure 2 | Correlates of the positive (PC1) and mismatching (PC2) combinations of juvenile survival
and longevity. Graphical summary of the main results from the backward model selections on
phylogenetically controlled linear mixed models investigating which life-history, ecological and social
traits characterised species with different combinations of juvenile survival (first-year survival) and
longevity (N=204).
The blue axis (PC1) represents combinations that concur with tETA's classical prediction (high juvenile
survival associated with high longevity or vice versa). The green axis (PC2) represents combinations
that deviate from tETA's classical prediction (deviation towards higher juvenile survival associated
with lower longevity or vice versa). Graphics of each independent results are provided in Figs. S8 and
S9. See Fig. S10 for species identification and Fig. S11 for order identification.
JS = juvenile survival, Coop. = cooperative breeding, bi. = biparental care, uni. = uniparental care, noct.
= nocturnal, diurn. = diurnal, F(…,…) = Conditional F statistic and its degrees of freedoms averaged
over the 300 models, p = averaged p value over the 300 models.

Discussion

Empirical studies often use longevity as a proxy of life-history pace, based on the assumption of tETA

that juvenile survival and longevity are positively correlated[2,8,12,94]. While this pattern is supported by

previous work[2,6,7,12] and is generally visible in our data, our analyses show that around 70% of bird

species significantly deviate from this overall juvenile survival-longevity positive relationship (Figs. 1

and S6). Our analyses demonstrate that a wide range of survivals' combinations evolved, some in

accordance with the classical prediction of tETA while others contrasting it, partly supporting recent

developments in this field[9,14-16]. Overall, this study raises awareness on the fact that the relationship

between juvenile survival and longevity is not a black or white concept, but a range of grey nuances,

and identifies key evolutionary forces driving the coevolution between juvenile survival and

longevity.

Correlates of juvenile survival and longevity. On average, a slow life-history pace (in our study

corresponding to: large body size, low annual reproductive investment, long incubation period,

Table 2) is associated with high juvenile survival and longevity (Table 3), supporting life-history

theory[12,19]. However, while juvenile survival and longevity are positively correlated (Figs. 1 and S5),

their individual variation are also associated with particular parameters (Table 3; [95]), supporting

findings from mammals[21]. Our analyses show that nest predation risk (index based on nest location

and nest type, Table 1) only influences juvenile survival while exposure to predators of adults (index

of habitat openness, Table 1 and 2) only influences longevity (Table 3). Consequently, these factors

are likely to play an important role in the evolution of diverse juvenile survival-longevity patterns.

Juveniles are often less conspicuous than adults due to more cryptic coloration and

behaviours[96-99], reducing their vulnerability to predation. Accordingly, a high exposure to predators

of adults is associated with decreased longevity only (Table 3). In contrast, a low nest predation risk

is associated with low juvenile survival only (Table 3). In this study, this latter association concerns

mainly cavity-breeding species (Table 2) known to often experience a lower nest predation risk than

open-nesting species[72]. However, in cavities, nestlings are often exposed to ectoparasites[100,101],

reducing their body condition[100,102,103], potentially explaining a reduced juvenile survival in these

species[95] (Table 3). Therefore, nesting habits that provide short-term benefits early on in life may

have negative down-streams effect on juvenile survival that so far were not anticipated (but see[42]).

Combinations of juvenile survival and longevity. Most species (78%) deviate significantly from the

positive juvenile survival-longevity regression revealing the existence of a continuum of patterns

(Figs. 1 and S6), challenging the classical assumption of tETA[2,6,7,12]. The degree of this deviation varies

considerably between species (Figs. 2 and S10), demonstrating that the association between juvenile

survival and longevity evolved towards multiple adaptive combinations in birds. Some part of this

mismatch may represent random variation and cannot be explained by consistent biological

patterns. However, variation in survival at different life stages is likely to represent distinct

strategies, shaped by natural selection to achieve the most optimal solutions in a given combination

of external and internal factors. Thus, instead of forcing the long-accepted pattern of tETA or

challenging it with opposing hypotheses, we should adopt a more diverse approach. Accordingly,

one should embrace that various possible juvenile survival-longevity combinations exist (including

the non-tETA compliant ones), and their actual values should be assumed to maximize population

viability. Our framework integrating ecological, life-history and social moderators clearly

demonstrates that such a heterogeneous picture is biologically more realistic.

Our analyses on the associations between juvenile survival and longevity do not allow us to

investigate unusual juvenile survival and longevity separately, limiting our ability to identify

underlying mechanisms. This would require an in-depth view of what is happening between

individuals, calling for more comparative studies and experiments on both juvenile survival and

longevity at the intra-species level. However, species-level deviations from the positive correlation

between juvenile survival and longevity likely reflect that certain selective factors only influence

specific life stages[35,104]. Patterns observed between different taxa can be thought of as averaged

outcomes of selective pressures, acting over long periods of time. Indeed, age-dependent changes in

body size, coloration, behaviour, or the onset of reproduction and senescence, can affect extrinsic

and intrinsic mortality differently at different life stages[35,104]. For example, juveniles early on in life

are often smaller than adults, making them more susceptible to predation[32,39]. Also, juvenile survival

may be low in species that live in challenging environments, have elaborate foraging techniques or a

specialised diet, as juveniles in those species seem to need more time to acquire adult skill

levels[62,68,105]. In contrast, only adults pay costs of reproduction, which may reduce their longevity

directly, or indirectly, for instance through increased exposure to predators as a consequence of

increased foraging effort[106], or displaying the own quality to potential partners[107].

Positive associations between juvenile survival and longevity. Positive JS-L combinations are in

accordance with the classical prediction of tETA, indicating that life-history, ecological, and social

parameters have similar effects on juvenile survival and longevity. Our analyses show that high

juvenile survival-high longevity combinations are found in species that are migratory, have a low

exposure to predators, a slow life-history pace or uniparental care (Fig. 2), and are mostly observed

in Accipitriformes, Anseriformes, Charadriiformes, and Pelicaniformes (Fig. S11). In contrast, low

juvenile survival-low longevity combinations are found in species that are sedentary, have a high

exposure to predators, a fast life-history pace, or have cooperative or biparental brood care (Fig. 2),

and are mostly observed in Galliformes and Passeriformes (Fig. S11).

Migration is regularly found in species breeding at higher latitudes or altitudes, allowing

them to escape harsh winter conditions[55]. In most of these species, juveniles and adults are

migratory, thus affecting both life stages. While previous research showed that migration can be

costly (i.e., being associated with smaller relative brain sizes;[108]), our results highlight that it has a

positive effect on survival in general. Moreover, a low exposure to predators is beneficial for both

juvenile and adults, making pelagic species particularly long-lived[32]. As predicted by life-history

theory, species with a slow life-history pace have increased juvenile survival and longevity[12,19,66].

Furthermore, parental care is costly[54,109]. To ensure the survival of their offspring, parents provide

them with food, thermoregulation, and protection from predators, which, on top of being energy

demanding, exposes the parents to an increased risk of predation[54,110]. Thus, it seems surprising that

species with uniparental care have combination of higher juvenile survival and longevity compared

to biparental and cooperatively breeding species. A possible explanation is that particularly species

with low costs of parental care evolved uniparental care, leading to increased juvenile survival and

longevity. Clearly, this finding calls for further studies to investigate both the drivers and

consequences of uniparental care.

Mismatches between juvenile survival and longevity. Mismatching combinations of juvenile

survival and longevity suggest that certain factors specifically act upon juvenile survival or longevity,

or have opposing effects on juvenile survival and longevity, leading to age-specific differences in

survival. Our results demonstrate that high juvenile survival-low longevity combinations are found in

species that live in stable environments with long growing seasons or are nocturnal (Fig. 2), and are

mostly observed in Apodiformes and Galliformes (Fig. S11). In contrast, low juvenile survival-high

longevity combinations are found in species that live in variable environments with short growing

seasons or have a slow life-history pace (Fig. 2), and are mostly observed in Pelicaniformes and

Procellariiformes (Fig. S11).

Conceivably, living in stable environments may particularly affect juvenile survival, reducing

their winter mortality, while the opposite is the case in variable environments. The high juvenile

survival-low longevity combinations found in nocturnal or crepuscular species is likely to reflect

reduced juvenile mortality, given that most predators of birds are diurnal bird species[32]. In contrast,

combinations of low juvenile survival-high longevity found in species with a slow life-history pace is

likely to reflect that long-lived species particularly invest in longevity, at the expense of high juvenile

survival in some species. Generally, interpreting those interactions is not straightforward. We urge

further studies, especially longitudinal ones, to improve our understanding of the interesting

interspecific patterns revealed here.

Conclusions. Our comparative study provides novel insights into interspecific variation in juvenile

survival, longevity and their combination in birds, and highlights the importance to consider age-

specific survival to understand the evolution of life-history traits[22,25,26,42,111]. It increases our

knowledge on the correlates of longevity and the under-studied juvenile survival and shows that

most species deviate from the classical prediction of tETA. Our findings show that multiple adaptive

combinations of juvenile survival and longevity evolved (more than commonly expected), some in

accordance with tETA's classical prediction while others contradict it. Accordingly, we call for a

novel, more diverse, approach to understand the link between juvenile survival and longevity, and to

move beyond the classical prediction of tETA. Our analyses demonstrate that positive JS-L

combinations co-vary along the pace of life continuum, and JS-L mismatches co-vary with the length

of the growing season, where long growing seasons promote juvenile survival, while short growing

seasons promote longevity. Interestingly, sociality (parental care) only explains positive JS-L

combinations, while ecological and life-history traits explain both positive JS-L combinations

(sedentariness, exposure to predators, pace of life) and JS-L mismatches (length of growing season,

period of activity, pace of life). Finally, our analysis emphasizes the need of not only studying typical

patterns, predicted by accepted hypotheses – but also looking at outlying cases, that may embody

genuine biological patterns rather than random deviations from assumed relationships.

Overall, this study reveals that the various combinations of juvenile survival and longevity

observed are shaped by a distinct and limited set of species-specific life-history, ecological and social

attributes. This may reflect divergent selection on each survival estimate, or that divergent age-

specific survival is at the origin of diversity in species attributes[112]. Finally, species with unexpected

age-specific survival relationships are more likely to evolve uncommon combination of life-history

traits[28]. Thus, insights into key factors associating with unusual age-specific survival (such as the one

found in this study) could contribute to a better understanding of life-history evolution[22,25-28,42,111].

Acknowledgements
We thank Carel van Schaik and Gretchen Wagner for discussions and comments on the manuscript.
Kate Mears and Katharine Bowgen for the help with data compilation. This study was financed by
the Swiss National Research Foundation (grant number PPOOP3_123520 and PPOOP3_150752, to
MG), the Polish National Science Centre (Sonata Bis program, grant agreement no UMO-
2015/18/E/NZ8/00505, to SMD) and the Australian Research Council (DECRA Fellowship number
DE180100202).

Authorship:
407 M.G. and E.M. compiled part of the data, E.M. performed all statistical analyses and wrote the first
408 draft of the manuscript. M.G. and S.M.D contributed suggestions and text to subsequent drafts. S.M.D
409 helped with the statistical methods. All authors contributed to revisions and gave final approval for
410 publication.

Competing Interests statement:
413 The research was conducted in the absence of any commercial or financial relationships that could be
414 construed as a potential conflict of interest.

Data accessibility:
417 The datasets supporting this article have been uploaded as part of the Supporting Information and
418 will be archived in Dryad. The data DOI will be included at the end of the article.

Figure Legend:
421 **Figure 1:** Correlation between juvenile survival (first-year survival) and maximum longevity on 293
422 species.
423 **Figure 2:** Graphical summary of the key results from the backward model selections on phylogenetically
424 controlled linear mixed models investigating which life-history, ecological and social traits characterised
425 species with different combinations of juvenile survival (first-year survival) and longevity.

Table Legend:
428 **Table 1.** Description and prediction of the parameters investigated in this study.
429 **Table 2.** Results of the Principal Component Analysis (PCA) with varimax rotation on the 12 continuous
430 predictors.
431 **Table 3.** Results from phylogenetically controlled linear mixed-effect models testing the influence of key
432 life-history, ecological and social traits on juvenile survival and longevity, respectively.

REFERENCES

Medawar, P. B. *An Unsolved Problem of Biology*. (Lewis, 1952).

Williams, G. C. Pleiotropy, natural selection, and the evolution of senescence. *Evolution* **11**, 398-411, doi:10.2307/2406060 (1957).

Kirkwood, T. B. L. Evolution of ageing. *Nature* **270**, 301-304 (1977).

Kirkwood, T. B. L. & Austad, S. N. Why do we age? *Nature* **408**, 233-238, doi:10.1038/35041682 (2000).

Abrams, P. A. Does Increased mortality favor the evolution of more rapid senescence? *Evolution* **47**, 877-887, doi:10.2307/2410191 (1993).

Kirkwood, T. B. L. Evolution of ageing. *Mech Ageing Dev* **123**, 737-745, doi:10.1016/s0047-6374(01)00419-5 (2002).

Ricklefs, R. E. Evolutionary theories of aging: Confirmation of a fundamental prediction, with implications for the genetic basis and evolution of life span. *Am. Nat.* **152**, 24-44, doi:10.1086/286147 (1998).

Ricklefs, R. E. & Scheuerlein, A. in *Lifespan: Evolutionary, Ecological and Demogrpahic Perspectives* Vol. A supplement to Population and Development Review, volume 29 (eds J. R. Carey & S. Tuljapurkar) 71-98 (Population Council, 2003).

Shokhirev, M. N. & Johnson, A. A. Effects of Extrinsic Mortality on the Evolution of Aging: A Stochastic Modeling Approach. *PLoS ONE* **9**, doi:10.1371/journal.pone.0086602 (2014).

Williams, P. D., Day, T., Fletcher, Q. & Rowe, L. The shaping of senescence in the wild. *Trends Ecol. Evol.* **21**, 458-463, doi:10.1016/j.tree.2006.05.008 (2006).

Caswell, H. Extrinsic mortality and the evolution of senescence. *Trends Ecol. Evol.* **22**, 173-174, doi:10.1016/j.tree.2007.01.006 (2007).

Charlesworth, B. *Evolution in Age Structured Populations*. (Cambridge University Press, 1994).

Williams, P. D. & Day, T. Antagonistic pleiotropy, mortality source interactions, and the evolutionary theory of senescence. *Evolution* **57**, 1478-1488, 1411 (2003).

Chen, H. & Maklakov, Alexei A. Longer Life Span Evolves under High Rates of Condition-Dependent Mortality. *Curr. Biol.* **22**, 2140-2143, doi:http://dx.doi.org/10.1016/j.cub.2012.09.021 (2012).

Dowling, D. K. Aging: Evolution of Life Span Revisited. *Curr. Biol.* **22**, R947-R949, doi:https://doi.org/10.1016/j.cub.2012.09.029 (2012).

Werfel, J., Ingber, D. E. & Bar-Yam, Y. Theory and associated phenomenology for intrinsic mortality arising from natural selection. *PLoS ONE* **12**, e0173677, doi:10.1371/journal.pone.0173677 (2017).

Reznick, D. N., Bryant, M. J., Roff, D., Ghalambor, C. K. & Ghalambor, D. E. Effect of extrinsic mortality on the evolution of senescence in guppies. *Nature* **431**, 1095-1099, doi:http://www.nature.com/nature/journal/v431/n7012/suppinfo/nature02936_S1.html (2004).

Roff, D. A. *The Evolution of Life Histories*. (Chapman and Hall, 1992).

Stearns, S. C. *The Evolution of Life Histories*. (Oxford University Press, 1992).

Martin, T. E. Age-related mortality explains life history strategies of tropical and temperate songbirds. *Science* **349**, 966-970, doi:10.1126/science.aad1173 (2015).

Promislow, D. E. L. & Harvey, P. H. Living fast and dying young - A comparative analysis of the life-history variation among mammals. *J. Zool.* **220**, 417-437 (1990).

Reznick, D. A., Bryga, H. & Endler, J. A. Experimentally induced life-history evolution in a natural population. *Nature* **346**, 357-359, doi:10.1038/346357a0 (1990).

Cole, L. C. The population consequences of life history phenomena. *Q. Rev. Biol.* **29**, 103-137, doi:10.1086/400074 (1954).

Williams, G. C. Natural selection, costs of reproduction, and a refinement of Lack's principle. *Am. Nat.* **100**, 687-690, doi:10.1086/282461 (1966).

Promislow, D. E. L. & Harvey, P. H. Mortality rates and the evolution of mammal life histories. *Acta Oecologica-International Journal of Ecology* **12**, 119-137 (1991).

Michod, R. E. Evolution of life histories in response to age-specific mortality factors. *Am. Nat.* **113**, 531-550, doi:10.1086/283411 (1979).

Martin, T. E. A new view of avian life-history evolution tested on an incubation paradox. *Proc. R. Soc. B* **269**, 309-316, doi:10.1098/rspb.2001.1879 (2002).

Kraus, C., Thomson, D. L., Kunkele, J. & Trillmich, F. Living slow and dying young? Life-history strategy and age-specific survival rates in a precocial small mammal. *J. Anim. Ecol.* **74**, 171-180, doi:10.1111/j.1365-2656.2004.00910.x (2005).

Briggs-Gonzalez, V. *et al.* Life histories and conservation of long-lived reptiles, an illustration with the American crocodile (Crocodylus acutus). *J. Anim. Ecol.* **86**, 1102-1113, doi:doi:10.1111/1365-2656.12723 (2017).

Jones, O. R. *et al.* Diversity of ageing across the tree of life. *Nature* **505**, 169, doi:10.1038/nature12789

https://www.nature.com/articles/nature12789#supplementary-information (2013).

Healy, K. *et al.* Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc. R. Soc. B* **281**, doi:10.1098/rspb.2014.0298 (2014).

Valcu, M., Dale, J., Griesser, M., Nakagawa, S. & Kempenaers, B. Global gradients of avian longevity support the classic evolutionary theory of ageing. *Ecography* **37**, 930-938, doi:10.1111/ecog.00929 (2014).

de Magalhaes, J. P., Costa, J. & Church, G. M. An analysis of the relationship between metabolism, developmental schedules, and longevity using phylogenetic independent contrasts. *Journals of Gerontology Series a-Biological Sciences and Medical Sciences* **62**, 149-160 (2007).

Wasser, D. E. & Sherman, P. W. Avian longevities and their interpretation under evolutionary theories of senescence. *J. Zool.* **280**, 103-155, doi:10.1111/j.1469-7998.2009.00671.x (2010).

Sullivan, K. A. Predation and Starvation: Age-Specific Mortality in Juvenile Juncos (Junco phaenotus). *Journal of Animal Ecology* **58**, 275-286 (1989).

Caughley, G. Mortality Patterns in Mammals. *Ecology* **47**, 906-918 (1966).

Charnov, E. L. Life History Evolution in a "Recruitment Population": Why Are Adult Mortality Rates Constant? *Oikos* **47**, 129-134, doi:10.2307/3566037 (1986).

Gaillard, J. M., Festa-Bianchet, M., Yoccoz, N. G., Loison, A. & Toigo, C. Temporal variation in fitness components and population dynamics of large herbivores. *Annu. Rev. Ecol. Syst.* **31**, 367-393, doi:10.1146/annurev.ecolsys.31.1.367 (2000).

Caro, T. *Antipredator Defenses in Birds and Mammals*. (University of Chicago Press, 2005).

Werner, E. E. & Hall, D. J. Ontogenetic habitat shifts in bluegill: the foraging rate-predation risk trade-off. *Ecology* **69**, 1352-1366 (1988).

Griesser, M. *et al.* Experience buffers extrinsic mortality in a group-living bird species. *Oikos* **126**, 1258-1268, doi:10.1111/oik.04098 (2017).

Martin, T. E. A Conceptual Framework for Clutch-Size Evolution in Songbirds. *Am. Nat.* **183**, 313-324, doi:10.1086/674966 (2014).

Ricklefs, R. E. Lack, Skutch, and Moreau: The early development of life-history thinking. *Condor* **102**, 3-8 (2000).

Skutch, A. F. Clutch size, nesting success, and predation on nests of Neotropical birds, reviewed. *Ornithol Monogr*, 575-594 (1985).

Grüebler, M. U. & Naef-Daenzer, B. Survival benefits of post-fledging care: experimental approach to a critical part of avian reproductive strategies. *J. Anim. Ecol.* **79**, 334-341, doi:10.1111/j.1365-2656.2009.01650.x (2010).

Drobniak, M. S., Wagner, G., Mourocq, E. & Griesser, M. Family living: an overlooked but pivotal social system to understand the evolution of cooperative breeding. *Behav. Ecol.* **26**, 805-811, doi:10.1093/beheco/arv015 (2015).

Griesser, M., Nystrand, M. & Ekman, J. Reduced mortality selects for family cohesion in a social species. *Proc. R. Soc. B* **273**, 1881-1886 (2006).

Robinson, W. D. *et al.* Diversification of Life Histories in New World Birds. *Auk* **127**, 253-262 (2010).

Møller, A. P. Sociality, age at first reproduction and senescence: comparative analysis of birds. *Journal for Evolutionary Biology* **19**, 682-689 (2006).

Carey, J. R. & Tuljapurkar, S. *Life Span: Evolutionary, Ecological, and Demographic Perspectives*. Vol. Population and development review. A Supplement to Volume 29, 2003 (Population council, 2003).

Maness, T. J. & Anderson, D. J. in *Predictors of Juvenile Survival in Birds   Ornithological Monographs*   1-55 (2013).

Cox, W. A., Thompson, F. R., Cox, A. S. & Faaborg, J. Post-fledging survival in passerine birds and the value of post-fledging studies to conservation. *The Journal of Wildlife Management* **78**, 183-193, doi:10.1002/jwmg.670 (2014).

Russell, E. M. Avian life histories: Is extended parental care the southern secret? *Emu* **100**, 377-399, doi:10.1071/mu0005s (2000).

Alonso-Alvarez, C. & Velando, A. in *The Evolution of parental care*   (eds N. J. Royle, P. T. Smiseth, & M. Kölliker) 40-61 (Oxford University Press, 2012).

Cox, G. W. The Evolution of Avian Migration Systems between Temperate and Tropical Regions of the New World. *The American Naturalist* **126**, 451-474, doi:10.1086/284432 (1985).

Sibly, R. M. *et al.* Energetics, lifestyle, and reproduction in birds. *Proc. Natl. Acad. Sci. U.S.A* **109**, 10937-10941, doi:10.1073/pnas.1206512109 (2012).

Whittingham, M. J. & Evans, K. L. The effects of habitat structure on predation risk of birds in agricultural landscapes. *Ibis* **146**, 210-220, doi:doi:10.1111/j.1474-919X.2004.00370.x (2004).

Lazarus, J. & Symonds, M. Contrasting effects of protective and obstructive cover on avian vigilance. *Anim. Behav.* **43**, 519-521, doi:https://doi.org/10.1016/S0003-3472(05)80110-1 (1992).

Lima, S. L. Protective Cover and the Use of Space: Different Strategies in Finches. *Oikos* **58**, 151-158, doi:10.2307/3545422 (1990).

Isler, K. & van Schaik, C. Costs of encephalization: the energy trade-off hypothesis tested on birds. *Journal of Human Evolution* **51**, 228-243 (2006).

Langer, P. Lactation, weaning period, food quality, and digestive tract differentiations in eutheria. *Evolution* **57**, 1196-1215 (2003).

Bautista, L. M., Tinbergen, J. & Kacelnik, A. To walk or to fly? How birds choose among foraging modes. *Proc. Natl. Acad. Sci. U.S.A* **98**, 1089-1094, doi:10.1073/pnas.98.3.1089 (2001).

Colles, A., Liow, L. H. & Prinzing, A. Are specialists at risk under environmental change? Neoecological, paleoecological and phylogenetic approaches. *Ecol. Lett.* **12**, 849-863, doi:doi:10.1111/j.1461-0248.2009.01336.x (2009).

Julliard, R., Jiguet, F. & Couvet, D. Common birds facing global changes: what makes a species at risk? *Global Change Biol.* **10**, 148-154, doi:doi:10.1111/j.1365-2486.2003.00723.x (2004).

Wilson, S. K. *et al.* Habitat utilization by coral reef fish: implications for specialists vs. generalists in a changing environment. *J. Anim. Ecol.* **77**, 220-228, doi:doi:10.1111/j.1365-2656.2007.01341.x (2008).

McNamara, J. M., Barta, Z., Wikelski, M. & Houston, A. I. A Theoretical Investigation of the Effect of Latitude on Avian Life Histories. *The American Naturalist* **172**, 331-345 (2008).

Blondel, J. *Evolution and ecology of birds on islands: Trends and prospects*. Vol. 50 (2000).

Heinsohn, R. G. Slow Learning of Foraging Skills and Extended Parental Care in Cooperatively Breeding White-Winged Choughs. *The American Naturalist* **137**, 864-881, doi:10.2307/2462405 (1991).

Blumstein, D. T. & Møller, A. P. Is sociality associated with high longevity in North American birds? *Biol. Lett.* **4**, 146-148, doi:10.1098/rsbl.2007.0606 (2008).

Van de Loock, D. *et al.* Cooperative breeding shapes post-fledging survival in an Afrotropical forest bird. *Ecology and Evolution*, n/a-n/a, doi:10.1002/ece3.2744 (2017).

Tarwater, C. E. & Brawn, J. D. The post-fledging period in a tropical bird: patterns of parental care and survival. *J. Avian Biol.* **41**, 479-487 (2010).

Martin, T. E. & Li, P. J. Life-history traits of open-nesting vs cavity-nesting birds. *Ecology* **73**, 579-592, doi:10.2307/1940764 (1992).

Martin, T. E. Nest predation and nest sites - New perspectives on old patterns. *Bioscience* **43**, 523-532, doi:10.2307/1311947 (1993).

Schmidtn.K. Locomotion: energy cost of swimming, flying, and running. *Science* **177**, 222-&, doi:10.1126/science.177.4045.222 (1972).

IUCN. IUCN Habitat Classification Scheme. (2007).

Botero, C. A. & Rubenstein, D. R. Fluctuating Environments, Sexual Selection and the Evolution of Flexible Mate Choice in Birds. *PLoS ONE* **7**, doi:10.1371/journal.pone.0032311 (2012).

Del Hoyo, J., Elliot, A., Sargatal, J. & Christie, D. A. *Handbook of the Birds of the World. Available via* http://www.hbw.com. (Lynx, 2011).

Poole, A. The birds of North America online. *Cornell Laboratory of Ornithology, Ithaca, NY. Available via* http://bna.birds.cornell.edu/BNA (2005).

Higgins, P. J. *et al. Handbook of Australian, New Zealand & Antarctic Birds*. (Oxford University Press, 1996-2006).

Maclean, G. L. & Robert, A. *Roberts' birds of southern Africa*. (Trustees of the John Voelcker Bird Book Fund Cape Town, 1985).

Commonwealth of Australia. *Australian Bird and Bat Banding Scheme Database, queried (May 2014). Available via* http://www.environment.gov.au/science/bird-and-bat-banding. (Department of the Environment, 2014).

de Magalhaes, J. P. & Costa, J. A database of vertebrate longevity records and their relation to other life-history traits. *J. Evol. Biol.* **22**, 1770-1774, doi:10.1111/j.1420-9101.2009.01783.x (2009).

Sankamethawee, W., Gale, G. A. & Hardesty, B. D. Post-Fledgling Survival of the Cooperatively Breeding Puff-Throated Bulbul (Alophoixus Pallidus). *The Condor* **111**, 675-683, 679 (2009).

Griesser, M., Drobniak, S. M., Nakagawa, S. & Botero, C. A. Family living sets the stage for cooperative breeding and ecological resilience in birds. *PLoS Biol* **15**, e2000483 (2017).

R: A language and environment for statistical computing. (R Foundation for Statistical Computing, Vienna, Austria. , 2015).

asreml: asreml() fits the linear mixed model. R package version 3.0. www.vsni.co.uk (2009).

Hadfield, J. D. & Nakagawa, S. General quantitative genetic methods for comparative biology: phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *J. Evol. Biol.* **23**, 494-508, doi:doi:10.1111/j.1420-9101.2009.01915.x (2010).

Huelsenbeck, J. P. & Crandall, K. A. Phylogeny estimation and hypothesis testing using maximum likelihood. *Annu. Rev. Ecol. Syst.* **28**, 437-466, doi:10.1146/annurev.ecolsys.28.1.437 (1997).

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. & Mooers, A. O. The global diversity of birds in space and time. *Nature* **491**, 444-448, doi:10.1038/nature11631 (2012).

Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods Ecol. Evol.* **1**, 103-113, doi:10.1111/j.2041-210X.2010.00012.x (2010).

Dormann, C. F. *et al.* Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* **36**, 27-46, doi:10.1111/j.1600-0587.2012.07348.x (2013).

Barton, K. MuMIn: Multi-model inference. *R package version 2.0.0.* http://R-Forge.R-project.org/projects/mumin/ (2013).

Burnham, K. P. & Anderson, D. R. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behav. Ecol. Sociobiol.* **65**, 23-35, doi:10.1007/s00265-010-1029-6 (2011).

Keller, L. & Genoud, M. Extraordinary lifespans in ants: a test of evolutionary theories of ageing. *Nature* **389**, 958-960, doi:10.1038/40130 (1997).

Francis, C. M., Richards, M. H., Cooke, F. & Rockwell, R. F. Long-term changes in survival rates of Lesser snow geese. *Ecology* **73**, 1346-1362, doi:10.2307/1940681 (1992).

Heinen, J. T. Cryptic behavior in juvenile toads. *J. Herpetol.* **19**, 524-527, doi:10.2307/1564206 (1985).

Ryer, C. H., Lemke, J. L., Boersma, K. & Levas, S. Adaptive coloration, behavior and predation vulnerability in three juvenile north Pacific flatfishes. *J. Exp. Mar. Biol. Ecol.* **359**, 62-66, doi:10.1016/j.jembe.2008.02.017 (2008).

Caro, T. The Adaptive Significance of Coloration in Mammals. *Bioscience* **55**, 125-136, doi:10.1641/0006-3568(2005)055[0125:TASOCI]2.0.CO;2 (2005).

Baker, R. R. & Parker, G. A. The evolution of bird coloration. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* **287**, 63-130, doi:10.1098/rstb.1979.0053 (1979).

Cantarero, A., Lopez-Arrabe, J., Redondo, A. J. & Moreno, J. Behavioural responses to ectoparasites in pied flycatchers Ficedula hypoleuca: an experimental study. *J. Avian Biol.* **44**, 591-599, doi:10.1111/j.1600-048X.2013.00134.x (2013).

Nilsson, S. G. Evolution of hole-nesting in birds: On balancing selection pressures. *Auk* **103**, 432-435 (1986).

Tomas, G. *et al.* Determinants of abundance and effects of blood-sucking flying insects in the nest of a hole-nesting bird. *Oecologia* **156**, 305-312, doi:10.1007/s00442-008-1001-6 (2008).

Brommer, J. E., Pitala, N., Siitari, H., Kluen, E. & Gustafsson, L. Body size and immune defense of nestling blue tits (Cyanistes caeruleus) in response to manipulation of ectoparasites and food supply. *Auk* **128**, 556-563, doi:10.1525/auk.2011.10284 (2011).

Lack, D. *Population Studies of Birds*. (Oxford University Press, 1966).

Schuppli, C., Isler, K. & van Schaik, C. P. How to explain the unusually late age at skill competence among humans. *J. Hum. Evol.* **63**, 843-850, doi:http://dx.doi.org/10.1016/j.jhevol.2012.08.009 (2012).

Zarybnicka, M., Korpimaki, E. & Griesser, M. Dark or Short Nights: Differential Latitudinal Constraints in Nestling Provisioning Patterns of a Nocturnally Hunting Bird Species. *PLoS ONE* **7**, doi:10.1371/journal.pone.0036932 (2012).

da Cunha, F. C. R., Fontenelle, J. C. R. & Griesser, M. The presence of conspecific females influences male-mobbing behavior. *Behavioral Ecology and Sociobiology* **71**, 52, doi:10.1007/s00265-017-2267-7 (2017).

Sol, D. *et al.* Evolutionary Divergence in Brain Size between Migratory and Resident Birds. *PLoS ONE* **5**, e9617, doi:10.1371/journal.pone.0009617 (2010).

Trivers, R. L. in *In: Sexual selection and the descent of man* (ed Aldine) 136-179 (Campbell, 1972).

Santos, E. S. A. & Nakagawa, S. The costs of parental care: a meta-analysis of the trade-off between parental effort and survival in birds. *J. Evol. Biol.* **25**, 1911-1917, doi:10.1111/j.1420-9101.2012.02569.x (2012).

Schaffer, W. M. Selection for optimal life histories: The effects of age structure. *Ecology* **55**, 291-303, doi:10.2307/1935217 (1974).

Healy, K., Ezard, T. H. G., Jones, O. R., Salguero-Gómez, R. & Buckley, Y. M. Animal life history is shaped by the pace of life and the distribution of age-specific mortality and reproduction. *Nature Ecology & Evolution*, doi:10.1038/s41559-019-0938-7 (2019).

Tables:

Table 1| Description and prediction of the parameters investigated in this study.

	parameter's name	description	prediction	source prediction
life-history	adult body mass*	mean adult body mass (g)	larger body size confers better ability to cope with temporary food shortages, climatic fluctuations and extreme weather than smaller body size; large body size may associate with higher juvenile survival and higher longevity	[39,52]
	incubation period*	number of days from laying to hatching	longer incubation period may associate with higher juvenile survival and higher longevity	[12,18]
	nestling period*	number of days from hatching to fledging	longer nestling period may associate with higher juvenile survival and higher longevity	[42]
	annual parental investment*	body-mass scaled annual reproductive investment (total mass of eggs produced annually divided by adult body mass)(a)	higher parental investment may associate with lower juvenile survival and lower longevity	[12,18,53]
	chick development mode	precocial vs. non precocial; semi-altricial or semi precocial species were categorised as non precocial	precocial species should have lower juvenile survival but higher longevity because of lower parental care after hatching while the opposite is expected for altricial species	[54]
	sedentariness	resident vs. migratory; based on the species maximum movement; sedentary species or with local movement were categorised as resident and the one with regional or inter-continental movement as migratory	costs associated with migration could translate into lower juvenile survival and lower longevity in migratory than in non-migratory species	[55]
	period of activity	diurnal vs. nocturnal; crepuscular species (i.e. active at dawn and dusk) were categorised as nocturnal	species that are active at night are likely to be harder for predators to detect and predators are more scarce at night thus, nocturnal species might have higher juvenile survival and live longer than diurnal species	[31]
	nest predation risk*	based on both most commonly used nest location and nest type; ordinally ranked: 1 = inaccessible nests in cavities, 2 = open nests in cliffs or tree, 3 = open nest in shrub-layer or the ground(b)	nest predation risk may alter the developmental phase of the nestling and the reproductive effort of the parents which may affect juvenile survival and longevity; greater nest predation risk may associate with lower juvenile survival and lower longevity	[20]
	foraging exposure*	level of exposure to predators during foraging time based on most commonly used foraging area; ordinally ranked: 1 = pelagic species, 2 = aerial foragers, 3 = terrestrial foragers	pelagic or aerial forager should have lower predation risk and be more capable of escaping from predators than species that feed on the ground; juvenile survival and longevity may be reduced in the latter more than in the formers	[56,57]
	vegetation cover*	cover of woody vegetation in habitat (%)	more open habitats provide less visual cover than habitats dense in vegetation, increasing the risk of being killed; thus, low vegetation density may associate with lower juvenile survival and lower longevity. The reverse may be true if vegetation cover, by obstructing the view of the prey, affects its survival	[57-59]
ecological	caloric content of food*	energy content of the food in kcal/100g(c)	food calory content can influence the energy available for maintenance; high calory diet may associate with higher juvenile survival and higher longevity	[60,61]

	fibre content of food*	fibre food content in g/100g[c]	food fibre content can influence digestion efficiency and thus the level of resource acquired and health; high fibre diet may associate with higher juvenile survival and higher longevity	[60,61]
	foraging cost*	level of energy demand for foraging based on most commonly used foraging technics; ordinally ranked: 1 = sit and wait hunters, 2 = swimming or short perch & short flights, 3 = aerial or under water foraging, 4 = terrestrial or gleaners [d]	species with highly energetically demanding foraging strategies may have lower juvenile survival and lower longevity than species with less energetically demanding technics	[62]
	diet specialisation	specialist (only one diet class) vs. generalist (more than one diet class)	a change in the food availability can have higher costs for specialist than generalist species as the later can deviate to other food resources; specialisation may associate with lower juvenile survival and lower longevity	[63]
	habitat specialisation	specialist (only one habitat type) vs. generalist (more than one habitat type)[e]	a change in habitat availability can have higher costs for specialist than generalist species as the latter can occupy other habitat types; specialisation is predicted to associate with lower juvenile survival and lower longevity	[64,65]
	MGS duration*	mean duration of the growing season in months[f] (i.e., month(s) of the yeuyuar in which temperature and rainfall allow significant plant productivity)	a short growing season implies changes in environmental conditions over the year, thus MGS duration can be seen as a proxy of environemental variability; less variable environments (long growing season) may associate with higher juvenile survival and longevity than highly variable environment (short growing season)	[66]
	region	breeding distribution range: northern or southern hemisphere, both hemispheres, island	southern hemisphere and island species may have higher juvenile survival and higher longevity compared to northern hemisphere species	[34,66,67]
	N avian predators*	number of sympatric adult's or independent juveniles' predator species[g]	a higher number of predators increases the risk of being predated; higher number of predators may associate with a lower juvenile survival and lower longevity	[32]
Social	parental care mode	uniparental, biparental, cooperative breeding	the presence of additional carer can reduce survival risks on young and survival costs on the other carer(s); biparental and cooperative breeding species may have higher juvenile survival and higher longevity than uniparental species	[68],but see[69,70]
	social system	family living (offspring remain at least 50 days beyond nutritional independence with parents) vs. non-family living[h]	species with prolonged post-fledging parental care or having a prolonged association with the parents beyond independence, as in family-living species, may have higher juvenile survival and lower longevity	[45,71]

Because experience (e.g., foraging, competition, reproductive strategies, anti-predation behaviours) differs between young individuals (inexperienced) and adults (experienced), we also assumed each of the abovementioned parameters to diferentially influence juvenile survival and longevity, and potentially explain variation in juvenile survival/longevity relationships.

* Included in the PCA (Table 2). The other parameters are categorical variables. (a)[56], (b)[72,73], (c)[60], (d)[74], (e)[75], (f)[76], (g)[32], (h)[46].

Table 2 | **Results of the Principal Component Analysis (PCA)** with varimax rotation on the 12 continuous predictors.

category	transformation	variable	nº component	life-history pace	exposure to predators	food fibre	nest predation risk	N avian predators	foraging cost	MGS duration	h2
				1	7	2	6	3	5	4	h2
life-history	ln	adult body mass		**0.90**	-0.25	-0.08	0.13	-0.01	-0.17	-0.12	0.95
	none	annual parental investment		**-0.89**	0.14	-0.17	0.13	0.14	-0.05	-0.02	0.87
	sqrt	incubation period		**0.76**	-0.35	0.23	-0.03	-0.10	-0.28	-0.04	0.84
	none	nestling period		0.51	0.04	0.44	-0.58	-0.13	0.01	0.20	0.85
ecological	none	nest predation risk		0.05	-0.10	-0.12	**0.94**	-0.12	0.09	-0.07	0.93
	none	foraging cost		-0.17	0.02	-0.11	0.08	-0.07	**0.96**	-0.05	0.98
	none	calorie content of food		0.15	0.49	0.62	0.02	0.44	0.01	-0.01	0.83
	ln	fibre content of food		-0.11	0.16	**-0.89**	0.22	-0.01	0.15	-0.10	0.92
	none	foraging exposure		-0.30	**0.86**	-0.04	-0.01	0.04	-0.02	0.00	0.83
	sqrt	vegetation cover		-0.21	**0.85**	-0.01	-0.13	-0.08	0.05	0.11	0.80
	sqrt	N avian predators		-0.18	-0.03	0.08	-0.07	**0.95**	-0.05	0.00	0.95
	none	MGS duration		-0.06	0.08	0.08	-0.11	-0.01	-0.05	**0.98**	0.99
		SS loadings		2.68	1.94	1.5	1.34	1.15	1.07	1.05	
		cumulative variance explained (%)		22	39	51	62	72	81	89	

We considered coefficients of correlation greater than 0.7 or less than -0.7 to be high loadings (highlighted in bold). h2 is the communality of the 7 components. ln: natural logarithm, sqrt : Square root

Table 3 | Correlates of juvenile survival and longevity. Results from phylogenetically controlled linear mixed-effect models testing the influence of key life-history, ecological and social traits on juvenile survival and longevity, respectively.

		juvenile survival (first-year survival rate)		longevity (maximum longevity)	
		estimates*	F_{s300}	estimates*	F_{s300}
(intercept)		0.20	0.02	**-0.70**	**1**
residual adult body mass (covariate)		-0.02	0	0.03	0
ln (research effort) (covariate)		**-0.14**	**1**	**0.18**	**1**
life-history pace PC		**0.50**	**1**	**0.42**	**1**
nest predation risk PC		**0.19**	**0.92**	-0.10	0
exposure to predators PC		-0.17	0	**-0.22**	**0.88**
N avian predators PC		0.07	0	-0.05	0
MGS duration PC		-0.01	0	-0.08	0
foraging cost PC		-0.14	0	-0.06	0
food fibre PC		-0.13	0	0.03	0
diet specialisation	generalist	0.00	0	0.00	0
	specialist	0.04		-0.07	
habitat specialisation	generalist	0.00	0	0.00	0
	specialist	0.10		-0.11	
period of activity	diurnal	0.00	0	0.00	0
	nocturnal	0.31		-0.38	
sedentariness	resident	0.00	0	0.00	0
	migratory	0.14		0.07	
region	both	0.00	0	0.00	0
	island	-0.32		-0.12	
	northern	0.39		-0.16	
	southern	0.35		-0.21	
chick development mode	non-precocial	0.00	0	0.00	0
	precocial	-0.10		-0.36	
parental care mode	biparental	0.00	0	0.00	0
	cooperation	-0.08		-0.17	
	uniparental	0.53		0.34	
social system	family-living	0.00	0	0.00	0
	non family-living	-0.23		-0.25	

Bold estimates correspond to predictors with significant effect.

PC: principal component from Table 2.

F_{s300}: frequency of trees for which p-values < 0.05.

*: reference level of categorical variables have an estimate of 0; estimates reflect difference in slope between the reference level and focal level.

Phylogenetic effect longevity model: likelihood ratio test: LRT = 29.52, df = 1, p < 0.001

Phylogenetic effect juvenile survival model: likelihood ratio test: LRT = 3.33, df = 1, p = 0.07

Figures:

Figure 1 | Correlation between juvenile survival (first-year survival) and maximum longevity on 293 bird species.
RMA slope = 53.15, 95% CI (34.13, 81.71); $r_{Spearman}$ = 0.28, S = 3003600, p <0.0001. 64 species (22%) are inside and 229 (78%) outside the 95%CI of the regression line (shaded area). See Fig. S6 for species identification.

Figure 2 | Correlates of the positive (PC1) and mismatching (PC2) combinations of juvenile survival and longevity. Graphical summary of the main results from the backward model selections on phylogenetically controlled linear mixed models investigating which life-history, ecological and social traits characterised species with different combinations of juvenile survival (first-year survival) and longevity (N=204).

The blue axis (PC1) represents combinations that concur with tETA's classical prediction (high juvenile survival associated with high longevity or vice versa). The green axis (PC2) represents combinations that deviate from tETA's classical prediction (deviation towards higher juvenile survival associated with lower longevity or vice versa). Graphics of each independent results are provided in Figs. S8 and S9. See Fig. S10 for species identification and Fig. S11 for order identification.

JS = juvenile survival, Coop. = cooperative breeding, bi. = biparental care, uni. = uniparental care, noct. = nocturnal, diurn. = diurnal, F(…,…) = Conditional F statistic and its degrees of freedoms averaged over the 300 models, p = averaged p value over the 300 models.