Identification of Six novel missense single nucleotide polymorphisms in MAOA gene predisposing to aggressive behavior. Bioinformatics study.

Abdelrahman H. Abdelmoenim[1*], Mujahed I. Mustafa[2], Naseem S. Murshed[3], Nosiba S. Omer[4], Alaa I. Mohammed[5], Abdelrafie M. Makhawi[2].

1-Faculty of medicine, Alneelain university, Khartoum, Sudan.

2-Department of Biotechnology, University of Bahri, Khartoum, Sudan.

3-Department of Bioinformatics, Africa City of Technology, Khartoum North, Sudan.

4-Faculty of nursing, Shandi university, Khartoum, Sudan.

5-Department of Hematology & Immunohematology, Faculty of Medical Laboratory Sciences, University of Khartoum, Khartoum, Sudan.

*Corresponding author email: abduhamza009@gmail.com.

Abstract:

Background: An astonishing observation is that aggressive behavior is actually highly heritable. Recent experimental work and behavior research has linked individual variation in a functional polymorphism of the monoamine oxidase-A gene (MAOA) to the occurrence of anger-driven aggression. Aggressive antisocial and violent behavior has become a regularly debated topic in the scientific community; the impending question is what is the source of aggressive behavior, is it genetic or environmental or is it just an individual choice. This study aim to analyses the SNPs found in MAOA gene and it is possible relation to aggressive behavior.

Method: Various bioinformatics software (soft, polphen2, provean, snap2,snp and go and pmut) is used to analyses the SNPs within MAOA gene to study the structural and functional implication on the associated protein , which is further clarified using chimera software. Then gene- gene interaction is studied with geneMANIA software. Furthermore conservation and annotation studies were done through ConSurf server and Variant Effect Predictor (VEP) respectively.

Result: 6 missense SNPs were found to affect the structural and functional prospect of MAOA protein. Conclusion: Genetic mutation within MAOA is likely to be associated with aggressive behavior; this will enrich future management and screening possibility for this behavior.

Keywords: aggressive behavior, antisocial behavior, MAOA, SNPs, bioinformatics.

Introduction :

Aggressive and antisocial behavior is highly heritable.(1) Recent experimental work and behavior research has linked individual variation in a functional polymorphism of the monoamine oxidase-A gene (MAOA) to the occurrence of anger-driven aggression.(2-5)

 Furthermore There is an extensive research examining the relation between a polymorphism in the promoter region of the MAOA gene and antisocial phenotypes ,as it has been found that 2-repeat allele- may have effects in modulating aggressive behaviors.(6, 7) Historically The clinical condition caused by MAOA deficiency, namely Brunner syndrome, was one of the first disorder to be described in relation to MAOA gene and it is characterized by overt antisocial and aggressive conduct.(8)studies has found that   interaction of low MAOA transcription polymorphisms with environmental factors such as early abuse , deprivation and iron deficiency anemia has been found to increase the occurrence of violent antisocial behavior.(9, 10)

The MAO is a mitochondrial enzyme . MAO family composed of two isoenzymes, termed A (Genbank Acc. No: BC008064.1, Gene ID: 4128, Xp11.3) and B (Genbank Acc. No: M69177, Gene ID: 4129, Xp21p11), which despite a substantial structural overlap, are remarkably different for substrate preference, inhibitor selectivity, regional distribution, and functional role in behavioral regulation.(11) The two monoamine oxidase (MAO) enzymes, A and B, preferentially deaminates monoamine neurotransmitters, such as serotonin, norepinephrine, and dopamine. The genes encoding for both types are mapped on chromosome Xp11.23 .(12)

It has been found that the termination of serotonin (5-hydroxytryptamine, 5-HT) neurotransmission is regulated by its uptake by the 5-HT transporter (5-HTT), as well as its degradation by monoamine oxidase (MAO)-A. MAO-A deficiency and hypo-methylations leads to a wide set of behavioral alterations, including perseverative behaviors ,social deficits and depression.(13-16) in addition to that cognitive functions like problem solving were found to be affected by mutations in this gene when it is combined with iron deficiency (ID) in rhesus monkey.(9, 17)

Previous evidence of gene-by-environment interactions associated with emotional and behavioral disorders is contradictory. Differences in findings may be related to variation in dose of the environmental factor, or failure to take account of gene-by-gene interactions.(18)

Furthermore, the association between MAOA gene with childhood maltreatment to the development of antisocial behaviours has been replicated, but not consistently. Several methodological issues can justify these discrepancies, including diverse sample , population genetic heterogeneity and exposure to various adverse environment.(19-21) The aim of this study is to analyze the missense mutations in MAOA gene that could result in the occurrence of antisocial and violent behaviors. Up to our knowledge, this will be the first bioinformatics study to address this topic.

Material and Methods:

2.1. Retrieving nsSNPs:

SNPs associated with MAOA gene were obtained from the Single Nucleotide Polymorphism database (dbSNP) in the National Center for Biotechnology Information (NCBI). (http://www.ncbi.nlm.nih.gov/snp/).

The sequence and natural variants of MAOA protein were obtained from the NCBI database as it considered as the most reliable database for protein sequences. [22] (https://www.uniprot.org/).

2.2. Identifying the most damaging nsSNPs and disease related mutations:

A total number of 342 nonsynonymous Single Nucleotide Polymorphisms (ns SNPs) were found from NCBI database, all of them were subjected to in silico analysis using nine different algorithms and softwares; SIFT, PROVEAN, PolyPhen-2, SNPs&GO, PhD-SNP, PMUT, I-mutant, GeneMANIA and Chimera.

2.2.1. SIFT Server:

Phenotypic effects of amino acid substitution on protein function were predicted by using Sorting Intolerant From Tolerant (SIFT) server, which is a powerful tool that use sequence homology. A list of nsSNPs (rsIDs) from NCBI's dbSNP database was submitted along with the (original) sequence to SIFT to predict tolerated and deleterious substitutions for every position of a protein sequence. The server divides the results into "Deleterious" and "Tolerated", nsSNPs with SIFT score ≤ 0.05 were classified as deleterious and are further analyzed to identify the damaging ones, and those > 0.05 were classified as tolerated and are not further analyzed. The server provide fast results but can not assess the result for SNPs that share the same position at the same time. (Available at: http://sift.bii.a-star.edu.sg/). [23-26]

2.2.2. Provean Server:

(Protein Variation Effect Analyzer) is the second software tool used. It predicts the effect of an amino acid substitution on the biological

function of a protein. It can predicts single amino acid substitutions, in-frame insertions, deletions, and multiple amino acid substitutions. The results are obtained as either "Deleterious" if the prediction score was ≤-2.5, while score > -2.5 indicates that the variant is predicted to have a "Neutral" effect. (Available at: http://provean.jcvi.org/index.php). [27, 28]

2.2.3. Polyphen-2 Server:

Polymorphism Phenotyping v2.0 (PolyPhen-2) is another online tool that predicts the possible effects of an amino acid substitution on the structure and function of the protein by analysis of multiple sequence alignment. The results are classified into "probably damaging" that is the most disease causing with a score near to 1 (0.7-1), "possibly damaging" with a less disease causing ability with a score of 0.5-0.8 and "benign" which does not alter protein functions with a score closer to zero; (0-0.4). We used patch query as it provide fast and accurate way to get the result. (Available at: http://genetics.bwh.harvard.edu/pph2/). [22, 29, 30]

2.2.4. SNAP2

It is another tool to predict Functional effects of mutations. SNAP2 is a trained classifier that is based on a machine learning device called "neural network". After entering the protein FASTA sequence as input query, It can distinguishes between effect and neutral variants/non-synonymous SNPs by taking a variety of sequence and variant features into account. The most important input signal for the prediction is the evolutionary information taken from an automatically generated multiple sequence alignment. Also structural features such as predicted secondary structure and solvent accessibility are considered. Although it take longer time than the previous software to get the result but it has a significant accuracy level (a sustained two-state accuracy (effect/neutral) of 82%).[31] It is available at https://rostlab.org/services/snap2web/.

2.2.4. SNPs & Go server:

An online web server that used to ensure the disease relationship with the studied single nucleotide polymorphisms SNPs. It gives three different results based on three different analytical algorithms; Panther result, PHD-SNP result, SNPs&GO result. Each one of these results is composed of three parts, the prediction which decides whether the mutation is neutral or disease related, reliability index (RI), and disease probability (if >0.5 mutation is considered as disease causing nsSNP). (Available at: http://snps-and-go.biocomp.unibo. it/snps-and-go/) . [32]

2.2.5. PMUT Server:

Is a neural network dependent web-based tool used for the prediction of pathological variants on proteins. The prediction results are classified as "Neutral" or "Disease". It is very fast and accurate, It is available at (http://mmb.irbbarcelona.org/PMut). [33]

2.3. Protein Structural analysis:

I-Mutant:

A online web-tool that is used for the prediction of protein stability changes upon single point mutations, determining whether the mutation

increases or decreases the protein stability. After insertion of reference sequencing and the required position and related mutant amino-acid, a result page will appears with the desired prediction. Acquiring result may take some time depending on the server load. (Available at: http://gpcr.biocomp.unibo.it/cgi/predictors/I-Mutant2.0/I-Mutant2.0.cgi). [34]

2.4. GeneMANIA:

A user-friendly web interface tool designed to asses protein function, analyzing submitted gene lists and prioritizing genes for functional assays. It provides lists with functionally similar genes that it identifies and recognizes using available genomics and proteomics data. It is accurate method to study gene –gene interaction. (Available at: (http://www.genemania.org/). [35]

2.5. Modeling nsSNP locations on protein structure:

2.5.1 Hope project:
It is a webserver that used to analyze the effect of single point mutation on structural level of protein. it best way to visualize the mutation since it is create a report consist of figures,animation,3d structure and mutation report just by submitting the protein sequence and mutation.[36] It is available at http://www.cmbi.ru.nl/hope/.

2.5.2 Chimera:

Chimera 1.8 software is used for visualization and editing of the three dimensional structure of protein. It visualizes the original amino acid with the mutated one to see the impact that can be produced. It is a highly extensible program for interactive visualization and analysis of molecular structures and related data, including density maps,

supramolecular assemblies, trajectories, and conformational ensembles. High-quality images and animations can be generated. (Availabe at: http://www.cgl.ucsf.edu/chimera). [37]

3. ConSurf server:

It is a web server suggests evolutionary conservation reviews for proteins of known structure in the PDB. ConSurf detect the similar amino acid sequences and run multi alignment approaches. The conserved amino acid across species detects its position using specific algorisms.(38)

4. Variant Effect Predictor (VEP):

The Ensembl Variant Effect Predictor software provides toolsets for an organized approach to annotate and assist for prioritization of mutations. The input data format was a list of variant identifiers while the output was filtered by choosing 1000 genome combined population to expend the population coverage.(39)

Result :

566 SNPs related to MAOA gene were downloaded from NCBI, of those 174 SNPs were missense mutation.these SNPS were futher analyzed by functional and structural software.

Table1:Deleterious nsSNPs associated variations predicted by SIFT, Polyphen, PROVEAN, SNAP2 softwares.

dbSNP rs#	Sub	SIFT PRE	SIFT SCORE	Prediction (cutoff= -2.5)	PROVEAN score	polyphen2 prediction	pph2_prob	snap2 prediction	snap2 score
rs1268810174	D15N	AFFECT	0	Deleterious	-3.01	probably damaging	0.999	effect	90
rs1131691720	L32S	AFFECT	0	Deleterious	-5.02	probably damaging	1	effect	30
rs796065312	R45W	AFFECT	0	Deleterious	-6.71	probably damaging	1	effect	69
rs201519600	D46G	AFFECT	0.03	Deleterious	-5.6	possibly damaging	0.662	effect	49
rs750916164	R79C	AFFECT	0.01	Deleterious	-6.88	probably damaging	1	effect	26
	T88I	AFFECT	0.01	Deleterious	-4.07	probably damaging	0.982	effect	34

rs140295792	R109W	AFFECT	0.01	Deleterious	-3.6	probably damaging	0.997	effect	47
rs1157474072	I138F	AFFECT	0.01	Deleterious	-3.66	probably damaging	0.999	effect	7
rs1450929926	P139Q	AFFECT	0.03	Deleterious	-6.97	possibly damaging	0.904	effect	5
rs199524208	A146V	AFFECT	0.03	Deleterious	-3.07	probably damaging	0.964	neutral	-12
rs1261565085	E159K	AFFECT	0.05	Deleterious	-2.88	possibly damaging	0.51	effect	51
rs574432879	I161T	AFFECT	0.02	Deleterious	-3.32	possibly damaging	0.782	neutral	-56
rs1390855194	R172W	AFFECT	0.01	Deleterious	-4.13	probably damaging	0.992	effect	55
rs77698881	E188K	AFFECT	0.02	Deleterious	-2.99	possibly damaging	0.804	effect	94
rs762184021	R217W	AFFECT	0.01	Deleterious	-6.68	probably damaging	1	effect	65
rs750174700	G221A	AFFECT	0	Deleterious	-5.66	probably damaging	1	effect	62
rs1181634890	Q225R	AFFECT	0.03	Deleterious	-3.43	possibly damaging	0.827	effect	28
rs779805250	R229W	AFFECT	0.02	Deleterious	-4.72	probably damaging	0.999	effect	36
	G235R	AFFECT	0.02	Deleterious	-5.39	probably damaging	0.999	neutral	-13
rs1216988286	P243S	AFFECT	0.04	Deleterious	-6.37	probably damaging	0.999	neutral	-31
rs587777457	C266F	AFFECT	0.01	Deleterious	-2.93	probably damaging	0.96	effect	92
	V269L	AFFECT	0.03	Deleterious	-2.82	probably damaging	0.996	effect	37
rs1452282305	A272V	AFFECT	0.01	Deleterious	-3.73	probably damaging	1	effect	68
rs749012045	P275L	AFFECT	0	Deleterious	-6.3	probably damaging	0.993	neutral	-19
rs780527254	L277S	AFFECT	0.01	Deleterious	-5.72	probably damaging	1	effect	56

Table 2: Disease effect nsSNPs associated variations predicted by Pmut, SNP&GO, PHD and I mutant softwares.

dbSNP rs#	sub	Prediction phd	RI	Probability	Prediction snp and go	RI	Probability	Pmut prediction	Pmut score
rs780647851	R297Q	AFFECT	0.03	Deleterious	-3.9	probably damaging	0.999	effect	58
rs61730725	L298P	AFFECT	0.02	Deleterious	-4.56	probably damaging	1	effect	77
rs867883669	P299S	AFFECT	0.02	Deleterious	-7.8	probably damaging	1	effect	22
rs1799835	F314V	AFFECT	0	Deleterious	-6.83	probably damaging	1	effect	89
rs1451367524	P342Q	AFFECT	0.05	Deleterious	-7.13	probably damaging	1	effect	43
rs1182312989	R356W	AFFECT	0.01	Deleterious	-4.92	probably damaging	0.999	effect	56
rs1428368525	A378T	AFFECT	0.01	Deleterious	-3.16	probably damaging	0.978	effect	10
rs1434727379	T417I	AFFECT	0.01	Deleterious	-5.23	probably damaging	0.998	effect	38
rs1181885385	A433T	AFFECT	0	Deleterious	-3.77	probably damaging	1	effect	21
rs770610370	A433V	AFFECT	0	Deleterious	-3.77	probably damaging	1	neutral	-3
rs764516141	T439R	AFFECT	0.03	Deleterious	-4.87	probably damaging	0.999	effect	47
rs1803986	M445I	AFFECT	0.03	Deleterious	-3.5	possibly damaging	0.925	neutral	-46
rs1359637885	G447R	AFFECT	0	Deleterious	-7.54	probably damaging	1	effect	93
rs775232342	G464R	AFFECT	0	Deleterious	-5.27	possibly damaging	0.529	effect	3
rs759671679	W472L	AFFECT	0.03	Deleterious	-9.44	possibly damaging	0.53	effect	19
rs1254032613	R526W	AFFECT	0	Deleterious	-3.21	probably damaging	0.998	effect	79

rs1131691720	L32S	Disease	7	0.87	Disease	4	0.725	Disease	0.90 (93%)
rs796065312	R45W	Disease	8	0.921	Disease	6	0.816	Disease	0.90 (93%)
rs750174700	G221A	Disease	4	0.679	Disease	2	0.581	Disease	0.72 (86%)
rs1799835	F314V	Disease	8	0.886	Disease	6	0.781	Disease	0.75 (88%)
rs1181885385	A433T	Disease	4	0.717	Disease	1	0.527	Disease	0.83 (90%)
rs1359637885	G447R	Disease	5	0.756	Disease	4	0.717	Disease	0.92 (94%)

Table (5) : Shows variant consequences, transcripts, and regulatory features by VEP tool:

Uploaded variation	Allele	Consequence	IMPACT	SYMBOL	Protein_position	Amino_acids
rs1131691720	C	missense variant	MODERATE	MAOA	32	L/S
rs1131691720	C	non coding transcript exon variant	MODIFIER	MAOA	-	-
rs1131691720	C	5 prime UTR variant	MODIFIER	MAOA	-	-
rs796065312	T	missense variant	MODERATE	MAOA	45	R/W
rs796065312	T	non coding transcript exon variant	MODIFIER	MAOA	-	-
rs796065312	T	5 prime UTR variant	MODIFIER	MAOA	-	-
rs750174700	C	missense variant	MODERATE	MAOA	221	G/A
rs750174700	C	non coding transcript exon variant	MODIFIER	MAOA	-	-
rs750174700	C	missense variant	MODERATE	MAOA	88	G/A
rs1799835	G	missense variant	MODERATE	MAOA	314	F/V
rs1799835	G	downstream gene variant	MODIFIER	MAOA	-	-
rs1799835	G	missense variant	MODERATE	MAOA	181	F/V
rs1181885385	A	missense variant	MODERATE	MAOA	433	A/T
rs1181885385	A	non coding transcript exon variant	MODIFIER	MAOA	-	-
rs1181885385	A	missense variant	MODERATE	MAOA	300	A/T
rs1359637885	A	missense variant	MODERATE	MAOA	447	G/R
rs1359637885	A	non coding transcript exon variant	MODIFIER	MAOA	-	-
rs1359637885	A	missense variant	MODERATE	MAOA	314	G/R

Reference sequence was submitted to RaptorX, to get the structural prediction of the protein, which was further studied by Chimera software; to visualize the changes in the wild and mutant amino acids.

Figure1:L32S: Leucine (green color) changed to Serine (red color) at position 32.

Figure 2:A45W:Arginine (green color) changed to Tryptophan (red color) at position 45.

Figure3:G221A:Glycine (green color) changed to Alanine(red color) at position 221.

Figure4:F314V:Phenylalanine (green color) changed to Valine(red color) at position 314.

Figure 5:A422T:Alanine (green color) changed to Threonine (red color) at position 433.

Figure6:G447A:Glycine (green color) changed to Arginine (red color) at position 447.

Figure7: shows gene-gene interaction by geneMANIA.

Figure(8): The conserved amino acids across species in MAOA protein were determined using ConSurf. e: exposed residues according to the neural-network algorithm are indicated in orange letters. b: residues predicted to be buried are demonstrated via green letters. f: predicted functional residues (highly conserved and exposed) are indicated with red letters. s: predicted structural residues (highly conserved and buried) are demonstrated in blue letters. I: insufficient data (the calculation for this site performed in less than 10% of the sequences) is demonstrated in yellow letters.

Discussion:

Six novel missense SNPs were found in MAOA gene to effect protein function and structure. Out of 566 SNPS downloaded from NCBI, 34 SNPs have been found to be damaging by four functional software. Which was further analyzed by phd ,snp and go and pmut, where 6 SNPs were found to be damaging, as illustrated in table (1 and 2).

Project hope was used to study the physiochemical changes inflicted by the missense mutations. All the SNPs were found to be located at Fad/Nad(P)-Binding Domain Superfamily IPR036188 and Flavin Amine Oxidase IPR001613 domain which indicate this SNPs is likely to be conserved . In all the SNPs the mutant amino acid were bigger in size than the wild type except in F314V and L32S SNPs . In addition all the mutant amino acids were found to be different in hydrophobicity when compared with the wild type amino acid. All these changes will affect the protein function.

Structural changes were further studied using chimera software , where changes in the amino acids Cleary change the structural of the MAOA protein as evident in figure(1-6).

The interaction and function of MAOA gene was studied using gene mania. where the predicted functions was: amine metabolic process, cellular amine metabolic process, cellular biogenic amine metabolic process, cellular response to xenobiotic stimulus, oxidoreductase activity, acting on the CH-NH2 group of donors, oxidoreductase activity, acting on the CH-NH2 group of donors, oxygen as acceptor, response to xenobiotic stimulus and xenobiotic metabolic process . The MAOA gene were found to have interactions with many other genes like CD50,F3,HGD and others as shown in Figure (7).

The Variant Effect Predictor annotates mutations using an extensive array of reference data from previously detected mutations, evidence based results, and estimation of biophysical consequences of mutations; and that is what makes VEP an accurate web based tool. .(39) VEP described regulatory consequences for several mutations, including 10 mutations within coding region, 5 mutations within a non-coding region. briefly, mutations within coding region affect the protein function, while mutations within non-coding regions can significantly affect disease and could contribute in the phenotypic feature and RNA-binding proteins (RBPs);(40, 41) while mutations in the upstream, downstream, 5⍰-, and 3⍰-UTRs might affect transcription or translation process.(42) The consequences are shown in Table(5).

We also used ConSurf web server; the nsSNPs that are located in highly conserved regions and predicted to cause structural and functional impacts on MAOA protein Figure (8)

Previous researches shows common regulatory variation in MAOA to moderate effects of childhood maltreatment on male antisocial behaviors.(21) while another research showed that certain SNPs polymorphism (1460 C>T) is associated with other conditions leading to the development of depressive symptoms in menopausal women.(43) In another research underwent in Chinese population the interaction between catechol-O-methyltransferase (COMT), Ala22/72Ser and MAOA T941G polymorphisms, and stressful life events (SLE) in the academic pressure and the relation to aggressive behavior development was also identified.(44) These SNP was not available in the data we acquired from NCBI, yet these studies does indicate the significant role of MAOA gene polymorphisms in aggrieve behavior development.

This study is limited, as it does not cover the environmental factor important for the development of antisocial behavior, nevertheless this study will lay the foundation for further clinical research in this phenomena.

Conclusion:

Six novel missense SNPs were found in MAOA gene to affect MAOA protein function and structure, which could predispose to violent and aggressive behavior.

Conflict of interest:

The authors decline any conflict of interest.

1.   Aslund C, Nordquist N, Comasco E, Leppert J, Oreland L, Nilsson KW. Maltreatment, MAOA, and delinquency: sex differences in gene-environment interaction in a large population-based cohort of adolescents. Behav Genet. 2011;41(2):262-72.
2.   Denson TF, Dobson-Stone C, Ronay R, von Hippel W, Schira MM. A functional polymorphism of the MAOA gene is associated with neural responses to induced anger control. J Cogn Neurosci. 2014;26(7):1418-27.
3.   Hwang IW, Lim MH, Kwon HJ, Jin HJ. Association of Monoamine Oxidase A (MAOA) Gene uVNTR and rs6323 Polymorphisms with Attention Deficit and Hyperactivity Disorder in Korean Children. Medicina (Kaunas). 2018;54(3).
4.   Klasen M, Wolf D, Eisner PD, Habel U, Repple J, Vernaleken I, et al. Neural networks underlying trait aggression depend on MAOA gene alleles. Brain Struct Funct. 2018;223(2):873-81.
5.   McDermott R, Tingley D, Cowden J, Frazzetto G, Johnson DD. Monoamine oxidase A gene (MAOA) predicts behavioral aggression following provocation. Proc Natl Acad Sci U S A. 2009;106(7):2118-23.
6.   Beaver KM, Barnes JC, Boutwell BB. The 2-repeat allele of the MAOA gene confers an increased risk for shooting and stabbing behaviors. Psychiatr Q. 2014;85(3):257-65.
7.   Craig IW. The role of monoamine oxidase A, MAOA, in the aetiology of antisocial behaviour: the importance of gene-environment interactions. Novartis Found Symp. 2005;268:227-37; discussion 37-41, 42-53.

8.   Bortolato M, Floris G, Shih JC. From aggression to autism: new perspectives on the behavioral sequelae of monoamine oxidase deficiency. J Neural Transm (Vienna). 2018;125(11):1589-99.

9.   Golub MS, Hogrefe CE, Unger EL. Influence of prenatal iron deficiency and MAOA genotype on response to social challenge in rhesus monkey infants. Genes Brain Behav. 2012;11(3):278-90.

10.   Lei H, Zhang X, Di X, Rao H, Ming Q, Zhang J, et al. A functional polymorphism of the MAOA gene modulates spontaneous brain activity in pons. Biomed Res Int. 2014;2014:243280.

11.   Bortolato M, Shih JC. Behavioral outcomes of monoamine oxidase deficiency: preclinical and clinical evidence. Int Rev Neurobiol. 2011;100:13-42.

12.   Edgnulu TG, Ozge A, Erdal N, Kuru O, Erdal ME. Association analysis of the functional MAOA gene promoter and MAOB gene intron 13 polymorphisms in tension type headache patients. Adv Clin Exp Med. 2014;23(6):901-6.

13.   Godar SC, Bortolato M, Castelli MP, Casti A, Casu A, Chen K, et al. The aggression and behavioral abnormalities associated with monoamine oxidase A deficiency are rescued by acute inhibition of serotonin reuptake. J Psychiatr Res. 2014;56:1-9.

14.   Kim SK, Park HJ, Seok H, Jeon HS, Chung JH, Kang WS, et al. Association study between monoamine oxidase A (MAOA) gene polymorphisms and schizophrenia: lack of association with schizophrenia and possible association with affective disturbances of schizophrenia. Mol Biol Rep. 2014;41(5):3457-64.

15.   Guo H, Ren Y, Zhao N, Wang Y, Li S, Cui H, et al. Synergistic effect of 5-HT2A receptor gene and MAOA gene on the negative emotion of patients with depression. Clin Physiol Funct Imaging. 2014;34(4):277-81.

16.   Schiele MA, Ziegler C, Kollert L, Katzorke A, Schartner C, Busch Y, et al. Plasticity of Functional MAOA Gene Methylation in Acrophobia. Int J Neuropsychopharmacol. 2018;21(9):822-7.

17.   Golub M, Hogrefe C. Prenatal iron deficiency and monoamine oxidase A (MAOA) polymorphisms: combined risk for later cognitive performance in rhesus monkeys. Genes Nutr. 2014;9(2):381.

18.   Nilsson KW, Comasco E, Hodgins S, Oreland L, Aslund C. Genotypes do not confer risk for delinquency but rather alter susceptibility to positive and negative environmental factors: gene-environmentinteractions of BDNF Val66Met, 5-HTTLPR, and MAOA-uVNTR [corrected]. Int J Neuropsychopharmacol. 2014;18(5).

19.   Ouellet-Morin I, Cote SM, Vitaro F, Hebert M, Carbonneau R, Lacourse E, et al. Effects of the MAOA gene and levels of exposure to violence on antisocial outcomes. Br J Psychiatry. 2016;208(1):42-8.

20.   Pingault JB, Cote SM, Booij L, Ouellet-Morin I, Castellanos-Ryan N, Vitaro F, et al. Age-dependent effect of the MAOA gene on childhood physical aggression. Mol Psychiatry. 2013;18(11):1151-2.

21.   Byrd AL, Manuck SB. MAOA, childhood maltreatment, and antisocial behavior: meta-analysis of a gene-environment interaction. Biol Psychiatry. 2014;75(1):9-17.

22.   Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen-2. Curr Protoc Hum Genet. 2013;Chapter 7:Unit7 20.

23.   Ng PC, Henikoff SJGr. Accounting for human polymorphisms predicted to affect protein function. 2002;12(3):436-46.

24.   Manickam M, Ravanan P, Singh P, Talwar P. In silico identification of genetic variants in glucocerebrosidase (GBA) gene involved in Gaucher's disease using multiple software tools. Front Genet. 2014;5:148.

25.   Ng PC, Henikoff SJARGHG. Predicting the effects of amino acid substitutions on protein function. 2006;7:61-80.

26.   Schneider G, Hu J, Sim N-L, Kumar P, Henikoff S, Ng PC.

27.   Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. PLoS One. 2012;7(10):e46688.

28.   Kumar P, Mahalingam K. In silico approach to identify non-synonymous SNPs with highest predicted deleterious effect on protein function in human obesity related gene, neuronal growth regulator 1 (NEGR1). 3 Biotech. 2018;8(11):466.

29.   Ma J, Wang S, Zhao F, Xu J.

30.   Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. Nature methods. 2010;7(4):248-9.

31.   Hecht M Fau - Bromberg Y, Bromberg Y Fau - Rost B, Rost B. Better prediction of functional effects for sequence variants. (1471-2164 (Electronic)).

32.   Calabrese R, Capriotti E, Fariselli P, Martelli PL, Casadio RJHm. Functional annotations improve the predictive score of human disease-related mutations in proteins. 2009;30(8):1237-44.

33.   López-Ferrando V, Gazzo A, de la Cruz X, Orozco M, Gelpí JLJNar. PMut: a web-based tool for the annotation of pathological variants on proteins, 2017 update. 2017;45(W1):W222-W8.

34.   Capriotti E, Fariselli P, Casadio RJNar. I-Mutant2. 0: predicting stability changes upon mutation from the protein sequence or structure. 2005;33(suppl_2):W306-W10.

35.   Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al. The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function. 2010;38(suppl_2):W214-W20.

36.   Venselaar H, Te Beek Ta Fau - Kuipers RKP, Kuipers Rk Fau - Hekkelman ML, Hekkelman Ml Fau - Vriend G, Vriend G. Protein structure analysis of mutations causing inheritable diseases. An e-Science approach with life scientist friendly interfaces. (1471-2105 (Electronic)).

37.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera—a visualization system for exploratory research and analysis. 2004;25(13):1605-12.

38.   Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, et al. ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in macromolecules. Nucleic acids research. 2016;44(W1):W344-50.

39.   McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, et al. The Ensembl Variant Effect Predictor. Genome biology. 2016;17(1):122.

40.   Rojano E, Seoane P, Ranea JAG, Perkins JR. Regulatory variants: from detection to predicting impact. Briefings in bioinformatics. 2018.

41.   Dhamija S, Menon MB. Non-coding transcript variants of protein-coding genes - what are they good for? RNA biology. 2018;15(8):1025-31.

42.   Tatarinova TV, Chekalin E, Nikolsky Y, Bruskin S, Chebotarov D, McNally KL, et al. Nucleotide diversity analysis highlights functionally important genomic regions. Scientific reports. 2016;6:35730.

43.   Rozycka A, Slopien R, Slopien A, Dorszewska J, Seremak-Mrozikiewicz A, Lianeri M, et al. The MAOA, COMT, MTHFR and ESR1 gene polymorphisms are associated with the risk of depression in menopausal women. Maturitas. 2016;84:42-54.

44.   Wang M, Li H, Deater-Deckard K, Zhang W. Interacting Effect of Catechol-O-Methyltransferase (COMT) and Monoamine Oxidase A (MAOA) Gene Polymorphisms, and Stressful Life Events on Aggressive Behavior in Chinese Male Adolescents. Front Psychol. 2018;9:1079.