Thermal cycling protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell injury through stress response mechanisms involving Akt pathway

Wei-Ting Chen[1,2,¶], Yu-Yi Kuo[1,2,¶], Guan-Bo Lin[1,2,¶], Chueh-Hsuan Lu[1,2],
Hao-Ping Hsu[3], Yi-Kun Sun[2], and Chih-Yu Chao[1,2,4,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National
Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of
Medicine, Taipei 10051, Taiwan

[3] Department of Medicine, National Taiwan University College of Medicine, Taipei
10051, Taiwan

[4] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan
University, Taipei 10617, Taiwan

* Corresponding author
E-mail: cychao@phys.ntu.edu.tw (CYC)
¶ These authors contributed equally to this work.

Abstract

Neurodegenerative diseases (NDDs) are becoming a major threat to public health, according to the World Health Organization (WHO). The most common form of NDDs is Alzheimer's disease (AD), boasting 60-70% share. Excessive aggregation of β-amyloid protein (Aβ) and neurofibrillary tangles has been deemed the major cause for the pathogenesis of AD. A growing number of evidences from studies, however, have suggested that reactive oxygen species (ROS) also play a key role in the onset and progression of AD. Although scientists have had some understanding of the pathogenesis of AD, the disease still cannot be cured, with existing treatment only capable of providing a temporary relief at best, partly due to the obstacle of blood-brain barrier (BBB). The study was aimed to ascertain the neuroprotective effect of thermal cycle hyperthermia (TC-HT) against hydrogen peroxide (H_2O_2) and Aβ-induced cytotoxicity in SH-SY5Y cells. Treating cells with this physical stimulation beforehand significantly improved the cell viability and decreased the ROS content. The underlying mechanisms were confirmed to be due to the activation of Akt pathway and the downstream antioxidant and prosurvival proteins. The findings manifest significant potential of TC-HT in neuroprotection, via inhibition of oxidative stress and cell apoptosis. It is believed that coupled with the use of drugs or natural compounds, this methodology can be even more effective in treating NDDs.

Introduction

According to the World Health Organization (WHO), the number of dementia-induced deaths more than doubled between 2000 and 2016, making it the 5th leading cause for deaths worldwide in 2016, up from 14th place in 2000. Among the various forms of dementia such as Alzheimer's disease (AD) and Parkinson's disease (PD), AD is the most common one, accounting for 60–70% of the cases. Typically, AD exhibits such features as deposition of cortical plaques caused by excessive aggregation of β-amyloid protein (Aβ) and neurofibrillary tangles, and progressive brain degeneration and deterioration of cognitive function among elderly people. Although the exact mechanism of AD pathogenesis remains unknown, it is believed that oxidative stress and activation of free radicals, induced by Aβ aggregation, play a key role in AD pathogenesis [1].

Reactive oxygen species (ROS) are reactive chemical species containing oxygen, which are generated as natural byproduct of oxygen metabolism. ROS play important roles in cell signalling and homeostasis, and their concentrations in cells are subtlety regulated by various antioxidant compounds and enzymes. However, with cells under continuing environmental stress (e.g. ultraviolet, inflammatory cytokines, or environmental toxins), the imbalance between prooxidants and antioxidants may cause chronic oxidative stress. Accumulation of ROS may cause cell death, accelerate

cell ageing, or induce age-related diseases [2]. More and more research evidences

have suggested that ROS plays a central role in the onset and progression of AD [3].

Therefore, the protection of neural cells against oxidative damage may be a potential

strategy to treat AD. Several in vitro or in vivo studies have explored the function of

antioxidant and antiapoptotic drugs in ameliorating AD [4-6] but the approach is

time-consuming and costly, plus the safety concern, which limits the use of these

drugs in AD treatment. Moreover, the blood-brain barrier (BBB) dampens the efficacy

of these drugs, since over 98% of small molecule drugs and ~100% of large molecule

drugs can not pass the BBB [7]. Therefore, a non-drug treatment may be more suited

to AD patients.

Scientists have long been interested in the profound effects of heat on cells, and

have utilized it in various types of thermotherapeutical applications such as

physiotherapy, urology, and cardiology [8]. One promising and effective thermal

therapy is the treatment of cancer by hyperthermia (HT) [9]. HT is used to kill cancer

cells directly or to potentiate the cytotoxicity of radiation and certain chemotherapy

drugs [10]. The ROS level increased by HT treatment has been identified to play an

important role as an intracellular mediator of HT-induced cell death [11]. On the

contrary, it has also been reported that heat shock (HS) will induce many cellular

defense, including the antioxidant effect. For example, Tchouagué *et al* demonstrated

that HS-generated ROS is involved in induction of cellular defense molecules Prxs,

GSH and G6PD through Nrf2 activation [12]. Mustafi *et al* also showed that heat

stress upregulates the HSP70 and MnSOD levels through ROS and p38MAPK [13].

In addition to the thermal treatment, the beneficial effects of light treatment were also

reported in literatures. The review article by Hamblin summarized some pre-clinical

studies and clinical trials by light therapy for brain disorders [14]. The physical

stimulation, therefore, holds great potential for AD or other neurodegenerative

diseases (NDDs).

The study employed a special thermal therapy, applied at high and low

temperatures alternately to achieve an effect similar to antioxidant and antiapoptotic

drugs. In our previous study, we used this novel method, thermal cycle hyperthermia

(TC-HT) for the treatment of pancreatic cancer, and found that this physical

stimulation greatly improved the anticancer effect of propolis and polyphenols on

PANC-1 cells without the heat-induced side effect [15, 16]. Traditional HS or HT

employs continuous heating to achieve curative effect, but it is likely to cause neuron

damage since damage to the central nervous system occurs within few minutes of

exposure to 42°C [17]. Nevertheless, the feature of TC-HT is that it can avoid the

damage caused by HT, which is crucial for neuroprotection.

In this study, we applied the TC-HT strategy to human neural cell line SH-SY5Y,

which has been extensively used in research on neurodegenerative damage in vitro,

and examined the prosurvival effect of TC-HT on preventing oxidative damage

induced by hydrogen peroxide (H_2O_2) and Aβ. The results showed that subjection of

the cells to heat at high and low temperatures alternately beforehand ameliorated the

H_2O_2 and Aβ-induced cytotoxicity in SH-SY5Y cells significantly. It was found that

TC-HT not only performed superior protective effect than the traditional HS but also

avoided thermal damage caused by continuous heating. Examination of the

underlying mechanism also showed that TC-HT could activate specific

neuroprotective proteins. These findings indicate that TC-HT is a promising thermal

therapy, which sheds light on novel treatment for AD or other NDDs.

Materials and methods

Cell culture and treatment

The human neuroblastoma SH-SY5Y cells purchased from Bioresource Collection

and Research Center (Hsinchu, Taiwan) were maintained in MEM/F-12 mixture

containing 10% fetal bovine serum (HyClone; GE Healthcare Life Sciences) and 1%

penicillin-streptomycin, supplemented with 1mM sodium pyruvate and 0.1 mM

non-essential amino acids. The cells were cultured at 37°C in a humidified incubator

composed of 5% CO_2. SH-SY5Y cells were seeded in 96-well plates overnight. Cells

were pretreated with HT or TC-HT by the Thermal Cycler (Applied Biosystems;

Thermo Fisher Scientific, Inc.) relative to the control at room temperature (RT). After

the treatment, the cells were maintained in a 37°C incubator for 4 h. Subsequently, the

pretreated cells were induced to undergo apoptosis by adding H_2O_2 to the culture

medium. Cell viability was measured 24 h after the H_2O_2 treatment. As for AD disease

model, the cells were treated with 25 or 50 μM $A\beta_{25\text{-}35}$ protein solution. The Aβ stock

solution was prepared by solubilizing the $A\beta_{25\text{-}35}$ peptide (Sigma-Aldrich; Merck

KGaA) in sterile deionized water to a concentration of 1 mM and then incubated at

37°C for 3 days to allow self-aggregation before treatment. The TC-HT therapy was

applied before or after the Aβ treatment, and the cell viability was determined 4 days

after the treatment. For the Akt inhibitor experiment, 25 μM LY294002 (Cell

Signaling Technology, Inc.) was treated 1 h before the TC-HT.

131   Cell Viability Assay

The cell viability was determined by

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich;

Merck KGaA) assay. In brief, the culture medium was replaced with MTT solution

(0.5 mg/mL in MEM/F12) and incubated at 37°C for 4 h. Equal volume of the

solubilizing buffer (0.01 M HCl and 10% SDS) was added to dissolve the formazan

crystals. The 96-well plates were analyzed on a FLUOstar OPTIMA microplate reader

(BMG Labtech, Ltd.) at 570 nm, and background absorbance at 690 nm was

subtracted.

Lactate dehydrogenase (LDH) assay

The assay measures a stable enzyme LDH, which is released into the cell culture

medium when cell membranes are damaged. LDH levels were measured using a

Cytoscan LDH cytotoxicity assay kit (G-Biosciences; Geno Technology Inc.)

according to the manufacturer's instructions. Briefly, the cells were seeded in 96-well

plates and then incubated with H_2O_2 with or without thermal cycle pretreatment. For

analysis, 50 µL culture mediums from all wells were transferred to a new 96-well

plate, and 50 µL of reaction mixtures were added to each well and incubated at 37°C

for 20 min. The absorbance was measured at 490 nm using a microplate reader (BMG

Labtech, Ltd.).

ROS level detection

ROS levels of cells were detected using the fluorescent dye dihydroethidium (DHE)

(Sigma-Aldrich; Merck KGaA). Cells pretreated with TC-HT or HT at 42.5°C high

temperature setting for 8 cycles or 2 h, respectively, were challenged with H_2O_2. After

the treatment of H_2O_2 for 24 h, cells were washed with PBS, and then incubated with

5 μM DHE dye for 20 min at 37°C in the dark. The fluorescence intensity emitted by

DHE was measured by flow cytometry in the PE channel, and ROS levels were

expressed as mean fluorescence intensity for comparison.

Mitochondrial membrane potential (MMP) measurement

The MMP was determined by flow cytometry using 3,3'-dihexyloxacarbocyanine

iodide ($DiOC_6(3)$) (Enzo Life Sciences International Inc.). $DiOC_6(3)$ is a lipophilic

cationic fluorescent dye which allows estimation of the percentage of cells with low

MMP. Cells were harvested and suspended at a density of 1 x 10^6 cells/mL in 1 μM

$DiOC_6(3)$ dye working solution. After incubation at 37°C for 15 min, $DiOC_6(3)$

intensity was analyzed by flow cytometry in the FITC channel.

Western blot analysis

170   The protein expression levels of SH-SY5Y cells were investigated by western blot

171   analysis. Cells treated with TC-HT, HT, H_2O_2, or Aβ were harvested and lysed in

172   RIPA lysis butter (EMD Millipore). After centrifugation and supernatant collection,

173   equal amounts of proteins were resolved by 10% SDS-PAGE and transferred to

174   polyvinylidene fluoride membranes. After drying for 1 h at RT, the membranes were

probed with p-Akt, Akt, Nrf2, p-CREB, PSMA3, PSMC3 (Cell Signaling Technology, Inc.), HO-1 (Enzo Life Sciences International Inc.), insulin-degrading enzyme (Abcam) and GAPDH (GeneTex, Inc.) antibodies overnight at 4˚C. The washed membranes were then incubated with horseradish peroxidase-conjugated goat anti-rabbit secondary antibodies (Jackson ImmunoResearch Laboratories, Inc.). Immunoreactivity was visualized with an enhanced chemiluminescence substrate (Advansta, Inc.) and detected with the Amersham Imager 600 imaging system (GE Healthcare Life Sciences). The images were analyzed with Image Lab software (Bio-Rad Laboratories, Inc.).

Statistical analysis

The results were presented as the mean ± standard deviation. Statistical analyses using one-way analysis of variance (ANOVA) followed by Tukey's post-hot test were performed by OriginPro 2015 software (OriginLab). P-value < 0.05 was considered to indicate a statistically significant difference.

Results

In vitro-applied TC-HT

We applied the thermal cycle (TC) treatment to SH-SY5Y cells by a modified

polymerase chain reaction (PCR) equipment (Fig 1C). Briefly, in our design, some

protruding parts of the PCR machine and plastic well were milled so that the bottom

of the well can touch the heat sink tightly. The basic TC settings were schemed as Fig

1A, where the temperature was elevated to the desired high temperature and sustained

for a period of time followed by a cooling process at body temperature. It's worth

noting that the adjustment of TC parameters (temperature and time) is based on the

needs and objectives of the experiments. In our experiments, the TC parameters were

set to high temperatures of 42.5°C or 41.5°C for 15 min, then lowered to 37°C and

lasted for 35 sec. This cycle was repeated 8 or 12 times. The HT group maintains the

same high temperature setting throughout the treatment in a continuous manner

without a break. To determine the actual temperatures of the cells, a needle

thermocouple located in the bottom of the well was used to monitor the temperatures.

Fig 1B represents the actual temperature measured by the thermocouple every 20 sec.

The actual cycle temperature sensed by the cells was ~42-40.3°C for the setting of

42.5-37°C and ~40.9-39.4°C for 41.5-37°C setting, as shown in Fig 1B.

Fig 1. The setup of TC-HT. (A) Schematic representation of TC parameter settings.

(B) Measurement of the cell temperature during the TC treatment by the

thermocouple. (C) Image of the TC controller setup.

Effect of TC-HT on H_2O_2 and Aβ-induced cytotoxicity in SH-SY5Y cells

We applied H_2O_2 to SH-SY5Y cells as an oxidative stress and the cell viability

was assessed by MTT assay. As shown in Fig 2A, treatment of H_2O_2 on SH-SY5Y

cells for 24 h induced significant decrease of cell viability in a dose-dependent manner.

Treatment of SH-SY5Y cells with 450 µM H_2O_2 for 24 h reduced the cell viability to

52.9% compared to control cells, which was used in the following experiments. To

investigate whether TC-HT or HT produced protection and increased cell survival

against the oxidative stress induced by H_2O_2, the SH-SY5Y cells were pretreated with

TC-HT or HT in the experiments. Two high temperature parameters (41.5°C or

42.5°C) were applied for 2 h or 3 h continuously in the HT group, and 8 or 12 cycles

(for 15 min heating time per cycle) in the TC group to make the total thermal doses of

the HT and TC groups equal. After the heat pretreatment, the treated SH-SY5Y cells

were kept in the incubator at 37°C for 4 h. They were then challenged with H_2O_2 to

generate the oxidative stress, and the cell viability was determined by MTT assay at

24 h after the H_2O_2 treatment. The results found that H_2O_2 treatment significantly

reduced the viability of SH-SY5Y cells to 53.7% of the control value, and showed

that the heat pretreatment conferred protective effect and increased the cell viability.

At 41.5°C for 2 h and 8 cycles, HT and TC treatments increased the cell viability to

61.9% and 68.5% of the control value, respectively (Fig 2B). As we increased the

high temperature setting to 42.5°C while maintaining the same heating time, TC

pretreatment greatly increased the cell viability to 83% of the control value under

H_2O_2 stress, while the HT group only exhibited a slight increase (Fig 2C). When we

further increased the heating time to 3 h, the treatment of HT alone was cytotoxic to

the cells, reducing the cell viability to 75.4%, and the viability was even lower after

the H_2O_2 treatment (Fig 2C). Interestingly, the TC treatment with the same total

thermal dose (12 cycles) had only a slight influence on the cell viability, and it still

conferred significant neuroprotective effect to the SH-SY5Y cells under H_2O_2 stress.

In particular, the study examined the influence of the cooling process on the

neuroprotective effect. As shown in Fig 2D, for low temperature period longer than 1

min, the TC pretreatment did not produce protective effect to the cells, indicating that

the low temperature period is a critical parameter to be determined for effective

neuroprotection. From molecular point of view, the neuroprotective effect would be

discontinued as long as there were at least one or few critical signalling pathways

being influenced as heating was halted, which coincided with the observation that the

low temperature period used for optimum neuroprotection was shorter than the

heating time. On the other hand, the LDH release rate was measured and found to

decrease significantly in the TC pretreatment group compared to the H_2O_2-treated

group (Fig 2E), further confirming its protective role in maintaining the cell

membrane integrity when cell was under oxidative stress. For AD disease model, Aβ

treatment was used to reduce the viability of SH-SY5Y cells due to the cytotoxicity of

the aggregated Aβ (Fig 3A). Our results showed that the TC treatment after Aβ

administration greatly improved the viability of SH-SY5Y cells (Fig 3B) for nearly

25%, indicating the curative effect of TC in AD disease model in vitro. The most

effective high temperature setting was 42.3°C for 8 cycles in the Aβ-treated cells. The

actual high temperature sensed by the cells was about 41.7°C. Interestingly, the

curative effect of TC pretreatment was less pronounced than that of TC post-treatment

(treated after Aβ administration), indicating the time point of TC application has great

impact on the curative effect. Moreover, there were no curative effects by both HT

pretreatment and post-treatment on SH-SY5Y cells, whose viability levels were even

slightly lower than that of the Aβ-treated cells (Fig 3B). Furthermore, the light

microscopy images showed that the integrity of the cells was destructed by Aβ, and

the TC treatment caused the protective effect to retain the cell morphology, as shown

in Fig 3C.

To conclude the result of the H_2O_2-induced oxidative stress on SH-SY5Y cells,

the protective effect of TC pretreatment was more pronounced than the HT

preconditioning. And the TC pretreatment at 42.5°C produced much superior

protection effect than at 41.5°C. Noteworthily, the HT pretreatment at 42.5°C for 3 h

decreased the cell viability which indicates that the continuous heating may induce

cell damage. The protective effect of TC was most remarkable at ~42.5°C for 8 cycles.

Therefore, we chose this parameter to evaluate the possible protective mechanism of

TC-HT in the following experiments.

Fig 2. Effect of TC-HT on H_2O_2-induced cytotoxicity in SH-SY5Y cells. (A)

Dose-response curve of SH-SY5Y cells treated with different concentrations of H_2O_2

for 24 h. (B) SH-SY5Y cells were pretreated at 41.5°C temperature setting and

challenged with or without 450 μM H_2O_2. The cell viability was measured by MTT

assay at 24 h after the H_2O_2 treatment. (C) SH-SY5Y cells were pretreated at 42.5°C

temperature setting with different thermal dosages and challenged with or without 450

μM H_2O_2. The cell viability was measured by MTT assay at 24h after the H_2O_2

treatment. (D) Comparison of the neuroprotective effect under different low

temperature period settings. (E) The LDH release was measured to confirm the

neuroprotective effect of TC treatment. Data represent the mean ± standard deviation

(n=3). [***]P < 0.001.

Fig 3. Effect of TC-HT on Aβ-induced cytotoxicity in SH-SY5Y cells. (A) The cell

viability of SH-SY5Y cells treated with 25 or 50 μM Aβ for 4 days. (B) The TC or HT

pretreatment and post-treatment were applied to the cells before or after 50 μM Aβ

administration, and the cell viability was measured by MTT assay 4 days after

treatment. (C) Representative light microscopy images of SH-SY5Y cells after

treatment. The integrity of the cells was destructed by Aβ, and the TC post-treatment

caused the protective effect and retained the cell morphology. Scale bar = 100 μm.

Data represent the mean ± standard deviation (n=3). ***$P < 0.001$.

TC-HT attenuates H_2O_2-induced ROS generation

The onset and progression of AD has been reported to be associated with ROS. We

further studied whether the H_2O_2-induced intracellular ROS production could be

attenuated by the TC pretreatment. The intracellular ROS levels were detected using

the fluorescent dye DHE (Fig 4A). Results showed that after exposure to 450 μM

H_2O_2 for 24 h, ROS level was increased to 160% of the control value. When the

SH-SY5Y cells were pretreated with TC-HT, the intracellular ROS level was

significantly ameliorated (Fig 4B). Compared to the TC group, the ROS level was not

changed in the HT group.

Fig 4. Effect of TC-HT on H_2O_2-induced ROS generation in SH-SY5Y cells. (A)

ROS level was measured by flow cytometry with DHE fluorescent dye. (B)

Quantification of the ROS levels after H_2O_2, TC+H_2O_2, or HT+H_2O_2 treatment. Data

represent the mean ± standard deviation (n=3). [**]$P < 0.01$.

TC-HT attenuates MMP loss in H_2O_2-treated cells

Many lines of evidence have suggested that mitochondria contribute to ageing-related

neurodegenerative diseases through the accumulation of net cytosolic ROS production

[18]. The levels of MMP are kept relatively stable in healthy cells. Apoptotic signals

are usually initiated by the disruption of normal mitochondrial function, especially the

collapse of MMP. To study the mechanism why H_2O_2-induced apoptosis is alleviated

by TC-HT pretreatment, the MMP is further analyzed by the flow cytometry with

$DiOC_6(3)$ fluorescent dye in the study. As shown in Fig 5A, the oxidative stress

induced by H_2O_2 caused mitochondrial dysfunction, decreasing the MMP and

therefore decreased the intensity of the $DiOC_6(3)$ fluorescent signal. From the

quantification results (Fig 5B), it was found that the percentage of cells with

decreased MMP drastically increased after the H_2O_2 treatment. In contrast to the

group of H_2O_2 alone, the TC-HT pretreatment noticeably suppressed the

H_2O_2-induced dissipation of MMP, so the cells with decreased MMP reduced

significantly, which was attributed to the preservation of the mitochondrial function

and thus caused the blockage of the apoptotic signaling transduction.

Fig 5. Effect of TC-HT on H_2O_2-induced MMP reduction in SH-SY5Y cells. (A)

MMP was analyzed by flow cytometry with $DiOC_6(3)$ fluorescent dye. (B)

Quantification of the cells with decreased MMP after H_2O_2, $TC+H_2O_2$, or $HT+H_2O_2$

treatment. Data represent the mean ± standard deviation (n=3). [***]$P < 0.001$.

Effect of heat treatment on heat shock protein (HSP) expressions in SH-SY5Y cells

It has been well known that HT triggers expression of HSPs so that cells are able to

protect themselves from stress. To investigate whether the protective effect of TC-HT

was related to the HSPs, we examined the protein expressions of HSP70 and HSP105,

both of which were reported to have protective effects. As shown in Fig 6, both HSP

expressions were enhanced significantly after HT and TC treatments. The activation

levels of TC, however, was only slightly higher than HT without significant difference.

This phenomenon points out that although TC treatment indeed protects SH-SY5Y

cells more from the oxidative stress of H_2O_2, the main mechanism could be apart from

the HSP70 or HSP105. Therefore, other heat-activated stress response proteins should

be considered in the experiments.

Fig 6. Effect of TC-HT on expressions of HSPs in SH-SY5Y cells. (A) Western blot

analysis of HSP70 and HSP105 expressions. (B) Quantification of HSP70 and

HSP105 expressions after H_2O_2, TC+H_2O_2, or HT+H_2O_2 treatment. The expression

levels were normalized to GAPDH. Data represent the mean ± standard deviation

(n=3). [***]P < 0.001.

Effect of heat treatment on insulin-degrading enzyme and

proteasome expressions in SH-SY5Y cells

Insulin-degrading enzyme (IDE) is a highly conserved zinc metallopeptidase which

was originally identified as the key enzyme for the degradation of insulin. Further

studies unraveled its ability to degrade several other polypeptides, including Aβ [19].

Recent studies demonstrated that IDE was up-regulated in a HSP-like fashion [20]. In

this study, we also examine the expression levels of IDE under the TC or HT

treatment. The results showed that H_2O_2 significantly decreased the expression level

of IDE and the heat treatment by TC remarkably recovered its expression level while

HT had notably weaker ability to regain the normal condition. Another important

protein quality control system which could degrade the Aβ peptide is the

ubiquitin-proteasome system (UPS).

Regulation of Aβ production by the UPS may have a role in AD pathogenesis and

therefore the UPS is regarded as potential therapeutic target for AD [21]. The

proteasome comprises a proteolytic 20S core particle capped by either one or two 19S

regulatory particles. In this study, the protein expression levels of 20S (PSMA3) and

19S (PSMC3) subunits were examined to verify the effect of thermal treatment on

proteasome expression. The results showed that both expressions of PSMC3 and

PSMA3 were weakened by H_2O_2 significantly. For the thermal treatment by TC, the

levels of PSMC3 and PSMA3 expressions were greatly recovered especially in the

PSMA3 level (return to 80% of the control value). For the HT treatment group, the

degree of recovery was much lower than TC treatment group in both PSMC3 and

PSMA3 expressions, while it had no significant difference compared to the

H_2O_2-treated group in PSMA3 protein expressions.

Fig 7. Effect of TC-HT on expressions of IDE and proteasome in SH-SY5Y cells.

(A) Western blot analysis of IDE and proteasome subunits (PSMC3 and PSMA3)

expressions. (B) Quantification of IDE, PSMC3, and PSMA3 expressions after H_2O_2,

TC+H_2O_2, or HT+H_2O_2 treatment. The expression levels were normalized to GAPDH.

Data represent the mean ± standard deviation (n=3). $^{***}P < 0.001$ and $^{**}P < 0.01$.

Effect of heat treatment on Akt-Nrf2/CREB signalling

pathways in H$_2$O$_2$-treated SH-SY5Y cells

We then examined the signalling pathways that could participate in the

neuroprotective mechanism of TC-HT. The Akt pathway is highly conserved in all

cells of higher eukaryotes and plays pivotal roles in cell survival and growth. Once

Akt is activated by phosphorylation, it positively regulates some transcription factors

to allow expression of prosurvival proteins. Therefore, in our study, we examined the

phosphorylation of Akt and the downstream transcription factors including cAMP

response element binding protein (CREB) and nuclear factor erythroid 2-related factor

2 (Nrf2) by western blot analysis. As shown in Fig 8, the expression of

phosphorylated Akt (p-Akt) was increased in H$_2$O$_2$-treated SH-SY5Y cells compared

to the untreated cells. The increased p-Akt level could be due to the stress response

induced by the oxidative stress of H$_2$O$_2$. Heat pretreatment further increased the p-Akt

levels in H$_2$O$_2$-treated SH-SY5Y cells, and the TC group had significantly higher

level of p-Akt than the HT group. We also examined the expression levels of p-CREB

and Nrf2. CREB is an important transcription factor that also triggers expression of

many prosurvival proteins, including Bcl-2 and brain derived neurotrophic factor

(BDNF). When CREB is phosphorylated, it starts to regulate downstream gene

expression. Besides, Nrf2 is also a key transcription factor that regulates expression of

antioxidant proteins, such as heme oxygenase-1 (HO-1). As shown in Fig 8, H_2O_2

treatment slightly influenced Nrf2 and p-CREB protein expression levels. In the

heat-treated groups, the TC treatment significantly increased the expressions of Nrf2

and p-CREB, while the HT treatment increased Nrf2 expression with less significant

level than TC treatment but in turn reduced the expression level of p-CREB. The

HO-1 protein is a heme-containing HSP which is crucial for neuroprotection against

oxidative stress. The result showed that the heat treatment significantly enhanced the

expression levels of HO-1 in TC and HT groups. Particularly, it is worth noting that

the TC treatment had a significant higher level of HO-1 than HT, which could account

for the antioxidant ability of the TC treatment in the ROS experiment.

Fig 8. Effect of TC-HT on Akt/Nrf2 and Akt/CREB signalling pathways and

related protein expressions. Western blot analysis of p-Akt, Nrf2, p-CREB, and

HO-1 proteins. Quantification of p-Akt, Nrf2, p-CREB, HO-1 expressions after H_2O_2,

TC+H_2O_2, or HT+H_2O_2 treatment. The expression level of p-Akt was normalized to

total Akt while other proteins were normalized to GAPDH. Data represent the mean $\pm$

standard deviation (n=3). [***]$P < 0.001$, [**]$P < 0.01$ and [*]$P < 0.05$.

Effect of LY294002 on the TC-HT activated Akt pathway and protective effect

In order to examine if the activation of the Akt pathway was involved in the

neuroprotective effect of TC treatment, we inhibited the phosphoinositide-3-kinase

(PI3K), the upstream kinase of Akt, by the inhibitor LY294002 [22] in the study. In

Fig 9A, we found that when H_2O_2 treatment reduced the viability of SH-SY5Y cells to

a little higher level, say, 61.1% of the control value, then TC treatment could greatly

increase the cell viability to ~94% of the control value under H_2O_2 stress. In the

presence of PI3K inhibitor, the p-Akt level was indeed reversed and the

neuroprotective effect of TC treatment against H_2O_2-induced cell apoptosis was

abrogated (Fig 9A and 9B), representing that the Akt pathway was involved to

mediate the neuroprotective effect of TC treatment against H_2O_2 toxicity. For the

downstream proteins of Akt, inhibition of the PI3K/Akt pathway was found to

significantly reduce the induction of Nrf2 by TC treatment, indicating that the

transcription factor Nrf2 was indeed activated by the Akt pathway. Collectively, these

data point out that TC treatment may exhibit neuroprotective effect via activation of

the Akt-Nrf2/CREB signalling pathways in the H_2O_2-treated SH-SY5Y neuron cells.

Fig 9. Effect of LY294002 on the neuroprotective effect of TC-HT and related protein expressions. (A) TC treatment conferred neuroprotective effect and significantly increased the cell viability of SH-SY5Y neuron cells under the H_2O_2 stress. The neuroprotective effect of TC treatment was abrogated by addition of the PI3K inhibitor LY294002. (B) Western blot analysis of p-Akt and Nrf2 proteins. The inhibitor LY294002 reversed the activated levels of p-Akt and Nrf2 induced by TC treatment. The expression level of p-Akt and Nrf2 was normalized to total Akt and GAPDH, respectively. We used the abbreviation "LY" to represent the PI3K inhibitor LY294002 in the figure. Data represent the mean ± standard deviation (n=3). $^{***}P < 0.001$.

Discussion

The study focused on investigating the neuroprotective effect of TC-HT on SH-SY5Y cells. It showed that subjection of the cells to heat at high and low temperatures alternately could produce significant protective effect against H_2O_2 and Aβ-induced apoptosis in SH-SY5Y human neural cells. The western blot results showed that TC-HT activated the PI3K/Akt signalling pathway and mediated the activation of Nrf2 and CREB proteins.

A human derived cell line SH-SY5Y has been extensively employed as general

in vitro model in evaluating neuronal damage or neurodegenerative diseases, such as

AD and PD [23, 24]. Therefore, the study used SH-SY5Y cells in the experiments to

determine the protective effect of TC-HT against H_2O_2 and Aβ-induced cytotoxicity.

Although the exact molecular mechanism of neurodegeneration pathogenesis is still

not clear, a common feature of these diseases is oxidative stress [25]. Many

researchers believe that oxidative stress may play a critical role in the etiology and

cause the exacerbation of disease progression. Cellular ROS are generated by both

extrinsic and intrinsic sources, with the former including ultraviolet, drugs, and

environmental toxins. Under normal physiological conditions, ROS generated

intrinsically from mitochondria and other enzymes are maintained at appropriate

levels by endogenous antioxidants. However, when mitochondria suffers from

decreased antioxidant defense or cell inflammation due to damage, excess ROS

production may occur. Since neuron cells are especially vulnerable to oxidative

damage due to their high oxygen demand and high polyunsaturated fatty acid contents

in membranes [26], the imbalance between ROS production and removal may cause

neuron damage or degeneration. A key cellular and molecular mechanism of several

neurodegenerative diseases, such as AD, PD, Huntington's disease (HD), amyotrophic

lateral sclerosis (ALS), is the accumulation of misfolded aggregation proteins in brain

[27]. The deposition of these misfolded protein aggregates can cause inflammation in

brain, inducing significant ROS production and leading to synaptic dysfunction,

neuronal apoptosis, and brain degeneration [28]. The composition of protein

aggregates is different among various neurodegenerative diseases. For AD, there are

two main kinds of protein aggregates, namely extracellular amyloid plaques deriving

from amyloid precursor protein and intracellular microtubule-associated protein tau,

known as neurofibrillary tangles [29]. With the complex pathogeneses of Aβ or tau

proteins still remaining largely unknown, several studies have proven that the

accumulation of Aβ increased oxidative stress and led to mitochondrial dysfunction

and DNA/RNA oxidation [30]. Matsuoka et al. showed that AD transgenic mouse

with Aβ accumulation can cause an increase in H_2O_2 level, suggesting that Aβ may

enhance oxidative stress in AD [31]. The oxidative stress can boost the

hyper-phosphorylation of tau protein and in turn facilitate the aggregation of Aβ [32].

Studies have also pointed out that overexpressed tau proteins make the cells more

vulnerable to oxidative stress [33]. Therefore, efforts have been underway to find

therapeutic strategies that can protect against oxidative stress and alleviate the

symptoms of neurodegenerative diseases. Among numerous ROS species, H_2O_2 can

be considered to be one of the most important ROS molecules involved in the

progression of AD. Many studies, therefore, used H_2O_2 to induce oxidative stress in

vitro, as a model in the investigation of the neuroprotective or neurodamage

mechanisms [34-36]. The study also found that H_2O_2 increased the ROS level in

SY-SY5Y cells and caused the cell death in a dose-dependent manner. Additionally,

Aβ was used to further investigate the neuroprotective effect of TC-HT in the AD

disease model. Both MTT and LDH release assays were employed in assessing the

neuroprotective effect of TC-HT. The results showed that TC-HT increased the cell

viability in the H_2O_2 and Aβ-treated neural cells. The detection of ROS level showed

that one possible mechanism of neuroprotective action of TC-HT exhibited the

antioxidant property, reducing the intracellular ROS level, perhaps due to the increase

of antioxidant enzymes in SH-SY5Y cells, induced by TC-HT. In the western blot

results, the study proved that such effect was through activation of Akt/Nrf2 pathway.

The Akt signaling pathway is a key regulator in the survival, proliferation, and

migration of cells, in response to extracellular signals [37]. Phosphorylation of Akt

further activates a set of downstream transcription factors, including Nrf2 and CREB,

which are considered to be mediators of neuroprotection by increasing the expression

of many antioxidant and prosurvival enzymes under oxidative stress. Nrf2 is one of

the most important transcription factors that regulate the expression of various

antioxidant proteins and maintain the redox state in mammalian cells [38, 39]. Among

various antioxidant proteins, HO-1 has been proven to be effective in promoting

antioxidant production and protection against neuronal injury [40]. The study found

that the physical stimulation, TC-HT, could activate the Akt/Nrf2 pathway and

upregulate the expression of the antioxidant protein HO-1, thereby reducing the ROS

generation by H_2O_2. Compared with the continuous heating of HT group, TC-HT

performed significantly better in activating Akt/Nrf2, HO-1 expression, and ROS

reduction, in agreement with the cell viability assay.

Another possible mechanism for neuroprotection of TC-HT may be related to the

activation of Akt/CREB pathway. CREB is the cellular transcription factor activated

by phosphorylation from Akt or other kinases, which triggers expression of

neurotrophins and antiapoptotic proteins. The critical role of CREB in neuronal

plasticity and long-term memory has also been well-documented [41]. In addition to

the neuroprotective effect of CREB by upregulating neurotrophins and antiapoptotic

proteins, recent studies have discovered that CREB also protects neurons via the

antioxidant pathways [42]. The downstream antioxidant genes include HO-1 and

manganese superoxide dismutase (MnSOD). Besides, CREB downregulation is

involved in the pathology of AD [43], and thus increasing the expression of CREB

has been considered a potential therapeutic strategy for AD [44]. In some

preconditioning experiments, CREB seemed to play a key role in the preconditioning

response which protected the cells from the subsequent stress [45]. In the study, it was

observed that TC-HT pretreatment significantly upregulated the activation of CREB

protein, compared with the control group in SH-SY5Y cells. The result further found

that HT significantly decreased the expression level of p-CREB compared with the

control cells. The decreased level of p-CREB in HT group indicates that HT may

exert too much heat stress and thus hinder the neuroprotective effect or even cause

damage to the cells. In the study, the result points out that the neuroprotective effect of

TC-HT could be attributed to the function of antioxidant protein HO-1 or other

prosurvival proteins. Further studies are necessary to identify the specific role of other

downstream neurotrophins and antiapoptotic or antioxidant proteins in the

neuroprotection effect.

Under stress amid a harsh environment or caused by external stimuli, cells or

whole organisms would trigger self-defense systems in response [46]. For example,

fever induced by viral or bacterial infections is a useful defense mechanism capable of

regulating the body temperature, as higher body temperature strengthens immune

cells and increases their ability to kill bacteria and viruses. It also inhibits the growth

of bacteria and viruses, thereby attaining anti-inflammatory effects. Depending on the

intensity and duration of stress, cells would activate cell repair or stress response to

adapt to the new environment or cause their death [47]. The stress response involves

the synthesis of some highly conserved proteins with protective function for cells [48].

For the thermal stress induced when cells or whole organisms are exposed to elevated

temperatures, they would respond by synthesizing an evolutionary highly conserved

family of proteins, known as HSPs [49]. The HS response is a universal phenomenon,

which appears in every tested organism. HSPs are named and classified, according to

their molecular weight. HSP70 (70 kilodaltons in size) is a typical group of HSPs,

which functions a molecular chaperone, to prevent the misfolding of nascent

polypeptide chains and facilitate refolding of misfolded proteins [50]. Many studies

have indicated that HSP70 sub-family is essential in protecting organisms from

various stresses [51]. The study also examined the effect of TC-HT and HT on the

expression of HSP70 and HSP105 [52], both of which have been reported to have

protective effects. The results showed that HT and TC-HT treatments enhanced the

expression of both HSPs significantly. However, the activation level of TC-HT was

only slightly higher than HT group, which could not explain the superior protective

effect of TC-HT compared to HT. Another novel HS-like protein, IDE, was identified

as the major Aβ degrading enzyme, and its protein expression level was found to

correlate negatively with the AD pathology [19]. Previous study had shown that IDE

expression was stress-inducible [20]. In the study, it was found that the TC treatment

greatly recovered the H_2O_2-induced IDE reduction, and the enhanced level was much

higher than HT treatment. Therefore, IDE could be one of the proteins participating in

the neuroprotective effect under Aβ administration.

The stress response enables cells to survive the stress-induced damage. Use of

cells' self-defense ability to adapt the stress environment or manipulation of the stress

response to increase the survival rate may be valuable in regenerative medicine or

disease curing. In fact, there have been examples involving preconditioning of cells

with non-lethal stress to minimize cell damage and improve the transplantation

outcome [53]. While there have been many experiments on thermotolerance and cell

survival [54], none have demonstrated the stress response in neuroprotection and

curation to the best of our knowledge. Besides, the challenge for the manipulation of

cellular stress response is how to fine-tune stress intensity and duration to suit specific

needs and prevent the stress-induced cell damage. The present study demonstrated a

delicate and efficient way to activate the stress response, leading to protection and

curation against H_2O_2 and Aβ-induced cytotoxicity on SH-SY5Y cells. The thermal

dosage was fine-tuned in a heat-and-cold cycling process, to maximize the protective

effect and minimize the heat stress-induced cell damage. Although traditional HS can

activate HSPs which may provide resistance to insults, such as hyperthermia, the

continuous exposure may cause cell damage or even death. Previous studies have

shown that the threshold for thermal damage is different among various tissue cells,

which have different levels of thermal sensibility [55]. For example, the skin cells can

withstand much higher thermal dosage (47°C for 20 min) than neural cells, because

the neural cells are more vulnerable. As the most thermally sensitive tissue, brain can

be damaged by low thermal dosage [55]. To attain optimal thermal dosage for various

tissues, the heating process should be divided into several parts, which is the basic

architecture of TC-HT.

In this study, the work presented an efficient and guaranteed way of controlling

the thermal dosage applied to cells. The novelty of the TC-HT strategy is periodically

interspersing the short period of cooling process in the continuous heating period. The

body temperature (37°C) was selected as the low temperature setting which mimics

the passive cooling process in the human body. Interestingly, our data demonstrated

that the application time of the low temperature period is a critical parameter to

determine the efficacy of neuroprotection for TC-HT. For too short cooling period, the

accumulation of the thermal dosage will cause damage to neural cells like HT. On the

other side, it was found that the TC-HT did not produce protective effect to the cells

when low temperature period was longer than 1 min (Fig 2D). It may be due to the

reason that the application of too long cooling period (> 1 min) could interrupt the

transmission of biochemical signals and protein expressions stimulated by thermal

stress, thus failing to achieve the neuroprotective effect. The results revealed that 35

sec cooling process at 37°C after 15 min heating at 42.5°C for 8 cycles produces the

optimum protective effect to SH-SY5Y cells. This protective effect employing TC-HT

was significantly much better than the continuous HT, which was due to the activation

of other thermal stress-associated proteins apart from HSPs and the prevention of heat

damage. From the LY294002 experiment, it was confirmed that the superior

protective effect resulted partially from the activation of Akt pathway. One fascinating

phenomenon is the antioxidant effect induced by TC-HT, which could be attributed to

the increased antioxidant protein HO-1. Besides, the Aβ-induced cytotoxicity was also

rescued in TC-HT treated cells, which points out the possibility that some

stress-induced proteins might have the ability to eliminate or refold the aggregated

protein or ease the symptom at least. The chaperone or proteasome systems are

possible candidates to be involved in this phenomenon [56]. Actually, previous studies

had shown that IDE will interact with 20S proteasome and modulate its proteolytic

activity [57]. Our results showed that TC treatment could not only enhance the

expression of IDE but also increase the proteasome expressions of 20S (PSMA3) and

19S (PSMC3) subunits. Although the examination was performed for only one

subunit of the 20S and 19S particles, respectively, one could not rule out the

possibility that TC treatment increased the abundance of proteasome complexes.

Further studies are needed to confirm if the proteolytic activity of 20S proteasome is

increased by the TC treatment and thus facilitates the degradation of Aβ polypeptides.

It is noteworthy that TC-HT applied after the Aβ administration gives better

curative effect than the pretreatment paradigm, which indicates that TC-HT not only

has the ability to prevent or protect the neurons against the Aβ-induced cytotoxicity

but also could have the potential to cure the disease. The possible explanation for the

superior effect of post-treatment could be that some biochemical signals and elevated

proteins could interact with the Aβ aggregation simultaneously, and these effects

could not maintain until the insult for the pretreated cells. Another advantage is that

the post-treatment paradigms for neuronal injury are also more typical in the clinical

condition. The present study employed PCR equipment to demonstrate the protective

effect of TC-HT in vitro. For in vivo or clinical applications, there are options of other

heating devices, such as high-intensity focused ultrasound (HIFU), which has been

widely used as a hyperthermal apparatus [58]. The thermal parameters can also be

fine-tuned by controlling the heating power and the size of heated volume to meet the

specific needs [59].

In summary, this is the first study on the manipulation of stress response by

TC-HT, which protects SH-SY5Y human neural cells from H_2O_2 and Aβ-induced

cytotoxicity. The results confirm that the physical stimulation by TC-HT provides

safer and much superior protective effect than continuous HS. The underlying

molecular mechanism for the protective effect is partially due to the activation of Akt

pathway and the function of some downstream proteins, such as Nrf2, CREB, and

HO-1. Further studies are necessary to confirm the involvement of other

stress-induced proteins activated by TC-HT in the neuroprotective effect or their

ability in degrading or disaggregating the aggregated proteins. It is also advisable for

coupling TC-HT with NDD drugs or natural compounds, in order to attain better

curative effect for NDD patients.

Funding

657 This work was supported by grants from Ministry of Science and Technology

658 (MOST 108-2112-M-002-016 and 105-2112-M-002-006-MY3 to CYC) and Ministry

659 of Education (MOE 106R880708 to CYC) of the Republic of China. The funders had

660 no role in study design, data collection and analysis, decision to publish, or

661 preparation of the manuscript.

Competing Interests

664 The authors have declared that no competing interests exist.

Acknowledgments

667   The authors would like to acknowledge the service provided by the Research

668 Core Facilities 3 Laboratory of the Department of Medical Research at National

669 Taiwan University hospital for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin, Chih-Yu Chao.

Formal analysis: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo-Lin, Chueh-Hsuan Lu,

Hao-Ping Hsu, Yi-Kun Sun, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin.

Writing – original draft: Wei-Ting Chen, Yu-Yi Kuo, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References:

1.   Kim GH, Kim JE, Rhie SJ, Yoon S. The role of oxidative stress in
neurodegenerative diseases. Exp Neurobiol. 2015; 24:325-340.

2. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing.
Nature. 2000; 408:239-247.

3. Liu Z, Zhou T, Ziegler AC, Dimitrion P, Zuo L. Oxidative stress in
neurodegenerative diseases: from molecular mechanisms to clinical applications.
Oxidative Med Cell Longev. 2017; 2017:2525967.

4. Bachurin SO, Bovina EV, Ustyugov AA. Drugs in clinical trials for Alzheimer's
disease: the major trends. Med Res Rev. 2017; 37:1186-1225.

5. Alghazwi M, Smid S, Musgrave I, Zhang W. In vitro studies of the
neuroprotective activities of astaxanthin and fucoxanthin against amyloid beta
(Aβ1-42) toxicity and aggregation. Neurochem Int. 2019; 124:215-224.

6. Hemming ML, Selkoe DJ. Amyloid beta-protein is degraded by cellular
angiotensin-converting enzyme (ACE) and elevated by an ACE inhibitor. J Biol Chem.
2005; 280:37644-37650.

7. Yang Y, Zhang X, Ye D, Laforest R, Williamson J, Liu Y, et al. Cavitation dose
painting for focused ultrasound-induced blood-brain barrier disruption. Sci Rep. 2019;
9:2840.

8. Habash RW, Bansal R, Krewski D, Alhafid HT. Thermal therapy, part 1: an
introduction to thermal therapy. Crit Rev Biomed Eng. 2006; 34:459-489.

9. Christophi C, Winkworth A, Muralihdaran V, Evans P. The treatment of
malignancy by hyperthermia. Surg Oncol. 1998; 7:83-90.

10. Schaaf L, Schwab M, Ulmer C, Heine S, Mürdter TE, Schmid JO, et al.
Hyperthermia synergizes with chemotherapy by inhibiting PARP1-dependent DNA
replication arrest. Cancer Res. 2016; 76:2868-2875.

11. Lord-Fontaine S, Averill-Bates DA. Heat shock inactivates cellular antioxidant
defenses against hydrogen peroxide: protection by glucose. Free Radic Biol Med.
2002; 32:752-765.

12. Tchouagué M, Grondin M, Glory A, Averill-Bates D. Heat shock induces the
cellular antioxidant defenses peroxiredoxin, glutathione and glucose 6-phosphate
dehydrogenase through Nrf2. Chem-Biol Interact. 2019; 310:108717.

13. Mustafi SB, Chakraborty PK, Dey RS, Raha S. Heat stress upregulates
chaperone heat shock protein 70 and antioxidant manganese superoxide dismutase
through reactive oxygen species (ROS), p38MAPK, and Akt. Cell Stress Chaperones.
2009; 14:579-589.

14. Hamblin MR. Shining light on the head: photobiomodulation for brain disorders.
BBA clinical. 2016; 6:113-124.

15. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal
therapy to synergistically enhance the anticancer effect of propolis on PANC-1 cells.
Int J Oncol. 2019; 55:617-628.

16. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal cycling-hyperthermia
in combination with polyphenols, epigallocatechin gallate and chlorogenic acid,
exerts synergistic anticancer effect against human pancreatic cancer PANC-1 cells.
PLoS One. 2019; 14:e0217676.

17. Sminia P, Zee JVD, Wondergem J, Haveman J. Effect of hyperthermia on the
central nervous system: a review. Int J Hyperthermia. 1994; 10:1-30.

18. Lin MT, Beal MF. Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature. 2006; 443:787-795.

19. Tundo GR, Sbardella D, Ciaccio C, Grasso G, Gioia M, Coletta A, et al. Multiple
functions of insulin-degrading enzyme: a metabolic crosslight? Crit Rev Biochem
Mol Biol. 2017; 52:554-582.

20. Tundo GR, Sbardella D, Ciaccio C, Bianculli A, Orlandi A, Desimio MG, et al.
Insulin-degrading enzyme (IDE): A novel heat shock-like protein. J Biol Chem. 2013;
288:2281-2289.

21. Gong B, Radulovic M, Figueiredo-Pereira ME, Cardozo C. The
ubiquitin-proteasome system: potential therapeutic targets for Alzheimer's disease and
spinal cord injury. Front Molec Neurosci. 2016; 9:4.

22. Sanchezmargalet V, Goldfine ID, Vlahos CJ, Sung CK. Role of
phosphatidylinositol-3-kinase in insulin receptor signaling: studies with inhibitor,
LY294002. Biochem Biophys Res Commun. 1994; 204:446-452.

23. Wu X, Kosaraju J, Zhou W, Tam KY. Neuroprotective effect of SLM, a novel
carbazole-based fluorophore, on SH-SY5Y cell model and 3xTg-AD mouse model of
Alzheimer's disease. ACS Chem Neurosci. 2017; 8:676-685.

24. Xicoy H, Wieringa B, Martens GJ. The SH-SY5Y cell line in Parkinson's disease
research: a systematic review. Mol Neurodegener. 2017; 12:10.

25. Barnham KJ, Masters CL, Bush AI. Neurodegenerative diseases and oxidative
stress. Nat Rev Drug Discov. 2004; 3:205-214.

26. Chen X, Guo C, Kong J. Oxidative stress in neurodegenerative diseases. Neural
Regen Res. 2012; 7:376-385.

27. Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. Nat
Med. 2004; 10:S10-S17.

28. Masters SL, O'Neill LAJ. Disease-associated amyloid and misfolded protein
aggregates activate the inflammasome. Trends Mol Med. 2011; 17:276-282.

29. Choi ML, Gandhi S. Crucial role of protein oligomerization in the pathogenesis
of Alzheimer's and Parkinson's diseases. FEBS J. 2018; 285:3631-3644.

30. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH.
Mitochondria are a direct site of $A\beta$ accumulation in Alzheimer's disease neurons:
implications for free radical generation and oxidative damage in disease progression.

Hum Mol Genet. 2006; 15:1437-1449.

31. Matsuoka Y, Picciano M, La Francois J, Duff K. Fibrillar β-amyloid evokes

oxidative damage in a transgenic mouse model of Alzheimer's disease. Neuroscience.

2001; 104:609-613.

32. Huang HC, Jiang ZF. Accumulated amyloid-β peptide and hyperphosphorylated

tau protein: relationship and links in Alzheimer's disease. J Alzheimers Dis. 2009;

16:15-27.

33. Zhao Y, Zhao B. Oxidative stress and the pathogenesis of Alzheimer's disease.

Oxidative Med Cell Longev. 2013; 2013:316523.

34. de Oliveira MR, de Bittencourt Brasil F, Fürstenau CR. Sulforaphane promotes

mitochondrial protection in SH-SY5Y cells exposed to hydrogen peroxide by an

Nrf2-dependent mechanism. Mol Neurobiol. 2017; 55:4777-4787.

35. Ismail N, Ismail M, Imam MU, Azmi NH, Fathy SF, Foo JB, et al. Mechanistic

basis for protection of differentiated SH-SY5Y cells by oryzanol-rich fraction against

hydrogen peroxide-induced neurotoxicity. BMC Complement Altern Med. 2014;

14:467.

36. Han SM, Kim JM, Park KK, Chang YC, Pak SC. Neuroprotective effects of

melittin on hydrogen peroxide-induced apoptotic cell death in neuroblastoma

SH-SY5Y cells. BMC Complement Altern Med. 2014; 14:286.

37. Jason S, Cui W. Proliferation, survival and metabolism: the role of

PI3K/AKT/mTOR signalling in pluripotency and cell fate determination.

Development. 2016; 143:3050-3060.

38. Huang Y, Li W, Su ZY, Kong ANT. The complexity of the Nrf2 pathway: beyond

the antioxidant response. J Nutr Biochem. 2015; 26:1401-1413.

39. Kubben N, Zhang W, Wang L, Voss TC, Yang J, Qu J, et al. Repression of the

antioxidant NRF2 pathway in premature aging. Cell. 2016; 165:1361-1374.

40. Wang Z, Guo S, Wang J, Shen Y, Zhang J, Wu Q. Nrf2/HO-1 mediates the

neuroprotective effect of mangiferin on early brain injury after subarachnoid

hemorrhage by attenuating mitochondria-related apoptosis and neuroinflammation.

Sci Rep. 2017; 7:11883.

41. Caracciolo L, Marosi M, Mazzitelli J, Latifi S, Sano Y, Galvan L, et al. CREB

controls cortical circuit plasticity and functional recovery after stroke. Nat Commun.

2018; 9:2250.

42. Sakamoto K, Karelina K, Obrietan K. CREB: a multifaceted regulator of

neuronal plasticity and protection. J Neurochem. 2011; 116:1-9.

43. Rosa E, Fahnestock M. CREB expression mediates amyloid β-induced basal

BDNF downregulation. Neurobiol Aging. 2015; 36:2406-2413.

44. Qin Z. Modulating nitric oxide signaling in the CNS for Alzheimer's disease

therapy. Future Med Chem. 2013; 5:1451-1468.

45. Baranova KA, Rybnikova EA, Churilova AV, Vetrovoy OV, Samoilov MO. The
adaptive role of the CREB and NF-κB neuronal transcription factors in post-stress
psychopathology models in rats. Neurochem J. 2014; 8:17-23.

46. Chovatiya R, Medzhitov R. Stress, Inflammation, and defense of homeostasis.
Mol Cell. 2014; 54:281-288.

47. Thomas MP, Lieberman J. Live or let die: posttranscriptional gene regulation in
cell stress and cell death. Immunol Rev. 2013; 253:237-252.

48. Blanco S, Bandiera R, Popis M, Hussain S, Lombard P, Aleksic J, et al. Stem cell
function and stress response are controlled by protein synthesis. Nature. 2016;
534:335-340.

49. Lindquist S. The heat-shock response. Annu Rev Biochem. 1986; 55:1151-1191.

50. Mosser DD, Caron AW, Bourget L, Meriin AB, Sherman MY, Morimoto RI, et al.
The chaperone function of hsp70 is required for protection against stress-induced
apoptosis. Mol Cell Biol. 2000; 20:7146-7159.

51. Yenari MA, Liu J, Zheng Z, Vexler ZS, Lee JE, Giffard RG. Antiapoptotic and

anti-inflammatory mechanisms of heat-shock protein protection. Ann NY Acad Sci.

2005; 1053:74-83.

52. Hatayama T, Yamagishi N, Minobe E, Sakai K. Role of hsp105 in protection
against stress-induced apoptosis in neuronal PC12 cells. Biochem Biophys Res
Commun. 2001; 288:528-534.

53. Matsumoto K, Honda K, Kobayashi N. Protective effect of heat preconditioning
of rat liver graft resulting in improved transplant survival. Transplantation. 2001;
71:862-868.

54. Mesihovic A, Iannacone R, Firon N, Fragkostefanakis S. Heat stress regimes for
the investigation of pollen thermotolerance in crop plants. Plant Reprod. 2016;
29:93-105.

55. Dewhirst MW, Viglianti BL, Lora-Michiels M, Hanson M, Hoopes PJ. Basic
principles of thermal dosimetry and thermal thresholds for tissue damage from
hyperthermia. Int J Hyperthermia. 2003; 19:267-294.

56. Kampinga HH, Bergink S. Heat shock proteins as potential targets for protective
strategies in neurodegeneration. Lancet Neurol. 2016; 15:748-759.

57. Sbardella D, Tundo GR, Coletta A, Marcoux J, Koufogeorgou EI, Ciaccio C, et
al. The insulin-degrading enzyme is an allosteric modulator of the 20S proteasome
and a potential competitor of the 19S. Cell Mol Life Sci. 2018; 75:3441-3456.

58. Frazier N, Payne A, de Bever J, Dillon C, Panda A, Subrahmanyam N, et al.
High intensity focused ultrasound hyperthermia for enhanced macromolecular

delivery. J Control Release. 2016; 241:186-193.

59. Partanen A, Tillander M, Yarmolenko PS, Wood BJ, Dreher MR, Kohler MO.

Reduction of peak acoustic pressure and shaping of heated region by use of multifoci

sonications in MR-guided high-intensity focused ultrasound mediated mild

hyperthermia. Med Phys. 2013; 40:013301.

Figure 1

A

B

C

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

B

Figure 8

Figure 9

A

Cell viability (%)

Ctrl H$_2$O$_2$ TC+H$_2$O$_2$ LY+TC+H$_2$O$_2$

***

B

p-Akt

Akt

GAPDH

Nrf2

GAPDH

p-Akt/Akt

***

Ctrl H$_2$O$_2$ TC+H$_2$O$_2$ LY+TC+H$_2$O$_2$

Nrf2/GAPDH

***

Ctrl H$_2$O$_2$ TC+H$_2$O$_2$ LY+TC+H$_2$O$_2$