Thermal cycling protects SH-SY5Y cells against hydrogen peroxide and β-amyloid-induced cell injury through stress response mechanisms involving Akt pathway

Wei-Ting Chen[1,2,¶], Yu-Yi Kuo[1,2,¶], Guan-Bo Lin[1,2,¶], Chueh-Hsuan Lu[1,2], Hao-Ping Hsu[3], Yi-Kun Sun[2], and Chih-Yu Chao[1,2,4,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Department of Medicine, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[4] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

¶ These authors contributed equally to this work.

Abstract

Neurodegenerative diseases (NDDs) are becoming a major threat to public health, according to the World Health Organization (WHO). The most common form of NDDs is Alzheimer's disease (AD), boasting 60-70% share. Excessive aggregation of β-amyloid protein (Aβ) and neurofibrillary tangles has been deemed the major cause for the pathogenesis of AD. A growing number of evidences from studies, however, have suggested that reactive oxygen species (ROS) also play a key role in the onset and progression of AD. Although scientists have had some understanding of the pathogenesis of AD, the disease still cannot be cured, with existing treatment only capable of providing a temporary relief at best, partly due to the obstacle of blood-brain barrier (BBB). The study was aimed to ascertain the neuroprotective effect of thermal cycle hyperthermia (TC-HT) against hydrogen peroxide (H_2O_2) and Aβ-induced cytotoxicity in SH-SY5Y cells. Treating cells with this physical stimulation beforehand significantly improved the cell viability and decreased the ROS content. The underlying mechanisms were confirmed to be due to the activation of Akt pathway and the downstream antioxidant and prosurvival proteins. The findings manifest significant potential of TC-HT in neuroprotection, via inhibition of oxidative stress and cell apoptosis. It is believed that coupled with the use of drugs or natural compounds, this methodology can be even more effective in treating NDDs.

Introduction

According to the World Health Organization (WHO), the number of dementia-induced deaths more than doubled between 2000 and 2016, making it the 5th leading cause for deaths worldwide in 2016, up from 14th place in 2000. Among the various forms of dementia such as Alzheimer's disease (AD) and Parkinson's disease (PD), AD is the most common one, accounting for 60–70% of the cases. Typically, AD exhibits such features as deposition of cortical plaques caused by excessive aggregation of β-amyloid protein (Aβ) and neurofibrillary tangles, and progressive brain degeneration and deterioration of cognitive function among elderly people. Although the exact mechanism of AD pathogenesis remains unknown, it is believed that oxidative stress and activation of free radicals, induced by Aβ aggregation, play a key role in AD pathogenesis [1].

Reactive oxygen species (ROS) are reactive chemical species containing oxygen, which are generated as natural byproduct of oxygen metabolism. ROS play important roles in cell signalling and homeostasis, and their concentrations in cells are subtlety regulated by various antioxidant compounds and enzymes. However, with cells under continuing environmental stress (e.g. ultraviolet, inflammatory cytokines, or environmental toxins), the imbalance between prooxidants and antioxidants may cause

chronic oxidative stress. Accumulation of ROS may cause cell death, accelerate cell

ageing, or induce age-related diseases [2]. More and more research evidences have

suggested that ROS plays a central role in the onset and progression of AD [3].

Therefore, the protection of neural cells against oxidative damage may be a potential

strategy to treat AD. Several in vitro or in vivo studies have explored the function of

antioxidant and antiapoptotic drugs in ameliorating AD [4-6] but the approach is time-

consuming and costly, plus the safety concern, which limits the use of these drugs in

AD treatment. Moreover, the blood-brain barrier (BBB) dampens the efficacy of these

drugs, since over 98% of small molecule drugs and ~100% of large molecule drugs can

not pass the BBB [7]. Therefore, a non-drug treatment may be more suited to AD

patients.

The study employed a special thermal therapy, applied at high and low

temperatures alternately to achieve an effect similar to antioxidant and antiapoptotic

drugs. In our previous study, we used this novel method, thermal cycle hyperthermia

(TC-HT) for the treatment of pancreatic cancer, and found that this physical stimulation

greatly improved the anticancer effect of propolis and polyphenols on PANC-1 cells

without the heat-induced side effect [8, 9]. Traditional heat shock (HS) or hyperthermia

(HT) employs continuous heating to achieve curative effect, but it is likely to cause

neuron damage since damage to the central nervous system occurs within few minutes

of exposure to 42°C [10]. Nevertheless, the feature of TC-HT is that it can avoid the

damage caused by HT, which is crucial for neuroprotection.

In this study, we applied the TC-HT strategy to human neural cell line SH-SY5Y,

which has been extensively used in research on neurodegenerative damage in vitro, and

examined the prosurvival effect of TC-HT on preventing oxidative damage induced by

hydrogen peroxide (H_2O_2) and Aβ. The results showed that subjection of the cells to

heat at high and low temperatures alternately beforehand ameliorated the H_2O_2 and Aβ-

induced cytotoxicity in SH-SY5Y cells significantly. It was found that TC-HT not only

performed superior protective effect than the traditional HS but also avoided thermal

damage caused by continuous heating. Examination of the underlying mechanism also

showed that TC-HT could activate specific neuroprotective proteins. These findings

indicate that TC-HT is a promising thermal therapy, which sheds light on novel

treatment for AD or other NDDs.

Materials and methods

Cell culture and treatment

The human neuroblastoma SH-SY5Y cells purchased from Bioresource Collection and

Research Center (Hsinchu, Taiwan) were maintained in MEM/F-12 mixture containing

10% fetal bovine serum (HyClone; GE Healthcare Life Sciences) and 1% penicillin-

streptomycin, supplemented with 1mM sodium pyruvate and 0.1 mM non-essential

amino acids. The cells were cultured at 37℃ in a humidified incubator composed of

5% CO_2. SH-SY5Y cells were seeded in 96-well plates overnight. Cells were pretreated

with HT or TC-HT by the Thermal Cycler (Applied Biosystems; Thermo Fisher

Scientific, Inc.) relative to the control at room temperature (RT). After the treatment,

the cells were maintained in a 37℃ incubator for 4 h. Subsequently, the pretreated cells

were induced to undergo apoptosis by adding H_2O_2 to the culture medium. Cell viability

was measured 24 h after the H_2O_2 treatment. As for AD disease model, the cells were

treated with 25 or 50 µM $A\beta_{25-35}$ protein solution. The Aβ stock solution was prepared

by solubilizing the $A\beta_{25-35}$ peptide (Sigma-Aldrich; Merck KGaA) in sterile deionized

water to a concentration of 1 mM and then incubated at 37℃ for 3 days to allow self-

aggregation before treatment. The TC-HT therapy was applied before or after the Aβ

treatment, and the cell viability was determined 4 days after the treatment. For the Akt

inhibitor experiment, 25 µM LY294002 (Cell Signaling Technology, Inc.) was treated

1 h before the TC-HT.

Cell Viability Assay

The   cell   viability   was   determined   by   3-(4,5-dimethylthiazol-2-yl)-2,5-

diphenyltetrazolium bromide (MTT) (Sigma-Aldrich; Merck KGaA) assay. In brief, the

culture medium was replaced with MTT solution (0.5 mg/mL in MEM/F12) and

incubated at 37℃ for 4 h. Equal volume of the solubilizing buffer (0.01 M HCl and

10% SDS) was added to dissolve the formazan crystals. The 96-well plates were

analyzed on a FLUOstar OPTIMA microplate reader (BMG Labtech, Ltd.) at 570 nm,

and background absorbance at 690 nm was subtracted.

Lactate dehydrogenase (LDH) assay

The assay measures a stable enzyme LDH, which is released into the cell culture

medium when cell membranes are damaged. LDH levels were measured using a

Cytoscan LDH cytotoxicity assay kit (G-Biosciences; Geno Technology Inc.) according

to the manufacturer's instructions. Briefly, the cells were seeded in 96-well plates and

then incubated with H_2O_2 with or without thermal cycle pretreatment. For analysis, 50

µL culture mediums from all wells were transferred to a new 96-well plate, and 50 µL

of reaction mixtures were added to each well and incubated at 37℃ for 20 min. The

absorbance was measured at 490 nm using a microplate reader (BMG Labtech, Ltd.).

ROS level detection

ROS levels of cells were detected using the fluorescent dye dihydroethidium (DHE)

(Sigma-Aldrich; Merck KGaA). Cells pretreated with TC-HT or HT at 42.5℃ high

temperature setting for 8 cycles or 2 h, respectively, were challenged with H_2O_2. After

the treatment of H_2O_2 for 24 h, cells were washed with PBS, and then incubated with 5

µM DHE dye for 20 min at 37℃ in the dark. The fluorescence intensity emitted by

DHE was measured by flow cytometry in the PE channel, and ROS levels were

expressed as mean fluorescence intensity for comparison.

Mitochondrial membrane potential (MMP) measurement

The MMP was determined by flow cytometry using 3,3'-dihexyloxacarbocyanine

iodide [$DiOC_6(3)$] (Enzo Life Sciences International Inc.). $DiOC_6(3)$ is a lipophilic

cationic fluorescent dye which allows estimation of the percentage of cells with low

MMP. Cells were harvested and suspended at a density of 1 x 10^6 cells/mL in 1 µM

$DiOC_6(3)$ dye working solution. After incubation at 37℃ for 15 min, $DiOC_6(3)$

intensity was analyzed by flow cytometry in the FITC channel.

Western blot analysis

The protein expression levels of SH-SY5Y cells were investigated by western blot

analysis. Cells treated with TC-HT, HT, H_2O_2, or Aβ were harvested and lysed in RIPA

lysis butter (EMD Millipore). After centrifugation and supernatant collection, equal

amounts of proteins were resolved by 10% SDS-PAGE and transferred to

polyvinylidene fluoride membranes. After drying for 1 h at RT, the membranes were

probed with p-Akt, Nrf2, p-CREB (Cell Signaling Technology, Inc.), HO-1 (Enzo Life

Sciences International Inc.), and GAPDH (GeneTex, Inc.) antibodies overnight at 4˚C.

The washed membranes were then incubated with horseradish peroxidase-conjugated

goat anti-rabbit secondary antibodies (Jackson ImmunoResearch Laboratories, Inc.).

Immunoreactivity was visualized with an enhanced chemiluminescence substrate

(Advansta, Inc.) and detected with the Amersham Imager 600 imaging system (GE

Healthcare Life Sciences). The images were analyzed with Image Lab software (Bio-

Rad Laboratories, Inc.).

Statistical analysis

172 The results were presented as the mean ± standard deviation. Statistical analyses using

173 one-way analysis of variance (ANOVA) followed by Tukey's post-hot test were

174 performed by OriginPro 2015 software (OriginLab). P-value < 0.05 was considered to

175 indicate a statistically significant difference.

Results

In vitro-applied TC-HT

We applied the thermal cycle (TC) treatment to SH-SY5Y cells by a modified polymerase chain reaction (PCR) equipment (Fig 1C). Briefly, in our design, some protruding parts of the PCR machine and plastic well were milled so that the bottom of the well can touch the heat sink tightly. The basic TC settings were schemed as Fig 1A, where the temperature was elevated to the desired high temperature and sustained for a period of time followed by a cooling process at body temperature. It's worth noting that the adjustment of TC parameters (temperature and time) is based on the needs and objectives of the experiments. In our experiments, the TC parameters were set to high temperatures of 42.5℃ or 41.5℃ for 15 min, then lowered to 37℃ and lasted for 35 sec. This cycle was repeated 8 or 12 times. The HT group maintains the same high temperature setting throughout the treatment in a continuous manner without a break. To determine the actual temperatures of the cells, a needle thermocouple located in the bottom of the well was used to monitor the temperatures. Fig 1B represents the actual temperature measured by the thermocouple every 20 sec. The actual cycle temperature sensed by the cells was ~42-40.3℃ for the setting of 42.5-37℃ and ~40.9-39.4℃ for 41.5-37℃ setting, as shown in Fig 1B.

Fig 1. The setup of TC-HT. (A) Schematic representation of TC parameter settings.

198   (B) Measurement of the cell temperature during the TC treatment by the thermocouple.

199   (C) Image of the TC controller setup.

201   Effect of TC-HT on H_2O_2 and Aβ-induced cytotoxicity in SH-
202   SY5Y cells

We applied H_2O_2 to SH-SY5Y cells as an oxidative stress and the cell viability

was assessed by MTT assay. As shown in Fig 2A, treatment of H_2O_2 on SH-SY5Y cells

for 24 h induced significant decrease of cell viability in a dose-dependent manner.

Treatment of SH-SY5Y cells with 450 μM H_2O_2 for 24 h reduced the cell viability to

52.9% compared to control cells, which was used in the following experiments. To

investigate whether TC-HT or HT produced protection and increased cell survival

against the oxidative stress induced by H_2O_2, the SH-SY5Y cells were pretreated with

TC-HT or HT in the experiments. Two high temperature parameters (41.5℃ or 42.5℃)

were applied for 2 h or 3 h in the HT group, and 8 or 12 cycles (for 15 min per cycle)

in the TC group to make the total thermal doses of the HT and TC groups equal. After

the heat pretreatment, the treated SH-SY5Y cells were kept in the incubator at 37℃ for

4 h. They were then challenged with H_2O_2 to generate the oxidative stress, and the cell

viability was determined by MTT assay at 24 h after the H_2O_2 treatment. The results

found that H_2O_2 treatment significantly reduced the viability of SH-SY5Y cells to

53.7% of the control value, and showed that the heat pretreatment conferred protective

effect and increased the cell viability. At 41.5℃ for 2 h and 8 cycles, HT and TC

treatments increased the cell viability to 61.9% and 68.5% of the control value,

respectively (Fig 2B). As we increased the high temperature setting to 42.5℃ while

maintaining the same heating time, TC pretreatment greatly increased the cell viability

to 83% of the control value under H_2O_2 stress, while the HT group only exhibited a

slight increase (Fig 2C). When we further increased the heating time to 3 h, the

treatment of HT alone was cytotoxic to the cells, reducing the cell viability to 75.4%,

and the viability was even lower after the H_2O_2 treatment (Fig 2C). Interestingly, the

TC treatment with the same total thermal dose (12 cycles) had only a slight influence

on the cell viability, and it still conferred significant neuroprotective effect to the SH-

SY5Y cells under H_2O_2 stress.

In particular, the study examined the influence of the cooling process on the

neuroprotective effect. As shown in Fig 2D, for low temperature period longer than 1

min, the TC pretreatment did not produce protective effect to the cells, indicating that

the low temperature period is a critical parameter to be determined for effective

neuroprotection. On the other hand, the LDH release rate was measured and found to

decrease significantly in the TC pretreatment group compared to the H_2O_2-treated group

(Fig 2E), further confirming its protective role in maintaining the cell membrane

integrity when cell was under oxidative stress. For AD disease model, Aβ treatment

was used to reduce the viability of SH-SY5Y cells due to the cytotoxicity of the

aggregated Aβ (Fig 3A). Our results showed that the TC treatment after Aβ

administration greatly improved the viability of SH-SY5Y cells (Fig 3B) for nearly

25%, indicating the curative effect of TC in AD disease model in vitro. The most

effective high temperature setting was 42.3℃ for 8 cycles in the Aβ-treated cells.

Interestingly, the curative effect of TC pretreatment was less pronounced than that of

TC post-treatment (treated after Aβ administration), indicating the time point of TC

application has great impact on the curative effect. Moreover, there were no curative

effects by both HT pretreatment and post-treatment on SH-SY5Y cells, whose viability

levels were even slightly lower than that of the Aβ-treated cells (Fig 3B). Furthermore,

the light microscopy images showed that the integrity of the cells was destructed by Aβ,

and the TC treatment caused the protective effect to retain the cell morphology, as

shown in Fig 3C.

To conclude the result of the H_2O_2-induced oxidative stress on SH-SY5Y cells, the

protective effect of TC pretreatment was more pronounced than the HT preconditioning.

And the TC pretreatment at 42.5℃ produced much superior protection effect than at

41.5℃. Noteworthily, the HT pretreatment at 42.5℃ for 3 h decreased the cell viability

which indicates that the continuous heating may induce cell damage. The protective

effect of TC was most remarkable at ~42.5℃ for 8 cycles. Therefore, we chose this

parameter to evaluate the possible protective mechanism of TC-HT in the following

experiments.

Fig 2. Effect of TC-HT on H₂O₂-induced cytotoxicity in SH-SY5Y cells. (A) Dose-

response curve of SH-SY5Y cells treated with different concentrations of H_2O_2 for 24

h. (B) SH-SY5Y cells were pretreated at 41.5℃ temperature setting and challenged

with or without 450 µM H_2O_2. The cell viability was measured by MTT assay at 24 h

after the H_2O_2 treatment. (C) SH-SY5Y cells were pretreated at 42.5℃ temperature

setting with different thermal dosages and challenged with or without 450 µM H_2O_2.

The cell viability was measured by MTT assay at 24h after the H_2O_2 treatment. (D)

Comparison of the neuroprotective effect under different low temperature period

settings. (E) The LDH release was measured to confirm the neuroprotective effect of

TC treatment. Data represent the mean ± standard deviation (n=3). [***]P < 0.001.

Fig 3. Effect of TC-HT on Aβ-induced cytotoxicity in SH-SY5Y cells. (A) The cell

viability of SH-SY5Y cells treated with 25 or 50 µM Aβ for 4 days. (B) The TC or HT

pretreatment and post-treatment were applied to the cells before or after 50 µM Aβ

administration, and the cell viability was measured by MTT assay 4 days after treatment.

(C) Representative light microscopy images of SH-SY5Y cells after treatment. The

integrity of the cells was destructed by Aβ, and the TC post-treatment caused the

protective effect and retained the cell morphology. Scale bar = 100 µm. Data represent

the mean ± standard deviation (n=3). [***]$P < 0.001$.

TC-HT attenuates H_2O_2-induced ROS generation

The onset and progression of AD has been reported to be associated with ROS. We

further studied whether the H_2O_2-induced intracellular ROS production could be

attenuated by the TC pretreatment. The intracellular ROS levels were detected using

the fluorescent dye DHE (Fig 4A). Results showed that after exposure to 450 µM H_2O_2

for 24 h, ROS level was increased to 160% of the control value. When the SH-SY5Y

cells were pretreated with TC-HT, the intracellular ROS level was significantly

ameliorated (Fig 4B). Compared to the TC group, the ROS level was not changed in

the HT group.

Fig 4. Effect of TC-HT on H_2O_2-induced ROS generation in SH-SY5Y cells. (A)

ROS level was measured by flow cytometry with DHE fluorescent dye. (B)

Quantification of the ROS levels after H_2O_2, TC+H_2O_2, or HT+H_2O_2 treatment. Data

represent the mean ± standard deviation (n=3). [**]$P < 0.01$.

TC-HT attenuates MMP loss in H_2O_2-treated cells

Many lines of evidence have suggested that mitochondria contribute to ageing-related

neurodegenerative diseases through the accumulation of net cytosolic ROS production

[11]. The levels of MMP are kept relatively stable in healthy cells. Apoptotic signals

are usually initiated by the disruption of normal mitochondrial function, especially the

collapse of MMP. To study the mechanism why H_2O_2-induced apoptosis is alleviated

by TC-HT pretreatment, the MMP is further analyzed by the flow cytometry with

$DiOC_6(3)$ fluorescent dye in the study. As shown in Fig 5A, the oxidative stress induced

by H_2O_2 caused mitochondrial dysfunction, decreasing the MMP and therefore

decreased the intensity of the $DiOC_6(3)$ fluorescent signal. From the quantification

results (Fig 5B), it was found that the percentage of cells with decreased MMP

drastically increased after the H_2O_2 treatment. In contrast to the group of H_2O_2 alone,

the TC-HT pretreatment noticeably suppressed the H_2O_2-induced dissipation of MMP,

so the cells with decreased MMP reduced significantly, which was attributed to the

preservation of the mitochondrial function and thus caused the blockage of the

apoptotic signaling transduction.

Fig 5. Effect of TC-HT on H_2O_2-induced MMP reduction in SH-SY5Y cells. (A)

MMP was analyzed by flow cytometry with $DiOC_6(3)$ fluorescent dye. (B)

Quantification of the cells with decreased MMP after H_2O_2, TC+H_2O_2, or HT+H_2O_2

treatment. Data represent the mean ± standard deviation (n=3). [***]$P < 0.001$.

Effect of heat treatment on heat shock protein (HSP) expressions in SH-SY5Y cells

It has been well known that HT triggers expression of HSPs so that cells are able to

protect themselves from stress. To investigate whether the protective effect of TC-HT

was related to the HSPs, we examined the protein expressions of HSP70 and HSP105,

both of which were reported to have protective effects. As shown in Fig 6, both HSP

expressions were enhanced significantly after HT and TC treatments. The activation

levels of TC, however, was only slightly higher than HT without significant difference.

This phenomenon points out that although TC treatment indeed protects SH-SY5Y cells

more from the oxidative stress of H_2O_2, the main mechanism could be apart from the

HSPs. Therefore, other heat-activated stress response proteins should be considered in

the experiments.

Fig 6. Effect of TC-HT on expressions of HSPs in SH-SY5Y cells. (A) Western blot

analysis of HSP expressions. (B) Quantification of HSP70 and HSP105 expressions

after H_2O_2, TC+H_2O_2, or HT+H_2O_2 treatment. The expression levels were normalized

to GAPDH. Data represent the mean ± standard deviation (n=3). [***]P < 0.001.

Effect of heat treatment on Akt-Nrf2/CREB signalling pathways in H_2O_2-treated SH-SY5Y cells

We then examined the signalling pathways that could participate in the neuroprotective

mechanism of TC-HT. The Akt pathway is highly conserved in all cells of higher

eukaryotes and plays pivotal roles in cell survival and growth. Once Akt is activated by

phosphorylation, it positively regulates some transcription factors to allow expression

of prosurvival proteins. Therefore, in our study, we examined the phosphorylation of

Akt and the downstream transcription factors including cAMP response element

binding protein (CREB) and nuclear factor erythroid 2-related factor 2 (Nrf2) by

western blot analysis. As shown in Fig 7, the expression of phosphorylated Akt (p-Akt)

was increased in H_2O_2-treated SH-SY5Y cells compared to the untreated cells. The

increased p-Akt level could be due to the stress response induced by the oxidative stress

of H_2O_2. Heat pretreatment further increased the p-Akt levels in H_2O_2-treated SH-SY5Y

cells, and the TC group had significantly higher level of p-Akt than the HT group. We

also examined the expression levels of p-CREB and Nrf2. CREB is an important

transcription factor that also triggers expression of many prosurvival proteins, including

Bcl-2 and brain derived neurotrophic factor (BDNF). When CREB is phosphorylated,

it starts to regulate downstream gene expression. Besides, Nrf2 is also a key

transcription factor that regulates expression of antioxidant proteins, such as heme

oxygenase-1 (HO-1). As shown in Fig 7, H_2O_2 treatment slightly influenced Nrf2 and

p-CREB protein expression levels. In the heat-treated groups, the TC treatment

significantly increased the expressions of Nrf2 and p-CREB, while the HT treatment

increased Nrf2 expression with less significant level than TC treatment but in turn

reduced the expression level of p-CREB. The HO-1 protein is a heme-containing HSP

which is crucial for neuroprotection against oxidative stress. The result showed that the

heat treatment significantly enhanced the expression levels of HO-1 in TC and HT

groups. Particularly, it is worth noting that the TC treatment had a significant higher

level of HO-1 than HT, which could account for the antioxidant ability of the TC

treatment in the ROS experiment.

Fig 7. Effect of TC-HT on Akt/Nrf2 and Akt/CREB signalling pathways and

related protein expressions. Western blot analysis of p-Akt, Nrf2, p-CREB, and HO-

1 proteins. Quantification of p-Akt, Nrf2, p-CREB, HO-1 expressions after H_2O_2,

TC+H_2O_2, or HT+H_2O_2 treatment. The expression levels were normalized to GAPDH.

Data represent the mean ± standard deviation (n=3). [***]$P < 0.001$, [**]$P < 0.01$ and [*]$P <$

0.05.

Effect of LY294002 on the TC-HT activated Akt pathway and protective effect

In order to examine if the activation of the Akt pathway was involved in the

neuroprotective effect of TC treatment, we inhibited the phosphoinositide-3-kinase

(PI3K), the upstream kinase of Akt, by the inhibitor LY294002 [12] in the study. In Fig

8A, we found that when H_2O_2 treatment reduced the viability of SH-SY5Y cells to a

little higher level, say, 61.1% of the control value, then TC treatment could greatly

increase the cell viability to ~94% of the control value under H_2O_2 stress. In the

presence of PI3K inhibitor, the p-Akt level was indeed reversed and the neuroprotective

effect of TC treatment against H_2O_2-induced cell apoptosis was abrogated (Fig 8A and

8B), representing that the Akt pathway was involved to mediate the neuroprotective

effect of TC treatment against H_2O_2 toxicity. For the downstream proteins of Akt,

inhibition of the PI3K/Akt pathway was found to significantly reduce the induction of

Nrf2 by TC treatment, indicating that the transcription factor Nrf2 was indeed activated

by the Akt pathway. Collectively, these data point out that TC treatment may exhibit

neuroprotective effect via activation of the Akt-Nrf2/CREB signalling pathways in the

H_2O_2-treated SH-SY5Y neuron cells.

Fig 8. Effect of LY294002 on the neuroprotective effect of TC-HT and related protein expressions. (A) TC treatment conferred neuroprotective effect and significantly increased the cell viability of SH-SY5Y neuron cells under the H_2O_2 stress. The neuroprotective effect of TC treatment was abrogated by addition of the PI3K inhibitor LY294002. (B) Western blot analysis of p-Akt and Nrf2 proteins. The inhibitor LY294002 reversed the activated levels of p-Akt and Nrf2 induced by TC treatment. The expression levels were normalized to GAPDH. We used the abbreviation "LY" to represent the PI3K inhibitor LY294002 in the figure. Data represent the mean ± standard deviation (n=3). [***]$P < 0.001$.

Discussion

The study focused on investigating the neuroprotective effect of TC-HT on SH-SY5Y cells. It showed that subjection of the cells to heat at high and low temperatures alternately could produce significant protective effect against H_2O_2 and Aβ-induced apoptosis in SH-SY5Y human neural cells. The western blot results showed that TC-HT activated the PI3K/Akt signalling pathway and mediated the activation of Nrf2 and CREB proteins.

A human derived cell line SH-SY5Y has been extensively employed as general in

vitro model in evaluating neuronal damage or neurodegenerative diseases, such as AD

and PD [13, 14]. Therefore, the study used SH-SY5Y cells in the experiments to

determine the protective effect of TC-HT against H_2O_2 and Aβ-induced cytotoxicity.

Although the exact molecular mechanism of neurodegeneration pathogenesis is still not

clear, a common feature of these diseases is oxidative stress [15]. Many researchers

believe that oxidative stress may play a critical role in the etiology and cause the

exacerbation of disease progression. Cellular ROS are generated by both extrinsic and

intrinsic sources, with the former including ultraviolet, drugs, and environmental toxins.

Under normal physiological conditions, ROS generated intrinsically from mitochondria

and other enzymes are maintained at appropriate levels by endogenous antioxidants.

However, when mitochondria suffers from decreased antioxidant defense or cell

inflammation due to damage, excess ROS production may occur. Since neuron cells are

especially vulnerable to oxidative damage due to their high oxygen demand and high

polyunsaturated fatty acid contents in membranes [16], the imbalance between ROS

production and removal may cause neuron damage or degeneration. A key cellular and

molecular mechanism of several neurodegenerative diseases, such as AD, PD,

Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), is the accumulation of

misfolded aggregation proteins in brain [17]. The deposition of these misfolded protein

aggregates can cause inflammation in brain, inducing significant ROS production and

leading to synaptic dysfunction, neuronal apoptosis, and brain degeneration [18]. The

composition of protein aggregates is different among various neurodegenerative

diseases. For AD, there are two main kinds of protein aggregates, namely extracellular

amyloid plaques deriving from amyloid precursor protein and intracellular

microtubule-associated protein tau, known as neurofibrillary tangles [19]. With the

complex pathogeneses of Aβ or tau proteins still remaining largely unknown, several

studies have proven that the accumulation of Aβ increased oxidative stress and led to

mitochondrial dysfunction and DNA/RNA oxidation [20]. Matsuoka et al. showed that

AD transgenic mouse with Aβ accumulation can cause an increase in H_2O_2 level,

suggesting that Aβ may enhance oxidative stress in AD [21]. The oxidative stress can

boost the hyper-phosphorylation of tau protein and in turn facilitate the aggregation of

Aβ [22]. Studies have also pointed out that overexpressed tau proteins make the cells

more vulnerable to oxidative stress [23]. Therefore, efforts have been underway to find

therapeutic strategies that can protect against oxidative stress and alleviate the

symptoms of neurodegenerative diseases. Among numerous ROS species, H_2O_2 can be

considered to be one of the most important ROS molecules involved in the progression

of AD. Many studies, therefore, used H_2O_2 to induce oxidative stress in vitro, as a model

in the investigation of the neuroprotective or neurodamage mechanisms [24-26]. The

study also found that H_2O_2 increased the ROS level in SY-SY5Y cells and caused the

cell death in a dose-dependent manner. Additionally, Aβ was used to further investigate

the neuroprotective effect of TC-HT in the AD disease model. Both MTT and LDH

release assays were employed in assessing the neuroprotective effect of TC-HT. The

results showed that TC-HT increased the cell viability in the H_2O_2 and Aβ-treated

neural cells. The detection of ROS level showed that one possible mechanism of

neuroprotective action of TC-HT exhibited the antioxidant property, reducing the

intracellular ROS level, perhaps due to the increase of antioxidant enzymes in SH-

SY5Y cells, induced by TC-HT. In the western blot results, the study proved that such

effect was through activation of Akt/Nrf2 pathway.

The Akt signaling pathway is a key regulator in the survival, proliferation, and

migration of cells, in response to extracellular signals [27]. Phosphorylation of Akt

further activates a set of downstream transcription factors, including Nrf2 and CREB,

which are considered to be mediators of neuroprotection by increasing the expression

of many antioxidant and prosurvival enzymes under oxidative stress. Nrf2 is one of the

most important transcription factors that regulate the expression of various antioxidant

proteins and maintain the redox state in mammalian cells [28, 29]. Among various

antioxidant proteins, HO-1 has been proven to be effective in promoting antioxidant

production and protection against neuronal injury [30]. The study found that the

physical stimulation, TC-HT, could activate the Akt/Nrf2 pathway and upregulate the

expression of the antioxidant protein HO-1, thereby reducing the ROS generation by

H_2O_2. Compared with the continuous heating of HT group, TC-HT performed

significantly better in activating Akt/Nrf2, HO-1 expression, and ROS reduction, in

agreement with the cell viability assay.

Another possible mechanism for neuroprotection of TC-HT may be related to the

activation of Akt/CREB pathway. CREB is the cellular transcription factor activated by

phosphorylation from Akt or other kinases, which triggers expression of neurotrophins

and antiapoptotic proteins. The critical role of CREB in neuronal plasticity and long-

term memory has also been well-documented [31]. In addition to the neuroprotective

effect of CREB by upregulating neurotrophins and antiapoptotic proteins, recent studies

have discovered that CREB also protects neurons via the antioxidant pathways [32].

The downstream antioxidant genes include HO-1 and manganese superoxide dismutase

(MnSOD). Besides, CREB downregulation is involved in the pathology of AD [33],

and thus increasing the expression of CREB has been considered a potential therapeutic

strategy for AD [34]. In some preconditioning experiments, CREB seemed to play a

key role in the preconditioning response which protected the cells from the subsequent

stress [35]. In the study, it was observed that TC-HT pretreatment significantly

upregulated the activation of CREB protein, compared with the control group in SH-

SY5Y cells. The result further found that HT significantly decreased the expression

level of p-CREB compared with the control cells. The decreased level of p-CREB in

HT group indicates that HT may exert too much heat stress and thus hinder the

neuroprotective effect or even cause damage to the cells. In the study, the result points

out that the neuroprotective effect of TC-HT could be attributed to the function of

antioxidant protein HO-1 or other prosurvival proteins. Further studies are necessary to

identify the specific role of other downstream neurotrophins and antiapoptotic or

antioxidant proteins in the neuroprotection effect.

Under stress amid a harsh environment or caused by external stimuli, cells or

whole organisms would trigger self-defense systems in response [36]. For example,

fever induced by viral or bacterial infections is a useful defense mechanism capable of

regulating the body temperature, as higher body temperature strengthens immune cells

and increases their ability to kill bacteria and viruses. It also inhibits the growth of

bacteria and viruses, thereby attaining anti-inflammatory effects. Depending on the

intensity and duration of stress, cells would activate cell repair or stress response to

adapt to the new environment or cause their death [37]. The stress response involves

the synthesis of some highly conserved proteins with protective function for cells [38].

For the thermal stress induced when cells or whole organisms are exposed to elevated

temperatures, they would respond by synthesizing an evolutionary highly conserved

family of proteins, known as HSPs [39]. The HS response is a universal phenomenon,

which appears in every tested organism. HSPs are named and classified, according to

their molecular weight. HSP70 (70 kilodaltons in size) is a typical group of HSPs,

which functions a molecular chaperone, to prevent the misfolding of nascent

polypeptide chains and facilitate refolding of misfolded proteins [40]. Many studies

have indicated that HSP70 sub-family is essential in protecting organisms from various

stresses [41]. The study also examined the effect of TC-HT and HT on the expression

of HSP70 and HSP105 [42], both of which have been reported to have protective effects.

The results showed that HT and TC-HT treatments enhanced the expression of both

HSPs significantly. However, the activation level of TC-HT was only slightly higher

than HT group, which could not explain the superior protective effect of TC-HT

compared to HT.

The stress response enables cells to survive the stress-induced damage. Use of

cells' self-defense ability to adapt the stress environment or manipulation of the stress

response to increase the survival rate may be valuable in regenerative medicine or

disease curing. In fact, there have been examples involving preconditioning of cells

with non-lethal stress to minimize cell damage and improve the transplantation

outcome [43]. While there have been many experiments on thermotolerance and cell

survival [44], none have demonstrated the antioxidant effect of stress response in

neuroprotection and curation to the best of our knowledge. Besides, the challenge for

the manipulation of cellular stress response is how to fine-tune stress intensity and

duration to suit specific needs and prevent the stress-induced cell damage. The present

study demonstrated a delicate and efficient way to activate the stress response, leading

to protection and curation against H_2O_2 and Aβ-induced cytotoxicity on SH-SY5Y cells.

The thermal dosage was fine-tuned in a heat-and-cold cycling process, to maximize the

protective effect and minimize the heat stress-induced cell damage. Although

traditional HS can activate HSPs which may provide resistance to insults, such as

hyperthermia, the continuous exposure may cause cell damage or even death. Previous

studies have shown that the threshold for thermal damage is different among various

tissue cells, which have different levels of thermal sensibility [45]. For example, the

skin cells can withstand much higher thermal dosage (47℃ for 20 min) than neural

cells, because the neural cells are more vulnerable. As the most thermally sensitive

tissue, brain can be damaged by low thermal dosage [45]. To attain optimal thermal

dosage for various tissues, the heating process should be divided into several parts,

which is the basic architecture of TC-HT.

In this study, the work presented an efficient and guaranteed way of controlling

the thermal dosage applied to cells. The novelty of the TC-HT strategy is the short

period of cooling process among the continuous heating period. The body temperature

(37℃) was selected as the low temperature setting which mimics the passive cooling

process in the human body. Interestingly, our data demonstrated that the application

time of the low temperature period is a critical parameter to determine the efficacy of

neuroprotection for TC-HT. For too short cooling period, the accumulation of the

thermal dosage will cause damage to neural cells like HT. On the other side, it was

found that the TC-HT did not produce protective effect to the cells when low

temperature period was longer than 1 min (Fig 2D). It may be due to the reason that the

application of too long cooling period (> 1 min) could interrupt the transmission of

biochemical signals and protein expressions stimulated by thermal stress, thus failing

to achieve the neuroprotective effect. The results revealed that 35 sec cooling process

at 37℃ after 15 min at 42.5℃ for 8 cycles produces the optimum protective effect to

SH-SY5Y cells. This protective effect was significantly much better than the

continuous HT, which is due to the activation of other thermal stress-associated proteins

apart from HSPs and the prevention of heat damage. From the LY294002 experiment,

it was confirmed that the superior protective effect resulted partially from the activation

of Akt pathway. One fascinating phenomenon is the antioxidant effect induced by TC-

HT, which could be attributed to the increased antioxidant protein HO-1. Besides, the

Aβ-induced cytotoxicity was also rescued in TC-HT treated cells, which points out the

possibility that some stress-induced proteins might have the ability to eliminate or

refold the aggregated protein or ease the symptom at least. The chaperone or

proteasome systems are possible candidates to be involved in this phenomenon [46]. It

is noteworthy that TC-HT applied after the Aβ administration gives better curative

effect than the pretreatment paradigm, which indicates that TC-HT not only has the

ability to prevent or protect the neurons against the Aβ-induced cytotoxicity but also

could have the potential to cure the disease. The possible explanation for the superior

effect of post-treatment could be that some biochemical signals and elevated proteins

could interact with the Aβ aggregation simultaneously, and these effects could not

maintain until the insult for the pretreated cells. Another advantage is that the post-

treatment paradigms for neuronal injury are also more typical in the clinical condition.

The present study employed PCR equipment to demonstrate the protective effect of TC-

HT in vitro. For in vivo or clinical applications, there are options of other heating

devices, such as high-intensity focused ultrasound (HIFU), which has been widely used

as a hyperthermal apparatus [47]. The thermal parameters can also be fine-tuned by

controlling the heating power and the size of heated volume to meet the specific needs

[48].

In summary, this is the first study on the manipulation of stress response by TC-

HT, which protects SH-SY5Y human neural cells from H_2O_2 and Aβ-induced

cytotoxicity. The results confirm that the physical stimulation by TC-HT provides safer

and much superior protective effect than continuous HS. The underlying molecular

mechanism for the protective effect is partially due to the activation of Akt pathway

and the function of some downstream proteins, such as Nrf2, CREB, and HO-1. Further

studies are necessary to confirm the involvement of other stress-induced proteins

activated by TC-HT in the neuroprotective effect or their ability in degrading or

disaggregating the aggregated proteins. It is also advisable for coupling TC-HT with

NDD drugs or natural compounds, in order to attain better curative effect for NDD

patients.

Funding

588 This work was supported by grants from Ministry of Science and Technology

589 (MOST 108-2112-M-002-016 and 105-2112-M-002-006-MY3 to CYC) and Ministry

590 of Education (MOE 106R880708 to CYC) of the Republic of China. The funders had

591 no role in study design, data collection and analysis, decision to publish, or preparation

592 of the manuscript.

Competing Interests

595 The authors have declared that no competing interests exist.

Acknowledgments

The authors would like to acknowledge the service provided by the Research Core Facilities 3 Laboratory of the Department of Medical Research at National Taiwan University hospital for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin, Chih-Yu Chao.

Formal analysis: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo-Lin, Chueh-Hsuan Lu, Hao-Ping Hsu, Yi-Kun Sun, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Wei-Ting Chen, Yu-Yi Kuo, Guan-Bo Lin.

Writing – original draft: Wei-Ting Chen, Yu-Yi Kuo, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References:

1. Kim GH, Kim JE, Rhie SJ, Yoon S. The role of oxidative stress in
neurodegenerative diseases. Exp Neurobiol. 2015; 24:325-340.

2. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing.
Nature. 2000; 408:239-247.

3. Liu Z, Zhou T, Ziegler AC, Dimitrion P, Zuo L. Oxidative stress in
neurodegenerative diseases: from molecular mechanisms to clinical applications.
Oxidative Med Cell Longev. 2017; 2017:2525967.

4. Bachurin SO, Bovina EV, Ustyugov AA. Drugs in clinical trials for Alzheimer's
disease: the major trends. Med Res Rev. 2017; 37:1186-1225.

5. Alghazwi M, Smid S, Musgrave I, Zhang W. In vitro studies of the neuroprotective
activities of astaxanthin and fucoxanthin against amyloid beta (Aβ1-42) toxicity and
aggregation. Neurochem Int. 2019; 124:215-224.

6. Hemming ML, Selkoe DJ. Amyloid beta-protein is degraded by cellular
angiotensin-converting enzyme (ACE) and elevated by an ACE inhibitor. J Biol Chem.
2005; 280:37644-37650.

7. Yang Y, Zhang X, Ye D, Laforest R, Williamson J, Liu Y, et al. Cavitation dose
painting for focused ultrasound-induced blood-brain barrier disruption. Sci Rep. 2019;
9:2840.

8. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal therapy
to synergistically enhance the anticancer effect of propolis on PANC-1 cells. Int J Oncol.
2019; 55:617-628.

9. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal cycling-hyperthermia
in combination with polyphenols, epigallocatechin gallate and chlorogenic acid, exerts
synergistic anticancer effect against human pancreatic cancer PANC-1 cells. PLoS One.
2019; 14:e0217676.

10. Sminia P, Zee JVD, Wondergem J, Haveman J. Effect of hyperthermia on the
central nervous system: a review. Int J Hyperthermia. 1994; 10:1-30.

11. Lin MT, Beal MF. Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature. 2006; 443:787-795.

12. Sanchezmargalet V, Goldfine ID, Vlahos CJ, Sung CK. Role of
phosphatidylinositol-3-kinase in insulin receptor signaling: studies with inhibitor,
LY294002. Biochem Biophys Res Commun. 1994; 204:446-452.

13. Wu X, Kosaraju J, Zhou W, Tam KY. Neuroprotective effect of SLM, a novel
carbazole-based fluorophore, on SH-SY5Y cell model and 3xTg-AD mouse model of
Alzheimer's disease. ACS Chem Neurosci. 2017; 8:676-685.

14. Xicoy H, Wieringa B, Martens GJ. The SH-SY5Y cell line in Parkinson's disease
research: a systematic review. Mol Neurodegener. 2017; 12:10.

15. Barnham KJ, Masters CL, Bush AI. Neurodegenerative diseases and oxidative

stress. Nat Rev Drug Discov. 2004; 3:205-214.

16. Chen X, Guo C, Kong J. Oxidative stress in neurodegenerative diseases. Neural Regen Res. 2012; 7:376-385.

17. Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. Nat Med. 2004; 10:S10-S17.

18. Masters SL, O'Neill LAJ. Disease-associated amyloid and misfolded protein aggregates activate the inflammasome. Trends Mol Med. 2011; 17:276-282.

19. Choi ML, Gandhi S. Crucial role of protein oligomerization in the pathogenesis of Alzheimer's and Parkinson's diseases. FEBS J. 2018; 285:3631-3644.

20. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a direct site of Aβ accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. Hum Mol Genet. 2006; 15:1437-1449.

21. Matsuoka Y, Picciano M, La Francois J, Duff K. Fibrillar β-amyloid evokes oxidative damage in a transgenic mouse model of Alzheimer's disease. Neuroscience. 2001; 104:609-613.

22. Huang HC, Jiang ZF. Accumulated amyloid-β peptide and hyperphosphorylated tau protein: relationship and links in Alzheimer's disease. J Alzheimers Dis. 2009; 16:15-27.

23. Zhao Y, Zhao B. Oxidative stress and the pathogenesis of Alzheimer's disease. Oxidative Med Cell Longev. 2013; 2013:316523.

24. de Oliveira MR, de Bittencourt Brasil F, Fürstenau CR. Sulforaphane promotes mitochondrial protection in SH-SY5Y cells exposed to hydrogen peroxide by an Nrf2-dependent mechanism. Mol Neurobiol. 2017; 55:4777-4787.

25. Ismail N, Ismail M, Imam MU, Azmi NH, Fathy SF, Foo JB, et al. Mechanistic basis for protection of differentiated SH-SY5Y cells by oryzanol-rich fraction against hydrogen peroxide-induced neurotoxicity. BMC Complement Altern Med. 2014; 14:467.

26. Han SM, Kim JM, Park KK, Chang YC, Pak SC. Neuroprotective effects of melittin on hydrogen peroxide-induced apoptotic cell death in neuroblastoma SH-SY5Y cells. BMC Complement Altern Med. 2014; 14:286.

27. Jason S, Cui W. Proliferation, survival and metabolism: the role of PI3K/AKT/mTOR signalling in pluripotency and cell fate determination. Development. 2016; 143:3050-3060.

28. Huang Y, Li W, Su ZY, Kong ANT. The complexity of the Nrf2 pathway: beyond the antioxidant response. J Nutr Biochem. 2015; 26:1401-1413.

29. Kubben N, Zhang W, Wang L, Voss TC, Yang J, Qu J, et al. Repression of the antioxidant NRF2 pathway in premature aging. Cell. 2016; 165:1361-1374.

30. Wang Z, Guo S, Wang J, Shen Y, Zhang J, Wu Q. Nrf2/HO-1 mediates the

neuroprotective effect of mangiferin on early brain injury after subarachnoid

hemorrhage by attenuating mitochondria-related apoptosis and neuroinflammation. Sci

Rep. 2017; 7:11883.

31. Caracciolo L, Marosi M, Mazzitelli J, Latifi S, Sano Y, Galvan L, et al. CREB

controls cortical circuit plasticity and functional recovery after stroke. Nat Commun.

2018; 9:2250.

32. Sakamoto K, Karelina K, Obrietan K. CREB: a multifaceted regulator of neuronal

plasticity and protection. J Neurochem. 2011; 116:1-9.

33. Rosa E, Fahnestock M. CREB expression mediates amyloid β-induced basal

BDNF downregulation. Neurobiol Aging. 2015; 36:2406-2413.

34. Qin Z. Modulating nitric oxide signaling in the CNS for Alzheimer's disease

therapy. Future Med Chem. 2013; 5:1451-1468.

35. Baranova KA, Rybnikova EA, Churilova AV, Vetrovoy OV, Samoilov MO. The

adaptive role of the CREB and NF-κB neuronal transcription factors in post-stress

psychopathology models in rats. Neurochem J. 2014; 8:17-23.

36. Chovatiya R, Medzhitov R. Stress, Inflammation, and defense of homeostasis.

Mol Cell. 2014; 54:281-288.

37. Thomas MP, Lieberman J. Live or let die: posttranscriptional gene regulation in

cell stress and cell death. Immunol Rev. 2013; 253:237-252.

38. Blanco S, Bandiera R, Popis M, Hussain S, Lombard P, Aleksic J, et al. Stem cell

function and stress response are controlled by protein synthesis. Nature. 2016; 534:335-

340.

39. Lindquist S. The heat-shock response. Annu Rev Biochem. 1986; 55:1151-1191.

40. Mosser DD, Caron AW, Bourget L, Meriin AB, Sherman MY, Morimoto RI, et al.

The chaperone function of hsp70 is required for protection against stress-induced

apoptosis. Mol Cell Biol. 2000; 20:7146-7159.

41. Yenari MA, Liu J, Zheng Z, Vexler ZS, Lee JE, Giffard RG. Antiapoptotic and

anti-inflammatory mechanisms of heat-shock protein protection. Ann NY Acad Sci.

2005; 1053:74-83.

42. Hatayama T, Yamagishi N, Minobe E, Sakai K. Role of hsp105 in protection

against stress-induced apoptosis in neuronal PC12 cells. Biochem Biophys Res

Commun. 2001; 288:528-534.

43. Matsumoto K, Honda K, Kobayashi N. Protective effect of heat preconditioning

of rat liver graft resulting in improved transplant survival. Transplantation. 2001;

71:862-868.

44. Mesihovic A, Iannacone R, Firon N, Fragkostefanakis S. Heat stress regimes for

the investigation of pollen thermotolerance in crop plants. Plant Reprod. 2016; 29:93-

105.

45.   Dewhirst MW, Viglianti BL, Lora-Michiels M, Hanson M, Hoopes PJ. Basic

principles of thermal dosimetry and thermal thresholds for tissue damage from

hyperthermia. Int J Hyperthermia. 2003; 19:267-294.

46.   Kampinga HH, Bergink S. Heat shock proteins as potential targets for protective

strategies in neurodegeneration. Lancet Neurol. 2016; 15:748-759.

47.   Frazier N, Payne A, de Bever J, Dillon C, Panda A, Subrahmanyam N, et al. High

intensity focused ultrasound hyperthermia for enhanced macromolecular delivery. J

Control Release. 2016; 241:186-193.

48.   Partanen A, Tillander M, Yarmolenko PS, Wood BJ, Dreher MR, Kohler MO.

Reduction of peak acoustic pressure and shaping of heated region by use of multifoci

sonications in MR-guided high-intensity focused ultrasound mediated mild

hyperthermia. Med Phys. 2013; 40:013301.

A

B

C

A

B

C

| Ctrl | Aβ | TC (post)+Aβ | HT (post)+Aβ |

B

A

Cell viability (%) — Ctrl, H_2O_2, TC+H_2O_2, LY+TC+H_2O_2

***

B

p-Akt / GAPDH

Nrf2 / GAPDH

p-Akt/GAPDH — Ctrl, H_2O_2, TC+H_2O_2, LY+TC+H_2O_2

Nrf2/GAPDH — Ctrl, H_2O_2, TC+H_2O_2, LY+TC+H_2O_2

***