Massively parallel, time-resolved single-cell RNA sequencing with scNT-Seq

Qi Qiu[1,2,#], Peng Hu[1,2,#], Kiya W. Govek[1,3], Pablo G. Camara[1,3], Hao Wu[1,2,*]

[1]Department of Genetics, University of Pennsylvania, Philadelphia, PA 19104

[2]Penn Epigenetics Institute, University of Pennsylvania, Philadelphia, PA 19104

[3]Institute of Biomedical Informatics, University of Pennsylvania, Philadelphia, PA 19104

[#]These authors contributed equally to this work

*Correspondence should be addressed to H.W. (haowu2@pennmedicine.upenn.edu).

ABSTRACT

Single-cell RNA sequencing offers snapshots of whole transcriptomes but obscures the temporal dynamics of RNA biogenesis and decay. Here we present single-cell nascent transcript tagging sequencing (scNT-Seq), a method for massively parallel analysis of nascent and pre-existing RNAs from the same cell. This droplet microfluidics-based method enables high-throughput chemical conversion on barcoded beads, efficiently marking metabolically labeled nascent transcripts with T-to-C substitutions. By simultaneously measuring nascent and pre-existing transcriptomes, scNT-Seq reveals neuronal subtype-specific gene regulatory networks and time-resolved RNA trajectories in response to brief (minutes) versus sustained (hours) neuronal activation. Integrating scNT-Seq with genetic perturbation reveals that DNA methylcytosine dioxygenases may inhibit stepwise transition from pluripotent embryonic stem cell state to intermediate and totipotent two-cell-embryo-like (2C-like) states by promoting global nascent transcription. Furthermore, pulse-chase scNT-Seq enables transcriptome-wide measurements of RNA stability in rare 2C-like cells. Time-resolved single-cell transcriptomic analysis thus opens new lines of inquiry regarding cell-type-specific RNA regulatory mechanisms.

INTRODUCTION

Dynamic changes in RNA levels are regulated by the interplay of RNA transcription, processing, and degradation[1, 2]. Understanding the mechanisms by which the transcriptome is regulated in functionally diverse cell-types within multi-cellular organisms thus requires cell-type-specific measurements of the kinetics of RNA biogenesis and decay. Recent advances in single-cell RNA sequencing (scRNA-Seq) technologies are leading to a more complete understanding of heterogeneity in cell types and states[3]. However, standard scRNA-Seq methods capture a mixture of newly synthesized and pre-existing RNAs without being able to directly reveal nascent RNA dynamics.

Accurately capturing changes in the nascent transcriptome over time at single-cell resolution is of particular interest because measuring nascent RNAs can reveal immediate regulatory changes in response to developmental, environmental, metabolic, and pathological signals[4]. Commonly used approaches for distinguishing nascent from pre-existing RNAs of the same population of transcripts rely on metabolic labeling that employs thiol-labeled nucleoside analogs such as 4-thiouridine (4sU) and subsequent biochemical enrichment of metabolically labeled RNAs[2]. Although these methods have yielded unprecedented insights into the regulation of RNA dynamics, they require ample starting material and present challenges for enrichment normalization. Several approaches have recently been developed to chemically convert 4sU into cytidine analogs, yielding uracil-to-cytosine (U-to-C) substitutions that label nascent RNAs after reverse transcription[5-7]. These chemical nucleotide conversion methods allow for direct measurement of temporal information about cellular RNAs in a sequencing experiment without biochemical enrichment. Recent studies integrated standard plate-based single-cell RNA-Seq method with one of these chemical methods (i.e. thiol(SH)-linked alkylation for the metabolic sequencing of RNA (SLAM)-Seq)[8, 9], demonstrating the feasibility of studying nascent transcriptomes at single-cell levels. However, these plate-based single-cell SLAM-Seq methods suffer from several limitations. First, they are costly, and the associated library preparation steps are time-consuming, prohibiting it for large-scale single-cell analysis of highly heterogeneous cell populations. Second, these methods lack unique molecular identifiers (UMIs), preventing accurate quantification of the nascent transcript levels.

To overcome these constraints, we developed single-cell nascent transcript tagging sequencing (scNT-Seq), a high-throughput and UMI-based nascent scRNA-Seq approach. In scNT-Seq, integration of metabolic RNA labeling, droplet microfluidics, and chemically induced recoding of 4sU to cytosine analog permits highly scalable and time-resolved single-cell analysis of cellular RNA dynamics. We demonstrate that the method is easy to set up and substantially improves the time and cost associated with simultaneous single-cell analysis of nascent and pre-existing transcriptomes. We show scNT-Seq enables more detailed characterization of gene regulatory networks and temporal RNA trajectories than single-cell whole-transcriptome measurements alone.

RESULTS

Development and validation of scNT-Seq. To enable nascent and pre-existing transcripts from the same cell to be analyzed in a scalable manner, we focused on the Drop-Seq platform because its unique barcoded bead design affords both immobilization of RNAs for efficient chemical conversion reactions and UMI-based high-throughput scRNA-Seq analysis, and this droplet microfluidics platform has been widely adopted[10-14]. The scNT-Seq consists of the following key steps (**Fig. 1a**): (1) metabolically labeling of cells with 4sU for a temporally defined time period; (2-3) co-encapsulating each individual cell with a barcoded oligo-dT primer coated bead in a nanoliter-scale droplet which captures both nascent and pre-existing RNAs; (4) performing 4sU chemical conversion on pooled barcoded beads in one reaction after droplet breakage; (5-8) reverse transcription, cDNA amplification, tagmentation, indexing PCR, and sequencing; and (9) using a UMI-based statistical model to analyze T-to-C substitutions within transcripts and infer the nascent transcript fraction at single-cell level.

To identify the optimal reaction conditions on barcoded beads, we first explored two independent chemical conversion methods (TimeLapse-Seq: 2,2,2-trifluoroethylamine (TFEA)/sodium periodate ($NaIO_4$)-based 4sU conversion; SLAM-Seq: iodoacetamide (IAA)-based conversion) and validated their performance with species-mixing experiments using cultured mouse embryonic stem cells (mESCs) and human K562 cells. This analysis indicates that TFEA/$NaIO_4$-based scNT-Seq substantially outperforms the IAA-based assay in terms of mRNA recovery rates (**Fig. 1b**). Furthermore, the collision rate is comparable between TFEA/$NaIO_4$-based scNT-Seq and standard Drop-Seq (**Fig. 1b**), demonstrating the specificity of scNT-Seq in analyzing single-cell transcriptomes. Shallow sequencing of mESCs under different treatment conditions showed that TFEA/$NaIO_4$-based scNT-Seq identified a similar number of genes or UMIs per cell compared to standard Drop-Seq (**Supplementary Fig. 1a**). Aggregated single-cell nascent or pre-existing transcriptomes were highly correlated between replicates (**Supplementary Fig. 1b**). Further analysis of K562 cells revealed that only 4sU labeling and TFEA/$NaIO_4$ treatment resulted in a substantial increase in T-to-C substitution (**Fig. 1c**) and in UMIs that contained one or more such substitutions (**Fig. 1d, e**). Notably, TFEA/$NaIO_4$ chemical treatment works efficiently with both freshly isolated and cryo-preserved (methanol-fixed) cells (**Supplementary Fig. 1c**), demonstrating the versatility of scNT-Seq. Collectively, these data indicate that TFEA/$NaIO_4$-based scNT-Seq is capable of efficiently detecting nascent transcripts at single-cell resolution.

Application of scNT-Seq to study neuronal activity-dependent nascent transcription and regulatory networks.

Neuronal activity induces expression of hundreds of activity-regulated genes (ARGs) in the vertebrate brain, leading to new protein synthesis and epigenetic changes necessary for short- and long-term memories of experiences. Thus, the coupling of synaptic activity to nascent transcription in the nucleus allows neurons to both respond dynamically to their immediate environment, and to store information

stably[15]. Recent studies suggest that different patterns of neuronal activity could induce a distinct set of

ARGs[16]. To investigate cell-type-specific nascent transcription in response to distinct activity-duration

patterns (brief versus sustained stimulation), we applied scNT-Seq to primary cortical neuronal cultures,

derived from mouse cortex (embryonic day 16 (E16)), which contains heterogeneous populations of

neuronal subtypes and undifferentiated neural progenitor cells.

To detect activity-regulated nascent transcriptional responses, we metabolically labeled primary mouse

cortical cultures (200 μM 4sU) for two hours and stimulated the cells with different durations of neuronal

activity (0, 15, 30, 60 and 120 min of potassium chloride (KCl)-mediated membrane depolarization) (**Fig.**

2a). After filtering low-quality cells and potential doublets, we retained 20,547 single-cell transcriptomes

from five time-points (**Fig. 2b** and **Supplementary Table 1**). After sample integration, principal

component analysis (PCA), graph-based clustering, and visualization by Uniform Manifold Approximation

and Projection (UMAP), we identified nine major cell-types based on known marker genes: *Neurod6*+

cortical excitatory neurons (Ex, 68.5%), four *Gad1*+ inhibitory neuronal subtypes (Inh1-4, 13.9% in total),

Dlx1/Dlx2+ inhibitory neuronal precursors (Inh-NP: 1.7%), two sub-populations of *Nes/Sox2*+ excitatory

neuronal precursors (Ex-NP1/2: 10.4% in total), and *Nes/Aldh1l1*+ radial glia (RG: 5.5%) (**Fig. 2b** and

Supplementary Fig. 2b, c).

We next sought to distinguish nascent from pre-existing RNAs by counting and statistically modeling T-to-

C substitutions in transcripts (UMIs), an approach that overcomes the problem of incomplete 4sU labeling

of nascent transcripts (up to 50% of all reads originated from new RNAs may not contain T-to-C

substitutions)[17]. The quantification accuracy is further improved by UMI-based analysis, which

substantially increases the number of uridines or T-to-C substitutions covered in each transcript

compared to the analysis of individual sequencing reads (**Supplementary Fig. 3a, b**). After statistical

correction, we obtained reliable measurements of nascent RNA faction for both activity-induced genes

(e.g., *Fos*, ~90% nascent/total) and slow turnover house-keeping genes (e.g., *Mapt*, <5% nascent/total) in

excitatory neurons (**Supplementary Fig. 3c, d**).

Furthermore, sub-clustering of nascent (upper) or pre-existing (lower) transcriptomes derived from

randomly sampled excitatory neurons could readily separate nascent, but not pre-existing single-cell

transcriptomes in an activity-pattern dependent manner (right panels in **Fig. 2b**). In contrast, nascent

transcriptomes of undifferentiated excitatory neural precursors (Ex-NP/RG) did not exhibit similar

distributions (**Supplementary Fig. 4a**). Next, we directly examined how nascent transcription of classic

ARGs are induced in different cell-types in response to distinct activity-patterns. While some ARGs, such

as *Jun* and *Btg2*, were specifically induced in Ex neurons upon activation, other ARGs (e.g., *Egr1*, *Fos*,

and *Npas4*), were broadly induced in many cell-types including non-neuronal cells, albeit with different

magnitudes and response curves (**Fig. 2c, Supplementary Fig. 4b**). There was little to no change at pre-

existing levels upon activation of Ex neurons (**Fig. 2c**). Together, these results suggest single-cell

nascent transcriptome analysis by scNT-Seq can accurately detect cell-type-specific, activity-induced changes in nascent transcription within the timescale of minutes to hours.

Recent advances in computational analysis enable identification of specific gene regulatory networks (GRNs, also known as regulon of a transcription factor (TF)) underlying stable cell states by linking *cis*-regulatory sequences to single-cell gene expression. Since the regulon is scored as a whole, instead of using the expression of individual genes, this approach is more robust against experimental dropouts or stochastic variation of gene expression due to transcriptional bursting. We reasoned that applying such an approach to nascent transcriptomes derived from scNT-Seq may allow for analysis of regulons underlying dynamic cell states induced by external stimuli such as neuronal activity. By applying single-cell regulatory network inference and clustering (SCENIC)[18] to statistically-corrected nascent and pre-existing single-cell transcriptomes, we identified 79 regulons with significant *cis*-regulatory motif enrichment showing significant changes in response to at least one activity-pattern (**Fig. 2d**, **Supplementary Fig. 5**). Many early-response ARGs encode TFs that regulate a subsequent wave of late-response gene expression. SCENIC analysis of nascent transcriptomes revealed neuronal activity-dependent increase in TF regulon activity of well-established IEG TFs (e.g., Fos, Jun, and Egr family of TFs) as well as constitutively expressed TFs such as Srf and Mef2, both of which are activated by multiple calcium-dependent signaling pathways and undergo post-translational modifications (e.g., phosphorylation) in response to neuronal activity (**Fig. 2d**). Thus, these TFs represent a group of TFs that are the main mediator of activity-dependent nascent transcription. This group also included several TFs that have not been previously implicated in neuronal activation, including Maff and Hspa5, both of which exhibited higher expression and regulon activity upon activation (**Fig. 2d**). In addition, both Rfx3, a ciliogenic TF[19], and the neural cell fate regulator Pou3f1[20] were identified as potential novel regulators induced by neuronal activity. Furthermore, we observed regulon activity of cell-type-specific TFs (Neurod1/2 for Ex, Sox2/3 for Ex-NP and Dlx1/2 for Inh), which are associated with stable cell-type identities and detected in both nascent and pre-existing transcriptomes (**Fig. 2d**). Collectively, these data suggest that scNT-Seq allows for the analysis of cell-type-specific TF regulons that underlie the dynamic cellular responses of both nascent and pre-existing transcriptomes to acute and sustained stimuli.

scNT-Seq enables nascent transcription-based RNA velocity analysis

A fundamental question in gene regulation is how transcriptional states in single cells change over time in response to both acute (minutes) and sustained (hours to days) external stimuli. Recent work showed that the time derivative of the gene expression state, termed "RNA velocity," can be estimated by distinguishing between unspliced ("new" intronic reads) and spliced ("old" exonic reads) mRNAs in scRNA-Seq datasets[21]. The RNA velocity measurement can predict the future state of individual cells on a timescale of hours. Because ultra-short metabolic labeling (5 minutes of 4sU pulse) can be used to identify transient RNAs[22] and scNT-Seq enables direct measurements of nascent and pre-existing

transcripts from the same cell, we next sought to augment the RNA velocity model to predict rapid cell

state changes by replacing unspliced/spliced read levels with UMI-based nascent and total RNA levels.

We focused on the excitatory neurons as this population of neurons robustly responds to neuronal

activation. To compare unspliced/spliced ratio-based RNA velocity (termed "splicing RNA velocity") and

nascent RNA fraction-based velocity (termed "nascent RNA velocity") directly, we projected and aligned

both velocity fields onto the same UMAP plot (**Fig. 3a**). While splicing RNA velocity showed a steady

directional flow (arrows) from 60 to 120 min (left in **Fig. 3a**), nascent RNA velocity revealed two distinct

phases of velocity flows (early phase: 15 to 30 min; late phase: 60 to 120 min) (right in **Fig. 3a**). Thus, our

analyses are consistent with recent reports that these two types of velocity measurements convey

different but complementary information on the future state of a cell[8, 21].

Nascent RNA velocity is grounded in direct measurements of metabolically labeled nascent transcription,

and this approach promises to improve quantitative analysis of the dynamics of cell states, particularly for

the analysis of transient and dynamic responses to rapid external stimulation. Indeed, nascent RNA

velocity (area under the receiver operating characteristic curve (AUC) value = 82.5%) outperformed

splicing RNA velocity (AUC=51.5%) analysis in predicting the activity-induced primary response genes

(PRGs, 137 genes) that were identified in targeted bulk RNA-seq analysis of primary cortical neurons[16]

(**Fig. 3b**). These results show that nascent RNA velocity is more reliable in revealing the rapid

transcriptional changes following acute neuronal activation.

To further explore the molecular basis underlying the two distinct phases of nascent RNA velocity flows,

we identified the TF regulons that are significantly associated with early- (n=24; induced at 15 or 30 min)

or late-response (n=73; induced at 60 or 120 min) genes (**Fig. 3c**, **Supplementary Fig. 4c**). We further

projected aggregated nascent RNA levels of early- and late-response genes onto respective RNA velocity

fields to visualize the relationship between distinct activity-patterns and ARG induction (**Fig. 3d**). Next,

we projected TF regulon activity onto nascent RNA velocity field, revealing distinct activity patterns of TFs

primarily associated with either early- (Jun) or late- (Mef2d) response genes in dynamic cell state

changes (**Fig. 3e**). Together, these results indicate that integrative analysis of nascent RNA velocity and

TF regulon activity can be a powerful approach to reveal molecular insights into dynamic cellular

processes.

scNT-Seq captures changes in nascent transcription during the pluripotent-to-totipotent stem cell
state transition.

Determining temporal RNA dynamics such as nascent transcriptional events within rare, transient cell

populations is critical to understanding cell state transition but has remained a significant challenge for

stem cell research. Recent advances in multiplex single-molecule imaging using sequential fluorescence

in situ hybridization (seqFISH) and intron probes allows for the detection of nascent transcription sites at

the single-cell level[23]; however, such imaging approaches are currently time-consuming and have not

been used to study rare stem cell populations at scale. Using mESC cultures as a model, we sought to determine whether scNT-Seq could be used to directly investigate changes in nascent transcription and cell state transition within rare stem cell populations.

Cultured mESCs are derived from the inner cell mass of pre-implantation blastocysts and exhibit a high level of transcriptional heterogeneity[24]. Interestingly, cells resembling 2-cell-stage embryos (2C-like cells), which are in a totipotent state and can differentiate into both embryonic and extraembryonic tissues, arise spontaneously in mESC cultures[25]. Nearly all mESCs reversibly cycle between pluripotent and totipotent 2C-like state in culture. However, the 2C-like cells are rare in mESC cultures (<1% in standard culture conditions)[25], making it challenging to study these totipotent cells with traditional methods. Recent studies using scRNA-Seq revealed changes in total RNAs during the transition to rare 2C-like state[26, 27] and identified an intermediate state during this low-frequency transition process, supporting a stepwise model of transcriptional reprogramming of the pluripotent to 2C-like state transition[26, 27]. However, these studies cannot distinguish nascent from pre-existing RNAs at each state. Thus, a more detailed understanding of direct gene regulatory changes in response to state transition is lacking.

To capture nascent transcriptomes and characterize GRNs in different states, mESCs were metabolically labeled with 4sU for 4 hours and were subjected to scNT-Seq analysis (**Fig. 4a**). After quality filtering, we obtained 4,633 single-cell transcriptomes from two biological replicates (**Supplementary Fig. 6a, b**). Unbiased clustering not only readily separated contaminating mouse feeder cells (*Col1a2*/*Thbs1*+) from mESCs, but also identified three principal states within mESCs (**Supplementary Fig. 6a-c**). The majority of mESCs (98.3%) belong to the pluripotent state as they are positive for pluripotency-genes such as *Sox2* and *Zfp42* (also known as Rex1) but are negative for 2C-embryo-specific transcripts such as the *Zscan4* gene family (**Fig. 4b**). Approximately 0.7% of cells express high levels of 2C-like state markers, including *Zscan4* genes, *Zfp352* and *Usp17lc*, but low levels of *Sox2*, suggesting that this rare population contains predominantly totipotent 2C-like cells. These results are consistent with previous FACS analysis of 2C-reporter lines[25, 28], indicating that scNT-Seq accurately captures this rare cell state. Interestingly, we also identified an intermediate state (~1.0%) which expresses low levels of *Zscan4* genes, but high levels of *Nanog* and *Tbx3* (**Supplementary Fig. 6c, d**). Unlike previous studies which overexpressed a master regulator, Dux, to induce 2C-like state[27], we did not observe a significant decrease in *Sox2* expression in the intermediate state (**Supplementary Fig. 6c**), highlighting the potential difference between the Dux-induced 2C-transitions and the naturally occurring pluripotent-to-totipotent transition process.

Comparative analysis of nascent and pre-existing transcriptomes revealed many state-specific genes (e.g., *Sox2* and *Zscan4d*) are associated with a higher proportion of nascent transcripts (consistent with their regulatory roles), whereas house-keeping genes, such as *Gapdh*, are of slower turnover rate (**Supplemental Fig. 7a, b**). For many state-specific genes (e.g., *Zfp42* and *Zscan4d*), their nascent RNA levels exhibited a more pronounced difference between states than the change of pre-existing RNAs (**Fig. 4b, c**), indicating that nascent transcription is a more sensitive indicator for the state transition. Gene

Ontology (GO) enrichment analysis further revealed that state-specific nascent and pre-existing
transcripts were enriched for different GO terms (**Fig. 4d**). For instance, many pluripotent-state-specific
nascent RNAs (but not pre-existing RNAs) are functionally related to post-transcriptional regulatory
processes, including translational initiation, mRNA processing, and RNA splicing. Furthermore, 2C-state-
specific nascent transcripts are preferentially enriched for deubiquitinating enzyme-related genes such as
Usp17lc/d/e (**Fig. 4b, d**). These results underscore that nascent transcriptomes are more robust than
steady-state transcriptomes to uncover genes related to state-specific biological processes within
heterogeneous stem cell populations.

DNA-binding proteins, such as TFs and epigenetic regulators, show rapid gene expression changes
during pluripotent-to-2C transition[27, 29]. Our analysis of changes in nascent transcription during the
pluripotent-to-2C transition identified 26 genes encoding DNA-binding proteins that showed a significant
difference at nascent RNA levels between states (**Fig. 4e**). Consistent with previous scRNA-Seq analysis
of sorted cell populations[29], 13 out of 14 commonly detected genes (*Tet1*, *Sox2*, *Nanog*, *Rex1/Zfp42*,
Sp110, *Zfp352,* and *Zscan4* family) showed similar patterns in our scNT-Seq analysis of nascent RNAs
(**Fig. 4e**). To further investigate TF/epigenetic regulator activity during the pluripotent-to-2C transition, we
applied SCENIC analysis to both nascent and pre-existing transcriptomes. Among all regulons uncovered
in mESCs (**Supplementary Fig. 8a**), we identified 25 TF/epigenetic regulators showing differential
nascent regulon activity between states (**Fig. 4f** and **Supplementary Fig. 8b**). The activity of several
positive regulators of the cell cycle, such as the E2F family of TFs, decreased in 2C-like states, which is
consistent with previous observations that 2C-like cells are associated with longer cell cycles[30].
Interesting, nascent RNA levels and regulon activity of Nanog peak in intermediate state compared to
pluripotent and totipotent 2C-like states, suggesting that this TF may play an unrecognized regulatory role
in intermediate state. In addition, several epigenetic regulators, including Phf8, Hdac2, Ezh2, and Tet1,
were associated with a decrease in regulon activity (**Fig. 4f**), suggesting a role of these enzymes in
promoting the pluripotent state. Notably, we did not identify the regulon activity of Zscan4 in 2C-like cells,
potentially due to the lack of Zscan4 motif information in the SCENIC TF database. Together, scNT-Seq
directly captures state-specific nascent transcriptomes in rare totipotent cells in heterogeneous mESC
cultures and allows analysis of dynamics of nascent regulon activity during state transitions.

TET-dependent regulation of the stepwise pluripotent-to-2C transition
The TET family of proteins (Tet1-3) are DNA dioxygenases that mediate active DNA demethylation at
many *cis*-regulatory elements such as promoters and distal enhancers[31, 32], thereby regulating gene
expression. During pluripotent-to-2C transition, both nascent RNA level and regulon activity of *Tet1*
rapidly decreased (**Fig. 4c, f**). The nascent transcript level of *Tet2* also decreased in both intermediate
and 2C-like states, while *Tet3* was undetected in all three states (**Supplementary Fig. 7c**). Interestingly,

genetic inactivation of all TET proteins in mESCs resulted in a substantially higher proportion of 2C-like

cells, and elevated expression level of 2C-specific transcripts[28].

To better understand the role of TET enzymes in state transition, we generated mESCs deficient for all

three Tet proteins (*Tet1/2/3* triple knockout, *Tet*-TKO) via CRISPR/Cas9 genome editing using previously-

tested sgRNAs[33] and analyzed isogenic WT and *Tet*-TKO mESCs (J1 strain) by scNT-Seq after 4 hours

of metabolic labeling. The genotype of *Tet*-TKO cells was confirmed by both Sanger sequencing and

reads from scNT-Seq (**Supplementary Fig. 9a**). Clustering and nascent RNA velocity analyses of

combined WT and *Tet*-TKO mESCs revealed two discrete phases of nascent RNA velocity flows (arrows).

The velocity flow within the pluripotent state is potentially driven by cell-cycle progression

(**Supplementary Fig. 9b, c**), whereas nascent velocity field of intermediate/2C-like states exhibited a

strong directional flow from intermediate to 2C-like state (**Fig. 5a, b**). These observations are consistent

with a stepwise model of the state-transition process and suggest that the pluripotent-to-intermediate

state transition is a rate-limiting step.

WT and *Tet*-TKO cells were separately clustered within the pluripotent state and *Tet*-TKO (but not WT)

cells were found immediately next to the intermediate/2C-like states (**Fig. 5a, b**), suggesting that *Tet*-TKO

cells are transcriptionally reprogrammed to adopt a more poised state to transition to intermediate/2C-like

states. Compared to WT cells, *Tet*-TKO cells exhibited a 2.2-fold or 3.6-fold increase in intermediate or

2C-like states (**Fig. 5c**), respectively, indicating that Tet enzymes negatively regulate the transition from

pluripotent to intermediate/2C-like state in WT cells. Because pluripotent and 2C-like cells are globally

distinct in levels of both major histone modifications[25] and DNA methylation[26], we next examined global

nascent transcription in each state. Interestingly, aggregating nascent transcriptomes within specific

states revealed that a totipotent 2C-like state is associated with markedly lower levels of global nascent

transcription compared to pluripotent and intermediate states in WT cells (**Fig. 5d**). In contrast, *Tet*-TKO

cells already exhibited a substantially lower level of global nascent transcription than WT cells in

pluripotent state (**Fig. 5d**), suggesting that Tet proteins may act as an epigenetic barrier for the

pluripotent-to-2C transition by maintaining a pluripotent state-specific nascent transcriptome.

Pulse-chase scNT-Seq enables transcriptome-wide measurement of mRNA stability in rare 2C-like
cells.

Pulse-chase assays combined with bulk SLAM-Seq have been used to measure mRNA stability in

mESCs[5], but rare intermediate and 2C-like cells have not been studied. Given the enhanced sensitivity of

scNT-Seq compared to bulk assays, combining a pulse-chase strategy with scNT-Seq may enable

transcriptome-wide measurement of mRNA stability in rare cell populations. To test this, we metabolically

labeled mESC cultures with 4sU for 24 hours (pulse), followed by a washout and chase using medium

containing a higher concentration of uridine. Cells from multiple chase time-points were harvested and

cryo-preserved first, and all samples were then re-hydrated and simultaneously analyzed by scNT-Seq

analysis. After computing the proportion of labeled transcripts for each gene at every time-point relative to

0 h (right after pulse), we calculated the half-life ($t_{1/2}$) of mRNAs by fitting a single-exponential decay

model in each cell state (**Fig. 6a**). In total, 20,190 cells were profiled by scNT-Seq from 7 time-points (**Fig.**

6b). After 24 hours of metabolic labeling, we observed a substantial accumulation of T-to-C substitution

(**Fig. 6c**), which is consistent with bulk assay observations[5]. The T-to-C substitution rate decreased over

time and returned to the baseline level 24 hours after chase (**Fig. 6c**). Inspection of two genes, *Sox2* and

Topa2a, revealed that total RNA levels did not change across the 24-hour time course, whereas the level

of metabolically labeled transcripts decreased over time (**Supplementary Fig. 10a**), confirming that this

strategy can accurately measure mRNA decay in a transcript-specific manner.

Based on expression of marker genes, clustering analysis readily separated mESCs into three states

(n=7, pluripotent: 97.4%+/-0.78%, intermediate: 1.5%+/-0.48%, and 2C-like: 1.1%+/-0.38%) (**Fig. 6b** and

Supplementary Fig 10b), suggesting that this pulse-chase approach did not significantly alter the state

transition. By filtering out genes expressed in less than 5% of cells, we were able to determine the half-life

of 2,310 genes in pluripotent state, and the RNA half-life determined by pulse-chase scNT-Seq is highly

concordant with previous observation derived from bulk SLAM-Seq assays[5] (R^2=0.81, **Fig. 6d**). RNA

stability can also be estimated by assuming simple exponential kinetics, which is calculated by measuring

the ratio of labeled and unlabeled transcripts after metabolic labeling for a specific time[34]. However, RNA

half-life estimated from one timepoint labeling (4sU, 4 hour) experiment is substantially less correlated

with measurements derived from bulk assays (R^2=0.49, **Fig. 6d**). These results suggest that a pulse-

chase strategy may more reliably measure RNA decay rate. Furthermore, the top 10% most stable and

unstable transcripts were enriched for similar GO terms that are uncovered by bulk SLAM-Seq assays[5]

(**Supplementary Fig. 10c**). Further analysis of cells in intermediate and 2C-like states revealed the half-

life of 1,743 and 821 transcripts in these rare cell states, respectively. Next, we analyzed commonly

detected transcripts between cell states to reveal state-specific regulation of mRNA stability

(**Supplementary Fig. 11**). Thus, scNT-seq enables transcriptome-wide measurement of RNA stability in

rare stem cell populations in heterogeneous mESC cultures.

Because dynamic changes in RNA levels are regulated by the interplay of RNA biogenesis and

degradation, we sought to investigate the relationship between mRNA stability and gene expression

during the pluripotent-to-2C transition (**Supplementary Fig. 12**). Consistent with previous findings[2], RNA

stability and total mRNA levels are not highly correlated, suggesting mRNA stability is not the major

contributor to total RNA level during stem cell state transition. Nevertheless, we identified a group of

genes showing coordinated changes of RNA stability and gene expression level between pluripotent and

2C-like states, suggesting mRNA stability may play a role in regulating total mRNA level for a subset of

genes (**Fig. 6e**). Further analysis showed that six RNA-binding protein (RBP) binding motifs were

enriched in 3'UTR of 15 genes showing coordinated changes (**Fig. 6f**). Among these RBPs, the Pabpc

family proteins (Pabpc1 and Pabpc4) are known regulators of mRNA stability and translation efficiency by

binding poly-A tails of mRNAs[35]. Collectively, these results indicate that scNT-Seq can serve as a

powerful approach to investigate RNA stability and post-transcriptional regulatory mechanisms in rare

cells within a heterogeneous population of cells.

DISCUSSION

By combining the widely accessible Drop-Seq platform and TimeLapse chemistry, scNT-Seq provides a

novel strategy for high-throughput analysis of nascent and pre-existing transcriptomes from the same cell.

A major advantage of scNT-Seq is its ability to accurately estimate nascent RNA levels using UMI-based

statistical correction, which not only reveals acute changes in the nascent transcription that are not

apparent in standard scRNA-seq, but also allows for the analysis of dynamic TF regulatory networks and

single-cell transcriptional trajectory. Our scNT-Seq is conceptually similar to sci-fate[36], a recently

developed method that integrates SLAM-Seq chemistry with single-cell combinatorial indexing RNA-Seq.

While scNT-Seq directly captures whole-cell total RNAs on barcoded beads in nanoliter droplets, sci-fate

requires paraformaldehyde fixation and permeabilization of cells for *in situ* chemical conversion and

reverse transcription, which may pose a challenge to unbiasedly recover both nuclear (enriched for

nascent) and cytoplasmic (enriched for pre-existing) RNAs for accurately estimating nascent RNA

fractions.

Recent studies demonstrate that the purine analog, 6-thioguanine (6tG), enables G-to-A conversions by

TimeLapse chemistry in bulk RNA-seq[37]. Thus, dual labeling of 4sU and 6tG followed by scNT-Seq may

allow two independent recordings of nascent transcription at single-cell levels. With new computational

approaches such as Dynamo[38], which can take advantage of controlled metabolic labeling to predict past

and future cell states over an extended time period, high-throughput single-cell nascent transcriptomic

analysis methods such as scNT-Seq and sci-fate can open new lines of inquiry regarding cell-type-

specific RNA regulatory mechanisms and provide a broadly applicable strategy to investigate dynamic

biological systems.

ACKNOWLEDGEMENTS

We are grateful to all members of the Wu lab for helpful discussion. This work was supported by the Penn Epigenetics Institute, the National Human Genome Research Institute (NHGRI) grants R00-HG007982 and R01-HG010646, National Heart Lung and Blood Institute (NHLBI) grant DP2-HL142044, and National Cancer Institute (NCI) grant U2C-CA233285 (to H.W.). The work of P.C. is partially supported by Stand-Up-To-Cancer Convergence 2.0.

AUTHOR CONTRIBUTIONS

H.W., Q.Q., and P.H. conceived of and developed the scNT-Seq approach. Q.Q. conducted most of the experiments. P.H. generated *Tet*1/2/3 TKO mESC line and performed most of the computational analysis. K.G. and P.C. contributed to the statistical modeling of T-to-C substitution. Q.Q., P.H., and H.W. analyzed the results and wrote the manuscript, with contributions from all of the authors. H.W. supervised the project.

COMPETING FINANCIAL INTERESTS

The authors declare no competing interest.

Online Methods

Mouse embryonic stem cell cultures and 4sU labeling

*Tet*1/2/3 triple-knock out (*Tet*-TKO) J1 mESCs were generated by CRISPR/Cas9 genome editing as previously described[33] and the genotype was confirmed by Sanger sequencing and scRNA-Seq. Wild-type (WT) and *Tet*-TKO J1 mESCs (ATCC, SCRC-1010) were initially cultured in presence of Mitomycin C inactivated mouse embryonic fibroblasts on 0.1% gelatin-coated (Millipore, ES-006-B) 6-well plates in Dulbecco's Modified Eagle's Medium (DMEM) (Gibco, 11965084) supplemented with 15% fetal bovine serum (Gibco, 16000044), 0.1 mM nonessential amino acid (Gibco, 11140050), 1 mM sodium pyruvate (Gibco, 11360070), 2 mM L-glutamine (Gibco, 25030081), 50 µM 2-mercaptoethanol (Gibco, 31350010), 1 µM MEK inhibitor PD0325901 (Axon Med Chem, Axon 2128) and 3 µM GSK3 inhibitor CHIR99021 (Axon Med Chem, Axon 2128), and 1,000 U/mL LIF (Gemini Bio-Products, 400-495-7).

4-thiouridine (4sU) (Alfa Aesar, J60679) were dissolved in DMSO to make 1 M stock. Before 4sU labeling experiments, WT and *Tet*-TKO mESCs were passaged and cultured in feeder-free conditions (0.1% gelatin-coated plates) for 48 hrs. For 4sU labeling in WT and Tet TKO mESC cultures, the medium was replaced with fresh mESC medium supplemented with 100 µM 4sU. After 4 h, mESCs were rinsed once with PBS and dissociated with TrypLE-Express (Gibco, 12605010) for 5 min at 37°C. After neutralizing with culture medium, cells were pelleted at 1,000 rpm for 3 min. After cell counting with the Countess II system, the single cell suspension was diluted to 100 cells/uL with DPBS containing 0.01% BSA for scNT-Seq analysis.

Human K562 cell cultures and species mixing experiments

Human K562 cells (ATCC, CCL-243) were cultured in RPMI media supplemented with 10% FBS (Sigma, F6178). For species mixing experiments, the mESC or K562 media was replaced with media supplemented with 4sU (100 µM). After 4 h, the mESCs and K562 cells were rinsed with PBS and harvested for scNT-Seq analysis.

Mouse primary neuronal culture and stimulation

Mouse cortices were dissected from embryonic day 16 (E16) C57BL/6 embryos of mixed sex (Charles River). Cortical neurons were dissociated with papain (Worthington) and plated on 6-well plates (at a density of ~600,000 cells/well) coated with poly-ornithine (30mg/mL, Sigma, P2533). Mouse cortical neuronal cultures were maintained in neurobasal media (Gibco, 21103049) supplemented with B27 (Gibco, 17504044), 2 mM GlutaMAX (Gibco, 35050061), and 1X Penicillin/streptomycin (Gibco, 15140122).

After 4 days *in vitro* culture, primary cortical cultures were stimulated with a final concentration of 55mM potassium chloride (KCl) for various time (0, 15, 30, 60, and 120 minutes). Before neuronal activation, the fresh media supplemented with 4sU (200 µM) was added. After 4sU labeling (2 h), the cells were washed with PBS once and were digested with 0.05% Trypsin-EDTA (Gibco, 25300054) for 20 min at 37°C, then replaced the buffer with 1 mL of DPBS and dissociated cells with a cell-scraper. After cell counting with Countess II system, the single cell suspension was diluted to 100 cells/µL with DPBS containing 0.01% BSA for scNT-Seq analysis.

Cell fixation, cryopreservation and rehydration for scNT-Seq

The cell fixation was performed as previously described[39]. Cultured mESCs in 6-well plates were digested with TrypLE-Express and harvested as aforementioned. After washing once with DPBS, the cells were resuspended with 0.4 mL of DPBS containing 0.01% BSA. Split the cells to two 1.5 mL LoBind tubes (Eppendorf) and add 0.8 mL methanol dropwise at final concentration of 80% methanol in DPBS. After mixing and incubating the cell suspension on ice for 1 hour, store the fixed cells in LoBind tubes at -80°C freezer for up to one month. For rehydration, cells were kept on ice after moved from –80 °C and kept in the cold throughout the procedure. After the cells were spun-down at 1,000 g for 5 min at 4°C, Methanol-

471   PBS solution was removed and cells were resuspended in 1 mL 0.01% BSA in DPBS supplemented with
472   0.5% RNase-inhibitor (Lucigen, 30281-2). After cell counting with the Countess II system, the single cell
473   suspension was diluted to 100 cells/µL and immediately used for scNT-Seq analysis.
474

Pulse-chase experiment for RNA stability analysis

Remove the medium from plates and add mESC medium supplemented with 200 µM 4sU[5]. The fresh
medium with 4sU were changed every 4 h to enhance 4sU incorporation and mESCs were labeled for 24
hrs. After 4sU-labeling, the 4sU-containing medium was removed and cells were washed once with
DPBS. Then mESC medium containing 10 mM Uridine (Sigma, U6381) was added to the culture before
cells were harvested at different time-points ranging from 0.5 hour to 24 hours. Cells were fixed with
methanol as aforementioned and store at -80°C for future use.
482

scNT-Seq library preparation and sequencing

Drop-Seq was performed as previously described with minor modifications[10]. Specifically, the single cell
suspension was diluted to a concentration of 100 cells/µL in DPBS containing 0.01% BSA. Approximately
1.5 mL of diluted single cell suspension was loaded for each scNT-Seq run. The single-nucleus
suspension was then co-encapsulated with barcoded beads (ChemGenes) using an Aquapel-coated
PDMS microfluidic device (uFluidix) connected to syringe pumps (KD Scientific) via polyethylene tubing
with an inner diameter of 0.38 mm (Scientific Commodities). Barcoded beads were resuspended in lysis
buffer (200 mM Tris-HCl pH8.0, 20 mM EDTA, 6% Ficoll PM-400 (GE Healthcare/Fisher Scientific), 0.2%
Sarkosyl (Sigma-Aldrich), and 50 mM DTT (Fermentas; freshly made on the day of run) at a concentration
of 120 beads/µL. The flow rates for cells and beads were set to 3,200 µL/hour, while QX200 droplet
generation oil (Bio-rad) was run at 12,500 µL/h. A typical run lasts ~20 min.

Droplet breakage with Perfluoro-1-octanol (Sigma-Aldrich). After droplet breakage, the beads were
treated with TimeLapse chemistry to convert 4sU to cytidine-analog[6]. Briefly, 50,000-100,000 beads were
washed once with 450 µL washing buffer (1 mM EDTA, 100 mM sodium acetate (pH 5.2)), then the beads
were resuspended with a mixture of TFEA (600 mM), EDTA (1 mM) and sodium acetate (pH 5.2, 100
mM) in water. A solution of 192 mM $NaIO_4$ was then added (final concentration: 10 mM) and incubated at
45°C for 1 hour with rotation. The beads were washed once with 1 mL TE, then incubated in 0.5 mL 1 X
Reducing Buffer (10 mM DTT, 100 mM NaCl, 10 mM Tris pH 7.4, 1 mM EDTA) at 37°C for 30 min with
rotation, followed by washing once with 0.3 mL 2X RT-buffer.

Reverse transcription and exonuclease I treatment were performed, as previously described, with minor
modifications[10]. Specifically, up to 120,000 beads, 200 µL of reverse transcription (RT) mix (1x Maxima
RT buffer (ThermoFisher), 4% Ficoll PM-400, 1 mM dNTPs (Clontech), 1 U/µL RNase inhibitor, 2.5 µM
Template Switch Oligo (TSO: AAGCAGTGGTATCAACGCAGAGTGAATrGrGrG)[10], and 10 U/ µL
Maxima H Minus Reverse Transcriptase (ThermoFisher)) were added. The RT reaction was incubated at
room temperature for 30 minutes, followed by incubation at 42°C for 120 minutes. To determine an
optimal number of PCR cycles for amplification of cDNA, an aliquot of 6,000 beads (corresponding to
~100 nuclei) was amplified by PCR in a volume of 50 µL (25 µL of 2x KAPA HiFi hotstart readymix (KAPA
biosystems), 0.4 µL of 100 µM TSO-PCR primer (AAGCAGTGGTATCAACGCAGAGT[10], 24.6 µL of
nuclease-free water) with the following thermal cycling parameter (95°C for 3 min; 4 cycles of 98°C for 20
sec, 65°C for 45 sec, 72°C for 3 min; 9 cycles of 98°C for 20 sec, 67°C for 45 sec, 72°C for 3 min; 72°C
for 5 min, hold at 4°C). After two rounds of purification with 0.6x SPRISelect beads (Beckman Coulter),
amplified cDNA was eluted with 10 µL of water. 10% of amplified cDNA was used to perform real-time
PCR analysis (1 µL of purified cDNA, 0.2 µL of 25 µM TSO-PCR primer, 5 µL of 2x KAPA FAST qPCR
readymix, and 3.8 µL of water) to determine the additional number of PCR cycles needed for optimal
cDNA amplification (Applied Biosystems QuantStudio 7 Flex). We then prepared PCR reactions per total
number of barcoded beads collected for each scNT-Seq run, adding 6,000 beads per PCR tube, and ran
the aforementioned program to enrich the cDNA for 4 + 10 to 12 cycles. We then tagmented cDNA using
the Nextera XT DNA sample preparation kit (Illumina, cat# FC-131-1096), starting with 550 pg of cDNA
pooled in equal amounts, from all PCR reactions for a given run. Following cDNA tagmentation, we
further amplified the library with 12 enrichment cycles using the Illumina Nextera XT i7 primers along with

the P5-TSO hybrid primer (AATGATACGGCGACCACCGAGATCTACACGCC

TGTCCGCGGAAGCAGTGGTATCAACGCAGAGT*A*C)[10]. After quality control analysis using a

Bioanalyzer (Agilent), libraries were sequenced on an Illumina NextSeq 500 instrument using the 75- or

150-cycle High Output v2 or v2.5 Kit (Illumina). We loaded the library at 2.0 pM and provided Custom

Read1 Primer (GCCTGTCCGCGGAAGCAGTGGTATCAACGCAGAGTAC) at 0.3 µM in position 7 of the

reagent cartridge. The sequencing configuration was 20 bp (Read1), 8 bp (Index1), and 60 or 130 bp

(Read2).

Then reverse transcription, exonuclease treatment, cDNA amplification and tagmentation were performed

as previously described[11]. The libraries were diluted to 2.0 pM, and subjected to paired-end sequencing

on Illumina NextSeq 500 sequencer as described previously[11].

SLAM-Seq reaction on barcoded Drop-Seq beads

After droplet breakage, the beads were washed once with $NaPO_4$ buffer with 30% DMSO (50 mM, pH

8.0), then incubate beads in 500 µL reaction-mix containing 10 mM IAA for either 15 min at 50°C

(standard SLAM-Seq) or 1 hour at 45°C (modified condition)[5]. Stop reaction by adding 10 µL 1 M DTT

(final concentration: 20 mM).

Bioinformatic analysis

Read mapping and quantification of labeled and unlabeled transcripts

Paired-end sequencing reads of scNT-Seq were processed as previous described[11] with some

modifications. Briefly, each mRNA read (read2) was tagged with the cell barcode (bases 1 to 12 of read

1) and unique molecular identifier (UMI, bases 13 to 20 of read 1), trimmed of sequencing adaptors and

poly-A sequences, and aligned using STAR v 2.5.2a to the mouse (mm10, Gencode release vM13),

human genome (GRCh38, Gencode release v23), or a concatenation of the mouse and human (for the

species mixing experiment) reference genome assembly. Both exonic and intronic reads mapped to

predicted strands of annotated genes were retained for the downstream analysis. To qualify the labeled

and unlabeled transcripts, uniquely mapped reads with mapping score > 10 were grouped by UMI

barcodes in every cell and were used to determine the T > C substitution using sam2tsv[40], T > C

substitutions with a base quality of Phred score > 27 were retained. For each experiment, locus with

background T > C substitutions (detected in the sample without TFEA/$NaIO_4$ treatment) was determined

and was excluded for T > C substitution identification. After background correction of T > C substitution, a

UMI was defined as labeled if there is a T > C substitution in any one of the reads belongs to that

particular UMI. To this end, every UMI will be assigned to labeled or unlabeled based on the existence of

T > C substitution (**Figs 1d and 5d**). For each transcript, the total number of uniquely labeled and

unlabeled UMI sequences were counted and finally were assembled into a matrix using gene name as

rows and cell barcode as columns.

Cell type clustering

For mouse cortical neurons and RNA-decay experiment (**Figs 2b** and **6b**), the raw digital expression

matrices of nascent and pre-existing UMI counts were added up and loaded into the R package Seurat[41]

(v 2.3.4). For normalization, UMI counts for all cells were scaled by library size (total UMI counts),

multiplied by 10,000 and transformed to log space. Only genes detected in >10 cells were retained. Cell

with a relatively high percentage of UMIs mapped to mitochondrial genes were discarded (QC metrics in

Supplementary Table 1). Moreover, cells with lower or higher detected genes were discarded (QC

metrics in **Supplementary Table 1**). As a result, 20,547 cells of mouse cortical culture and 20,190 cells of

mESC (in pulse-chase assay) were retained, respectively. The highly variable genes (HVGs) were

identified using the function *FindVariableGenes* with the parameters: *x.low.cutoff* = .05, *y.cutoff* = .5 in

Seurat, resulting in 2,290 HVGs of primary cortical culture sample and 2,165 HVGs of mESC sample. The

expression level of highly variable genes in the cells was scaled and centered along each gene and was conducted to principal component analysis (PCA). The most significant 30 PCs were selected and used for 2-dimension reduction by uniform manifold approximation and projection[42] (UMAP), implemented by the Seurat software with the default parameters. Clusters were identified using the function *FindCluster* in Seurat with the resolution parameter set to 1. To obtain high level of cell type classification, we merge the adjacent clusters in UMAP which highly expressed excitatory markers (*Neurod2* and *Neurod6*) and define it as "Ex" cluster in mouse cortical neurons, while several close clusters highly expressed *Sox2* were combined to "pluripotent" cluster in mESC Pulse-chase experiment. Cell type specific markers were identified using function *FindMarkers* in Seurat with wilcoxon rank sum test with default parameters.

Cell clustering will be affected both by genotype[28] and by cell type. To enable directly comparative analyses within cell types between WT and *Tet*1/2/3 triple-knock out (*Tet*-TKO) mESCs, we used Seurat 3 (v. 3.0.0.9000) which was demonstrated as an effective strategy to perform joint analyses[43] (**Supplementary Fig. 6a**). The raw digital expression matrices of nascent and pre-existing UMI counts were added up and loaded into the Seurat 3. For normalization, UMI counts for all cells were scaled by library size (total UMI counts), multiplied by 10,000 and transformed to log space. Only genes detected in >10 cells were retained. Cell with a relatively high percentage of UMIs mapped to mitochondrial genes (>=0.05) were discarded. Moreover, cells with fewer than 500 or more than 5,000 detected genes were discarded, resulting in 4,633 WT cells and 2,319 *Tet*-TKO cells. Top 2,000 HVGs were identified using the function *FindVariableFeatures* with "vst" method. Canonical correlation analysis (CCA) was used to identify common sources of variation between WT and *Tet*-TKO cells. The first 20 dimensions of the CCA was chosen to integrate the two datasets. After integration, the expression level of HVGs in the cells was scaled and centered along each gene and was conducted to PCA analysis. The 20 most significant PCs were selected and used for 2-dimension reduction by UMAP. Clusters were identified using the function *FindCluster* in Seurat with the resolution parameter set to 3. As above mentioned, adjacent clusters highly expressed *Sox2* were combined to "pluripotent" cluster.

Estimation of the fraction of nascent transcripts

We implemented a statistical modeling strategy as previously described with some modifications for UMI-based scNT-Seq analysis[6]. For each experiment, a binomial mixture model, was used to approximate the number of T-to-C substitutions y_i for each gene transcript i:

$$f(y_i|\theta, p, q) = \theta \, \text{Binom}(y_i; p, n_i) + (1 - \theta) \, \text{Binom}(y_i; q, n_i)$$

In this expression, θ is the fraction of nascent transcripts in each experiment, p and q are the probabilities of a T-to-C substitution at each nucleotide for nascent and pre-existing transcripts, respectively, and n_i is the number of uridine nucleotides observed in the transcript i. A consensus sequence for each transcript is built by pooling reads with the same UMI barcode and taking the most frequent variant at each position. 10,000 UMIs were randomly sampled and the global substitution probabilities p and q were estimated based on the above mixture model. The model was fit by maximizing the likelihood function using the Nelder-Mead algorithm. The optimization was repeated ten times with random initialization values for θ, p, and q in the range [0,1], keeping the best fit with $\theta \in [0,1]$.

For mouse cortical neuronal culture datasets, besides Ex and Inh1, we respectively combined 4 inhibitory neuronal clusters (Inh 2-4 and Inh-NP) and 3 non-neuronal clusters (Ex-NP1, Ex-NP2 and RG) to obtain enough UMI for global parameters estimation at each time-point. By doing this, Ex, Inh, and 2 combined clusters were subjected to statistical modelling. For mESC data sets, we assumed that *Tet*-TKO will not affect 4sU utilization efficiency and thus combined WT and TKO data sets to estimate unified global parameters, *p* and *q*, for 3 cell states (pluripotent, intermediate and 2C-like). To this end, 20 sets of *p* and *q* (5 time-points X 4 combined clusters) were determined in cortical neuronal data sets and 3 sets of *p* and *q* (pluripotent, intermediate and 2C-like clusters) were estimated for mESC data sets. Once these global parameters were determined, they were used to estimate the fraction of nascent transcripts.

1.   *Computing aggregate nascent transcript for cell clusters*

For Figs 2C and S3d, we aggregate all the UMIs belongs to the same cell-type and estimate the fraction
of nascent transcripts θ for each gene with more than 100 UMIs in that cell type at each time-point. The
likelihood function for the mixture model above was maximized using the Brent algorithm with the
constraint $\theta \in [10^{-8}, 1]$. The 95% confidence interval was calculated from the Hessian matrix, and θ
estimates for genes with a confidence interval greater than 0.2 were thrown out. The nascent transcript
(N) was then estimated:

$$N = \theta(L + U)$$

Where θ is the nascent fraction of a gene in a cell type, *L* is labeled transcripts of a gene, *U* is unlabeled
transcripts of a gene. The pre-existing transcript was calculated by: $(1 - \theta)(L + U)$.

2. Computing nascent transcript for each individual cell

In theory, for every cell the fraction of nascent transcripts should be calculated for each gene, which
requires adequate gene coverage of each individual cells as bulk RNA-seq analysis. However, for data
sets generated by high-throughput droplet-based scRNA-Seq methods and shallow sequencing, it is not
feasible to obtain sufficient coverage for every gene in individual cells. Moreover, modeling every gene in
thousands of cells were computationally intensive. To obtain the single-cell level estimation of nascent
transcripts in each cell, we introduced detection rate α to estimate the nascent transcript for each cell.
Since 4sU incorporation is random but each cell may vary in 4sU incorporation and most genes have the
highly similar detection rate α (**Supplementary Fig. 3d**), we used the strategy above to estimate the
fraction of nascent transcripts for individual cell. The detection rate α was then computed by diving all the
labeled transcripts of a cell by estimated nascent transcript of that cell. We discarded the cells without-
range values ($\alpha > 1$). We computed the detection rate α for 88.3% (18,133/20,547) of mouse cortical
cells, whereas 95.1% (6,609/6,952) of mESC cells were retained. The mean of detection rates α were
60% and 66% in cortex and mESC, respectively. For each individual gene, the nascent transcript was
computed according to the formula:

$$N = \min\left(\frac{L}{\alpha}, (L + U)\right)$$

Where α is the T-to-C detection rate of a cell, *L* is the number of labeled transcripts of a gene in that cell,
U is the number of unlabeled transcripts of a gene. The number of pre-existing transcripts was calculated
by: $L + U - N$. The computed nascent and pre-existing transcripts were used for all downstream single-
cell-based analysis, including stacked UMAP (**right panel of Fig. 2b** and **Supplementary Fig. 4a**),
SCENIC (**Figs. 2d, 3c, 4f, Supplementary Fig. 5** and **Supplementary Fig. 8a**) and nascent RNA
velocity analysis (**Figs. 3a** and **5b**).

Gene ontology (GO) enrichment analysis

GO enrichment analysis was performed as previously described[12]. To identify functional categories
associated with defined gene lists, the GO annotations were downloaded from the Ensembl database. An
enrichment analysis was performed via a hypergeometric test. The *P* value was calculated using the
following formula:

$$P = 1 - \sum_{i=0}^{m-1} \frac{\binom{M}{i}\binom{N-M}{n-i}}{\binom{N}{n}}$$

where N is the total number of background genes, n is the total number of selected genes, M is the
number of genes annotated to a certain GO term, and i is the number of selected genes annotated to a
certain GO term. *P* value was corrected by function *p.adjust* with false discovery rate (FDR) correction in
R. GO terms with FDR<0.05 were considered enriched. All statistical calculations were performed in R.

For enrichment analysis of stable/unstable mRNAs (**Supplementary Fig.10c**), genes were ranked by the
RNA half-life. Top 10% genes with longest half-lives were defined as stable genes, whereas top 10%
genes with shortest half-life were considered as unstable. Then the stable and unstable gene sets were

subjected into GO enrichment analysis. For **Fig. 4d**, genes showed >1.5-fold changes between pluripotent and 2C-like state were selected and subjected into GO enrichment analysis.

Identification of differentially expressed genes (DEGs)

Differential gene expression analysis of nascent transcripts between different time-points of neuronal activation (15, 30, 60 and 120 min) and control (0 min) was performed using the function *FindMarkers* in Seurat, using a Wilcoxon rank sum test. Nascent transcripts with a fold-change of more than 1.5 and an adjust *P*-value less than 0.05/4 were considered to be differentially expressed (**Supplementary Fig. 4b**). Neuronal induction genes were defined if a nascent transcript was significantly increased in at least one time-point with KCl stimulation in any one of cell-types (**Fig. 2c and Supplementary Fig. 4b**). For comparison among 3 cell states within WT mESCs (**Fig. 4e**), genes encoding DNA binding proteins with a fold-change of nascent transcript expression level more than 0.25 (log-scale) and an adjust *P*-value less than 0.2 were listed.

Estimation of RNA half-life

For each gene, we separately aggerate labeled and unlabeled UMI counts in each cell state (**Fig. 6** and **Supplementary Fig.11**). Then the fraction of labeled transcripts was calculated with summed labeled UMI counts divided by total UMI counts (labeled + unlabeled). The fraction of labeled transcripts was normalized to chase-onset (0 min). R function *nls* was used to perform curve fitting with the parameters setting: "*y ~ I(a*exp(-b*x)+c)*", "*start=list(a=1, b=1, c=1)*" and "*na.action=na.exclude*". We kept the fit with the goodness of $R^2 > 0.7$.

RNA velocity analysis

For standard RNA velocity analysis (splicing RNA velocity), we started with the bam files which were generated by the Drop-Seq computational analysis pipeline. The reads were demultiplexed using dropEst[44] (version 0.8.5) pipeline, using "-m -V -b -f -L eiEIBA" options to annotate bam files. The genome annotations (mm10, Gencode release vM13) were used to count spliced and unspliced molecules for each experiment. The python package scVelo[21, 45, 46] (https://github.com/theislab/scvelo, version 0.1.19) were employed to perform RNA velocity analysis. Default parameter settings were used, unless stated otherwise. After loading spliced and unspliced molecules to scVelo, for analysis of excitatory neuron data sets **(left panel of Fig 3a)**, genes with less than 5 counts of spliced or unspliced molecules were filtered out. Spliced and unspliced counts of top 1,000 highly variable genes were KNN-imputed in a PCA reduced space with 10 components, using 15 neighbors and 'distances' mode. For nascent velocity analysis (using the ratio of nascent over total transcripts), all the parameters are the same as splicing RNA velocity analysis except that we loaded computed nascent transcripts (as spliced counts) and total transcript (as unspliced counts) into scVelo (**right panel of Fig 3a**). For nascent RNA velocity analysis of mESCs (**Fig 5a**), top 3,000 highly variable genes were KNN-imputed in with 10 PCs, using 15 neighbors and 'distances' mode as well. The gene-specific velocities are obtained by fitting a ratio between nascent and total mRNA by function '*scv.tl.velocity*'. Finally, function '*scv.pl.velocity_embedding_stream*' was used to project the velocities onto UMAP.

To directly compare splicing and nascent RNA velocity, we asked which of them could best predict whether well-known activity induced genes upregulated with different durations of KCl stimulation in excitatory neurons. The list of 137 primary response genes was previously defined by bulk RNA-Seq[16]. We calculated the average induction fold of genes (expressed in >1% cells) at 30 min of KCl stimulation compared to 0 min (**Fig 3b**). For splicing RNA velocity, we used unspliced counts to calculate the induction fold while estimated nascent transcript were used in nascent RNA velocity analysis. Finally, *auc* function in R package pROC was conducted to calculate the area under receiver operating characteristic curve (AUC), which reflects the accuracy of predicting the known PRGs by gene induction fold computed in different method. The plots were generated by function *plot.roc* in pROC package[47].

Gene regulatory network (GRN) analysis by SCENIC

To assess the regulatory activity of transcript factors associated with different cell states or cell-types, we used SCENIC[18] (version 1.1.2.2) to perform gene regulatory network (GRN or regulon) analysis. Regulatory modules are identified by inferring co-expression between TFs and genes containing TF binding motif in their promoters. We separate expression matrix into two parts based on the expression level of nascent and pre-existing transcripts, then combined them as inputs to SCENIC analysis, which enabled us to identify specific regulatory modules associated with either nascent or pre-existing transcriptomes from the same cell. Two gene-motif rankings, 10kb around the TSS and 500 bp upstream, were loaded from RcisTarget databases (mm9). Gene detected in > 1% of all the cells and listed in the gene-motif ranking databases were retained. To this end, 8,744 genes in cortical culture data and 9,388 mESC genes were subjected into the downstream analysis. Then GRNBoost, which was implemented in pySCENIC, was used to infer the co-expression modules and quantify the weight between TFs and target genes. Targets genes that did not show a positive correlation (> 0.03) in each TF-module were filtered out. SCENIC found 4,944 and 5,406 TF-modules in mouse cortical neuronal culture and mESC data sets, respectively. A *cis*-regulatory motif analysis on each of the TF-modules with RcisTarget revealed 277 and 325 regulons in cortical culture and mESC data, respectively. The top 1 percentile of the number of detected genes per cell was used to calculate the enrichment of each regulon in each cell. For **Figs. 2d, 4f, and Supplementary Fig. 8**, we computed the mean AUC of all cells belongs to defined groups, then scaled the mean AUC by function *scale* in R. R package pheatmap[48] was used to draw the heatmap.

For **Fig. 4f and Supplementary Fig. 8**, AUC values of TFs were obtained and then subjected to Wilcoxon rank sum test to access significance of the difference of TF activity. TFs with a fold-change of mean AUC values more than 1.5 and an adjust *P*-value (Bonferroni corrected) less than 0.05 were considered differentially regulated.

RNA binding protein motif enrichment analysis

For **Fig. 6f**, 3' UTR sequences of 552 genes were retrieved from Ensembl Biomart[49]. All the 3' UTR sequences of a gene were obtained to explore all possible binding site. The position weight matrix (PWM) of 188 mouse RNA binding protein motifs were downloaded from CISBP-RNA database[50] (http://cisbp-rna.ccbr.utoronto.ca/). FIMO[51] (version 5.0.5) motif scanning software in MEME Suite was used to search for the RNA binding motifs in the 3' UTR sequences. 3' UTR sequences with *P*- values smaller than the default threshold (0.0001) were considered to contain a binding motif. As a result, 533 genes were identified to have RNA binding motif in their 3' UTR. Hypergeometric test was performed in R by function *phyper*, which was used to access significance of RNA binding motif enrichment in defined gene sets. The pairs of RNA binding protein and targets were visualized in Cytoscape.

Data availability

All data will be available through from the Gene Expression Omnibus (GEO) database.

Code availability

The analysis source code underlying the final version of the paper will be available on GitHub repository (https://gibhub.com/wulabupenn/scNT-seq).

REFERENCES

1. Rabani, M. et al. High-resolution sequencing and modeling identifies distinct dynamic RNA regulatory strategies. *Cell* **159**, 1698-1710 (2014).
2. Rabani, M. et al. Metabolic labeling of RNA uncovers principles of RNA production and degradation dynamics in mammalian cells. *Nat Biotechnol* **29**, 436-442 (2011).
3. Tanay, A. & Regev, A. Scaling single-cell genomics from phenomenology to mechanism. *Nature* **541**, 331-338 (2017).
4. Wissink, E.M., Vihervaara, A., Tippens, N.D. & Lis, J.T. Nascent RNA analyses: tracking transcription and its regulation. *Nat Rev Genet* **20**, 705-723 (2019).
5. Herzog, V.A. et al. Thiol-linked alkylation of RNA to assess expression dynamics. *Nature methods* **14**, 1198-1204 (2017).
6. Schofield, J.A., Duffy, E.E., Kiefer, L., Sullivan, M.C. & Simon, M.D. TimeLapse-seq: adding a temporal dimension to RNA sequencing through nucleoside recoding. *Nat Methods* **15**, 221-225 (2018).
7. Riml, C. et al. Osmium-Mediated Transformation of 4-Thiouridine to Cytidine as Key To Study RNA Dynamics by Sequencing. *Angew Chem Int Ed Engl* **56**, 13479-13483 (2017).
8. Erhard, F. et al. scSLAM-seq reveals core features of transcription dynamics in single cells. *Nature* **571**, 419-423 (2019).
9. Hendriks, G.J. et al. NASC-seq monitors RNA synthesis in single cells. *Nature communications* **10**, 3138 (2019).
10. Macosko, E.Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).
11. Hu, P. et al. Dissecting Cell-Type Composition and Activity-Dependent Transcriptional State in Mammalian Brains by Massively Parallel Single-Nucleus RNA-Seq. *Molecular cell* **68**, 1006-1015 e1007 (2017).
12. Hu, P. et al. Single-nucleus transcriptomic survey of cell diversity and functional maturation in postnatal mammalian hearts. *Genes Dev* **32**, 1344-1357 (2018).
13. Lake, B.B. et al. Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. *Nat Biotechnol* **36**, 70-80 (2018).
14. Habib, N. et al. Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nature methods* **14**, 955-958 (2017).
15. Yap, E.L. & Greenberg, M.E. Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018).
16. Tyssowski, K.M. et al. Different Neuronal Activity Patterns Induce Different Gene Expression Programs. *Neuron* **98**, 530-546 e511 (2018).
17. Duffy, E.E., Schofield, J.A. & Simon, M.D. Gaining insight into transcriptome-wide RNA population dynamics through the chemistry of 4-thiouridine. *Wiley interdisciplinary reviews. RNA* **10**, e1513 (2019).
18. Aibar, S. et al. SCENIC: single-cell regulatory network inference and clustering. *Nat Methods* **14**, 1083-1086 (2017).
19. Laclef, C. et al. The role of primary cilia in corpus callosum formation is mediated by production of the Gli3 repressor. *Hum Mol Genet* **24**, 4997-5014 (2015).
20. Machado, C.B. et al. Reconstruction of phrenic neuron identity in embryonic stem cell-derived motor neurons. *Development* **141**, 784-794 (2014).
21. La Manno, G. et al. RNA velocity of single cells. *Nature* **560**, 494-498 (2018).
22. Schwalb, B. et al. TT-seq maps the human transient transcriptome. *Science* **352**, 1225-1228 (2016).

23. Shah, S. et al. Dynamics and Spatial Genomics of the Nascent Transcriptome by Intron seqFISH. *Cell* **174**, 363-376 e316 (2018).

24. Kolodziejczyk, A.A. et al. Single Cell RNA-Sequencing of Pluripotent States Unlocks Modular Transcriptional Variation. *Cell stem cell* **17**, 471-485 (2015).

25. Macfarlan, T.S. et al. Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature* **487**, 57-63 (2012).

26. Eckersley-Maslin, M.A. et al. MERVL/Zscan4 Network Activation Results in Transient Genome-wide DNA Demethylation of mESCs. *Cell reports* **17**, 179-192 (2016).

27. Fu, X., Wu, X., Djekidel, M.N. & Zhang, Y. Myc and Dnmt1 impede the pluripotent to totipotent state transition in embryonic stem cells. *Nature cell biology* **21**, 835-844 (2019).

28. Lu, F., Liu, Y., Jiang, L., Yamaguchi, S. & Zhang, Y. Role of Tet proteins in enhancer activity and telomere elongation. *Genes Dev* **28**, 2103-2119 (2014).

29. Rodriguez-Terrones, D. et al. A molecular roadmap for the emergence of early-embryonic-like cells in culture. *Nature genetics* **50**, 106-119 (2018).

30. Nakai-Futatsugi, Y. & Niwa, H. Zscan4 Is Activated after Telomere Shortening in Mouse Embryonic Stem Cells. *Stem cell reports* **6**, 483-495 (2016).

31. Wu, H. & Zhang, Y. Reversing DNA methylation: mechanisms, genomics, and biological functions. *Cell* **156**, 45-68 (2014).

32. Pastor, W.A., Aravind, L. & Rao, A. TETonic shift: biological roles of TET proteins in DNA demethylation and transcription. *Nature reviews. Molecular cell biology* **14**, 341-356 (2013).

33. Wang, H. et al. One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* **153**, 910-918 (2013).

34. Russo, J., Heck, A.M., Wilusz, J. & Wilusz, C.J. Metabolic labeling and recovery of nascent RNA to accurately quantify mRNA stability. *Methods* **120**, 39-48 (2017).

35. Yi, H. et al. PABP Cooperates with the CCR4-NOT Complex to Promote mRNA Deadenylation and Block Precocious Decay. *Molecular cell* **70**, 1081-1088 e1085 (2018).

36. Cao, J., Zhou, W., Steemers, F., Trapnell, C. & Shendure, J. Characterizing the temporal dynamics of gene expression in single cells with sci-fate. *bioRxiv*, 666081 (2019).

37. Kiefer, L., Schofield, J.A. & Simon, M.D. Expanding the Nucleoside Recoding Toolkit: Revealing RNA Population Dynamics with 6-Thioguanosine. *Journal of the American Chemical Society* **140**, 14567-14570 (2018).

38. Qiu, X. et al. Mapping Vector Field of Single Cells. *bioRxiv*, 696724 (2019).

39. Alles, J. et al. Cell fixation and preservation for droplet-based single-cell transcriptomics. *BMC biology* **15**, 44 (2017).

40. Lindenbaum, P. Jvarkit: java utilities for bioinformatics. (2015).

41. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature Biotechnology* **36**, 411 (2018).

42. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).

43. Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* (2019).

44. Petukhov, V. et al. dropEst: pipeline for accurate estimation of molecular counts in droplet-based single-cell RNA-seq experiments. *Genome Biology* **19**, 78 (2018).

45. Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biology* **19**, 15 (2018).

46. Bergen, V., Lange, M., Peidli, S., Wolf, F.A. & Theis, F.J. Generalizing RNA velocity to transient cell states through dynamical modeling. *bioRxiv*, 820936 (2019).

47. Robin, X. et al. pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* **12**, 77 (2011).
48. Kolde, R. & Kolde, M.R. Package 'pheatmap'. *R Package* **1** (2015).
49. Kinsella, R.J. et al. Ensembl BioMarts: a hub for data retrieval across taxonomic space. *Database* **2011** (2011).
50. Ray, D. et al. A compendium of RNA-binding motifs for decoding gene regulation. *Nature* **499**, 172 (2013).
51. Grant, C.E., Bailey, T.L. & Noble, W.S. FIMO: scanning for occurrences of a given motif. *Bioinformatics* **27**, 1017-1018 (2011).

 FIGURE LEGENDS

Figure 1. Development and validation of scNT-Seq. (**a**) Overview of <u>s</u>ingle-<u>c</u>ell <u>N</u>ascent transcript
<u>T</u>agging sequencing (scNT-Seq). (**b**) Multi-species mixing experiment measures scNT-Seq specificity.
Mouse ESCs and human K562 cells were mixed at 1:1 ratio after 4sU labeling (100 µM, 4 hrs).
Scatterplot shows the number of transcripts (UMIs) mapped to mouse (y-axis) or human (x-axis) genome
for each cell (dot). Cells with mostly mouse transcripts are labeled as mouse (red), while cells with mostly
human transcripts are labeled as human (blue). Cells with a relatively high percentage of both mouse and
human transcripts are labeled as mixed (green). (**c**) Transcriptome-wide nucleotide substitutions rates
demonstrate specific T-to-C conversion in 4sU labeled K562 cells after TEFA/NaIO$_4$-treatment compared
to untreated control. (**d**) Box plots showing percentage of T-to-C containing transcripts (UMIs) in individual
K562 cells. The box plots display the median (center line) and interquartile range (IQR, from the 25th to
75th percentile), the whiskers represent 1.5 times the interquartile range, and the circles represent
outliers. (**e**) All UMIs of the *ACTG1* gene are shown for one K562 cell with or one without TFEA/NaIO$_4$-
treatment. Grey circles stand for T without T-to-C substitution, while crosses ("X"s) stand for sites of T-to-
C substitution in at least one read. The color of "X" indicates the read coverage in a UMI containing T-to-
C substitution at a particular site. All reads from one exemplary transcript (2nd UMI from TFEA/NaIO$_4$-
treated sample) are further shown in the lower panel.

Figure 2. scNT-Seq captures cell-type-specific nascent transcriptional dynamics and TF activity in
response to distinct neuronal activity patterns. (**a**) Schematics of applying scNT-Seq analysis to study
neuronal activation of primary mouse cortical cultures. (**b**) Uniform manifold approximation and projection
(UMAP) visualization of 20,547 cells from primary mouse cortical cultures. Relative percentage of each
cell-type is listed on the left. Randomly sampled 1,000 excitatory neurons from 5 time points (200 cells
from each time point) are sub-clustered based on nascent (top) or pre-existing transcripts (bottom). Cells
from each time point are color-coded and projected into the stacked UMAP on the right panel. The
nascent and pre-existing transcriptomes of same cell were connected by a black line. Ex, excitatory
neurons; Inh, inhibitory neurons; NP, neural progenitors; RG, radial glia. (**c**) Cell-type-specific responses
of activity-regulated genes (ARGs) upon neuronal activation. The mean nascent and pre-existing RNA
expression levels (transcripts per 10k, TP10k) were shown in line plot. (**d**) Heat map showing cell-type-
specific TF regulon activity (AUC value quantified by SCENIC) in response to distinct activity durations.
Red color highlights known regulators of ARGs, while blue color highlights potential novel regulators
induced by neuronal activity.

Figure 3. scNT-Seq reveals neuronal activity-induced nascent RNA velocity. (**a**) RNA velocity
analysis of excitatory neurons using ratios of unspliced-to-spliced transcripts (splicing RNA velocity, left
panel) or nascent-to-total transcripts (nascent RNA velocity, right panel) as inputs. The cells are color-
codeed by different neuronal activity durations. The arrow indicates the projection from the observed state
to extrapolated future state. (**b**) Comparison between splicing and nascent RNA velocity. Spliced counts
or nascent RNA counts (30 min KCl stimulation) were used to predict induction of 137 known primary
response genes[16]. The area under the curve (AUC) values of predictions from these two methods are
shown. (**c**) Enrichment analysis of TF regulons for early- or late-response genes. Bar plots showing the
significance of enriched TF regulons by hypergeometric test. TF regulons that are enriched for both
groups, only early- or only late-response genes are highlighted by three boxes (left to right). The cut off of
significance was indicated by red dash line (adjusted *P*-value < 0.001). (**d**) UMAP plots showing clustered
Ex neurons from five time-points (left), expression level of early-response (middle), and late-response
genes (right). (**e**) UMAP plots showing the regulon activity of two representative TFs regulating early-
(Jun) or late-response genes (Mef2d), respectively. The UMAP plots in Fig. 3d-e are the same as in Fig.
3a.

Figure 4. scNT-Seq captures both nascent and pre-existing transcripts in different states of
mESCs. (**a**) Schematics of the scNT-Seq analysis of mESCs. (**b**) MA-plot showing differential gene
expression of nascent and pre-existing transcripts between pluripotent and 2C-like states. Dashed line

denotes 1.5-fold differences. (**c**) The line plots showing nascent and pre-existing transcript levels of *Tet1* and *Zscan4d* in pluripotent, intermediate, and 2C-like state. (**d**) Heat map showing enriched GO terms of state-specific genes. Significance of enrichment (FDR) is scaled by colors. (**e**) Heat map showing nascent transcript levels of state-specific genes encoding DNA-binding proteins. (**f**) Heat map showing the regulon activity of state-specific TFs or epigenetic regulators. The color scales depict the regulon activity.

Figure 5. scNT-Seq reveals TET-dependent regulation of the plutipotent-to-2C transition. (*a*) Nascent RNA velocity analysis of combined wild-type (WT) and *Tet*-TKO mESCs. The arrow indicates the projection from the observed state to extrapolated future state. (**b**) Projection of cell state and the nascent RNA level of the *Zscan4a* gene on the same UMAP plot as in Fig. 5a. (**c**) Relative composition of three stem cell states in WT (4,633 cells) and *Tet*-TKO (2,319 cells) mESCs. (**d**) The fraction of labeled transcripts in WT and *Tet*-TKO mESCs across three states. The box plots display the median (center line) and interquartile range (IQR, from the 25th to 75th percentile), the whiskers represent 1.5 times the interquartile range, and the circles represent outliers.

Figure 6. Pulse-chase scNT-Seq reveals cell state-specific mRNA decay in mESCs. (**a**) Schematics of pulse-chase scNT-Seq experiment. (**b**) UMAP plots of 20,190 mESCs profiled in the pulse-chase scNT-Seq analysis. Cells are colored by three states (left) or by 7 time-points (right). Cell numbers of each state across different time-points are also shown. (**c**) Line plots showing temporal changes of nucleotide substitution rates. (**d**) Scatterplots showing the correlation of RNA half-life measurements between this study (top: one timepoint inference analysis; bottom: pulse-chase analysis) and bulk SLAM-Seq[5]. (**e**) Heat map (left) and bar plot (right) showing concordant change of RNA half-life (left) and gene expression (right) between pluripotent and 2C-like states, respectively. (**f**) Network of RNA-binding proteins (pink quadrate) and their binding targets (green circle). Enriched RNA-binding proteins were inferred by motif sequenced enriched in 3'UTR of target genes.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figure 1: Quality control of scNT-Seq in mESCs.

a. Violin plot comparing the number of genes and number of UMIs detected in individual cells among different treatment conditions. Cell number and sequencing depth were shown on the top.
b. Scatter plot showing reproducibility between two replicates.
c. Transcriptome-wide nucleotide substitution rates reveal the effect of labeling time (100 μM 4sU/4hr or 200 μM 4sU/24hr) and methanol fixation (two rehydration buffers, PBS versus SSC) on T-to-C conversion rate. The control sample (100 μM 4sU/4hr) was not treated with TFEA/NaIO$_4$.

Supplementary Figure 2: scNT-Seq identifies different cell types in primary mouse cortical cultures.

a. UMAP visualization of 20,547 cells from mouse cortical cultures (same UMAP plot as Fig. 2b). The cells are colored by different durations of neuronal activation.

b. Violin-plot showing gene expression levels of representative marker genes in different cell-types.

c. Marker gene expression is scaled by colors in the same UMAP as (a). Ex, excitatory neurons; Inh, inhibitory neurons; NP, neural progenitors; RG, radial glial progenitor cells.

Supplementary Figure 3: UMI-based statistical correction of the nascent RNA levels.

a. Density plot of the number of covered uridine sites per read or UMI. Data set of excitatory neurons with 60 min KCl stimulation were shown.

b. Bar plot of the number of T-to-C substitutions per read or UMI. Data set of excitatory neurons with 60 min KCl stimulation were shown.

c. All UMIs of the *Fos* and *Mapt* genes from one excitatory neuron with 60 min KCl stimulation were shown. Grey circles stand for T without T-to-C conversion, while crosses ("X"s) denote sites of T-to-C conversion in at least one read.

d. Comparison of nascent RNA levels of each gene in excitatory neurons (with 60 min KCl stimulation) before and after statistical correction. Ten neuron activity-dependent genes and two house-keeping genes are highlighted with red dots.

Supplementary Figure 4: Different cell types show different neuron activity-dependent gene expression programs.

a. UMAP visualization of the expression profiles of 20,547 cells from primary mouse cortical cultures with or without KCl stimulation (bottom panel, same UMAP plot as Figure 2b). Cells from non-neuronal cell clusters (RG/Ex-NP) are sub-clustered based on nascent or pre-existing mRNAs (top). The nascent and pre-existing mRNAs of same cell were connected by black line. Ex, excitatory neurons; Inh, inhibitory neurons; NP, neural progenitors; RG, radial glial progenitor cells.

b. Heat map showing nascent transcript levels of genes with significantly increased expression level after KCl stimulation in at least one cell type.

c. Heat map showing nascent transcript levels of early- and late-response genes in excitatory neurons with different durations of KCl stimulation. 97 significantly induced genes were clustered into 2 groups (early- and late-response).

Supplementary Figure 5: Heat map of activity-regulated TF regulons in mouse cortical cultures.

Heat map showing correlated transcription factor activity (quantified by SCENIC) during neuronal activity-dependent gene expression.

Supplementary Figure 6: scNT-Seq reveals different stem cell states in mESC cultures.

a. UMAP visualization of 4,633 WT mESCs from two replicates. The cells are colored by different cell-types or cell-states. Feeders (purple dots) are contaminated mouse embryonic fibroblasts in cell culture.

b. UMAP plots were colored by replicate.

c. Violin plot showing gene expression level of marker genes in different stem cell states or cell-types.

d. Feature-plots showing marker gene expression is scaled by colors in the same UMAP as (a).

Supplementary Figure 7: scNT-Seq reveals nascent and pre-existing RNA level of representative genes in mESCs.

a. Violin plots showing expression levels of nascent and pre-existing transcripts of selected genes in three cell-states of mESCs.

b. Scatterplots showing nascent and pre-existing RNA levels of representative genes, including pluripotent genes (*Sox2, Nanog*), 2C-like state specific genes (*Zscan4d, Gm8300*), and house-keeping gene (*Gapdh*).

c. Nascent and pre-existing RNA levels of major DNA methylation regulators in three stem cell states.

Supplementary Figure 8: scNT-Seq reveals TF activity changes during pluripotent-to-2C transition in mESCs.

a. Heat maps showing TF regulon activity (left) and the similarity of transcription factors activities in three stem cell states (right).

b. Volcano-plot showing state-specific TF regulon activity. TFs with a fold-change of mean AUC values more than 1.5 and adjust *P*-value less than 0.05 were highlighted in red.

Supplementary Figure 9: scNT-Seq analysis of TET-dependent regulation of the plutipotent-to-2C transition in mESCs.

a. Validation of genotypes of the *Tet1* (-11bp/+1bp) and *Tet2* (-7bp/-1bp) genes in *Tet*-TKO cells by aligning scNT-Seq reads to the CRISPR/Cas9 genome editing sites.
b. Projection of cell states onto the same UMAP plot as Fig. 5a.
c. Projection of genotypes and the cell-cycle states onto the same UMAP plot as Fig. 5a.

Supplementary Figure 10: Pulse-chase scNT-Seq reveals transcript-specific mRNA decay of mESCs.

a. Violin plots showing levels of labeled and total transcripts of the *Sox2* and *Top2a* genes during pulse-chase assay.
b. Marker gene expression is scaled by colors in the same UMAP plot as Fig. 6b.
c. Enrichment analysis of GO terms within stable (top 10% genes with longest half-lives) and unstable genes (top 10% genes with shortest half-life) in pluripotent state mESCs.

Supplementary Figure 11: Pulse-chase scNT-Seq reveals cell state-specific mRNA decay in mESCs.

Scatterplots comparing the RNA half-life of commonly detected transcripts between two stem cell states (**a**: pluripotent vs. intermediate; **b**: intermediate vs. 2C; **c**: pluripotent vs. 2C) and number of genes showing >1.5-fold change in RNA half-life (indicated by blue and red dashed line) between two states are shown (left panels). Representative genes showing state-specific RNA half-life are shown in the right panels.

Supplementary Figure 12: Characterization of mRNA stability and gene expression levels in different states of mESCs.

a. Scatterplot showing the correlation between gene expression levels (total transcripts) and RNA half-life in pluripotent and 2C-like states. Representative GO terms enriched in genes in each quadrant, as well as genes belong to these GO terms, are highlighted. The mRNA decay kinetics and expression level of selected 4 genes were also shown in the middle panel. Representative state-specific genes with different RNA half-life are shown in the right panel. 2C-like state enriched genes are highlighted in blue, while genes preferentially expressed in the pluripotent state are highlighted in red.

b-c. Scatterplot showing the correlation between gene expression levels (total transcripts) and RNA half-life in different states (b: Pluripotent vs. intermediate; c: intermediate vs. 2C-like). Genes that are differentially expressed in one stem cell state are highlighted in red or blue.