Incipient resistance to an effective pesticide results from genetic adaptation and the canalization of gene expression

Xiaoshen Yin[a,†], Alexander S. Martinez[a,†], Abigail Perkins[a,b], Morgan M. Sparks[a], Avril M. Harder[a], Janna R. Willoughby[a,c], Maria S. Sepúlveda[d], Mark R. Christie[a,d,1]

[a]Department of Biological Sciences, Purdue University; 915 W. State St., West Lafayette, Indiana 47907-2054 USA

[b]Department of Anatomy, Cell Biology & Physiology, Indiana University School of Medicine; 320 W. 15[th] St., Indianapolis, Indiana 46202-2266 USA

[c]School of Forestry and Wildlife Sciences, Auburn University; 602 Duncan Dr., Auburn, Alabama 36849 USA

[d]Department of Forestry and Natural Resources, Purdue University; 715 W. State St., West Lafayette, Indiana 47907-2054 USA

[†]Authors contributed equally

[1]To whom correspondence should be addressed to: Mark Christie, 915 W. State Street, Department of Biological Sciences, Purdue University, West Lafayette, IN 47907-2054. Phone: (765) 494-2070, E-mail: markchristie@purdue.edu.

Keywords: contemporary evolution, genetic resistance, incipient resistance, pesticides, sea lamprey, RNA-seq

<u>Abstract</u>

The resistance of bacteria, disease vectors, and pests to chemical controls has vast ecological,

economic, and human-health costs. In most cases, resistance is only detected after non-

susceptible phenotypes have spread throughout the entire population. Detecting resistance in its

incipient stages, by comparison, provides time to implement preventative strategies. Incipient

resistance (IR) can be detected by coupling standard toxicology assays with large-scale gene

expression experiments. We apply this approach to a system where an invasive parasite, sea

lamprey (*Petromyzon marinus*), has been treated with the highly-effective pesticide 3-

trifluoromethyl-4-nitrophenol (TFM) for 60 years. Toxicological experiments revealed that

lamprey from treated populations did not have higher survival to TFM exposure than lamprey

from their native range, demonstrating that full-fledged resistance has not yet evolved. In stark

contrast, we find hundreds of genes differentially expressed in response to TFM in the

population with the longest history of exposure, many of which relate to TFM's primary mode of

action, the uncoupling of oxidative phosphorylation. One gene critical to oxidative

phosphorylation, *ATP5PB*, which encodes subunit b of ATP synthase, was nearly fixed for

alternative alleles in comparisons between native and treated populations (F_{ST} > 9 SD from the

mean). A gene encoding an additional subunit of ATP synthase, *ATP5F1B*, was canalized for

high expression in treated populations, but remained plastic in response to treatment in sea

lamprey from the native range. These combined genomic and transcriptomic results illustrate that

an adaptive, genetic response to TFM is driving incipient resistance in a damaging pest species.

Introduction

The evolution of resistance often occurs in response to the continued and large-scale application of chemical compounds such as antibiotics, herbicides, and pesticides (1-4). The costs associated with the evolution of resistance measure in the billions of dollars per year and cut across diverse fields ranging from healthcare to agriculture (5-7). Increased rates of human mortality, decreased food security, and large-scale environmental costs represent only some of the negative effects when microbes, pathogens, and pests can no longer be effectively controlled (4, 8). In light of these costs, there is a considerable need to prevent or delay the evolution of resistance. Unfortunately, resistance is often detected too late, after most individuals in a population have acquired a resistant phenotype (9). Once this stage is reached, there is little to be done except to implement alternative control measures, which are often less effective (10, 11), or stop employing control measures until resistant genotypes are eliminated from the population, which can take tens to hundreds of generations (12-14).

One solution to mitigating the costs of resistance is early detection; identifying the evolution of resistance in its incipient stages can allow time for developing alternative control measures or implementing management actions that can prevent the onset of population-wide resistance (e.g., adaptive and integrative management strategies (15-17)). One approach to detecting incipient resistance is to screen a large fraction of the population for resistant genotypes or phenotypes. However, such measures can be costly and impractical because of the large sample sizes required to identify small numbers of fully resistant individuals. Detectability issues (e.g., false positives and negatives) compound this problem. Alternatively, the coupling of experimental, eco-toxicological assays with the identification of genomic and transcriptomic responses in populations with varied treatment histories can provide key insights into how and

whether resistance is beginning to evolve. We applied this design to an invasive vertebrate pest

species that has been controlled with a chemical pesticide for 60 years.

Throughout the Laurentian Great Lakes, invasive sea lamprey (*Petromyzon marinus*)

have negatively impacted native fish communities by feeding parasitically on adult fishes (18).

With the construction and improvement of shipping canals in the early 1900s, sea lamprey

expanded their range from the northwestern Atlantic Ocean to the Great Lakes (19) (Fig. 1).

Adult sea lamprey are hematophagous parasites of other fishes and are not host specific (Fig.

1D). The lack of host specialization, coupled with ample spawning habitat found throughout the

Great Lakes region, spurred the growth of parasitic sea lamprey populations into the millions of

individuals (19). High abundance of sea lamprey in the Great Lakes contributed to rapid and

substantial declines in native lake trout (*Salvelinus namaycush*) and other ecologically and

commercially important fishes (20, 21). In response to the effects of sea lamprey on native fish

and fisheries, there was an immediate and concerted effort to develop efficient means for

controlling sea lamprey populations in the Great Lakes.

One effort, initiated in the 1950s, involved testing over 4000 chemical compounds on sea

lamprey and other fish species (22). The organic compound 3-trifluoromethyl-4-nitrophenol

(hereafter, TFM) was found to effectively kill larval sea lamprey, but had no detectable effects

on other fishes at low concentrations (23, 24). The larval stage of sea lamprey (Fig. 1C) are

targeted because they are small (typically < 120 mm) and aggregate at high densities. The

primary mode of action for TFM is to uncouple oxidative phosphorylation (25, 26), and because

sea lamprey have high sensitivity to this mechanism, TFM is a very effective pesticide (Fig. 1B).

Since the application of TFM to locations with abundant larval lamprey, invasive sea lamprey

populations have declined by up to 90% (19, 27). However, TFM does not kill all larval sea

lamprey (28) and it is possible that resistant individuals could survive and reproduce. If this

process is repeated over enough generations, then resistant individuals could increase in

frequency – a scenario documented in many systems where pests have been controlled by

chemical means (29). The evolution of resistance would greatly reduce the effectiveness of TFM,

which is currently the primary sea lamprey control method (28), and decreased pesticide

effectiveness would likely hamper the restoration of native and commercially-important fish

populations throughout the Great Lakes.

To test for the evolution of resistance, we first collected larval sea lamprey from three

populations with varied histories of TFM treatment (Fig. 1A): Lake Michigan (56 years of TFM

treatment), Lake Champlain (26 years of TFM treatment), and Connecticut River (native range; 0

years of TFM treatment). Larvae were acclimated in a common environment for 4 months before

being experimentally exposed to TFM at lethal, sublethal, and control (no TFM) levels. Detailed

survival and toxicological analyses were coupled with tissue-specific RNA-seq. This procedure

was followed by the identification of differentially expressed genes, network analysis, and

identification of genomic regions under selection. Survival analyses revealed that the outright

resistance of sea lamprey to lethal concentrations of TFM could not be detected, but genomic and

transcriptomic analyses suggest that the early, incipient stages of resistance are well under way

in populations with a history of pesticide exposure.

Results

We performed multiple independent toxicological assays over a two-year period using larvae

collected from historically untreated (native range) and treated tributaries (Fig. 1A; Table S1).

The larvae collected from treated tributaries were not previously exposed to TFM prior to the

toxicological assays (i.e., any resistance, if detected, could not be due to inducible tolerance

(30)). If outright resistance has evolved, we would expect to observe higher survivorship in sea

lamprey collected from populations with a history of treatment (i.e., Lake Michigan or Lake

Champlain) in comparison to sea lamprey collected from areas that have never been treated (i.e.,

the native range, Connecticut River). We found no evidence to support this prediction. In 2016,

the mortality for treated individuals was highly similar across all populations over a 12-hour

exposure period (LM 96.4%; LC 100%; CT 91.1%; Fig. 2A, *SI Text*). In 2017, survivorship

through time again did not vary substantially among populations (Fig. S1) and estimates of lethal

concentrations required to kill 50% of the population (LC50) varied only slightly among

populations and trials (Fig. 2B, *SI Text*). Furthermore, no individuals from Lake Michigan

survived TFM concentrations greater than 2 mg/L (Fig. S1). When TFM is applied in the field,

stream concentrations of TFM range from 5.5 to 9 mg/L for the pH and alkalinity values used in

our experimental treatments (Fig. 2B, *SI Text*). In total, we performed 5 independent tests with 6

tested concentrations and never detected any meaningful differences in survival across

populations (Table S2, *SI Text*). From these analyses, we conclude that outright resistance to

TFM has not yet evolved, as no treated individuals survived TFM exposure at concentrations

they would be exposed to under ordinary stream treatment conditions.

By contrast, we observed remarkably large differences in patterns of gene expression

among sea lamprey from different populations exposed to the same, sublethal concentration of

TFM. Briefly, we exposed 168 larvae to either 0.2 or 0.3 mg/L of TFM for 6 hours (Table S2).

An equal number of control individuals per population were included (i.e., individuals not

exposed to TFM), individuals from all populations had the exact same acclimation history (4

months), and all populations were simultaneously included in each of two large experiments to

avoid confounding circadian (31) or environmental effects (see Methods, *SI Text,* and Tables S2-

S3). Remarkably, when comparing samples treated with 0.3 mg/L TFM to control samples, we

found 336 genes that were differentially expressed (275 upregulated, 61 downregulated in

comparison to control individuals) in muscle tissue in the Lake Michigan population, dwarfing

the number of differentially expressed genes (DEGs) found in both the Lake Champlain (n = 21)

and the Connecticut River (n = 68) populations (Fig. 3A-C; Table S4). No DEGs were shared

among all three populations and different population responses were observed when lower

concentrations (i.e., 0.2 mg/L) of TFM were applied and when different tissues were examined

(Figures S2-S5).

121   The primary mode of action for TFM is to uncouple oxidative phosphorylation in the

mitochondria resulting in severe ATP depletion and eventual death (25, 26, 32). Of the 336

DEGs identified in Lake Michigan, several genes were directly related to this mode of action

including *CRCM1* (LogFC = 2.60), a calcium release-activated channel protein that controls the

influx of calcium into cells when depleted, and *PLCD4* (LogFC = 2.41), an enzyme responsible

for hydrolyzing phosphatidylinositol 4,5-bisphosphate into two secondary messenger molecules,

one of which (inositol 1,4,5-trisphosphate or 'IP3') is known to release cellular calcium stores

and increase calcium concentrations within the cell (33). Upregulated expression of genes that

increase intracellular concentrations of calcium could enhance oxidative phosphorylation in Lake

Michigan larvae and partially compensate for the uncoupling effects of TFM (Fig. 4A). In

addition, calcium plays an integral role in activating and driving the oxidative phosphorylation

cascade in mitochondria, and studies have demonstrated a positive relationship between

nanomolar free calcium and the rate of phosphorylation (34, 35). In gene ontology (GO)

hierarchy networks, we also found regulation of calcium ion transport, regulation of calcium ion

transmembrane transport, regulation of calcium ion transmembrane transporter activity, and

regulation of voltage-gated calcium channel activity as significant biological processes (Fig. S6).

In Lake Champlain, a gene encoding ADP/ATP translocase3, *SLC25A6*, was significantly

downregulated in treated individuals (LogFC = -3.50). The downregulation of *SLC25A6* may

disrupt the exchange of ADP and ATP across mitochondrial membranes and inhibit apoptosis,

suggesting that individuals from Lake Champlain are retaining ATP within their mitochondria in

response to TFM (36, 37).

We next aligned our reads to the sea lamprey genome (38, 39), called single nucleotide

polymorphisms (SNPs), and identified outliers based on Z-scores of genetic divergence (Weir

and Cockerham's F_{ST}) (40, 41). Out of 28,315 SNPs, the single highest outlier, *ATP5PB*, was

found in comparisons between Lake Michigan and Connecticut River and had an F_{ST} value

greater than 9 standard deviations from the mean (Fig. 3D). *ATP5PB* encodes a subunit of ATP

synthase (subunit b) and, similar to many of the genes identified by expression analyses, appears

to be responding to strong selection in order to restore normal oxidative phosphorylation in

response to TFM's primary mode of action (Fig. 4A). Closer examination of the genotypes at the

tri-allelic SNP found in this gene illustrates that a single allele is nearly fixed in both Lake

Michigan and Lake Champlain, but only occurs rarely in the native range (Fig. 4B). This outlier

was consistently identified across genotype calling pipelines and tissue types (muscle and liver;

Methods, Figures S7-S9) and was also observed in comparisons between Lake Champlain and

Connecticut River (Figures S10-S11). Although the native-range, Connecticut River lamprey are

not fixed at the genetic level for this variant, they were able to respond plastically with respect to

TFM at ATP synthase. The gene *ATP5F1B*, which codes for subunit beta of ATP synthase, was

significantly upregulated in Connecticut River lamprey in response to TFM treatment (LogFC =

4.68, FDR-corrected *p*-value < 0. 0072, Fig. 4C). Interestingly, this gene was not upregulated in

control individuals from Connecticut River, suggesting a plastic response, but was upregulated in

both control and treatment individuals from Lake Michigan and Lake Champlain in comparison

to Connecticut River control individuals, suggesting a genetic response to constitutively increase

expression of this gene. Thus, here we show both genomic and transcriptomic evidence of

among-population variation in response to TFM; critically, the two populations with a lengthy

history of TFM treatment show adaptive responses in ATP synthase at the genomic level, while

the naïve, native population shows plastic responses in transcription of the same enzyme in

response to TFM.

Discussion

Here we provide evidence for the evolution of incipient resistance in invasive sea lamprey

treated with a pesticide. We found 336 genes differentially expressed in response to TFM in

larvae from Lake Michigan, the population with the longest history of using TFM (56 years).

This transcriptional response represents a nearly 5-fold greater number of differentially

expressed genes in comparison to larvae collected from the native range. The pesticide TFM's

primary mode of action is to serve as a protonophore, translocating protons across the inner

mitochondrial membrane, thus uncoupling oxidative phosphorylation and depleting ATP. Many

of the DEGs identified in Lake Michigan represent direct or indirect responses to ATP depletion,

TFM's primary mode of action. We also identified genomic evidence of a response to selection

imposed by TFM. The gene coding for ATP synthase subunit b, *ATP5PB*, had an F_{ST} value

greater than 9 standard deviations from the mean in comparisons between Lake Michigan and

Connecticut River (Fig. 3D) and greater than 7 standard deviations from the mean in

comparisons between Lake Champlain and Connecticut River (Fig. S10). These large shifts in

allele frequencies indicate a response to selection in a gene that is critical to successful oxidative

phosphorylation. We also identified one gene, *ATP5F1B*, that was differentially expressed in the

native Connecticut River population, but not the other, treated populations. Interestingly, this

gene is constitutively expressed at high levels in treated populations, suggesting that an

additional component of ATP synthase has responded to pesticide-induced selection.

Lake Michigan and Lake Champlain have been treated with TFM for 56 and 26 years,

respectively. Both show evidence of selection at *ATP5PB* and *ATP5F1B*, however, both

responded to TFM differently at the gene expression level with 336 and 21 genes differentially

expressed in Lake Michigan and Lake Champlain, respectively. These large differences in the

number of genes differentially expressed may reflect different evolutionary responses to TFM,

one of which is mediated exclusively at the genetic level and one that is mediated at both genetic

and transcriptomic levels. Alternatively, these differences may reflect the deeper evolutionary

histories between these two populations or the different duration and intensities of pesticide

usage (42). We also observed large differences in gene expression depending on the

concentration of TFM used and the tissue examined (Figures S2-S5). These similarities in

adaptive, genetic responses and differences in broad patterns of gene expression suggest that the

TFM-mediated selection results in a convergent, genetic response while the different

evolutionary and treatment histories of these populations affect overall patterns of gene

expression.

We identified large genetic and transcriptomic differences among populations but did not

detect any individuals able to survive concentrations of TFM applied to streams in practice (Fig.

2). Thus, we may be documenting the earliest stages of resistance evolving in treated populations

(i.e., incipient resistance). Survival and reproductive advantages by larvae exposed to sublethal

concentrations of TFM, for example individuals at the peripheries of TFM treatment or

individuals buried deeper in the sediment, are inherited by their offspring and have spread

throughout the treated populations. It may only be a matter of time before further mutations

allow individuals to survive higher concentrations of TFM. In fact, recent modelling efforts have

revealed that given the estimated strength of selection imposed by lamprey control, full-fledged

resistance to TFM is predicted to occur soon (12). Because of the high gene flow and lack of

population structure found throughout the Great Lakes (i.e., Lake Michigan is one, large

panmictic population, see *SI Text*), fully-resistant individuals may already be spreading

throughout the population even though it may take many more years for successful detection

(12). Alternatively, it may be that our toxicological assays were simply not sensitive enough to

detect resistance to lethal concentrations of TFM. From a pragmatic standpoint, the detection of

incipient resistance (IR) means that the development of alternative control measures and the

implementation of adaptive management strategies focusing on delaying resistance should be a

priority for the continued control of invasive sea lamprey. The detection of resistance in its

earliest stages, as documented here, allows time for such actions to assist with the continued

restoration of native and commercially-important fishes. Regardless of the system, the detection

of incipient resistance can greatly assist in efforts to mitigate and delay the costly consequences

of resistance.

Methods

Sample Collection and Experimental Design

We collected a total of 2,035 larval sea lamprey (ammocoetes) from five locations during the

summers of 2016 and 2017, focusing our sampling efforts on populations with different

durations of exposure to TFM (Fig. 1; Table S1). In 2016, we obtained 565 ammocoetes from the

Manistee River in Lake Michigan, 517 ammocoetes from Corbeau Creek and the LaPlatte River

in Lake Champlain, and 404 ammocoetes from the Connecticut River (Fig. 1). Lake Michigan

ammocoetes have been treated with TFM since 1960 (43), Lake Champlain ammocoetes have

been treated with TFM since 1990 (44), and the Connecticut River is part of the sea lamprey's

native range and has never been treated with TFM. All ammocoetes were collected using pulsed-

DC backpack electrofishers. To examine lamprey of similar ages and developmental stages, we

only collected ammocoetes from a narrow size range (80-120 mm). All ammocoetes were

collected within five calendar days (July 11-July 15, 2016) of each other and were transported to

Purdue's Aquaculture Research Lab (ARL) immediately after collection (see *SI Text* for housing

and acclimation details). In 2017, we collected an additional 284 larval lamprey from the Platte

River in Lake Michigan and 265 larval lamprey from higher up the Connecticut River (Turner

Falls, Massachusetts) to independently validate our 2016 toxicological analyses (Table S1-S2; *SI*

Text).

Lethal Exposure Trials

After a 4-month acclimation period, we constructed an experimental array to conduct exposure

trials in a temperature-controlled environmental chamber (12.8°C) consisting of 36, 2.5-gallon

glass aquaria (see *SI Text* for details). In 2016, we conducted two toxicological exposure trials to

directly assess whether sea lamprey populations exhibited differential mortality in response to

TFM exposure (see *SI Text* for details). To analyze survivorship data, we conducted a Kaplan-

Meier survival analysis that employed the log-rank test to quantify time to death for ammocoetes

in different populations (45). The Kaplan-Meier method generates an estimate of survival

probability (*S*) at time t_i and is calculated as:

249
$$S(t_i) = S(t_i - 1)(1 - \frac{d_i}{n_i})$$

where $S(t_i - 1)$ is the probability of being alive at time $t_i - 1$, n_i is the number of individuals

alive just prior to t_i, and d_i is the number of deaths occurring at t_i (45). We performed the survival

analysis in R 3.5.1, using both the SURVIVAL and OIsurv packages to create survival curves and

assess statistical differences between the survival curves of the three populations (46, 47). *P*-

values associated with pairwise tests used to determine which populations differ in their

survivorship were FDR adjusted using the Benjamini-Hochberg method (48).

To confirm these results, we conducted a second set of three toxicological assays in 2017,

using a modified design. These trials were organized to measure lethal concentration (or dose), a

different measure of phenotypic response to TFM. The 20-hour trials were conducted with a

single population at five different dosage levels (0, 0.5, 1.2, 1.5, and 2.0 mg/L). Trials were

limited to a single population to increase power for the detection of lethal concentrations in a

given population (see *SI Text* and Table S2 for details). To compare differences in survival,

lethal concentrations for each trial were computed in R using the DRC package (v 3.0-1) (49),

and comparisons were made among trials at different dosage levels using Kaplan-Meier survival

analyses in the SURVIVAL package (v 2.43-3) (47) with visualization using the SURVMINER

package (v 0.4.3) (50).

Sublethal Exposure Trials

To assess whether resistance was evolving at a sublethal level, we conducted an exposure trial to

assess *(i)* how gene expression profiles changed in response to TFM, and *(ii)* whether TFM

exposure altered patterns of gene expression differently among the three populations with

varying histories of exposure. The experimental design for sublethal exposure was similar to the

lethal exposure trials (outlined above) with two key differences: 1.) we exposed ammocoetes to

sublethal levels of TFM (Table S2) and 2.) we preserved tissue from ammocoetes at a

predetermined timepoint (t = 6 hr) during the 12-hour gene expression exposure trial for RNA-

seq. Sampling consisted of removing a single ammocoete per replicate tank and immediately

euthanizing it in a lethal dose of MS-222 before weighing, measuring, and flash-freezing in

liquid nitrogen. At the completion of the experiment, cryovials were directly transferred to a -

80°C freezer (see *SI Text* for details).

RNA-seq and Differential Gene Expression Analysis

We used tissue-specific RNA-seq to compare patterns of gene expression in ammocoetes from

our three populations with varying histories of exposure to TFM. Comparing tissue-specific

patterns of expression required that we first dissect specific tissue samples from the body

segments collected during the gene expression trial. Given TFM's established mode of action,

we targeted ammocoete muscle, liver, and brain tissues for downstream analyses of gene

expression profiles (see *SI Text* for details). RNA extractions were performed using a Qiagen

RNeasy Mini Kit. In total, we performed 71 RNA extractions across three populations (Lake

Michigan, Lake Champlain, and Connecticut River), and three tissue types (muscle, liver, and

brain) (Table S3). Library preparation and RNA sequencing (RNA-seq) were performed by the

Purdue Genomics Core Facility. We obtained a total of 3.25 billion 150bp paired-end reads that

we used to create a high-quality, annotated transcriptomes (see *SI Text* for details).

We aligned all reads to our assembled sea lamprey transcriptome and estimated transcript

abundances using RSEM (51). Differentially expressed genes (hereafter, DEGs) were identified

in R using edgeR's GLM functionality (46, 52, 53). We compared gene expression profiles of

treated versus untreated ammocoetes within each population to identify genes that were

significantly up- and down-regulated in response to TFM treatment and were specific to each

population. For genes to be classified as differentially expressed, we set minimum thresholds for

significance (FDR correction via the Benjamini-Hochberg method; *p*-value < 0.05). While we

did not filter DEGs by log fold change, the minimum log2-fold changes we observed for DEGs

were > 2 across all three populations (2.04, 3.50, and 2.74 for Lake Michigan, Lake Champlain,

and Connecticut River, respectively). Finally, GOseq was used to determine which gene

ontology functional categories were overrepresented among differentially expressed genes in

TFM treated individuals for each population-tissue combination (54). Only functionally enriched

categories significant below an FDR-adjusted *p*-value threshold of 0.05 were retained. We

constructed gene ontology (GO) hierarchy networks of the top 100 significant GO terms

identified with differentially expressed genes (FDR-corrected *p*-value = 0.01) using the

metacoder package in R (55).

Genomic Analysis

We called SNPs from our RNA-seq data (Table S3) following the methods and algorithms

provided by the Genome Analysis Toolkit (GATK – Calling Variants in RNA-seq, (56)) for

muscle (n = 44 samples) and liver (n = 19 samples) tissues separately. Variant calling details are

reported in *SI Text*. Since the GATK pipeline called variants for each sample separately, we

removed indels, and combined variants for all samples into a single variant-call-format file. We

filtered out loci genotyped at fewer than 80% of samples followed by samples genotyped at

fewer than 80% of loci. After filtering, we used the remaining variants as high-quality SNPs and

we kept 42 and 19 samples for muscle and liver tissues, respectively. In each population, we

further excluded loci out of Hardy-Weinberg Equilibrium (HWE), which we identified using the

R package HWxtest (likelihood ratio *p*-value < 0.05, (57)). A locus fixed for a single allele

within a population could represent a genotyping error that would not be removed with HWE

tests (e.g., misalignment, insufficient read depth). Thus, we next employed two genotype-calling

pipelines to create 4 SNP data sets, based on whether loci fixed for a single homozygote within a

population ("fixed homozygotes") were included or not: 1.) muscle – without fixed

homozygotes, 2.) muscle – with fixed homozygotes, 3.) liver – without fixed homozygotes, and

4.) liver – with fixed homozygotes. We calculated pairwise F_{ST} for each locus among all three

populations (i.e., LM vs. LC, LM vs. CT, and LC vs. CT populations) (40). To detect significant

outliers, we z-transformed F_{ST} in each comparison and defined loci with $Z(F_{ST})$ greater than five

standard deviations from the mean as outlier loci. We then identified the corresponding genes on

which outlier loci were located by mapping outlier loci to the sea lamprey genome (38, 39) and

calculated the frequency of all alleles at each outlier locus.

<u>Acknowledgments</u>

We thank B.J. Allaire, Bryan Apell, William Ardren, Steve Gephard, Nick Johnson, Robert Stira,

Bradley Young, the Conte Anadromous Fish Research Laboratory, Michigan DNR, Connecticut

Department of Energy and Environmental Protection, Hammond Bay Biological Station, and

U.S. Fish & Wildlife for assistance with collecting larval lamprey. We thank Lori Criger, Sam

Guffey, Elizabeth LaRue, Phillip San Miguel, Bob Rode, Mike Siefkes, Jeramiah Smith, Benson

Solomon, and the Purdue Genomics Core for assistance with experiments and analyses. Matthew

Byrnes, Matthew Campbell, Lindsey Dice, Jessica Elliott, Megan Gannon, Nathan Greiling,

Kimberly Gulbranson, Jason Jaworski, Tara Novak, Ashe Owens, Emily Reverman, and Cassidy

Robinson aided in the second set of toxicological trials. This research was funded by the Great

Lakes Fishery Commission Project ID: 2016_CHR_54053.

Author Contributions

MRC, ASM, and MSS designed the project, all authors provided logistic support, ASM

performed molecular analyses, XY, AMH, and ASM performed gene expression and genomic

analyses, ASM and MMS performed survival analyses, and XY, ASM, and MRC wrote the

manuscript with input from all authors.

Competing Financial Interests

The authors declare no competing financial interests.

References

1. G. P. Georghiou, C. E. Taylor, Genetic and biological influences in the evolution of insecticide
resistance. *Journal of economic entomology* **70**, 319-323 (1977).
2. C. Délye, M. Jasieniuk, V. Le Corre, Deciphering the evolution of herbicide resistance in weeds.
Trends in Genetics **29**, 649-658 (2013).
3. J. Davies, D. Davies, Origins and evolution of antibiotic resistance. *Microbiol. Mol. Biol. Rev.* **74**,
417-433 (2010).
4. F. Gould, Z. S. Brown, J. Kuzma, Wicked evolution: Can we address the sociobiological dilemma
of pesticide resistance? *Science* **360**, 728-732 (2018).
5. R. Smith, J. Coast, The true cost of antimicrobial resistance. *Bmj* **346**, f1493 (2013).
6. S. B. Levy, B. Marshall, Antibacterial resistance worldwide: causes, challenges and responses.
Nature medicine **10**, S122 (2004).
7. D. Pimentel, M. Burgess, "Environmental and economic costs of the application of pesticides
primarily in the United States" in Integrated pest management. (Springer, 2014), pp. 47-71.
8. H. Ranson, N. Lissenden, Insecticide resistance in African Anopheles mosquitoes: a worsening
situation that needs urgent action to maintain malaria control. *Trends in parasitology* **32**, 187-
196 (2016).
9. M. McKenna, Antibiotic resistance: the last resort. *Nature News* **499**, 394 (2013).
10. C. Ghosh, P. Sarkar, R. Issa, J. Haldar, Alternatives to conventional antibiotics in the era of
antimicrobial resistance. *Trends in microbiology* (2019).
11. M. A. Peterson, A. Collavo, R. Ovejero, V. Shivrain, M. J. Walsh, The challenge of herbicide
resistance around the world: a current summary. *Pest Management Science* **74**, 2246-2259
(2018).
12. M. Christie, M. Sepulveda, E. Dunlop, Rapid resistance to pesticide control is predicted to evolve
in an invasive fish. *bioRxiv*, 718593 (2019).
13. D. I. Andersson, B. R. Levin, The biological cost of antibiotic resistance. *Current opinion in
microbiology* **2**, 489-493 (1999).
14. D. I. Andersson, D. Hughes, Antibiotic resistance and its cost: is it possible to reverse resistance?
Nature Reviews Microbiology **8**, 260 (2010).
15. J. L. Apple, R. F. Smith (1976) Integrated pest management. (Springer).
16. J. A. Stenberg, A conceptual framework for integrated pest management. *Trends in plant science*
22, 759-769 (2017).
17. D. Andow, A. Ives, Monitoring and adaptive resistance management. *Ecological Applications* **12**,
1378-1390 (2002).
18. A. Lawrie, The sea lamprey in the Great Lakes. *Transactions of the American Fisheries Society* **99**,
766-775 (1970).
19. B. Smith, J. Tibbles, Sea lamprey (Petromyzon marinus) in Lakes Huron, Michigan, and Superior:
history of invasion and control, 1936–78. *Canadian Journal of Fisheries and Aquatic Sciences* **37**,
1780-1801 (1980).
20. N. Baldwin, R. Saalfeld, M. Dochoda, H. Buettner, R. Eshenroder, Commercial fish production in
the Great Lakes 1867–2000. *Ann Arbor, Michigan: Great Lakes Fishery Commission. Available:
www. glfc. org.(September 2006)* (2002).
21. C. R. Bronte *et al.*, Fish community change in Lake Superior, 1970–2000. *Canadian Journal of
Fisheries and Aquatic Sciences* **60**, 1552-1574 (2003).
22. V. C. Applegate, *Toxicity of 4,346 chemicals to larval lampreys and fishes* (US Fish and Wildlife
Service, 1957).

23.   C. R. Middaugh, M. S. Sepúlveda, T. O. Höök, Growth and behavioral effects of the lampricide
395   TFM on non-target fish species. *Journal of Great Lakes Research* **40**, 1010-1015 (2014).
396   24.   T. D. Hubert, Environmental fate and effects of the lampricide TFM: a review. *Journal of Great*
397   *Lakes Research* **29**, 456-474 (2003).
398   25.   O. Birceanu, G. B. McClelland, Y. S. Wang, J. C. Brown, M. P. Wilkie, The lampricide 3-
399   trifluoromethyl-4-nitrophenol (TFM) uncouples mitochondrial oxidative phosphorylation in both
400   sea lamprey (Petromyzon marinus) and TFM-tolerant rainbow trout (Oncorhynchus mykiss).
401   *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **153**, 342-349
402   (2011).
403   26.   O. Birceanu, G. B. McClelland, Y. S. Wang, M. P. Wilkie, Failure of ATP supply to match ATP
404   demand: The mechanism of toxicity of the lampricide, 3-trifluoromethyl-4-nitrophenol (TFM),
405   used to control sea lamprey (Petromyzon marinus) populations in the Great Lakes. *Aquatic*
406   *toxicology* **94**, 265-274 (2009).
407   27.   J. W. Heinrich *et al.*, Sea lamprey abundance and management in Lake Superior, 1957 to 1999.
408   *Journal of Great Lakes Research* **29**, 566-583 (2003).
409   28.   E. S. Dunlop *et al.*, Rapid evolution meets invasive species control: the potential for pesticide
410   resistance in sea lamprey. *Canadian Journal of Fisheries and Aquatic Sciences* **75**, 152-168
411   (2017).
412   29.   M. Whalon, D. Mota-Sanchez, R. Hollingworth, Analysis of global pesticide resistance in
413   arthropods. *Global pesticide resistance in arthropods* **5**, 31 (2008).
414   30.   J. Hua, N. I. Morehouse, R. Relyea, Pesticide tolerance in amphibians: induced tolerance in
415   susceptible populations, constitutive tolerance in tolerant populations. *Evolutionary applications*
416   **6**, 1028-1040 (2013).
417   31.   A. Conesa *et al.*, A survey of best practices for RNA-seq data analysis. *Genome biology* **17**, 13
418   (2016).
419   32.   O. Birceanu *et al.*, The effects of the lampricide 3-trifluoromethyl-4-nitrophenol (TFM) on fuel
420   stores and ion balance in a non-target fish, the rainbow trout (Oncorhynchus mykiss).
421   *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **160**, 30-41 (2014).
422   33.   M. J. Berridge, Inositol trisphosphate and calcium signalling. *Nature* **361**, 315 (1993).
423   34.   B. D. Fink, F. Bai, L. Yu, W. I. Sivitz, Regulation of ATP production: dependence on calcium
424   concentration and respiratory state. *American Journal of Physiology-Cell Physiology* **313**, C146-
425   C153 (2017).
426   35.   B. Glancy, W. T. Willis, D. J. Chess, R. S. Balaban, Effect of calcium on the oxidative
427   phosphorylation cascade in skeletal muscle mitochondria. *Biochemistry* **52**, 2793-2809 (2013).
428   36.   A. Dabbaghizadeh *et al.*, Identification of proteins interacting with the mitochondrial small heat
429   shock protein Hsp22 of Drosophila melanogaster: Implication in mitochondrial homeostasis. *PloS*
430   *one* **13**, e0193771 (2018).
431   37.   O. Lytovchenko, E. R. Kunji, Expression and putative role of mitochondrial transport proteins in
432   cancer. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **1858**, 641-654 (2017).
433   38.   J. J. Smith *et al.*, Sequencing of the sea lamprey (Petromyzon marinus) genome provides insights
434   into vertebrate evolution. *Nature genetics* **45**, 415 (2013).
435   39.   J. J. Smith *et al.*, The sea lamprey germline genome provides insights into programmed genome
436   rearrangement and vertebrate evolution. *Nature genetics* **50**, 270 (2018).
437   40.   B. S. Weir, C. C. Cockerham, Estimating F-statistics for the analysis of population structure.
438   *evolution* **38**, 1358-1370 (1984).
439   41.   J. R. Willoughby, A. M. Harder, J. A. Tennessen, K. T. Scribner, M. R. Christie, Rapid genetic
440   adaptation to a novel environment despite a genome-wide reduction in genetic diversity.
441   *Molecular ecology* **27**, 4041-4051 (2018).

442   42.   A. Whitehead, B. W. Clark, N. M. Reid, M. E. Hahn, D. Nacci, When evolution is the solution to
443   pollution: Key principles, and lessons from rapid repeated adaptation of killifish (Fundulus
444   heteroclitus) populations. *Evolutionary applications* **10**, 762-783 (2017).
445   43.   D. S. Lavis, M. P. Henson, D. A. Johnson, E. M. Koon, D. J. Ollila, A case history of sea lamprey
446   control in Lake Michigan: 1979 to 1999. *Journal of Great Lakes Research* **29**, 584-598 (2003).
447   44.   J. E. Marsden *et al.*, Sea lamprey control in Lake Champlain. *Journal of Great Lakes Research* **29**,
448   655-676 (2003).
449   45.   E. L. Kaplan, P. Meier, Nonparametric estimation from incomplete observations. *Journal of the*
450   *American statistical association* **53**, 457-481 (1958).
451   46.   R. C. Team, R: A Language and Environment for Statistical Computing. *R Foundation for*
452   *Statistical Computing*, Vienna, Austria (2019).
453   47.   T. Therneau (2015) A Package for Survival Analysis in S. version 2.38.
454   48.   Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: a practical and powerful
455   approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*
456   **57**, 289-300 (1995).
457   49.   C. Ritz, F. Baty, J. C. Streibig, D. Gerhard, Dose-response analysis using R. *PloS one* **10**, e0146021
458   (2015).
459   50.   A. Kassambara, M. Kosinski, survminer: Drawing Survival Curves using 'ggplot2'. 2018. *URL*
460   *https://CRAN. R-project. org/package= survminer. R package version 0.4* **3**, 15.
461   51.   B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or without a
462   reference genome. *BMC bioinformatics* **12**, 323 (2011).
463   52.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential
464   expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).
465   53.   D. J. McCarthy, Y. Chen, G. K. Smyth, Differential expression analysis of multifactor RNA-Seq
466   experiments with respect to biological variation. *Nucleic acids research* **40**, 4288-4297 (2012).
467   54.   M. D. Young, M. J. Wakefield, G. K. Smyth, A. Oshlack, Gene ontology analysis for RNA-seq:
468   accounting for selection bias. *Genome biology* **11**, R14 (2010).
469   55.   Z. S. Foster, T. J. Sharpton, N. J. Grünwald, Metacoder: An R package for visualization and
470   manipulation of community taxonomic diversity data. *PLoS computational biology* **13**, e1005404
471   (2017).
472   56.   A. McKenna *et al.*, The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
473   generation DNA sequencing data. *Genome research* **20**, 1297-1303 (2010).
474   57.   W. R. Engels, Exact Tests for Hardy–Weinberg Proportions. *Genetics* **183**, 1431-1441 (2009).
475

Figure Legends

Figure 1: Sample collection sites, sample sizes, and pesticide (TFM) treatment histories for

larval sea lamprey used in this study (A). Connecticut River sea lamprey were collected from

their native range whereas sea lamprey from Lake Michigan and Lake Champlain are considered

invasive. Sea lamprey abundance and metric tons of TFM applied to Lake Michigan from 1987

onwards (B); notice that TFM is an effective pesticide with decrease in lamprey abundance

following increases in TFM application. The larval stage of sea lamprey remain buried in the

sediment (C), aggregate in high densities, and are targeted with the pesticide TFM. After

transformation, the juveniles are generalist parasites, using their circularly-arrayed teeth to attach

to hosts and their rasping tongues to feed (D).

Figure 2: Results of toxicology assays illustrating survival through time and lethal

concentrations. In 2016, survival analyses revealed that previously unexposed larvae collected

from treated populations did not survive at higher rates than larvae collected from the native

population during exposure to lethal concentrations of TFM (A). In 2017, independent

toxicology assays from additional larvae collected from Lake Michigan (MI) and Connecticut

River (native range; CT) again revealed that larvae collected from populations with a history of

exposure did not survive lethal doses of TFM at higher rates than larvae collected from the native

range (B). Lethal concentrations for 25% and 50% mortality were calculated and two separate

assays were performed for Connecticut River individuals (CT1 and CT2, see *SI Text* for

additional analyses). Dashed green lines border the range of TFM values larvae would be

exposed to under typical stream application conditions. Collectively, these results illustrate that

resistance to lethal concentrations of TFM has not yet evolved.

Figure 3: Transcriptomic and genomic evidence of incipient resistance (IR) in sea lamprey. A

total of 336 genes were differentially expressed in response to sublethal concentrations of TFM

in larval sea lamprey collected from Lake Michigan, the population with the longest history of

TFM treatment (A). By contrast, only 21 and 68 genes were identified as differentially expressed

in lamprey from Lake Champlain (B) and Connecticut River (C), respectively. After aligning

reads back to the sea lamprey reference genome (sea lamprey have 99 chromosomes, 90 of

which are assembled), calling SNPs and calculating F_{ST}, the largest outlier SNP (9.1 standard

deviations greater than the mean) was found to belong to *ATP5PB*, a gene encoding subunit b of

ATP synthase (D), a putatively adaptive response to TFM's primary mode of action.

Figure 4: The pesticide TFM's primary mode of action is to act as a protonophore, translocating

protons from the inter-membrane space back into the inner mitochondrial matrix, thus disrupting

oxidative phosphorylation (A). Several genes upregulated in Lake Michigan, the population with

the longest history of exposure, may play a role in responding to TFM's mode of action. For

example, the upregulation of *CRCM1* and *PLCD4* increases intracellular concentrations of

calcium and could enhance oxidative phosphorylation in Lake Michigan ammocoetes (A). The

gene *ATP5PB*, which encodes subunit b of ATP synthase, was identified as an outlier (Fig. 3D),

is nearly fixed at a tri-allelic SNP in the treated Lake Michigan and Lake Champlain populations,

but segregates for largely different alleles in the native-range sea lamprey (B). A gene encoding

an additional subunit of ATP synthase, *ATP5F1B*, is canalized for high expression in treated

populations (i.e., the gene is expressed at high levels in both treatment (TFM+) and control

(TFM-) individuals from the treated Lake Michigan (LM) and Lake Champlain (LC)

populations) (C). By contrast, this gene is not constitutively upregulated in native-range,

Connecticut River (CT) lamprey, but responds plastically to treatment. Error bars depict standard

deviation, *p*-value calculated from FDR-corrected tests of differential gene expression.

Figure 1

Figure 2

Figure 3

Figure 4

(A) Upregulated Genes Increase ETS Activity

(B) ATP5PB

(C) ATP5F1B