Title: The evolutionary origins of extreme halophilic Archaeal lineages

Authors: Yutian Feng[1], Uri Neri[2], Sean Gosselin[1], Artemis S. Louyakis[1], R. Thane Papke[1], Uri

Gophna[2], J. Peter Gogarten[1,3]

[1]Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT, 06268, USA

[2]School of Molecular Cell Biology and Biotechnology, Tel Aviv University, P.O. Box 39040,

Tel Aviv 6997801, Israel

[3]Institute for Systems Genomics, University of Connecticut, Storrs, CT, 06268, USA

Correspondence: gogarten@uconn.edu

Running Head: Evolutionary origins of halophilic Archaea

Keywords: Nanohaloarchaea, Methanonatronarchaeia, gene concordance, metagenomic-

assembled genome (MAG), single amplified genome (SAG)

Significance Statement

Many recent analyses have considered large groups of Bacteria and Archaea composed

exclusively of environmentally assembled genomes as deep branching taxonomic groups in their

respective domains. These groups display characteristics distinct from other members of their

domain, which can attract unrelated lineages into those groups. This manuscript evaluates the

case of the Nanohaloarchaea, and their inclusion in the DPANN Archaea, through careful

analysis of the genes that compose the core of the Nanohaloarchaea. Analyses without inspection

of the genes that compose a phylogenomic marker set increases the potential for the inclusion of

artifacts and confuses the tree/web of life. Due to horizontal gene transfer and phylogenetic

reconstruction artifacts, the placement of divergent archaeal classes into larger groups remains

uncertain.

Abstract

Interest and controversy surrounding the evolutionary origins of extremely halophilic Archaea

has increased in recent years, due to the discovery and characterization of the Nanohaloarchaea

and the Methanonatronarchaeia. Initial attempts in explaining the evolutionary placement of the

two new lineages in relation to the classical Halobacteria (also referred to as Haloarchaea)

resulted in hypotheses that imply the new groups share a common ancestor with the Haloarchaea.

However, more recent analyses have led to a shift: the Nanohaloarchaea have been largely

accepted as being a member of the DPANN superphylum, outside of the euryarchaeota; while

the Methanonatronarchaeia have been placed near the base of the Methanotecta (composed of the

class II methanogens, the Halobacteriales, and Archaeoglobales). These opposing hypotheses

have far-reaching implications on the concepts of convergent evolution (unrelated groups evolve

similar strategies for survival), genome reduction, and gene transfer. In this work, we attempt to

resolve these conflicts with phylogenetic and phylogenomic data. We provide a robust

taxonomic sampling of Archaeal genomes that spans the Asgardarchaea, TACK Group,

euryarchaeota, and the DPANN superphylum. In addition, we assembled draft genomes from

seven new representatives of the Nanohaloarchaea from distinct geographic locations.

Phylogenies derived from these data imply that the highly conserved ATP synthase

catalytic/non-catalytic subunits of Nanohaloarchaea share a sisterhood relationship with the

Haloarchaea. We also employ a novel gene family distance clustering strategy which shows this

sisterhood relationship is not likely the result of a recent gene transfer. In addition, we present

and evaluate data that argue for and against the monophyly of the DPANN superphylum, in

particular, the inclusion of the Nanohaloarchaea in DPANN.

Introduction

Recent studies discovered several new archaeal lineages in hypersaline environments, including the nanosized Nanohaloarchaea and the methanogenic Methanonatronarchaeia. The exact placement of these lineages within the archaeal phylogeny remains controversial; consequently, the number of independent acquisitions of key adaptations to a halophilic lifestyle remains to be determined. Dissecting the evolutionary relationships between these new lineages and the Haloarchaea may inform on the origins of halophily and the role of genome streamlining. To thrive in extreme hypersaline environments (>150 g/L^{-1}), Haloarchaea employ a "salt-in" strategy through the import of potassium ions, in which the intracellular salt concentration equalizes with the external environmental condition (Oren, 2008). This acts to balance the cellular osmotic pressure but also has caused significant changes in amino acid usage, leading to an overabundance of acidic residues, aspartate and glutamate (D/E) in all Haloarchaea (Lanyi, 1974; Madern et al., 2000).

The evolutionary origins of the Nanohaloarchaea have remained uncertain since their discovery (Ghai *et al.*, 2011; Narasingarao *et al.*, 2012). The composition of their proteome indicates that Nanohaloarchaea also use the "salt-in" strategy similar to Haloarchaea (Narasingarao *et al.*, 2012). It was originally suggested that the Nanohaloarchaea are euryarchaeota that form a clade with the Haloarchaea, based on phylogenies of the 16S rRNA gene and ribosomal proteins (Narasingarao *et al.*, 2012; Petitjean *et al.*, 2014). Additional data obtained from individual cells via cell sorting followed by genome amplification and 16S rRNA sequencing analysis confirmed the original observations of the Nanohaloarchaea as a sister taxon to the Haloarchaea (Zhaxybayeva *et al.*, 2013). More recently, based on analyses of concatenated conserved protein sequences, the Nanohaloarchaea were placed in a group together with similarly nanosized organisms, the Diapherotrites, Parvarachaeota, Aenigmarchaeota, and Nanoarchaeota, forming the DPANN superphylum (Andrade *et al.*, 2015; Castelle *et al.*, 2015; Rinke *et al.*, 2013).

Past analyses of this superphylum (Brochier-Armanet *et al.*, 2011; Petitjean *et al.*, 2014; Raymann *et al.*, 2014; Williams *et al.*, 2015) suggested that the DPANN grouping may not reflect shared ancestry but rather an artifact due to long branches and/or small genomes. However, more recent analyses supported a monophyletic DPANN clade (Williams et al., 2017).

Aouad *et al.* performed a multi-locus analysis using various models, which did not include

DPANN sequences, and placed the Nanohaloarchaea with the Methanocellales and the

Haloarchaea with the Methanomicrobiales (Aouad *et al.*, 2018); *i.e.*, the Nanohaloarchaea were

recovered as a member of the euryarchaeota, but not as a sister-group to the Haloarchaea. We

note that a similar controversy surrounds the phylogenetic position of the Nanoarchaeota.

Nanoarchaeum equitans was first considered a representative of a new deep branching archaeal

phylum (Huber *et al.*, 2002), *i.e.,* an archaeon not a member of the euryarchaeotes or

crenarchaeotes. However, later analyses of ribosomal proteins, phylogenetically informative

HGTs, and signature genes led to the conclusion that *N. equitans* may represent a fast-evolving

euryarchaeote instead of an early branching novel phylum (Brochier *et al.*, 2005; Dutilh *et al.*,

2008; Urbonavičius *et al.*, 2008). Several more recent analyses placed the Nanoarchaeota inside

of the DPANN ((*Adam et al., 2017; Dombrowski et al*., 2019; Spang *et al*., 2017), reflecting the

ongoing controversy in the phylogenetic placement of these groups.

Recently, another group of extreme halophiles, the Methanonatronarchaeia (also spelled

as Methanonatronarcheia), were discovered and predicted to also use the "salt-in" strategy

(Sorokin *et al*., 2017). Initial multi-locus phylogenetic analyses placed these methanogenic

halophiles in a monophyletic clade with the Haloarchaea*,* suggesting they are an evolutionary

intermediate between methanogens and modern halophiles. However, several recent studies have

contested this placement: a multi-locus dataset placed the Methanonatronarchaeia basal to a

superclass named Methanotecta, a group that includes the Archaeoglobales, class II methanogens

and Haloarchaea (Adam *et al.*, 2017; Aouad *et al.*, 2019; Martijn *et al*., 2020).In addition, to the

three extreme halophiles mentioned, the recently characterized Hikarchaeia has been identified

as a non-halophilic sister-group to the Haloarchea (Martijn *et al*., 2020). Temporal analysis of

the Hikarchaeia divergence from the Haloarchaea may shed light on the genomic events that

prelude the Haloarchaea's adaptation to hypersalinity (see discussion).

Several conclusions can be drawn from these latter results with regard to adaptation to a

halophilic lifestyle, most note-worthy of which is the convergent evolution of the "salt-in"

strategy among these three lineages. Independent adaptation to hypersalinity in extreme

halophiles is certainly a viable evolutionary hypothesis; this is seen in the case of the

Salinibacter and *Salinicoccus* (Mongodin *et al*., 2005). However, if Nanohaloarchaea,

Haloarchaea, and Methanonatronarcheia form a monophyletic group, as seen with some analyses

of 16S rRNA and ribosomal proteins, the hypothesis of common ancestral origins can more

easily account for the evolutionary development of the salt-in strategy.

Fig 1. **Summary of proposed placements of halophilic lineages mapped on an Archaeal reference tree.** This reference tree mostly depicts the positions of various euryarchaea. Individual taxa have been collapsed into higher taxonomic groups. The red (R) indicators represent the different placements proposed for the Nanohaloarchaea, while the purple (P) indicators are used for the Methanonatronarchaeia. Sources for each placement: R1 (Narasingarao *et al*., 2012), R2 (Andrade *et al*., 2015), R3 (Aouad *et al*., 2018); P1 (Sorokin *et al*., 2017), and P2 (Aouad *et al.,* 2019).

123   The evolutionary relationships of the three extreme halophilic archaeal lineages remain

124 unresolved; Figure 1 summarizes the current controversies. This lack of resolution can, at least in

125 part, be due to biases that are known to complicate phylogenetics. The genomes of the

126 Methanonatronarchaeia and Nanohaloarchaea are comparatively small with average genome

127 sizes of <2.1Mb and ~1.1 Mb, respectively. Furthermore, most genome entries in public

128 databases are incomplete. The Haloarchaea are known to be highly recombinogenic (Boucher *et*

129 *al*., 2004; Méheust *et al*., 2018; Mohan *et al*., 2014; Naor *et al*., 2012; D. Williams *et al*.,2012)

130 and are physically associated with at least some of the Nanohaloarchaea (Andrade *et al*., 2015;

131 Cono *et al*., 2019; Hamm *et al*., 2019).

132   Phylogenies based on the concatenation of many genes face many problems: 1) Genes

133 have different evolutionary histories (e.g. duplication and transfer) and forcing the histories of all

134 the genes on a single tree does not reflect the complex evolutionary history of the genomes

135 (Lapierre *et al*., 2014). In particular, genes acquired from outside the group under consideration

136 may create a strong signal for placing the recipient of the transferred gene at the base of the

137 group. 2) Genes experience differing levels of purifying selection, especially between different

138 lineages. This can lead to long branch attraction (LBA) artifacts (Felsenstein, 1978), even if the

139 individual genes evolved along the same history as the host species (Philippe *et al.*, 2005). 3)

140 Substitution bias may create convergent signals in unrelated groups.

141   The work reported here was guided by the hypothesis that the phylogenetic

142 reconstruction of a single, slowly evolving gene might be more robust against artifacts of

143 phylogenetic reconstructions compared to analyses that are based on large sets of genes that may

144 represent different evolutionary histories, include missing data, and contain genes with high

145 substitution rates. We reconstruct single gene alongside multi-locus phylogenies to correct for

146 these sources of bias and to critically assess the evolutionary relationships of the Haloarchaea,

147 Nanohaloarchaea, and Methanonatronarchaeia. We also cluster and dissect the evolutionary

148 relationships of the gene families in the Nanohaloarchaeal core genome, using a gene family

149 clustering technique.

150   The ATP synthase catalytic and non-catalytic subunits, AtpA and AtpB, represent

151 extremely slow evolving genes (Gogarten, 1994) conserved throughout Archaea and are among

152 the slowest evolving genes in cellular organisms (Table S5). The evolution of these subunits may

be slow enough to ameliorate rate signal bias and minimize compositional heterogeneity that

otherwise plague reconstructions that includes DPANN and Haloarchaeal sequences. ATP

synthase subunits have been used successfully as a phylogenetic marker for large scale

reconstructions (Gogarten & Taiz, 1992); however, a drawback of the ATPases is that they are

known to have been transferred between divergent phyla (Olendzenski *et al*., 2000). Recently,

Wang *et al.* convincingly showed the transfer of this operon lead to the adaptation of

Thaumarchaeota to more acidic environments (Wang *et al.,* 2019). The same authors drew a

similar conclusion when the Nanohaloarchaea-Haloarchaea sister-group was recovered, which

the authors interpreted as suggesting HGT of the ATPase genes in the Nanohaloarchaea-

Haloarchaea. To shed light on this HGT hypothesis, we cluster and correlate the gene families in

the Nanohaloarchaea and contrast the position of the ATPase genes in these clusters to the same

genes in the Thaumarchaeota. We also provide a more robust sampling of the Nanohaloarchaea;

we include seven newly sequenced and assembled nanohaloarchaeal genomes together with

existing genomes mined from the NCBI database. Robust sampling of the taxa of interest, like

the one offered here, has the potential to improve the recovery of evolutionary relationships

without adding more sites (genes) (Graybeal, 1998).

In maximum likelihood and Bayesian phylogenies, we find that the Nanohaloarchaea

group robustly with the Haloarchaea in the single gene phylogenies, while the

Methanonatronarchaeia were placed as a deeper branching euryarchaeal lineage, most likely at

the base of the Methanotecta superclass. In large, concatenated datasets, we recover a

monophyletic DPANN (including the Nanohaloarchaea). We also provide evidence that the

ATPase genes have likely not been transferred in the case of the Nanohaloarchaea-Haloarchaea,

and contrast this specific relationship with the clearly transferred ATPases in the

Thaumarchaeota.

Results

Increased genomic representation of the Nanohaloarchaea

We obtained five new Nanohaloarchaea single amplified genomes (SAGs) from solar salterns in

Spain and two metagenome assembled genomes (MAGs) from Israel. The summary statistics

and accompanying information of these genomes can be found in Table S2. While the SAGs are

of poor assembly quality and completeness, enough genes were recovered from them for

phylogenomic analyses; furthermore, they unequivocally group with the other Nanohaloarchaea

in all analyses. These seven genomes expand the number of Nanohaloarchaea assemblies

available for analyses (18 at time of writing). Total average nucleotide identity (tANI) was used

to delineate taxonomy amongst the newly described Nanohaloarchaea. Figure S2 is a distance-

based tree calculated from corrected tANI distance (see Figure S1 for distance matrix) between

the previously described and newly described Nanohaloarchaea. Using conservative cutoffs, it

appears that SAGs SCGC AAA188-M06 and M04 may belong to the genus *Ca.* Nanosalina.

SAG M21 seems to be a member of *Ca.* Nanosalinarium, while the remaining new genomes

(SAGs and MAGs) do not belong to any previously described candidate genera.

The genome described as Nanohaloarchaea archaeon PL-Br10-U2g5 (Vavourakis *et al.*,

2016) was likely miss-identified as a Nanohaloarchaeon. We find that this strain unequivocally

groups within *Halorubrum* species in ribosomal (protein and rRNA), whole genome, and single

gene phylogenies (Figure S2).

While the fragmented nature of the SAGs is useful for phylogenetic analyses, there are

not enough genes to paint a comprehensive picture of their inferred metabolisms; the focus of the

metabolic analyses was therefore centered on the two MAGs described above as they are almost

complete. The two MAGs, M322 and AT22, have metabolic capabilities comparable to

previously described Nanohaloarchaea (Narasingarao *et al.*, 2012). Both MAGs are deficient for

enzymes in their nitrogen incorporation and lipid biosynthesis pathways. Both genomes encode

key enzymes involved in glycolysis and sugar metabolism; the presence of a number of sugar

dehydrogenases indicates a possible fermentative lifestyle. AT22 encodes a membrane bound

domain and the jellyroll fold LamG, which have been implicated in host cell interactions in

DPANN archaea (Golyshina *et al.*, 2017; Hamm *et al.*, 2019). The comprehensive list of genes in

their respective genomes (including the SAGs) described here are provided in Table S3.

Phylogenetic placement of halophilic lineages

210   To shed light on the evolutionary origins of the Methanonatronarchaeia and the

211   Nanohaloarchaea, we have produced three sets of trees from distinct markers that contain

212   differing phylogenetic signals. A marker set composed of the AtpAB proteins (Figure 2, S3) was

213   used to calculate phylogenies; these are slowly evolving single copy genes. The phylogenies

214   calculated from this marker set were compared to phylogenies calculated by large concatenates:

215 a concatenate of 44 ribosomal proteins (Figure S6), and a concatenate of 282 genes calculated to

216 be within the core genome of the Nanohaloarchaea (Figures 5, S7,9). All three tree sets contain

217 >150 taxa, representing Archaea that span the euryarchaeota, TACK group (including Asgard

218 archaea), and the candidate DPANN superphylum. The phylogenies are depicted as rooted with

219 the TACK Group, but should be considered as **unrooted**, as the root of the Archaeal tree remains

220 an open question with the emergence of Eukaryotes from the Archaea likely having rendered

221 them paraphyletic (Fournier & Poole, 2018; Gribaldo *et al.*, 2010; Spang et al., 2018; Williams *et*

222 *al.*, 2020; Williams *et al.,* 2013). In addition, a recent study places the root inside of the

223 euryarchaeota (Raymann *et al*., 2015). However, the placement of the Archaeal root does not

224 impact the conclusions drawn from our phylogenies, presuming the Archaeal root is placed

225 outside of the euryarchaeal crown group.

227 **ATP synthase catalytic and non-catalytic subunit phylogenies**

228 The ATP synthase catalytic and non-catalytic subunits are slow evolving, essential genes. Single

229 protein phylogenies of these subunits may ameliorate LBA and deletion-transfer-loss (DTL:

230 evolutionary conflicts driven by gene transfer, loss, or deletion that may mislead the

231 interpretation of a phylogeny) conflicts; both of which have plagued large scale, multi-locus

232 attempts at reconstructing the Archaeal phylogeny. A drawback of the ATP synthase is that it

233 has been suggested to have been horizontally transferred, and thus its phylogeny, although less

234 prone to artifacts of phylogenetic reconstruction, may not represent the whole genome (see

235 discussion below). Maximum likelihood phylogenies (see Table S4) of the AtpA and AtpB

236 proteins were created using site-homogeneous and site-heterogeneous substitution models. All

237 tree reconstructions based on the original, unaltered multiple sequence alignments confidently

238 placed the Nanohaloarchaea as a sister-group to the Haloarchaea ($\geq$ 91 Bootstrap Value (BV)). A

239 representative example tree constructed with the AtpA+B concatenate is shown below (Figure

240 2). Nanohaloarchaea and Haloarchaea are grouped together and are positioned inside the

241 Methanotecta**.** Typically, Haloarchaea are often seen as a sister-group to the Class II

242 methanogens (group including the Methanosarcinales, Methanomicrobiales, and

243 Methanocellales), in ATPases subunit phylogenies this relationship was interrupted by the

244 placement of other archaeal groups: in case of the AtpA+B concatenate and AtpA, the

245 Nanohaloarchaea-Haloarchaea sistergroup is separated from the Class II methanogens by Marine

Group II archaea and a Woesearchaeon (Figure 2, Table S4, Figure S3a). In case of the AtpB

protein the Nanohaloarchaea-Haloarchaea sistergroup is also sometimes (dependent on type of

substitution model, % of sites retained, and alterations to the alignment matrix, i.e., recoding)

recovered at the base of the Methanotecta (a group including the class II methanogens and the

Archaeoglobales) (Table S4, Figure S3b).

Curiously, the Methanonatronarchaeia are placed as a deeper branching euryarchaeal

lineage, suggesting either gene transfer or convergent evolution in regard to the extreme

halophilic "salt-in" strategy. In only one analysis did all three lineages group together, with poor

support inside the Methanotecta, (Table S4, LG+C50 AtpA d4). The newly described

Hikarchaeia's (non-halophilic sister-group of the Haloarchaea; Martijn *et al*., 2020) sisterhood

with the Haloarchaea is recovered in the AtpA and AtpA+B sequences (Figure 2, Figure S3a). In

the case of the AtpB tree, the Hikarchaeia emerged from within the Haloarchaea (Figure S3b),

but with marginal support (BV= 68). In all of these ATPase subunit phylogenies the

Nanohaloarchaeota branch before the split between Haloarchaea and Hikarchaeia.

The placement of the remaining DPANN taxa (i.e., without Nanohaloarchaea) appears

erratic. However, it is worth noting the groups considered as members of DPANN fail to form a

monophyletic clade in all of the ATPase based trees and the branches breaking the DPANN

group apart are supported by high BVs (Table S4).

266
267
Fig 2. **Maximum likelihood phylogeny calculated from AtpAB proteins.** The depicted tree
contains most features of the other calculated ATP synthase phylogenies. Several taxa were collapsed into higher
taxonomic ranks. Important taxa including the halophilic lineages and DPANN (teal) sequences have been colored;
Nanohaloarchaea (red), *Haloarchaea* (blue), *Methanonatronarchaeia* (purple), Methanomada (brown),
Methanotecta methanogens (orange), and the Hikarchaeia (magenta). The tree is drawn as rooted by the TACK
Group but should be considered as unrooted. This tree was calculated using the LG+C60 model.
274

Compositional bias

Compositional bias in encoded amino acids can generate artifacts in large, domain-wide
phylogenies (Aouad *et al.,* 2018; Aouad *et al.,* 2019). However, due to the slow rate of evolution
in these ATPase subunits, compositional bias has been minimized. A chi-squared test of
composition for both protein alignments revealed only 10% and 6% of taxa fail the composition
test in AtpA and AtpB sequences, respectively. None of the sequences that failed this
composition test belong to a member of the halophilic lineages barring one sequence that
belonged to a Nanohaloarchaeon with an incompletely sequenced *atpA*. To minimize
compositional bias, both alignments were recoded into 4 and 6 Dayhoff groups (Susko & Roger,
2007). These recoded alignments were used to create maximum likelihood and Bayesian
phylogenies, which mostly recapitulated the groupings discussed earlier (Table S4). The only
difference was that in several instances Methanonatronarchaeia moved either to the base of the
Methanotecta*,* grouped with the Haloarchaea and Nanohaloarchaea, or with the TACK group.

A reason for bias in extreme halophilic lineages is an acidic proteome, *i.e.,* increased
presence of aspartic and glutamic acid (D/E) in their protein sequence. This may lead to
"compositional attraction", where those taxa that have an abundance of D/E sites are more likely
to cluster together in a phylogeny. Sites that contained a conserved D/E residue among the
Haloarchaea, Nanohaloarchaea, and the Methanonatronarchaeia were deleted from the AtpA and
AtpB alignments. Maximum likelihood phylogenies were created from these new alignments
(Table S4), and the topology discussed above was recovered, albeit with lower support due to the
loss of phylogenetically informative sites. We recognize that this method limits compositional
attraction, but cannot completely rule out the possibility other residues have evolved
independently in similar hypersaline environments.

Suitability of the ATP synthase subunits as a phylogenetic marker

Although the ATP synthase subunits are an attractive phylogenetic marker, these genes have
been shown to be horizontally transferred (Wang *et al*., 2019), sometimes between distantly
related organisms (Lapierre, *et al*., 2006; Olendzenski *et al*., 2000), and are presumably adaptive.
Wang *et al.* has recently shown, convincingly, that Thaumarchaeota have adapted to more acidic
environments, with a gene transfer of V-Type ATPases from other acidophilic archaea. A similar

transfer has also been suggested for the Nanohaloarchaea, which would complicate the

interpretation of the ATPase phylogenies. To shed light on whether this is the case, we regressed

and correlated the pairwise distance matrices of the gene families in the Nanohaloarchaea and the

Thaumarchaeota (using a modification of the approach described in (Rangel *et al*., 2019; Rangel

et al., 2020) see methods for details). Two analyses were performed with this gene family

distance method, with different marker sets. The GTDB Archaeal 122 marker set (Parks *et al*.,

2018) with the addition of ATPase genes was used to compare the correlation of gene families

within the Thaumarchaeota and the Nanohaloarchaea. These markers have been found to

produce consistent phylogenies, and should be commonly found in most Archaeal genomes, thus

they are appropriate for making comparisons between two groups. Relationships (via correlation

distance $1-r^2$) between gene families in the Nanohaloarchaea and Thaumarcheota were

calculated, and further ordinated using nMDS (non-metric multi-dimensional scaling) and

statistically evaluated using a categorical Mantel test at the 95% confidence level (Figure 3).

321
Fig 3. **nMDS plots of the gene families in the Thaumarchaeota and Nanohaloarchaea.** Shows
the ordination of various gene families (from the Archaea 122 marker set) in the Thaumarchaeota and
Nanohaloarchaea. A categorical Mantel test with two defined categories, ATPase genes (colored in blue) and non-
ATPase genes (colored in red), was used to determine significance with the 95% confidence ellipse. A) The gene
families in the Thaumarchaeota, the ATPase genes clearly fall outside of the 95% confidence ellipse, with a p =
0.001. B) The gene families in the Nanohaloarchaea, where the ATPase genes clearly fall inside the ellipse, with a p
= 0.182.

The ordination plots (Figure 3) clearly show the contrast of the relative evolutionary

trajectories of the ATPase genes in the Thaumarchaeota versus Nanohaloarchaea. In the case of

the Thaumarchaeota, whose ATPase genes are known to be transferred (Wang *et al.,* 2019), the

ATPase genes clearly fall outside of the 95% confidence ellipse. The 95% confidence ellipse, in

this analysis, comprises most of the gene families belonging to a common evolutionary trajectory

and are likely not the result of a recent horizontal gene transfer between divergent species. The

stark contrast between the ATPase genes in these two groups, highlights the difference in

circumstance around the evolutionary trajectory of the ATPase genes relative to other gene

families in both groups. For reference, this analysis was also performed on subsets of the

Thaumarchaeota genomes (Wang *et al*., 2019); one subset containing the acidophiles (received V

Type ATPases via HGT) and another subset containing only the neutrophilic Thaumarchaeota

(vertically inherited A type ATPases). The acidophilic Thaumarchaeota's ATPases also stands

out as atypical in this analysis (transfer detected, p-value = 0.024; Figure S14a), while the

ATPases in the neutrophiles falls comfortably inside the 95% confidence ellipse (p-value =

0.576).

In addition to the ordinations, the correlations between the gene family distance matrices

and their associated clustering diagrams for both the Archaeal 122 and the 282 core gene marker

sets revealed, that in the Nanohaloarchaea (Figures. 4, S4), the ATPase genes fall into a large

cluster of genes (Figure 4, genes highlighted by purple rectangle from 282 core genes of the

Nanohaloarchaea). Every gene family enclosed by these purple rectangles, including the AtpAB

genes, share a broadly similar evolutionary trajectory. This cluster is also clearly distant to other

large clusters (i.e., genes enclosed by the blue rectangle in Figure 4) and the genes located on

deep, long branches that fail to form a significantly large cluster. The genes on deep, long

branches in these clustering diagrams represent those that likely have been horizontally

transferred or follow an unconventional (i.e., not strictly vertical) evolutionary history, reflected

in the pairwise distance matrices of the gene family. The genes within the large clusters share a

similar evolutionary history, which might be explained with predominantly vertical inheritance.

In the gene families of the Thaumarchaeota (Figure S5), the ATPase genes (which were

identified as having been transferred, Wang *et al.,* 2019) form a separate cluster distant from

three other clusters. The Thaumarchaeota ATPase subunits likely share an evolutionary

trajectory with each other that differs from the trajectories in the other two clusters and other

individual genes on long branches. A description of this clustering implementation can be found

in the methods section, and in an interactive Jupyter notebook script with instructions

(https://github.com/Gogarten-Lab-Team/NanoH_GBE_2020).

The 282 core gene families of the Nanohaloarchaea were also ranked on their speed of

evolution; based on their average slopes of pairwise distance matrices (1 gene family vs. the 281

other gene families) and their gamma shape parameters (Table S5). The AtpA+B proteins both

fall in the top 10 percentile of the slowest evolving genes, furthering their case for inclusion in

phylogenetic analyses, considering they have likely not been transferred recently between the

Nanohaloarchaea and the Haloarchaea. Furthermore, the clustering of slowly evolving genes

(like the ATPases) and fast evolving genes (other genes, see Figure 4 and Table S5 for speed

ranks) into the same evolutionary trajectory indicates the gene family clustering method

described here may not be very sensitive to LBA artifacts. This is likely due to the analyses

focusing on the correlations of individual pairwise distance matrices of each gene family.

Analysis of Ribosomal Data

To explore the possible synergy of using multiple loci in reconstructing the history of these

halophilic lineages, we constructed supermatrices of core genes and ribosomal proteins from a

taxonomic sampling similar to the ATP synthase trees. We first created a ribosomal supermatrix

containing 44 concatenated ribosomal proteins (RBS supermatrix). The placement of the

Haloarchaea and the Nanohaloarchaea calculated from the ribosomal supermatrix (Figure S6b)

resemble many previously calculated large-scale phylogenies based on RBS proteins (Andrade *et*

al., 2015; Sorokin *et al.,* 2017). In contrast to the single protein trees, the RBS tree confidently

places the Nanohaloarchaea in a monophyletic clade with the rest of the DPANN members (in

both the original and recoded supermatrices). The phylogeny of the 16S+23S rRNA genes

recovers the Nanohaloarchaea as a sister-group to the Haloarchaea (Figure S10); however this

sister-group is placed outside of the euryarchaeota. The reliability of the rRNA phylogeny is

questionable, as several other groups (i.e., several DPANN members, Methanococcales, etc.)

have moved from their accepted positions. These placements may reflect rRNA HGT between

various Archaeal lineages (Boucher, *et al*., 2004) (Table S4), and compositional biases (in the

rRNA genes) driven by adaptation to various environmental pressures.

Analysis of Nanohaloarchaeal core genome

We also created a core genome matrix composed of 282 loci (called the Large Core supermatrix)

of all genes represented in every single Nanohaloarchaeal genome considered complete. In the

previously described analysis, we clustered the gene families based on the correlation between

the distance matrices for these 282 gene families. From these clusters we created 3 other

supermatrices which are subsets of the Large Core supermatrix, these are called the Left, Right

and Center supermatrices (corresponding to the positions of the clusters on the correlated gene

family dendrogram; Figure 4). Although all 282 gene families in the Large Core are represented

in each Nanohaloarchaeal genome, these gene families must be analyzed for DTL conflicts and

suitability in a concatenated alignment. For an example, the AtpA+B genes recover the

Nanohaloarchaea and Haloarchaea as a sister-group, and falls within the Left supermatrix.

However, not all large clusters of gene families (Figure 4) should fall under the assumption of

vertical inheritance (one possible evolutionary trajectory). These dendrograms show differential

transfer patterns of gene family clusters, and it is possible a large cluster (*i.e.,* Left or Right)

could be an entire ensemble of transferred genes (another possible evolutionary trajectory).

Splitting up the Large Core families into subdivisions (like the Left, Right, and Center

supermatrices) may be a method to dissects common evolutionary trajectories contained in the

Nanohaloarchaeal core genome.

All phylogenies calculated using the Large Core supermatrix (282 genes), and its

derivatives (Left, Right and Center) recover the Nanohaloarchaea with other members of

DPANN (Figure 5). In all of these supermatrices, except for the entire Large Core supermatrix,

the Methanonatronarchaeia is resolved at the base of the Methanotecta group, as reported by

(Aouad *et al.*, 2019; Martijn *et al.*, 2020). In addition, the Center and Left supermatrices recover

the Haloarchaea as a sister-group to the Methanomicrobiales*,* similar to (Aouad *et al.*, 2018).

Curiously, using the Right supermatrix (Figure 5d) recovers the Haloarchaea as a sister-group to

the Nanohaloarchaea. However, the Haloarchaea are relocated to outside of the euryarchaeota,

and the Nanohaloarchaea are not placed outside the DPANN. This indicates that the Right cluster

of genes (94 total) may contain genes frequently transferred from the Nanohaloarchaea to the

Haloarchaea, representing traces of the Haloarchaea-Nanohaloarchaea symbiotic interaction. One

of the pitfalls of using large concatenates is illustrated in Figure 5bc, when the genes from the

Left cluster and the Right cluster are combined into a single concatenate. These clusters have

clearly different phylogenetic signals (Figure 5c for the Left, Figure 5d for the Right), and

forcing them on the same tree (Figure 5b) clearly leads to the one of the signals being

overwhelmed by the other, with little effect on overall bootstrap support (the Right cluster is

overwhelmed in this case).

The conflict of the Nanohaloarchaeal core gene families is revealed once again when the

Large Core and the Left supermatrices were recoded into 4 Dayhoff groups to reduce

compositional biases (Figure S7). The recoded Large Core phylogeny recovers the monophyly of

all three halophilic lineages inside of the Methanotecta, with high support. However, the recoded

Left core subdivision recovers the Nanohaloarchaea inside of a monophyletic DPANN. In

recoded Left supermatrix, the Methanonatronarchaeia were recovered at the base of the

Methanotecta. We also note that while the gene family clustering shows a better correlation for

families belonging to the same cluster, the reconstructed phylogenies for individual gene families

within each cluster are not identical (most individual gene families have poor phylogenetic

resolution) and not identical to the phylogeny calculated from the concatenation, e.g., the AtpA

and AtpB proteins are part of cluster the left cluster (Figure 5c), whose phylogeny reconstructed

from the concatenation is different from the ATPase phylogeny.

442
443
444

445
Figure 4. **Clustering diagram of 282 gene families that form the core to the Nanohaloarchaea**, clustered by the
pairwise correlation between distance matrices calculated for individual gene families. Families clustered together
share similar (although not identical) evolutionary trajectories as assessed by their distance matrices calculated using
maximum likelihood models (see materials and methods). Gene families enclosed by the rectangles share broadly
similar evolutionary trajectories (with the same members of their cluster), and likely not have been transferred
between divergent lineages, while gene families on deep, long branches likely have an unconventional evolutionary
trajectory. Subdivisions of the Large Core supermatrix were defined using the clusters (rectangles) in the
dendrogram, called the Left (gene families enclosed by the purple rectangle), Right (blue rectangle), and Center (a
combination of Left and Right clusters). The blue tip labels indicates where the AtpAB genes fall in the clusters.

Fig 5. **Maximum likelihood phylogenies of Archaeal Large Core genome supermatrices**. All

phylogenies were calculated with the LG+C60 mixture model A) Maximum likelihood phylogeny calculated using

all 282 gene families. B) Phylogeny calculated using the Center supermatrix. C) Phylogeny calculated using the Left

supermatrix (95 gene families). D) Phylogeny calculated using the Right supermatrix (94 gene families). Colored

node circles indicate bootstrap support value magnitude.

In an attempt to assess the impact of long branch attraction, the DPANN sequences

(excluding the Nanohaloarchaea) were removed from the entire Large Core genome supermatrix.

The phylogenetic tree calculated from this new supermatrix (Figure S8) places the

Nanohaloarchaea, and Haloarchaea together as sister-groups, but this sister-group has been

moved out of their accepted position in the Methanotecta, and possibly out of the euryarchaeota

altogether (see discussion).

Topology Tests of Halophile Monophyly

To further investigate the possible monophyly of the three lineages, we used trees that

recovered the Nanohaloarchaea, Haloarchaea, and Methanonatronarchaeia inside of the

euryarchaeota as a monophyletic group (topology from Figure S7a). The explanatory power of

these constrained monophyletic trees were tested against the Large and Left supermatrices using

the approximately unbiased test (AU-test, Shimodaira, 2002). The topology test revealed that

these constrained trees differed significantly in their likelihood landscape as compared to signals

in both supermatrices, with p-values near 0. The AU-test reveals that the explanatory power of

the trees with a monophyletic grouping of halophilic archaea is not compatible with the

supermatrices and thus the hypothesis of halophile monophyly should be rejected under these

datasets.

We also employed gene concordance factor analysis (gCF (Ané *et al*., 2007; Gadagkar *et*

al., 2005)) to our Large Core supermatrix to dissect which topology each individual gene family

supported. Reference trees were used to reflect conflicting evolutionary hypotheses (a tree

grouping the Nanohaloarchaea and Haloarchaea together (Figure S7a) vs. a tree that placed the

Nanohaloarchaea in the DPANN (Figure 5c)). Using a reference tree that was constrained to

group the Nanohaloarchaea with the Haloarchaea, 16 genes were found to recover the internode

that supports these groups' monophyly. These 16 genes include highly conserved proteins such

as the ATP synthase operon, ribosomal proteins, and an elongation initiation factor. In contrast,

we also used an alternative reference tree that was constrained to group the Nanohaloarchaea

within the DPANN superphylum and found 15 different genes in support of this hypothesis.

These DPANN supporting genes include RNA polymerase subunits, FtsY, and ribosomal

proteins. It is worth noting that the genes supporting the Nanohaloarchaea+Haloarchaea group

consist of genes that evolve significantly slower (average rank ~74) than those of the DPANN

support set (average rank ~93), based on the rate of evolution rankings discussed previously. We

also applied the gCF method to locate the best supported placement of the

Methanonatronarchaeia, and found the highest concordance was at the base of Methanotecta

super class. The full table of concordant genes, as well as reference trees can be found in Figures

S11,12 and Table S6.

Discussion

Sisterhood of Nanohaloarchaea and Haloarchaea

Analysis of the catalytic and non-catalytic subunits of the archaeal ATP synthase group the enzyme from Nanohaloarchaea as a sister-group to the Halobacterial (Haloarchaeal) subunits (Figure 2; Wang *et al.,* 2019). This strongly supported grouping is also recovered when the data are recoded to reduce compositional bias, when alignment columns containing acidic residues in both the Nanohaloarchaea and the Haloarchaea are deleted, and when the CAT-GTR model (a model that is less sensitive to compositional effects and long branch attraction artifacts) is used in phylogenetic reconstruction. None of these analyses recovered the DPANN clan, however, this may not be strong evidence against the existence of DPANN, as HGT in members of DPANN is largely unexplored in this work.

Placement of the individual DPANN groups in the ATP synthase phylogenies and the absence of an ATPase in some ectoparasitic Nanoarachaeota (Wurch *et al.*, 2016) may be interpreted as evidence questioning whether the ancestor of the DPANN even possessed a functional ATP synthase. The *Nanoarchaeum equitans* genome is an example of a DPANN member which encodes the typical archaeal ATPase headgroup (3 A and 3 B subunits), although it may lack ATP synthesis activity (Mohanty *et al.*, 2015). However, homology between the F (from Bacteria) and A type (from Archaea) ATPases demonstrates that these ATPases are older than the origin of Archaea (Gogarten *et al.*, 1989; Gogarten & Taiz, 1992). While ATPases are known to have been horizontally transferred, the described findings suggest that the Nanoarchaeal ancestor possessed an ATPase. It is certainly possible that modern DPANN genomes replaced their ATP synthases with homologs from their hosts or from other organisms occupying the same environment.

Given the consistent support for the Nanohaloarchaea-Haloarchaea clade in the AtpA and AtpB phylogenies, it is unlikely that this finding is due to compositional bias or long branch attraction. Two conflicting hypotheses can reconcile our findings with those of previous analyses based on concatenation of several genes or on gene tree/species tree reconciliations: (1) the ATP synthase was acquired by the ancestor of the Nanohaloarchaea from a relative of the Haloarchaea or (2) the previous multi-locus analyses do not reflect evolutionary history, but are artifacts due

to high substitution rates, gene transfer, and small genomes; and the Nanohaloarchaea and

Haloarchaea share a common ancestor.

The recent study by Wang *et al*. (2019) includes a phylogeny derived from the entire

ATPase operon in Archaea, that also recovered the sisterhood between the Nanohaloarchaea and

Haloarchaea. Wang *et al*. consider horizontal transfer of the operon as explanation for this

grouping, and also observe an identical operon structure in both groups, which supports the

monophyly of nanohaloarchaeal and haloarchaeal ATPases. Those authors recognized clear

conflicts between a DPANN supergroup and the ATPases phylogeny, and reconciled this conflict

by invoking ATPase horizontal gene transfer. In correlations of Nanohaloarchaea gene families,

it was revealed the ATPase genes are more closely clustered to a large set of genes; in contrast

Thaumarchaeota ATPase genes form their own evolutionary cluster distant from other genes in

the genome. These results can be reconciled by considering that ATPase genes were indeed

transferred into Thaumarchaeota, but not in the case of Nanohaloarchaea. The gene family

correlation method would be highly sensitive to single and multiple recent gene transfers, as the

distance matrices analyzed for each gene family are vectorized (i.e., taxon specific information is

saved and kept consistent throughout the entire clustering process, so if a single gene is

transferred into only a single taxon, it will be recorded and thus affect the clustering of the entire

gene family). While the clustering based on gene distance correlations does well in recovering

ATPase transfer in Thaumarchaeota, the absence of detected transfer events in Nanohaloarchaea

cannot be considered proof that no transfer has happened. A suspected transfer would have

occurred greater than ~1 BY ago before the deepest splits within the Nanohaloarchaea and

Haloarchaea, respectively. Within-niche transfer of ATPase operons is certainly possible and is

supported in the case of the Thaumarchaeota (Wang *et al.,* 2019, Figure S5) and Deinococcaceae

(Lapierre *et al*., 2006); however, we are not aware of any evidence that extends this logic to the

Nanohaloarchaea.

We provide evidence that ATPases do not stand out as atypical in their evolutionary

history as compared to other genes found in Nanohaloarchaea (Figures 3-4, S4-5). Our

interpretation of these data is that the sisterhood relationship of the Nanohaloarchaea and

Haloarchaea ATPases should not be immediately discarded as resulting from HGT. The analyses

presented here and by Rayman *et al.,* 2014 and Aouad *et al.,* 2018-9 suggest that

Nanohaloarchaeal genomes have been shaped by a complex evolutionary history. Many gene

families support inclusion of the Nanohaloarchaea into DPANN, while the aforementioned

studies suggest of a placement within Haloarchaea or other euryarchaeota. The totality of support

for a Nanohaloarchaea-Haloarchaea sister-group contained within our analyses include: ATPase

phylogenies (and that these gene are unlikely to have been transferred (Figures 2-4, S4-5));

recoded nanohaloarchaeal Large core genome; and gCF analyses which identified several slowly

evolving genes also supporting a Nanohaloarchaea-Haloarchaea sister-group relation.

Due to the observation of radically different phylogenetic signals present in the

nanohaloarchaeal core, we consider an analogy between Nanohaloarchaea and Thermotoga. The

Thermotoga core genome is extremely chimeric: its evolutionary history indicates genes

comprising the "informational" functionality (i.e., genes involved in replication, repair, etc) are

bacterial in origin, while genes that contribute to metabolism are of Archaeal or Clostridial origin

(Logsdon & Faguy, 1999; Zhaxybayeva *et al*., 2009). In comparison, genes that compose the

Left cluster of genes in Nanohaloarchaea are enriched for proteins that encode for translational

functions, while the Right cluster is enriched for proteins that serve transcription purposes (Table

S5, K= transcription, J=translation), indicating that Nanohaloarchaea are highly chimeric too.

Phylogenies calculated from concatenated datasets support the existence and monophyly

of the DPANN superphylum (including the Nanohaloarchaea). When genomes from DPANN

members were included, the Nanohaloarchaea were recovered as part of the DPANN group. In

the absence of the other DPANN genomes, Nanohaloarchaea formed a clade with Haloarchaea

(Figure S8), even after removing potential biases. However, the sister-group moved out of

Methanotecta, and possibly the euryarchaeota too. As to whether this sister-group is located in

the euryarchaeota depends on where one places the archaeal root. If one expects the root to be

inside the euryarchaeota (Raymann *et al*., 2015), this sister-group has a possibility of falling

outside the euryarchaeota, as it falls outside Methanotecta and may lead to a branch where the

DPANN superphylum could attach. The observation that a monophyletic Nanohaolarchaeal-

Haloarchaeal grouping is recovered from the Large core concatenation but at the base or even

outside the euryarchaeota illustrates observed evolutionary relationships between archaeal

classes obtained from gene family concatenations have to be interpreted with caution.

Phylogenetic reconstruction that constrained Nanohaloarchaea to group with Haloarchaea

resulted in a maximum likelihood phylogeny that the AU-test (Shimodaira, 2002) evaluated as

incompatible with the best tree for this Large Core genome dataset, revealing a strong

phylogenetic signal, either due to shared ancestry or systematic artifact, that does contradict the

sister-group relationship between Nanohaloarchaea and Haloarchaea.

Radically different placements of Nanohaloarchaea (Figure 1, red indicators) can be at

least partially attributed to the taxonomic sampling of the DPANN superphylum. In instances

where the Nanohaloarchaea were recovered inside the euryarchaeota (Narasingarao *et al.*, 2012,

Zhaxybayeva *et al.*, 2013; Aouad *et al.,* 2018; Aouad *et al.,* 2019), DPANN sequences were not

included in the tree. However, including a robust sampling of DPANN sequences in the

alignment (Andrade *et al.,* 2015; Sorokin *et al.,* 2017; Wang *et al.,* 2019, Figure 5) generally

attracts the Nanohaloarchaea into that superphylum.

The gCF analysis revealed 16 core genes in support of Nanohaloarchaea-Haloarchaea

sister-group; however, 15 genes support Nanohaloarchaea inclusion in DPANN. In support of

Nanohalo+Haloarchaea group are 16 genes that evolve significantly slower than those in support

of the opposing hypothesis (Table S6). Previous analyses have indicated high bootstrap support

for including the Nanohaloarchaea within DPANN (Sorokin *et al.*, 2017; Wang *et al.*, 2019).

This support may reflect the strong but artifactual signal in fast evolving genes, phylogenetic

signals created through gene transfers, and forcing all genes with different histories onto the

same tree -- conflicting signals are likely abundant of in all concatenated marker sets. Our gCF

analysis dissected the concatenation based on individual gene trees, revealing opposing

phylogenetic signals present in the original concatenated dataset. It is important to supplement

the sampling variance measure for the singular branch (*i.e.,* bootstrap), with a measure of

variance in the overall dataset with metrics like the concordance factors. The concordance factors

can reveal variance (conflict) within the multi-locus alignment datasets.

In an attempt to dissect the Large Core genome concatenation even further, we

subdivided it into 3 subdivisions; the Left, Right and Center supermatrices (Figure 4). These

subdivisions are based on pairwise distance matrices of each individual gene family. Phylogenies

of the Right supermatrix reveals that a large ensemble of genes (94) that are a part of the

Nanohaloarchaea core genome may have been transferred from the Nanohaloarchaea to

Haloarchaea, as Haloarchaea moved from euryarchaeota into DPANN. Concatenations involving

these genes may calculate a phylogeny with high artifact potential due to their possible transfer

or unconventional evolutionary trajectory. It is worth noting again, that the AtpA+B genes fall

into the Left supermatrix, even though their individual signal robustly groups the

Nanohaloarchaea and Haloarchaea as sister-groups. This demonstrates that genes evolving

through a similar evolutionary trajectory (Left cluster), can recover evolutionary placements and

may be convoluted by concatenating gene families with disparate rates of evolution.

628   Monophyly of extreme halophilic archaea

The Methanonatronarchaeia did not reveal a well-supported association with any particular

Archaeal group in any of these phylogenies. In the ATP synthase-based phylogenies, homologs

from three members of this group were recovered as a deeper branching euryarchaeal lineage

without well supported affinity to any other euryarchaeal group. Sequences from the

Methanonatronarchaeia were, however, separated by at least one well supported bipartition from

other halophilic archaea grouping with non-halophilic methanogens (Figure 2, S3).

A concatenation of Nanohaloarchaeal core genes reliably placed Methanonatronarchaeia

(Figure 5bc) basal to the Methanotecta super-class, as proposed by Aouad *et al.,* 2019. When

using the entire Large Core genome supermatrix (Figure 5a), Methanonatronarchaeia appeared as

a sister-group to Haloarchaea (BV= 88). Aouad and colleagues provided evidence for three

independent adaptations to high salt environments (through the salt-in strategy) in Haloarchaea,

Nanohaloarchaea, and Methanonatronarchaeia (Aouad *et al.,* 2018; Aouad *et al.,* 2019). While

we consider convergent evolution events rare, independent adaptations to hypersalinity resulting

from salt-in strategy pressures and revealed through shifts in protein isoelectric points (Oren,

2008) have been observed in *Salinibacter* (Bacteroidetes) and *Salinicoccus* (Firmicutes) (see

Figure S13), with minimal reliance on HGT from haloarchaea (Mongodin *et al*., 2005).

Methanonatronarchaeia have been deduced to employ a salt-in strategy, using

intracellular potassium ion concentrations (Sorokin *et al.,* 2017), the same adaptation present in

Nanohaloarchaea and Haloarchaea. However, a proteomic analysis of theoretical isoelectric point

(pI) distributions reveals a less biased distribution of pIs in these methanogens compared to other

proteomes of organisms that use a salt-in strategy (Haloarchaea, Nanohaloarchaea, *etc*.) (Figure

S13). This distribution of theoretical pIs in Methanonatronarchaeia resembles that found in

marine archaea (Figure S13), and *Halanaerobiales*. The *Halanaerobiales* follow an

experimentally confirmed salt-in strategy without an acidic proteome. Instead, they hydrolyze

Glutamine (Q) and Asparagine (N) to compensate for the lack of acidic amino acids (Bardavid &

Oren, 2012). However, the genome of *Acethalobium arabaticum*, a member of the

Halanaerobiales, encodes a more acidic proteome, similar to *Salinibacter* and *Salinicoccus*

(Figure S13). Methanonatronarchaeia may be a similar example of independent adaptation to

hypersalinity. In some Methanonatronarchaeia the concentration of intracellular potassium did

not yet have a significant impact on the distribution of pIs of encoded proteins or possibly, they

may also hydrolyze their N/Q residues to make their acidic conjugates, like *Halanaerobiales*.

The AtpAB dataset robustly recovers the Nanohaloarchaea-Haloarchaea sister-group.

Furthermore, we provide evidence that these genes are slow evolving (Table S5), and unlikely to

have been transferred recently between the groups (Fig 3-5, S4-5). An obvious caveat is that the

better resolved single gene phylogeny represents only a single gene or operon, and that its

phylogeny is embedded in the net-like, reticulated genome phylogeny. Data from concatenated

datasets robustly recovers the Nanohaloarchaea group within DPANN (the exception being

recoded Large core phylogeny, Figure S7a). However, these datasets are rife with conflict

(transferred genes, genes with differing rates of evolution; gCF and Figure 4) and forcing them

on a single tree likely is inappropriate. We consider phylogenetic placement of the

Nanohaloarchaea an open question. A plethora of analyses using large concatenates support

inclusion of Nanohaloarchaea in DPANN (Andrade *et al*., 2015; Castelle *et al*., 2015; Rinke *et*

al., 2013; Dombrowski et al., 2020), but the same can be said for the opposite (Brochier-

Armanet, Forterre, & Gribaldo, 2011; Petitjean *et al*., 2014; Raymann *et al*., 2014; Williams *et*

al., 2015, Aouad *et al*., 2018, Aouad *et al*., 2019). Conflict between these analyses (Figure 1)

may, at least in part, be due to reliance on large concatenates and forcing disparate evolutionary

signals onto the same tree. Dissecting these evolutionary signals and evaluating their suitability

for such analyses could be a way forward for resolving the debate of the nanohaloarchaeal

placement and the existence of DPANN.

Recently, Hikarchaeia (Martijn *et al*., 2020) were found to be more closely related to

Haloarchaea (than Nanohaloarchaea) in the ATPase phylogenies. A hallmark of a salt-in

strategist can be found in the Hikarchaeia's proteomes, as they decidedly favor acidic residues

found in other salt-in strategists (Figure S13mn; Oren *et al*., 2008, Paul *et al*., 2008). Results of

ATPase phylogenetic analysis (Figure 2; Figure S3) also raises the possibility that both the

Hikarchaeia and Haloarchaea evolved from an extreme halophilic ancestor, and that Hikarchaeia

lost this adaptation after their divergence. If the ATPase phylogeny topology results from HGT

or gene sharing, acquisition of the Haloarchaeal ATPase by Nanohaloarchaea predates the split

between Hikarchaeia and Haloarchaea, again suggesting that extreme halophily might have been

an ancestral character of the Hikarchaeia. The alternative assumption that the Hikarchaeia and

Haloarchaea ancestor was not an extreme halophile would imply that transfer of the ATPase

operon occurred before Haloarchaea and Nanohaloarchaea convergently adapted to

hypersalinity. The inclusion of the Hikarchaeia in future phylogenetic analyses (once there is a

larger sampling of this lineage) may further elucidate the genomic events that lead to hypersaline

adaptation.

While our analyses do not prove that Nanohaloarchaea are not part of a DPANN

grouping, our findings indicate that when they are strongly supported in concatenated datasets

this might be the result of an artifact, and that the phylogenies of conserved slowly evolving

genes (ATPases, ribosomal proteins, and an elongation initiation factor) may better reflect the

origin of the Nanohaloarchaea. In most gene families the phylogenetic signal regarding

relationships between different archaeal classes is weak, and single gene phylogenies are poorly

resolved. The popular solution of data set concatenation (Lapierre *et al*., 2014) to amplify a

weak phylogenetic signal comes with the possibility that systematic artifacts and not a combined

phylogenetic signal dominate the resulting phylogenies (Bapteste *et al*., 2008). Resolving deep

divergences remains a hard problem. Due to horizontal gene transfer and phylogenetic

reconstruction artifacts the placement of divergent archaeal classes into larger groups remains

uncertain.

Methods

Sample collection, DNA extraction, and sequencing of new genomes

Two hypersaline environments in Israel were sampled for metagenomic sequences: the Dead Sea

and hypersaline pools at the Mediterranean coast in Atlit. Briefly, water samples from the Dead

Sea (31°30'07.2"N 35°28'37.2"E) were extracted using Niskin bottles in late July 2018. To create

the enriched media, the Dead Sea water (DSW) was diluted with autoclaved double distilled

water (DDW) (final ratio ⅕ [DDW/DSW]), amended with 0.1% glycerol, 1 uM KH_2PO_4, 1 g/L

peptone (Bacto, New South Wales, Australia), 1 g/L casamino acids (Difco, Detroit, MI USA).

The media was incubated at 30 °C for 42 days.

The Atlit environmental samples were collected from high salt tide-pools on the coast of

Israel (32°42′37.3″N 34°56′32.0″E) in mid-October 2018. Harvesting of the microbial

communities was performed by serial passage through filters (0.45um, 0.22um, 0.1um) (Merck

KGaA, Darmstadt, Germany). Environmental samples (Atlit) were first prefiltered using filter

paper No. 1 (11um pore size) (Munktell & Filtrak, Bärenstein, Germany). The filters were then

kept in -80 °C until DNA extraction. DNA was extracted from the filters using DNeasy

PowerLyzer PowerSoil kit (QIAGEN, Hilden, Germany) following the manufacturer's protocol.

For Dead Sea and Atlit samples, DNA purified from the 0.22um filters was used for library

preparation (NuGen Celero enzymatic with UDI indexing). The libraries were ran on Illumina

NovaSeq with SP flow cell, generating paired end reads (2x150bp).

Single amplified genomes (SAGs) were generated using fluorescence-activated cell

sorting and multiple displacement amplification, as previously described (Zhaxybayeva *et al.,*

2013), from hypersaline salterns located in Santa Pola (Spain). Low coverage shotgun

sequencing of SAGs was performed using Nextera library preparation and NextSeq 500

sequencers (Stepanauskas *et al*. 2017), resulting in an average of 377k, 2x150 bp reads per SAG.

Although this number of reads is sub-optimal for high-quality genome reconstruction

(Stepanauskas *et al*. 2017), they were sufficient to perform the specific analyses of this study.

SAG generation and raw sequence generation were performed at the Bigelow Laboratory for

Ocean Sciences Single Cell Genomics Center (scgc.bigelow.org).

Sequence quality control

Raw reads obtained from single cell sequencing were trimmed and quality assured using Sickle

v1.33 (Joshi & Fass, 2011) and FastQC v0.115 (Andrews, 2010). SPAdes v3.10.1 (Bankevich *et*

al., 2012) was used to complete initial assemblies of single cell genomes, using option -sc.

Contigs from the initial assembly were polished and bridged using the post-assembly Unicycler

v0.4.7 pipeline (Wick *et al*., 2017), using normal and bold settings. Conflicts between normal

and bold assemblies were investigated and reconciled in Bandage v0.8.1 (Wick *et al*., 2015). For

the metagenome assembled genomes (MAGs), raw reads were trimmed using Trimmomatic-0.36

(Bolger *et al*., 2014) and quality assured using FastQC v0.10.1. SPAdes v3.11.0 was used to

assemble the MAGs, using option -meta. Assembly Graphs were manually investigated using

Bandage v0.8.1. Binning was conducted with MetaBat2, the bins that contained the

nanohaloarchaeal MAGs were comprised of a single contig. The taxonomy and completeness of

the MAGs and SAGs were checked with CheckM v1.0.7 (Parks *et al*., 2015), on default settings

using a custom lineage marker developed specifically for Nanohaloarchaea (available on

request). The assembled genomes were annotated with Archaeal mode Prokka v1.13.3 (Seemann,

2014). Sequences annotated as the ATP synthase alpha and beta subunits were retrieved from

these genomes manually. The two MAGs in addition to ten high quality assemblies on NCBI

were compiled in a library to identify the 282 core genes of the Nanohaloarchaea.

Get_Homologues v03012018 (Contreras-Moreira & Vinuesa, 2013) with the COGtraingles v2.1

(Kristensen *et al*., 2010) and orthoMCL v1.4 (Li *et al*., 2003) algorithms (-t 0/1 option, -e option

to exclude paralogs) were used to identify these "bona-fide" core genes used to comprise the

core genome marker set.

Whole genome distance analysis

Average nucleotide identity (ANI) was calculated using a slight modification of the JSpecies

method (Richter & Rosselló-Móra, 2009). Genomes were divided into 1,020 nt fragments and

used as the query for pairwise BLAST searches. A 70% identity and 70% coverage cutoff was

implemented in a manner akin to the global ANI (gANI) filtering method (Varghese *et al*., 2015).

The filtered BLASTN (Camacho *et al*., 2009) searches were also used to calculate a modified

gANI and alignment fraction (AF), which were used to construct a phylogenetic tree as per the

tANI method (Gosselin *et al.*, 2020). The entire method and standalone script can be found at:

https://github.com/SeanGosselin/tANI_Matrix.git.

Assembly of datasets

169 high quality genomes spanning the Archaea domain were collected through NCBI's ftp site,

and were supplemented with the seven newly assembled Nanohaloarchaea genomes (Table S1).

AtpA and AtpB protein sequences were found in these genomes and gathered with BLASTP

v2.7.1, using default parameters. Similarly, protein sequences of 282 Nanohaloarchaea core

proteins and forty-four ribosomal proteins were found and gathered from these genomes using

TBLASTN using default parameters. Sequences hits from each protein were categorized into

their own respective files and aligned with Mafft-linsi (Katoh & Standley, 2013). Alignments

were trimmed by BMGE (Criscuolo & Gribaldo, 2010). Each alignment file of the core and

ribosomal protein dataset was concatenated using FASconCAT-G (Kück & Longo, 2014) to

generate supermatrices and the associated nexus partition files. A description of the core

supermatrices is available in the supplemental material (Table S5).

The core supermatrices and the ATPase dataset were recoded into Dayhoff groups (4 and

6) based on functional classes of amino acids, using PhyloBayes v4.1 (Lartillot *et al.*, 2009). We

also manually curated alternative alignments which had removed alignment columns if they

contained an Aspartate or Glutamate (D/E) residue that was conserved in the Nanohaloarchaea,

Haloarchaea, and Methanonatronarchaeia, to minimize compositional attraction in the ATPase

dataset.

Phylogenetic Estimation

IQTREE v1.6.9 (Nguyen *et al.*, 2015) was used to calculate maximum likelihood phylogenies for

all alignments and supermatrices. The best site homogeneous models were used for the

estimation as determined by the Bayesian Information Criterion using ModelFinder

(Kalyaanamoorthy *et al.*, 2017), for single gene phylogenies and guide tree calculation. The

ATPase and concatenated alignments (also recoded versions) were also analyzed by the LG+C60

(Le *et al.*, 2008) mixture model, all trees reported in the main text and supplemental text were

calculated using this model. Bayesian inference of Dayhoff recoded ATPase alignments were

conducted within PhyloBayes v4.1(Lartillot *et al.*, 2009; Quang *et al.*,2008) using the

CAT+GTR +G4 model in two independent chains for each alignment. These chains ran until

convergence (maxdiff < 0.25), >400,000 trees sampled, with a burn-in of the first 10% of the

trees, to calculate a majority rule consensus tree. All trees in this paper were drawn and

editorialized with Figtree v1.4.3 (Rambaut, 2016). The approximately unbiased tests were also

carried out in IQTREE (parameters: -zb 10000, -n 0) with multi-treefiles that contained

phylogenies from the hypotheses of interest (Nanohaloarchaea in DPANN or as a sister-group to

Haloarchaea) and 1000 bootstrap trees from opposing hypotheses (either from ATPases or core

genome dataset). gCF analyses of the large core supermatrix was carried out in the

IQTREEv1.7.17 beta.

Clustering of Gene Families

Individual alignments of encoded proteins were gathered for two marker sets, in 3 separate

clustering analyses: the 282 core gene families (which focused on the Nanohaloarchaea), and the

Archaea 122 (Parks *et al*., 2018) marker set (which was compiled for the Thaumarchaeota and

the Nanohaloarchaea, separately), in addition to the ATPase genes. For each taxon of interest , a

sampling of genomes from the suspected transfer partners for each taxon was also included (i.e.,

for the Nanohaloarchaeal analysis Haloarchaeal genomes were also included; for the

Thaumarchaeota genomes from Micrarchaeota and Thermoplasmatales were included (based on

Wang *et al.,* 2019)). For the Thaumarchaeota, no distinction was made between those species

that have received their ATPases via HGT and those that have not, both types of genomes were

included. Phylogenies were calculated for all the alignments in IQTREE (using the best model

determined by BIC), and pairwise distance matrices (taxon vs taxon) were generated. These

pairwise distance matrices contain distances from pairwise sequence comparisons, calculated

with maximum likelihood based on model parameter estimates from an initial tree for each

protein alignment. Using pairwise distance matrices built from sequence comparisons avoids

relying on reconstructing phylogenetic trees where the placement of our groups of interest are

already uncertain. This correlation of pairwise distance matrices of gene families is based on co-

evolution implementations found in (Gueudré, *et al.*, 2016, Rangel *et al.,* 2020)   These pairwise

distance matrices were regressed (in the sklearn Python module) against all other distance

matrices in each respective dataset (*i.e.,* each gene *vs* the 281 other genes, or 1 gene *vs* 121

genes). The ability of one distance matrix to predict the values of another distance matrix was

defined by $1\text{-}r^2$, and a summary pairwise distance matrix (gene family vs gene family) was

calculated. These values were the basis of agglomerative clustering implemented in Agnes

(cluster v2.1.0), which computes all pairwise dissimilarities between gene families and considers

the average of a pair the distance on the clustering diagram, and generated the clustered gene

families in Figure S4-5. The full implementation and instructions of this method can be found in

the gene_fam_dist.ipynb file in https://github.com/Gogarten-Lab-Team/NanoH_GBE_2020.

Acknowledgments and Funding

This work was supported by grants from the Binational Science Foundation [BSF 2013061 to

U.G., J.P.G., and R.T.P.]; the National Science Foundation [NSF/MCB 1716046 to J.P.G.,

R.T.P., and U.G.] within the BSF-NSF joint research program; and the National Aeronautics and

Space Administration exobiology [NNX15AM09G and 80NSSC18K1533 to R.T.P.]. We thank

the staff of the Bigelow Laboratory for Ocean Sciences' Single Cell Genomics Center for the

generation of single cell genomic data. We specifically thank Ramunas Stepanauskas for helpful

critical discussions and overseeing the single cell sequencing. We also thank Dr. Thiberio Rangel

for his insightful input and the code base for the gene family clustering analyses, and Dr. Jan

Gogarten for his input with the nMDS plots.

Author Contributions

The project was conceived by RTP, JPG, and UG. Sampling, sequencing, and genome

reconstruction of Dead Sea samples were conducted by UN and UG. Single cell samples were

collected by RTP, and assemblies performed by ASL and YF. Phylogenies were calculated by

YF, whole genome distance by SG. All authors contributed to writing and editing the

manuscript.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability

The newly assembled Nanohaloarchaeal genomes and accompanying information have been

deposited into GenBank under BioProject PRJNA587522 for the SAGs, and PRJNA588232 for

the MAGs. Alignments and raw newick treefiles are archived in https://github.com/Gogarten-

Lab-Team/NanoH_GBE_2020.

Supplementary Materials Table of Contents

Supplementary Figures (supp_files.pdf):

S1. tANI Distance Matrix to classify new Nanohaloarchaea.

S2. Classification of new Nanohaloarchaea genomes with tANI distances.

S3. Selected phylogenies of the Archaeal ATP synthase subunits.

S4. Clustering diagram of clustered core gene families in the Nanohaloarchaea.

S5. Diagram of clustered gene families in the Nanohaloarchaea, using the Archaeal 122

marker set.

S6. Diagram of clustered gene families in the Thaumarchaeota.

S7. Phylogenetic trees calculated from a supermatrix of ribosomal proteins.

S8. Phylogeny of recoded core supermatrices.

S9. Phylogeny of the Large core supermatrix, with DPANN sequences removed.

S10. Phylogenies of the Large core supermatrix with fast evolving sites stripped from the

alignment.

S11. Phylogeny of the 16S+23S rRNA concatenate.

S12. Constrained reference tree used to test the monophyly of the halophiles in gCF

analyses.

S13. Constrained reference tree used to test the hypothesis of the Nanohaloarchaea falling

in DPANN in gCF analyses.

S14. Protein theoretical isoelectric point distribution in selected proteomes.

S15. Figure S4 with tip labels**.**

S16. Clustering of gene families from Thaumarchaeota subsets**.**

Table S6. Concordant gene trees with each evolutionary hypothesis.

Supplementary Tables (Supplementary_tables1-5.xlsx):

Table S1. Genome files and associated statistics used in this study.

Table S2. Assembled genome statistics.

Table S3. Assembled genome gene content.

Table S4. Summary of phylogenetic analyses.

Table S5. Description of large core genome markers.

References

Adam, P. S., Borrel, G., Brochier-Armanet, C., & Gribaldo, S. (2017). The growing tree of

Archaea: New perspectives on their diversity, evolution and ecology. *ISME Journal*.

https://doi.org/10.1038/ismej.2017.122

Andrade, K., Logemann, J., Heidelberg, K. B., Emerson, J. B., Comolli, L. R., Hug, L. A.,

Probst, A., Keillar, A., Thomas B. J., Miller, C. S., Allen, E., Moreau, J., Brocks, J. J., &

Banfield, J. F. (2015). Metagenomic and lipid analyses reveal a diel cycle in a hypersaline

microbial ecosystem. *ISME Journal*. https://doi.org/10.1038/ismej.2015.66

Andrews, S. (2010). FastQC. *Babraham Bioinformatics*. https://doi.org/citeulike-article-

id:11583827

Ané, C., Larget, B., Baum, D. A., Smith, S. D., & Rokas, A. (2007). Bayesian estimation of

concordance among gene trees. *Molecular Biology and Evolution*.

https://doi.org/10.1093/molbev/msl170

Aouad, M., Borrel, G., Brochier-Armanet, C., & Gribaldo, S. (2019). Evolutionary placement of

Methanonatronarchaeia. *Nature Microbiology*. https://doi.org/10.1038/s41564-019-0359-z

Aouad, M., Taib, N., Oudart, A., Lecocq, M., Gouy, M., & Brochier-Armanet, C. (2018).

Extreme halophilic archaea derive from two distinct methanogen Class II lineages.

Molecular Phylogenetics and Evolution. https://doi.org/10.1016/j.ympev.2018.04.011

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V.,

Nikolenko, S., Pham, S., Prjibelski, S., Psyhkin, A., Sirotkin, A., Vyahhi, N., Tesler, G.,

Alekseyev, M., & Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and

its applications to single-cell sequencing. *Journal of Computational Biology : A Journal of*

Computational Molecular Cell Biology. https://doi.org/10.1089/cmb.2012.0021

Bapteste, E., Susko, E., Leigh, J., Ruiz-Trillo, I., Bucknam, J., & Doolittle, W. F. (2008).

Alternative methods for concatenation of core genes indicate a lack of resolution in deep

nodes of the prokaryotic phylogeny. *Molecular Biology and Evolution*.

https://doi.org/10.1093/molbev/msm229

Bardavid, R. E., & Oren, A. (2012). The amino acid composition of proteins from anaerobic

halophilic bacteria of the order Halanaerobiales. *Extremophiles*.

https://doi.org/10.1007/s00792-012-0455-y

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btu170

Boucher, Y., Douady, C. J., Sharma, A. K., Kamekura, M., & Doolittle, W. F. (2004).

Intragenomic heterogeneity and intergenomic recombination among haloarchaeal rRNA

genes. *Journal of Bacteriology*. https://doi.org/10.1128/JB.186.12.3980-3990.2004

Brochier-Armanet, C., Forterre, P., & Gribaldo, S. (2011). Phylogeny and evolution of the

Archaea: One hundred genomes later. *Current Opinion in Microbiology*.

https://doi.org/10.1016/j.mib.2011.04.015

Brochier, C., Gribaldo, S., Zivanovic, Y., Confalonieri, F., & Forterre, P. (2005). Nanoarchaea:

representatives of a novel archaeal phylum or a fast-evolving euryarchaeal lineage related to

Thermococcales? *Genome Biology*. https://doi.org/10.1186/gb-2005-6-5-r42

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T.

L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*.

https://doi.org/10.1186/1471-2105-10-421

Castelle, C. J., Wrighton, K. C., Thomas, B. C., Hug, L. A., Brown, C. T., Wilkins, M. J.,

Frischkorn, K., Tringe, S., Markillie, L., Taylor, R. C., Williams, K. H., & Banfield, J. F.

(2015). Genomic expansion of domain archaea highlights roles for organisms from new

phyla in anaerobic carbon cycling. *Current Biology*.

https://doi.org/10.1016/j.cub.2015.01.014

Cono, V. La, Messina, E., Rohde, M., Arcadi, E., Ciordia, S., Crisafi, F., Denaro, R., Ferrer, M.,

Giuliano, L., Golyshin, P., Golyshina, O., Hallsworth, J., Spada, G., Mena, M., Shevcenko,

M., Smedile, F., Sorokin, D. Y., Toshchakov, S., Mushegian, A., & Yakimov, M. M.

(2019). Differential polysaccharide utilization is the basis for a nanohaloarchaeon :

haloarchaeon symbiosis. *BioRxiv*. https://doi.org/10.1101/794461

Contreras-Moreira, B., & Vinuesa, P. (2013). GET_HOMOLOGUES, a Versatile Software

Package for Scalable and Robust Microbial Pangenome Analysis. *Applied and*

Environmental Microbiology. https://doi.org/10.1128/aem.02411-13

Criscuolo, A., & Gribaldo, S. (2010). BMGE (Block Mapping and Gathering with Entropy): A

new software for selection of phylogenetic informative regions from multiple sequence

alignments. *BMC Evolutionary Biology*. https://doi.org/10.1186/1471-2148-10-210

Dombrowski, N., Lee, J. H., Williams, T. A., Offre, P., & Spang, A. (2019). Genomic diversity,

lifestyles and evolutionary origins of DPANN archaea. *FEMS Microbiology Letters*.

https://doi.org/10.1093/femsle/fnz008

Dombrowski, N., Williams, T. A., Sun, J., Woodcroft, B. J., Lee, J. H., Minh, B. Q., Rinke, C.,

& Spang, A. (2020). Undinarchaeota illuminate DPANN phylogeny and the impact of gene

transfer on archaeal evolution. *Nature Communications*. https://doi.org/10.1038/s41467-

020-17408-w

Dutilh, B. E., Snel, B., Ettema, T. J. G., & Huynen, M. A. (2008). Signature genes as a

phylogenomic tool. *Molecular Biology and Evolution*.

https://doi.org/10.1093/molbev/msn115

Felsenstein, J. (1978). Cases in which Parsimony or Compatibility Methods will be Positively

Misleading. *Systematic Biology*. https://doi.org/10.1093/sysbio/27.4.401

Fournier, G. P., & Poole, A. M. (2018). A briefly argued case that Asgard Archaea are part of the

eukaryote tree. *Frontiers in Microbiology*. https://doi.org/10.3389/fmicb.2018.01896

Gadagkar, S. R., Rosenberg, M. S., & Kumar, S. (2005). Inferring species phylogenies from

multiple genes: Concatenated sequence tree versus consensus gene tree. *Journal of*

Experimental Zoology Part B: Molecular and Developmental Evolution.

https://doi.org/10.1002/jez.b.21026

Ghai, R., Pašić, L., Fernández, A. B., Martin-Cuadrado, A. B., Mizuno, C. M., McMahon, K. D.,

Papke, R. T., Stepanauskas, R., Rodriguez-Brito, B., Rohwer, F., Sanchez-Porro, C.,

Ventosa, A., & Rodríguez-Valera, F. (2011). New abundant microbial groups in aquatic

hypersaline environments. *Scientific Reports*. https://doi.org/10.1038/srep00135

Gogarten, J. P. (1994). Which is the most conserved group of proteins? Homology-orthology,

paralogy, xenology, and the fusion of independent lineages. *J Mol Evol*, *39*(5), 541–543.

Gogarten, J.P., Kibak, H., Dittrich, P., Taiz, L., Bowman, E. J., Bowman, B. J., Manolson, M. F.,

Poole, R., Date, T., & Oshima, T. (1989). Evolution of the vacuolar H+-ATPase:

implications for the origin of eukaryotes. *PNAS*, *86*(17), 6661–6665.

Gogarten, Johann Peter, & Taiz, L. (1992). Evolution of proton pumping ATPases: Rooting the

tree of life. *Photosynthesis Research*. https://doi.org/10.1007/BF00039176

Golyshina, O. V., Toshchakov, S. V., Makarova, K. S., Gavrilov, S. N., Korzhenkov, A. A., La

Cono, V., Arcadi, E., Nechitaylo, T., Ferrer, M., Kublanov, I., Wolf, Y., Yakimov, M., &

Golyshin, P. N. (2017). "ARMAN" archaea depend on association with euryarchaeal host in

culture and in situ. *Nature Communications*. https://doi.org/10.1038/s41467-017-00104-7

Graybeal, A. (1998). Is It Better to Add Taxa or Characters to a Difficult Phylogenetic Problem?

Systematic Biology. https://doi.org/10.1080/106351598260996

Gribaldo, S., Poole, A. M., Daubin, V., Forterre, P., & Brochier-Armanet, C. (2010). The origin

of eukaryotes and their relationship with the Archaea: Are we at a phylogenomic impasse?

Nature Reviews Microbiology. https://doi.org/10.1038/nrmicro2426

Gueudré, T., Baldassi, C., Zamparo, M., Weigt, M., & Pagnani, A. (2016). Simultaneous

identification of specifically interacting paralogs and interprotein contacts by direct

coupling analysis. *PNAS*. https://doi.org/10.1073/pnas.1607570113

Hamm, J. N., Erdmann, S., Eloe-Fadrosh, E. A., Angeloni, A., Zhong, L., Brownlee, C.,

Williams, T., Barton, K., Carswell, S., Smith, M., Brazendale, S., Hancock, A., Allen, M.,

Raftery, M., & Cavicchioli, R. (2019). Unexpected host dependency of Antarctic

Nanohaloarchaeota. *PNAS*. https://doi.org/10.1073/pnas.1905179116

Huber, H., Hohn, M. J., Rachel, R., Fuchs, T., Wimmer, V. C., & Stetter, K. O. (2002). A new

phylum of Archaea represented by a nanosized hyperthermophilic symbiont. *Nature*.

https://doi.org/10.1038/417063a

Joshi, N., & Fass, J. (2011). sickle - A windowed adaptive trimming tool for FASTQ files using

quality. *(Version 1.33)*. https://doi.org/10.1088/0022-3727/13/9/001

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A., & Jermiin, L. S. (2017).

ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods*.

https://doi.org/10.1038/nmeth.4285

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. *Molecular Biology and Evolution*.

https://doi.org/10.1093/molbev/mst010

Kristensen, D. M., Kannan, L., Coleman, M. K., Wolf, Y. I., Sorokin, A., Koonin, E. V., &

Mushegian, A. (2010). A low-polynomial algorithm for assembling clusters of orthologous

groups from intergenomic symmetric best matches. *Bioinformatics*.

https://doi.org/10.1093/bioinformatics/btq229

Kück, P., & Longo, G. C. (2014). FASconCAT-G: Extensive functions for multiple sequence

alignment preparations concerning phylogenetic studies. *Frontiers in Zoology*.

https://doi.org/10.1186/s12983-014-0081-x

Lanyi, J. K. (1974). Salt-dependent properties of proteins from extremely halophilic bacteria.

Bacteriological Reviews, *38*(3), 272–290. Retrieved from

https://pubmed.ncbi.nlm.nih.gov/4607500

Lapierre, P., Lasek-Nesselquist, E., & Gogarten, J. P. (2014). The impact of HGT on

phylogenomic reconstruction methods. *Briefings in Bioinformatics*.

https://doi.org/10.1093/bib/bbs050

Lapierre, P., Shial, R., & Gogarten, J. P. (2006). Distribution of F- and A/V-type ATPases in

Thermus scotoductus and other closely related species. *Systematic and Applied*

Microbiology. https://doi.org/10.1016/j.syapm.2005.06.004

Lartillot, N., Lepage, T., & Blanquart, S. (2009). PhyloBayes 3: A Bayesian software package

for phylogenetic reconstruction and molecular dating. *Bioinformatics*.

https://doi.org/10.1093/bioinformatics/btp368

Le, S. Q., Lartillot, N., & Gascuel, O. (2008). Phylogenetic mixture models for proteins.

Philosophical Transactions of the Royal Society B: Biological Sciences.

https://doi.org/10.1098/rstb.2008.0180

Li, L., Stoeckert, C. J., & Roos, D. S. (2003). OrthoMCL: Identification of ortholog groups for

eukaryotic genomes. *Genome Research*. https://doi.org/10.1101/gr.1224503

Logsdon, J. M., & Faguy, D. M. (1999). Evolutionary genomics: Thermotoga heats up lateral

gene transfer. *Current Biology*. https://doi.org/10.1016/S0960-9822(99)80474-6

Madern, D., Ebel, C., & Zaccai, G. (2000). Halophilic adaptation of enzymes. *Extremophiles :*

Life under Extreme Conditions, *4*(2), 91–98. https://doi.org/10.1007/s007920050142

Martijn, J., Schön, M. E., Lind, A. E., Vosseberg, J., Williams, T. A., Spang, A., & Ettema, T. J.

G. (2020). Hikarchaeia demonstrate an intermediate stage in the methanogen-to-halophile

transition. *Nature Communications*. Department of Medical Biochemistry and

Microbiology, Uppsala University, Uppsala, Sweden. https://doi.org/10.1038/s41467-020-

19200-2

Méheust, R., Watson, A. K., Lapointe, F. J., Papke, R. T., Lopez, P., & Bapteste, E. (2018).

Hundreds of novel composite genes and chimeric genes with bacterial origins contributed to

haloarchaeal evolution. *Genome Biology*. https://doi.org/10.1186/s13059-018-1454-9

Mohan, N. R., Fullmer, M. S., Makkay, A. M., Wheeler, R., Ventosa, A., Naor, A., Gogarten, J.

P., & Papke, R. T. (2014). Evidence from phylogenetic and genome fingerprinting analyses

1054  suggests rapidly changing variation in Halorubrum and Haloarcula populations. *Frontiers in*

1055   *Microbiology*. https://doi.org/10.3389/fmicb.2014.00143

1056 Naor, A., Lapierre, P., Mevarech, M., Papke, R. T., & Gophna, U. (2012). Low species barriers

1057   in halophilic archaea and the formation of recombinant hybrids. *Current Biology*.

1058   https://doi.org/10.1016/j.cub.2012.05.056

1059 Narasingarao, P., Podell, S., Ugalde, J. A., Brochier-Armanet, C., Emerson, J. B., Brocks, J. J.,

1060   Hiedelberg, K., Banfield, J., & Allen, E. E. (2012). De novo metagenomic assembly reveals

1061   abundant novel major lineage of Archaea in hypersaline microbial communities. *ISME*

1062   *Journal*. https://doi.org/10.1038/ismej.2011.78

1063 Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and

1064   effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*

1065   *Biology and Evolution*. https://doi.org/10.1093/molbev/msu300

1066 Olendzenski, L, Liu, L., Zhaxybayeva, O., Murphey, R., Shin, D. G., & Gogarten, J. P. (2000).

1067   Horizontal transfer of archaeal genes into the deinococcaceae: detection by molecular and

1068   computer-based approaches. *Journal of Molecular Evolution*, *51*(6), 587–599.

1069   https://doi.org/10.1007/s002390010122

1070 Olendzenski, Lorraine, Liurid, L., Zhaxybayeva, O., Murphey, R., Shin, D. G., & Gogarten, J. P.

1071   (2000). Horizontal transfer of archaeal genes into the deinococcaceae: Detection by

1072   molecular and computer-based approaches. *Journal of Molecular Evolution*.

1073   https://doi.org/10.1007/s002390010122

1074 Oren, A. (2008). Microbial life at high salt concentrations: Phylogenetic and metabolic diversity.

1075   *Saline Systems*. https://doi.org/10.1186/1746-1448-4-2

1076 Parks, D. H., Chuvochina, M., Waite, D. W., Rinke, C., Skarshewski, A., Chaumeil, P. A., &

Hugenholtz, P. (2018). A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. *Nature Biotechnology*. https://doi.org/10.1038/nbt.4229

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM:

Assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Research*. https://doi.org/10.1101/gr.186072.114

Paul, S., Bag, S. K., Das, S., Harvill, E. T., & Dutta, C. (2008). Molecular signature of

hypersaline adaptation: insights from genome and proteome composition of halophilic

prokaryotes. *Genome Biology*, *9*(4), R70. https://doi.org/10.1186/gb-2008-9-4-r70

Petitjean, C., Deschamps, P., López-Garciá, P., & Moreira, D. (2014). Rooting the domain

archaea by phylogenomic analysis supports the foundation of the new kingdom

Proteoarchaeota. *Genome Biology and Evolution*. https://doi.org/10.1093/gbe/evu274

Philippe, H., Zhou, Y., Brinkmann, H., Rodrigue, N., & Delsuc, F. (2005). Heterotachy and

long-branch attraction in phylogenetics. *BMC Evolutionary Biology*.

https://doi.org/10.1186/1471-2148-5-50

Quang, L. S., Gascuel, O., & Lartillot, N. (2008). Empirical profile mixture models for

phylogenetic reconstruction. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btn445

Rambaut, A. (2016). FigTree v1.4.3.

Rangel, L. T., Marden, J., Colston, S., Setubal, J. C., Graf, J., & Gogarten, J. P. (2019).

Identification and characterization of putative Aeromonas spp. T3SS effectors. *PLoS ONE*.

https://doi.org/10.1371/journal.pone.0214035

Raymann, K., Brochier-Armanet, C., & Gribaldo, S. (2015). The two-domain tree of life is

linked to a new root for the Archaea. *PNAS*. https://doi.org/10.1073/pnas.1420858112

Raymann, K., Forterre, P., Brochier-Armanet, C., & Gribaldo, S. (2014). Global phylogenomic

analysis disentangles the complex evolutionary history of DNA replication in Archaea.

GBE. https://doi.org/10.1093/gbe/evu004

Richter, M., & Rosselló-Móra, R. (2009). Shifting the genomic gold standard for the prokaryotic

species definition. *PNAS*. https://doi.org/10.1073/pnas.0906412106

Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N. N., Anderson, I. J., Cheng, J. F., Darling, A.,

Malfatti, S., Swan, B., Gies, E., Dodsworth, J., Hedlund, B., Tsiamis, G., Sievert, S., Liu,

W. T., Eisen, J., Hallam S. J., Kyrpides, N., Stepanauskas, R., Rubin, E., Hugenholtz, P., &

Woyke, T. (2013). Insights into the phylogeny and coding potential of microbial dark

matter. *Nature*. https://doi.org/10.1038/nature12352

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*.

https://doi.org/10.1093/bioinformatics/btu153

Sorokin, Di. Y., Makarova, K. S., Abbas, B., Ferrer, M., Golyshin, P. N., Galinski, E. A.,

Ciordia, S., Mena, M., Merkel, A., Wolf, Y., van Loosdrecht, M., & Koonin, E. V. (2017).

Discovery of extremely halophilic, methyl-reducing euryarchaea provides insights into the

evolutionary origin of methanogenesis. *Nature Microbiology*.

https://doi.org/10.1038/nmicrobiol.2017.81

Spang, A., Caceres, E. F., & Ettema, T. J. G. (2017). Genomic exploration of the diversity,

ecology, and evolution of the archaeal domain of life. *Science*, *357*(6351), eaaf3883.

https://doi.org/10.1126/science.aaf3883

Spang, A., Eme, L., Saw, J. H., Caceres, E. F., Zaremba-Niedzwiedzka, K., Lombard, J., Guy,

L., & Ettema, T. J. G. (2018). Asgard archaea are the closest prokaryotic relatives of

eukaryotes. *PLoS Genetics*. https://doi.org/10.1371/journal.pgen.1007080

Susko, E., & Roger, A. J. (2007). On reduced amino acid alphabets for phylogenetic inference.

Molecular Biology and Evolution. https://doi.org/10.1093/molbev/msm144

Urbonavičius, J., Auxilien, S., Walbott, H., Trachana, K., Golinelli-Pimpaneau, B., Brochier-

Armanet, C., & Grosjean, H. (2008). Acquisition of a bacterial RumA-type tRNA(uracil-54,

C5)-methyltransferase by Archaea through an ancient horizontal gene transfer. *Molecular*

Microbiology. https://doi.org/10.1111/j.1365-2958.2007.06047.x

Varghese, N. J., Mukherjee, S., Ivanova, N., Konstantinidis, K. T., Mavrommatis, K., Kyrpides,

N. C., & Pati, A. (2015). Microbial species delineation using whole genome sequences.

Nucleic Acids Research. https://doi.org/10.1093/nar/gkv657

Vavourakis, C. D., Ghai, R., Rodriguez-Valera, F., Sorokin, D. Y., Tringe, S. G., Hugenholtz, P.,

& Muyzer, G. (2016). Metagenomic insights into the uncultured diversity and physiology of

microbes in four hypersaline soda lake brines. *Frontiers in Microbiology*.

https://doi.org/10.3389/fmicb.2016.00211

Wang, B., Qin, W., Ren, Y., Zhou, X., Jung, M.-Y., Han, P., Eloe-Fadrosh, E., Li, M., Zheng,

Y., Lu, L., Yan, X., Ji, J., Liu, Y., Liu, L., Heiner, C., Hall, R., Martens-Habbena, W.,

Herbold, C., Rhee, S., Bartlett, D., Huang, L., Ingalls, A., Wagner, M., Stahl, D., & Jia, Z.

(2019). Expansion of Thaumarchaeota habitat range is correlated with horizontal transfer of

ATPase operons. *ISME J*. https://doi.org/10.1038/s41396-019-0493-x

Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. *PLoS Computational Biology*.

https://doi.org/10.1371/journal.pcbi.1005595

Wick, R. R., Schultz, M. B., Zobel, J., & Holt, K. E. (2015). Bandage: Interactive visualization

of de novo genome assemblies. *Bioinformatics*.

https://doi.org/10.1093/bioinformatics/btv383

Williams, D., Gogarten, J. P., & Papke, R. T. (2012). Quantifying homologous replacement of

loci between haloarchaeal species. *Genome Biology and Evolution*.

https://doi.org/10.1093/gbe/evs098

Williams, T. A., Cox, C. J., Foster, P. G., Szöllősi, G. J., & Embley, T. M. (2020).

Phylogenomics provides robust support for a two-domains tree of life. *Nature Ecology and*

Evolution. https://doi.org/10.1038/s41559-019-1040-x

Williams, T. A., & Embley, T. M. (2014). Archaeal "dark matter" and the origin of eukaryotes.

GBE. https://doi.org/10.1093/gbe/evu031

Williams, T. A., Foster, P. G., Cox, C. J., & Embley, T. M. (2013). An archaeal origin of

eukaryotes supports only two primary domains of life. *Nature*.

https://doi.org/10.1038/nature12779

Williams, T. A., Heaps, S. E., Cherlin, S., Nye, T. M. W., Boys, R. J., & Embley, T. M. (2015).

New substitution models for rooting phylogenetic trees. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *370*(1678), 20140336.

https://doi.org/10.1098/rstb.2014.0336

Williams, T. A., Szöllősi, G. J., Spang, A., Foster, P. G., Heaps, S. E., Boussau, B., Ettema, T.,

& Embley, T. M. (2017). Integrative modeling of gene and genome evolution roots the

archaeal tree of life. *Proceedings of the National Academy of Sciences*.

https://doi.org/10.1073/pnas.1618463114

Zhaxybayeva, O., Stepanauskas, R., Mohan, N. R., & Papke, R. T. (2013). Cell sorting analysis

of geographically separated hypersaline environments. *Extremophiles*.

https://doi.org/10.1007/s00792-013-0514-z

Zhaxybayeva, O., Swithers, K. S., Lapierre, P., Fournier, G. P., Bickhart, D. M., DeBoy, R. T.,

Nelson, K., Nesbo, C., Doolittle, W. F., Gogarten, J. P., & Noll, K. M. (2009). On the

chimeric nature, thermophilic origin, and phylogenetic placement of the Thermotogales.

PNAS. https://doi.org/10.1073/pnas.0901260106

1172

1173

1174