Neural modelling of the semantic predictability gain under challenging listening conditions

Anna Rysop [1]*, Lea-Maria Schmitt [2]*, Jonas Obleser [2]#, & Gesa Hartwigsen [1]#

[1] Lise Meitner Research Group Cognition and Plasticity, Max Planck Institute for Human Cognitive and Brain Sciences, Stephanstrasse 1a, 04103 Leipzig, Germany

[2] Department of Psychology, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

* both authors contributed equally

shared senior authorship

Data availability: All data are available from the corresponding authors upon reasonable request.

Funding: This work was supported by the German Research Foundation (DFG; HA 6314/4-1 and OB 352/2-1). JO is supported by an ERC Consolidator Grant (ERC-CoG-2014; grant number 646696).

Conflict of interest: The authors declare no conflicts of interest.

Ethics approval: All experimental procedures were approved by the local ethics committee at the Medical Faculty of the University of Leipzig (reference number 281/16-ek).

Patient consent: All participants gave written informed consent.

Permission to reproduce material from other sources: Not applicable.

Clinical trial registration: Not applicable.

Correspondence:
Anna Rysop (rysop@cbs.mpg.de; https://orcid.org/0000-0002-9853-358X)
Lea-Maria Schmitt (l.schmitt@uni-luebeck.de; https://orcid.org/0000-0002-9356-2234)
Jonas Obleser (jonas.obleser@uni-luebeck.de; https://orcid.org/0000-0002-7619-0459)
Gesa Hartwigsen (hartwigsen@cbs.mpg.de; https://orcid.org/0000-0002-8084-1330)

Abstract

When speech intelligibility is reduced, listeners exploit constraints posed by semantic context to facilitate comprehension. The left angular gyrus (AG) has been argued to drive this semantic predictability gain. Taking a network perspective, we ask how the connectivity within language-specific and domain-general networks flexibly adapts to the predictability and intelligibility of speech. During continuous functional magnetic resonance imaging (fMRI), participants repeated sentences, which varied in semantic predictability of the final word and in acoustic intelligibility. At the neural level, highly predictable sentences led to stronger activation of left-hemispheric semantic regions including subregions of the AG (PGa, PGp) and posterior middle temporal gyrus when speech became more intelligible. The behavioural predictability gain of single participants mapped onto the same regions but was complemented by increased activity in frontal and medial regions. The facilitatory influence of PGp on PGa (quantified by dynamic causal modelling) increased for highly predictable sentences. In contrast, cingulo-opercular regions showed stronger activation for sentences with low predictability under increasing intelligibility when compared to high predictability. Critically, better individual performance in highly predictable sentences correlated with a stronger inhibitory influence of pre-supplementary motor area on right insula. These results suggest that successful comprehension in noisy listening conditions relies on an interplay of semantic regions and concurrent inhibition of cognitive control regions when semantic cues are available.

Keywords: angular gyrus, cingulo-opercular network, semantic network, predictability gain, speech in noise comprehension.

Introduction

In everyday life, we are remarkably successful in following a conversation even when background noise degrades the speech signal. One important strategy facilitating comprehension in demanding listening scenarios is the prediction of upcoming speech. For example, the sentence fragment "She made the bed with new" provides rich semantic context to inform an accurate prediction of the sentence continuation "sheets". However, poor semantic context like "We are very pleased with the new" provides little information to build up the prediction of the word "sheets". This *semantic predictability gain* is a well-documented behavioural phenomenon. It can be observed in healthy young and older listeners (Obleser et al. 2007; Obleser and Kotz 2010; Vaden et al. 2013; Vaden et al. 2015), in hearing-impaired listeners (Holmes et al. 2018), as well as in cochlear implant users (Winn 2016).

To study the interactive effects of intelligibility and semantic predictability in the listening brain, previous neuroimaging studies used sentences with different occurrence probabilities of the final word while varying the spectral resolution of speech using noise-vocoding (see Shannon et al. (1995)). In these studies, the left angular gyrus (AG) has been identified as a key region sensitive to strong semantic constraints in degraded speech (Obleser et al. 2007; Obleser and Kotz 2010; Golestani et al. 2013). When highly predictable speech was degraded to a moderate extent, activity in AG increased. However, when degradation rendered speech either easily or, if it all, hardly comprehensible, activity in AG dropped irrespective of semantic constraints. Finally, Hartwigsen and colleagues (2015) used focal perturbations induced by transcranial magnetic stimulation to demonstrate that the ventral portion of left AG is functionally relevant for the behavioural predictability gain. Together, these studies provide converging evidence for a role of the left AG in the top-down use of semantic information when comprehension is challenged by degraded speech signals.

The AG is considered a heterogeneous region and has structural connections to language-specific fronto-temporal regions as well as domain-general cingulo-opercular regions (Binder et al. 2009, Seghier 2013). Based on its cytoarchitectonics, AG can be subdivided into a rostral part (PGa) and a caudal part (PGp; Caspers et al., 2006, 2008). These

subregions have been associated with different functional roles: PGa has been associated with the automatic and domain-general allocation of goal-directed attention and episodic memory. In contrast, PGp was (inconsistently) linked to more controlled and complex semantic-specific processes irrespective of control demands (Noonan et al. 2013; Bonnici et al. 2016; Bzdok et al. 2016; Jung et al. 2017). These findings indicate that the AG might play a central role in language-specific but also in domain-general cognitive operations. Yet, the functional interplay of PGa and PGp within left AG as well as their differential roles in the processing of semantic context in degraded speech are unknown.

As comprehension of degraded speech poses a challenging task on listeners, the brain recruits additional resources beyond the semantic network (Peelle 2018). These additional resources are thought to be provided by domain-general regions that typically do not show a functional specification, such as the cingulo-opercular network. The cingulo-opercular network has been implicated in adaptive control processes subserving the flexible allocation of attentional resources (Dosenbach et al. 2008; Duncan 2010). Indeed, previous studies on degraded speech processing have found not only fronto-temporo-parietal regions of the semantic network (Adank 2012; Alain et al. 2018) but also regions of the cingulo-opercular network (Erb et al. 2013; Vaden et al. 2015; Vaden et al. 2017). Additionally, using a word recognition task under challenging listening conditions, Vaden and colleagues (2013) demonstrated that the magnitude of activation within the cingulo-opercular network predicted the word recognition success in the following trial. This upregulation might reflect the activation of additional cognitive resources when no external linguistic cues are available (Peelle 2018). In contrast, the recruitment of the semantic network is thought to improve comprehension most effectively when semantic context is available. Here, we aim to identify network-level descriptors of the semantic predictability gain to comprehension and investigate how changing task demands shape the adaptive interplay of language-specific and domain-general resources in successful speech comprehension.

Previous studies manipulating predictability and intelligibility of speech suffered from small samples and mainly relied on few levels of noise fixed across participants. Moreover, most of

these studies used sparse temporal sampling, precluding the investigation of effective connectivity. Consequently, it remains unclear how the identified regions influence each other during successful comprehension of speech in noise. We aimed to overcome some of the previous shortcomings by performing a continuous event-related functional magnetic resonance imaging (fMRI) experiment with six individualized, varying levels of intelligibility, ranging from largely unintelligible to easily intelligible. Thereby, we accounted for individual differences in auditory perception and assured comparable task difficulty across participants.

Based on previous studies, we expected a facilitatory behavioural effect of semantic predictability on speech comprehension, which should be strongest at intermediate levels of intelligibility (i.e. when the auditory signal is somewhat degraded but still intelligible) and least pronounced for unintelligible and intelligible sentences. We hypothesized to find increased activation in left AG and other semantic regions as a neural correlate of the semantic predictability gain at intermediate levels of intelligibility. This interaction should be driven by increased effective connectivity between the AG and other semantic regions. Moreover, we expected increasing activity and within-network connectivity in the cingulo-opercular network to improve comprehension in challenging listening conditions (i.e., intermediate intelligibility and low predictability).

As a main finding of our study, highly predictable sentences drove activity in core regions of the semantic network and increased the functional influence between subregions of the left AG. The behavioural comprehension gain was correlated with higher activity in extended regions of the semantic network at intermediate intelligibility levels. Processing of less predictable sentences drew on cognitive control regions and strengthened effective connectivity between regions within the cingulo-opercular network, while higher predictability decreased connectivity within this network. The individual degree of inhibitory influence of higher predictability on the connection from the pre-supplementary motor area (pre-SMA) to the right insula was associated with the individual predictability gain.

1 Materials and Methods

2 Participants

Thirty healthy young German native speakers took part in this study. After excluding three

participants due to excessive head movement during scanning (i.e. range of motion > 1.5 times

the voxel size; see Supplementary Figure 1 for an illustration of head movements) and one

participant due to reduced task-related activity across the whole brain in the global F-test, our

final sample yielded 26 participants (age range: 19–29 years, M = 25 years; 15 females). All

participants were right-handed (mean lateralization index > 90; Oldfield, 1971) and reported

no history of neurological or psychiatric disorders as well as no hearing difficulties or disorders.

Participants gave written informed consent prior to participation and received

reimbursement of €10 per hour of testing. The study was performed according to the guidelines

of the Declaration of Helsinki and approved by the local ethics committee at the Medical Faculty

of the University of Leipzig.

14 Experimental Procedures

Participants took part in one fMRI session. During this session, we first assessed the individual

ability of participants to comprehend speech in noise by tracking the speech reception

threshold (SRT). Second, participants performed a repetition task on sentences varying in

intelligibility and predictability (Figure 1). The task had a 2 x 6 full factorial design (predictability:

high vs. low; intelligibility: -9, -4, -1, +1, +4, +9 dB sentence intensity relative to the individual

SRT) with 12 experimental conditions and 18 trials per condition.

Experimental task design

High Predictability
Sie bezieht das Bett mit frischen **Laken**.
She made the bed with new **sheets**

Low Predictability
Wir freuen uns sehr über die **Laken**.
We please ourselves very with the **sheets**

Intelligibility (dB)
SRT 50

Figure 1. Design of the sentence repetition task. In each trial, participants listened to a
sentence embedded in distracting background noise while viewing a fixation cross on the
screen. With the onset of the sentence's final keyword, a green traffic light prompted
participants to orally repeat the whole sentence as accurately as possible within a 5 sec
recording period. The preceding sentence context was either predictive (green) or non-
predictive of the keyword (orange). Sentences varied orthogonally in intelligibility (ratio of
speech intensity to SRT).

Adaptive tracking procedure of the speech reception threshold

To account for inter-individual differences in speech perception acuity in the main experiment,

we determined each participant's SRT beforehand, that is the signal-to-noise ratio (SNR)

required to correctly repeat a sentence with a probability of 50%. In an adaptive up-down

staircase procedure (Kollmeier et al. 1988), participants listened to 20 sentences with highly

predictable keywords energetically masked by speech-shaped noise. After the presentation of

each sentence, participants repeated the whole sentence as accurately as possible and the

investigator directly rated the response as either correct or incorrect. A correct repetition was

followed by an SNR decrease (i.e. the following sentence was less intelligible), an incorrect

repetition was followed by an SNR increase (i.e. the following sentence was more intelligible).

The tracking procedure started with an initial SNR of 5 dB and a trial-to-trial step size of 6 db.

A turning point (i.e., an increase in SNR is followed by a decrease or vice versa) reduced the

step size by a factor of 0.85. We determined SRT's as the average SNR on the last five trials

leading to a turn point. SRTs had an average of +1.7 dB SNR (SD = 2.3, range: -2.1 to +6.4

dB). All sentences were presented in the MR scanner while running the fMRI sequence later

used in the experiment. We used the hereby determined SRTs as a reference for the

manipulation of intelligibility in the experiment to account for inter-individual differences in

speech-in-noise comprehension and effectively modulate behavioural performance.

Stimulus material

To manipulate predictability of speech in the sentence repetition task, we used the German

version of the speech in noise (SPIN) corpus (Kalikow, Stevens, & Elliott, 1977; for a detailed

description of the German version, see Erb, Henry, Eisner, & Obleser, 2012). The German

SPIN corpus contains pairs of spoken sentences which have the same final word (i.e. keyword)

but differ in cloze probability (i.e. the expectancy of a word given the preceding sentence;

Taylor 1953) of the keyword. One sentence of a pair provides poor semantic context for the

prediction of the keyword ("We are very pleased with the new <u>sheets</u>."; low predictability),

whereas the complementary sentence provides rich context for the same keyword ("She made

the bed with new <u>sheets</u>."; high predictability). More specifically, the predictability of the

keyword varies with the number of pointer words in a sentence that semantically link to the

sentence-final word (e.g., "made", "bed", and "new" in the above example). Keywords from

sentences with rich context had a mean cloze probability of 0.85 (SD = 0.14), low predictable

keywords had a mean cloze probability of 0.1 (SD = 0.02).

In addition to the 100 original sentence pairs, we extended the corpus by another 36

sentences which were constructed and recorded during the development of the original

German SPIN corpus. Initially, these sentences were not included in the corpus, as a smaller

set of sentences was sufficient for the purposes of the developers. To obtain predictability

ratings for the keywords in the new sentences, 10 participants who did not take part in the fMRI

experiment (age range: 24–28 years; M = 26 years; 4 females) performed a written sentence

completion test (low predictability: < 5 participants reported the correct keyword; high

predictability: > 5 participants). Together, 8 new sentence pairs and the original German SPIN

sentence pairs yielded 216 experimental sentences (M = 2143 ms, SD = 256 ms). The

remaining 20 new sentences were used in the adaptive tracking procedure prior to the

experiment (see Supplementary Figure 2 for a detailed illustration of how sentences were

assigned to the experiment and the adaptive tracking procedure).

Intelligibility of speech was manipulated by varying the intensity of the experimental sentences relative to a constant signal intensity of speech-shaped noise in the background. We created spectrally speech-shaped noise by filtering white noise with the long-term average spectrum of all experimental sentences (e.g. Nilsson, Soli, & Sullivan, 1994). The noise stream preceded and ensued single sentences by 250 ms. Our six experimental intelligibility levels were symmetrically spaced around individual SRTs and logarithmically increased towards the extremes. With our intelligibility levels, we aimed to cover the full range of behavioural performance in each participant and sample intermediate intelligibility in smaller steps as SNR changes in this range affect performance more strongly.

Sentence repetition task

Participants performed an overt sentence repetition task during fMRI. They listened to single sentences while viewing a white fixation cross on a black screen. With the onset of the keyword, a green traffic light appeared on the screen and indicated the start of a 5000 ms recording period. During the recording period, participants orally repeated the sentence as accurately as possible. Whenever participants did not comprehend the complete sentence, they repeated the sentence fragments they could grasp. If participants could not repeat any word, they said that they did not understand the sentence. The intertrial interval was randomly set between 2000 and 7000 ms and served as an implicit baseline for the fMRI analysis. Sentences were presented in a pseudorandomized order avoiding the repetition of one intelligibility level more than three times in a row to prevent adaptation to specific intelligibility levels. The experiment was split in six blocks of approximately 8 minutes which were intermitted by pauses of 20 s each.

All sentences were presented at a comfortable volume. Sound was played via MR-Confon headphones (Magdeburg, Germany) and recorded via a FOMRI-III microphone (Optoacoustics, Yehuda, Israel). Stimulus presentation was controlled by Presentation software (version 18.0, Neurobehavioral Systems, Berkeley, USA, www.neurobs.com). The experiment lasted 50 minutes.

MRI acquisition

Whole brain fMRI data were acquired in one continuous run with a 3 Tesla Siemens Prisma

Scanner and 32-channel head coil, using a dual gradient-echo planar imaging multiband

sequence (Feinberg et al., 2010) with the following scanning parameters: TR = 2,000 ms; TE

= 12 ms, 33 ms; flip angle = 90°; voxel size = 2.5 x 2.5 x 2.5 mm with an interslice gap of 0.25

mm; FOV = 204 mm; multiband acceleration factor = 2. In sum, 1,500 volumes were acquired

per participant, each consisting of 60 slices in axial direction and interleaved order. To increase

coverage of anterior temporal lobe regions, slices were tilted by 10° off the AC-PC line. Field

maps were acquired for later distortion correction (TR = 620 ms; TE = 4 ms, 6.46 ms).

Additionally, high-resolution T1-weighted images were either obtained from the in-house

database if available or were acquired prior to the functional scans with an MPRAGE sequence

(whole brain coverage, TR = 1300 ms, TE = 2.98 ms, voxel size = 1 x 1 x 1 mm, matrix size =

256 x 240 mm, flip angle = 9°).

Data Analysis

Behavioural Data Analysis

All of participants' spoken response recordings from the main experiment were cleaned from

scanner noise using Audacity (version 2.2.2, https://www.audacityteam.org/). For

transcriptions, we split the recordings in two independent sets and assigned each recording

set to one of two raters. To validate the quality of transcripts, a third rater additionally

transcribed half of the recordings in each set. Cohen's kappa coefficient (Cohen 1960)

indicated very good inter-rater reliability for keywords (rater 1 and 3: κ = .94; rater 2 and 3: κ =

.97). Finally, one rater manually determined onset and duration of oral responses.

As only predictability of the final keyword but not its preceding context is explicitly

manipulated in the German SPIN corpus, we limited behavioural analyses to keywords. After

rating keywords as incorrect (including missing, incomplete and inaccurately inflected

keywords) or correct, we calculated the proportion of correctly repeated keywords for each

participant and condition.

In the behavioural analysis, we quantified the effects of predictability and intelligibility on speech comprehension. For each participant, we fitted psychometric curves to the proportion of correct keywords across intelligibility levels, separately for sentences with low and high predictability. In detail, we fitted cumulative Gaussian sigmoid functions as beta-binomial observer models using the Psignifit toolbox (Fründ et al. 2011) in MATLAB (version R2018b, MathWorks). To estimate all five parameters describing the function (guess and lapse rate, threshold, width and overdispersion), we applied Psignifit's default priors for experiments with binary single-trial outcome (here: correctly vs. incorrectly repeated keyword; Schütt et al. 2016). As indicated by model R^2 for the proportion correct data, the goodness of fit was very good for all remaining psychometric curves of sentences with high ($M = 0.97$, $SD = 0.03$, range: 0.88-0.997) and low predictability ($M = 0.96$, $SD = 0.04$, range: 0.85-0.998).

As all 12 experimental conditions thus become subsumed in two psychometric curves, interaction effects can be evaluated by comparing parameter estimates between curves for sentences with low and high predictability. The threshold represents the intelligibility level at which participants correctly repeat half of the keywords. Keeping all other parameters constant, a threshold shift to the left (i.e. a smaller value) indicates a comprehension gain at intermediate but not high and low intelligibility levels. We hypothesized that the interaction effect of intelligibility and predictability manifests in a smaller threshold for sentences with high predictability when compared to low predictability. Additionally, we expected that an increase in intelligibility leads to a stronger comprehension gain for sentences with high relative to low predictability. The sensitivity to changes in intelligibility is reflected by the slope which is inversely related to the width and here describes the steepness of a psychometric curve at a proportion correct of 0.5. To explore whether also other parameters of psychometric curves were modulated by the predictability of sentences, we compared guess and lapse rate (i.e., lower and upper asymptote) as well as width between sentences with high and low predictability. Parameter estimates of the two psychometric curves were contrasted using a paired-sample *t*-test (or a Wilcoxon signed-rank test whenever the assumption of normality was not met).

We quantified the strength of evidence in favour of our hypotheses by calculating the

Bayes Factor (BF; Jeffreys 1961) for all t-tests and correlations carried out in the behavioural

analyses using the default settings in JASP (i.e., comparing against a Cauchy prior with scale

$= \sqrt{2} / 2$; version 0.8.6.0). Whereas $BF_{10} = 1$ supports neither the alternative nor the null

hypothesis, $BF_{10} > 3$ indicates increasingly substantial support for the alternative hypothesis

(Kass and Raftery 1995). For $BF_{10} = 3$, the data are 3 times more likely to have occurred under

the alternative hypothesis than the null hypothesis. As an additional effect size measure, we

report Pearson's correlation coefficient *r* (Rosenthal and Rubin 1994).

Functional MRI Data

Functional MRI data were analysed using the Statistical Parametric Mapping software

(SPM12, version 7219, Wellcome Department of Imaging Neuroscience, London, UK) and

MATLAB (version R2017b). Preprocessing steps were applied in line with SPM's default

settings and comprised realignment, distortion correction, segmentation, co-registration to the

individual high-resolution structural T1-scan and spatial normalization to the standard template

by the Montreal Neurological Institute (MNI) while keeping the original voxel size. Volumes

were smoothed using a 5 mm full width at half maximum Gaussian kernel to allow statistical

inference based on Gaussian random-field theory (Friston et al. 1994). Realignment

parameters were obtained from the first echo. Temporal signal-to-noise ratio maps were

calculated for each echo based on the first 30 volumes of each participant. Echoes were

combined using the information from the corresponding temporal signal-to-noise ratio maps as

weighting factors and realigned to the first scan. This approach has been proposed to increase

SNR in brain regions typically suffering from signal loss (e.g., anterior temporal lobe regions;

see Halai, Welbourne, Embleton, & Parkes, 2014 for a similar approach). Motion parameters

were calculated from the parameters obtained by realignment. Additionally, framewise

displacement as described by Power et al. 2012 was calculated as an index of head

movement. Volumes that exceeded a framewise displacement of 0.9 (Siegel et al. 2014) were

included as nuisance regressors.

At the single-participant level, we set up a general linear model (GLM). For each experimental condition, onset and duration of the sentence presentation period were modelled with a stick function and convolved with SPM's canonical hemodynamic response function. Additionally, we modelled onset and duration of the sentence repetition periods using individual speech onset times and speech durations. We further included motion parameters obtained from the realignment step and framewise displacement as regressors of no interest. Data were high-pass filtered with a cut-off at 128 s and serial correlations were taken into account using a first-order autoregressive model. Each experimental regressor was contrasted against the implicit baseline (intertrial interval), thus yielding 12 contrast images of interest, one for each experimental condition. Additionally, an *F*-contrast containing the experimental regressors was set up to capture experimental effects of interest and exclude effects of no interest (i.e. sentence repetition periods, motion-related activity).

At the group level, these contrasts were submitted to a random-effects flexible factorial design with the experimental factors predictability (high, low) and intelligibility (-9 to +9 dB) entered as interaction term, and an additional factor modelling the subjects' constant. First, we set up a *t*-contrast modelling a linear increase in intelligibility (-9 > -4 > -1 > +1 > +4 > +9 dB SNR) to reveal brain regions that are implicated in speech processing irrespective of predictability (i.e. main effect of intelligibility).

Further, we were interested in those brain regions showing a linear or quadratic interaction of intelligibility and predictability. The interaction contrasts were set up as separate t-contrasts for linearly (high > low, -9 > -4 > -1 > +1 > +4 > +9 dB SNR relative to SRT) and quadratically modelled intelligibility (high > low, -9 < -4 < -1 = +1 > +4 > +9 dB). These parametric interaction contrasts were set up in both directions (high > low / low > high).

Single-participant's parameter estimates were extracted from significant peak voxels and transformed into percent signal change using the rfx toolbox (Gläscher 2009) for illustration purposes. Figures were created using MRIcroGL (www.mricro.com, version v1.0.20180623).

All effects for the whole-brain group analysis are reported at a voxel-level family-wise error (FWE) correcting threshold of $p < 0.05$ to control for multiple comparisons. We do not report

clusters with less than 20 voxels for the main effect and 10 voxels for the interaction effects, as we consider such small clusters biologically implausible. Local maxima are reported in MNI space. We used the SPM Anatomy Toolbox (Eickhoff et al., 2005, version 2.2c) and the Harvard-Oxford cortical structural atlas (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki) to classify the anatomical correspondence of significant voxels and clusters.

Brain–Behaviour Correlation

We identified brain regions reflecting the behavioural predictability gain across intelligibility levels by correlating neural response patterns with task performance. On the behavioural level, we extracted the proportion of correctly repeated keywords in each experimental condition and participant as modelled by psychometric curves.

To quantify how strongly the proportion correct in each intelligibility level was affected by the predictability of sentences, we projected task performance to the interaction subspace: the condition-wise proportion correct of each participant was multiplied with the *F*-contrast of the interaction effect between intelligibility and predictability (for a detailed description see Henson, 2015). This procedure yielded the proportion of variance in performance explained by the interaction, i.e. the predictability gain (or loss) in each intelligibility level.

On the neural level, we limited our analysis to all voxels consistently implicated in task-related activity as indicated by a significant *F*-contrast ($p_{uncorrected} < 0.05$) calculated across single-participant parameter estimates of all experimental conditions (see section 2.4.2.2). For each voxel within the mask, we correlated single-participant behavioural predictability gain and neural parameter estimates across conditions. After applying Fisher's z-transformation to all Pearson product-moment correlation coefficients, single-participant maps were submitted to the Local Indicators of Spatial Association (LISA) group-level one-sample *t*-test (Lohmann et al. 2018). LISA is a threshold-free framework that allows to find consistent effects in small brain regions by applying a non-linear filter to statistical maps before controlling for multiple comparisons at a false discovery rate (FDR) < 0.05. Voxel-wise FDR scores were obtained from a Bayesian two-component mixture model in which filtered statistical values are

compared to a null distribution based on 5000 permutations (i.e., randomly switching signs of statistical maps in participants).

Dynamic Causal Modelling (DCM)

To investigate condition-specific interactions between brain areas, we conducted two separate bilinear one-state deterministic Dynamic Causal Modelling (DCM) analyses using the DCM 12.5 implementation in SPM 12. DCM is a forward-model-based method that models the dynamics of hidden neuronal states within a predefined set of regions and relates them to the measured blood-oxygen-level-dependent (BOLD) signal (Friston et al. 2003). Using this Bayesian framework, modulations of effective connectivity by experimental conditions can be assessed.

Seed region selection

According to our main hypothesis, we were interested in the neural network dynamics underlying the predictability gain in speech in noise processing and the role of left angular gyrus. In the GLM analysis, we found two distinct networks implicated in the neural interaction of predictability and intelligibility: the semantic network and the cingulo-opercular network (CO). Here, we test effective connectivity in a subset of regions within each network. The semantic model was based on the GLM interaction contrast (high > low, linearly modelled intelligibility) and included left posterior middle temporal gyrus (pMTG), PGa and PGp as seed regions. This model specification allowed to investigate differential contributions of subregions within AG, namely PGa and PGp. Based on the opposite GLM interaction contrast (low > high, linearly modelled intelligibility), we specified the CO model including pre-SMA/paracingulate gyrus as well as left and right anterior insula as seed regions.

For each subject, we identified the peak voxel nearest to the group maximum of each seed region within a sphere of 10 mm using the appropriate first-level contrast. Individual timeseries of each region (summarized as the first eigenvariate) were extracted from all voxels within 6 mm spheres centered on the individual maximum and exceeding the liberal threshold of $p_{uncorrected} < 0.05$. The extracted timeseries were adjusted to the respective effects-of-interest

F-contrast to exclude variance of no interest. The exact anatomical localization of individual seed regions was verified using SPM's Anatomy Toolbox. For the semantic model, we made sure that seed regions were not overlapping within each subject.

The summarized timeseries together with information about the onsets of the relevant conditions served as input for the DCM. The design matrix for the DCM analyses differed from the design matrix used in the GLM analysis in the following way: three regressors were defined, modelling the onsets of (1) all experiment stimuli, (2) all high predictable sentences at intermediate-to-good intelligibility levels (-1, +1, +4, +9 dB SNR relative to SRT), and (3) all low predictable sentences at intermediate-to-good intelligibility levels. The first regressor served as sensory input (i.e. driving input) of the stimuli into the model. Regressors (2) and (3) were used as modulatory inputs and encoded whether the stimulus had high or low predictability. As inputs were not mean-centered, intrinsic parameter estimates can be interpreted as average connection strength in the absence of experimental manipulations (Marreiros et al. 2008).

In the DCM analysis, we were mainly interested in the neural network dynamics underlying *context-dependent* changes in *successful* speech comprehension. Therefore, we excluded all intelligibility levels that showed no apparent second-level difference between sentences with high and low predictability in behavioural repetition accuracies as well as in parameter estimates of the interaction contrast. We also excluded intelligibility levels with average task performance below 50% in both predictability conditions. This procedure resulted in an exclusion of the two most unintelligible SNR levels (-9 and -4 dB SNR relative to SRT).

DCM model architecture

At the first-level, 84 models were specified for each subject. All models consisted of three types of parameters: (i) intrinsic connections between two nodes (unidirectional or bidirectional), (ii) changes in coupling strength between regions dependent on the experimental manipulation (modulatory effects) and (iii) direct influence of external input on a region (driving input).

For both DCM analyses, we modelled full reciprocal intrinsic connections as well as (inhibitory) self-connections. The driving input was set to either one region at a time, or all possible combinations between regions (resulting in 7 possible combinations). We allowed for predictability-dependent modulation of connections in the following way: either one connection was modulated separately or two afferent or efferent connections were modulated by predictability (12 possible combinations; see Supplementary Figure 3 for a schematic overview of the model space). Since family-level inference has been found especially useful in case of large model spaces (Penny et al. 2010), the resulting model space of 7 x 12 models was partitioned into 12 families, grouping models according to the modulatory input (i.e. within a family, the modulating connection was kept constant while the driving input was varied). We used the same model architecture for both DCM analyses.

All 84 models per participant were inverted using a Variational Bayes approach to estimate parameters that provide the best trade-off between accuracy and complexity quantified by free energy (Friston et al. 2006; Friston et al. 2007). At the second level, a random-effects Bayesian Model Selection procedure was applied to identify the most probable family of models given the fMRI data quantified by the respective exceedance probability (Stephan et al. 2009; Penny et al. 2010). Single-subject parameter estimates were averaged across all models within the winning family using Bayesian Model Averaging and submitted to a one-sample *t*-test. The strength of the effects was quantified using the Bayes Factor. For *t*-tests and Bayes Factor we used JASP (version 0.9.1).

Finally, we investigated the between-subject relationship of behavioural comprehension and functional connectivity in the brain. For single participants, we summarized behavioural performance in sentences with low and high predictability as the area under the psychometric curve spanning all intelligibility levels included in the DCM analysis. The coupling strength of connections significantly modulated by predictability was correlated with behavioural performance in the according predictability condition.

1 Results

2 Semantic predictability benefits speech comprehension at intermediate levels of

3 intelligibility

On average, participants repeated 67.4% of keywords correctly (SD = 10.52). Psychometric

curves fitted to the proportion of correctly repeated keywords across intelligibility levels shifted

leftwards for sentences with high predictability when compared to low predictability (Figure

2A). As reflected by a decrease of the threshold for sentences with high predictability relative

to low predictability (t_{25} = -9.87, p < 0.001, r = 0.57, BF_{10} > 1,000; Figure 2B), high semantic

predictability had a facilitatory effect on speech comprehension at intermediate intelligibility

levels. Moreover, the lapse rate was smaller for sentences with high compared to low

predictability (Z = -2.78, p = 0.005, r = 0.55), indicating a predictability gain that persisted (to a

smaller degree) beyond intermediate intelligibility in the highest intelligibility level.

There was no significant difference between sentences with low and high predictability for

slope (t_{25} = 1.6, p = 0.122, r = -0.11, BF_{10} = 0.64) and width (t_{25} = 1.84, p = 0.077, r = 0.11,

BF_{10} = 0.896) or guess rate (Z = 0.27, p = 0.79, r = 0.05) of psychometric curves.

Figure 2. Behavioural results of the sentence repetition task. (A) Psychometric functions
(coloured lines) were fitted to the proportion of correctly repeated keywords (coloured dots)
across intelligibility levels for sentences with low (orange) and high predictability (green); grand
average is displayed for illustration purposes. **(B)** The threshold of sentences with low
predictability shifted towards more intelligible levels when compared to sentences with high
predictability ($p < 0.001$; top). There was no evidence for a slope difference between sentences
with low and high predictability ($p = 0.122$; bottom). Density plots illustrate the difference
between the proportion correct of sentences with high vs. low predictability.

A fronto-temporo-parietal network is tuned towards intelligible speech

First, we were interested in brain regions showing stronger activity for increasing intelligibility

of speech stimuli, irrespective of sentence predictability. The parametric variation of

intelligibility showed increased activation in a set of regions comprising anterior-to-posterior

bilateral superior temporal gyri (STG), left precentral and postcentral gyrus and left inferior

frontal gyrus (IFG) as well as a large left-hemispheric temporo-parietal cluster and left

precuneus (Figure 3). These regions have previously been reported to support auditory speech

comprehension and suggested to be tuned to perceptual clarity (e.g. Rauschecker & Scott,

2009).

Figure 3. Main effect of increasing speech intelligibility across the whole cortex. Top:
Activation map thresholded at p < 0.05 (FWE-corrected). STG = superior temporal gyrus, IFG
= inferior frontal gyrus, AG = angular gyrus, SFG = superior frontal gyrus, prec = precuneus.
Bottom: Parameter estimates (percent signal change) for each experimental condition at peak
voxels of left AG (MNI: x = -47, y = -67, z = 25), left IFG (MNI: x = -54, y = 33, z = 8) and right
STG (MNI: x = 53, y = -12, z = 2). Error bars represent the standard error of the mean (SEM).

Increasing intelligibility of high predictable sentences enhances recruitment of left-hemispheric temporo-parietal regions

Further, we were interested in brain regions that were differentially affected by predictability at different levels of intelligibility. Higher activation for sentences with high compared to low predictability under linearly increasing intelligibility was found in left-hemispheric regions encompassing pMTG, middle to posterior cingulate cortex, two separate clusters in left AG (PGa and PGp), inferior temporal gyrus, precuneus and supramarginal gyrus (SMG; Figure 4, Table 1). Similar but weaker activation patterns were found for the quadratically modulated interaction contrast. Interestingly, the two separate left-hemispheric AG clusters fell into different cytoarchitectonic subregions: the larger cluster was mainly located in the anterior dorsal portion of AG (PGa & PFm) at the boundary to SMG and the smaller cluster was mainly located in the posterior portion of AG (PGp) in close proximity to the middle occipital gyrus (MOG). The AG clusters and the pMTG cluster showed a pattern of deactivation relative to the fixation baseline with relatively less deactivation for high predictable as compared to low predictable sentences (Figure 4B). As both, the AG and pMTG are considered key regions of semantic processing, their time series were submitted to later connectivity analyses.

Increasing intelligibility of low predictable sentences enhances recruitment of domain-general regions

The opposite interaction contrast revealed greater activity for low compared to high predictable sentences under linearly increasing intelligibility and encompassed pre-SMA/paracingulate gyrus, left-hemispheric IFG (pars triangularis; BA 45) and the bilateral anterior insula (see Figure 4). These regions overlap with the cingulo-opercular network (Dosenbach et al., 2008), which is frequently reported in conditions associated with high task demands. To investigate context-dependent changes within this network, the pre-SMA/paracingulate gyrus and bilateral insulae were analysed in a later DCM analysis. Note that the interaction contrasts modelling intelligibility quadratically (i.e. inverted u-shape with the highest activation at intermediate

levels of intelligibility) resulted in a similar pattern of brain regions and therefore are not shown here.

Table 1. Results from the context-dependent interaction.

Location	Hemisphere	MNI			*t*	Cluster Size
		x	y	z		
High > Low predictable sentences						
Middle cingulate cortex	R	3	−34	38	7.49	159
Middle cingulate cortex	L	−4	−30	42	7.17	subcluster
Middle temporal gyrus	**L**	**−60**	**−52**	**−5**	**6.85**	**150**
Middle temporal gyrus	L	−54	−62	2	6.6	subcluster
Angular gyrus (PFm/PGa)	**L**	**−42**	**−62**	**50**	**6.47**	**135**
Angular gyrus (PGa)	L	−42	−70	42	6.31	subcluster
Angular gyrus (PGp)	L	−34	−77	42	6.02	subcluster
Angular Gyrus (PGp)/ middle occipital gyrus	**L**	**−34**	**−77**	**30**	**6.34**	**51**
Angular gyrus (PGp)	L	−42	−77	28	5.63	subcluster
Supramarginal gyrus	L	−60	−30	30	5.91	26
Inferior temporal gyrus	L	−47	−52	−25	5.82	21
Angular gyrus (PGp)	R	46	−67	42	5.71	8
Precuneus	L	−10	−47	55	5.62	15
Low > High predictable sentences						
Presupplementary motor area/paracingulate gyrus	**R/L**	**0**	**16**	**50**	**7.72**	**138**
Inferior frontal gyrus (p. orbitalis, area 45)	L	−42	26	−10	6.47	106
Insular cortex	**L**	**−30**	**26**	**−2**	**6.43**	**subcluster**
Inferior frontal gyrus (p. triangularis, area 45)	L	−50	23	5	5.9	subcluster
Inferior frontal gyrus (p. triangularis, area 45)	L	−44	20	0	5.88	subcluster
Insular cortex	**R**	**30**	**26**	**−2**	**6.13**	**24**
Inferior frontal gyrus (p. triangularis, BA 45)	L	−57	20	15	5.71	16
Inferior frontal gyrus (p. triangularis, BA 45)	L	−50	20	18	5.07	subcluster

High > low predictable sentences: Regions showing greater activity for high as compared to low predictable sentences under increasing intelligibility; Low > high predictable sentences: Regions showing greater activity for low as compared to high predictable sentences under increasing intelligibility. Regions are reported at a peak-level FWE-correcting threshold at an alpha of .05. Seed regions for DCM analyses are highlighted in bold.

Figure 4. Group-level fMRI results illustrating brain regions that are sensitive to the interaction of intelligibility and predictability. (A) Parameter estimates illustrate the interaction effect, extracted from peak voxels of DCM seed regions for the semantic network. **(B)** Brain regions showing a significant interaction; IFG = inferior frontal gyrus, SMG = supramarginal gyrus, pMTG = posterior middle temporal gyrus, PGa = anterior angular gyrus, PGp = posterior angular gyrus, MCC/PCC = middle and posterior cingulate cortex, pre-SMA = pre-supplementary motor area, aIns = anterior insula; green = high > low predictable sentences, orange = low > high predictable sentences. The activation map is thresholded at $p_{FWE-corrected} < 0.05$. **(C)** Parameter estimates illustrate the interaction effect, extracted from peak voxels of DCM seed regions for the cingulo-opercular (CO) network. Error bars represent the standard error of the mean (SEM).

The extended semantic system scales to the behavioural predictability gain dependent on speech intelligibility

The proportion of individual task performance explained by the interaction of predictability and intelligibility in each experimental condition correlated positively with BOLD activity in a broad semantic network (Figure 5; Binder et al., 2009). At intermediate levels of intelligibility, neural activity increased for sentences with high predictability in those participants exploiting a stronger behavioural predictability gain.

In line with the linear interaction effect (see Table 1, high > low predictability), modelling

the behavioural interaction effect revealed significant clusters in parietal and lateral temporal

cortex. A broad left-hemispheric and a smaller right-hemispheric cluster were observed in PGp

(Table 2). Additional clusters were found in bilateral pMTG.

Figure 5. Projection of activity onto the behavioural predictability gain across levels of
intelligibility. (A) The condition-specific proportion of behavioural performance explained by
the interaction of intelligibility and predictability in each participant (middle line plot) was
correlated with individual parameter estimates of single-voxel BOLD activity (left and right bar
plots; exemplary grand average parameter estimates for two peak voxels). Thin lines in the
line plot represent single participants; fat lines represent grand average. Error bars represent
±1 SEM. **(B)** Group statistics revealed brain areas modulated by single-participant interaction
effect of intelligibility and predictability on speech comprehension. FP = frontal pole; PGp =
posterior angular gyrus; pMTG = posterior middle temporal gyrus; PHG = parahippocampal
gyrus; SCC = subcallosal cortex; SFG = superior frontal gyrus; PCG = paracingulate gyrus (for
an exhaustive list of clusters see Table 2).

More importantly, the behavioural interaction effect mapped onto frontal and ventromedial

brain regions not implicated in the linear interaction effect. The frontal pole encompassed one

broad medial cluster spanning from the left to the right hemisphere as well as a right-

hemispheric cluster at the border with IFG and a left-hemispheric cluster extending into the

superior frontal gyrus (Table 2). Near the inferior-posterior frontal pole, we observed bilateral clusters in IFG. The limbic lobes contained a bilateral cluster in the posterior subcallosal cortex that was complemented by a bilateral cluster in inferior paracingulate gyrus extending into anterior subcallosal cortex as well as cingulate gyrus and frontal medial cortex. Additional clusters comprised the anterior division of bilateral superior parahippocampal gyrus and left medial-posterior insular cortex. In the medial temporal lobe, we found a left-hemispheric cluster in the posterior division of the temporal fusiform gyrus. No significant cluster in the brain showed a negative correlation with the interaction effect on task performance.

Table 2. Brain areas sensitive to the interaction effect of intelligibility and predictability on task performance.

Location	Hemisphere	MNI			t	Cluster Size
		x	y	z		
Brain Behaviour Correlation						
Frontal pole	L	–5	56	5	3.53	161
	R	48	38	3	3.96	103
	L	–12	56	35	3.71	26
Angular gyrus (PGp)	L	–50	–70	33	3.85	100
	R	46	–67	25	3.45	55
Subcallosal cortex	L	–2	6	–13	4.22	78
Middle temporal gyrus (temporo-occipital)	L	–60	–55	–5	3.65	80
	R	58	–45	–5	4.04	25
	L	–57	–65	10	2.76	2
Parahippocampal gyrus (anterior)	R	21	1	–18	3.74	55
	L	–25	–2	–20	4.31	51
Paracingulate gyrus	R	1	41	-8	3.96	54
Insular cortex	L	–33	–17	5	3.74	17
Temporal fusiform gyrus (posterior)	L	–42	–30	–23	3.53	5
Inferior frontal gyrus (p. orbitalis)	R	28	31	–15	3.94	13
	L	–37	36	–10	3.71	3

Each cluster ($p_{FDR} < 0.05$) is described by hemisphere, MNI coordinates and t-value of its peak voxel as well as the number of voxels it contains.

Semantic predictability increases connectivity between subregions of the angular gyrus

Next, to investigate context-dependent changes in effective connectivity between regions identified in the mass univariate GLM analysis, we performed two separate DCM analyses, one within the semantic system and one within the cingulo-opercular network.

The family of models with a context-dependent modulation of the connection from PGp to pMTG and PGa was identified as the winning family by means of Bayesian model selection (exceedance probability: 0.62; see Supplementary Figure 4A for an overview of all exceedance probabilities). Note that the second-best family exhibited an exceedance probability of 0.09. The intrinsic connection from PGa to pMTG was negative. The remaining intrinsic connections were positive (see Supplementary Table 1), although only the outgoing connections from pMTG reached statistical significance. Within the winning family, the only significant modulatory effect was the effect of high predictable sentences on the connection from PGp to PGa ($t_{25} = 2,53$, $p = 0.02$, $d = 0.5$, $BF_{10} = 0.51$). This modulatory effect was positive ($M = 0.38$, $SD = 0.77$), indicating an excitatory influence of PGp on PGa for high predictable sentences. None of the connections was significantly modulated by low predictability, suggesting that connectivity within this network is modulated by semantic context.

Within the semantic model, we found no significant correlations between individual task performance and predictability-induced changes in connectivity strength from PGp to PGa (high predictability: $r = -0.07$, $p = 0.742$, $BF_{10} = 0.26$).

Semantic predictability decreases connectivity within the cingulo-opercular network

For the cingulo-opercular network model, the family of models with a modulatory effect on the connections from pre-SMA to bilateral insulae was identified as the winning family, with an exceedance probability of 0.35 (see Supplementary Figure 4B). Note that the family with modulations of the connections from pre-SMA to left insula as well as from right insula to left insula was identified as second probable family of models with an exceedance probability of 0.29.

In the winning model, both connections from pre-SMA to insula were significantly

modulated by predictability (see Figure 6 and Supplementary Table 2). Low predictable

sentences had on average a facilitatory effect on the connection strength, indicating increased

coupling when predictability was low. Highly predictable sentences further decreased the

weakly negative intrinsic connection, indicating a decrease in coupling when predictability was

high.

When listening to sentences with high predictability, participants with stronger inhibitory

influence from pre-SMA to right insula showed increased behavioural performance (r = -0.65,

p < 0.001, BF_{10} = 109.79; Figure 6C). No significant correlations were observed between

individual task performance and predictability-induced changes in connectivity strength from

pre-SMA to left insula (high predictability: r = -0.09, p = 0.645, BF_{10} = 0.27; low predictability: r

= -0.22, p = 0.279, BF_{10} = 0.42).

Figure 6. DCM results. Effective connectivity averaged across models within the winning
family for **(A)** the semantic network and **(B)** the cingulo-opercular network. Grey arrows with
parameter estimates indicate significant intrinsic connections. Black arrows indicate intrinsic
connections modulated by high (green) and low (orange) predictability. **(C)** Between-subject
correlation of behavioural performance (area under the psychometric curve; see illustration on
the left) and modulatory connectivity strength in three cingulo-opercular connections
significantly modulated by predictability; n.s. = not significant, * $p < 0.05$, ** $p < 0.01$, *** $p <$
0.001. PGa = anterior angular gyrus; PGp = posterior angular gyrus; pMTG = posterior middle
temporal gyrus; pre-SMA = pre-supplementary motor area; l INS = left anterior insula; r INS =
right anterior insula.

Discussion

In the current fMRI study, we investigated the neural underpinnings of the semantic predictability gain during speech processing in noisy listening situations at the network level. First, we showed that highly predictable but acoustically degraded speech engages left-hemispheric semantic regions and modulates the individual comprehension gain in behaviour via additional frontal and medial semantic regions. Second, we found that highly predictable speech strengthens effective connectivity between two subregions of a key semantic network node in the left angular gyrus. Third, we showed that highly predictable speech decreases connectivity from pre-SMA to bilateral insulae within the cingulo-opercular network. Notably, stronger decoupling of right insula predicted the individual comprehension gain. These findings suggest that semantic predictability facilitates speech comprehension in challenging listening conditions via upregulation of the semantic network and active inhibition of domain-general regions.

Our sentence repetition task had been designed to overcome the experimental shortcomings of previous studies and investigate speech comprehension in noise on the basis of rich neural and behavioural data. Hitherto, fMRI studies on the neural underpinnings of degraded speech processing had mainly used sparse temporal sampling to avoid interference of scanner noise with experimentally manipulated signal-to-noise ratio of speech stimuli (Obleser and Kotz 2010; Erb et al. 2013; Golestani et al. 2013). As we aimed to investigate speech processing directly during (compared to after) listening and to improve sampling density for our connectivity analyses, BOLD activity in the present study was continuously recorded throughout the whole experiment. In line with studies using temporal sparse sampling (Davis and Johnsrude 2003; Rauschecker and Scott 2009; Abrams et al. 2013), we found activity in regions associated with auditory comprehensibility (e.g., bilateral STG as well as left IFG) to be stronger with increasing intelligibility of speech. This replication of the intelligibility effect speaks for continuous scanning as a feasible technique to investigate *online* speech processing.

Notably, in contrast to previous studies with only few and fixed experimental levels of

speech degradation, we covered the full range of intelligibility from hardly to easily

comprehensible speech in the individual participant. At the behavioural level, we were thus

able to fit full psychometric curves (i.e., speech comprehension as a function of intelligibility,

see Figure 2). We found the psychometric curve to shift towards less intelligible levels for

sentences with high compared to low semantic predictability. This threshold shift is an

established effect in more challenging comprehension tasks (Shannon et al. 2004) and reflects

a comprehension benefit for predictable speech at intermediate intelligibility. As left AG has

been found to show a strong differentiation between sentences with high and low predictability

at intermediate intelligibility (Obleser et al. 2007; Obleser and Kotz 2010) and to exhibit

functional relevance in mediating the comprehension gain (Hartwigsen et al. 2015), this region

has been proposed to serve top-down activation of semantic concepts that facilitate speech

comprehension (Obleser and Kotz 2010).

However, as an unexpected finding, two separate AG clusters showed stronger activation

for sentences with high compared to low predictability under linearly increasing intelligibility: a

larger cluster mainly falling into PGa at the boundary to the supramarginal gyrus, and a smaller

cluster at the boundary between PGp and middle occipital gyrus. Importantly, these AG

subregions have been discussed to subserve different functions (Seghier et al. 2010; Noonan

et al. 2013; Bonnici et al. 2016). PGa is implicated in domain-general attention and converges

with coordinates commonly reported as AG activation during speech in noise processing

(Obleser et al. 2007; Obleser and Kotz 2010; Clos et al. 2014; Guediche et al. 2014). In

contrast, PGp activation has often been labelled as MOG (McGettigan et al. 2012; Guediche

et al. 2016). In these studies, MOG was interpreted to provide free resources otherwise

recruited by visual processing to the auditory domain during resource-demanding speech in

noise comprehension. Other studies have associated PGp activation with "pure" semantic

processing (Seghier et al., 2010; Bonnici et al., 2016). Our findings implicate that both AG

subregions contribute to successful speech comprehension under challenging listening

conditions when semantic cues are available.

To further characterize the functional contributions of anterior and posterior AG, we

analysed the connectivity between these subregions and observed that coupling from posterior

to anterior AG was strengthened by high but not low predictability. This finding suggests that

an interplay of AG subregions enables integration of semantic information. As a semantic key

region, left PGp might play an important role in integrating activated semantic information.

Depending on the availability of the semantic context, PGp might recruit additional attentional

resources from PGa to shift attention to semantic constraints. Aside from AG, we largely found

the same regions (e.g., left pMTG, left SMG, left inferior temporal gyrus, posterior cingulate

gyrus and precuneus) that were previously implicated in processing of sentences with high

predictability under increasing intelligibility (Obleser et al. 2007; Obleser and Kotz 2010;

Golestani et al. 2013). These regions overlap with the previously described core semantic

(control) regions (Binder et al. 2009, Jefferies 2013) and form a functional network during

degraded speech processing (Obleser et al. 2007).

To complement the linear interaction contrast, we also modelled task performance in the

brain and expected to find the left-hemispheric semantic network strengthened by this contrast

fine-tuned to inter-subject variation. Indeed, the semantic core regions extended to their

homologous regions in the right hemisphere. Strikingly, additional semantic regions scaled in

activation with the behavioural predictability gain at intermediate intelligibility. These regions

largely pertained to two subsets of the semantic network: *lateral and ventral temporal cortex*

includes left temporal fusiform gyrus and parahippocampal gyrus, whereas *left ventromedial*

prefrontal cortex includes subcallosal cortex and frontal pole (Binder et al. 2009). The temporal

cortex has been shown to communicate with the hippocampus via a connection from temporal

fusiform gyrus to parahippocampal gyrus evident in structural (Powell et al. 2004) and

functional imaging (Teipel et al. 2010). It has been suggested that this pathway is used to

encode semantic information (Levy et al. 2004; Martin 2007). Critically, ventral temporal

activation has been implicated in the processing of semantically unambiguous speech in a

memory-demanding word recognition task (Rodd et al. 2005) and processing of visually

presented words that were correctly repeated during later recall (Strange et al. 2002). In the

sentence repetition task employed here, memory formation is crucial to correctly repeat the

sentence after listening. The association of more accurate recall with stronger ventral temporal

activation for predictable but degraded sentences suggests that memory engagement is most

crucial for successful speech comprehension when semantic information is available for

integration and must be protected against competing but irrelevant information (i.e.,

background noise). In line with this notion, prefrontal cortex regions implicated in executive

control functioning (Duncan and Owen 2000) show increased activation when efficient

management of cognitive resources promises the strongest gain in performance. While we will

refrain from overinterpreting these exploratory results, it is worth highlighting that the role of

executive control and memory formation might have been often overlooked in previous studies

on speech comprehension despite their face validity when it comes to understanding individual

hearing difficulties.

As a last key finding of our study, challenging speech comprehension in the absence of

semantic constraints is accompanied by an increase of activity in regions within the cingulo-

opercular network. This finding is broadly in line with previous reports on activation of cingulo-

opercular regions during effortful speech processing (Eckert et al. 2009; Hervais-Adelman et

al. 2012; Erb and Obleser 2013; Vaden et al. 2013; Clos et al. 2014). The cingulo-opercular

network has been associated with domain-general control or executive processes such as task

monitoring (e.g. Dosenbach et al., 2008; Duncan, 2010; Vaden et al., 2013, 2015), executive

control (Erb et al. 2013) and attentional control (Fitzhugh et al. 2019). Specifically, in the

speech in noise literature, the increased reliance on this network in difficult listening conditions

has been interpreted as an adaptive control mechanism to enable speech comprehension

when little or no semantic information is available. This is what we show in our study.

Within these cingulo-opercular regions, we found an inverted u-shaped response, with

highest activation at intermediate levels of intelligibility and less activation for the conditions

that were either hardly or easily comprehensible irrespective of predictability. This activation

pattern suggests that domain-general regions contribute to effortful speech processing as long

as *any* information can be extracted. Analogously, recruitment of these regions is less

important when the auditory signal is too bad or when speech is clear. Further, we observed that connectivity between these regions was differentially modulated by high and low predictability. Whereas within-network connectivity was facilitatory under low semantic predictability, highly predictable speech covaried with an inhibition in the functional interaction between these regions. These findings suggest that domain-general cognitive control regions may become actively suppressed when language-specific cues are available. An alternative interpretation outside the scope of the present methodology might be that acoustic degradation leads to an activation of this network without the assessment of alternative facilitatory cues such as semantic context, which is only taken into account in a subsequent stage.

Interestingly, the individual degree of inhibitory influence from pre-SMA to the right insula was associated with individual task performance. Those subjects who had higher comprehension scores, showed the strongest inhibitory modulation of this connection. This result converges with and extends previous task-based fMRI studies that found increased activity in the right insula when auditory input was degraded (Erb et al. 2013). While older adults with well-preserved hearing abilities and younger participants recruited the right insula under challenging listening conditions, recruitment of this region was already observed in clear speech conditions for older adults with hearing loss (Erb and Obleser 2013).

Our results are further in line with a recent study that reported a decrease in the functional connectivity between left IFG and right insula under facilitatory neurostimulation that was associated with improved word learning (Fiori et al. 2018). Together, the previous and present findings suggest that recruitment of the right insula is beneficial in challenging conditions (e.g. in the beginning of a learning phase or when auditory input is deteriorated) but becomes unnecessary as soon as other factors can be exploited (such as semantic context). Moreover, the present findings suggest that better behavioural performance may require an active inhibition of this region to ensure efficient processing.

Recently, the pre-SMA has been suggested to be relevant during high task demands (Hertrich et al. 2016) and to play a role in the wider semantic control network (Hallam et al. 2016). Specifically, Hallam and colleagues (2016) found increased activity in pre-SMA and

pMTG after inhibitory TMS to left IFG, which was interpreted as upregulation of the relative contribution to semantic control in the presence of a TMS-induced disruption. Further, Dietrich and colleagues (2018) reported decreased performance in a sentence repetition task after inhibitory TMS over the pre-SMA, which was selectively evident for the most challenging task condition. Note that pre-SMA activation has also been reported in sentence comprehension tasks in the degraded speech literature before (e.g. Clos et al., 2014: sentence matching task). Therefore, it is unlikely that our pre-SMA finding mainly reflects the choice of the task alone. Taken together, our results replicate the previously reported involvement of domain-general regions during effortful speech processing and extend these findings by demonstrating predictability-dependent modulation of functional connectivity within this network.

However, it is important to bear in mind that the effective connectivity results are restricted to the limited set of regions included in the network models. Furthermore, the current study investigated modulatory effects of semantic predictability on effective connectivity across intermediate to good intelligibility within the respective network and therefore did not operationalize the interactive effect of intelligibility and predictability per se. Future studies are needed to directly address the interactive effect and additionally investigate the dynamic interplay *between* the semantic and cingulo-opercular network.

Conclusion

Our results demonstrate that predictability in challenging listening conditions not only modulates the semantic network but critically extends to modulations in the domain-general network in a behaviourally relevant fashion. The inhibitory modulation of connectivity within the domain-general network was associated with increased behavioural speech comprehension. Further, we demonstrated that predictability modulated connectivity between two subregions of left AG. This underscores the functional heterogeneity of the AG region and emphasizes the importance of reporting findings within this region with greater regional specificity.

Acknowledgements

The authors would like to thank Manuela Hofmann, Sylvie Neubert and Simone Wipper for support with the data acquisition and Dagny Kühner, Isabel Gebhardt, Anna Ruhe and Ricarda Vielhauer for support with recording transcription. We further thank the University of Minnesota Center for Magnetic Resonance Imaging for providing the fMRI multiband sequence software.

References

Abrams DA, Ryali S, Chen T, Balaban E, Levitin DJ, Menon V. 2013. Multivariate activation and connectivity patterns discriminate speech intelligibility in Wernicke's, Broca's, and Geschwind's areas. Cereb Cortex. 23(7):1703–1714. doi:10.1093/cercor/bhs165.

Adank P. 2012. The neural bases of difficult speech comprehension and speech production: Two Activation Likelihood Estimation (ALE) meta-analyses. Brain Lang. 122(1):42–54. doi:10.1016/j.bandl.2012.04.014.

Alain C, Du Y, Bernstein LJ, Barten T, Banai K. 2018. Listening under difficult conditions: An activation likelihood estimation meta-analysis. Hum Brain Mapp. 39(7):2695–2709.

Binder JR, Desai RH, Graves WW, Conant LL. 2009. Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. Cereb Cortex. 19(12):2767–2796. doi:10.1093/cercor/bhp055.

Bonnici HM, Richter FR, Yazar Y, Simons JS. 2016. Multimodal feature integration in the angular gyrus during episodic and semantic retrieval. J Neurosci. 36(20):5462–5471. doi:10.1523/JNEUROSCI.4310-15.2016.

Bzdok D, Hartwigsen G, Reid A, Laird AR, Fox PT, Eickhoff SB. 2016. Left inferior parietal lobe engagement in social cognition and language. Neurosci Biobehav Rev. 68:319–334. doi:10.1016/j.neubiorev.2016.02.024.

Caspers S, Eickhoff SB, Geyer S, Scheperjans F, Mohlberg H, Zilles K, Amunts K. 2008. The human inferior parietal lobule in stereotaxic space. Brain Struct Funct. 212(6):481–495. doi:10.1007/s00429-008-0195-z.

Caspers S, Geyer S, Schleicher A, Mohlberg H, Amunts K, Zilles K. 2006. The human inferior parietal cortex: Cytoarchitectonic parcellation and interindividual variability. NeuroImage. 33(2):430–448. doi:10.1016/j.neuroimage.2006.06.054.

Clos M, Langner R, Meyer M, Oechslin MS, Zilles K, Eickhoff SB. 2014. Effects of prior information on decoding degraded speech: An fMRI study. Hum Brain Mapp. 35(1):61–74. doi:10.1002/hbm.22151.

Cohen J. 1960. A Coefficient of Agreement for Nominal Scales. Educ Psychol Meas. 10(1):37–46.

Davis MH, Johnsrude IS. 2003. Hierarchical Processing in Spoken Language Comprehension. 23(8):3423–3431.

Dietrich S, Hertrich I, Müller-Dahlhaus F, Ackermann H, Belardinelli P, Desideri D, Seibold VC, Ziemann U. 2018. Reduced performance during a sentence repetition task by continuous theta-burst magnetic stimulation of the pre-supplementary motor area. Front Neurosci. 12:1–13. doi:10.3389/fnins.2018.00361.

Dosenbach NUF, Fair DA, Cohen AL, Schlaggar BL, Petersen SE. 2008. A dual-networks architecture of top-down control. (February):99–105. doi:10.1016/j.tics.2008.01.001.

Duncan J. 2010. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. Trends Cogn Sci. 14(4):172–179. doi:10.1016/j.tics.2010.01.004.

Duncan J, Owen AM. 2000. Common regions of the human frontal lobe recruited by diverse cognitive demands. Trends Neurosci. 23(10):475–483.

Eckert MA, Menon V, Walczak A, Ahlstrom J, Denslow S, Horwitz A, Dubno JR. 2009. At the
heart of the ventral attention system: The right anterior insula. Hum Brain Mapp.
30(8):2530–2541. doi:10.1002/hbm.20688.

Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, Zilles K. 2005. A new
SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging
data. NeuroImage. 25(4):1325–1335. doi:10.1016/j.neuroimage.2004.12.034.

Erb J, Henry MJ, Eisner F, Obleser J. 2012. Auditory skills and brain morphology predict
individual differences in adaptation to degraded speech. Neuropsychologia. 50(9):2154–
2164. doi:10.1016/j.neuropsychologia.2012.05.013.

Erb J, Henry MJ, Eisner F, Obleser J. 2013. The Brain Dynamics of Rapid Perceptual
Adaptation to Adverse Listening Conditions. J Neurosci. 33(26):10688–10697.
doi:10.1523/JNEUROSCI.4596-12.2013.

Erb J, Obleser J. 2013. Upregulation of cognitive control networks in older adults' speech
comprehension. Front Syst Neurosci. 7:1–13. doi:10.3389/fnsys.2013.00116.

Fiori V, Kunz L, Kuhnke P, Marangolo P, Hartwigsen G. 2018. Transcranial direct current
stimulation (tDCS) facilitates verb learning by altering effective connectivity in the healthy
brain. NeuroImage. 181:550–559. doi:10.1016/j.neuroimage.2018.07.040.

Fitzhugh MC, Hemesath A, Schaefer SY, Baxter LC. 2019. Functional Connectivity of Heschl
's Gyrus Associated With Age-Related Hearing Loss : A Resting-State fMRI Study. Front
Psychol. 10(November). doi:10.3389/fpsyg.2019.02485.

Friston KJ, Harrison L, Penny W. 2003. Dynamic causal modelling. NeuroImage.
19(4):1273–1302. doi:10.1016/S1053-8119(03)00202-7.

Friston KJ, Kilner J, Harrison L. 2006. A free energy principle for the brain. J Physiol-Paris.
100(1–3):70–87. doi:10.1016/j.jphysparis.2006.10.001.

Friston KJ, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W. 2007. Variational free
energy and the Laplace approximation. NeuroImage. 34(1):220–234.
doi:10.1016/j.neuroimage.2006.08.035.

Friston KJ, Worsley KJ, Frackowiak RSJ, Mazziotta JC, Evans AC. 1994. Assessing the
significance of focal activations using their spatial extent. Hum Brain Mapp. 1(3):210–
220. doi:10.1002/hbm.460010306.

Fründ I, Haenel NV, Wichmann FA. 2011. Inference for psychometric functions in the
presence of nonstationary behavior. J Vis. 11(6):1–19. doi:10.1167/11.6.16.

Gläscher J. 2009. Visualization of group inference data in functional neuroimaging.
Neuroinformatics. 7(1):73–82. doi:10.1007/s12021-008-9042-x.

Golestani N, Hervais-Adelman A, Obleser J, Scott SK. 2013. Semantic versus perceptual
interactions in neural processing of speech-in-noise. NeuroImage. 79:52–61.
doi:10.1016/j.neuroimage.2013.04.049.

Guediche S, Blumstein SE, Fiez JA, Holt LL. 2014. Speech perception under adverse
conditions: insights from behavioral, computational, and neuroscience research. Front
Syst Neurosci. 7:1–16. doi:10.3389/fnsys.2013.00126.

Guediche S, Reilly M, Santiago C, Laurent P, Blumstein SE. 2016. An fMRI study
investigating effects of conceptually related sentences on the perception of degraded
speech. Cortex. 79:57–74. doi:10.1016/j.cortex.2016.03.014.

Halai AD, Welbourne SR, Embleton K, Parkes LM. 2014. A comparison of dual gradient-
echo and spin-echo fMRI of the inferior temporal lobe. Hum Brain Mapp. 35(8):4118–
4128. doi:10.1002/hbm.22463.

Hallam GP, Whitney C, Hymers M, Gouws AD, Jefferies E. 2016. Charting the effects of
TMS with fMRI: Modulation of cortical recruitment within the distributed network
supporting semantic control. Neuropsychologia. 93:40–52.
doi:10.1016/j.neuropsychologia.2016.09.012.

Hartwigsen G, Golombek T, Obleser J. 2015. Repetitive transcranial magnetic stimulation
over left angular gyrus modulates the predictability gain in degraded speech
comprehension. Cortex. 68:100–110. doi:10.1016/j.cortex.2014.08.027.

Henson R. 2015. Analysis of variance (ANOVA). Brain Mapp Encycl Ref Elsevier.:477–481.

Hertrich I, Dietrich S, Ackermann H. 2016. The role of the supplementary motor area for
speech and language processing. Neurosci Biobehav Rev. 68:602–610.
doi:10.1016/j.neubiorev.2016.06.030.

Hervais-Adelman AG, Carlyon RP, Johnsrude IS, Davis MH. 2012. Brain regions recruited
for the effortful comprehension of noise-vocoded words. Lang Cogn Process. 27(7–
8):1145–1166. doi:10.1080/01690965.2012.662280.

Holmes E, Folkeard P, Johnsrude IS, Scollie S. 2018. Semantic context improves speech
intelligibility and reduces listening effort for listeners with hearing impairment. Int J
Audiol. 57:483–492. doi:10.1097/PSY.0000000000000183.

Jefferies E. 2013. The neural basis of semantic cognition: Converging evidence from
neurophysiology, neuroimaging and TMS. Cortex. 49:611-625.
doi:10.1016/j.cortex.2012.10.008

Jeffreys H. 1961. The Theory of Probability. 3rd Edition. Oxford: Oxford University Press.

Jung JY, Cloutman LL, Binney RJ, Lambon Ralph MA. 2017. The structural connectivity of
higher order association cortices reflects human functional brain networks. Cortex.
97:221–239. doi:10.1016/j.cortex.2016.08.011.

Kalikow DN, Stevens KN, Elliott LL. 1977. Development of a test of speech intelligibility in
noise using sentence materials with controlled word predictability. J Acoust Soc Am.
61(5):1337–1351. doi:10.1121/1.381436.

Kass RE, Raftery AE. 1995. Bayes Factors. J Am Stat Assoc. 90(430):773–795.
doi:10.1080/01621459.1995.10476572.

Kollmeier B, Gilkey RH, Sieben UK. 1988. Adaptive staircase techniques in psychoacoustics:
A comparison of human data and a mathematical model. J Acoust Soc Am. 83(5):1852–
1862. doi:10.1121/1.396521.

Levy DA, Bayley PJ, Squire LR. 2004. The anatomy of semantic knowledge: Medial vs.
lateral temporal lobe. Proc Natl Acad Sci U S A. 101(17):6710–6715.
doi:10.1073/pnas.0401679101.

Lohmann G, Stelzer J, Lacosse E, Kumar VJ, Mueller K, Kuehn E, Grodd W, Scheffler K. 2018. LISA improves statistical analysis for fMRI. Nat Commun. 9(1):1–9. doi:10.1038/s41467-018-06304-z.

Marreiros AC, Kiebel SJ, Friston KJ. 2008. Dynamic causal modelling for fMRI: A two-state model. NeuroImage. 39(1):269–278. doi:10.1016/j.neuroimage.2007.08.019.

Martin A. 2007. The Representation of Object Concepts in the Brain. Annu Rev Psychol. 58(1):25–45. doi:10.1146/annurev.psych.57.102904.190143.

McGettigan C, Faulkner A, Altarelli I, Obleser J, Baverstock H, Scott SK. 2012. Speech comprehension aided by multiple modalities: Behavioural and neural interactions. Neuropsychologia. 50(5):762–776. doi:10.1016/j.neuropsychologia.2012.01.010.

Nilsson M, Soli SD, Sullivan JA. 1994. Development of the Hearing in Noise Test for the measurement of speech reception thresholds in quiet and in noise. J Acoust Soc Am. 95(2):1085–1099.

Noonan KA, Jefferies E, Visser M, Lambon Ralph MA. 2013. Going beyond inferior prefrontal involvement in semantic control: evidence for the additional contribution of dorsal angular gyrus and posterior middle temporal cortex. J Cogn Neurosci. 25(11):1824–1850.

Obleser J, Kotz SA. 2010. Expectancy constraints in degraded speech modulate the language comprehension network. Cereb Cortex. 20(3):633–640. doi:10.1093/cercor/bhp128.

Obleser J, Wise RJS, Dresner AM, Scott SK. 2007. Functional Integration across Brain Regions Improves Speech Perception under Adverse Listening Conditions. J Neurosci. 27(9):2283–2289. doi:10.1523/JNEUROSCI.4663-06.2007.

Peelle JE. 2018. Listening Effort: How the Cognitive Consequences of Acoustic Challenge Are Reflected in Brain and Behavior. Ear Hear. 39(2):204–214. doi:10.1097/AUD.0000000000000494.

Penny W, Stephan KE, Daunizeau J, Rosa MJ, Friston KJ, Schofield TM, Leff AP. 2010. Comparing families of dynamic causal models. PLoS Comput Biol. 6(3). doi:10.1371/journal.pcbi.1000709.

Powell HWR, Guye M, Parker GJM, Symms MR, Boulby P, Koepp MJ, Barker GJ, Duncan J. 2004. Noninvasive in vivo demonstration of the connections of the human parahippocampal gyrus. NeuroImage. 22(2):740–747. doi:10.1016/j.neuroimage.2004.01.011.

Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. NeuroImage. 59(3):2142–2154. doi:10.1016/j.neuroimage.2011.10.018.

Rauschecker JP, Scott SK. 2009. Maps and streams in the auditory cortex: nonhuman primates illuminate human speech processing. Nat Neurosci. 12(6):718.

Rodd JM, Davis MH, Johnsrude IS. 2005. The Neural Mechanisms of Speech Comprehension: fMRI studies of Semantic Ambiguity. Cereb Cortex. 15(8):1261–1269. doi:10.1093/cercor/bhi009.

Rosenthal R, Rubin DB. 1994. The counternull value of an effect size: A New Statistic. Psychol Sci. 5(6):329–334. doi:10.1111/j.1467-9280.1994.tb00281.x.

Schütt HH, Harmeling S, Macke JH, Wichmann FA. 2016. Painfree and accurate Bayesian estimation of psychometric functions for (potentially) overdispersed data. Vision Res. 122:105–123. doi:10.1016/j.visres.2016.02.002.

Seghier ML. 2013. The Angular Gyrus. The Neuroscientist. 19(1):43–61. doi:10.1177/1073858412440596.

Seghier ML, Fagan E, Price CJ. 2010. Functional Subdivisions in the Left Angular Gyrus Where the Semantic System Meets and Diverges from the Default Network. J Neurosci. 30(50):16809 – 16817. doi:10.1523/JNEUROSCI.3377-10.2010.

Shannon R, Fu Q-J, Galvin J. 2004. The number of spectral channels required for speech recognition depends on the difficulty of the listening situation. Acta Otolaryngol (Stockh). 124:50–54. doi:10.1080/03655230410017562.

Siegel JS, Power JD, Dubis JW, Vogel AC, Church JA, Schlaggar BL, Petersen SE. 2014. Statistical improvements in functional magnetic resonance imaging analyses produced by censoring high-motion data points. Hum Brain Mapp. 35(5):1981–1996. doi:10.1002/hbm.22307.

Stephan KE, Penny W, Daunizeau J, Moran RJ, Friston KJ. 2009. Bayesian model selection for group studies. NeuroImage. 46(4):1004–1017. doi:10.1016/j.neuroimage.2009.03.025.

Strange BA, Otten LJ, Josephs O, Rugg MD, Dolan RJ. 2002. Dissociable Human Perirhinal, Hippocampal, and Parahippocampal Roles during Verbal Encoding. J Neurosci. 22(2):523–528. doi:10.1523/JNEUROSCI.22-02-00523.2002.

Taylor, W. L. 1953. "Cloze Procedure": A New Tool for Measuring Readability. Journalism Quarterly. 30:415-433.

Teipel SJ, Bokde ALW, Meindl T, Amaro E, Soldner J, Reiser MF, Herpertz SC, Möller H-J, Hampel H. 2010. White matter microstructure underlying default mode network connectivity in the human brain. NeuroImage. 49(3):2021–2032. doi:10.1016/j.neuroimage.2009.10.067.

Vaden KI, Kuchinsky SE, Ahlstrom J, Dubno JR, Eckert MA. 2015. Cortical Activity Predicts Which Older Adults Recognize Speech in Noise and When. J Neurosci. 35(9):3929–3937. doi:10.1523/JNEUROSCI.2908-14.2015.

Vaden KI, Kuchinsky SE, Cute SL, Ahlstrom J, Dubno JR, Eckert MA. 2013. The Cingulo-Opercular Network Provides Word-Recognition Benefit. J Neurosci. 33(48):18979–18986. doi:10.1523/JNEUROSCI.1417-13.2013.

Vaden KI, Teubner-Rhodes S, Ahlstrom J, Dubno JR, Eckert MA. 2017. Cingulo-opercular activity affects incidental memory encoding for speech in noise. NeuroImage. 157(June):381–387. doi:10.1016/j.neuroimage.2017.06.028.

Winn M. 2016. Rapid Release from Listening Effort Resulting from Semantic Context, and Effects of Spectral Degradation and Cochlear Implants. Trends Hear. 20:1–17. doi:10.1177/2331216516669723.