Postnatal Expansion, Maturation and Functionality of MR1T Cells in Humans

Gwendolyn M. Swarbrick[1,4][¶][*], Anele Gela[5][¶], Meghan E. Cansler[1], Megan D. Null[1], Rowan B.
Duncan[1], Elisa Nemes[5], Muki Shey[5, 6], Mary Nsereko[7], Harriet Mayanja-Kizza[7, 8], Sarah Kiguli[7,
9], Jeffrey Koh[3], Willem A. Hanekom[5], Mark Hatherill[5], Christina Lancioni[1], David M.
Lewinsohn[2, 4], Thomas J. Scriba[5][&], Deborah A. Lewinsohn[1][&]

[1]Division of Infectious Diseases, Department of Pediatrics, Oregon Health & Science University,
Portland, Oregon, USA

[2]Division of Pulmonary and Critical Care Medicine, Department of Medicine, Oregon Health &
Science University, Portland, Oregon, USA

[3]Department of Anesthesiology and Perioperative Medicine, Oregon Health & Science University,
Portland, Oregon, USA

[4]Portland Veterans Administration Healthcare System, Portland, Oregon, USA

[5]South African Tuberculosis Vaccine Initiative, Institute of Infectious Disease and Molecular
Medicine and Division of Immunology, Department of Pathology, University of Cape Town, Cape
Town, South Africa

[6]Department of Medicine & Wellcome Centre for Infectious Disease Research in Africa (CIDRI-
Africa), Institute of Infectious Disease and Molecular Medicine, University of Cape Town, South
Africa

[7]Uganda-CWRU Research Collaboration, Kampala, Uganda.

[8]Dept of Medicine, Makerere University, Kampala, Uganda.

[9]Dept of Paediatrics, Makerere University, Kampala, Uganda.

*Correspondence
Gwendolyn Swarbrick
swarbric@ohsu.edu

[¶]These authors contributed equally to this work.

[&]These authors are Joint Senior Authors.

Keywords: innate T cells, human mucosal immunology, MAIT cells, infant, tuberculosis.

Abstract

MR1-restricted T (MR1T) cells are a T cell subset that recognizes and mediates host
immune responses to a broad array of microbial pathogens, including *Mycobacterium*
tuberculosis. Here, we sought to characterize development of circulating human MR1T cells
defined by MR1-5-OP-RU tetramer labelling and expression of TRAV1-2/CD26/CD161. We
analysed postnatal expansion, maturation and functionality of peripheral blood MR1T cells in
cohorts from three different geographic settings with different tuberculosis (TB) vaccination
practices, levels of exposure to and infection with *M. tuberculosis.* Early after birth, frequencies
of MR1-5-OP-RU tetramer-defined MR1T cells increased rapidly by several fold. This coincided
with marked phenotypic changes, from a predominantly $CD4^+$ and $TRAV1-2^-$ phenotype in
neonates, to predominantly $TRAV1-2^+CD8^+$ MR1T cells that also expressed CD26 and CD161.
We also observed that $tetramer^+$ MR1T cells that expressed TNF upon mycobacterial stimulation
were very low in neonates, but increased ~10-fold in the first year of life. These functional MR1T
cells in all age groups were MR1-5-OP-RU $tetramer^{+,}$ $TRAV1-2^+$ cells and expressed CD26 and
CD161, markers that appeared to signal phenotypic and functional maturation of this cell subset.
This age-associated maturation was also marked by the loss of naïve T cell markers on $tetramer^+$
$TRAV1-2^+$ MR1T cells more rapidly than $tetramer^+TRAV1-2^-$ MR1T cells and non-MR1T cells.
These data suggest that neonates have infrequent populations of MR1T cells with diverse
phenotypic attributes; and that exposure to the environment rapidly and preferentially expands the
MR1-5-OP-RU $tetramer^+TRAV1-2^+$ population of MR1T cells, which becomes the predominant
population of functional MR1T cells early during childhood.

Introduction

MR1-restricted T (MR1T) cells, also known as mucosal-associated invariant T (Paul K Sue) cells, are a highly conserved population of T-cells that exhibit innate-like functions similar to those described for NKT cells. MR1T cells are defined by their recognition of metabolite antigens presented by the monomorphic MHC class 1-related molecule, MR1, the most highly conserved MHC class I related molecule in mammalian species (Harriff et al., 2018; Godfrey et al., 2019). MAIT cells, the predominant subset of MR1T cells, are also defined by a highly restricted T-cell-receptor (TCR) repertoire, comprising an invariant TCR α-chain (TRAV1.2-TRAJ33/20/12 in humans; TRAV1-TRAJ33 in mice) with a CDR3 of constant length paired with a limited number of Vβ segments (Vβ2/13 in humans; Vβ6/8 in mice) (Treiner et al., 2003; Martin et al., 2009; Reantragoon et al., 2013; Lepore et al., 2014). MAIT cells have been defined by different methods. Prior to availability of the MR1 tetramer, MAIT cells were defined by expression of TRAV1-2 and other phenotypic markers, like CD161 and CD26. Now the MR1 tetramer can also be more broadly used to define MR1T cells. In peripheral blood, MAIT cells defined by MR1 tetramer or phenotype are highly concordant in adults (Sharma et al., 2015). As we will show in this paper, the MR1 tetramer allows detection of a much broader repertoire of MR1Ts that are not only restricted to TRAV1-2$^+$ (MAIT cells).

MR1T cell development is a stepwise process, with an intra-thymic selection followed by peripheral expansion. Koay *et al* reported a three-stage development pathway for mouse and human MR1T cell populations, where three populations of human MR1T precursors were defined based on differential expression of CD27 and CD161. The initial two stages of MR1-5-OP-RU tetramer$^+$TRAV1-2+CD161$^-$ cells, which were either CD27- or CD27+, occurred in the thymus and these cells were low in the periphery. The more mature MR1 tetramer$^+$TRAV1-2$^+$CD161$^+$CD27$^+$ cells, designated as stage 3, have been shown to represent only a small fraction of T cells but form a more abundant population in adult blood (Martin et al., 2009; Dusseaux et al., 2011; Koay et al., 2016). Such studies indicate that MR1T cell thymopoeisis is complemented by an important postnatal peripheral expansion and maturation, which was recently shown to be dependent upon acquisition of the intestinal microbiome (Constantinides et al., 2019; Legoux et al., 2019).

MR1T cells recognize microbial derived riboflavin (Vitamin B2) biosynthesis intermediate derivatives, such as 5-OP-RU (5-(2-oxopropylideneamino)-6-d-ribitylaminouracil) and a broad spectrum of microbes that synthesize riboflavin, including *S. aureus, E. coli, and C. albicans* (Gold et al., 2010; Kjer-Nielsen et al., 2012; Gold et al., 2013; Corbett et al., 2014; Meermeier et al., 2016). While 5-OP-RU was the first MR1 ligand defined, several other ligands have subsequently been defined, including additional riboflavin-derived ligands, including both stimulatory and inhibitory ligands (Meermeier et al., 2016; Harriff et al., 2018). Upon recognition of microbial antigens or activation through cytokine signaling, MR1T cells display immediate effector function by secreting inflammatory cytokines and cytotoxic molecules, which can kill infected cells (Gold et al., 2010; Dusseaux et al., 2011; Le Bourhis et al., 2013; Kurioka et al., 2015). In a mouse model of pulmonary legionella infection, MR1T cells exhibited enhanced protection in an MR1-dependent manner, mediated by the effector molecules IFN-γ and GM-CSF (Wang et al., 2018). MR1T cells have also been shown to facilitate control of *Klebsiella pneumonia, Mycobacterium*

bovis BCG, and *Francicella tularensis* infections (Georgel et al., 2011; Meierovics et al., 2013;
Sakala et al., 2015; Meierovics and Cowley, 2016). Many of the bacterial pathogens that are
important causes of infant morbidity and mortality, including *S. aureus*, *S. pneumoniae*, and enteric
pathogens (e.g. *E. coli*, *Klebsiella*, *Salmonella*), are recognized by MR1T cells. MR1T cells
possess immediate innate-like, effector functions, and can therefore mediate immune functions
before priming, clonal expansion and maturation of antigen-specific MHC-restricted T cells.
Therefore, MR1T cells could play an important role in host defense to bacterial pathogens,
especially in infancy.

MR1T cells are known to also recognize and respond to *Mycobacterium tuberculosis,* the
leading global cause of mortality from a single infectious agent. Infants and young children suffer
disproportionately from disseminated and severe forms of tuberculosis (TB) disease and are
significantly more susceptible to disease than adults (Marais et al., 2004a; Marais et al., 2004b;
Marais et al., 2006). Here, we sought to characterize development of MR1T cells (defined by MR1-
tetramer labeling) and their subset of phenotypically defined MAIT cells (TRAV1-
$2^+CD26^+CD161^+$ T cells) by analyzing postnatal expansion, maturation and functionality in
humans. We compared MR1T/MAIT cell development in cohorts from three different settings
with differing TB vaccination practices and levels of exposure and infection with *M. tuberculosis.*

Results

Human participants

We enrolled nine cohorts across three different countries to study MR1T cell development. The cohorts in South Africa (SA) were neonates (cord blood, $n = 10$), infants (aged 10 weeks, BCG vaccinated at birth, $n = 10$) and adolescents [aged 14-18 years, BCG vaccinated at birth, QuantiFERON-TB Gold (QFT)-negative, $n = 10$]. The cohorts enrolled in the United States (US) were neonates (cord blood, $n = 10$), infants (aged 9-15 months, BCG unvaccinated, $n = 10$) and adults (aged 29-60 years, $n = 10$). The cohorts in Uganda (UG) were infants [2-24 months old, BCG vaccinated at birth +/-, tuberculin skin test (TST) +/-, $n = 36$], children (24 - 60 months, BCG vaccinated at birth +/-, TST +/-, $n = 23$) and adults (aged 18-60 years, BCG vaccinated at birth +/-, TST +/-, $n = 91$).

Frequencies of tetramer-defined MR1T cell and phenotypically-defined MAIT cell populations increase with age irrespective of BCG vaccination

We defined MR1T cells as $CD3^+$ T cells positively labeled with the MR1-5-OP-RU tetramer. MR1/6FP tetramers were used as a negative control. The MR1T cell subset, MAIT cells, were defined phenotypically as $CD3^+$ T cells that stained positive for antibodies against TRAV1-2, CD161 and CD26. This phenotypic definition of MAIT cells is concordant with the vast majority of tetramer-defined MR1T cells in adults (Sharma et al., 2015). We determined frequencies of MR1T cells and the phenotypically-defined MAIT cell subset in all cohorts (Fig 1). Because of limited availability of the MR1-5-OP-RU tetramer, determination of tetramer-defined MR1T cell frequencies in the Ugandan cohorts was performed on a subset of 10 donors (Fig 1B and S1 Fig). In all cohorts, both tetramer-defined MR1T cell and phenotypically-defined MAIT cell frequencies increased with age (Fig 1B).

Tetramer-defined MR1T cell and phenotypically-defined MAIT cell populations are concordant in adults, adolescents and infants but discordant in neonates

We next determined the degree of concordance between tetramer-defined MR1T cell and phenotypically-defined MAIT cell populations. The majority of phenotypically-defined MAIT cells population, defined by $TRAV1-2^+CD26^+CD161^+$ co-expression, also stained with MR1 tetramer in infants, adolescents and adults, demonstrating high concordance between tetramer-defined MR1T cells and phenotypically-defined MAIT cells in these cohorts. (Fig 1C). However, in cord blood from both SA and US neonates, these two populations were discordant as the minority of $TRAV1-2^+CD26^+CD161^+$ MAIT cells also expressed the MR1T tetramer (Fig 1C). The expression of CD4 and CD8 across the discordant, phenotypically-defined MAIT cell populations that do not bind MR1 tetramer cell populations varied between cohorts.

The MR1-5-OP-RU tetramer$^+$ subset of phenotypically-defined MAIT cells contain the functional population

To investigate whether tetramer-defined MR1T cells and phenotypically-defined MAIT cells have different functional capacity, we measured TNF expression in the US cohort by intracellular cytokine staining (ICS) and flow cytometry following incubation with *M. smegmatis*-infected A549 cells. Among all phenotypically defined by $TRAV1-2^+CD26^+CD161^+$ MAIT cells, only the MR1/5-OP-RU tetramer$^+$ subset expressed TNF; MR1-5-OP-RU tetramer-negative cells,

148 by contrast, did not appear to respond to *M. smegmatis* stimulation (Fig 2). Because of this result,
149 we focused on the functional, tetramer-defined MR1T cells for subsequent experiments.

Neonates have higher proportions of CD4$^+$ and TRAV1-2$^-$ MR1T populations compared to infants, adolescents and adults

To explore the diversity of MR1T cells and how this may change during early development,
we applied the dimensionality reduction method, t-distributed stochastic neighbor embedding
(tSNE), on tetramer-defined MR1T cells. Five discrete phenotypic MR1T cell clusters were
identified in both the SA and US cohorts (Figure 3A-D), the relative proportions of which varied
with age (Figure 3B and D). Specifically, the CD4$^+$ (Clusters 3 and 4) and TRAV1-2$^-$CD26$^-$
CD161$^-$ (Clusters 1, 2 and 3) MR1T cell clusters predominated in neonates, while CD4$^-$CD8$^{+/-}$
TRAV1-2$^+$CD26$^+$CD161$^+$ cluster (Cluster 5) was predominant in infant, adolescent, and adult
cohorts from both the US and SA (Figure 3B and D). Relative proportions of MR1T cell clusters
in 10 week old (SA) and 12 month old (US) infants were more similar to those in adults and
adolescents than to those in neonates, consistent with the hypothesis that MR1T cells mature
rapidly upon exposure to MR1T cell antigens (Figure 3E and F). It was also noteworthy that CD4$^-$
CD8$^{+/-}$TRAV1-2$^+$CD26$^+$CD161$^+$ MR1T cells that fell into cluster 5 comprised a much lower
percentage of the MR1Ts in cord blood, while forming the dominant phenotype in 12-month-old
(US) infants. In the 10 week old (SA) infants, this cluster had started to appear but was not yet
"fully developed", suggesting rapid development during the first months of life, from a
predominantly TRAV1-2 negative to a TRAV1-2 positive population with loss of expression of
CD4 and acquisition of CD26 and CD161. We also note that in the US cohort only there is a
population that is positive for all markers. Because it was only seen in one cohort, it was excluded
from the comparison of clusters.

MR1T cells become more functional with age

Next, we explored how tetramer-defined MR1T cells developed functionally with age in
the US cohort by tracking the TNF-expressing population following *M. smegmatis* or
PMA/Ionomycin stimulation. TNF-expressing tetramer$^+$ MR1T cells in neonates were present at
very low frequencies (Figure 2B) and represented a much smaller proportion of the total tetramer$^+$
MR1T cell population than in infants and adults (Figure 4A and B). Proportions of tetramer$^+$ MR1T
cells responding to either mycobacteria or PMA/Ionomycin were higher in infants than adults
(p=0.0007 and p=0.0021, respectively) or neonates (p<0.0001 for both stimulations). However,
because tetramer$^+$ MR1T cell frequencies were higher in adults than infants (Figure 1B), overall
frequencies of TNF-producing MR1T cells were highest in adults. To explore the functional
capacity of MR1T cells relative to other T cell subsets within each cohort, we compared
proportions of TNF-expressing MR1T cells (CD3$^+$TCRγδ$^-$MR1-5-OP-RU$^+$) with TCRγδ T cells
(CD3$^+$TCRγδ$^+$ MR1-5OPRU$^-$), CD4$^+$ T cells (CD3$^+$TCRγδ$^-$MR1-5OPRU$^-$CD4$^+$) and CD8$^+$ T cells
(CD3$^+$TCRγδ$^-$MR1-5OPRU$^-$CD4$^+$) after PMA/Ionomycin stimulation (Figure 4C). In neonates,
proportions of TNF-expressing MR1T cells were significantly higher than proportions of TCRγδ
T cells and CD8$^+$ T cells. In infants and adults, proportions of TNF-expressing MR1T cells were
significantly higher than TCRγδ$^+$, CD4$^+$ and CD8$^+$ T cells (Figure 4C).

Functional CD3$^+$MR1-5-OP-RU$^+$ MR1T cells are TRAV1-2$^+$

188    Next, we investigated TRAV1-2 expression among functional, TNF-expressing
190 CD3$^+$MR1-5-OP-RU$^+$ MR1T cells. Only TRAV1-2$^+$ MR1 T cells produced TNF in response to
191 mycobacterial stimulation while TRAV1-2$^-$ MR1 T cells were not functional (Figure 4D). This
192 pattern was unique to mycobacterial stimulation since TNF-expressing TRAV1-2$^-$ MR1T cells
193 were observed in response to PMA/Ionomycin stimulation, particularly in infants and adults
194 (Figure 4E). Interestingly, the vast majority of these functional, TNF-expressing TRAV1-2$^+$
195 MR1T cells also expressed the phenotypic markers, CD26 and CD161 (Figure 4F-G). Thus, the
196 CD3$^+$MR1-5-OP-RU$^+$TRAV1-2$^+$CD26$^+$CD161$^+$ MR1T cell population contains the functional
197 MR1T cells that respond to mycobacteria. The frequency of this population increases very quickly
198 with age.

TRAV1-2$^+$ MR1T cells lose naïve markers more rapidly with age than TRAV1-2$^-$ MR1T cells
or non-MR1T cells

201    To characterize the development of MR1T cells further, we also measured expression of
202 CD27 and CD161, markers previously described to indicate MR1T cell maturation, and CD45RA,
203 CCR7, CD38 and CD56 on tetramer-defined MR1Ts (Figures 5-6) (Koay et al., 2018). A
204 population of immature MR1T cells, defined as CD27$^+$CD161$^-$ cells, was observed in neonates but
205 not in infants and adults, as previously described (Koay et al., 2016). This was also reflected by
206 expression of the more traditional T cell maturation markers, CD45RA, CCR7 and CD38 (Figure
207 6 A, C, E). Most tetramer-defined MR1T cells in neonates displayed a CD45RA$^+$CCR7$^+$CD38$^+$
208 "naïve" phenotype, but this phenotype was lost more rapidly with age on TRAV1-2$^+$ MR1Ts than
209 on TRAV1-2$^-$ MR1T cells or non-MR1T cells (Figure 6A-F). The inverse was true for the
210 activation marker, CD56, which was increased in neonatal, infant and adult TRAV1-2$^+$ tetramer-
211 defined MR1T cells compared to TRAV1-2$^-$ tetramer-defined MR1T cells and non-MR1T cells
212 (Figure 6G-H). These results are consistent with our finding that TRAV1-2$^+$ tetramer-defined
213 MR1T cells display functional capacity in all age groups, even though they represent the minority
214 of neonatal MR1T cells.

Discussion

We characterized the development of MR1T cells during early childhood in three cohorts, two from Africa and one in the USA, to elucidate phenotypic and functional changes that occur after birth and during the first few years of life. A better understanding of MR1T cell development is critical given the likely role of MR1T cells as innate-like effectors that act as key mediators of immunity in newborns, who are at very high risk of infection, including *M. tuberculosis.* We found that neonates had relatively low frequencies of MR1T cells with high phenotypic diversity, which included considerable proportions of CD4$^+$ and TRAV1-2$^-$ cells. These patterns changed as early as 10 weeks after birth, and became more prominent at 1-2 years of age, adolescence and ultimately by adulthood. This included higher frequencies of MR1T cells in peripheral blood, characterized by a preferential expansion in TRAV1-2$^+$CD8$^+$ MR1T cells such that TRAV1-2$^+$CD8$^+$MR1T cells constituted the vast majority of the tetramer$^+$ MR1T cell population. The development also included changes in expression of maturation markers and MR1T cell effector function, since MR1T cells from older individuals had lower expression of the naïve T cell markers, CD45RA, CCR7 and CD38, and an increased ability to express TNF.

Our finding that MR1T cells are found at low frequency in cord blood and increase in frequency with age confirms our previous observations, as well as those of others, who have utilized either the MR1-5-OP-RU tetramer or phenotypic characterization of MAIT cells to demonstrate the expansion of these cell populations from birth through adulthood (Martin et al., 2009; Gold et al., 2013; Walker et al., 2014; Koay et al., 2016; Ben Youssef et al., 2018; Gherardin et al., 2018). The very rapid changes in MR1T cells seen after birth are striking and were clearly seen in both African and American settings. This finding rules out the possibility that BCG substantially contributes to these changes, since BCG is not routinely administered at birth in the USA. Rather, it is more likely that changes occur upon the transition from the limited microbial exposure in the womb to the postnatal environment, accompanied by the rapid colonization of the gut microbiome. MR1T cells are absent in germ-free mice and depend on the colonization of the gut microbiome with normal intestinal flora, which produce riboflavin (Le Bourhis et al., 2013; Harriff et al., 2018; Legoux et al., 2019). We suggest that this colonization of the gut microbiome is the driving force that accounts for the increase and phenotypic changes in MR1T cells in blood after birth and the gain in functional capacity.

While we confirm our previous work that phenotypic definition of MAIT cells accurately identifies functional MR1T cells in adults, we find that there is a high level of discordance in neonates in part due to MR1-5-OP-RU tetramer$^+$ cells that do not express TRAV1-2, CD26 or CD161. These cells are by definition MR1-restricted T cells as they bind the MR1-5-OP-RU tetramer. However, these cells are poorly reactive to mycobacteria, suggesting that they either recognize a different antigen or represent immature MR1T cells. The poor PMA/Ionomycin-induced TNF secretion of these cells supports the latter. However, because our only measure of MR1T cell responses was TNF-α production, we may have failed to detect other responses characterized by the production of alternative cytokines or cytolytic markers.

We also show that neonates possess MR1-5-OP-RU tetramer$^-$ cells that phenotypically resemble MAIT cells (the discordant MR1-5-OP-RU tetramer$^-$ phenotypically defined (CD26$^+$, CD161$^+$, and TRAV1-2$^+$) MAIT cell population). As Fergusson *et al* suggested that distinct T

subsets that express CD161 share a common transcriptional profile characteristic of typical MR1T
cells (Fergusson et al., 2014), it is certainly possible that these CD161$^+$ cells are MR1-restricted or
share common attributes with MR1T cells. Further characterization of this neonatal T cell
population is required to determine whether or not these cells are indeed MR1-restricted and if so,
if these cells are immature or recognize distinct microbes, or produce a different cytokine not
detected in our functional assay. Given that, in the mouse model, 5-OP-RU produced by the
microbiome is necessary and sufficient to drive the expansion of typical MR1T cells (Legoux et
al., 2019), it is perhaps expected that with age tetramer-positive MR1T TRAV1-2$^+$ cells would
quickly become the predominant MR1T cell population, while tetramer negative MR1T cells
might persist as a minority population of naïve T cells.

Consistent with our prior findings (Gold et al., 2013) and those of Youssef *et al*. (Ben
Youssef et al., 2018), our phenotypic analysis of MR1-5-OP-RU tetramer$^+$ cells in neonates
demonstrates that MR1T cells expressed a naïve cell surface phenotype (CD45RA$^+$CCR7$^+$CD38$^+$)
and included a prominent subset of CD4$^+$ cells.   It is noteworthy that these MR1T cells rapidly
lost expression of CD45RA, CCR7 and CD38 and acquired a phenotype consistent with those
observed in antigen-experienced memory T cells much more rapidly than other types of T cells
(Chua et al., 2012; Tsukamoto et al., 2013; Lindestam Arlehamn et al., 2014).

While Koay *et al* showed the functional potential of neonatal MR1T cells that produced
TNF following stimulation with PMA/Ionomycin, we show for the first time the presence of
functional microbe-reactive MR1T cells in neonates, albeit at low frequency. Virtually all of the
mycobacteria-reactive TNF-producing MR1T cells were contained within the TRAV1-2$^+$CD26$^+$
CD161$^+$ subset of 5-OP-RU tetramer$^+$ cells, which are also concordant with "Stage 3" MR1T cells
defined by Koay *et al*. Although the immature CD4$^+$, CD4-CD8- and/or TRAV1-2- MR1T cells
were maintained through to adulthood, we observed that CD8$^+$TRAV1-2$^+$ MR1T cells expanded
preferentially over TRAV1-2$^-$ MR1T cells. This expansion coincided with the acquisition of CD26
and CD161 as markers of mature functional MR1 T cells. In addition, we show that the frequency
of these functional mycobacterial-reactive MR1T cells increases ~10-fold in the first year of life.
A surprising finding was that given a comparable stimulus (PMA/Ionomycin and *M. smegmatis*),
the infant MR1Ts exhibited a higher functional capacity than the adults. Interestingly, we
demonstrated that relative to other T cell subsets, MR1T cells had the highest capacity to secrete
the pro-inflammatory cytokine, TNF-α, in infants. Thus, MR1T cells may represent an important
source of pro-inflammatory effectors in infants, who are more vulnerable to serious bacterial
infections in the first two years of life, relative to older children and adults.

Regarding newborns, it is not known to what degree the naïve phenotype and low
functionality of MR1T cells contributes to the very high susceptibility to bacterial infection in the
first two months of life (Paul K Sue, 2018). Since MR1T cells recognize important childhood
bacterial pathogens, including *S. aureus, S. pneumoniae and enteric pathogens (e.g. E. coli,*
Klebsiella, Salmonella species, it is plausible that the low frequency of functional MR1T cells
during the neonatal period contributes to neonatal susceptibility to infection. At the very least, our
findings add to the well-described differences in adaptive and innate immunity between neonates
and adults (Paul K Sue, 2018).

298   With regard to *M. tuberculosis*, there is mounting evidence that supports an important host
299 defense role for MR1T cells, especially during early infection. Upon encounter with
300 mycobacterial-infected cells, MR1T cells produce pro-inflammatory cytokines such as IFN-γ and
301 TNF (Gold et al., 2010), known to be essential mediators to host defense against *M. tuberculosis*
302 (Lindestam Arlehamn et al., 2014). MR1-deficient mice control BCG less well than wild type
303 mice (Chua et al., 2012). In humans, MR1 polymorphisms are associated with susceptibility to TB
304 meningitis (Seshadri et al., 2017). It is tempting to speculate, in light of the high susceptibility of
305 infants to TB, and the phenotypic and functional attributes of MR1T cells we observed, that MR1T
306 cells contribute to control of early infection with *M. tuberculosis* through their early and robust
307 anti-mycobacterial response, particularly in the lung.

Methods

Study participants

University of Cape Town, Western Cape, South Africa. All participants were enrolled either at the South African Tuberculosis Vaccine Initiative Field Site, Well-Baby Clinics or Maternity Wards in the Worcester region, near Cape Town, South Africa. Pregnant mothers were enrolled for collection of cord blood immediately after childbirth by Cesarean section. Mothers who may have been exposed to *M. tuberculosis*, HIV infected, had any other acute or chronic disease, and receiving medication other than anesthetic given during delivery were excluded. Infants born to HIV negative mothers, and who had received BCG at birth, had no perinatal complications, or household contact with any person with TB, and did not have any acute or chronic disease were enrolled for collection of blood. Healthy adolescents of 12-18 years of age without evidence of latent Mtb infection as determined by positivity negativity for the Tuberculin Skin Test (<10mm) or QuantiFERON (<0.35IU/mL) were enrolled for blood collection. Heparinized blood was collected for cord blood mononuclear cell (CBMC) and peripheral blood mononuclear cells (PBMC) isolation by means of density gradient centrifugation.

Oregon Health and Science University, Portland, OR, USA. Umbilical cord blood was obtained from the placenta of uncomplicated term pregnancies after delivery into citrate CPT tubes (BD). Blood was processed per manufacturer's instructions within 24hr of collection, and resulting CBMC cryopreserved. Infants aged 1-24 months undergoing elective surgery at OHSU Doernbecher Children's Hospital were enrolled. Infants had not received BCG vaccination, which is not part of the childhood immunization series in the United States. Blood was drawn into citrate CPT tubes (BD) and processed per manufacturer's instructions to isolate PBMC within 24hr of collection, and PBMC subsequently cryopreserved. Adult PBMC were obtained by leukapheresis from healthy adult donors and cryopreserved. Adult participants lacked any detectable responses to *Mycobacterium tuberculosis* (Mtb) immunodominant antigens CFP-10 and ESAT-6 and therefore were assumed to be Mtb uninfected.

Makerere University, Kampala, Uganda. Children from age 2-60 months hospitalized with lower respiratory infection (LRTI) from etiologies other than TB were recruited from Mulago Hospital Paediatric Wards. All children were HIV-negative and had no known recent TB contacts. Children were reassessed two months following enrollment to establish vital status and ensure clinical improvement. Following completion of the study, data from all participating children was reviewed by study investigators (DAL, CLL, SK) to confirm diagnosis of non-TB LRTI. Blood was drawn into citrate CPT tubes (BD) and processed per manufacturer's instructions to isolate PBMC within 24hr of collection, and PBMC subsequently cryopreserved. Adult participants were recruited from the National TB Treatment Center at Mulago Hospital, hospital staff, and the community surrounding Kampala, Uganda, between 2001 and 2014. Participants were HIV uninfected and included individuals with (tuberculin skin test, TST $\geq$ 10) or without (TST < 10 mm) evidence of Mtb infection. All had the absence of signs or symptoms of TB (prolonged cough, hemoptysis, fever, weight loss, and night sweats). Heparinized blood was collected for PBMC isolation by means of density gradient centrifugation.

Study approvals

The studies of all participants were conducted in accordance with the World's Medical
Association's Declaration of Helsinki guidelines. The studies from the Western Cape, South Africa
were approved by the Human Research Ethics Committee (HREC) of the University of Cape
Town. Written informed consent was obtained from parents or legal guardians of adolescents and
infants. Adolescents also provided written informed assent. The Institutional Review Board of
Oregon Health & Science University (Portland, Oregon) approved this study of participants from
Portland, OR. Written, informed consent was obtained from all adult participants and from a parent
or legal guardian of infants before enrollment. For participants from Kampala, Uganda, the
National Council for Science and Technology, HIV/AIDS Research Committee in Uganda, the
Institutional Review Board of Oregon Health & Sciences University (Portland, OR) and the
Institutional Review Board of Makerere University (Kampala, Uganda) approved this study.
Written, informed consent was obtained from all adult participants and from a parent or legal
guardian of infants and children before enrollment.

Infection of antigen presenting cells

A549 cells (ATCC CCL-185) were used as stimulators for direct ex vivo determination of
mycobacterial-reactive T cells. *Mycobacterium smegmatis* (strain mc^{2}155, ATCC) was grown to
late log phase and functionally titered in this assay to find the optimal volume of *M. smegmatis*
that elicited a strong response in an adult control donor but did not reduce the viability of the
antigen presenting cells. A549 cells were seeded in T25 flasks at 1.25e6 in 7.5ml of antibiotic free
media. After allowing cells to adhere for 3 hours, the flasks were infected with 31μl of *M.*
smegmatis. After 18 h, the cells were harvested and washed twice in RPMI/10% human serum
containing antibiotic (gentamicin) before being resuspended in RPMI/10% human serum
containing antibiotic (gentamicin).

Flow cytometry assays

To perform ICS, cryopreserved PBMC were thawed in the presence of DNAse,
resuspended, in 10% heat inactivated human serum with RPMI (Lonza) at a concentration of 2 x
10^7/ml. PBMC (2 x 10^6 cells/well) were stimulated with anti-CD28 (1 μg/ml) and anti-CD49d (1
μg/ml) (FastImmune, BD Biosciences) and 1e5 *M. smegmatis*-infected or uninfected A549 cells
in RPMI/10% Human Serum for 18 h at 37°C with 5% C0$_2$. During the last 12 h of this incubation,
Brefeldin-A (5 μg/ml) was added. PBMC incubated with uninfected A549 cells were used to assess
for background cytokine production. To determine the total functional capacity of T cells, PBMC
were incubated with uninfected A549 cells and PMA/Ionomycin (PMA, 20ng/ml, Sigma;
Ionomycin, 1μM, Sigma) for 3 hr. PBMC were then harvested and stained with MR1-5-OP-RU or
MR1-6FP tetramers (Corbett et al., 2014) (NIH Tetramer Core) at a 1:500 dilution for 45 minutes
at room temperature. All tetramers used were conjugated to R-phycoerythrin (PE) or Brilliant
Violet (BV) 421. After 45 minutes, an antibody cocktail containing LIVE/DEAD Fixable Dead
Cell Stain Kit (Thermo Fisher), and the surface antibodies listed in S1 Table were added for 30
minutes at 4°C. After washing, cells were fixed and permeabilized using Cytofix/Cytoperm (BD
Biosciences) per manufacturer's instructions. Cells were then stained for CD3 and TNF, then fixed
in 1% PFA. Acquisition was performed using a (Beckman Coulter) Cytoflex flow cytometer with

CytExpert software (BC). All flow cytometry data was analyzed using FlowJo software (TreeStar)
and Prism (Graphpad).

To perform cell surface staining for phenotyping MR1T cells, cryopreserved PBMC were
thawed as described above, and 1-2 x 10^6 PBMC were stained with MR1/5-OP-RU or MR1/6FP
tetramers at 1:500 for 45 minutes at room temperature. Cells were then subsequently stained with
an antibody cocktail containing LIVE/DEAD Fixable Dead Cell Stain Kit (Thermo Fisher), and
surface stained with the antibodies listed in S2-S4 Tables for 30 minutes at 4°C. Samples were
washed and fixed with 1% PFA. Acquisition of the US samples was performed using a Cytoflex
flow cytometer (Beckman) with CytExpert software (BC). For experiments with the Ugandan
samples, acquisition was performed on a BD LSRFortessa. For experiments performed at SATVI,
samples were acquired on a BD LSR-II flow cytometer configured with 4 lasers: solid state blue
(488nm; 100mW; 3 detectors), solid state violet (405nm; 25mW; 8 detectors), HeNe gas red
(635nm; 70mW; 3 detectors), and diode-pumped Coherent Compass yellow-green (532nm;
150mW; 8 detectors). Flow cytometric analysis was performed using FlowJo version 10.5.3
(Treestar, Ashland, OR, USA).

Data analysis and Statistics

Adjusted frequency (raw frequency of cytokine production following stimulation minus
the frequency of cytokine production in resting/unstimulated condition), was used to account for
the potential presence of background, or non-specific, responses. A positive cytokine response was
defined as the detection of at least 0.05% cytokine-positive T cells (following background
subtraction). Prism version 7 (GraphPad, La Jolla, CA, USA) was used for data analysis. A p-value
< 0.05 was considered as statistically significant.

To characterize phenotypic diversity within MR1T cell populations at the different ages
we performed t-Stochastic Neighbor Embedding (tSNE) on MR1-5-OP-RU tetramer[+] defined
CD3[+] T cells. Data from MR1-5-OP-RU tetramer[+] cells from all individuals across the age groups
were concatenated into a single FCS file to perform tSNE analysis in FlowJo. Equal event sampling
was not performed due to low events per individual in the cord blood and infant samples.
Clustering analysis was performed based on the expression of five surface markers (CD4, CD8,
TRAV1-2, CD26, and CD161). Parameter settings were fixed at 2000 iterations with a learning
rate (Eta) of 10 and perplexity 30, respectively (van der Maaten and Hinton, 2008).

Conflict of Interest statement

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Author Contributions

GMS, AG, DML, TJS and DAL contributed to the conception and/or design of the
work. GMS and AG are joint first authors. TJS and DAL jointly supervised the work. MN, HMK,
SK, JK, CL, EN, MS, WAH, MH and TJS contributed to clinical activities. EN, WAH, MH, DML,
DAL and TJS raised grants to fund the research. GMS, AG, MEC, MDN, RBD, DML, TJS and
DAL substantially contributed to the acquisition, analysis or interpretation of data and drafting of
the manuscript. All authors substantially contributed to revising and critically reviewing the
manuscript for important intellectual content. All authors approved the final version of this
manuscript to be published and agree to be accountable for all aspects of the work.

Funding

This project has been funded in whole or in part with Federal funds from the National Institutes of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, under grant no R01 AI04229 and contract HHSN272200900053C.

Work at SATVI was supported by Aeras, the National Institutes of Health Grant R01-AI087915 and R01-AI065653 and the European and Developing Countries Clinical Trial Partnership. Anele Gela was supported by a Postdoctoral Fellowship from the Claude Leon Foundation.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We would like to thank the participants who gave time and dedication to this health research; W. Henry Boom, LaShaunda Malone and Keith Chervenak, Case Western Reserve University; and Erin Merrifield, Department of Pediatrics, OHSU, for their contributions to this study.

The MR1 tetramer technology was developed jointly by Dr. James McCluskey, Dr. Jamie Rossjohn, and Dr. David Fairlie, and the material was produced by the NIH Tetramer Core Facility as permitted to be distributed by the University of Melbourne.

References

Ben Youssef, G., Tourret, M., Salou, M., Ghazarian, L., Houdouin, V., Mondot, S., et al. (2018).

Ontogeny of human mucosal-associated invariant T cells and related T cell subsets. *Journal*

of Experimental Medicine 215(2)**,** 459-479. doi: 10.1084/jem.20171739.

Chua, W.J., Truscott, S.M., Eickhoff, C.S., Blazevic, A., Hoft, D.F., and Hansen, T.H. (2012).

Polyclonal Mucosa-Associated Invariant T Cells Have Unique Innate Functions in

Bacterial Infection. *Infection and Immunity* 80(9)**,** 3256-3267. doi: 10.1128/Iai.00279-12.

Constantinides, M.G., Link, V.M., Tamoutounour, S., Wong, A.C., Perez-Chaparro, P.J., Han,

S.J., et al. (2019). MAIT cells are imprinted by the microbiota in early life and promote

tissue repair. *Science* 366(6464). doi: 10.1126/science.aax6624.

Corbett, A.J., Eckle, S.B.G., Birkinshaw, R.W., Liu, L.G., Patel, O., Mahony, J., et al. (2014).

T-cell activation by transitory neo-antigens derived from distinct microbial pathways.

Nature 509(7500)**,** 361-+. doi: 10.1038/nature13160.

Dusseaux, M., Martin, E., Serriari, N., Peguillet, I., Premel, V., Louis, D., et al. (2011). Human

MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17-secreting T cells.

Blood 117(4)**,** 1250-1259. doi: 10.1182/blood-2010-08-303339.

Fergusson, J.R., Smith, K.E., Fleming, V.M., Rajoriya, N., Newell, E.W., Simmons, R., et al.

(2014). CD161 Defines a Transcriptional and Functional Phenotype across Distinct Human

T Cell Lineages. *Cell Reports* 9(3)**,** 1075-1088. doi: 10.1016/j.celrep.2014.09.045.

Georgel, P., Radosavljevic, M., Macquin, C., and Bahram, S. (2011). The non-conventional

MHC class I MR1 molecule controls infection by Klebsiella pneumoniae in mice. *Mol*

Immunol 48(5)**,** 769-775. doi: 10.1016/j.molimm.2010.12.002.

Gherardin, N.A., Souter, M.N.T., Koay, H.F., Mangas, K.M., Seemann, T., Stinear, T.P., et al.

(2018). Human blood MAIT cell subsets defined using MR1 tetramers. *Immunology and*

Cell Biology 96(5)**,** 507-525. doi: 10.1111/imcb.12021.

Godfrey, D.I., Koay, H.F., McCluskey, J., and Gherardin, N.A. (2019). The biology and

functional importance of MAIT cells. *Nat Immunol* 20(9)**,** 1110-1128. doi:

10.1038/s41590-019-0444-8.

Gold, M.C., Cerri, S., Smyk-Pearson, S., Cansler, M.E., Vogt, T.M., Delepine, J., et al. (2010).

Human mucosal associated invariant T cells detect bacterially infected cells. *PLoS Biol*

8(6)**,** e1000407. doi: 10.1371/journal.pbio.1000407.

Gold, M.C., Eid, T., Smyk-Pearson, S., Eberling, Y., Swarbrick, G.M., Langley, S.M., et al.

(2013). Human thymic MR1-restricted MAIT cells are innate pathogen-reactive effectors

that adapt following thymic egress. *Mucosal Immunology* 6(1)**,** 35-44. doi:

10.1038/mi.2012.45.

Harriff, M.J., McMurtrey, C., Froyd, C.A., Jin, H.H., Cansler, M., Null, M., et al. (2018). MR1

displays the microbial metabolome driving selective MR1-restricted T cell receptor usage.

Science Immunology 3(25). doi: UNSP eaao2556

10.1126/sciimmunol.aao2556.

Kjer-Nielsen, L., Patel, O., Corbett, A.J., Le Nours, J., Meehan, B., Liu, L., et al. (2012). MR1

presents microbial vitamin B metabolites to MAIT cells. *Nature* 491(7426)**,** 717-723. doi:

10.1038/nature11605.

Koay, H.F., Gherardin, N.A., Enders, A., Loh, L., Mackay, L.K., Almeida, C.F., et al. (2016). A

three-stage intrathymic development pathway for the mucosal-associated invariant T cell

lineage. *Nature Immunology* 17(11)**,** 1300-1311. doi: 10.1038/ni.3565.

Koay, H.F., Godfrey, D.I., and Pellicci, D.G. (2018). Development of mucosal-associated

invariant T cells. *Immunology and Cell Biology* 96(6)**,** 598-606. doi: 10.1111/imcb.12039.

Kurioka, A., Ussher, J.E., Cosgrove, C., Clough, C., Fergusson, J.R., Smith, K., et al. (2015).

MAIT cells are licensed through granzyme exchange to kill bacterially sensitized targets.

Mucosal Immunology 8(2)**,** 429-440. doi: 10.1038/mi.2014.81.

Le Bourhis, L., Dusseaux, M., Bohineust, A., Bessoles, S., Martin, E., Premel, V., et al. (2013).

MAIT Cells Detect and Efficiently Lyse Bacterially-Infected Epithelial Cells. *Plos*

Pathogens 9(10). doi: ARTN e1003681

10.1371/journal.ppat.1003681.

Legoux, F., Bellet, D., Daviaud, C., El Morr, Y., Darbois, A., Niort, K., et al. (2019). Microbial

metabolites control the thymic development of mucosal-associated invariant T cells.

Science 366(6464)**,** 494-499. doi: 10.1126/science.aaw2719.

Lepore, M., Kalinicenko, A., Colone, A., Paleja, B., Singhal, A., Tschumi, A., et al. (2014).

Parallel T-cell cloning and deep sequencing of human MAIT cells reveal stable oligoclonal

TCR beta repertoire. *Nature Communications* 5. doi: ARTN 3866

10.1038/ncomms4866.

Lindestam Arlehamn, C.S., Lewinsohn, D., Sette, A., and Lewinsohn, D. (2014). Antigens for

CD4 and CD8 T cells in tuberculosis. *Cold Spring Harb Perspect Med* 4(7)**,** a018465. doi:

10.1101/cshperspect.a018465.

Marais, B.J., Gie, R.P., Schaaf, H.S., Hesseling, A.C., Enarson, D.A., and Beyers, N. (2006).

The spectrum of disease in children treated for tuberculosis in a highly endemic area.

International Journal of Tuberculosis and Lung Disease 10(7)**,** 732-738.

Marais, B.J., Gie, R.P., Schaaf, H.S., Hesseling, A.C., Obihara, C.C., Nelson, L.J., et al. (2004a).

The clinical epidemiology of childhood pulmonary tuberculosis: a critical review of

literature from the pre-chemotherapy era. *International Journal of Tuberculosis and Lung*

Disease 8(3)**,** 278-285.

Marais, B.J., Gie, R.R., Schaaf, H.S., Hesseling, A.C., Obihara, C.C., Starke, J.J., et al. (2004b).

The natural history of childhood intra-thoracic tuberculosis: a critical review of literature

from the pre-chemotherapy era. *International Journal of Tuberculosis and Lung Disease*

8(4)**,** 392-402.

Martin, E., Treiner, E., Duban, L., Guerri, L., Laude, H., Toly, C., et al. (2009). Stepwise

Development of MAIT Cells in Mouse and Human. *Plos Biology* 7(3)**,** 525-536. doi:

ARTN e1000054

10.1371/journal.pbio.1000054.

Meermeier, E.W., Laugel, B.F., Sewell, A.K., Corbett, A.J., Rossjohn, J., McCluskey, J., et al.

(2016). Human TRAV1-2-negative MR1-restricted T cells detect S-pyogenes and

alternatives to MAIT riboflavin-based antigens. *Nature Communications* 7. doi: ARTN

12506

10.1038/ncomms12506.

Meierovics, A., Yankelevich, W.J.C., and Cowley, S.C. (2013). MAIT cells are critical for

optimal mucosal immune responses during in vivo pulmonary bacterial infection.

Proceedings of the National Academy of Sciences of the United States of America 110(33)**,**

E3119-E3128. doi: 10.1073/pnas.1302799110.

Meierovics, A.I., and Cowley, S.C. (2016). MAIT cells promote inflammatory monocyte

differentiation into dendritic cells during pulmonary intracellular infection. *Journal of*

Experimental Medicine 213(12)**,** 2793-2809. doi: 10.1084/jem.20160637.

Paul K Sue, M.M., and Maite de la Morena (2018). "Immunologic Development and

Susceptibility to Infection.," in *Principles and Practice of Pediatric Infectious Diseases.*

5th ed., ed. G.C.P. S. Sarah Long, M. Fischer.: Elsevier), p85-92.

Reantragoon, R., Corbett, A.J., Sakala, I.G., Gherardin, N.A., Furness, J.B., Chen, Z.J., et al.

(2013). Antigen-loaded MR1 tetramers define T cell receptor heterogeneity in mucosal-

associated invariant T cells. *Journal of Experimental Medicine* 210(11)**,** 2305-2320. doi:

10.1084/jem.20130958.

Sakala, I.G., Kjer-Nielsen, L., Eickhoff, C.S., Wang, X., Blazevic, A., Liu, L., et al. (2015).

Functional Heterogeneity and Antimycobacterial Effects of Mouse Mucosal-Associated

Invariant T Cells Specific for Riboflavin Metabolites. *J Immunol* 195(2)**,** 587-601. doi:

10.4049/jimmunol.1402545.

Seshadri, C., Thuong, N.T.T., Mai, N.T.H., Bang, N.D., Chau, T.T.H., Lewinsohn, D.M., et al.

(2017). A polymorphism in human MR1 is associated with mRNA expression and

susceptibility to tuberculosis. *Genes and Immunity* 18(1)**,** 8-14. doi: 10.1038/gene.2016.41.

Sharma, P.K., Wong, E.B., Napier, R.J., Bishai, W.R., Ndung'u, T., Kasprowicz, V.O., et al.

(2015). High expression of CD26 accurately identifies human bacteria-reactive MR1-

restricted MAIT cells. *Immunology* 145(3)**,** 443-453. doi: 10.1111/imm.12461.

Treiner, E., Duban, L., Bahram, S., Radosavljevic, M., Wanner, V., Tilloy, F., et al. (2003).

Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1 (vol

422, pg 164, 2003). *Nature* 423(6943)**,** 1018-1018. doi: 10.1038/nature01700.

Tsukamoto, K., Deakin, J.E., Graves, J.A.M., and Hashimoto, K. (2013). Exceptionally high

conservation of the MHC class I-related gene, MR1, among mammals. *Immunogenetics*

65(2)**,** 115-124. doi: 10.1007/s00251-012-0666-5.

van der Maaten, L., and Hinton, G. (2008). Visualizing Data using t-SNE. *Journal of Machine*

Learning Research 9**,** 2579-2605.

Walker, L.J., Tharmalingam, H., and Klenerman, P. (2014). The Rise and Fall of MAIT Cells

with Age. *Scandinavian Journal of Immunology* 80(6)**,** 462-463. doi: 10.1111/sji.12237.

Wang, H.M., D'Souza, C., Lim, X.Y., Kostenko, L., Pediongco, T.J., Eckle, S.B.G., et al. (2018).

MAIT cells protect against pulmonary Legionella longbeachae infection. *Nature*

Communications 9. doi: ARTN 3350

10.1038/s41467-018-05202-8.

 Figure Legends

Figure 1. Frequencies of tetramer-defined MR1T cell and phenotypically defined MAIT cell
populations in peripheral blood from individuals of different ages. PBMC or CBMC were
stained with a live/dead discriminator, antibodies to CD3, CD4, CD8, TRAV1-2, CD26, CD161,
and either the MR1-5-OP-RU or MR1-6FP tetramers. Live, CD3$^+$ lymphocytes were gated and the
frequencies of MR1-5-OP-RU$^+$ or TRAV1-2$^+$CD26$^+$CD161$^+$ cells as a percentage of CD3$^+$
lymphocytes were determined (gating strategy in Figure S2). **A.** Flow cytometry plots showing
CD3$^+$ T cell staining with MR1-5-OP-RU tetramer, MR1-6FP tetramer, TRAV1-2 or
CD26/CD161 of samples from a representative US adult, infant and neonate. **B.** Frequencies of
tetramer-defined (CD3$^+$MR1-5-OP-RU$^+$) and phenotypically-defined (CD3$^+$TRAV1-
2$^+$CD26$^+$CD161$^+$) MAIT cells in neonates, 10 week-old infants and adolescents from South Africa,
neonates, 12 month-old infants and adults from the United States and infants (0-2 years old),
children (2-5 years old) and adults from Uganda (all cohorts, *n*=10). Mann-Whitney u-tests were
used to test differences between groups. Horizontal lines depict the median and the error bars the
95% confidence interval. **C.** Relative proportions of median phenotypically-defined
(CD3$^+$TRAV1-2$^+$CD26$^+$CD161$^+$) MAIT cells that are MR1-5-OP-RU$^+$ (Blue) and MR1-5-OP-
RU$^-$ (Red shades) for each age group at each site. In each doughnut chart the MR1-5-OP-RU$^-$
population is further divided into CD8$^+$, CD8$^+$CD4$^+$, CD4$^+$, and CD8$^-$CD4$^-$ populations.

Figure 2. Functional analysis of MAIT cells in individuals of different ages. PBMC or CBMC
from the US cohort were incubated overnight with *M. smegmatis*-infected A549 cells or uninfected
A549 cells and then stained with the MR1-5-OP-RU or MR1/6FP tetramers, followed by staining
with a live/dead discriminator and antibodies to TCRγδ, CD3, CD4, CD8, TRAV1-2, CD26 and
CD161. ICS was then performed and the cells stained for TNF. **A.** Dot plots showing representative
co-staining of live, TCRγδ$^-$CD3$^+$TRAV1-2$^+$CD26$^+$CD161$^+$ cells with MR1-5-OP-RU and TNF in
M. smegmatis stimulated or unstimulated samples. The gating strategy is shown in Figure S3.
Examples of the TNF response in a neonate, infant and adult are shown. **B.** Frequencies of
functional, phenotypically-defined (CD3$^+$TRAV1-2$^+$CD26$^+$CD161$^+$) MAIT cells that are MR1-5-
OP-RU$^+$ or MR1-5-OP-RU$^-$ as a proportion of CD3$^+$ TCRγδ$^-$ T cells. Horizontal lines depict the
median and the error bars the 95% confidence interval. Wilcoxon-rank sum was used to test
differences within the same cohort.

Figure 3. Phenotypic diversity of MR1T cells in individuals of different ages. t-Distributed
stochastic neighbor embedding (tSNE) plots showing MR1-5-OP-RU tetramer-defined MR1T
cells (gating strategy in Figure S2) and the phenotypic clusters of MR1T cells identified in the
South African (SA) or United States (US) cohorts. Plots represent cells from 10 samples from each
age group. **A.** and **C.** The five clusters identified in both sites are shown in **A** (SA) and **C** (US).
B. and **D.** Dot plots colored by the relative expression level of each marker (CD4, CD8, TRAV1-
2, CD161 or CD26) in tetramer-defined MR1T cells for each age group in **B** (SA) and **D** (US).
Red = highest expression, blue = lowest expression. **E.** and **F.** The proportion of each cluster of
tetramer-defined MR1T cells by age group is shown in **E** (SA) and **F** (US). Horizontal lines depict
the median and the error bars the 95% confidence interval. Mann-Whitney u tests were used to test
differences between groups.

Figure 4. Phenotypic analysis of functional MR1T cells at different ages. PBMC or CBMC from the US cohort were incubated overnight with *M. smegmatis*-infected A549 cells, uninfected A549 cells, or uninfected A549 cells and PMA/ionomycin. All cells were then stained with the MR1-5-OP-RU or MR1-6FP tetramers, followed by a live/dead discriminator and antibodies to TCRγδ, CD3, CD4, CD8, TRAV1-2, CD26 and CD161. ICS was then performed and the cells stained for TNF. Live, TCRγδ⁻CD3⁺MR1-5-OP-RU⁺ cells were gated and the TNF⁺ percentage determined in both the *M. smegmatis* stimulated and unstimulated conditions. The gating strategy is shown in Figure S3. **A.** Dot plots showing TNF expression by live, TCRγδ⁻MR1-5-OP-RU⁺ CD3⁺ cells in the same representative neonate, infant and adult as Figure 2A. **B.** Background subtracted frequencies of TNF⁺ MR1-5-OP-RU⁺ CD3⁺ cells in response to *M. smegmatis*-infected A549 cells as a percentage of total MR1/5-OP-RU⁺ CD3⁺ cells are shown. Mann-Whitney u-tests were used to test differences between groups. **C.** Background subtracted frequencies of TNF⁺ cells to PMA/ionomycin among different T cell subpopulations, 1) MR1T cells (CD3⁺TCRγδ⁻MR1-5-OP-RU⁺ cells); 2) γδ T cells (CD3⁺TCRγδ⁺MR1-5-OP-RU⁻ cells); 3) CD8⁺ T cells (CD3⁺TCRγδ⁻MR1-5-OP-RU⁻CD8⁺ cells); and 4) CD4⁺ T cells (CD3⁺TCRγδ⁻MR1-5-OP-RU⁻CD4⁺ cells). Wilcoxon-rank sum was used to test differences within the same cohort. **D.** and **E.** Frequencies of TNF⁺TRAV1-2⁺ MR1-5-OP-RU⁺ CD3⁺ and TNF⁺TRAV1-2⁻ MR1-5-OP-RU⁺ CD3⁺ cells in response to *M. smegmatis* infected A549 cells (**D**) or PMA/ionomycin stimulation (**E**) minus the background frequencies of TNF⁺TRAV1-2⁺/⁻ MR1-5-OP-RU⁺ CD3⁺ cells. Wilcoxon-rank sum was used to test differences within the same cohort.

Figure 5. Analysis of MR1T cell development by measuring maturation markers. PBMC or CBMC from the US were stained as in Figure 1 with the addition of an antibody to CD27. Live, CD3⁺MR1-5-OP-RU⁺ cells (gating strategy in Figure S2) were gated for the maturation pattern of S1, S2 and S3 described in (Koay et al., 2016). S1 = CD161⁻CD27⁻, S2=CD161⁻CD27⁺, S3=CD161⁺CD27⁺/⁻. **A.** Gating of MR1T cells for S1, S2 and S3 in a representative neonate, infant and adult. **B.** Frequencies of S1, S2, and S3 MR1T cells as a percentage of CD3⁺ MR1-5-OP-RU⁺ cells are shown for U.S. neonates, infants and adults. Bars represent the median and the error bars the 95% confidence interval.

Figure 6. Comparative phenotypic analysis of MR1T and non-MR1T cells or TRAV1-2⁺ and TRAV1-2⁻ MR1T cells at different ages. PBMC or CBMC from the US were stained as in Figure 1 with the addition of antibodies to either CD45RA (**A, B**), CCR7 (**C, D**), CD38 (**E, F**) or CD56 (**G, H**). The gating strategy is shown in Figure S2 and examples of the staining of each marker is shown in Figure S4. Frequencies of cells expressing each marker in non-MR1T cells (CD3⁺MR1-5-OP-RU⁻) or MR1T cells (CD3⁺MR1-5-OP-RU⁺) in each age group (**A, C, E, G**). The frequency of each marker expressed in TRAV1-2⁺ or TRAV1-2⁻ MR1T cells is shown in each age group (**B, D, F, H**). The median and 95% confidence intervals are shown in each graph. When the 95% confidence intervals do not overlap between conditions, those conditions are considered statistically significant.

A

	MR1/5-OP-RU	MR1/6FP	TRAV1-2	CD161
US Neonate	0.043%	0.016%	1.51%	4.07%
US Infant	0.53%	0.02%	3.45%	15.1%
US Adult	1.39%	0.018%	3.83%	41.6%

B

South Africa (SA)

United States (US)

Uganda (Ug)

MR1T among CD3+ T cells (%)

Phenotypically-defined   Tetramer-defined

● Neonates ● Infants ● Adolescents
● Neonates ● Infants ● Adults
● Infants ● Children ● Adults

C

South Africa (SA)

Neonates   Infants (10 weeks)   Adolescents

United States (US)

Neonates   Infants (1 year)   Adults

Uganda (Ug)

Infants <2 years   Children 2-5 years   Adults

■ MR1/5-OP-RU+   ■ MR1/5-OP-RU- CD8+   ■ MR1/5-OP-RU- CD8+ CD4+   ■ MR1/5-OP-RU- CD4+   ■ MR1/5-OP-RU- CD8- CD4-

A

	Neonate	Infant	Adult

M. smegmatis

Neonate: 19.4% | 13.9% | 66.7% | 0%
Infant: 54.4% | 32.9% | 9.71% | 2.94%
Adult: 73.4% | 15.1% | 11.0% | 0.5%

No Stimulation

Neonate: 35.2% | 0% | 64.8% | 0%
Infant: 91.3% | 0% | 8.66% | 0%
Adult: 91.1% | 0.55% | 8.3% | 0%

MR1/5-OP-RU

TNF

B

TNF+ Phenotypically-defined MAIT cells (%)

Neonates: p=0.0117

Infants: p=0.0020   Adults: p=0.0020

● MR1/5-OP-RU+   □ MR1/5-OP-RU-

A — CD45RA
B — CD45RA
C — CCR7
D — CCR7
E — CD38
F — CD38
G — CD56
H — CD56

Proportion of CD3 T cells (%)

Proportion of MR1T cells (%)

Neonates   Infants   Adults

• Non-MR1T Cells   □ MR1T Cells

• TRAV1-2-   □ TRAV1-2+