Skin microbiota dynamics of dairy cows affected by digital dermatitis

Juan Manuel Ariza[1*], Dörte Döpfer[2], Kenny Oberle[1], Kelly Anklam[2], Sophie Labrut[3],

Nathalie Bareille[1], Anne Relun[1], and Raphaël Guatteo[1]

[1] *INRAE, Oniris, BIOEPAR, 101 route de Gachet, 44300, Nantes, France*

[2] *School of Veterinary Medicine, University of Wisconsin, Madison 53706-1102.*

[3] Laboniris, Oniris, Nantes Atlantic College of Veterinary Medicine, Food Science and

Engineering, Nantes, France

*Corresponding author: Juan Manuel Ariza. Email: juan-manuel.ariza-

chacon@oniris-nantes.fr

Abstract

Bovine digital dermatitis (DD), is a highly prevalent disease among dairy cattle

characterized by ulcerative and painful lesions. While multiple management factors

are involved in the disease, its precise etiology remains uncertain and the

effectiveness of current control strategies remains highly variable. The major role of

Treponema spp. in the development of the disease is consistently recognized.

Nevertheless, it remains unclear how other bacterial communities are relevant to the

onset and progression of the disease, and how the skin microbiota is affected by the

environment during the course of the disease. The objective of this study was to

describe the dynamics of microbiota recovered from DD affected feet under field

conditions. This study described the diversity, structure, and composition of DD

lesion microbiota over 45 days according to different clinical and management

factors. The results of this investigation confirmed the existence of a specific skin

microbiota associated with DD lesions, dominated by *Treponema* spp. and very

different from the microbiota of healthy skin. Interestingly, the diversity and structure

of the microbiota in DD lesions did not vary with the footbath disinfectant or the

individual topical antibiotic treatments used. In addition, microbiotas from proliferative

lesions evidenced a different structure and diversity in comparison to non-proliferative

lesions. Our results confirm the major role of *Treponema* spp. And highlight the

potential role of *Mycoplasmopsis* spp. in the DD lesion onset. Further studies are

needed to confirm whether the clinical course of DD lesions is driven by a particular

microbiota and how that microbiota may induce disease.

Key words: Bovine digital dermatitis, Microbiota, Polymicrobial, Dynamics, Control

strategies.

Highlights

Multiple bacteria have been identified in DD lesions. However, many of these

microorganisms are inhabitants of the foot skin and the farm environment. For the

first time, the microbiota of DD lesions was monitored for 45 days under field

conditions to describe its evolution over time. The results of this investigation

highlighted a particular microbiota dominated by *Treponema* spp. present on the skin

of DD affected animals and highly different from those microbiotas of healthy skin.

The microbiota of DD lesions evolved over the study period and differential bacteria

were identified. Further studies are warranted to determine the role of the bacteria

composing these microbiotas on lesion onset and outcome.

Introduction

The increased prevalence of lameness and its impact on the welfare and productivity

of animals reflects an important concern facing modern dairy farming. One of the

main causes of infectious lameness is bovine digital dermatitis (DD)(1), a

multifactorial and polymicrobial disease spread across the world and characterized

by ulcerative and painful lesions which might persist as a chronic condition. The DD

lesions progress in dynamic ways conditioned by multiple environmental factors that

influence the spread of associated-pathogens and the integrity of the foot skin.

Currently, the control strategies implemented in dairy farms such as footbaths or

individual treatments, have demonstrated variable effectiveness in the healing and/or

prevention of DD lesions(2, 3).

Although the precise etiology of DD remains unclear, the previous maceration of the

foot skin and the presence of *Treponema* spp. have been consistently identified as

the major etiological components of DD(4–7). However, multiple other bacteria have

also been associated with DD infection in different studies, such as *Mycoplasma*

spp.(8, 9); *Fusobacterium necrophorum*(10); some Bacteroides, specifically

Porphyromonas levii(11, 12); different Proteobacteria, such as *Campylobacter* spp.

and *Dichelobacter nodosus*(13–15), and finally a broad range of other Firmicutes(16).

Nevertheless, the large part of these DD-associated pathogens (including the

Treponema spp.) are inhabitants of the normal foot skin, the rumen, and the

gastrointestinal tract of ruminants, and thereby, of the farm environment(17–19).

Together, all these findings have reinforced the concept of DD as a polymicrobial

disorder suggesting that a particular microbiota may drive the lesion environment and

affect its outcome.

The development of collective and individual treatment strategies has been focused

on the incriminate DD associated-pathogens, mainly against *Treponema* spp.

Nevertheless, these treatments may influence the structure of the microbial

communities that constitute the foot skin barrier. The disruption of the normal skin

microbiota can lead to dysbiosis and consequently induce colonization of DD-

associated pathogens and the consequent apparition of DD lesions. Hypothetically, a

successful control strategy should regulate microbiota for preventing or decreasing

the proliferation of the DD-associated pathogens.

Next-generation sequencing (NGS) had enlarged the perspectives to explore deeply

the dynamics of the entire skin microbiota, beyond the restrictions imposed by the

culture of pathogens or the pre-selection of targeted species. Through the

sequencing analysis of the 16S ribosomal RNA (rRNA), this investigation explored

the dynamics of DD microbiotas over 45 days using skin biopsies collected from 10

cows of 5 different dairy farms, and submitted to recognized treatment strategies

(individual topical treatment and/or footbath disinfectant)

Results

A clinical and microbial description of the lesions studied is presented in Table 1

according to the sample time. During the 45 days of follow-up, all 10 initial DD lesions

studied remained among the clinical stages M2, M4 or M4.1. None of the DD lesions

evidenced a transition to a healing or healthy status (M3-M0). Regarding the lesion

type recorded, only two additional proliferative lesions were recorded during the study

period. During the study period, 4 feet were treated with topic antibiotic treatment at

days 17 (n=2), 35 and 39, respectively. According to the histologic evaluation of

epidermal damage, the 10 samples of healthy skin were considered at score 1 and

most of the DD lesions were considered at score 3 (n=32), as expected (Table 1).

Considering the epidermal damage, while the highest score (EDS=3) was

systematically associated with bacterial invasion of spirochaetal origin (Figure 1),

lesion with medium score (EDS=2) where less frequently associated with bacterial

invasion (BIS=1 (n=3), BIS=2 (n=3), and BIS=3 (n=1)) (Table 1). Finally, lesions with

low score (EDS=1) were never associated with bacterial invasion.

Regarding the raw reads processing, 1571 non-chimeric unique amplicon sequence

variants (ASV) were identified. After removing low prevalence ASV occurring in less

than two samples, 1398 ASV were conserved. Finally, the phylogenetic tree-based

agglomeration conducted to the identification of 139 unique ASV including 8 uniquely

named taxa on the phylum level and 24 on the genus level (Figure S1). The set of

microbial proxies studied according to every sampling time are presented in Table 1.

For the within day of sample comparisons, no differences were found in any of the

microbial proxies studied at the start of the study (Healthy skin at day 0, DD lesion at

day 0), or during the study (day 15, day 30, and day 45). Otherwise, microbiota from

healthy skin showed significantly increased evenness (p = 0.001), diversity (p =

0.001), and dissimilarity (p = 0.027) compared to microbiota from DD lesions

sampled at day 0. The evenness of DD lesion microbiota on days 15 and 30 was

significantly increased compared with that on day 0. Furthermore, compared to

microbiota of DD lesions at day 0, microbiotas collected at day 45 have significantly

reduced dissimilarity and increased richness, evenness and diversity.

According to the clinical variables explored and the histological evaluation, no

differences in the richness were detected for any of the comparisons (Table 2). In this

study, we were unable to detect any significant impact on the evenness, diversity or

dissimilarity of the microbiota according to the usage or not of the individual topical

treatment and/or the footbath disinfectant. Otherwise, proliferative lesions presented

decreased dissimilarity and increased evenness and diversity in comparison to non-

proliferative lesions. The DD lesions considered at M2, M4 or M4.1 stages did not

differ between them in any of the microbial proxies studied. As expected, the

microbiota from healthy skin showed significantly increased evenness, diversity, and

dissimilarity compared to any other lesion. Similarly, microbiota from samples

considered at the lowest BIS and EDS scores, had noticeably higher evenness,

diversity, and dissimilarity (Table 2).

Regarding the composition of the microbiota and in terms of relative abundance, the

most frequent phylum found among the healthy skin samples were Firmicutes (49%),

Actinobacteria (15%), Bacteroidetes (13%), Spirochaetes (11%), and Proteobacteria

(5%). Otherwise, the most frequent phylum found in DD lesions at day 0 were

Spirochaetes (54%), Bacteroidetes (25%), Firmicutes (14%), and Tenericutes (3%)

(Figure S1).

A visual description of the structure of DD lesion microbiota over time is provided in

Figure S2. The microbiota from healthy skin (day 0) and DD lesion at day 45 showed

a more dispersed pattern compared to the other time points (Figure S2 A).

Nevertheless, when comparing all-time points together, no major visual differences or

cluster formation were evidenced (Figure S2 B).

Figure 2A shows the microbiota according to whether or not the footbath disinfectant

was used during the study period (DD samples from days 15, 30 and 45). No

distinctive clustering is observed. Regarding the use of individual topical treatment,

the treated samples seem to cluster separately with an obvious reduction of

Spirochaetes and, in particular, Fusobacteria were placed completely outside the

clusters (figure 2B). The scatter plot of the lesion type suggests a distinctive cluster

for proliferative and non-proliferative lesions (Figure 2C). Finally, based on the clinical

score, only the microbiota of healthy skin appears to separate from the cluster formed

by DD lesions. Nevertheless, each stage of the lesion seems to draw distinctive

scatter plots (Figure 2D). Otherwise, according to the histological evaluation, scores

0 and 1 seems to cluster separately from scores 2 and 3 when comparing bacterial

invasion scores (Figure S3 A). Likewise, score 1 and 2 seems to cluster separately

from score 3 when comparing epidermal damage scores (Figure S3 B).

Finally, the dynamics of DD microbiota composition were explored through a

differential abundance analysis (Figure 3). For the baseline comparison between

healthy skin and DD lesions at day 0, 40 ASVs significantly reduced their abundance

and 58 increased their abundance. Indeed, compared to healthy skin, the microbiota

of DD lesions at day 0 have increased abundance of *Treponema* spp. (~4 to ~28 FC),

Fretibacterium spp. (~5 FC), *Mycoplasmopsis* spp. (~6 FC) and *Proteus* spp. (~21

FC). Decreased abundances of *Campylobacter* spp. (~17 FC), other unidentified

Spirochaetes (~17 FC), and different genera of Proteobacteria (~4 to ~18 FC) and

Tenericutes (~5 FC). Changes in both directions were detected on different genera

and other unidentified Bacteroidetes (~-20 to ~10 FC) and Firmicutes (~-22 to ~22

FC) abundances. When comparing the abundances of DD samples from day 15 to

day 0, the abundance of only 18 ASV changed significantly. In detail, there was a

reduction in *Treponema* spp. (~16 FC) and *Proteus* spp. (~21 FC), and an increase in

Actinobacteria (~16 FC). Different genera and other unidentified Bacteriodetes (~ -5

to ~18 FC) and Firmicutes (~ -28 to ~17 FC) change in both directions. For the

comparison between day 30 and day 15, the abundance of 34 ASV changed

significantly. We noticed a reduction in *Oligella* spp. (~19 FC) and an increase in

Campylobacter spp. (~4 to ~26 FC) and other unidentified Spirochaetes (~34 FC),

Proteobacteria (~24 FC) and ASV (~40 FC). In particular, changes in both directions

were observed in *Treponema* spp. (~-25 to ~24 FC) and for different genera and

other unidentified Bacteriodetes (~ -24 to ~33 FC) and Firmicutes (~-20 to ~34 FC).

Finally, the abundance of 22 ASV changed significantly when comparing form day 45

to day 30 microbiota. We noticed a reduction in *Treponema* spp. (~6 to ~25 FC) and

an increase in *Oligella* spp. (~21 FC) and other unidentified Proteobacteria (~7 FC).

Changes in both directions were observed for different genera and other unidentified

Bacteriodetes (~-10 to ~26 FC) and Firmicutes (~ -34 to ~24 FC).

Discussion

The present investigation describes the differences between microbiotas from healthy

skin and those affected by DD lesions under field conditions. Additionally, for the first

time, the microbiota dynamics of DD lesions were investigated during a period of 45

days exploring the putative impact of recognized factors, such as the use of individual

and collective treatments. As expected, DD lesions microbiota were dominated by

Treponema spp. and their configuration was largely different from healthy skin in

terms of diversity, structure, and composition. The DD lesions microbiota also

evidenced significant changes in its structure, in particular the populations of

Treponema spp., *Oligella* spp., Spirochaetes, Bacteriodetes and Firmicutes

constantly changed over time. Although the footbath disinfectant and the individual

topical treatment used in this study have recognized DD healing potential based on

their antimicrobial properties, no major impact on microbiota characteristics was

associated with their use over time. Other important determinants, such as foot

hygiene or the farm environment, may have altered the effectiveness of these

strategies. In addition, we detect significant differences between non-proliferative and

proliferative lesions microbiotas. We recorded a partial recovery of the microbiota

diversity trough the 45 days of follow-up. Our findings reveal important dynamic

changes on the abundances of *Treponema* spp. during the study period. For the first

time, this investigation provided insight into how the microbiota of DD lesions may

change over time under field conditions.

Our findings support the polymicrobial nature of DD highlighted in previous

investigations using sequencing and culture–based approaches. It is in line with

findings from previous research signaling *Treponema* spp. as the main

microorganism related to DD lesions(5, 8, 9). As expected, the microbiotas gathered

in this study share a similar structure with microbiotas previously described on

healthy skin and DD lesions from dairy cattle, respectively(5, 20). In detail, the main

bacteria linked to DD lesions in previous investigations that were again identified by

our findings include: *Treponema* spp.*, Acholeplasma* spp.(5, 8, 9), *Porphyromonas*

spp., *Prevotella* spp.(12), *Corynebacterium* spp.(9), *Tissierella* spp., *Mycoplasmopsis*

spp.(8, 21) and some unclassified Bacteriodetes, Firmicutes and Proteobacteria(11,

12, 20). On the contrary, some bacteria previously associated with DD, such as

Dichelobacter nodosus(14, 22) was not linked to lesions in this study. Although

Fusobacterium spp. have been previously related to DD lesions(5, 8, 9), our findings

indicate that its presence seems to be related to a particular farm environment rather

than to the disease microbiota dynamics. The present study also found an over-

representation of *Fretibacterium* spp. in DD lesions compared to healthy skin, an

anaerobic bacterium not previously implicated DD but, like different *Treponema* spp,

related to human periodontitis(23). Our findings coincide with the results of a recent

meta-analysis revealing the main role of *Treponema* spp., *Mycoplasmopsis* spp. on

DD pathogenesis(24). Indeed, we recorded significant changes in the abundances of

a large number and diversity of genera in DD lesions compared to healthy skin.

Nevertheless, future studies are needed to elucidate which specific genera might be

related to disease occurrence, regional particularities of the sampled farms, or

sampling contamination. Similarly, we recorded consistent changes in the abundance

of *Oligella* spp. over time, a bacterium previously reported in ruminant milk

microbiota(25) but not yet associated with DD and which might be an active part of

the farm environment. However, as these are the first skin microbiota assembled

from dairy cattle in France, no regional comparison can be performed. Additionally,

the fact that previous studies have not linked this bacteria genus to DD lesions might

be explained by the original statistical procedures implemented in the analyses of our

present study. Indeed, the relative abundances of every sample were calculated after

the standardization of the observed taxonomic groups according to their variance and

not through procedures implemented previously such as the rough normalization

which may entail the loss of valuable information(26). More in detail, when using

rough normalization procedures for data analyses, information from samples with a

scarce number of sequences is excluded from the dataset. In specific cases, such as

healthy skin samples, the exclusion of this information may conduct to neglect the

healthy nature of the sample which could potentially have low numbers of sequences

in comparison to diseased samples. The within-cow design of this study offered the

possibility of exploring the dynamics of microbiotas according to the clinical evolution

of each animal. Indeed, the simple comparison between microbiotas from DD lesions

obtained from different animals at diverse states may underestimate or neglect the

impact of individual and environmental factors (i.e. foot conformation or barn hygiene)

on the clinical outcome of lesions and their microbial configuration.

One of the main limitations of this study was related to the difficulties encountered in

interpreting microbiotas obtained from an anatomical region which inevitably is in

close contact with the ground and thereby with dirtiness. Indeed, the distinction of

pathogens among environmental microorganisms is a challenge. Furthermore, due to

the nature of our molecular approach, our study not provide evidence of bacterial

viability. We have tried to reduce this limitation by the meticulous washing and

scrubbing procedures performed during the biopsy collection, which in turn may have

altered the skin microbiota. However, colonization of tissues by opportunistic

pathogens could be favored by the successive incisional biopsies performed during

the follow-up of the lesions. Moreover, depending on the lesion size successive

biopsies in a same location   could restraint the expected healing of the skin. Another

limitation of the present investigation was related to the small sample size studied.

Larger samples may highlight with more precision the benefit of control strategies for

DD in increasing the microbiota diversity of the skin. As several risk factors at the

farm level are involved with the disease, such as hygienic conditions, differences in

the microbial profile according to the farms were expected. However, the small

number of farms (n=5) and the small number of individuals sampled within farm

restrain the exploration of factors related to farm hygiene. In this study, the changes

in the microbiota allowed a clear distinction between the diseased (M2-4-4.1) and

healthy (M0) states, but the potential impact of individual or collective treatments on

the microbiota was not detected. In contrast to previous studies(27), for the present

investigation the potential utility of NGS tools to measure the potential effectiveness

of control strategies seemed inferior or at least different from standard observation

methodologies such as DD stages or lameness scoring. In terms of ethics, feasibility,

and more importantly precision, invasive methodologies (biopsies) are limited in large

samples and, as pointed above, entails a challenge for the interpretation. Another

point concerns the fact that the effect of both, the footbath disinfectant and the

individual topical treatment, could be highly affected by the exacerbated anti-

inflammatory response of the animals after each biopsy. Therefore, measuring local

immune responses may enhance the precision of the true effect of control

interventions. Regarding the histological evaluation of the samples, this additional

assessment clearly allowed the distinction of the bacterial invasion advance between

the samples. Although bacterial invasion (BIS) was in concordance with the clinical

evaluation of the samples and the dynamic evolution of the microbiota abundances,

the epidermal damage (EDS) was more difficult to interpret due to the potential effect

of the recurrent biopsy procedures.

Among the different clinical factors studied, the structure and diversity of microbiotas

from DD lesions were not affected by the usage of individual topical treatment. This

could be explained by the short and not persistent effect of this strategy over

time(28). Otherwise, the findings of this study indicate noticeable differences between

the microbiotas of non-proliferative and proliferative DD lesions. The presence of

proliferative tissues is associated with chronical processes and thereby to lesions

prone to relapse(29). The hyper-proliferation of tissues surrounding the lesion may

alter the environment of the skin and the topical activity of footbath and individual

topical treatment, thus influencing a distinctive microbiota(30–32). Recently,

Treponemas spp. have also been involved in some non-infectious diseases inducing

lameness(33). Further studies are needed to evaluate the role of microbiota in the

dynamics of those non-infectious lesions that are related to the same risk factors

recognized for DD. Other recognized factors influencing the occurrence of DD lesions

at the individual level, such as parity or days in milk, were not studied due to the

small number of cows evaluated at every time point and the large variability among

them.

Beyond the recognition of the pathogenic role of some specific bacteria, exploring the

microbiotas from DD lesions at the community level may highlight how the dynamics

of skin microbiota determine the persistence and the occurrence of DD lesions.

Indeed, the 16*S* rRNA gene analyses may capture broad shifts in community

diversity over time, but with limited resolution and lower sensitivity compared to

metagenomic data. Nevertheless, the results of this investigation reveal the dynamic

nature of microbiotas from DD lesions and highlight the potential importance of

specific bacteria in the progress of the lesions over time. Future studies could benefit

from shotgun sequencing tools, improving sensitivity in recognizing pathogens and

inhabitants and defining their specific role in disease. Therefore, further investigations

linking the skin microbiota to different herd management practices might enhance the

current understanding of the disease and promote the development of successful

control strategies.

In conclusion, we have explored the dynamics of the skin microbiota in feet affected

with DD lesions. The diversity and structure of microbiotas from DD lesions were

different from those of healthy skin but did not vary according to the usage of the

footbath disinfectant or the individual topical treatment. The microbiota of proliferative

lesions displayed a different structure and diversity compared to non-proliferative

lesions. Concerning the dynamics of DD lesion progression, we confirmed the

putative role of *Treponema* spp. and open new questions about the potential role of

Mycoplasmopsis spp., *Oligella* spp. and other Spirochaetes, Bacteriodetes and

Firmicutes. Further studies are needed to understand how risk factors associated

with DD at the farm and individual level may affect the interactions between the

bacteria comprising the skin microbiota, and how these interactions may determine

the clinical evolution of DD lesions.

Methods

Ethics statement

The research protocol was reviewed and approved by the Ethics Veterinary

Committee in Clinical Research and Epidemiology from the Veterinary School of

Nantes, ONIRIS, France (CERVO, France) (registered number: CERVO-2016-12-V).

Study population

Five dairy farms participating simultaneously in a controlled clinical trial were included

in the study(34). The herds were composed exclusively of Holstein cows to limit the

putative breed effect. The herds were housed in cubicles and on average 90 cows

were milked twice a day in a rotatory or conventional milking-parlor. The prevalence

of active DD lesions at the start of the study ranged from 20% to 40%.

During the study period (45 days), farmers were allowed to detect and then treat

severe cases of DD by using 2 applications of oxytetracycline (30 mg/ml)

(Oxytetrin™, MSD Animal Health/Intervet, France) 2 days apart. For the current

investigation, from each farm, the owners have chosen two animals suffering from a

rear DD lesion. Therefore, the study population consisted of 10 different cows

affected by DD. For each cow, a single lesion was recorded and follow-up during the

study period. Five of the cows did not receive any collective disinfectant treatment on

the foot suffering the DD lesion recorded and the remaining 5 feet received a

collective disinfectant treatment through a footbath (Pink-Step™, Qalian). The

frequency of administration of the footbath was of 2 days (4 consecutive milkings)

every week of the study period. The cows included in the study were between their

first and fifth parity, and their days in milk ranged from 3 to 378 at the start of the

study.

Follow-Up and Data Collection

Farms were visited by 2 investigators trained by practical lessons to practice the

biopsies and score the lesions. All lesions sampled were photographed and scored

using the M5 score system(32). At every sample time, we recovered information

about the eventual administration of any concomitant treatment between the

samplings. To perform the biopsies every cow was carefully restrained in a trimming

chute. The skin area was washed with water and then local anesthesia was provided

using Procaine 2% (Procamidor™, Axience, France). Thereafter, the area was rinsed

and brushed with a PBS solution before performing the sample with a sterile biopsy

punch (6 mm). The incisional samples were again washed with sterile PBS solution,

cut tangentially into two approximately equal parts. One half was stored in 10%

neutral buffered formalin solution for histological evaluation, and the other half was

stored in sterile 2.0 ml microcentrifuge tubes at -20°C for DNA extraction and

sequencing analysis. Finally, an aerosol bandage of aluminum (Aluspray™,

Vetoquinol, France) was sprayed directly over the incisional biopsy wound of each

cow. Additionally, in the case of pain detected by farmers, NSAIDs were provided

after the biopsy sampling.

For every cow, five skin biopsies of the affected foot were performed at 4 different

times. Only for the first sampling, two biopsies were taken at the same foot, one

sample on the healthy skin, and the second on the DD lesion. The biopsy sampling

start immediately before the application of first footbath. Thereafter, 3 subsequent

biopsies of DD lesions were taken every 15 days. This time interval was chosen to

allow the partial healing of the skin. The location of each biopsy lesion intended to be

taken as close as possible to the previous sampling sites. In total 50 biopsies were

recovered.

Histopathology

The paraffin sections of every tissue sample were stained with *Hematoxylin-Eosin-*

Safran (HES) for the general histopathological evaluation and with *Warthin Starry*

Silver (WSS) to evaluate the presence and degree of spirochaetal invasion. The

degree of epidermal damage (EDS) was assessed according to a modified score

system described in a previous study(14). A score of 1 was defined as normal skin or

presenting mild acanthosis and hyperkeratosis without any other significant

microscopic lesions associated and without bacteria. Score 2 was defined as skin

with marked acanthosis and hyperkeratosis, without erosion or ulceration, with or

without pustules, with or without hemorrhages and erythrocytic crusts with or without

dermal fibrosis. Finally, score 3 was defined as skin presenting a segmental

epidermis, localized to extensive, necrotizing to necrosuppurative epidermis with

balloon degeneration of epithelial cells, necrotic vasculitis and intralesional bacteria

(spirochaetes, bacilli, and coccobacillus). Otherwise, the total bacterial invasion score

(BIS) encompassed also the spirochaetes identified by the WWS staining. The score

0 represented the absence of invasive bacteria; score 1, a low number of invasive

bacteria; score 2, a moderate number of invasive bacteria; score 3, a high number of

invasive bacteria.

DNA Extraction, 16S rRNA amplification, illumina MiSeq sequencing.

Bacterial DNA extraction was enhanced adding to every sample 20 µl of proteinase

K, 180 µl of tissue lysis buffer, and 40 µl of lysozyme (Qiagen, Valencia, CA, USA).

Thereafter, all the samples were incubated for 12 h at 56°C and then processed

directly for DNA extraction using the Powerlyzer Powersoil Kit (Quiagen, Valencia,

CA) according to the manufacturer's instructions. Extractions were performed in

rounds of 10 samples and in every round, an additional empty tube was processed in

parallel to serve as a negative extraction control. Finally, the resulting supernatant of

each sample was transferred to a labeled microcentrifuge tube and stored at -20°C

The Quant-iT dsDNA Assay Kit (Thermo Fisher Scientific, MA, USA) was used to

measure DNA concentrations prior to PCR amplification. The V4 region of the

bacterial 16S rRNA gene was amplified by PCR, using Illumina Nextera XTIndex

Primer 1 (N7xx) and Primer 2 (S5xx), where barcodes were unique to each sample.

The PCR protocol consisted of an initial denaturation at 95 □C for 10 minutes

followed by 40 cycles of 95 □C for 30 seconds and 60 □C for 1 minute and the real-

time fluorescence data acquisition occurred at the end of each annealing/extension

phase. All PCR assays use the same PCR cycling conditions allowing parallel testing

of all PCR assays. The amplicons were extracted and purified from the gel using the

Zymo gel extraction kit according to the manufacturer's instructions and quantified

using the Quant-iT dsDNA Assay Kit (Thermo Fisher Scientific, MA, USA). Finally,

the purified amplicons were sequenced using the MiSeq reagent kit v2 (13 cycles) on

the MiSeq platform (Illumina, Inc., CA, USA).

Sequencing data processing

The generated 16S rRNA raw gene sequences were preprocessed to remove any

remaining, barcodes or adapters using Trimmomatic(35). Raw reads were then

processed using R(36), implementing the established DADA2 workflow(37). The

amplicon sequence variants (ASV) in each sample were inferred using the "divisive

amplicon denoising algorithm", a parameterized model of substitution errors to

distinguish sequencing errors from real biological variation(38,39). The inferred

forward and reverse sequences were then merged, removing sequences that did not

perfectly overlap and subsequently chimeras were removed. To each ASV a

taxonomic label was assigned through a Bayesian classifier using the Ribosomal

Database Project (RDP) v18 training set(40, 41). The phylogenetic relatedness of the

ASV identified was calculated using the R package *decipher* (42) and, spurious ASV

not occurring in two or fewer samples were removed. Finally, the ASV were

agglomerated according to a cophenetic distance inferior to 0.03. Sequencing data

have been deposited in the MG-RAST (https://www.mg-rast.org) database and are

publically available at: https://www.mg-rast.org/linkin.cgi?project=mgp80659.

Statistical analyses

The statistical analyses were performed using R (36). The analysis strategy

consisted of the following 4 steps: i) Describe the dynamics of the skin microbiota on

feet with DD lesions using different microbial proxies of diversity and community

structure, ii) explore whether the studied microbiotas were affected by different

factors that have been previously identified as impacting the clinical course of DD

lesions (clinical variables), and iii) identify differential ASV that may affect the course

of DD lesions over time.

The clinical variables explored were the usage of the footbath disinfectant, the usage

or not of individual topical antibiotic treatments, the M5 stages recorded, the lesion

type (proliferative or not), and the histopathologic evaluation (degree of bacterial

invasion and lesion scoring). The microbiotas were charcaterized using four different

microbial proxies including: richness (count of observed ASV), evenness (Pielou

index), diversity (Shannon index), and dissimilarity (43). More in detail, while the

Pielou index measures the evenness of ASV abundances and how abundances are

distributed, the Shannon index determines the microbial diversity controlling by the

richness (44). The dissimilarity was explored by calculating the Bray-Curtis distances

according to the different clinical variables studied. All microbial proxies were compared between the clinical variables studied using Kruskal-Wallis and pairwise Mann–Whitney U tests. The impact of the clinical variables on the structure of microbiotas over time was visualized using the Bray-Curtis distances and nonmetric multidimensional scaling ordination (NMDS). In the scatter plots created, high dissimilarity of microbiotas was represented by long distances between dots, whereas microbiotas with a similar composition were clustered together.

Finally, the dynamic on DD microbiota composition was explored by identifying specific ASV changing over time through a differential abundance analysis using the R package *DEseq2* (45). *DEseq2* uses a negative binomial distribution and implement the variance stabilization transformations of the ASV read counts identified in the raw data set to account for overdispersion. A single model was fitted including the main effects of the sample time (day 0, 15, 30 and 45) and independent hypothesis weighting (IHW)(46) was used to corrected for multiple comparisons such as the dependencies between the multiple samples within the same cow. Therefore, significant log 2 fold change (FC) (p-value ≤ 0.1) on the ASV counts were explored between the microbiotas of: i) healthy skin Vs. of DD lesions at day 0, ii) DD lesions at day 0 Vs. DD lesions after 15 days, iii) DD lesions at day 15 Vs. DD lesions after 30 days and, iv) DD lesions at day 30 Vs. DD lesions after 45 days. All figures were created with the R packages *ggplot2* and *plotly* (47, 48).

References

1. Laven RA, Lawrence KR. 2006. An Evaluation of the Seasonality of Veterinary Treatments for Lameness in UK Dairy Cattle. J Dairy Sci 89:3858–3865.
2. Apley MD. 2015. Clinical Evidence for Individual Animal Therapy for Papillomatous Digital Dermatitis (Hairy Heel Wart) and Infectious Bovine Pododermatitis (Foot Rot). Vet Clin North Am Food Anim Pract 31:81–95.

3.　Ariza JM, Relun A, Bareille N, Oberle K, Guatteo R. 2017. Effectiveness of collective treatments in the prevention and treatment of bovine digital dermatitis lesions: A systematic review. J Dairy Sci 100:7401–7418.

4.　Capion N, Boye M, Ekstrøm C, Dupont K, Jensen TK. 2013. A Study into Digital Dermatitis Transmission and Bacterial Associated Pathological Changes Involved in the Disease. Open J Vet Med 03:192–198.

5.　Zinicola M, Lima F, Lima S, Machado V, Gomez M. 2015. Altered Microbiomes in Bovine Digital Dermatitis Lesions , and the Gut as a Pathogen Reservoir 1–23.

6.　Krull AC, Cooper VL, Coatney JW, Shearer JK, Gorden PJ, Plummer PJ. 2016. A Highly Effective Protocol for the Rapid and Consistent Induction of Digital Dermatitis in Holstein Calves. PLoS One 11:e0154481.

7.　Wilson-Welder JH, Nally JE, Alt DP, Palmer M V., Coatney J, Plummer P. 2017. Experimental Transmission of Bovine Digital Dermatitis to Sheep: Development of an Infection Model. Vet Pathol 030098581773657.

8.　Krull AC, Shearer JK, Gorden PJ, Cooper VL, Phillips GJ, Plummer PJ. 2014. Deep Sequencing Analysis Reveals the Temporal Microbiota Changes Associated with the Development of Bovine Digital Dermatitis. Infect Immun 82:3359–3373.

9.　Nielsen MW, Strube ML, Isbrand A, Al-Medrasi WDHM, Boye M, Jensen TK, Klitgaard K. 2016. Potential bacterial core species associated with digital dermatitis in cattle herds identified by molecular profiling of interdigital skin samples. Vet Microbiol 186:139–149.

10.　Moe KK, Yano T, Misumi K, Kubota C, Nibe K, Yamazaki W, Muguruma M, Misawa N. 2010. Detection of antibodies against fusobacterium necrophorum and porphyromonas levii-like species in dairy cattle with papillomatous digital dermatitis. Microbiol Immunol 54:338–346.

11.　Yano T, Moe KK, Yamazaki K, Ooka T, Hayashi T, Misawa N. 2010. Identification of candidate pathogens of papillomatous digital dermatitis in dairy cattle from quantitative 16S rRNA clonal analysis. Vet Microbiol 143:352–362.

12.　Berry SL, Read DH, Walker RL, Famula TR. 2010. Clinical, histologic, and bacteriologic findings in dairy cows with digital dermatitis (footwarts) one month after topical treatment with lincomycin hydrochloride or oxytetracycline hydrochloride. J Am Vet Med Assoc 237:555–560.

13. Döpfer D, Koopmans a, Meijer F a, Szakáll I, Schukken YH, Klee W, Bosma
RB, Cornelisse JL, van Asten a J, ter Huurne a a. 1997. Histological and
bacteriological evaluation of digital dermatitis in cattle, with special reference to
spirochaetes and Campylobacter faecalis. Vet Rec 140:620–623.

14. Rasmussen M, Capion N, Klitgaard K, Rogdo T, Fjeldaas T, Boye M, Jensen
TK. 2012. Bovine digital dermatitis: Possible pathogenic consortium consisting
of Dichelobacter nodosus and multiple Treponema species. Vet Microbiol
160:151–161.

15. Knappe-Poindecker M, Gilhuus M, Jensen TK, Vatn S, Jørgensen HJ, Fjeldaas
T. 2014. Cross-infection of virulent Dichelobacter nodosus between sheep and
co-grazing cattle. Vet Microbiol 170:375–382.

16. Santos TM a, Pereira R V., Caixeta LS, Guard CL, Bicalho RC. 2012. Microbial
diversity in bovine Papillomatous digital dermatitis in Holstein dairy cows from
upstate New York. FEMS Microbiol Ecol 79:518–529.

17. Klitgaard K, Nielsen MW, Ingerslev HC, Boye M, Jensen TK. 2014. Discovery
of bovine digital dermatitis-associated Treponema spp. in the dairy herd
environment by a targeted deep-sequencing approach. Appl Environ Microbiol
80:4427–4432.

18. Zinicola M, Higgins H, Lima S, Machado V, Guard C, Bicalho R. 2015. Shotgun
Metagenomic Sequencing Reveals Functional Genes and Microbiome
Associated with Bovine Digital Dermatitis. PLoS One 10:e0133674.

19. Nascimento LV, Mauerwerk MT, dos Santos CL, Filho IB, Birgel Júnior EH,
Sotomaior CS, Madeira HMF, Ollhoff RD. 2015. Treponemes detected in digital
dermatitis lesions in Brazilian dairy cattle and possible host reservoirs of
infection. J Clin Microbiol JCM.03586-14.

20. Santos TMA, Pereira R V., Caixeta LS, Guard CL, Bicalho RC. 2012. Microbial
diversity in bovine Papillomatous digital dermatitis in Holstein dairy cows from
upstate New York. FEMS Microbiol Ecol 79:518–529.

21. Caddey B, Orsel K, Naushad S, Derakhshani H, Buck J De. Identi fi cation and
Quanti fi cation of Bovine Digital Dermatitis- Associated Microbiota across
Lesion Stages in Feedlot Beef.

22. Knappe-Poindecker M, Gilhuus M, Jensen TK, Klitgaard K, Larssen RB,
Fjeldaas T. 2013. Interdigital dermatitis, heel horn erosion, and digital
dermatitis in 14 Norwegian dairy herds. J Dairy Sci 96:7617–7629.

23.   Szafranski SP, Wos-Oxley ML, Vilchez-Vargas R, Jáuregui R, Plumeier I,
Klawonn F, Tomasch J, Meisinger C, Kühnisch J, Sztajer H, Pieper DH,
Wagner-Döbler I. 2015. High-Resolution Taxonomic Profiling of the Subgingival
Microbiome for Biomarker Discovery and Periodontitis Diagnosis. Appl Environ
Microbiol 81:1047–1058.

24.   Caddey B, De Buck J. 2021. Meta-Analysis of Bovine Digital Dermatitis
Microbiota Reveals Distinct Microbial Community Structures Associated With
Lesions. Front Cell Infect Microbiol 11:1–12.

25.   Li Z, Wright ADG, Yang Y, Si H, Li G. 2017. Unique bacteria community
composition and co-occurrence in the milk of different ruminants. Sci Rep 7:1–
9.

26.   McMurdie PJ, Holmes S. 2014. Waste Not, Want Not: Why Rarefying
Microbiome Data Is Inadmissible. PLoS Comput Biol 10.

27.   Gotoh Y, Chiba K, Sekiyama Y, Okada K, Hayashi T, Misawa N. 2020. 16S
rRNA-based amplicon analysis of changes in the bacterial population in the
lesions of papillomatous digital dermatitis in dairy cattle after topical treatment
with allyl isothiocyanate. Microbiol Immunol 64:416–423.

28.   Jacobs C, Orsel K, Mason S, Barkema W. 2018. Comparison of effects of
routine topical treatments in the milking parlor on digital dermatitis lesions. J
Dairy Sci 101:5255–5266.

29.   Biemans F, Bijma P, Boots NM, de Jong MCM. 2017. Digital Dermatitis in dairy
cattle: The contribution of different disease classes to transmission. Epidemics
https://doi.org/10.1016/j.epidem.2017.12.007.

30.   Gomez A, Cook NB, Bernardoni ND, Rieman J, Dusick a F, Hartshorn R,
Socha MT, Read DH, Döpfer D. 2012. An experimental infection model to
induce digital dermatitis infection in cattle. J Dairy Sci 95:1821–30.

31.   Gomez A, Cook NB, Rieman J, Dunbar KA, Cooley KE, Socha MT, Dopfer D.
2015. The effect of digital dermatitis on hoof conformation. J Dairy Sci 98:927–
936.

32.   Berry SL, Read DH, Famula TR, Mongini A, Döpfer D. 2012. Long-term
observations on the dynamics of bovine digital dermatitis lesions on a
California dairy after topical treatment with lincomycin HCl. Vet J 193:654–658.

33.   Bay V, Griffiths B, Carter S, Evans NJ, Lenzi L, Bicalho RC, Oikonomou G.
2018. 16S rRNA amplicon sequencing reveals a polymicrobial nature of

complicated claw horn disruption lesions and interdigital phlegmon in dairy
cattle. Sci Rep 8:1–12.

34.   Ariza JM, Bareille N, Lehebel A, Oberle K, Relun A, Guatteo R. 2019.
Evaluation of a biocide footbath solution in the occurrence and healing of digital
dermatitis lesions in dairy cows: A clinical trial. Prev Vet Med 163:58–67.

35.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for
Illumina sequence data. Bioinformatics 30:2114–2120.

36.   R Core Team. 2022. R: A Language and Environment for Statistical
Computing. Vienna, Austria.

37.   Callahan BJ, Sankaran K, Fukuyama JA, Mcmurdie PJ, Holmes SP, Charlop-
powers Z. 2018. Bioconductor workflow for microbiome data analysis: from
raw reads to community analyses [version 1; referees: 3 approved]
Referee Status: 1–44.

38.   Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should
replace operational taxonomic units in marker-gene data analysis. ISME J
11:2639–2643.

39.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP.
2016. DADA2: High-resolution sample inference from Illumina amplicon data.
Nat Methods 13:581–583.

40.   Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should
replace operational taxonomic units in marker-gene data analysis. ISME J
11:2639–2643.

41.   Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-
Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. 2009. The
Ribosomal Database Project: Improved alignments and new tools for rRNA
analysis. Nucleic Acids Res 37:141–145.

42.   Wright ES. 2015. DECIPHER: Harnessing local sequence context to improve
protein multiple sequence alignment. BMC Bioinformatics 16:1–14.

43.   McMurdie PJ, Holmes S. 2013. Phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. PLoS One 8.

44.   Jost L. 2010. The relation between evenness and diversity. Diversity 2:207–
232.

45.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol 15:1–21.

46.   Ignatiadis N, Klaus B, Zaugg JB, Huber W. 2016. Data-driven hypothesis
612   weighting increases detection power in genome-scale multiple testing. Nat
613   Methods 13:577–580.

614   47.   Wickham H. 2009. Elegant graphics for data analysisMedia. Springer New
615   York, New York, NY. http://had.co.nz/ggplot2/book.

616   48.   Sievert C, Parmer C, Hocking T, Chamberlain S, Ram K, Corvellec M, Pedro D.
617   2016. plotly: Create Interactive Web Graphics via "plotly.js."

Acknowledgments

The authors acknowledge the farmers and veterinarians who have contributed to this study. Additionally, we thank G. Puel (Oniris, Nantes, France) for helpful participation as investigator, for his time and motivation. This study was supported as part of a Ph.D. studentship by Qalian (Neovia group, Segré, France) and the "Association Nationale de la Recherche et de la Technologie" (ANRT, Paris, France).

Author contributions

J.M.A., D.D., K.O., A.R., N.B., and R.G. designed the study. J.M.A., S.L., D.D., and K.A. analyzed the data. J.M.A. and R.G. interpreted results. J.M.A. and R.G. drafted the manuscript. All authors revised critically the manuscript for important intellectual content and approved the final version to be published. The authors acknowledge the farmers and veterinarians who have contributed to this study.

Figure Legends

Figure 1. Skin biopsy from a dairy cow with an active bovine digital dermatitis lesion (M2). Colonizing bacteria within surface lesions and defined intra-lesional spirochetes within inflammatory loci are easily evidenced with the Warthin Starry silver staining.

Figure 2. Nonmetric multidimensional scaling (NMDS) scatterplot of Bray-Curtis dissimilarities of the skin microbiotas from feet suffering bovine digital dermatitis (DD)

lesions. Microbiotas are visualized by dots representing the amplicon sequence variants (ASV) present in the samples according to the usage of the footbath disinfectant (A), the usage of individual topical antibiotic treatments (B), the lesion type (C) and the M5 score lesion score (D). The size of the dots represents the mean relative ASV abundance and are colored according to their taxonomic classification at the Phylum level.

Figure 3. Differential abundance of amplicon sequence variants (ASV) present in skin microbiotas from feet suffering bovine digital dermatitis (DD) lesions. (**A**) DD lesions versus healthy skins at day 0, (**B**) DD lesions at day 15 versus DD lesions at day 0, (**C**) DD lesions at day 30 versus DD lesions at day 15, (**D**) DD lesions at day 45 versus DD lesions at day 30. Bubbles represent the ASV classified at the genus level (x-axis) that showed noticeably changes ($p\ value$ < 0.1) on a log2-scale (y-axis). The size of the bubbles represents the mean relative ASV abundance and are colored according to their phylum-level classification. Unclassified ASV at the genus level are gruoped at the Phylum level (*).

Figure S1. Relative abundances of the ASV present in the samples according to the sampling day. Taxonomic classification at the phylum level (A) and at the genus level (B).

Figure S2. The composition of skin microbiotas from feet suffering bovine digital dermatitis (DD) before and across the 45 days of study follow-up. Microbiotas are visualized by dots representing the amplicon sequence variants (ASV) identified in the samples. The size of the dots represents the mean relative ASV abundance and are colored according to their taxonomic classification at the Phylum level. The ordination was constructed using a Bray-Curtis matrix and Nonmetric multidimensional scaling (NMDS).

Figure S3. Nonmetric multidimensional scaling (NMDS) scatterplot of Bray-Curtis

dissimilarities of the skin microbiotas from feet suffering bovine digital dermatitis (DD)

lesions. Microbiotas are visualized by dots representing the amplicon sequence

variants (ASV) present in the samples according to the histological evaluation for

bacterial invasion (A) and epidermal damage (B). The size of the dots represents the

mean relative ASV abundance and are colored according to their taxonomic

classification at the Phylum level.

Table 1. Clinical variables and microbial proxies studied in 10 dairy cows with bovine digital dermatitis lesions followed over 45 days.

Variables Studied	All samples (n=10)				
	Healthy Skin	DD Lesions	DD Lesions		
	Day 0		Day 15	Day 30	Day 45
Clinical evolution					
Lesion stage					
M0	10	-	-	-	-
M1	-	-	-	-	-
M2	-	7	7	7	3
M3	-	-	-	-	-
M4	-	3	-	-	3
M4.1	-	-	3	3	4
Lesion Type					
Proliferative	-	4	5	6	5
Histopathology					
Bacterial Invasion Score					
0	10	-	1	2	1
1	-	3	2	1	1
2	-	4	3	1	-
3	-	3	4	6	8
Epidermal Damage Score					
1	9	-	1	-	1
2	1	7	1	-	2
3	-	3	8	10	7
Microbial proxies					
Richness	476.5 (368.57)	262.30 (116.90)	291.10 (141.08)	320.50 (170.92)	404.90§ (126.68)
Evenness	0.80 (0.08)	0.43* (0.13)	0.53§ (0.16)	0.51§ (0.14)	0.60§ (0.10)
Diversity	4.66 (0.72)	2.39* (0.84)	2.94 (0.98)	2.92 (0.98)	3.58§ (0.76)

Dissimilarity	0.37 (0.14)	0.26* (0.06)	0.24 (0.05)	0.25 (0.08)	0.19§ (0.05)

[a]All variables are presented as mean (standard deviation).
*Noticeable differences between the DD lesions and healthy skin from day 0.
§ Noticeable differences between the DD lesions from days 15, 30 or 45 and DD lesions on day 0.

Table 2. Effect of different clinical variables on the microbial proxies studied in 10 dairy cows with bovine digital dermatitis lesions followed over 45 days.

Microbial proxies[1]	[2]Footbath disinfectant treatment			[2]Individual topical treatment			[2]Lesion Type			Lesion stage				Bacterial Invasion Score				Epidermal Damage Score		
	Untreated (n=20)	Treated (n=20)	p	Untreated (n=34)	Treated (n=6)	p	Non-proliferative (n=20)	Proliferative (n=20)	p	Healthy Skin (n=10)	M2 (n=23)	M4 (n=6)	M4.1 (n=11)	0 (n=14)	1 (n=7)	2 (n=8)	3 (n=21)	1 (n=11)	2 (n=7)	3 (n=32)
Richness	291.87 (128.92)	385.80 (159.78)	0.08	309.00 (148.33)	380.33 (119.31)	0.16	300.50 (148.30)	338.9 (143.37)	0.31	476.50 (368.57) a	315.22 (150.61) a	413.17 (174.67) a	278.09 (99.85) a	481.93 (320.75) a	322.86 (167.15) a	312.88 (152.41) a	287.76 (108.09) a	480.91 (358.41) a	346.29 (150.05) a	307.47 (136.36) a
Evenness	0.53 (0.13)	0.56 (0.15)	0.71	0.52 (0.14)	0.49 (0.16)	0.51	0.46 (0.16)	0.57 (0.10)	0.02	0.80 (0.08) a	0.49 (0.16) b	0.56 (0.17) b	0.55 (0.09) b	0.73 (0.17) a	0.46 (0.25) ab	0.48 (0.10) b	0.54 (0.10) b	0.75 (0.16) a	0.56 (0.18) ab	0.52 (0.14) b
Diversity	2.99 (0.89)	3.30 (0.98)	0.36	2.97 (0.96)	2.89 (1.02)	0.95	2.61 (1.06)	3.30 (0.72)	0.01	4.66 (0.72) a	2.80 (1.01) b	3.37 (1.22) b	3.06 (0.65) b	4.29 (1.07) a	2.67 (1.59) ab	2.72 (0.63) b	3.06 (0.73) b	4.38 (1.11) a	3.26 (1.17) ab	2.93 (0.91) b
Dissimilarity	0.25 (0.06)	0.25 (0.06)	0.90	0.27 (0.06)	0.22 (0.05)	0.07	0.28 (0.06)	0.24 (0.06)	0.02	0.38 (0.13) a	0.27 (0.06) b	0.23 (0.06) b	0.23 (0.06) b	0.35 (0.13) a	0.25 (0.04) ab	0.26 (0.05) ab	0.25 (0.06) b	0.38 (0.13) a	0.24 (0.04) b	0.26 (0.06) b

[1]All measures are presented as mean (standard deviation).
[2]Excluding initial samples of heathy skin

Means within the same comparison with different subscripts are noticeably differences from one another (p < 0.05).

Figure 1. Skin biopsy from a dairy cow with an active bovine digital dermatitis lesion (M2). Colonizing bacteria within surface lesions and defined intra-lesional spirochetes within inflammatory loci are easily evidenced with the Warthin Starry silver staining.

Figure 2. Nonmetric multidimensional scaling (NMDS) scatterplot of Bray-Curtis dissimilarities of the skin microbiotas from feet suffering bovine digital dermatitis (DD) lesions. Microbiotas are visualized by dots representing the amplicon sequence variants (ASV) present in the samples according to the usage of the footbath disinfectant (A), the usage of individual topical antibiotic treatments (B), the lesion type (C) and the M5 score lesion score (D). The size of the dots represents the mean relative ASV abundance and are colored according to their taxonomic classification at the Phylum level.

Figure 3. Differential abundance of amplicon sequence variants (ASV) present in skin microbiotas from feet suffering bovine digital dermatitis (DD) lesions. (**A**) DD lesions versus healthy skins at day 0 ,(**B**) DD lesions at day 15 versus DD lesions at day 0, (**C**) DD lesions at day 30 versus DD lesions at day 15, (**D**) DD lesions at day 45 versus DD lesions at day 30. Bubbles represent the ASV classified at the genus level (x-axis) that showed noticeably changes (*p value* < 0.1) on a log2-scale (y-axis). The size of the bubbles represents the mean relative ASV abundance and are colored according to their phylum-level classification. Unclassified ASV at the genus level are gruoped at the Phylum level (*).