1   **Title:** Susceptible host availability modulates climate effects on dengue dynamics

2   **Authors:** Nicole Nova[1*], Ethan R. Deyle[2,3], Marta S. Shocket[1,4],

3   Andrew J. MacDonald[1,5], Marissa L. Childs[6], Martin Rypdal[7],

4   George Sugihara[2], and Erin A. Mordecai[1]

5   **Affiliations:** [1]Department of Biology, Stanford University, Stanford, CA, USA.

6   [2]Scripps Institution of Oceanography, University of California San

7   Diego, La Jolla, CA, USA.

8   [3]Department of Biology, Boston University, Boston, MA, USA.

9   [4]Department of Ecology and Evolutionary Biology, University of

10   California, Los Angeles, CA, USA.

11   [5]Earth Research Institute & Bren School of Environmental Science and

12   Management, University of California Santa Barbara, Santa Barbara,

13   CA, USA.

14   [6]Emmett Interdisciplinary Program in Environment and Resources,

15   Stanford University, Stanford, CA, USA.

16   [7]Department of Mathematics and Statistics, UiT The Arctic University

17   of Norway, Tromsø, Norway.

18   ***Correspondence email:** nicole.nova@stanford.edu

19   **Keywords:** Arbovirus, climate, dengue, empirical dynamic modeling, forecasting,

20   vector-borne disease, rainfall, susceptible population size, temperature

Abstract

Experiments and models suggest that climate affects mosquito-borne disease transmission. However, disease transmission involves complex nonlinear interactions between climate and population dynamics, which makes detecting climate drivers at the population level challenging. By analyzing incidence data, estimated susceptible population size, and climate data with methods based on nonlinear time series analysis (collectively referred to as empirical dynamic modeling), we identified drivers and their interactive effects on dengue dynamics in San Juan, Puerto Rico. Climatic forcing arose only when susceptible availability was high: temperature and rainfall had net positive and negative effects, respectively. By capturing mechanistic, nonlinear, and context-dependent effects of population susceptibility, temperature, and rainfall on dengue transmission empirically, our model improves forecast skill over recent, state-of-the-art models for dengue incidence. Together, these results provide empirical evidence that the interdependence of host population susceptibility and climate drive dengue dynamics in a nonlinear and complex, yet predictable way.

INTRODUCTION

Ecological systems are complex, nonlinear, and dynamic. Understanding their mechanistic drivers is increasingly important in a rapidly changing world. Because of this complexity, mechanistic studies are often conducted in simplified, controlled environments at small scales. Connecting these experimental results with larger-scale emerging patterns remains a challenge but is nonetheless imperative for understanding the impacts of ecological change.

In concert with globalization and climate change, mosquito-borne diseases, and dengue in particular, are (re)emerging globally and spreading to higher latitudes (Kilpatrick & Randolph 2012; Ryan *et al.* 2019). Dengue virus—vectored primarily by urban *Aedes aegypti* (Kraemer *et al.* 2015)—places half of the global human population in 128 countries at risk of infection (Brady *et al.* 2012; Kraemer *et al.* 2019). In the absence of effective vaccines or treatments (Katzelnick *et al.* 2017a; Sridhar *et al.* 2018), public health agencies rely on vector control to reduce dengue transmission (Erlanger *et al.* 2008). Effective vector control interventions require understanding the mechanisms linking climate, vector ecology, disease transmission, and host population susceptibility to better predict disease outbreaks—a major challenge.

Since *Aedes* spp. mosquitoes are sensitive to climate, including temperature and rainfall (Stewart Ibarra *et al.* 2013; Mordecai *et al.* 2019), we expect temperature and rainfall to be important drivers of dengue outbreaks. Although

temperature affects mosquito and viral traits in laboratory experiments (Watts *et al.*

1987; Lambrechts *et al.* 2011; Mordecai *et al.* 2017), the relationship between

temperature and dengue incidence in the field has been ambiguous (Caldwell *et al.*

2020). Thus, temperature-dependent models have had mixed success predicting the

timing and magnitudes of epidemics (Hii *et al.* 2012; Johansson *et al.* 2016; Johnson

et al. 2018). Similarly, the rainfall–dengue relationship is complex, because the

effect of rainfall on mosquitoes depends on local breeding habitat and human

behavior. In some settings, rainfall fills container-breeding habitats for mosquitoes,

increasing mosquito abundance and dengue incidence (Stewart Ibarra *et al.* 2013).

By contrast, low rainfall can also facilitate dengue transmission by promoting water

storage that serves as standing-water habitat for mosquitoes (Oliveira-lima *et al.*

2000). Further, heavy rainfall can reduce mosquito abundance by flushing out

larvae (Koenraadt & Harrington 2008). The net effect of climate on dengue

therefore depends on many different mechanisms and is highly context-dependent.

Disease incidence also depends nonlinearly on susceptible availability,

because epidemic growth slows as the population of susceptible individuals is

exhausted (Anderson & May 1979; Dushoff *et al.* 2004; Mina *et al.* 2015; Pitzer *et al.*

2015; Rypdal & Sugihara 2019). Further, susceptible availability may influence the

effects of climate on dengue dynamics. However, such interactive effects are difficult

to detect since susceptibility is difficult to observe, especially in endemic settings

where multiple serotypes circulate and create a complex landscape of time-

dependent and serotype-dependent immunity (Katzelnick *et al.* 2017b). Specifically,

four serotypes of dengue regularly circulate in many regions: each provides long-

term serotype-specific (homologous) immunity and short-term (heterologous)

cross-protection against other serotypes (dos Santos *et al.* 2017; Jiménez-Silva *et al.*

2018; Hamel *et al.* 2019). Following a brief period of cross-protection, antibodies at

a mid-range of titers can cause antibody-dependent enhancement of disease

following heterologous, secondary infection, until titers decay to the point of nearly

full heterologous susceptibility (Katzelnick *et al.* 2017b). Given this complex and

dynamic immune landscape, directly detecting population susceptibility to

circulating dengue virus at any point in time is difficult without longitudinal

serology studies, which are not widely available (Gordon *et al.* 2013; Katzelnick *et*

al. 2017b).

Previous prediction models of dengue outbreaks used phenomenological

(Johansson *et al.* 2009b; Hii *et al.* 2012; Johnson *et al.* 2018) and mechanistic

equation-based approaches (Tran *et al.* 2013; Liu-Helmersson *et al.* 2014; Morin *et*

al. 2015; Mordecai *et al.* 2017), which may not fully capture interdependence

between climate and susceptible availability. Phenomenological models may

underperform when extrapolating past observed contexts, and equation-based

mechanistic models rely on parameter estimates from laboratory studies

engineered to isolate single mechanisms producing separate relationships between

drivers and outcome, eliminating the complex interdependence at the population

level. While laboratory studies provide robust validation of mechanisms

(Lambrechts *et al.* 2011), the fixed relationships obtained from them do not

necessarily translate into robust causal understanding for the complexity of field

systems (Sugihara *et al.* 2012). Even if causality exists between two variables in such

a system, their correlation can switch signs during different time periods, resulting

in a net correlation of zero (Deyle *et al.* 2016b). This temporal variation in the

direction of correlation results from the nonlinear, state-dependent relationship

between the variables (i.e., the effect depends on another variable's state).

Conversely, even if two variables are consistently correlated, the association could

be spurious due to a confounder.

To overcome these challenges, we used empirical dynamic modeling (EDM)

(Sugihara *et al.* 2012)—a mechanistic, equation-free, data-driven approach that

accounts for the context-dependence of ecological drivers to identify and model

mechanisms driving dengue epidemics. EDM is based on reconstructing the

essential system dynamics evident in time series, without assuming fixed

relationships. This means relationships among variables can change through time to

reflect that interactions among variables depend on context and system state. EDM

does not require assumptions about the functional form of the model, but instead

derives dynamic relationships empirically by constructing an attractor—a

geometric object (i.e., curve or manifold) that embodies the rules for how

relationships among variables change with respect to each other through time

depending on system state (specific location on the attractor)—from time-series

observations. Like a set of equations, the attractor encompasses the dynamics of a

system, and thus can provide a mechanistic understanding of the system that is

derived empirically, without requiring an *a priori* assumed set of equations.

Here, we used EDM and a proxy for susceptible population size (Rypdal &

Sugihara 2019) to answer three questions: (1) Do temperature, rainfall, and/or

inferred susceptible availability drive population-level dengue incidence? (2) Can

we predict dengue dynamics using temperature and rainfall data and inferred

susceptible availability? (3) What is the functional form of each climate–dengue

relationship at the population level, and how is this relationship influenced by

susceptible availability?

METHODS

Time series data

We obtained time series of weekly observations of dengue incidence (total number

of new cases of all serotypes), average temperature (°C), and total rainfall (mm) in

San Juan, Puerto Rico, for 19 seasons (1990/1991–2008/2009) spanning calendar

week 18, 1990 to week 17, 2009 (Figure 1a–c) from the National Oceanic and

Atmospheric Administration in November 2016

(http://dengueforecasting.noaa.gov/). We obtained data for four additional seasons

(2009/2010–2012/2013) from Johnson *et al.* (2018) in April 2020

(https://github.com/lrjohnson0/vbdcast). Although dengue incidence data were

also available for Iquitos, Peru (Johansson *et al.* 2019), we chose to focus on San

Juan because the time series was longer, and therefore more amenable to EDM

analyses (Munch *et al.* 2020).

Direct measurements of susceptible availability are not available, so from

weekly incidence data $I(t)$, we estimated time-dependent growth rates: $\lambda =$

$I(t + \Delta t)/ I(t)$. The growth rate, λ, is proportional to the effective reproduction

number, R_{eff}, and equivalent to R_{eff} if Δt equals the average time between primary

and secondary host infections. Vector-borne disease models show that R_{eff} is

proportional to the geometric mean of the susceptible host population and the

susceptible vector population: $R_{\mathrm{eff}} = \sqrt{S_{\mathrm{h}}S_{\mathrm{v}}}\, R_0$, where R_0 is the basic reproduction

number (Zhao *et al.* 2020). Hence, $\lambda \propto \sqrt{S_{\mathrm{h}}S_{\mathrm{v}}}$ and λ can be used as a proxy for the

susceptible population size at least during inter-outbreak periods where the

transmission rate and R_0 can be assumed to vary very little (Rypdal & Sugihara

2019).

We estimated λ by linear regression using the model $I(t + \Delta t) = \lambda\, I(t)$ for

12 time points in a 12-week running window ($\Delta t = 1$ week). The model is robust to

the window size (Rypdal & Sugihara 2019). In the discrete case, when $\lambda < 1$ the

system is stable (inter-outbreak period) and when $\lambda \geq 1$ then the system is unstable

(outbreak period) (Supporting Information). We treated the resulting time series of

λ, hereafter "susceptibles index" (Figure 1d), as a proxy for the susceptible

population size when $\lambda < 1$, and a proxy for the combined effects of susceptible

availability and R_0 when $\lambda \geq 1$.

Empirical dynamic modeling (EDM)

EDM infers a system's mechanistic underpinnings and predicts its dynamics using

time series data of one or more variables to construct an attractor in state space

(Figure S1). This procedure is called univariate (using lagged versions of a single

variable time series) or multivariate state-space reconstruction (SSR). Properties of

the attractor are assessed to examine characteristics of the system (Deyle &

Sugihara 2011). We normalized each time series to zero mean and unit variance to

remove measurement unit bias, ensuring the variables would be comparable and

the attractor would not be distorted. All analyses were conducted in R version 3.5.1

(R Development Core Team 2018) and all EDM analyses were performed using

package rEDM (Park *et al.* 2020).

To infer mechanisms, EDM should be applied in systems where there is

evidence of underlying low-dimensional deterministic dynamics (Cummins *et al.*

2015). EDM assumptions are met when stochasticity is present (e.g., due to

measurement noise, stochastic drivers, or unexplained variability) (Cenci *et al.*

2019; Munch *et al.* 2020), but the system cannot be entirely stochastic. To test for

low-dimensional deterministic dynamics we performed univariate SSR for each

variable, and used *simplex projection* (Sugihara & May 1990)—a type of nearest

neighbor regression performed on an attractor—to check whether the system is

forecastable beyond the skill of an autoregressive model—an indicator of

underlying deterministic dynamics (Figures S2a and S4; Supporting Information).

To test for nonlinear state dependence of a variable—the motivation behind EDM—

we used the *S-map* test for nonlinearity (Sugihara 1994) (Figures S2b, c and S5;

Supporting Information).

EDM—Convergent cross-mapping

We used an EDM approach called *convergent cross-mapping* (CCM) (Sugihara *et al.*

2012) to identify drivers of dengue incidence. If two variables are causally related,

then a multivariate attractor—where each variable in the system represents a

dimension that traces the dynamics of the system—can be reconstructed (up to a

practical limit) using lagged versions of just one of the variables (Figure S1). Based

on Takens' Theorem, this univariate "shadow attractor" preserves the structural and

dynamic properties of the original multivariate attractor (Takens 1981; Sugihara *et*

al. 2012). The concept behind CCM is that if temperature causes dengue incidence,

then information about past temperature will be embedded in the dynamics of

dengue, such that the shadow attractor produced using only incidence data allows

us to accurately reconstruct temperature in the past. However, the converse

scenario would not be true: since dengue does not cause temperature, the shadow

attractor constructed using temperature data should not contain information to

accurately reconstruct past dengue incidence (Supporting Information).

The critical criterion for estimating causal (directional) associations between

two variables using CCM is checking that the cross-mapping skill (i.e., Pearson's

correlation coefficient, ρ, between predicted driver values using the univariate SSR

of the response variable, and the observed driver values) monotonically increases

and plateaus (i.e., converges) with the length of the response variable time series

used in cross-mapping. We used the Kendall's τ test as a significance test for

convergence of cross-mapping skill using the Kendall package (McLeod 2011). If

$\tau > 0$ then there is convergence (Grziwotz *et al.* 2018).

We performed pairwise cross-correlations on the time series to investigate

time-lagged relationships between potential drivers (i.e., temperature, rainfall, and

susceptibles index) and dengue incidence using the tseries package (Trapletti &

Hornik 2018). Based on these analyses (Figure S6), we applied a 9-week time lag

between temperature and incidence, an averaged lag of 3–9 weeks for rainfall (i.e.,

the average rainfall over the preceding 3–9 weeks) to resemble standing water as

mosquito breeding habitat over a longer time period, and a 5-week lag for the

susceptibles index. These lags are proxies for the time delays of potential cause-and-

effects and are consistent with results from field studies (Chen *et al.* 2010; Stewart

Ibarra *et al.* 2013).

We assessed the strength of evidence for effects of potential drivers on

dengue by comparing the CCM performance using the data with the performance of

two null models that control for the seasonal trend (i.e., interannual mean) observed

in all variables (Figure 2). These null models address the sensitivity of CCM to

periodic fluctuations (i.e., seasonality), which can make two variables appear to be

causally linked when instead they are simply synchronized by a seasonal

confounder (Cobey & Baskerville 2016; Deyle *et al.* 2016a). In the first "seasonal"

null model, we preserved the seasonal signal, but randomized the interannual

anomalies (Deyle *et al.* 2016a). In the second, more conservative "Ebisuzaki" null

model, we conserved any periodicity (beyond seasonal) and randomized the phases

of Fourier-transformed time series (Ebisuzaki 1997). We tested for statistically

significant differences in cross-mapping skill between the model that used the data

versus the null models by performing Kolmogorov-Smirnov (K-S) tests after

convergence.

We also repeated CCM in the nonsensical, reverse-causal direction (e.g., to

test whether incidence drives climate) as a control for potential spurious

relationships generated by non-causal covariation (e.g., due to seasonality). This

addresses the issue of synchrony, in which CCM can indicate bidirectional causality

when one direction is false or nonsensical (Baskerville & Cobey 2017; Sugihara *et al.*

2017).

EDM—Forecast improvement

We examined the predictive power of the drivers on dengue incidence by assessing

how well we can predict dengue dynamics using temperature, rainfall, susceptibles

index, and their combined effects. We used a combination of univariate SSR (i.e.,

with incidence data) and multivariate SSR to build forecasting models and to

determine the improvement of forecasting using simplex projection when including

different combinations of drivers (Dixon *et al.* 1999; Deyle *et al.* 2013, 2016a)

(Supporting Information). We built the SSR forecasting models/attractors using the

1990/1991–2008/2009 season data (Figure 1) and made forecasts 8 weeks ahead.

We assessed model forecasting performance using leave-one-out cross-validation.

Next, we evaluated out-of-sample forecasting performance of these models

using testing data from four additional seasons (2009/2010–2012/2013).

Predictions made on week zero for the first forecast of the 2009/2010–2012/2013

period (8 weeks ahead) came only from SSR using the 1990/1991–2008/2009 data.

All subsequent weekly forecasts (8 weeks ahead) were made from updated SSR

using all previous data, including past observations from the testing dataset.

Forecast uncertainty was evaluated by taking the density and morphology of

the attractor into account. The more compact a simplex was and the less its starting

position on the attractor mattered for the simplex projection, the more certain we

were about our point estimate. Forecast variance was obtained from a distribution

of weighted nearest neighbor regression from edges of simplexes constructed at

various starting positions in the past.

Finally, we compared our top model performance with performance of

previous models from 16 teams that participated in a dengue forecasting challenge

(Johansson *et al.* 2019) and had access to the same data. To make a fair comparison,

we followed the procedure as directed in the challenge (Supporting Information).

EDM—Scenario exploration

In nonlinear systems, drivers generally have an effect that is state-dependent: the strength and direction of the effect depends on the current state of the system. Scenario exploration with multivariate EDM allowed us to assess the effect of a small change in temperature or rainfall on dengue incidence, across different states of the system. The outcome of these small changes allowed us to deduce the relationship between each climate driver and dengue incidence and how they depend on the system state. For each time step t we used S-maps (Sugihara 1994; Deyle *et al.* 2016a) to predict dengue incidence using a small increase ($+\Delta X$) and a small decrease ($-\Delta X$) of the observed value of driver $X(t)$ (temperature or rainfall). For each putative climate driver, the difference in dengue predictions between these small changes is $\Delta Y = Y(t+1)\left[X(t) + \frac{\Delta X(t)}{2}\right] - Y(t+1)\left[X(t) - \frac{\Delta X(t)}{2}\right]$, where $Y(t+1)$ is a function of X and all other state variables, and we used $\Delta Y/\Delta X$ to approximate the effect of driver X at time t. We repeated this over all time steps in our time series for both temperature and rainfall to recover their approximate relationships with dengue incidence at different states of the system. Scenario exploration analyses were repeated across several model parameterizations to address potential sensitivity to parameter settings (Supporting Information).

RESULTS

Drivers of dengue dynamics

EDM showed that temperature, rainfall, and the susceptibles index drive dengue incidence since the convergence criterion was met (Kendall's $\tau > 0$, $P < 0.01$) in all three cases (Figure 3a–c). Rainfall and susceptibles index were significant drivers of dengue incidence beyond seasonality, as their effects were distinguishable from seasonal and Ebisuzaki null models (Figures 3b–c and S8b–c; K-S $P < 0.0001$). This implies statistically significant effects of both rainfall and the susceptibles index on dengue, which are not obscured by a periodic confounder. However, temperature was not a significant driver beyond seasonality (Figures 3a and S8a; K-S $P = 0.90$). We cannot rule out the possibility that the apparent forcing of temperature on dengue is due to a seasonal confounder. However, if no such confounder exists, then the seasonal trend in temperature, which accounts for most temperature variation in San Juan, drives the seasonal trend observed in dengue incidence. Compared to the other drivers, the converging cross-mapping skill of the temperature null models were relatively high (Figures 3 and S8), suggesting that temperature seasonality in each null model was a strong driver. Thus, seasonal temperature may be driving dengue dynamics, a result consistent with other studies (Huber *et al.* 2018; Robert *et al.* 2019).

As expected, EDM tests for putative causality in the nonsensical directions—incidence driving temperature or rainfall—were not significant (i.e., no

convergence; Figure S7, black lines). This result further supports the finding that

temperature and rainfall drive dengue incidence, because their causal relationships

were not confounded by spurious bidirectionality. The null models for the

nonsensical directions of causality (Figure S7, grey lines) also displayed no

convergence (completely flat), as expected (i.e., seasonality of dengue incidence

does not drive seasonality of temperature or rainfall). However, seasonality (or any

periodicity) of temperature, rainfall and susceptibles index drive dengue dynamics,

shown by convergence of the seasonal and Ebisuzaki null models (grey lines in

Figures 3 and S8).

Predictive power of drivers

The multivariate SSR model using only temperature and rainfall data did not predict

dengue incidence very well (ρ = 0.3839, RMSE = 47.72) although it captured the

seasonality of the epidemics (Figure 4a). Forecasting skill doubled when the

susceptibles index was included along with rainfall and temperature (ρ = 0.7547,

RMSE = 37.40; Figure 4c), where timing and magnitude of epidemics were captured

reasonably well. Dengue incidence prediction improved even further when

incidence was added into the model with all drivers (ρ = 0.7662, RMSE = 37.14;

Figure 4e). Dengue incidence was somewhat predictable using univariate SSR of

incidence data alone (ρ = 0.4459, RMSE = 46.75; Figure 4g), suggesting that the

dengue incidence time series contains information about its drivers, although

limited. This points to some additional value of including the driver variables.

We also evaluated the performance of the SSR models (Figure 4a, c, e, g)

constructed using data from seasons 1990/1991–2008/2009 on external, testing

data from 2009/2010–2012/2013 that were not used in SSR (Figure 4b, d, f, h). The

average out-of-sample forecasting skill for each model for the testing seasons was

higher than that of the 1990/1991–2008/2009 forecasts, although the errors were

larger. The model using only temperature and rainfall displayed predictability (ρ =

0.8989, RMSE = 52.30; Figure 4b), the model that also included the susceptibles

index improved predictions (ρ = 0.9475, RMSE = 52.12; Figure 4d), and the model

that also included past incidence made highly accurate predictions (ρ = 0.9697,

RMSE = 46.75; Figure 4f). The model that only included dengue incidence without

the drivers was also predictive, although more error-prone (ρ = 0.9044, RMSE =

57.34; Figure 4h). All SSR models (Figure 4a–h) had significant forecasting skill (ρ)

values (Fisher's z-transformation P < 0.001).

The model with the highest prediction skill for the testing seasons

(2009/2010–2012/2013), which included past climate, susceptibles index, and

incidence data as predictors (Figure 4f), also outperformed models from the dengue

forecasting challenge, including the ensemble model (Johansson *et al.* 2019) for

predicting peak incidence, peak week, and seasonal incidence for all seasons on

average (Tables S1–S2, Figures S9–S12). This demonstrates the benefit of the EDM

approach for capturing the mechanistic, nonlinear, interdependent relationships

among drivers over both equation-based mechanistic models and phenomenological

models.

State-dependent functional responses

As expected, we find state-dependent effects of temperature and rainfall with non-zero median effects. We found that temperature had a small positive median effect (2.88 cases/°C, Wilcox $P < 0.001$) on dengue incidence (Figure 5a). A positive effect is expected for the temperature range in Puerto Rico (Mordecai *et al.* 2017) (Figure 6e, black dashed lines), although the effect was occasionally much stronger, both positive and negative (Figure 5a, b). The large negative effects occurred only at the highest temperature values (as predicted by mechanistic models of temperature-dependent transmission), reinforced by a lower quantile regression with a strongly negative slope (Figure 5b, bottom dashed red line). However, positive effects occurred across the whole temperature range, which is limited to temperatures below the 29°C optimal temperature for transmission estimated from mathematical models and laboratory data (Mordecai *et al.* 2017).

Rainfall had a small negative median effect (−0.12 cases/mm, Wilcox $P < 0.001$), but occasionally had very large negative effects (Figure 5a, c). These large, negative effects of rainfall on dengue occurred when there was less than 100 mm of rain per week (Figure 5c), consistent with expectations that drought could lead to a high number of dengue cases due to water storage, which can provide mosquito breeding habitat (Oliveira-lima *et al.* 2000). There are also small positive effects of rainfall on dengue (Figure 5c), suggesting that overall the results showed competing effects of low to moderate rain providing standing water for mosquito breeding and

humans storing water where mosquitoes can breed when there is drought or low

rainfall.

These results suggest the strength and direction of the effects of climate on

dengue dynamics depend on the state of the system. In addition to the nonlinear

effects of climate drivers themselves on dengue incidence, another potential cause

of state-dependent climate effects on dengue dynamics is the variation in the

susceptible population size over time (Figure 6a, b). Outbreaks do not occur when

there are too few susceptible people in the population. As expected, when the

susceptibles index was small ($\lambda < 0.85$) incidence was insensitive to climate (Figure

6c, f). By contrast, when the susceptibles index was large ($\lambda > 0.85$), temperature

and rainfall effects on dengue incidence appeared (Figure 6d, g). The gradual

increase and decrease of the rate of change of dengue as a function of temperature

(Figure 6d, red solid lines) aligned well with the changes in slope over the

increasing part (Figure 6e, black dashed lines representing the temperature range in

our study) of the unimodal temperature response curve for dengue transmission by

Ae. aegypti developed previously (Mordecai *et al.* 2017). This is an important

finding, since evidence of climate functional responses for disease dynamics is rare

due to the difficulty of obtaining appropriately informative field data. It is possible

that if we had temperature data ranging across a larger spectrum—possibly by

assembling data across multiple climates—that the empirical functional response

derived from EDM would also look unimodal. Further, when the susceptibles index

was high, the slope of the relationship between rainfall and dengue incidence

became more negative as rainfall increased, suggesting a concave-down effect of

rainfall on incidence (Figure 6g, h). This relationship has been difficult to

characterize in the field because of multiple, possibly context-dependent and lagged,

mechanisms linking rainfall to dengue.

DISCUSSION

High host susceptibility allows seasonal climate suitability to fuel large dengue

epidemics in San Juan, Puerto Rico. The effects of climate and susceptibility are

nonlinear, interdependent, and state-dependent, which makes inference from

controlled mechanistic experiments, equation-based mechanistic models, or

phenomenological models difficult. EDM provides an essential toolkit for identifying

these drivers, quantifying their predictive power, and approximating their

functional responses. In Puerto Rico, the causes of extensive interannual variability

in dengue incidence have remained a mystery, despite hypotheses that climate and

host susceptibility were involved. Here, we used EDM and a proxy for susceptible

availability to disentangle nonlinear and interactive mechanisms driving disease

dynamics.

We found that rainfall, susceptible availability, and plausibly temperature

(via its seasonality) interact to drive dengue incidence. Combined, these three

drivers predicted dengue incidence with high accuracy (Figure 4c). The EDM-based

forecasting model outperformed 16 models and an ensemble model in a recently

published dengue forecasting challenge (Johansson *et al.* 2019), suggesting that it

could enhance dengue control efforts if surveillance efforts continue to report

weekly case data. Finally, as expected from epidemiological theory, climate effects

on dengue only appeared when susceptible availability exceeded a threshold ($\lambda >$

0.85; Figure 6).

The fact that climate effects are first observed when $\lambda \approx 0.85$ (before the

onset of an outbreak, $\lambda = 1$), suggests that rainfall, and possibly temperature, have

an effect on the timing of an impending epidemic. Climate could drive the

transmission rate, thus influencing λ (which is proportional to both susceptible

population size and R_0 when λ is close to 1), and therefore the timing of an outbreak

could be attributed to the changes in transmission caused by seasonal climatic

drivers (Rypdal & Sugihara 2019). The seasonality of temperature and rainfall had

higher predictive skill than seasonality of susceptibles index (Figures 3 and S8, grey

lines), further supporting that seasonality of incidence was associated with climate.

However, the susceptibles index was critical for predicting dengue epidemic

magnitudes (Figure 4c–f). Using the same data, Johnson *et al.* (2018) found that

mechanistic models could predict the timing of seasonal epidemics, but that a

phenomenological machine learning component was needed to capture interannual

variation in epidemic magnitude. Our work suggests that the unobserved size of the

susceptible population was a key missing link for predicting magnitude variation

across years.

Previous studies have built models accounting for both susceptible

availability and climate on dengue by reconstructing time series of susceptibles

from a compartmental modeling framework (Metcalf *et al.* 2017). However, no

previous studies on dengue have explored the interdependence between climate

and susceptible population size. We showed that susceptible availability modifies

climate effects on dengue: climate has negligible effects unless the susceptible

population size is large enough (Figure 6). The interdependence of climate and

population susceptibility has also been studied in diseases where the opposite effect

was found. For example, climate effects on SARS-CoV-2 are expected to be negligible

when susceptible availability is high in the early stage of the emerging pandemic

(Baker *et al.* 2020). For influenza dynamics, population density in cities—potentially

a proxy for susceptible availability—also modulated climate effects on disease

transmission: climate effects were negligible in cities with high population densities

(Dalziel *et al.* 2018).

Because dengue susceptibility is so complex—due to the serotype dynamics

and time- and antibody titer-dependent cross-protection and enhancement

(Katzelnick *et al.* 2017b)—total population density or size may not be a reasonable

proxy for susceptible availability in dengue dynamics, and a direct mechanistic

estimate of population susceptibility will likely never be widely available for most

populations. Accordingly, it has been difficult for previous mechanistic models to

capture susceptible dynamics for dengue and their interactions with climate.

However, our approach provides a useful proxy that captures the susceptible

population dynamics even in the absence of more detailed immunological

information. By inferring the susceptibles index from incidence data, we were able

to capture the strong influence of the susceptible availability on dengue dynamics,

which in turn moderated the effect of climate on dengue dynamics. This result is

expected from theory (Kermack & McKendrick 1927; Xu *et al.* 2017), but

demonstrating it empirically is a unique contribution of this study.

Even when accounting for susceptible availability, the effects of temperature

and rainfall on dengue were strongly state-dependent (Figure 6d, g). This result is

potentially due to nonlinear effects of each climate driver (Figure 6e, h), interactions

and correlations between temperature and rainfall, microclimate variation over

space and time that is not captured by weekly averages, and complex lagged effects

that are not captured by a single fixed lag (e.g., 9 weeks). In Puerto Rico, mosquitoes

also breed in septic tanks year-round, allowing transmission to occur independently

from rainfall (Mackay *et al.* 2009), thus weakening the rainfall–dengue negative

relationship (Figure 6g). Some of this additional variation may be captured in the

dengue incidence time series itself, which may explain why including it improves

forecast skill over climate and susceptibility alone (Figure 4e, f). Despite this

additional variation, our results are consistent with previous studies suggesting that

dengue dynamics in Puerto Rico are positively associated with temperature

(Johansson *et al.* 2009b; Barrera *et al.* 2011; Morin *et al.* 2015), and possibly

negatively associated with rainfall (Johansson *et al.* 2009a; Morin *et al.* 2015), since

most *Ae. aegypti* pupae in Puerto Rico are produced in human-made containers

during periods of drought (Barrera *et al.* 2011).

The climate and incidence data used here have been used in multiple

forecasting efforts, where ensemble approaches and approaches that incorporated

mechanisms outperformed purely statistical approaches (Johansson *et al.* 2019).

However, even the high-performing forecasting methods using the same dataset,

and including (experimentally-derived) assumed mechanisms for the joint influence

of climate and susceptibility on dengue dynamics, are still error-prone to the timing

(on the order of weeks) and the magnitude (on the order of 50 cases) of intra-

annual epidemics (Morin *et al.* 2015; Johansson *et al.* 2019). Mechanisms isolated

independently in controlled experiments do not necessarily scale up to the

population level, and susceptible dynamics derived from compartmental models

may be too simple to properly capture true susceptibility at the population level for

dengue (Katzelnick *et al.* 2017b). EDM allowed us to infer mechanisms empirically

from population-level data, and accounted for the population-level interdependence

between climate and susceptible availability for forecasting, which probably

contributed to our model outperforming previous forecasting models and

ensembles (Table S1).

Connecting climate and dengue at the population level is challenging,

because causal relationships are likely to be nonlinear and state-dependent. A

toolkit of methods for testing hypotheses, understanding mechanisms, and making

predictions is essential for understanding disease dynamics in complex, natural

populations. Ultimately, understanding how climate-driven vector-borne diseases

are influenced by other variables, such as susceptible population size, is important

for optimizing vector control under critical conditions where climate might spark

epidemics. Our approach, using EDM and an inferred proxy for the susceptible

population size from data, confirmed that climate has nonlinear, seasonal effects on

dengue epidemics in San Juan, Puerto Rico, but only above a certain threshold of

susceptible availability. The mechanisms inferred from EDM could be applied to

understand and predict future ecological responses to changing environments,

including dengue epidemics in a world undergoing rapid environmental change.

ACKNOWLEDGEMENTS

We thank Giulio De Leo, Marcus Feldman, Dmitri Petrov, and members of the Fukami, Mordecai, Peay, and Sugihara labs for helpful feedback. NN was supported by the Bing Fellowship in Honor of Paul Ehrlich and the Stanford Data Science Scholars program. ERD and GS were supported by the National Science Foundation (NSF) DEB-1655203, NSF-ABI-1667584, DoD-Strategic Environmental Research and Development Program (SERDP) 15 RC-2509; Lenfest Foundation Award 00028335 and the McQuown Chair in Natural Sciences, University of California, San Diego. MSS and EAM were supported by an NSF Ecology and Evolution of Infectious Diseases grant (DEB-1518681). EAM was also supported by an NSF Rapid Response Research grant (RAPID 1640780), an NIH NIGMS R35 MIRA award (R35GM133439), the Stanford University Woods Institute for the Environment Environmental Ventures Program, the Hellman Faculty Fellowship, a Stanford King Center Seed Grant, and the Terman Fellowship. AJM was supported by an NSF Postdoctoral Research Fellowship in Biology (1611767). MLC was supported by the Lindsay Family E-IPER Fellowship and Illich-Sadowsky Interdisciplinary Graduate Fellowship.

REFERENCES

Anderson, R.M. & May, R.M. (1979). Population biology of infectious diseases: Part I. *Nature,* 280, 361–367.

Baker, R.E., Yang, W., Vecchi, G.A., Metcalf, C.J.E. & Grenfell, B.T. (2020). Susceptible supply limits the role of climate in the early SARS-CoV-2 pandemic. *Science*, 369, 315–319.

Barrera, R., Amador, M. & MacKay, A.J. (2011). Population dynamics of *Aedes aegypti* and dengue as influenced by weather and human behavior in san juan, puerto rico. *PLoS Negl. Trop. Dis.*, 5, e1378.

Baskerville, E.B. & Cobey, S. (2017). Does influenza drive absolute humidity? *Proc. Natl. Acad. Sci.*, 114, 201700369.

Brady, O.J., Gething, P.W., Bhatt, S., Messina, J.P., Brownstein, J.S., Hoen, A.G., *et al.* (2012). Refining the Global Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. *PLoS Negl. Trop. Dis.*, 6, e1760.

Caldwell, J., LaBeaud, D., Lambin, E., Stewart-Ibarra, A., Ndenga, B., Mutuku, F., *et al.* (2020). Climate explains geographic and temporal variation in mosquito-borne disease dynamics on two continents. *bioRxiv*, 2020.02.07.938720.

Cenci, S., Sugihara, G. & Saavedra, S. (2019). Regularized S-map for inference and forecasting with noisy ecological time series. *Methods Ecol. Evol.*, 2041–210X.13150.

Chen, S.C., Liao, C.M., Chio, C.P., Chou, H.H., You, S.H. & Cheng, Y.H. (2010). Lagged

temperature effect with mosquito transmission potential explains dengue

variability in southern Taiwan: Insights from a statistical analysis. *Sci. Total*

Environ., 408, 4069–4075.

Cobey, S. & Baskerville, E.B. (2016). Limits to causal inference with state-space

reconstruction for infectious disease. *PLoS One*, 11, 1–22.

Cummins, B., Gedeon, T. & Spendlove, K. (2015). On the Efficacy of State Space

Reconstruction Methods in Determining Causality. *SIAM J. Appl. Dyn. Syst.*, 14,

335–381.

Dalziel, B.D., Kissler, S., Gog, J.R., Viboud, C., Bjørnstad, O.N., Metcalf, C.J.E., *et al.*

(2018). Urbanization and humidity shape the intensity of influenza epidemics

in U.S. cities. *Science*, 362, 75–79.

Deyle, E.R., Fogarty, M., Hsieh, C., Kaufman, L., Maccall, A.D. & Munch, S.B. (2013).

Predicting climate effects on Pacific sardine. *Proc. Natl. Acad. Sci. U. S. A.*, 110,

6430–6435.

Deyle, E.R., Maher, M.C., Hernandez, R.D., Basu, S. & Sugihara, G. (2016a). Global

environmental drivers of influenza. *Proc. Natl. Acad. Sci.*, 113, 13081–13086.

Deyle, E.R., May, R.M., Munch, S.B. & Sugihara, G. (2016b). Tracking and forecasting

ecosystem interactions in real time. *Proc. R. Soc. B Biol. Sci.*, 283.

Deyle, E.R. & Sugihara, G. (2011). Generalized theorems for nonlinear state space

reconstruction. *PLoS One*, 6.

Dixon, P.A., Milicich, M.J. & Sugihara, G. (1999). Episodic fluctuations in larval supply.

Science, 283, 1528–1530.

Dushoff, J., Plotkin, J.B., Levin, S.A. & Earn, D.J.D. (2004). Dynamical resonance can

account for seasonality of influenza epidemics. *Proc. Natl. Acad. Sci. U. S. A.*, 101,

16915–16916.

Ebisuzaki, W. (1997). A method to estimate the statistical significance of a

correlation when the data are serially correlated. *J. Clim.*, 10, 2147–2153.

Erlanger, T.E., Keiser, J. & Utzinger, J. (2008). Effect of dengue vector control

interventions on entomological parameters in developing countries: A

systematic review and meta-analysis. *Med. Vet. Entomol.*

Gordon, A., Kuan, G., Mercado, J.C., Gresh, L., Avilés, W., Balmaseda, A., *et al.* (2013).

The Nicaraguan Pediatric Dengue Cohort Study: Incidence of Inapparent and

Symptomatic Dengue Virus Infections, 2004-2010. *PLoS Negl. Trop. Dis.*, 7,

e2462.

Grziwotz, F., Strauß, J.F., Hsieh, C. & Telschow, A. (2018). Empirical Dynamic

Modelling Identifies different Responses of *Aedes Polynesiensis* Subpopulations

to Natural Environmental Variables. *Sci. Rep.*, 8, 16768.

Hamel, R., Surasombatpattana, P., Wichit, S., Dauvé, A., Donato, C., Pompon, J., *et al.*

(2019). Phylogenetic analysis revealed the co-circulation of four dengue virus

serotypes in Southern Thailand. *PLoS One*, 14, e0221179.

Hii, Y.L., Zhu, H., Ng, N., Ng, L.C. & Rocklöv, J. (2012). Forecast of Dengue Incidence

Using Temperature and Rainfall. *PLoS Negl. Trop. Dis.*, 6, e1908.

Huber, J.H., Childs, M.L., Caldwell, J.M. & Mordecai, E.A. (2018). Seasonal

temperature variation influences climate suitability for dengue, chikungunya,

and Zika transmission. *PLoS Negl. Trop. Dis.*, 12, e0006451.

Jiménez-Silva, C.L., Carreño, M.F., Ortiz-Baez, A.S., Rey, L.A., Villabona-Arenas, C.J. &

Ocazionez, R.E. (2018). Evolutionary history and spatio-temporal dynamics of

dengue virus serotypes in an endemic region of Colombia. *PLoS One*, 13,

e0203090.

Johansson, M.A., Apfeldorf, K.M., Dobson, S., Devita, J., Buczak, A.L., Baugher, B., *et al.*

(2019). An open challenge to advance probabilistic forecasting for dengue

epidemics. *Proc. Natl. Acad. Sci.*, 116, 24268–24274.

Johansson, M.A., Cummings, D.A.T. & Glass, G.E. (2009a). Multiyear climate

variability and dengue - El Niño southern oscillation, weather, and dengue

incidence in Puerto Rico, Mexico, and Thailand: A longitudinal data analysis.

PLoS Med., 6, e1000168.

Johansson, M.A., Dominici, F. & Glass, G.E. (2009b). Local and global effects of climate

on dengue transmission in Puerto Rico. *PLoS Negl. Trop. Dis.*, 3, e382.

Johansson, M.A., Reich, N.G., Hota, A., Brownstein, J.S. & Santillana, M. (2016).

Evaluating the performance of infectious disease forecasts: A comparison of

climate-driven and seasonal dengue forecasts for Mexico. *Sci. Rep.*, 6, 33707.

Johnson, L.R., Gramacy, R.B., Cohen, J., Mordecai, E., Murdock, C., Rohr, J., *et al.*

(2018). Phenomenological forecasting of disease incidence using

heteroskedastic gaussian processes: A dengue case study. *Ann. Appl. Stat.*, 12,

27–66.

Katzelnick, L.C., Coloma, J. & Harris, E. (2017a). Dengue: knowledge gaps, unmet

610  needs, and research priorities. *Lancet Infect. Dis.*, 17, e88–e100.

611 Katzelnick, L.C., Gresh, L., Halloran, M.E., Mercado, J.C., Kuan, G., Gordon, A., *et al.*

612  (2017b). Antibody-dependent enhancement of severe dengue disease in

613  humans. *Science*, 358, 929–932.

614 Kermack, W.O. & McKendrick, A.G. (1927). A Contribution to the Mathematical

615  Theory of Epidemics. *Proc. R. Soc. A Math. Phys. Eng. Sci.*, 115, 700–721.

616 Kilpatrick, A.M. & Randolph, S.E. (2012). Drivers, dynamics, and control of emerging

617  vector-borne zoonotic diseases. *Lancet.*

618 Koenraadt, C.J.M. & Harrington, L.C. (2008). Flushing effect of rain on container-

619  inhabiting mosquitoes *Aedes aegypti* and *Culex pipiens* (Diptera: *Culicidae*). *J.*

620  *Med. Entomol.*, 45, 28–35.

621 Kraemer, M.U.G., Reiner, R.C., Brady, O.J., Messina, J.P., Gilbert, M., Pigott, D.M., *et al.*

622  (2019). Past and future spread of the arbovirus vectors *Aedes aegypti* and *Aedes*

623  *albopictus*. *Nat. Microbiol.*, 4, 854–863.

624 Kraemer, M.U.G., Sinka, M.E., Duda, K.A., Mylne, A.Q.N., Shearer, F.M., Barker, C.M., *et*

625  *al.* (2015). The global distribution of the arbovirus vectors *Aedes aegypti* and

626  *Ae. Albopictus*. *Elife*, 4.

627 Lambrechts, L., Paaijmans, K.P., Fansiri, T., Carrington, L.B., Kramer, L.D., Thomas,

628  M.B., *et al.* (2011). Impact of daily temperature fluctuations on dengue virus

629  transmission by *Aedes aegypti*. *Proc. Natl. Acad. Sci. U. S. A.*, 108, 1–6.

630 Liu-Helmersson, J., Stenlund, H., Wilder-Smith, A. & Rocklöv, J. (2014). Vectorial

631  capacity of *Aedes aegypti*: Effects of temperature and implications for global

dengue epidemic potential. *PLoS One*, 9.

Mackay, A.J., Amador, M., Diaz, A., Smith, J. & Barrera, R. (2009). Dynamics of *Aedes*

aegypti and *Culex quinquefasciatus* in Septic Tanks. *J. Am. Mosq. Control Assoc.*,

25, 409–416.

McLeod, A.I. (2011). Kendall: Kendall rank correlation and Mann-Kendall trend test.

Metcalf, C.J.E., Walter, K.S., Wesolowski, A., Buckee, C.O., Shevliakova, E., Tatem, A.J.,

et al. (2017). Identifying climate drivers of infectious disease dynamics: Recent

advances and challenges ahead. *Proc. R. Soc. B Biol. Sci.*

Mina, M.J., Metcalf, C.J.E., De Swart, R.L., Osterhaus, A.D.M.E. & Grenfell, B.T. (2015).

Long-term measles-induced immunomodulation increases overall childhood

infectious disease mortality. *Science*, 348, 694–699.

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., *et*

al. (2019). Thermal biology of mosquito-borne disease. *Ecol. Lett.*, ele.13335.

Mordecai, E.A., Cohen, J.M., Evans, M. V., Gudapati, P., Johnson, L.R., Lippi, C.A., *et al.*

(2017). Detecting the impact of temperature on transmission of Zika, dengue,

and chikungunya using mechanistic models. *PLoS Negl. Trop. Dis.*, 11,

e0005568.

Morin, C.W., Monaghan, A.J., Hayden, M.H., Barrera, R. & Ernst, K. (2015).

Meteorologically driven simulations of dengue epidemics in San Juan, PR. *PLoS*

Negl. Trop. Dis., 9, e0004002.

Munch, S.B., Brias, A., Sugihara, G. & Rogers, T.L. (2020). Frequently asked questions

about nonlinear dynamics and empirical dynamic modelling. *ICES J. Mar. Sci.*,

77, 1463–1479.

Oliveira-lima, W., Hodgson, J.C., Pontes, R.J.S. & Freeman, J. (2000). Vector densities

that potentiate dengue outbreaks in a Brazilian city. *Trop. Med.*, 62, 378–383.

Park, J., Smith, C., Sugihara, G. & Deyle, E.R. (2020). rEDM: Empirical Dynamic

Modeling ('EDM'). R package version 1.6.1.

Pitzer, V.E., Viboud, C., Alonso, W.J., Wilcox, T., Metcalf, C.J., Steiner, C.A., *et al.* (2015).

Environmental Drivers of the Spatiotemporal Dynamics of Respiratory

Syncytial Virus in the United States. *PLoS Pathog.*, 11, 1–14.

Robert, M.A., Christofferson, R.C., Weber, P.D. & Wearing, H.J. (2019). Temperature

impacts on dengue emergence in the United States: Investigating the role of

seasonality and climate change. *Epidemics*, 28.

Ryan, S.J., Carlson, C.J., Mordecai, E.A. & Johnson, L.R. (2019). Global expansion and

redistribution of *Aedes*-borne virus transmission risk with climate change. *PLoS*

Negl. Trop. Dis., 13, e0007213.

Rypdal, M. & Sugihara, G. (2019). Inter-outbreak stability reflects the size of the

susceptible pool and forecasts magnitudes of seasonal epidemics. *Nat.*

Commun., 10, 2374.

dos Santos, T.P., Cruz, O.G., da Silva, K.A.B., de Castro, M.G., de Brito, A.F., Maspero,

R.C., *et al.* (2017). Dengue serotype circulation in natural populations of *Aedes*

aegypti. *Acta Trop.*, 176, 140–143.

Sridhar, S., Luedtke, A., Langevin, E., Zhu, M., Bonaparte, M., Machabert, T., *et al.*

(2018). Effect of Dengue Serostatus on Dengue Vaccine Safety and Efficacy. *N.*

Engl. J. Med., 379, 327–340.

Stewart Ibarra, A.M., Ryan, S.J., Beltrá N, E., Mejía, R.L., Silva, M. & Muñ Oz, N. (2013).

Dengue Vector Dynamics (*Aedes aegypti*) Influenced by Climate and Social

Factors in Ecuador: Implications for Targeted Control. *PLoS One*, 8.

Sugihara, G. (1994). Nonlinear forecasting for the classification of natural time

series. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 348, 477–495.

Sugihara, G., Deyle, E.R. & Ye, H. (2017). Reply to Baskerville and Cobey:

Misconceptions about causation with synchrony and seasonal drivers. *Proc.*

Natl. Acad. Sci., 114, 201700998.

Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M., *et al.* (2012). Detecting

Causality in Complex Ecosystems. *Science*, 338, 496–500.

Sugihara, G. & May, R.M. (1990). Nonlinear forecasting as a way of distinguishing

chaos from measurement error in time series. *Nature,* 344, 734–741.

Takens, F. (1981). Detecting strange attractors in turbulence. In: *Dynamical Systems*

and Turbulence, Warwick 1980, Lecture Notes in Mathematics (eds. Rand, D. &

Young, L.-S.). Springer, Berlin, Heidelberg., pp. 366–381.

Tran, A., L'Ambert, G., Lacour, G., Benoît, R., Demarchi, M., Cros, M., *et al.* (2013). A

rainfall- and temperature-driven abundance model for *Aedes albopictus*

populations. *Int. J. Environ. Res. Public Health*, 10, 1698–1719.

Trapletti, A. & Hornik, K. (2018). tseries: Time Series Analysis and Computational

Finance.

Watts, D.M., Burke, D.S., Harrison, B.A., Whitmire, R.E. & Nisalak, A. (1987). Effect of

temperature on the vector efficiency of *Aedes aegypti* for dengue 2 virus. *Am. J.*

Trop. Med. Hyg., 36, 143–152.

Xu, L., Stige, L.C., Chan, K.-S., Zhou, J., Yang, J., Sang, S., *et al.* (2017). Climate variation

drives dengue dynamics. *Proc. Natl. Acad. Sci.*, 114, 113–118.

Zhao, S., Musa, S.S., Hebert, J.T., Cao, P., Ran, J., Meng, J., *et al.* (2020). Modelling the

effective reproduction number of vector-borne diseases: The yellow fever

outbreak in Luanda, Angola 2015-2016 as an example. *PeerJ*, 2020.

Figure 1. Dengue incidence, climate, and susceptibles index data. Time series

(seasons 1990/1991–2008/2009) of (a) weekly dengue incidence (i.e., total number

of cases per week), (b) weekly average temperature, (c) total weekly rainfall, and

(d) a proxy for susceptible population size (see Supporting Information for details)

in San Juan, Puerto Rico.

Figure 2. **Seasonal trends and lags of dengue incidence and its drivers.** The strong seasonal signal of dengue cases and other variables suggests potential causal lags between dengue incidence and temperature, rainfall, or the proxy for the susceptible population size. The lines represent interannual averages for each week of the year (i.e., calendar week) of dengue incidence (black), temperature shifted 9 weeks forward in time (red), average rainfall over the preceding 3–9 weeks and shifted 3 weeks forward in time (blue), and susceptibles index shifted 5 weeks forward in time (purple).

Figure 3. Climate and susceptibles index drive dengue incidence. Cross-

mapping between dengue incidence and temperature (a; red), rainfall (b; blue), or

susceptibles index (c; purple) display significant (Kendall's $\tau > 0$; $P < 0.01$)

convergence in cross-mapping skill (i.e., ρ increases and reaches an asymptote) as

the length of the time series increases (a signal of putative causality). Red, blue and

purple shaded regions represent the 0.025 and 0.975 quantiles of bootstrapped

time series segments. Grey shaded regions represent the 0.025 and 0.975 quantiles

of the seasonal null distributions obtained from 500 runs of randomized time series

with conserved seasonal trends (Deyle *et al.* 2016a). Solid lines represent medians

of distributions. Rainfall and susceptibles index showed significant forcing above

and beyond seasonal signal (K-S $P < 0.0001$), because cross-mapping of the true

time series (blue and purple) are distinguishable from their respective null models

(grey), whereas temperature forcing was not distinguishable from the null (K-S $P =$

0.90).

Figure 4. Predictive power of climate and susceptibles index (λ) on in-sample (left) and out-of-sample (right) dengue incidence. Forecasting results of incidence (8 weeks ahead) are shown in turquoise (solid lines represent the mean; shaded regions represent 90% confidence intervals) and observed incidence in black. (a, c, e, g) Time series for seasons 1990/1991–2008/2009 were used to construct SSR models for forecasts using leave-one-out cross-validation. (b, d, f, h) Data for seasons 2009/2010–2012/2013 were used to evaluate the SSR models constructed in a, c, e, and g, respectively, for out-of-sample forecasts. All SSR models (a–h) had significant forecasting skill (ρ) values (Fisher's z-transformation $P <$ 0.001).

Figure 5. Temperature and rainfall show mixed effects on dengue incidence.

Scenario exploration quantified the variable effect of changes in drivers on dengue.

Boxplots show that the median effects of rainfall (Rain) and temperature (Temp) are

small (close to zero), but drivers occasionally have strong impacts (a). To investigate

climate driver functional responses, we plotted the rate of change of dengue

incidence as a function of temperature (b) and rainfall (c). Red and blue lines

represent regression on the median for temperature and rainfall, respectively, in a

quantile regression. The dashed red and blue lines represent regression on the 0.05

and 0.95 quantiles of temperature and rainfall, respectively. Temperature has an

overall positive effect on dengue incidence (median regression line of the rate of

change is positive), but can also have large negative and positive effects (a, b).

Rainfall has an overall negative effect (median regression line of the rate of change

is negative), but can also have small positive and large negative effects (a, c).

Figure 6. Temperature and rainfall effects on dengue incidence vary depending on the susceptible population size (λ). The effect of changes in temperature (a) and rainfall (b) against λ shows that driver effects are split around the threshold $\lambda \approx 0.85$ (purple dashed line). The red and blue lines represent the median regression of temperature and rainfall effects, respectively, in a quantile regression (a–d, f, g). The dashed red and blue lines represent the 0.05 and 0.95 quantile regressions of temperature and rainfall effects, respectively (a–d, f, g). Neither driver has an effect on dengue incidence when susceptible availability is low

($\lambda < 0.85$; c, f). However, when $\lambda > 0.85$ climate effects are observed: temperature

has mostly a positive effect (d), possibly sigmoidal in that temperature range (e),

and rainfall has a negative effect (g), and conceptually a concave down functional

response (h; black lines represent tangents, where the slope of the tangent is the

rate of change). The effect of temperature on relative R_0 of dengue assuming

transmission via *Aedes aegypti* mosquitoes is unimodal (Mordecai *et al.* 2017) over a

large temperature range (e; dashed lines indicate the minimum and maximum

temperature values in the data of our study, black lines represent tangents, where

the slope of the tangent is the rate of change of relative R_0 of dengue as a function of

temperature). Assuming that relative R_0 is proportional to dengue incidence, our

results suggest that the rate of change of dengue incidence is increasing until

reaching a maximum and then decreasing (d; red median regression lines).

However, even when driver effects are split at the evident threshold of $\lambda = 0.85$ (c,

d, f, g), there are still many occurrences when the susceptible population size is

sufficient large ($\lambda > 0.85$) but temperature and rainfall have no effect. In certain

cases, temperature has even a negative effect on dengue (d).