Expansions of adaptive-like NK cells with a tissue-resident phenotype in human lung and blood

Nicole Marquardt[1][*], Marlena Scharenberg[1], Jeffrey E. Mold[2], Joanna Hård[2], Eliisa Kekäläinen[3,4], Marcus Buggert[1], Son Nguyen[5,6], Jennifer N. Wilson[1], Mamdoh Al-Ameri[7], Hans-Gustaf Ljunggren[1], Jakob Michaëlsson[1]

Running title: Adaptive-like human lung tissue-resident NK cells

Affiliations:

[1]Center for Infectious Medicine, Department of Medicine, Karolinska Institutet, Stockholm, Sweden

[2]Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm, Sweden

[3]Translational Immunology Research Program & Department of Bacteriology and Immunology, University of Helsinki, Finland

[4]HUSLAB, Division of Clinical Microbiology, Helsinki University Hospital, Helsinki, Finland,

[5]Department of Microbiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[6]Institute for Immunology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[7]Thoracic Surgery, Department of Molecular Medicine and Surgery, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden

*Corresponding author/Lead contact: Nicole Marquardt, Center for Infectious

Medicine, Department of Medicine, Karolinska Institutet, 141 52 Huddinge, Sweden;

E-mail: nicole.marquardt@ki.se, Phone: +46-8-58589791 Fax: +468-7467637.

ORCID: https://orcid.org/0000-0003-3186-4752

Abstract

Human adaptive-like "memory" $CD56^{dim}CD16^+$ NK cells in peripheral blood from

cytomegalovirus-seropositive individuals have been extensively investigated in recent

years and are currently explored as a treatment strategy for hematological cancers.

However, treatment of solid tumors remains limited due to insufficient NK cell tumor

infiltration, and it is unknown whether large expansions of adaptive-like NK cells that

are equipped for tissue-residency and tumor-homing exist in peripheral tissues. Here,

we show that human lung and blood contains adaptive-like $CD56^{bright}CD16^-$ NK cells

with hallmarks of tissue-residency, including expression of CD49a. Expansions of

adaptive-like lung tissue-resident (tr)NK cells were found to be present independently

of adaptive-like $CD56^{dim}CD16^+$ NK cells and to be hyperresponsive towards target

cells. Together, our data demonstrate that phenotypically, functionally, and

developmentally distinct subsets of adaptive-like NK cells exist in human lung and

blood. Given their tissue-related character and hyperresponsiveness, human lung

adaptive-like trNK cells might represent a suitable alternative for therapies targeting

solid tumors.

Key words: NK cells, adaptive, memory, lung, tissue-resident

Abbreviations:

Eomes: Eomesodermin

FITC: Fluorescein isothiocyanate

HCMV: Human cytomegalovirus

ILC: Innate lymphoid cell

KIR: Killer cell immunoglobulin-like receptor

NK: Natural killer

PE: Phycoerythrin

ROS: Reactive oxygen species

Introduction

Natural killer (NK) cells are a crucial component of the innate immune system. They

target and eliminate virus-infected and malignant cells, and boost immunity through the

production of proinflammatory cytokines including IFN-γ and TNF. In recent years,

the concept of adaptive-like or "memory" NK cells has emerged from studies in mice

(1-4) and humans (5-10). These adaptive-like NK cells share a distinct phenotype and

increased target cell responsiveness as well as having features of longevity and superior

recall potential reminiscent of memory T cells (11).

Most studies of human adaptive-like NK cells have focused on subsets of

NKG2C$^+$(KIR$^+$)CD56^{dim}CD16$^+$ NK cells, originally found to be expanded and stably

imprinted in peripheral blood of approximately 30-40% of human CMV (HCMV)

seropositive individuals (5,10). Adaptive-like NKG2C$^+$(KIR$^+$)CD56^{dim}CD16$^+$ NK cells

in human peripheral blood have a distinctive phenotypic (5,10), epigenetic (8,9), and

functional (8-10) profile compared to conventional NK cells and have been suggested

to contribute to immunity against HCMV (1,12). Importantly, adaptive-like peripheral

blood-derived   CD56^{dim}CD16$^+$ NK   cells   (herein   defined   as   adaptive-like

CD56^{dim}CD16$^+$ pbNK cells) are currently explored for improving NK cell-mediated

cancer therapies. While adoptive NK cell transfer showed optimistic results in the

treatment of hematological malignancies (reviewed in (13)), targeting solid tumors was

less successful due to poor migration to and infiltration into the tumor (reviewed in

(13)). In these cases, adaptive-like NK cells with an increased capacity to infiltrate

tissues e.g. through co-expression of tissue-specific ligands might be desirable. In fact,

similar approaches for targeting solid tumors have recently been suggested for T$_{RM}$ cells

(14).

We   and   others   recently   identified   a   subset   of   tissue-resident

CD49a$^+$CD56^{bright}CD16$^-$ NK (trNK) cells in the human lung (15,16). The human lung

is a frequent site of acute infections, including infections with viruses such as influenza

virus, respiratory syncytial virus (RSV), and HCMV, as well as serving as a reservoir

for latent HCMV infection (17). Although human $CD56^{dim}CD16^+$ lung NK cells are

hyporesponsive to *ex vivo* target cell stimulation (18), exposure of human lung NK cells

to virus-infected cells is likely to result in functional NK cell priming and expansion of

distinct NK cell subsets. Indeed, increased polyfunctional responses have been

observed in $CD16^-$ lung NK cells following *in vitro* infection with IAV (16,19).

However, the presence of expansions of functional adaptive-like trNK cells in the

human lung is to date unknown.

Here, we identify and examine a $CD49a^+KIR^+NKG2C^+CD56^{bright}CD16^-$ NK

cell population with features of tissue-resident NK cells in human lung and blood,

which is distinct from adaptive-like $CD56^{dim}CD16^+$ pbNK cells. In donors with

expansions of adaptive-like $CD49a^+$ lung trNK cells, small but detectable frequencies

of adaptive-like $CD49a^+$ NK cells were observed in paired peripheral blood. While

adaptive-like $CD56^{dim}CD16^+$ pbNK cells (as commonly identified in peripheral blood

of HCMV-seropositive donors) and adaptive-like $CD49a^+$ trNK cells in lung and blood

shared a common core gene signature, we identified several unique features of each

population indicating that they may represent developmentally distinct populations.

Notably, NK cells from donors with an adaptive-like trNK cell expansion in the lung

were hyperresponsive towards target cells. Thus, we provide evidence indicating that

$CD49a^+KIR^+NKG2C^+CD56^{bright}CD16^-$ trNK cells in the human lung represent a

distinct population of adaptive-like NK cells with potential implications in lung

surveillance and future treatment options of solid tumors.

Results

Adaptive-like NK cells can be identified in human lung

We first set out to investigate whether expansions of adaptive-like

KIR^+NKG2C^+ NK cells could be identified in the human lung. The majority of NK

cells in the lung are phenotypically similar to pbNK cells ($CD69^-CD56^{dim}CD16^+$),

suggesting that these cells may recirculate between the lungs and peripheral blood (18).

Accordingly, circulating populations of expanded adaptive-like $CD56^{dim}CD16^+$ NK

cells could also be identified in both the peripheral blood and lungs from patients

undergoing surgery for suspected lung cancer (Fig. 1A). Surprisingly, KIR and NKG2C

were also co-expressed on less differentiated $CD56^{bright}CD16^-$ lung NK cells in many

of the patients included in the study, with varying frequencies between donors (Fig. 1B,

C) (see Supplementary Fig. 1A for the gating strategy to identify KIR^+NKG2C^+

$CD56^{bright}CD16^-$ and $CD56^{dim}CD16^+$ NK cells). In several donors the frequencies of

$KIR^+NKG2C^+CD56^{bright}CD16^-$ NK cells in human lung vastly exceeded those

previously described for $CD16^-$ NK cells in the liver (7), with up to 98% of

$CD56^{bright}CD16^-$ lung NK cells co-expressing KIR and NKG2C (Fig. 1 A, B).

Next, we assessed phenotypic features of $KIR^+NKG2C^+CD56^{bright}CD16^-$ lung

NK cells in an unbiased manner using high dimensional flow cytometry. Uniform

manifold approximation and expression (UMAP) analysis revealed a distinct subset of

cells with an expression pattern consistent with adaptive-like NK cells found in

peripheral blood and liver, including low expression of Siglec-7 and CD161, and high

expression of NKG2C, KIRs, and CD2 (7,8,20,21) (Fig. 1D). Unlike adaptive-like

$CD56^{dim}CD16^+$ pbNK cells, $KIR^+NKG2C^+CD56^{bright}CD16^-$ NK cells expressed lower

levels of CD45RA and NKp80, and higher levels of CD8 (Fig. 1D). In addition, to these

expression patterns, manual analysis of individual samples additionally confirmed low

expression of ILT2 and FcεR1γ, as compared to $KIR^+NKG2C^+CD56^{dim}CD16^+$ lung NK

cells (Fig. 1E, F). Furthermore, KIR$^+$NKG2C$^+$CD56^{bright}CD16$^-$ lung NK cells displayed

low expression of ILT2 and FcεR1γ and high expression of Eomes and NKG2A, but

similar levels in T-bet expression, revealing a phenotype distinct from human adaptive-

like KIR$^+$NKG2C$^+$CD16$^-$ NK cells in the liver (7) (Fig. 1E, F). Together, our data reveal

the presence of a previously uncharacterized population of an adaptive-like NK cell

subset, herein identified as KIR$^+$NKG2C$^+$CD56^{bright}CD16$^-$, in the human lung, which

is distinct from adaptive-like CD56^{dim}CD16$^+$ pbNK cells.

Adaptive-like CD56^{bright}CD16$^-$ NK cells in human lung are tissue-resident

TrNK cells in human lung are characterized by expression of CD69 and the

integrins CD49a and CD103 (15,16), while adaptive-like NK cells have not yet been

described within this subset. Strikingly, the vast majority of the distinct population of

NKG2C$^+$CD56^{bright}CD16$^-$ NK cells identified by UMAP analysis co-expressed CD69

(80%) and CD49a (77%), and a substantial proportion (38%) also co-expressed CD103

(Fig. 2A, B), suggesting tissue-residency of adaptive-like CD56^{bright}CD16$^-$ NK cells in

the lung. KIR$^+$NKG2C$^+$ NK cells co-expressing CD49a, CD69, or CD103, were mainly

CD56^{bright}CD16$^-$ (Fig. 2C, D), further demonstrating that they are clearly distinct from

adaptive-like CD56^{dim}CD16$^+$ pbNK cells.

To further characterize adaptive-like trNK cells in the lung, we compared the

gene expression profiles of sorted adaptive KIR$^+$NKG2C$^+$ and non-adaptive KIR$^-$

NKG2C$^-$ trNK cells in human lung using RNA sequencing (Fig. 2E, F; see sorting

strategy in Supplementary figure 1B, C). 102 genes were differentially expressed

(p<0.05, log2FC>1), including several *KIR* genes, *CD8A*, *GPR183* (EBI2)*, IRF8* and

SH2D1B (EAT2), and genes encoding for the transcription factors MafF (*MAFF*) and

ZNF498 (*ZSCAN25*). GPR183 has been demonstrated to be crucial for tissue-specific

migration of innate lymphoid cells (22), while EAT2 expression has previously been

shown to be downregulated in adaptive-like CD56^{dim}CD16^{+} pbNK cells (8).

Interestingly, while upregulation of both *IRF8* and *THEMIS2* has been reported to be

essential for NK cell-mediated responses against MCMV infection (23,24), gene

expression of both molecules was low in adaptive-like trNK cells in human lung (Fig.

2E), indicating different activation pathways in adaptive-like NK cells in mice and

humans. Furthermore, we compared differentially expressed genes in adaptive-like

lung trNK cells *versus* CD56^{dim}CD16^{+} pbNK cells and observed an overlap of

approximately one third of the differentially expressed genes (GSE117614) (25). These

genes included *KLRC2* (NKG2C), *KLRC3* (NKG2E), *IL5RA*, *GZMH*, *ITGAD* (CD11d),

RGS9, *RGS10* (upregulated), and *KLRB1* (CD161), *KLRC1* (NKG2A), *KLRF1*

(NKp80), *TMIGD2*, *IL18RAP*, *FCER1G*, *MLC1*, *CLIC3* and *TLE1* (downregulated)

(Fig. 2F). These results demonstrate that adaptive-like lung trNK and CD56^{dim}CD16^{+}

pbNK cells share a common core gene set specific for adaptive-like NK cells.

Adaptive-like NK cells in peripheral blood and in the human liver commonly

have a distinct KIR expression profile which is dominated by KIRs that bind to self-

HLA class I (self-KIRs) (7,10,26). In the lung, analysis of single KIR expression on

CD16^{-} and CD16^{+} NK cell subsets in donors with high frequencies of adaptive-like

lung trNK cells revealed that the former subset displayed unique KIR expression

patterns compared to CD16^{+} NK cells in paired blood or lung (Fig. 2G-I;

Supplementary Fig. 1D for the gating strategy). High expression of KIRs on the

adaptive-like trNK cells was limited to self-KIR, identical to the phenotype described

for adaptive-like NK cells in liver and peripheral blood. Notably, even in a donor with

adaptive-like NK cell expansions of both trNK cells and pbNK cells (Fig. 2G), the KIR-

expression profile differed between the two subsets. These results suggest a subset-

specific development and/or differentiation of adaptive-like NK cells in blood and lung.

Together, CD49a$^+$KIR$^+$NKG2C$^+$CD56^{bright}CD16$^-$ trNK cells in the human lung

exhibit a unique profile of activating and inhibitory NK cell receptors (e.g. NKG2A,

KIR, NKp80), as well as adaptor, signaling, and effector molecules (FcεR1γ, SH2D1B,

granzyme H). This indicates that these are bona fide adaptive-like trNK cells which are

distinct from adaptive-like CD56^{dim}CD16$^+$ pbNK cells, demonstrating the presence of

an as of yet unexplored NK cell population in the human lung.

Lung adaptive-like trNK cells are hyperresponsive to target cells

In order to determine whether adaptive-like lung trNK cells differed

functionally from non-adaptive CD56^{bright}CD16$^-$ lung trNK cells, we first compared

expression levels of genes associated with functional competence (Fig. 3A). Gene

expression in adaptive-like lung trNK cells was higher for *IFNG*, *IL32*, *XCL1* and

GZMH (granzyme H), and lower for *GNLY* (granulysin), *GZMA* (granzyme A), *GZMK*

(granzyme K), *IL2RB* (CD122) and *IL18RAP* as compared to non-adaptive lung trNK

cells (Fig. 3A). These results suggest a potential cytokine-mediated priming of

adaptive-like trNK cells *in vivo*, e.g. by IL-12 and IL-18, which are potent inducers of

IFN-γ and IL-32 in NK cells (27). While expression levels of the effector molecules

PRF1 (perforin) and *GZMB* (granzyme B) did not differ between adaptive- and non-

adaptive-like CD56^{bright}CD16$^-$ NK cells at transcriptome level (Fig. 3A), *ex vivo* protein

expression was increased for granzyme B in adaptive-like trNK cells (Fig. 3B, C). This

indicated a potential cytotoxic function in this particular subset, despite lung NK cells

generally being characterized as hyporesponsive to target cell stimulation (16,18).

Intriguingly, NK cells from donors with an expansion of adaptive-like trNK cells in the

lung degranulated stronger and produced more TNF compared to those from donors

without such expansions (Fig. 3D). In particular NK cells co-expressing CD49a and

KIR degranulated strongly and produced high levels of TNF upon target cell

stimulation (Fig. 3D-F). This hyperresponsiveness of adaptive-like lung trNK cells was

independent from co-expression of CD103, since similar activation levels were

observed between CD49a$^+$CD103$^-$ and CD49a$^+$CD103$^+$ KIR$^+$NKG2C$^+$ NK cells (Fig.

3F-H). Furthermore, blood NK cells from donors with expanded adaptive-like lung

trNK cells also responded stronger to target cells as compared to donors without such

expansions (Fig. 3F-H). Taken together, these results revealed that the presence of

expanded adaptive-like trNK cells in the lung is linked to hyperresponsivity towards

target cells, implicating a role of these cells in active immune regulation within the

lung.

Adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ NK cells can be identified in matched patient

peripheral blood

As a hallmark of tissue-resident cells, CD49a is commonly expressed on subsets of T

cells and NK cells in non-lymphoid compartments such as the lung (15,16,19,28), liver

(7), skin (29), uterus (30), and intestine (31), but not in peripheral blood. Intriguingly,

however, we identified a small subset of CD49a$^+$KIR$^+$NKG2C$^+$ NK cells within the

CD16$^-$ NK cell population in paired peripheral blood of donors harboring expansions

of adaptive-like trNK cells in the lung (Fig. 4A, B, see gating strategy in Supplementary

Fig. 1E). The frequencies of CD49a$^+$KIR$^+$NKG2C$^+$CD16$^-$ NK cells in peripheral blood

(herein identified as CD49a$^+$ pbNK cells) were overall considerably lower as compared

to either adaptive-like lung trNK cells or CD56^{dim}CD16$^+$ pbNK cells, respectively (Fig.

4B). We observed that 18.6% and 25.6% of all donors had an expansion (identified as

outliers) of adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ NK cells in lung and paired

peripheral blood, respectively. In comparison, 20.9% and 15.1% of all donors had an

expansion of adaptive-like CD56^{dim}CD16$^+$ pbNK cells in lung and paired blood,

respectively   (Fig.   4B).   Interestingly,   expansions   of   adaptive-like

CD49a$^+$CD56^{bright}CD16$^-$ NK and CD56^{dim}CD16$^+$ pbNK cells were virtually mutually

exclusive in donors (Fig. 4C). However, there was a substantial overlap within each of

these subsets between lung and peripheral blood (Fig. 4C), that is, e.g. a high likelihood

of having an expansion of adaptive-like trNK cells in the lung if an expansion of

adaptive-like CD49a$^+$ NK cells was present in the paired blood, and *vice versa*. This

distinct distribution of adaptive-like NK cell populations per donor suggests adaptive-

like CD49a$^+$CD56^{bright}CD16$^-$ and CD56^{dim}CD16$^-$ pbNK cells are developing

independently from each other. We next analyzed the phenotype of adaptive-like

CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells and found intermediate expression of CD57 with

relatively low co-expression of NKG2A (Fig. 4D, E), in contrast to lower expression

of CD57 and higher expression of NKG2A in the adaptive-like lung trNK cells (Fig.

1E, F). Furthermore, adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells differed from

their counterpart in lung by low expression of both CD69 and CD103 (Fig. 4E, D),

consistent with known phenotypic differences between tissue-resident and circulatory

lymphocyte populations (15,32-34).

Taken together, these results demonstrate the presence of an adaptive-like

CD49a$^+$CD56^{bright}CD16$^-$ NK cell subset in the peripheral blood of a subset of donors,

which is linked to adaptive-like trNK cells in the human lung and emerging

independently from CD56^{dim}CD16$^+$ adaptive-like pbNK cells.

Peripheral blood adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells from healthy

donors share features with lung trNK cells and adaptive-like CD56^{dim}CD16$^+$ pbNK

cells

The presence of adaptive-like lung trNK cells in patients undergoing surgery for

suspected lung cancer did not significantly correlate with any demographical or clinical

parameters including age, gender, cigarette smoking, COPD, the type of lung tumor,

survival, lung function, or HCMV IgG concentrations in plasma (Supplementary Fig.

2A-F). Since we could identify circulating CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells in the

majority of patients with an expansion of adaptive-like trNK cells in the lung, we next

sought to determine if such cells could also be detected in the peripheral blood of

unrelated healthy donors. Indeed, we found KIR$^+$NKG2C$^+$ NK cells co-expressing

CD49a in the CD56^{bright}CD16$^-$ NK cell subset in 16% of healthy blood donors (Fig. 5A,

B). The frequencies of CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells out of CD16$^-$ NK cells were

lower in healthy peripheral blood (up to 5%, mean 0.3%) as compared to those found

in patients with suspected lung cancer (up to 21%, mean 1.2%) (Fig. 5B). Within the

CD56^{bright}CD16$^-$ NK cell subset, KIR$^+$NKG2C$^+$ NK cells were almost exclusively

detected in the CD49a$^+$ population (Fig. 5C). UMAP analysis of CD56^{bright}CD16$^-$ NK

cells from healthy donors with CD49a$^+$CD56^{bright}CD16$^-$ NK cells in the blood revealed

a strong separation of the CD49a$^+$ NK cell subset co-expressing KIR and NKG2C based

on lower expression or lack of CD69, CD45RA, CD57, CD38, NKp80, and TIM-3 as

well as high expression of CD8, CXCR3 and granzyme B on CD49a$^+$KIR$^+$NKG2C$^+$

NK cells (Fig. 5D). This phenotype could be confirmed in individual samples (Fig. 5E).

Interestingly, strong expression of CXCR6 could be identified on CD69$^+$, but not on

adaptive-like CD49a$^+$ CD56^{bright}CD16$^-$ pbNK cells, indicating that the latter NK cell

subset depends on other chemokine receptors such as CXCR3 for tissue homing.

To gain further insight into the adaptive-like CD49a$^+$ pbNK cells, we sorted this

subset and compared it to sorted blood non-adaptive CD49a$^-$CD56^{bright}CD16$^-$ NK cells

using RNAseq (Fig. 6A, see Supplementary Fig. 1C for gating strategy). We next

investigated whether gene expression differences in adaptive-like CD49a$^+$ pbNK cells

indicated a different functional profile. Adaptive-like CD49a$^+$ pbNK cells expressed

particularly higher levels of *CCL5, LAMP1, GZMH* and *GNLY*, and lower levels of

XCL1, *HIF1A*, *IL2RB*, and *L18RAP* (Fig. 5F). Hence, adaptive-like CD49a$^+$ pbNK cells

and adaptive-like lung trNK cells from different donors (Fig. 3A) shared a common

gene expression pattern for some (*CCL5*, *GZMH*, *IL2RB* and *IL18RAP)*, but not all (i.e.,

GNLY) genes, indicating that they are functionally distinct from each other but also

from their non-adaptive counterparts.

To assess whether adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells segregate

further at the transcriptome level, we analyzed differentially expressed genes between

adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells and non-adaptive CD56^{bright}CD16$^-$

NK cells in peripheral blood from healthy blood donors. A total of 351 genes were

differentially expressed (padj<0.01, log2FC>1) and clearly segregated both subsets

(Fig. 6A). Since adaptive-like CD49a$^+$ pbNK cells resembled to some extent adaptive-

like trNK cells in the lung, we next sought to identify similarities to non-adaptive and

adaptive-like trNK cells also at transcriptome level. For this, we compared

differentially expressed genes in adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells

(compared to CD49a$^-$KIR$^-$CD56^{bright}CD16$^-$ NK cells in peripheral blood) and in non-

adaptive lung trNK cells (defined as CD69$^+$CD49a$^+$CD103$^+$NKG2A$^+$NKG2C$^-$CD16$^-$

NK cells) (compared to non-tissue-resident CD69$^-$CD56^{bright}CD16$^-$ NK cells in lung)

(Fig. 6B). Adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells shared 73 DEGs with

non-adaptive trNK cells in lung, including high expression of *ITGA1* (CD49a), *ZNF683*

(Hobit), *PRDM1* (Blimp-1), *CCL5*, *PIK3R1*, *PLA2G16*, *ATXN1*, as well as lower

expression of *SELL* (CD62L), *GPR183*, *IL18R1*, *IL18RAP*, *SOX4*, *RAMP1*, and

IFITM3 (Fig. 6B). All of these genes have also been shown to be differentially

expressed in trNK cells in the bone marrow and/or CD8$^+$ T$_{RM}$ cells in lung (32,34). It

should however be noted that other core-genes associated with tissue-resident

lymphocytes (e.g. *S1PR1*, *S1PR5*, *CXCR6*, *ITGAE*, *RGS1*, *KLF2*, *KLF3*, and *RIPOR2*)

were not differentially expressed between adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK

cells and non-adaptive CD56^{bright}CD16$^-$NK cells, indicating that they only partially

have a tissue-resident phenotype.

Next, we determined whether adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK

share a common gene signature with adaptive-like trNK cells and/or CD56^{dim}CD16$^+$

pbNK cells. Indeed, adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells shared

differentially expressed genes with adaptive-like trNK cells in lung (Fig. 6C) and/or

CD56^{dim}CD16$^+$ pbNK cells (8,25), including increased expression of *KIRs*, *KLRC2*,

GZMH, *ITGAD*, *CCL5*, *IL32*, *ZBTB38*, *CD3E*, *ARID5B*, *MCOLN2*, and *CD52*, and

decreased expression of *KLRB1*, *FCER1G*, *IL18RAP*, *IL2RB2*, *TLE1*, *AREG*, and

KLRF1 (Fig. 6A, C, Supplementary Fig. 3).

Taken together, adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells share traits with

both non-adaptive lung trNK cells, and adaptive-like trNK and CD56^{dim}CD16$^+$ pbNK

cells.

Discussion

Human adaptive-like NK cells have been described within the $CD56^{dim}CD16^+$ subset in peripheral blood (5,6,8,10,35) and the $CD56^{bright}CD16^-$ subset in liver (7). Here, we identified and characterized a yet unexplored and unique subset of adaptive-like $CD49a^+KIR^+NKG2C^+CD56^{bright}CD16^-$ trNK cells in the human lung, paired blood, and in unrelated healthy human blood. Lung adaptive-like trNK cells shared several phenotypic features with other adaptive-like NK cell subsets both in blood and/or liver, including high expression of CD49a (liver), CD69 (liver), CD2 (blood) and lack of, or decreased expression of CD57 (liver), CD45RA (liver) and perforin (liver), as well as low expression of FcεR1γ and Siglec-7 (blood) (5,7-10,21,35,36). However, lung adaptive-like trNK cells segregate from liver adaptive-like trNK cells on the basis of high expression of Eomes and CD103, and from adaptive-like $CD56^{dim}CD16^+$ pbNK cells by lack of CD57 and a $CD56^{bright}CD16^-$ phenotype (7). Transcriptome analysis revealed shared core genes in adaptive-like trNK cells, and $CD56^{dim}CD16^+$ and $CD56^{bright}CD16^-$ pbNK cells, underlining common features between all adaptive-like NK cell populations. Intriguingly, lung adaptive-like trNK cells were highly target cell-responsive, and the overall paired blood and lung NK cell populations were hyperresponsive in donors with expansions of adaptive-like trNK cells in the lung. These findings indicate *in vivo* priming akin to what has been observed previously in human antigen-dependent (3,37), antigen-independent, and cytokine-dependent (2,38-40) NK cell recall responses. Furthermore, adaptive-like $CD56^{dim}CD16^+$ pbNK cells have recently been shown to be functionally primed against target cells following IL-12 and IL-18 stimulation, while showing only a poor response to these cytokines alone (41). In line with our results, this emphasizes a role of potential cytokine-mediated priming of adaptive-like trNK cells in the human lung.

Despite the indications of *in vivo* priming, the presence of expansions of

adaptive-like CD49a⁺CD56^bright^CD16⁻ NK cells in the lung donors did not correlate

with presence or kind of tumor, HCMV serostatus, or clinical and demographic

parameters. In fact, we could identify adaptive-like CD49a⁺CD56^bright^CD16⁻ NK cells

also in the peripheral blood of healthy donors. These adaptive-like

CD49a⁺CD56^bright^CD16⁻ pbNK cells shared a gene signature with trNK cells in the

human lung, indicating tissue imprinting. These findings suggest re-entry of adaptive-

like trNK cells from tissue into circulation and, hence, potential seeding of tissues with

adaptive-like trNK cells via peripheral blood. In mice, CD8⁺ T effector cells egress

from infected lung in a tightly regulated manner following infection with influenza A

virus (42), and CD8⁺ T_{RM} cells wane over time in self-limiting viral infections of the

respiratory tract (43). MCMV-specific CD8⁺ T cells convert to CD103⁺ T_{RM} cells, with

small numbers of new T_{RM} cells deriving from the circulation (44), and memory

inflation is required for retention of CD8⁺ T_{RM} cells in the lungs after intranasal

vaccination with MCMV (45). This indicates a dynamic retention of T_{RM} cells by a

persistent infection. It remains to be determined whether virus-dependent expansion

and maintenance of T_{RM} cells is analogous in adaptive-like trNK cells in the lung.

However, our data indicate that T_{RM} cells and adaptive-like NK cells differ at least in

their recruitment to the lung, with T_{RM} cells being dependent on CXCR6 (46) while

adaptive-like CD49a⁺CD56^bright^CD16⁻ pbNK cells lacked CXCR6 but expressed high

levels of CXCR3.

Adaptive-like CD49a⁺CD56^bright^CD16⁻ NK cell expansions were rarely

observed in donors with adaptive-like CD56^dim^CD16⁺ pbNK cell expansions, indicating

that these two distinct subsets have different developmental cues. Indeed, even in the

rare cases where we could detect expansions of both adaptive-like

CD49a⁺CD56^bright^CD16⁻ NK cells and CD56^dim^CD16⁺ pbNK cells in the same

individual, these populations displayed specific and individual KIR repertoires.

Furthermore, expansions of adaptive-like lung trNK cells were detected in HCMV-

seronegative individuals (Supplementary Fig. 2E, F), while expansions of adaptive-like

CD56^{dim}CD16$^+$ pbNK cells were restricted to HCMV-seropositive individuals

(Supplementary Fig. 2E) (5,35). It thus remains possible that other viral infections than

CMV in humans and mice (1,47,48) could drive the expansion of adaptive-like trNK

cells, as has previously been suggested for the generation of cytokine-induced memory

NK cells, e.g. after influenza virus infection in humans (49) and mice (4,50,51), as well

as vesicular stomatitis virus (VSV) (4), vaccinia virus (52), HIV-1 (4), and herpes

simplex virus 2 (HSV-2) (51), and after immunization with simian immunodeficiency

virus (SIV) in rhesus macaques (53). Taken together, our data support a model where

adaptive-like trNK cells and CD56^{dim}CD16$^+$ pbNK cells develop independently from

each other, possibly due to distinct environmental requirements for their expansion.

We observed increased gene expression levels of *GZMH* in adaptive-like

CD49a$^+$CD56^{bright}CD16$^-$ lung and blood NK cells, and levels for *CCL5* were higher in

both conventional and adaptive-like lung trNK cells as compared to CD69$^-$

CD56^{bright}CD16$^-$ lung NK cells, and particularly highly expressed in adaptive-like

CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells. An antiviral activity has been proposed for

granzyme H (54,55), however, a direct association of this effector molecule with

adaptive-like NK cells remains to be determined. In contrast, *CCL5* and *XCL1*, which

are both upregulated in human CD49a$^+$CD56^{bright}CD16$^-$ adaptive-like NK cells, were

predominantly produced by mouse Ly49H$^+$ NK cells upon stimulation with MCMV-

derived m157 protein (56), and CCL5 has been shown to be specifically expressed by

CD8$^+$ memory T$_{EM}$ cells (57). Thus, adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ blood and

lung NK cells share functional characteristics with other memory lymphocyte

populations. Furthermore, we showed that adaptive-like lung trNK cells were

hyperresponsive against target cells, hence, they might be clinically relevant e.g. in

disease progression in respiratory viral infections and/or the defense against malignant

tumor cells. Similarly, lung CD8$^+$ T$_{RM}$ cells have previously been shown to be able to

control tumor growth and to correlate with increased survival in lung cancer patients

(58). Since adaptive-like trNK cells likely exceed their circulating pbNK cell

counterpart in tissue-homing and tumor-infiltration based on their expression of tissue-

specific receptors such as CD49a and CXCR3, these cells could be harnessed for future

treatment options of solid tumors.

Together, our data reveal the presence of a yet unexplored and distinct adaptive-

like trNK cell subset in the human lung, indicating that adaptive-like NK cells are not

confined to peripheral blood and/or liver and that different lineages of adaptive-like NK

cells potentially exist. Expansions of adaptive-like trNK cells in the lung were

commonly accompanied by the presence of adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ NK

cells in paired peripheral blood, enabling the non-invasive identification of donors with

potential adaptive-like lung trNK cell expansions as well as the isolation of adaptive-

like NK cells with tissue-resident characteristics. Finally, adaptive-like NK cells with

tissue-resident features and excessive functional responsiveness in the human lung and

blood could be an attractive source for tailored cancer immunotherapies, in particular

for targeting solid tumors.

Acknowledgements

The authors acknowledge support from the National Genomics Infrastructure in

Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg

Foundation, the Swedish Research Council, and SNIC/Uppsala Multidisciplinary

Center for Advanced Computational Science for assistance with massively parallel

sequencing and access to the UPPMAX computational infrastructure. The authors also

acknowledge the MedH Core Flow Cytometry Facility (Karolinska Institutet),

supported by Karolinska Institutet and Region Stockholm, for providing cell-sorting

services. Furthermore, the authors want to thank all donors for participating in the study

and A.-C. Orre, V. Jackson, and S. Hylander for administrative and clinical help as well

as E. Yilmaz for assistance in the laboratory. **Funding:** This work was supported by

the Swedish Research Council, the Strategic Research Foundation, the Swedish

Foundation for Strategic Research, the Swedish Cancer Society, Sweden's Innovation

Agency, the Eva and Oscar Ahréns Research Foundation, Stockholm, the Åke Wiberg

Foundation and the Tornspiran Foundation.

Author contribution: Conceptualization: N.M., H.G.L., Ja.M.; Methodology: N.M.,

Ja.M.; Investigation: N.M., M.S., Je.M., J.H., E.K., M.B., S.N., J.N.W.; Resources:

M.A.-A.; Writing – original draft: N.M., Ja.M.; Writing – review and editing: N.M.,

Ja.M., H.G.L; Visualization: N.M., Ja.M.; Funding acquisition: N.M., Ja.M., H.G.L.

Competing Interests: The authors declare no competing financial interest.

     Figure legends

Figure 1: Adaptive-like KIR$^+$NKG2C$^+$ NK cells exist in the CD56^{bright}CD16$^-$ NK

cell subset in the human lung. (A) Representative overlay displaying pan-KIR and

NKG2C expression on CD56^{dim}CD16$^+$ NK cells in paired blood (black contour) and

lung (orange). **(B)** Representative dot plots displaying pan-KIR and NKG2C expression

on CD56^{bright}CD16$^-$ NK cells in the lungs of three different donors. **(C)** Summary of

data showing the frequencies of KIR$^+$NKG2C$^+$ NK cells in CD56^{dim}CD16$^+$ and

CD56^{bright}CD16$^-$ NK cells in paired blood and lung (n = 77). Friedman test, Dunn's

multiple comparisons test. ***p<0.001, ****p<0.0001. **(D)** UMAP analysis of

CD56^{bright}CD16$^-$ lung NK cells from four donors with 2,000 events/donor (942 events

in one of the donors). UMAPs were constructed using expression of Siglec-7, CD8,

CD2, CD57, CD161, NKG2C, CD56, CD45RA, NKG2A and NKp80. Color scale

shows log2(normalized expression + 1). **(E)** Representative histograms and **(F)**

summary of data showing surface expression of NKG2A (n = 27), CD57 (n = 27),

Siglec-7 (n = 7), CD161 (n = 12), CD2 (n = 5), ILT2 (n = 6), CD8 (n = 20), NKp80 (n

= 6), CD45RA (n = 5), and intracellular expression of FcεR1γ (n = 4), Eomes (n = 7)

and T-bet (n = 6) in KIR$^+$NKG2C$^+$ NK cells in CD56^{dim}CD16$^+$ blood (grey) and lung

(orange) NK cells and CD56^{bright}CD16$^-$ lung NK cells (blue). Friedman test, Dunn's

multiple comparisons test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 2: Adaptive-like CD56^{bright}CD16$^-$ lung NK cells are tissue-resident. (A)

Overlay of UMAPs from data from Figure 1, showing the position of NKG2C$^+$ (blue)

and NKG2C$^-$ (grey) populations among CD56^{bright}CD16$^-$ NK cells. **(B)** Expression of

the tissue-residency markers CD69, CD49a, and CD103 within the UMAP of

CD56^{bright}CD16$^-$ lung NK cells. **(C)** Representative histograms and **(D)** summary of

data showing the expression of the tissue-residency markers CD69 (n = 23), CD49a (n=

21) and CD103 (n = 21) on CD56^{dim}CD16$^+$ blood (grey) and lung (orange) NK cells

and CD56^{bright}CD16$^-$ lung NK cells (blue), respectively. Friedman test, Dunn's multiple

comparisons test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. **(E)** Heatmap

showing 102 differentially expressed genes between KIR$^+$NKG2C$^+$ trNK cells and

KIR$^-$NKG2C$^-$ trNK cells in the human lung. Differentially expressed genes shared with

CD57$^+$NKG2C$^+$CD56^{dim}CD16$^+$ adaptive-like NK cells in blood (from GSE117614) are

highlighted in red. **(F)** Log2 fold-change for non-adaptive (NKG2C$^-$) trNK cells vs

adaptive-like trNK cells in lung against log2 fold change for CD57$^+$NKG2C- vs

adaptive-like CD57$^+$NKG2C$^+$ CD56^{dim} NK cells in blood. Data for CD56^{dim} NK cells

in peripheral blood are from GSE117614 (25). **(G-I)** Single KIR expression analysis on

CD56^{dim}CD16$^+$ NK cells from peripheral blood (red), CD49a$^-$CD103$^-$CD56^{dim}CD16$^+$

or CD103$^-$CD56^{dim}CD16$^+$ (black) and CD49a$^+$CD103$^+$CD56^{bright}CD16$^-$ or

CD103$^+$CD56^{bright}CD16$^-$ (blue) NK cells in matched lung of three different donors.

Educating KIR are highlighted in red. **(G)** Donor with expansions of self-KIR$^+$ NK cells

both in the CD56^{dim}CD16$^+$ subset in paired blood and lung and in the CD56^{bright}CD16$^-$

NK cell subset in the lung. **(H)** Donor with an expansion of self-KIR$^+$ NK cells

exclusively in the CD56^{bright}CD16$^-$ NK cell subset in the lung. **(I)** Donor with

expansions of self-KIR$^+$ NK cells both in the CD56^{dim}CD16$^+$ and CD56^{bright}CD16$^-$

subsets in paired blood and lung, respectively.

Figure 3: Adaptive-like lung trNK cells are highly functional. (A) Gene expression

levels (counts per million reads) for selected genes associated with functional capacity

are shown for non-adaptive CD49a$^+$KIR$^-$NKG2C$^-$ and adaptive-like

CD49a$^+$KIR$^+$NKG2C$^+$ lung trNK cells (clear circles: CD49a$^+$CD103$^-$ NK cells, filled

circles: CD49a$^+$CD103$^+$ NK cells). Mean ± SEM is shown. **(B)** Representative

histograms and **(C)** summary of data displaying expression of perforin and granzyme

B (GzmB) (n = 4) in $CD56^{dim}CD16^+$ and in non-adaptive $CD49a^-CD56^{bright}CD16^-$ as

well as adaptive-like $CD49a^+CD56^{bright}CD16^-$ NK cells, respectively, in human lung *ex*

vivo. $CD14^-CD19^-CD3^-CD45^+CD127^+CD161^+$ cells were gated as controls in (B).

Friedman test, Dunn's multiple comparisons test. *p<0.05. **(D)** Representative dot plots

showing expression of CD107a and CD49a on KIR^-NKG2C^- and KIR^+NKG2C^+ NK

cells in a donor without (upper panel) and with (lower panel) expansion of adaptive-

like trNK cells in the lung (expression KIR and NKG2C are displayed in the left panel

for each of the two donors). **(E)** Summary of data showing the frequency of K562 target

cell-induced $CD107a^+$ (left) and TNF^+ (right) NK cell subsets from donors with NK

cell expansions in the human lung. Responses by unstimulated controls were subtracted

from stimulated cells (n = 4). Mean ± SD is shown. **(F)** Representative dot plots

showing expression of CD107a and TNF vs CD103 on non-adaptive $CD49a^+KIR^-$

$NKG2C^-$ (upper panel) or adaptive-like $CD49a^+KIR^+NKG2C^+$ (lower panel) bulk NK

cells in a donor with an expansion of adaptive-like trNK cells in the lung. **(G, H)**

Summary of data showing the frequencies of **(G)** $CD107a^+$ and **(H)** TNF^+ NK cells in

blood NK cells and in subsets of lung NK cells ($CD49a^-CD103^-$, expressing either

CD49a or CD103, or $CD49a^+CD103^+$) from donors without (left panels, n = 5 for

CD107a, n= for TNF) or with (right panels, n = 5) expansions of KIR^+NKG2C^+ trNK

cells in the lung. Responses by unstimulated controls were subtracted from stimulated

cells. (G, H) Violin plots with quartiles and median are shown. Friedman test, Dunn's

multiple comparisons test. *p<0.05.

Figure 4: Expansions of adaptive-like trNK cell in the lung indicate presence of

adaptive-like $CD49a^+CD56^{bright}CD16^-$ NK cells in paired blood. (A) Representative

dot plots displaying expression of CD49a and NKG2C on NK cells in lung and paired

peripheral blood. **(B)** Summary of data of frequencies of adaptive-like $CD49a^+$ NK cells

of CD16⁻ trNK cells and of adaptive-like pbNK cells in the CD16⁺ NK cell subset in

paired lung and peripheral blood. Adaptive-like NK cell "expansions" were identified

as outliers (filled circles) using the Robust regression and Outlier removal (ROUT)

method (ROUT coefficent Q=1). Error bars show the median with interquartile range

(n = 86). Median with interquartile range is shown. **(C)** Euler diagram indicating

overlaps and relationships between adaptive-like trNK and pbNK cell expansions in

peripheral blood and lung. The number of individuals with overlaps between the subsets

and compartments are indicated in the circles. **(D)** Representative overlays and **(E)**

summary of data showing phenotypic differences between adaptive-like CD49a⁺ (blue)

and non-adaptive CD49a⁻ (grey) NK cells within the CD56^{bright}CD16⁻ NK cell subset

in blood. (NKG2A, n=6; CD57, n=5; CD69, n=6; CD103, n=6; CD127, n=3; CD161,

n=4). Violin plots with quartiles and median are shown.

Figure 5: Expansions of adaptive-like CD49a⁺CD56^{bright}CD16⁻ NK cells in healthy

blood donors. (A) Representative dot plot (left plot) and overlay (right plot) showing

expression of KIR and NKG2C (left plot), and CD49a on adaptive-like KIR⁺NKG2C⁺

NK cells (blue) versus non-adaptive KIR⁻NKG2C⁻ NK cells (grey) (right plot) within

CD16⁻ blood NK cells of healthy blood donors. **(B)** Identification of expansions (filled

circles) of adaptive-like CD56^{dim}CD16⁺ pbNK cells (21 outliers, 20%) and adaptive-

like CD49a⁺CD56^{bright}CD16⁻ pbNK cells (17 outliers, 16%) via the ROUT method (see

also Figure 4E). Error bars show the median with interquartile range. (n=95). Median

with interquartile range is shown. **(C)** Frequencies of KIR⁺NKG2C⁺ cells of CD49a⁻

CD16⁻ or CD49a⁺CD16⁻ NK cells in healthy blood. The respective maternal population

comprised at least 45 cells (n=13). Wilcoxon matched-pairs signed rank test. **p<0.005

(D) UMAPs based on CD56^{bright}CD16⁻ NK cells from three donors with

KIR⁺NKG2C⁺CD56^{bright}CD16⁻ NK cells. UMAPs were constructed using expression

of CXCR3, CD161, Ki67, NKG2C, CD103, TIGIT, perforin, granzyme B, NKG2A,

CD16, CD56, CD49a, CD38, CD8, CXCR6, CD4, CD57, CD45RA, NKp80, CD69,

GL183/EB6 (KIR), and CD127. Color scale indicates log2(normalized protein

expression +1) for each parameter. **(E)** Summary of protein expression on adaptive-like

CD49a$^+$ NK cells from peripheral blood from healthy donors. (CD69, n=11; CD103,

n=7; CD57, n=11; NKG2A, n=11; CD127, n=7; CD161, n=7; CD8, n=11; CD38, n=5;

CD45RA, n=4; NKp80, n=5; TIM-3, n=5; CXCR3, n=4; CXCR6, n=5; Ki67, n=5;

perforin, n=7; granzyme B, n=5). Violin plots with quartiles and median are shown. **(F)**

Gene expression levels (counts per million reads) for selected genes associated with

functional capacity are shown for CD49a$^-$KIR$^-$ and CD49a$^+$KIR$^+$ blood

CD56^{bright}CD16$^-$ NK cells. Mean ± SEM is shown.

Figure 6: Adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ pbNK cells in healthy blood

donors share traits with both trNK cells and with adaptive-like trNK cells. (A)

Heatmap showing 138 differentially expressed genes (padj<0.001, log2FC>2) between

adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ NK cells and non-adaptive CD49a$^-$

CD56^{bright}CD16$^-$ NK cells in peripheral blood from unrelated healthy donors (n=4).

Genes shared with trNK cells in the lung were highlighted in dark blue, shared with

adaptive-like trNK cells in bright blue, and genes shared with adaptive-like

CD56^{dim}CD16$^+$ pbNK cells in orange. **(B)** Log2 fold-change for trNK cells vs non-

tissue-resident CD56^{bright}CD16$^-$ NK cells in lung against log2 fold-change for adaptive-

like CD49a$^+$CD56^{bright}CD16$^-$ vs non-adaptive CD49a$^-$CD56^{bright}CD16$^-$ NK cells in

blood. **(C)** Log2 fold-change for KIR$^+$NKG2C$^+$ trNK cells vs NKG2C$^-$ trNK cells in

lung against log2 fold-change for adaptive-like CD49a$^+$CD56^{bright}CD16$^-$ vs non-

adaptive CD49a$^-$CD56^{bright}CD16$^-$ NK cells in blood.

Table 1. Clinical and demographic details of the 103 patients included in the study.

	Never-smoker (n = 16)	Current smoker (n = 25)	Ex-smoker (n = 62)
Female / male	8 / 8	14 / 11	41 / 21
Age (y), mean ± SD	67 ± 13.4	67 ± 8.0	69 ± 7.1
FEV1/FVC (% of predicted) mean ± SD	100 ± 10.3	88 ± 13.7[*]	93 ± 14.9[#]
Pathology	*% (n)*	*% (n)*	*% (n)*
Non-malignant	6 (1)	4 (1)	2 (1)
Adenocarcinoma	44 (7)	52 (13)	76 (47)
Large cell carcinoma	0 (0)	12 (3)	3 (2)
Squamous cell carcinoma	6 (1)	16 (4)	5 (3)
Metastasis	0 (0)	4 (1)	3 (2)
Carcinoid	31 (5)	8 (2)	6 (4)
Adenosquamous carcinoma	6 (1)	0 (0)	3 (2)
Other[§]	6 (1)	4 (1)	2 (1)
Medication	*% (n)*	*% (n)*	*% (n)*
Inhaled corticosteroids	19 (3)	0 (0)	2 (1)
Statins	38 (6)	28 (7)	31 (19)
Systemic immunosuppression	0 (0)	4 (1)	5 (3)
Beta-agonists or anti-cholinergics	0 (0)	12 (3)	13 (8)
Inhaled corticosteroid and long-acting beta-agonist combination	0 (0)	8 (2)	6 (4)
Diagnoses affecting lung function	*% (n)*	*% (n)*	*% (n)*
Asthma	19 (3)	0 (0)	5 (3)
COPD	0 (0)	32 (8)	10 (6)
Other lung parenchyme disease	6 (1)	0 (0)	2 (1)

575   * n = 24

576   # n = 59

577   § = uncertain histopathological diagnosis, data missing, combined small cell carcinoma

Material and methods

Lung patients and healthy blood

A total of 103 patients undergoing lobectomy for suspected lung cancer were included in this study for collection of lung tissue and paired peripheral blood. None of the patients received preoperative chemotherapy and/or radiotherapy. Patients with records of strong immunosuppressive medication and/or hematological malignancy were excluded from the study. Clinical and demographic details are summarized in Table 1. Furthermore, healthy blood was collected from regular, unrelated blood donors. The regional review board in Stockholm approved the study, and all donors gave informed written consent prior to collection of samples.

Processing of tissue specimens and peripheral blood

Lung tissue was processed as previously described (18). Briefly, a small part of macroscopically tumor-free human lung tissue from each patient was transferred into ice-cold Krebs-Henseleit buffer and stored on ice for less than 18 h until further processing. The tissue was digested using collagenase II (0.25 mg/ml, Sigma-Aldrich) and DNase (0.2 mg/ml, Roche), filtered and washed in complete RPMI 1640 medium (Thermo Scientific) supplemented with 10% FCS (Thermo Scientific), 1 mM L-glutamine (Invitrogen), 100 U/ml penicillin, and 50 µg/ml streptomycin (R10 medium). Finally, mononuclear cells from the lung cell suspensions and peripheral blood were isolated by density gradient centrifugation (Lymphoprep).

RNA-sequencing and RNAseq data analysis

RNA of sorted NK cell subsets from blood and lung were sequenced and analyzed as described previously (15). Briefly, RNAseq was performed using a modified version of the SMART-Seq2 protocol (59). For analysis of lung adaptive-like NK cells, live

NKG2C$^+$KIR$^+$CD3$^-$CD14$^-$CD19$^-$CD56$^+$CD16$^-$ NK cells were sorted from two donors

and were compared to previously published data on CD69$^+$CD49a$^+$CD103$^-$ and

CD69$^+$CD49a$^+$CD103$^+$ NKG2A$^+$CD16$^-$ trNK cells (GSE130379) (15). For analysis of

KIR$^+$CD49a$^+$ CD56^{bright}CD16$^-$ NK cells, we sorted KIR$^+$CD49a$^+$ and KIR$^-$CD49a$^-$ live

CD14$^-$CD19$^-$CD3$^-$CD56^{bright}CD16$^-$ NK cells from cryopreserved PBMCs from 4

donors. Duplicates of 100 cells from each population from two individual donors were

sorted into 4.2ul of lysis buffer (0.2% Triton X-100, 2.5uM oligo-dT (5′-

AAGCAGTGGTATCAACGCAGAGTACT30VN-3′), 2.5mM dNTP, RNAse

Inhibitor (Takara), and ERCC RNA spike in controls (Ambion)) in a 96-well V-bottom

PCR plate (Thermo Fisher). Sorted cells were then frozen and stored at -80°C until they

could be processed. Subsequent steps were performed following the standard SMART-

Seq2 protocol with 22 cycles of cDNA amplification and sample quality was

determined using a bioanalyzer (Agilent, High Sensitivity DNA chip). 5ng of amplified

cDNA was taken for tagmentation using a customized in-house protocol (60) and

Nextera XT primers. Pooled samples were sequenced on a HiSeq2500 on high output

mode with paired 2x125bp reads.

Transcriptome analysis

Following sequencing and demultiplexing, read pairs were trimmed from Illumina

adapters using cutadapt (version 1.14) (61), and UrQt was used to trim all bases with a

phred quality score below 20 (62). Read pairs were subsequently aligned to the protein

coding sequences of the human transcriptome (gencode.v26.pc_transcripts.fa) using

Salmon (version 0.8.2) (63), and gene annotation using gencode.v26.annotation.gtf.

DeSeq2 (64) was used to analyze RNA-seq data in R studio version 1.20. Briefly, raw

count values were used as input into deSeq2, and variance stabilizing transformation

was used to transform data. Data were batch- and patient corrected using Limma (65).

A cut-off of >100 counts across the samples was used to filter out low expressed genes.

Genes with an adjusted p-value<0.05 and a log2-fold change greater than 1 were

considered as differentially expressed between paired samples. Similarly, previously

published data sets on adaptive-like NKG2C$^+$CD57$^+$CD56^{dim}CD16$^+$ NK cells and

conventional NKG2C$^-$CD57$^+$CD56^{dim} NK cells (GSE117614) (25) were analyzed using

deSeq2 to identify differentially expressed genes. Heatmaps of gene expression were

generated using Pheatmap in R and show the z-score for differentially expressed genes

(as determined above in deSeq2) for all donors and replicates.

Flow cytometry

Antibodies and clones reactive against the following proteins were used: CD2 (TS1/8,

BV421 or Pacific Blue, Biolegend), CD3 (UCHT1, PE-Cy5, Beckman Coulter), CD8

(RPA-T8, Brilliant Violet 570, Biolegend, or RPA-8, BUV395 or SK1, BUV737, BD

Biosciences), CD14 (MφP9, Horizon V500, BD Biosciences), CD16 (3G8, Brilliant

Violet 570 or Brilliant Violet 711, or Brilliant Violet 785, Biolegend), CD19 (HIB19,

Horizon V500, BD Biosciences), CD38 (HIT2, Brilliant Violet 711 or BUV661, BD

Biosciences), CD45 (HI30, Alexa Fluor 700, Biolegend, or BUV805, BD Biosciences),

CD45RA (HI100, Brilliant Violet 785, Biolegend), CD49a (TS2A, AlexaFluor 647,

Biolegend, or HI30, BUV615, or 8R84, Brilliant Violet 421, BD Biosciences), CD56

(N901, ECD, Beckman Coulter, or HCD56, Brilliant Violet 711, Biolegend, or

NCAM16.2, PE-Cy7, or BUV563, BD Biosciences), CD57 (TB01, purified,

eBioscience, or HNK-1, Brilliant Violet 605, Biolegend), CD103 (APC, B-Ly7,

eBioscience, or biotin, 2G5, Beckman Coulter, or Ber-ACT8, Brilliant Violet 711,

BUV395, BD Biosciences, or Ber-ACT8, PE-Cy-7, Biolegend), KIR2DL1

(FAB1844F, biotin, R&D Systems), KIR2DL3 (180701, FITC, R&D Systems),

KIR3DL1 (DX9, Brilliant Violet 421, Biolegend), KIR3DL2 (DX-31, Brilliant Violet

711, Biolegend), KIR2DL2/S2/L3 (GL183, PE-Cy5.5, Beckman Coulter),

KIR2DL1/S1 (EB6, PE-Cy5.5 or PE-Cy7, Beckman Coulter), NKG2A (Z1991.10,

APC-A780, or PE, Beckman Coulter, or 131411, BUV395, BD Biosciences), NKG2C

(134591, Alexa-Fluor 488 or PE, R&D Systems), CD69 (TP1.55.3, ECD, Beckman

Coulter, or FN50, PE-CF594, BD Biosciences, or FN50, Brilliant Violet 786,

Biolegend), CD127 (Brilliant Violet 421, HIL-7R-M21, BD Biosciences or PE-Cy7,

R34.34, Beckman Coulter), CD161 (HP3-3G10, Brilliant Violet 605 or APC/Fire 750,

Biolegend), CXCR3 (Alexa Fluor 647, G025H7, Biolegend), CXCR6 (K041E5,

Brilliant Violet 421, Biolegend), CD85j/ILT2 (HP-F1, Super Bright 436, Invitrogen),

NKp80 (5D12, PE, BD Biosciences, or 4A4.D10, PE-Vio770, Miltenyi), Siglec-7 (5-

386, Alexa Fluor 488, Bio-Rad), TIM-3 (7D3, Brilliant Violet 711, BD Biosciences).

After two washes, cells were stained with streptavidin Qdot 605 or Qdot 585 (both

Invitrogen), anti-mouse IgM (II/41, eFluor 650NC, eBioscience) and Live/Dead Aqua

(Invitrogen). After surface staining, peripheral blood mononuclear cells (PBMC) were

fixed and permeabilized using FoxP3/Transcription Factor staining kit (eBioscience).

For intracellular staining the following antibodies were used: Eomes (WF1928, FITC,

eBioscience), FcεR1γ (polyclonal, FITC, Merck), granzyme B (GB11, BB790, BD

Biosciences), Ki67 (B56, Alexa Fluor 700, BD Biosciences), perforin (dG9, BB755,

BD Biosciences, or B-D48, Brilliant Violet 421, Biolegend), T-bet (4B10, Brilliant

Violet 421, BD Biosciences), and TNF (MAb11, Brilliant Violet 421, Biolegend, or

Brilliant Violet 650, BD Biosciences). Purified NKG2C (134591, R&D Systems) was

biotinylated using a Fluoreporter Mini-biotin XX protein labeling kit (Life

Technologies) and detected using streptavidin-Qdot 585, 605 or 655 (Invitrogen).

Samples were analyzed on a BD LSR Fortessa equipped with four lasers (BD

Biosciences) or a BD FACSymphony A5 equipped with five lasers (BD Biosciences),

and data were analyzed using FlowJo version 9.5.2 and version 10.6.1 (Tree Star Inc).

UMAPs were constructed in FlowJo 10.6.1 using the UMAP plugin. UMAP

coordinates and protein expression data were subsequently exported from FlowJo, and

protein expression for each parameter was normalized to a value between 0 and 100.

UMAP plots were made in R using ggplot, and color scale show log2(normalized

protein expression +1).

For sorting of NK cells from lung and peripheral blood for RNA sequencing,

thawed cryopreserved mononuclear cells were stained with anti-human CD57 (NK-1,

FITC, BD Biosciences), CD16 (3G2, Pacific Blue, BD Biosciences), CD14 (MφP9,

Horizon V500, BD Biosciences), CD19 (HIB19, Horizon V500, BD Biosciences),

CD103 (Ber-ACT8, Brilliant Violet 711, BD Biosciences), CD49a (TS2/7, Alexa Fluor

647, Biolegend), CD45 (HI30, A700, Biolegend), CD8 (RPA-T8, APC/Cy7, BD

Biosciences), NKG2A (Z199.10, PE, Beckman Coulter), CD69 (TP1.55.3, ECD,

Beckman Coulter), CD3 (UCHT1, PE/Cy5, Beckman Coulter), KIR2DL1/S1 (EB6,

PE/Cy5.5, Beckman Coulter), KIR2DL2/3/S2 (GL183, PE/Cy5.5, Beckman Coulter,),

NKG2C (134591, biotin, R&D Systems, custom conjugate), CD56 (NCAM16.1,

PE/Cy7, BD Biosciences), streptavidin Qdot655 (Invitrogen), and Live/Dead Aqua

(Invitrogen).

DNA isolation and KIR/HLA-ligand genotyping

Genomic DNA was isolated using a DNeasy Blood & Tissue Kit (Qiagen) from 100 µl

of whole blood. KIR genotyping and KIR ligand-determination were performed using

PCR-SSP technology with a *KIR* typing kit and a *KIR HLA* ligand kit (both Olerup-

SSP) according to the manufacturer's instructions.

CMV IgG ELISA

Concentrations of anti-CMV IgG relative to a standard curve and internal negative and

positive control were determined by ELISA (Abcam, UK) and read in a microplate

spectrophotometer (Bio-Rad xMark) at 450nm with a 620nm reference wavelength.

Activation assay

Degranulation and TNF production of fresh blood and lung NK cells were assessed as

previously described (18,19). In brief, fresh lung and blood mononuclear cells were

resuspended in R10 medium and rested for 15 to 18 hours at 37°C. Subsequently, the

cells were co-cultured in R10 medium alone or in presence of K562 cells for 2 hours in

the presence of anti-human CD107a (FITC or Brilliant Violet 421, H4A3, BD

Biosciences, San Jose, Calif.).

Statistics

GraphPad Prism 6 and 7 (GraphPad Software) was used for statistical analysis.

For each analysis, measurements were taken from distinct samples. The statistical

method used is indicated in each figure legend.

Data Availability

The dataset generated for this study can be found in the Gene Expression Omnibus with

accession no. xxxx (data will be deposited and made available before publication).

References

1. Sun JC, Beilke JN, Lanier LL. Adaptive immune features of natural killer cells. Nature. 2009 Jan 29;457(7229):557–61.

2. Cooper MA, Elliott JM, Keyel PA, Yang L, Carrero JA, Yokoyama WM. Cytokine-induced memory-like natural killer cells. Proceedings of the National Academy of Sciences of the United States of America. 2009 Feb 10;106(6):1915–9.

3. O'Leary JG, Goodarzi M, Drayton DL, Andrian von UH. T cell- and B cell-independent adaptive immunity mediated by natural killer cells. Nature immunology. 2006 May;7(5):507–16.

4. Paust S, Gill HS, Wang B-Z, Flynn MP, Moseman EA, Senman B, et al. Critical role for the chemokine receptor CXCR6 in NK cell-mediated antigen-specific memory of haptens and viruses. Nature immunology. 2010 Dec;11(12):1127–35.

5. Guma M, Angulo A, Vilches C, Gomez-Lozano N, Malats N, Lopez-Botet M. Imprint of human cytomegalovirus infection on the NK cell receptor repertoire. Blood. 2004 ed. 2004 Dec 1;104(12):3664–71.

6. Lopez-Vergès S, Milush J, Schwartz B, Pando M, Jarjoura J, York V, et al. Expansion of a unique CD57$^+$NKG2Chi natural killer cell subset during acute human cytomegalovirus infection. Proceedings of the National Academy of Sciences of the United States of America. 2011;108(36):14725–32.

7. Marquardt N, Béziat V, Nyström S, Hengst J, Ivarsson MA, Kekäläinen E, et al. Cutting Edge: Identification and Characterization of Human Intrahepatic CD49a+ NK Cells. J Immunol. 2015 Feb 11;194(6):2467–71.

8. Schlums H, Cichocki F, Tesi B, Theorell J, Béziat V, Holmes TD, et al. Cytomegalovirus infection drives adaptive epigenetic diversification of NK cells with altered signaling and effector function. Immunity. 2015 Mar 17;42(3):443–56.

9. Lee J, Zhang T, Hwang I, Kim A, Nitschke L, Kim M, et al. Epigenetic modification and antibody-dependent expansion of memory-like NK cells in human cytomegalovirus-infected individuals. Immunity. 2015 Mar 17;42(3):431–42.

10. Béziat V, Liu LL, Malmberg J-A, Ivarsson MA, Sohlberg E, Björklund AT, et al. NK cell responses to cytomegalovirus infection lead to stable imprints in the human KIR repertoire and involve activating KIRs. Blood. 2013 Apr 4;121(14):2678–88.

11. Cerwenka A, Lanier LL. Natural killer cell memory in infection, inflammation and cancer. Nature reviews Immunology. 2016 Feb;16(2):112–23.

12. Kuijpers TW, Baars PA, Dantin C, van den Burg M, van Lier RAW, Roosnek E. Human NK cells can control CMV infection in the absence of T cells. Blood. 2008 Aug 1;112(3):914–5.

13. Chen Z, Yang Y, Liu LL, Lundqvist A. Strategies to Augment Natural Killer (NK) Cell Activity against Solid Tumors. Cancers (Basel). Multidisciplinary Digital Publishing Institute; 2019 Jul 23;11(7):1040.

14. Menares E, Gálvez-Cancino F, Cáceres-Morgado P, Ghorani E, López E, Díaz X, et al. Tissue-resident memory CD8+ T cells amplify anti-tumor immunity by triggering antigen spreading through dendritic cells. Nat Commun. Nature Publishing Group; 2019 Sep 27;10(1):4401–12.

15.   Marquardt N, Kekäläinen E, Chen P, Lourda M, Wilson JN, Scharenberg M, et al.
Unique transcriptional and protein-expression signature in human lung tissue-resident
NK cells. Nat Commun. Nature Publishing Group; 2019 Aug 26;10(1):3841–12.

16.   Cooper GE, Ostridge K, Khakoo SI, Wilkinson TMA, Staples KJ. Human CD49a+ Lung
Natural Killer Cell Cytotoxicity in Response to Influenza A Virus. Front Immunol.
Frontiers; 2018;9:1671.

17.   Gordon CL, Miron M, Thome JJC, Matsuoka N, Weiner J, Rak MA, et al. Tissue
reservoirs of antiviral T cell immunity in persistent human CMV infection. The Journal
of experimental medicine. 2017 Mar 6;214(3):651–67.

18.   Marquardt N, Kekäläinen E, Chen P, Kvedaraite E, Wilson JN, Ivarsson MA, et al.
Human Lung NK Cells are Predominantly Comprised of Highly Differentiated
Hypofunctional CD69(-)CD56(dim) Cells. J Allergy Clin Immunol. 2016 Sep
23;139(4):1321–4.

19.   Scharenberg M, Vangeti S, Kekäläinen E, Bergman P, Al-Ameri M, Johansson N, et
al. Influenza A Virus Infection Induces Hyperresponsiveness in Human Lung Tissue-
Resident and Peripheral Blood NK Cells. Front Immunol. Frontiers; 2019;10:1116.

20.   Chiesa Della M, Falco M, Podestà M, Locatelli F, Moretta L, Frassoni F, et al.
Phenotypic and functional heterogeneity of human NK cells developing after umbilical
cord blood transplantation: a role for human cytomegalovirus? Blood. 2012 Jan
12;119(2):399–410.

21.   Liu LL, Landskron J, Ask EH, Enqvist M, Sohlberg E, Traherne JA, et al. Critical Role
of CD2 Co-stimulation in Adaptive Natural Killer Cell Responses Revealed in NKG2C-
Deficient Humans. Cell Rep. 2016 May 3;15(5):1088–99.

22.   Emgård J, Kammoun H, García-Cassani B, Chesné J, Parigi SM, Jacob J-M, et al.
Oxysterol Sensing through the Receptor GPR183 Promotes the Lymphoid-Tissue-
Inducing Function of Innate Lymphoid Cells and Colonic Inflammation. Immunity. 2018
Jan 16;48(1):120–8.

23.   Adams NM, Lau CM, Fan X, Rapp M, Geary CD, Weizman O-E, et al. Transcription
Factor IRF8 Orchestrates the Adaptive Natural Killer Cell Response. Immunity. 2018
Jun 19;48(6):1172–6.

24.   Lau CM, Adams NM, Geary CD, Weizman O-E, Rapp M, Pritykin Y, et al. Epigenetic
control of innate and adaptive immune memory. Nature immunology. Nature
Publishing Group; 2018 Sep;19(9):963–72.

25.   Cichocki F, Wu C-Y, Zhang B, Felices M, Tesi B, Tuininga K, et al. ARID5B regulates
metabolic programming in human adaptive NK cells. The Journal of experimental
medicine. Rockefeller University Press; 2018 Sep 3;215(9):2379–95.

26.   Foley B, Cooley S, Verneris MR, Pitt M, Curtsinger J, Luo X, et al. Cytomegalovirus
reactivation after allogeneic transplantation promotes a lasting increase in educated
NKG2C+ natural killer cells with potent function. Blood. 2012 Mar 15;119(11):2665–
74.

27.   Kim S-H, Han S-Y, Azam T, Yoon D-Y, Dinarello CA. Interleukin-32: a cytokine and
inducer of TNFalpha. Immunity. 2005 Jan;22(1):131–42.

28.   Ray SJ, Franki SN, Pierce RH, Dimitrova S, Koteliansky V, Sprague AG, et al. The
collagen binding alpha1beta1 integrin VLA-1 regulates CD8 T cell-mediated immune
protection against heterologous influenza infection. Immunity. 2004 Feb;20(2):167–79.

29.   Cheuk S, Schlums H, Gallais Sérézal I, Martini E, Chiang SC, Marquardt N, et al.
CD49a Expression Defines Tissue-Resident CD8(+) T Cells Poised for Cytotoxic
Function in Human Skin. Immunity. 2017 Feb 21;46(2):287–300.

30.   Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, et al.
Single-cell reconstruction of the early maternal-fetal interface in humans. Nature.
Nature Publishing Group; 2018 Nov;563(7731):347–53.

31.   Masopust D, Choo D, Vezys V, Wherry EJ, Duraiswamy J, Akondy R, et al. Dynamic T
cell migration program provides resident memory within intestinal epithelium. The
Journal of experimental medicine. 2010 Mar 15;207(3):553–64.

32.   Kumar BV, Ma W, Miron M, Granot T, Guyer RS, Carpenter DJ, et al. Human Tissue-
Resident Memory T Cells Are Defined by Core Transcriptional and Functional
Signatures in Lymphoid and Mucosal Sites. Cell Rep. 2017 Sep 19;20(12):2921–34.

33.   Sojka DK, Plougastel-Douglas B, Yang L, Pak-Wittel MA, Artyomov MN, Ivanova Y, et
al. Tissue-resident natural killer (NK) cells are cell lineages distinct from thymic and
conventional splenic NK cells. Elife. 2014;3:e01659.

34.   Melsen JE, Lugthart G, Vervat C, Kielbasa SM, van der Zeeuw SAJ, Buermans HPJ,
et al. Human Bone Marrow-Resident Natural Killer Cells Have a Unique
Transcriptional Profile and Resemble Resident Memory CD8+ T Cells. Front Immunol.
Frontiers; 2018;9:1829.

35.   Béziat V, Dalgard O, Asselah T, Halfon P, Bedossa P, Boudifa A, et al. CMV drives
clonal expansion of NKG2C+ NK cells expressing self-specific KIRs in chronic
hepatitis patients. Eur J Immunol. 2012;42(2):447–57.

36.   Hammer Q, Rückert T, Romagnani C. Natural killer cell specificity for viral infections.
Nature immunology. Nature Publishing Group; 2018 Aug;19(8):800–8.

37.   Nikzad R, Angelo LS, Aviles-Padilla K, Le DT, Singh VK, Bimler L, et al. Human
natural killer cells mediate adaptive immunity to viral antigens. Sci Immunol. Science
Immunology; 2019 May 10;4(35):eaat8116.

38.   Romee R, Schneider SE, Leong JW, Chase JM, Keppel CR, Sullivan RP, et al.
Cytokine activation induces human memory-like NK cells. Blood. 2012 Dec
6;120(24):4751–60.

39.   Romee R, Rosario M, Berrien-Elliott MM, Wagner JA, Jewell BA, Schappe T, et al.
Cytokine-induced memory-like natural killer cells exhibit enhanced responses against
myeloid leukemia. Sci Transl Med. American Association for the Advancement of
Science; 2016 Sep 21;8(357):357ra123–3.

40.   Ni J, Hölsken O, Miller M, Hammer Q, Luetke-Eversloh M, Romagnani C, et al.
Adoptively transferred natural killer cells maintain long-term antitumor activity by
epigenetic imprinting and CD4+ T cell help. Oncoimmunology. Taylor & Francis;
2016;5(9):e1219009.

41.   Hammer Q, Rückert T, Dunst J, Romagnani C. Adaptive Natural Killer Cells Integrate
Interleukin-18 during Target-Cell Encounter. Front Immunol. Frontiers; 2017;8:1976.

42.   Jennrich S, Lee MH, Lynn RC, Dewberry K, Debes GF. Tissue exit: a novel control
point in the accumulation of antigen-specific CD8 T cells in the influenza a virus-
infected lung. Journal of virology. 2012 Apr;86(7):3436–45.

43.   Wu T, Hu Y, Lee Y-T, Bouchard KR, Benechet A, Khanna K, et al. Lung-resident
memory CD8 T cells (TRM) are indispensable for optimal cross-protection against
pulmonary virus infection. J Leukoc Biol. Wiley-Blackwell; 2014 Feb;95(2):215–24.

44.   Smith CJ, Caldeira-Dantas S, Turula H, Snyder CM. Murine CMV Infection Induces
865   the Continuous Production of Mucosal Resident T Cells. Cell Rep. 2015 Nov
866   10;13(6):1137–48.

867   45.   Morabito KM, Ruckwardt TJ, Bar-Haim E, Nair D, Moin SM, Redwood AJ, et al.
868   Memory Inflation Drives Tissue-Resident Memory CD8+ T Cell Maintenance in the
869   Lung After Intranasal Vaccination With Murine Cytomegalovirus. Front Immunol.
870   Frontiers; 2018;9:1861.

871   46.   Takamura S, Kato S, Motozono C, Shimaoka T, Ueha S, Matsuo K, et al. Interstitial-
872   resident memory CD8+ T cells sustain frontline epithelial memory in the lung. The
873   Journal of experimental medicine. Rockefeller University Press; 2019 Sep
874   26;159:jem.20190557.

875   47.   Orr M, Murphy W, Lanier L. "Unlicensed" natural killer cells dominate the response to
876   cytomegalovirus infection. Nature immunology. 2010;11(4):321–7.

877   48.   Smith HRC, Heusel JW, Mehta IK, Kim S, Dorner BG, Naidenko OV, et al. Recognition
878   of a virus-encoded ligand by a natural killer cell activation receptor. Proceedings of the
879   National Academy of Sciences of the United States of America. 2002 Jun
880   25;99(13):8826–31.

881   49.   Goodier MR, Rodríguez-Galán A, Lusa C, Nielsen CM, Darboe A, Moldoveanu AL, et
882   al. Influenza Vaccination Generates Cytokine-Induced Memory-like NK Cells: Impact
883   of Human Cytomegalovirus Infection. J Immunol. 2016 May 27.

884   50.   Li T, Wang J, Wang Y, Chen Y, Wei H, Sun R, et al. Respiratory Influenza Virus
885   Infection Induces Memory-like Liver NK Cells in Mice. J Immunol. American
886   Association of Immunologists; 2017 Jan 23;198(3):1242–52.

887   51.   Dou Y, Fu B, Sun R, Li W, Hu W, Tian Z, et al. Influenza vaccine induces intracellular
888   immune memory of human NK cells. PLoS ONE. 2015;10(3):e0121258.

889   52.   Gillard GO, Bivas-Benita M, Hovav A-H, Grandpre LE, Panas MW, Seaman MS, et al.
890   Thy1+ NK [corrected] cells from vaccinia virus-primed mice confer protection against
891   vaccinia virus challenge in the absence of adaptive lymphocytes. PLoS Pathog. 2011
892   Aug;7(8):e1002141.

893   53.   Reeves RK, Li H, Jost S, Blass E, Li H, Schafer JL, et al. Antigen-specific NK cell
894   memory in rhesus macaques. Nature immunology. 2015 Sep;16(9):927–32.

895   54.   Tang H, Li C, Wang L, Zhang H, Fan Z. Granzyme H of cytotoxic lymphocytes is
896   required for clearance of the hepatitis B virus through cleavage of the hepatitis B virus
897   X protein. J Immunol. American Association of Immunologists; 2012 Jan
898   15;188(2):824–31.

899   55.   Andrade F, Fellows E, Jenne DE, Rosen A, Young CSH. Granzyme H destroys the
900   function of critical adenoviral proteins required for viral DNA replication and granzyme
901   B inhibition. EMBO J. 2007 Apr 18;26(8):2148–57.

902   56.   Dorner BG, Smith HRC, French AR, Kim S, Poursine-Laurent J, Beckman DL, et al.
903   Coordinate expression of cytokines and chemokines by NK cells during murine
904   cytomegalovirus infection. J Immunol. 2004 Mar 1;172(5):3119–31.

905   57.   Swanson BJ, Murakami M, Mitchell TC, Kappler J, Marrack P. RANTES production by
906   memory phenotype T cells is controlled by a posttranscriptional, TCR-dependent
907   process. Immunity. 2002 Nov;17(5):605–15.

58. Nizard M, Roussel H, Diniz MO, Karaki S, Tran T, Voron T, et al. Induction of resident
memory T cells enhances the efficacy of cancer vaccine. Nat Commun. Nature
Publishing Group; 2017 May 24;8:15221.

59. Picelli S, Faridani OR, Björklund ÅK, Winberg G, Sagasser S, Sandberg R. Full-length
RNA-seq from single cells using Smart-seq2. Nat Protoc. Nature Publishing Group;
2014 Jan;9(1):171–81.

60. Picelli S, Björklund ÅK, Reinius B, Sagasser S, Winberg G, Sandberg R. Tn5
transposase and tagmentation procedures for massively scaled sequencing projects.
Genome Res. 2014 Dec;24(12):2033–40.

61. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnetjournal. 2011 Aug 2;17(1):10.

62. Modolo L, Lerat E. UrQt: an efficient software for the Unsupervised Quality trimming of
NGS data. BMC Bioinformatics. 2015 Apr 29;16(1):175.

63. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
aware quantification of transcript expression. Nature Methods. 2017 Mar 6;14(4):417–
9.

64. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. BioMed Central; 2014;15(12):550.

65. Weigt SS, Wang X, Palchevskiy V, Gregson AL, Patel N, DerHovanessian A, et al.
Gene Expression Profiling of Bronchoalveolar Lavage Cells Preceding a Clinical
Diagnosis of Chronic Lung Allograft Dysfunction. PLoS ONE. Public Library of
Science; 2017;12(1):e0169894.