Determining insulin sensitivity from glucose tolerance tests in Iberian and

Landrace pigs

J. M. Rodríguez-López[1,a], M. Lachica[1], L. González-Valero[1] and I. Fernández-Fígares[1]

[1]*Department of Physiology and Biochemistry of Animal Nutrition, Estación*

Experimental del Zaidín, CSIC, Profesor Albareda 1, 18008 Granada, Spain

[a]*Present address: Unité Transformations & Agro-Ressources - EA 7519, Institut*

Polytechnique UniLaSalle, 19 rue Pierre Waguet - BP 30313 - F-60026 Beauvais

Cedex, France

Corresponding author: Ignacio Fernández-Fígares. Email: ifigares@eez.csic.es

Short title: Insulin sensitivity in Iberian and Landrace pigs

Abstract

As insulin sensitivity may help to explain divergences in growth and body composition

between native and modern breeds, metabolic responses to glucose infusion were

measured using an intra-arterial glucose challenge (IAGTT). Iberian (n = 4) and

Landrace (n = 5) barrows (47.0 ± 1.2 kg BW), fitted with a permanent carotid artery

catheter were injected with glucose (500 mg/kg BW) and blood samples collected at -

10, 2, 5, 10, 15, 20, 25, 30, 45, 60, 90, 120 and 180 min following glucose infusion.

Plasma samples were analysed for insulin, glucose, lactate, triglycerides, cholesterol,

creatinine, albumin and urea. Insulin sensitivity indices were calculated and analysed.

Mean plasma glucose, creatinine and cholesterol concentrations were lower (0.01 < *P*

< 0.001) in Iberian (14, 68 and 22%, respectively) compared with Landrace pigs during

the IAGTT. However, mean plasma insulin, lactate, triglycerides and urea

concentrations were greater (0.01 $<P<$ 0.001) in Iberian (50, 35, 18 and 23%,

respectively) than in Landrace pigs. Iberian pigs had larger area under the curve (AUC)

of insulin ($P < 0.05$) and lactate ($0.1 < P < 0.05$), and smaller ($P < 0.05$) AUC for glucose

0-60 min compared with Landrace pigs. QUICKI index decreased while HOMA-%B

index increased indicating improved β-cell function in Iberian compared with Landrace

pigs, but no difference ($P > 0.10$) in calculated insulin sensitivity index was found

between breeds. A time response ($P < 0.05$) was obtained for insulin, glucose and

lactate so that maximum concentration was achieved 10 and 15 min post-infusion for

insulin (Iberian and Landrace, respectively), immediately post-infusion for glucose, and

20 min post-infusion for lactate, decreasing thereafter until basal levels. There was no

time effect for the rest of metabolites evaluated. In conclusion, growing Iberian pigs

challenged with an IAGTT showed changes in biochemical parameters and insulin

response that may indicate an early stage of insulin resistance.

Keywords: insulin resistance, metabolism, obesity, *Sus scrofa domestica, Sus scrofa*

mediterraneus

Implications

Intense selective breeding and selection for increased growth rate and leanness has

favourable consequences for the health of consumers and, from an economic point of

view, for the livestock industry. Nevertheless, the metabolic repercussion of this

selection may involve differences on insulin -a central hormone regulator of nutrient

uptake and metabolism- sensitivity between modern and local breeds. The Iberian pig is an autochthonous breed showing insulin resistance during the growing phase.

Introduction

The Iberian pig is a slow growing native breed of the Mediterranean basin with much greater whole body fat content than lean-type pigs (Nieto *et al.*, 2002). Compared with conventional breeds, Iberian pigs show a lower efficiency of energy utilisation for protein deposition in the growing period (Barea *et al.*, 2007). The greater relative viscera weight (Rivera-Ferre *et al.*, 2005) and total heat production (González-Valero *et al.*, 2016) associated in part with the greater rate of muscle protein turnover (Rivera-Ferre *et al.*, 2005) in Iberian compared with lean-type pigs help to explain the low energy efficiency for growth. In fact, Rivera-Ferre *et al.* (2005) showed that muscle protein degradation was increased in Iberian pigs resulting in decreased muscle protein accretion compared with Landrace pigs. Interestingly, insulin resistance at the muscle level could explain an increased protein degradation (Wang *et al.*, 2006) affecting overall protein accretion. In a previous study using balanced or lysine deficient diets at two CP levels, Iberian had greater fasting serum insulin concentration than Landrace pigs (Fernández-Fígares *et al.*, 2007), suggesting the possibility of insulin resistance in Iberian pigs. We hypothesised that compared with modern lean breeds Iberian pigs have decreased insulin sensitivity, which could explain differences on growth, body composition and metabolic characteristics compared with modern breeds. The objective of the present study was to evaluate differences on insulin sensitivity between Iberian and Landrace pigs using an intra-arterial glucose tolerance test (IAGTT).

Material and methods

Animals

All procedures used in this study were approved by the Bioethical Committee of the Spanish National Research Council (CSIC, Spain), and the animals cared for in accordance with the Royal Decree No. 1201/2005 (Spain). The experiment was performed with five Landrace and four Iberian (*Silvela* strain) barrows supplied by Granja El Arenal (Córdoba, Spain) and Sánchez Romero Carvajal (Jabugo S.A., Puerto de Santa María, Cádiz, Spain), respectively. The pigs were allowed *ad libitum* access to a standard diet (160 g CP/kg; 14 MJ metabolizable energy/kg DM) with free access to water in a controlled-environment room (21 ± 1.5°C). After acclimatization, each animal was surgically fitted with a chronic catheter in carotid artery following the procedure described previously (Rodríguez-López et al., 2013). After surgery, the pigs were fed at 2.4 × metabolizable energy for maintenance (444 kJ/kg$^{0.75}$ BW; National Research Council (NRC) 1998). On the day of the experiment, pigs (47.0 ± 1.2 kg BW) were given an intra-arterial bolus (500 mg/kg BW) of glucose (50% sterile dextrose; glucosado 50% Braun, B. Braun Medical S.A., Rubi, Barcelona, Spain) over a 1 min period after an overnight fast. The catheter was immediately flushed with 5 ml of sterile saline. Blood samples (5 ml) were collected at -10, 0, 5, 10, 15, 20, 25, 30, 45, 60, 90, 120 and 180 min following glucose infusion.

Biochemical analysis and calculations

Plasma was obtained by centrifugation (4ºC, 1820 x g for 30 min; Eppendorf 5810 R, Hamburg, Germany) and stored in aliquots at -20ºC until insulin and metabolites (glucose, lactate, triglycerides, cholesterol, creatinine, albumin and urea) were

analysed. All samples were assayed in duplicate except for insulin which was assayed

in triplicate.

Insulin was measured using commercially-available radioimmuno assay (RIA) kit

following the directions of the manufacturer (Millipore porcine insulin RIA kit; Cat. PI-

12K). Radioactivity in samples was measured using a gamma counter (Behring 1612;

Nuclear Enterprises Ltd, Edinburgh, Scotland). Human insulin was used as standard,

and the assay was validated for use in porcine plasma samples (Fernández-Fígares

et al., 2007). The intra- and inter-assay CVs for plasma insulin were 4.4 and 9.1%,

respectively.

Plasma glucose, lactate, triglycerides, cholesterol, creatinine, albumin and urea

were measured colorimetrically using an automated Cobas Integra 400® analyser

(Roche Diagnostics GmbH, Mannheim, Germany).

Responses of plasma insulin, glucose and lactate were evaluated separately by

computing total area under the response curve (AUC) determined using trapezoidal

geometry for the time period indicated following intra-arterial glucose infusion (e.g.

AUC5 stands for the integrated area between 0-5 min post-infusion, AUC10 between

0-10 min post-infusion, etc.).

The rates of decline in plasma insulin and glucose concentrations were calculated

based on the slope in the linear portion of the response curve (0 to 30 min post-

challenge) and expressed as a fractional rate constant determined from the slope of

the natural logarithm of plasma concentrations *vs.* time (Shipley and Clark, 1972). The

fractional turnover rates (*k*), or disappearance rates, of plasma insulin and glucose

(%/min) were calculated using the relationship (Kaneko *et al.*, 2008):

$k = (Ln1 - Ln2)/(T2 - T1)$

where Ln1 and Ln2 are the natural logarithms of plasma insulin (or glucose)

concentrations at times T1 (0 min) and T2 (30 min), respectively.

From the k value, the half-life, $T_{\frac{1}{2}}$ (min), may be calculated as:

$T_{\frac{1}{2}} = 100 \times 0.693/k$

For insulin sensitivity, indices used in human medicine were utilized.

The so-called homeostasis model assessment (HOMA; Matthews *et al.*, 1985) was

calculated for estimating insulin resistance (HOMA-IR) and β-cell function (HOMA-%B)

at fasting conditions, as follows:

HOMA-IR: fasting plasma insulin $\times$ fasting plasma glucose/22.5

HOMA-%B: (20 $\times$ fasting plasma insulin)/(fasting plasma glucose - 3.5)

It is assumed that non-insulin-resistant individuals have 100% β-cell function and an

insulin resistance of 1.

The quantitative insulin sensitivity check index (QUICKI; Katz *et al.*, 2000) was

computed as:

$1/[Ln(I_0) + Ln(G_0)]$

where I_0 is the fasting insulin (μU/ml), and G_0 is the fasting glucose (mg/dl).

Finally, the insulin sensitivity index (CSI; Tura *et al.*, 2010) was calculated as:

$CSI = K_G/(\Delta AUC_{INS}/T)$

where K_G is the slope of Ln glucose in the linear portion of the response curve, ΔAUC_{INS}

is the AUC of insulin above basal value, and T is the time interval (between 0 and 30

min) when K_G and ΔAUC_{INS} are calculated.

Statistical analyses

Plasma metabolites were evaluated using a mixed ANOVA with repeated measures

(Version 9.4; PROC MIXED, SAS Institute Inc., Cary, NC, USA) with the fixed effects

of breed, time of sampling and their interaction in the model statement. The

experimental unit was the pig. The covariance-structure used was the

autoregressive(1). To compare differences among the means, the pdiff (piecewise

differentiable) option was used. Assumptions that are required for a ANOVA were

tested following the protocol from Zuur et al. (2010).Homogenity of variance was

assured by applying the Levene's-Test. No transformation was required.

Fractional turnover rates, indices of insulin resistance and AUC corrected for baseline

(0 to 180 min or fractions) were analysed using the GLM procedure of SAS with the

pig as the experimental unit. Least square means and pooled SEM are presented.

Differences were considered significant at $P < 0.05$ and trends approaching

significance were considered for $0.05 < P < 0.10$.

Results

Fasting plasma insulin was greater in Iberian compared with Landrace pigs (15.6 *and*

8.10 µU/ml, respectively; sem=1.55, $P < 0.05$) whereas fasting plasma glucose was

similar for both breeds (4.68 and 5.85 mmol/l for Iberian and Landrace, respectively;

sem=0.92, $P > 0.10$).

Average plasma metabolites and insulin concentrations after the IAGTT are shown

in Table 1. Mean plasma glucose, cholesterol and creatinine concentrations were lower

in Iberian (14, 22 and 68%, respectively; $0.01 < P < 0.001$) compared with Landrace

pigs. However, mean plasma insulin, lactate, triglycerides and urea concentrations

were greater in Iberian (50, 35, 18 and 23%, respectively; $0.01 < P < 0.001$) than in

Landrace pigs. No differences ($P > 0.1$) were found between breeds for albumin levels.

Indices of insulin sensitivity are shown in Table 2. QUICKI index decreased ($P <$

0.05) while HOMA-%B index increased ($P < 0.01$) in Iberian compared with Landrace

pigs. No differences ($P > 0.10$) were found for HOMA-IR and CSI.

Only plasma insulin (Figure 1), glucose (Figure 2) and lactate (Figure 3)

concentrations changed throughout time ($P < 0.001$; Table1) after the IAGTT.

An interaction between breed and time was found for plasma insulin, such that

concentration of insulin was greater in Iberian pigs from -10 to 15 min and from 90-180

min and lower between 25-45 min (Figure 1). In both breeds, plasma insulin levels

increased 7-fold, reaching a peak concentration 10 and 15 min after glucose infusion

for Iberian (113.6 ± 7.1 μU/ml) and Landrace (55.7 ± 6.4 μU/ml) pigs, respectively.

Insulin remained well above fasting levels until 20 and 45 min after glucose infusion

for Iberian and Landrace pigs, respectively; thereafter insulin levels rapidly decreased

until fasting levels were attained. Fractional turnover rate tended to increase in Iberian

compared with Landrace pigs (6.2 and 4.7 ± 0.46%/min, respectively; $0.05 < P < 0.1$)

while half-life tended to decrease (11.3 and 15.2 ± 1.30 min for Iberian and Landrace

pigs, respectively; $0.05 < P < 0.1$).

Glucose peaked (Figure 2) immediately after glucose infusion (288% on average (n

= 9) compared with mean basal levels) and thereafter gradually decreased to values

below fasting levels after 20 min for Iberian and 25 min for Landrace pigs. After glucose

nadir (3.3 ± 0.6 mM), glucose concentration gradually increased again to reach values

comparable to fasting levels at 180 min. No differences were found between breeds

for glucose fractional turnover rate (4.1 and 3.3 ± 0.37%/min for Iberian and Landrace

pigs, respectively; $P > 0.10$) or glucose half-life (17.1 and 22.1 ± 2.67 min for Iberian

and Landrace pigs, respectively; $P > 0.10$).

Lactate increased after the IAGTT, peaked at 20 min for both breeds and declined

progressively until reaching basal concentrations at 180 min (Figure 3).

The AUC values for each sampling time of insulin, glucose and lactate are shown in

Figure 4, 5 and 6, respectively. Insulin AUC was greater ($P < 0.05$) for Iberian

compared with Landrace pigs at all times. Conversely, glucose AUC between 0-15, 0-

20, 0-25, 0-30, 0-45 and 0-60 min were lower ($P < 0.05$) for Iberian than Landrace pigs.

Glucose AUC tended to be greater ($0.05 < P < 0.10$) between 0-90 and 0-120 min. No

differences ($P < 0.05$) in glucose AUC was found between 0-5, 0-150 and 0-180 min.

Plasma lactate AUC was greater ($P < 0.05$) for Iberian for 0-10 and 0-15 min and

tended to be greater ($0.05 < P < 0.10$) for 0-5, 0-20, 0-25, 0-90, 0-120, 0-150 and 0-

180 min after glucose infusion.

Discussion

The IAGTT method allowed the comparison of the insulin responsiveness of an obese

(Iberian) and a lean (Landrace) pig breed. It has previously been shown that Iberian

pigs are fatty pigs with slow growth rate compared to modern breeds (Nieto et al.,

2012). As arterial blood represents the metabolites concentration to which the tissues

are exposed (Brouns *et al.*, 2005), chronic catheters were inserted in carotid artery for

glucose infusion and blood sampling.

In the current study we have shown increased fasting plasma insulin and insulin

response (higher plasma concentrations and AUC) after glucose infusion in Iberian

compared with Landrace pigs. Greater postprandial serum levels of insulin have been

described in Iberian (Fernández-Fígares *et al.*, 2007) and Ossabaw (Wangsness *et al.*,

1981) pigs (both obese pigs) compared with lean pigs after ingestion of feed or glucose

infusion. However, other comparative studies found increased insulin secretion in

Large White boars than in Meishan (Weiler *et al.*, 1998) boars (obese breed). The

limited growth and development of slow growing pigs could result at least partly from

lower concentrations of plasma insulin, because most cells of the body require insulin

for adequate uptake of glucose and amino acids (Claus and Weiler, 1994).

Nevertheless, decreased insulin concentration in blood does not seem to explain the

poor growth performance of Iberian compared to Landrace pigs. If the concentration

of insulin is compared among animals of different breeds, the sensitivity of each breed

to insulin should be considered.

After glucose ingestion, blood concentration rapidly returns to preprandial values as

shown in the present experiment, which indicates that exogenous glucose was

efficiently metabolized, stored as glycogen, or both. As expected, when glucose was

infused, plasma glucose levels were rapidly increased and a subsequent insulin

response was observed. The elevated insulin lowered plasma glucose below fasting

values within 20 min and insulin levels returned to baseline as plasma glucose

declined. In our study, glucose concentration and glucose AUC during the glucose

challenge were lower in Iberian compared with Landrace pigs, with no differences in

fasting plasma glucose, may be due to the limited number of pigs. When interpreting

the individual glucose curves, a monophasic pattern was identified for both breeds.

The lower glucose AUC of Iberian pigs (-19% on average) may be related to the greater

insulin AUC (+33% on average), a common pattern in many models of obesity.

However, the reasons for the unequal physiological response between breeds are not

well understood and must be discussed.

As it has been proved that the energy needs of portal-drained viscera are fulfilled

by the oxidation of glucose, glutamate, and glutamine in pigs (Stoll *et al.*, 1999), a

larger gastrointestinal tract of Iberian pigs compared to Landrace (Rivera-Ferre *et al.*,

2005) is in line with the decreased AUC of glucose in the present experiment.

However, despite the larger size of the gastrointestinal tract and lower portal blood

flow (González-Valero *et al.*, 2016) of Iberian compared with Landrace pigs, no

differences on net portal flux of glucose after ingestion of the same diet were found

(Rodríguez-López *et al.*, 2013). Differences on insulin stimulated glucose transport at

portal-drained viscera level may help to explain these results. Iberian have lower

glucose concentrations than Landrace pigs after an intravenous adrenaline challenge

(Fernández-Fígares *et al.*, 2016), suggesting a decreased response of Iberian pigs to

sympathetic nervous system stimuli which is in line with the lower glucose AUC

reported here.

Lactate appearance after an intravenous glucose test is positively associated to

insulin sensitivity in humans (Lovejoy *et al.*, 1992), as it is related to lactate production

by insulin sensitive tissues (mainly muscle and fat). Because only limited amounts of

lactate are produced by muscle after glucose loading (Ykijarvinen *et al.*, 1990), the

source of lactate appearance should predominantly be adipose tissue (Lovejoy *et al.*,

1992), with a large capacity to convert glucose to lactate (Marin *et al.*, 1987). We report

a delay of 20 min in plasma lactate elevation relative to glucose peak following IAGTT,

which may reflect the time lag in adipose tissue uptake of glucose and subsequent

lactate production under the stimulation of insulin. Compared with Landrace, the

increased lactate AUC in Iberian pigs after the IAGTT could therefore be a

consequence of a larger adipose tissue (Nieto *et al.*, 2002) instead of greater insulin

sensitivity. On the other hand, insulin resistance was associated with elevated basal

lactate levels in obese humans (Lovejoy *et al.*, 1990), so increased basal lactate

concentrations in Iberian pigs (1.040 vs. 0.730 mmol/l sem=0.063 could also indicate

insulin resistance or reduced insulin sensitivity. Although inhibition of insulin action on

glycogenolysis in fasting conditions may lead to increased glucose release from

glycogen and subsequent conversion of glucose to lactate, there is no direct evidence

of this. There is indirect evidence, though, that elevated lactate levels reflect a glucose

sparing effect (decreased glucose utilisation) in muscle (Pearce and Connett, 1980).

Obesity is frequently associated to different degrees of dyslipidemia manifested as

increased triglyceridemia and low HDL-cholesterol. In our experiment, we found lower

plasma cholesterol but greater plasma triglycerides concentration in Iberian compared

with Landrace pigs, although we did not separate LDL and HDL fractions. Reduced

total cholesterol concentration could be due to reduced hepatic insulin sensitivity as

insulin stimulates cholesterol synthesis (Nelson and Cox, 2017).

Previous studies in our lab have shown the low genetic potential of growing Iberian

pigs for muscle protein deposition in comparison to lean breeds (Nieto *et al.*, 2002),

possibly due to the greater muscle protein degradation and turnover of the former

(Rivera-Ferre *et al.*, 2005). In line with this, plasma urea level (an indirect protein

degradation indicator) was 23% greater in Iberian compared with Landrace pigs in the

present study. Differences on circulating insulin or the capacity of insulin release

between breeds may explain differences in lean tissue deposition, as insulin has an

important role in skeletal muscle metabolism. In obese db/db mice (a model of insulin

deficiency) higher muscle protein degradation in comparison with control mice (normal

plasma insulin concentration) was reported; the authors concluded that insulin

resistance was associated with accelerated muscle protein degradation (Wang *et al.*,

2006). The elevated protein degradation reported in Iberian compared with Landrace

pigs (Rivera-Ferre *et al.*, 2005) suggests the possibility of insulin resistance at this

level. The lower plasma creatinine level (indicator of muscle mass) found in this study

for Iberian pigs, is in accordance with previous studies (Fernández-Fígares *et al.*,

2007) and also with the low muscle protein deposition and muscle size described

previously (Nieto *et al.*, 2002; Rivera-Ferre *et al.*, 2005). As insulin resistance is

associated with decreased muscle mass, plasma creatinine levels can also be used

as an indicator of insulin signalling disorders as reported by Kashima *et al.* (2017) in

humans. Further research regarding amino acid concentration after an IAGTT may

help to explain differences in the effect of insulin on muscle protein metabolism

between breeds.

When insulin sensitivity indices used in human medicine were applied to the

conditions of the present experiment, QUICKI and HOMA-%B were more sensitive

detecting differences between breeds. Indeed, QUICKI index decreased in Iberian

compared with Landrace pigs, pointing out an incipient insulin sensitivity impairment in

fasting Iberian pigs. Similarly, reduced QUICKI index was found in Bama miniature

pigs fed a high sucrose high fat diet compared with a control diet (0.5 *vs.* 0.6,

respectively; Liu *et al.*, 2017). The QUICKI index has been shown to provide

reasonable approximations of insulin efficiency in minipigs (Christoffersen *et al.*, 2009).

When we used the homeostasis model assessment (HOMA), differences on hepatic

insulin resistance (HOMA-IR index) were negligible between breeds. However, Iberian

had improved β-cell function compared with Landrace pigs according to HOMA-%B

index, which may be due to enhanced sensitivity of the β-cells to glucose during the

fasting period, consistent with QUICKI index. As a consequence, Iberian β-cell insulin

synthesis increases in accordance with the increased insulin release after the glucose

challenge and the elevated basal insulin concentrations reported for Iberian pigs.

Previous studies from our lab indicate that growing Iberian pigs are prone to insulin

resistance compared with modern breeds as denoted by increased hepatic

gluconeogenesis (González-Valero *et al.*, 2014), greater plasma free fatty acid

concentration (Fernández-Fígares *et al.*, 2016) and lower plasma creatinine and

QUICKI index (Fernández-Fígares *et al.*, 2007). Additionally, in this experiment we

show greater HOMA-%B and increased plasma insulin and lactate concentrations after

an IAGTT. The increased plasma insulin AUC after an IAGTT suggests insulin

resistance in comparison to values obtained for lean pigs, although the concentration

of glucose remained low which could indicate the absence of a peripheral insulin

resistance. Although Iberian pigs may be considered an obese breed in terms of body

composition (Nieto *et al.*, 2002; Barea *et al.*, 2007), at the development stage of this

experiment, insulin resistance has not yet been fully established. Nevertheless, insulin

resistance and impaired glucose tolerance has been shown in Iberian sows (2.5 years

old) *ad libitum* fed a saturated fat enriched diet for three months (Torres-Rovira *et al.*,

2012). The utilization of the hyperinsulinemic euglycemic clamp, the most definitive

approach to determine whole-body insulin action, may provide conclusive evidence

regarding the establishment of insulin resistance in growing Iberian pigs.

Acknowledgements

The authors thank the company Sánchez Romero Carvajal (Jabugo S.A., Puerto de

Santa María, Spain) for their helpful collaboration, Dr Luis Lara for statistical advice

and Dr Thomas J. Caperna for critically reading the manuscript. This research was

supported by grant AGL2006-5937 from the Ministry of Science and Education of

Spain.

Declaration of interest

There are no conflicts of interest.

Ethics statement

The procedures used in this study were approved by the Bioethical Committee of the

Spanish National Research Council (CSIC, Spain), and the animals cared for in

accordance with the Royal Decree No. 1201/2005 (Spain).

Software and data repository resources

Data and models are not deposited in an official repository.

References

Barea R, Nieto R and Aguilera JF 2007. Effects of the dietary protein content and the feeding

level on protein and energy metabolism in Iberian pigs growing from 50 to 100 kg body weight.

Animal: an International Journal of Animal Bioscience 1, 357-365.

Brouns F, Bjorck I, Frayn KN, Gibbs AL, Lang V, Slama G and Wolever TMS 2005. Glycaemic

index methodology. Nutrition research reviews 18, 145-171.

Claus R and Weiler U 1994. Endocrine regulation of growth and metabolism in the pig - a

review. Livestock Production Science 37, 245-260.

Christoffersen B, Ribel U, Raun K, Golozoubova V and Pacini G 2009. Evaluation of different

methods for assessment of insulin sensitivity in Gottingen minipigs: introduction of a new,

simpler method. American Journal of Physiology-Regulatory Integrative and Comparative

Physiology 297, R1195-R1201.

Fernández-Fígares I, Rodríguez-López JM, González-Valero L and Lachica M 2016. Metabolic

responses to adrenaline challenge in Iberian compared to Landrace pigs. In 9th International

Symposium on Mediterranean Pig, Portalegre, Portugal, p. 67.

Fernández-Fígares I, Lachica M, Nieto R, Rivera-Ferre MG and Aguilera JF 2007. Serum

profile of metabolites and hormones in obese (Iberian) and lean (Landrace) growing gilts fed

balanced or lysine deficient diets. Livestock Science 110, 73-81.

González-Valero L, Rodríguez-López JM, Lachica M and Fernández-Fígares I 2014. Metabolic

differences in hepatocytes of obese and lean pigs. Animal 8, 1873-1880.

González-Valero L, Rodríguez-López JM, Lachica M and Fernández-Fígares I 2016.

Contribution of portal-drained viscera to heat production in Iberian gilts fed a low protein diet:

comparison to Landrace. Journal of the Science of Food and Agriculture 96, 1202-1208.

Kaneko JJ, Harvey JW and Bruss, ML 2008. Clinical Biochemistry of Domestic Animals, 6th

Edition.

Kashima S, Inoue K, Matsumoto M and Akimoto K 2017. Low serum creatinine is a type 2

diabetes risk factor in men and women: The Yuport Health Checkup Center cohort study.

Diabetes & Metabolism 43, 460-464.

Katz A, Nambi S, Mather K, Baron A, Follmann D, Sullivan G and Quon M 2000. Quantitative

Insulin Sensitivity Check Index: A Simple, Accurate Method for Assessing Insulin Sensitivity In

Humans. The Journal of clinical endocrinology and metabolism 85, 2402-2410.

Liu YQ, Yuan JF, Xiang L, Zhao YQ, Niu MM, Dai X and Chen H 2017. A high sucrose and

high fat diet induced the development of insulin resistance in the skeletal muscle of Bama

miniature pigs through the Akt/GLUT4 pathway. Experimental Animals 66, 387-395.

Lovejoy J, Mellen B and Digirolamo M 1990. Lactate generation following glucose-ingestion -

relation to obesity, carbohydrate-tolerance and insulin sensitivity. International Journal of

Obesity 14, 843-855.

Lovejoy J, Newby FD, Gebhart SSP and Digirolamo M 1992. Insulin resistance in obesity is

associated with elevated basal lactate levels and diminished lactate appearance following

intravenous glucose and insulin. Metabolism-Clinical and Experimental 41, 22-27.

Marin P, Rebuffescrive M, Smith U and Bjorntorp P 1987. Glucose-uptake in human adipose-

tissue. Metabolism-Clinical and Experimental 36, 1154-1160.

Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF and Turner RC 1985.

Homeostasis model assessment - insulin resistance and beta-cell function from fasting

plasma-glucose and insulin concentrations in man. Diabetologia 28, 412-419.

Nelson DL and Cox MM 2017. Lehninger principles of biochemistry. W.H. Freeman and

Company ; Houndmills, Basingstoke : Macmillan Higher Education, New York, NY.

National Research Council (NRC) 1998. Nutrient requirements of swine,10th edition. National

Academy Press, Washington, DC.

Nieto R, Lara L, Barea R, Garcia-Valverde R, Aguinaga MA, Conde-Aguilera JA and Aguilera

JF 2012. Response analysis of the Iberian pig growing from birth to 150 kg body weight to

changes in protein and energy supply. Journal of Animal Science 90, 3809-3820.

Nieto R, Miranda A, Garcia MA and Aguilera JF 2002. The effect of dietary protein content and

feeding level on the rate of protein deposition and energy utilization in growing Iberian pigs

from 15 to 50 kg body weight. British Journal of Nutrition 88, 39-49.

Pearce FJ and Connett RJ 1980. Effect of lactate and palmitate on substrate utilization of

isolated rat soleus. American Journal of Physiology 238, C149-C159.

Rivera-Ferre MG, Aguilera JF and Nieto R 2005. Muscle fractional protein synthesis is higher

in Iberian than in landrace growing pigs fed adequate or lysine-deficient diets. Journal of

Nutrition 135, 469-478.

RodrÍguez-López JM, Lachica M, González-Valero L and Fernández-Fígares I 2013.

Approaches for quantifying gastrointestinal nutrient absorption and metabolism in a native and

a modern pig breed. Journal of Agricultural Science 151, 434-443.

Shipley RA and Clark RE 1972. Tracer methods for In vivo kinetics. Theory and applications.

Academic Press, New York.

Stoll B, Burrin DG, Henry J, Yu H, Jahoor F and Reeds PJ 1999. Substrate oxidation by the

portal drained viscera of fed piglets. American Journal of Physiology-Endocrinology and

Metabolism 277, E168-E175.

Torres-Rovira L, Astiz S, Caro A, Lopez-Bote C, Ovilo C, Pallares P, Perez-Solana ML,

Sanchez-Sanchez R and Gonzalez-Bulnes A 2012. Diet-Induced Swine Model with

Obesity/Leptin Resistance for the Study of Metabolic Syndrome and Type 2 Diabetes.

Scientific World Journal.

Tura A, Sbrignadello S, Succurro E, Groop L, Sesti G and Pacini G 2010. An empirical index

of insulin sensitivity from short IVGTT: validation against the minimal model and glucose clamp

indices in patients with different clinical characteristics. Diabetologia 53, 144-152.

Wang X, Hu Z, Hu J, Du J and Mitch W 2006. Insulin Resistance Accelerates Muscle Protein

Degradation: Activation of the Ubiquitin-Proteasome Pathway by Defects in Muscle Cell

Signaling. Endocrinology 147, 4160-4168.

Wangsness PJ, Acker WA, Burdette JH, Krabill LF and Vasilatos R 1981. Effect of fasting on

hormones and metabolites in plasma of fast-growing, lean and slow-growing obese pigs.

Journal of Animal Science 52, 69-74.

Weiler U, Claus R, Schnoebelen Combes S and Louveau I 1998. Influence of age and

genotype on endocrine parameters and growth performance: a comparative study in Wild

boars, Meishan and Large White boars. Livestock Production Science 54, 21-31.

Ykijarvinen H, Bogardus C and Foley JE 1990. Regulation of plasma lactate concentration in

resting human-subjects. Metabolism-Clinical and Experimental 39, 859-864.

Zuur AF, Ieno EN and Elphick CS 2010. A protocol for data exploration to avoid common

statistical problems. Methods in Ecology and Evolution 1, 3-14.

Table 1 *Average plasma metabolites and insulin concentrations in Iberian (n = 4) and Landrace (n = 5) pigs during an intra-arterial glucose challenge (IAGTT; 500 mg/kg BW, 0-180 min)[1]*

	Breed			P-value[2]		
	Iberian	Landrace	SEM	Breed	Time	Breed × Time
Insulin (µU/ml)	41	27	1.9	***	***	***
Glucose (mmol/l)	6.8	7.7	0.26	**	***	ns
Lactate (mmol/l)	1.3	1.0	0.039	***	***	ns
Triglycerides (mmol/l)	0.28	0.24	0.009	**	ns	ns
Cholesterol (mmol/l)	1.5	1.8	0.033	***	ns	ns
Creatinine (µmol/l)	54	90	1.2	**	ns	ns
Albumin (mmol/l)	0.48	0.50	0.009	ns	ns	ns
Urea (mmol/l)	3.0	2.4	0.102	***	ns	ns

[1]Average (n = 9) basal level of each metabolite: Glucose 5.33 mM, Insulin 11.43 µU/ml, Lactate 0.87 mM, Triglycerides 0.25 mM, Albumin 0.46 mM, Cholesterol 1.63 mM; Creatinine 76.2 µM, Urea 2.92 mM.

[2]ns = non-significant; **$P < 0.01$; ***$P < 0.001$.

Table 2 *Indices of glucose tolerance and insulin sensitivity in Iberian (n = 4) and Landrace (n = 5) pigs subjected to an intra-arterial glucose tolerance test[1]*

	Iberian	Landrace	SEM	*P*-value[2]
QUICKI	0.31	0.33	0.007	*
HOMA-IR	3.3	2.3	0.58	ns
HOMA-%B	267	100	25.6	**
CSI ($\times 10^{-4}$)	-12	-13	1.8	ns

[1]QUICKI: quantitative insulin sensitivity check index; HOMA-IR: homeostasis model assessment for estimating insulin resistance; HOMA-%B: homeostasis model assessment for estimating β-cell function; CSI: calculated insulin sensitivity index.

[2]ns = non-significant; *$P < 0.05$; **$P < 0.01$.

Figure 1 Plasma insulin concentrations during intra-arterial glucose challenge (500 mg/kg BW) in growing Iberian (n = 4) and Landrace (n = 5) pigs. Comparisons *versus* basal or control treatment: [†]$0.05 < P < 0.1$, *$P < 0.05$, ***$P < 0.001$. Average basal level of insulin 11.43 µU/ml.

469 **Figure 2** Plasma glucose concentration during intra-arterial glucose challenge (500

470 mg/kg BW) in growing Iberian (n = 4) and Landrace (n = 5) pigs. Comparisons *versus*

471 basal or control treatment: †0.05 < P < 0.1, $^*P$ < 0.05, $^{**}P$ < 0.01, *** P < 0.001; ns, not

472 significant (P > 0.10). Average basal level of glucose 5.33 mM.

483 **Figure 3** Plasma lactate concentration during intra-arterial glucose challenge (500

484 mg/kg BW) in growing Iberian (n = 4) and Landrace (n = 5) pigs. Comparisons *versus*

485 basal or control treatment: †0.05 < P < 0.1, *P < 0.05, ***P < 0.001; ns, not significant

486 (P > 0.10). Average basal level of lactate 0.87 mM.

498 **Figure 4** Area under the curve (AUC) of plasma insulin concentration during intra-
499 arterial glucose challenge (500 mg/kg BW) between minute 0 and the time points
500 indicated in growing Iberian (n = 4) and Landrace (n = 5) pigs. Differences between
501 breeds: * $P < 0.05$.

513   **Figure 5** Areas under the curve (AUC) of plasma glucose concentration during intra-

514   arterial glucose challenge (500 mg/kg BW) between minute 0 and the time points

515   indicated in growing Iberian (n = 4) and Landrace (n = 5) pigs. Differences between

516   breeds:[†]$0.05 < P < 0.1$, *$P < 0.05$.

527　**Figure 6** Area under the curve (AUC) of plasma lactate concentration during an intra-

528　arterial glucose challenge (500 mg/kg BW) between minute 0 and the time points

529　indicated in growing Iberian (n = 4) and Landrace (n = 5) pigs. Differences between

530　breeds:[†]$0.05 < P < 0.1$, [*]$P < 0.05$.