Modelling the emergence of whisker barrels

Sebastian S. James[a,1]**, Leah A. Krubitzer**[b]**, and Stuart P. Wilson**[a]

[a] Department of Psychology, The University of Sheffield, Sheffield, United Kingdom.; [b] Center for Neuroscience, The University of California, Davis, United States.

Brain development relies on an interplay between genetic specification and self-organisation. Striking examples of this relationship can be found in the somatosensory brainstem, thalamus, and cortex of rats and mice, where the arrangement of the facial whiskers is preserved in the arrangement of cell aggregates to form precise somatotopic maps. We show in simulation how realistic whisker maps can self-organize, by assuming that information is exchanged between adjacent cells only, under the guidance of gene expression gradients. The resulting model provides a simple account of how patterns of gene expression can constrain spontaneous pattern formation to faithfully reproduce functional maps in subsequent brain structures.

Barrel cortex | Self-organization | Somatotopic map | Axon guidance

Spatial patterns in neural connectivity provide clues about the constraints under which brains evolve and develop (1). Perhaps the most distinctive pattern can be found in the barrel cortex of many rodent species (2). The barrels are identifiable soon after birth in layer 4 of primary somatosensory cortex as dense clusters of thalamocortical axons, which are enclosed by borders a few neurons thick from postnatal day 3 (3). In the plane tangential to the cortical surface the barrels constitute a somatotopic map of the whiskers, with cells within adjacent barrels responding most strongly and quickly to deflection of adjacent whiskers (4). Barrel patterning reflects subcortical whisker maps comprising cell aggregates called barrelettes in the brainstem and barreloids in the thalamus.

Barrel formation requires afferent input from whisker stimulation and thalamic calcium waves (5), and depends on a complex network of axon guidance molecules such as ephrin-A5 and A7 and adhesion molecules such as cadherin-6 and 8 (6, 7). This network is orchestrated by interactions between morphogens Fgf8 and Fgf17 and transcription factors Emx2, Pax6, Sp8, and Coup-tf1 (8), which are expressed in gradients that mark orthogonal axes and can be manipulated to stretch, shrink, shift, and even duplicate barrels (9).

The barrel boundaries form a Voronoi tessellation across the cortical sheet (10) (Fig. 1A), suggesting that barreloid topology is preserved in the projection of thalamocortical axons into the cortex, and that a barrel forms by lateral axon branching from an initial centre-point that ceases upon contact with axons branching from adjacent centres. But the assumption of pre-arranged centres is difficult to resolve with the observation that axons arrive in the cortical plate as an undifferentiated bundle, prior to barreloid formation (11).

Alternatively, reaction-diffusion dynamics could generate a Voronoi tessellation without pre-arranged centres, by amplifying characteristic modes in a noisy initial distribution of axon branches, as a net effect of short-range cooperative and longer-range competitive interactions. Accordingly, the barrel pattern would be determined by the relative strength of these interactions and by the shape of the cortical field boundary. However, intrinsic cortical dynamics alone cannot account for the topographic correspondence between thalamic and cortical domains, the irregular sizes and specific arrangement of the barrels in rows and arcs, or the influence of gene expression gradients.

The centre-point and reaction-diffusion models are not mutually exclusive. Pre-organised centres could bias reaction-diffusion processes to generate specific arrangements more reliably, and mechanisms of lateral axon branching may constitute the tension between cooperation and competition required for self-organization. However, proof that barrel patterning can emerge from an undifferentiated bundle of axons, based only on local interactions, would show that a separate stage and/or extrinsic mechanism for pre-organising thalamocortical connections need not be assumed. To this end, we ask whether barrel maps can emerge in a system with reaction-diffusion dynamics, under the guidance of signalling gradients, and in the absence of pre-defined centres.

Models

Karbowski & Ermentrout (12) developed a reaction-diffusion style model of how extrinsic signalling gradients can constrain the emergence of distinct fields from intrinsic cortical dynamics. Their model defines how the density of connections $c(x, t)$ and axon branches $a(x, t)$ interact at time t, along a 1D anterior-posterior axis x, for N thalamocortical projections indexed by i. The model was derived from the assumption that the rates at which a_i and c_i grow are reciprocally coupled. Extending the original 1D model to simulate arealization on a 2D cortical sheet, we use $a_i(\mathbf{x}, t)$ and $c_i(\mathbf{x}, t)$, and model synaptogenesis as

$$\frac{\partial c_i}{\partial t} = -\alpha c_i + \beta \left(1 - \sum_{j=1}^{N} c_j \right) [a_i]^k. \quad [1]$$

Accordingly, where the total density of synaptic connections sums to one, connections decay at rate α. Otherwise the connection density increases non-linearly ($k > 1$) with the density of axon branching. Axon branching is modelled as

$$\frac{\partial a_i}{\partial t} = \nabla \cdot \left(D \nabla a_i - a_i \sum_{j=1}^{M} \gamma_{i,j} \nabla \rho_j(\mathbf{x}) \right) - \frac{\partial c_i}{\partial t} + \chi_i. \quad [2]$$

The first term on the right describes the divergence (indicated by $\nabla \cdot$) of the quantity in parentheses, which is referred to as the 'flux' of axonal branching. The flux represents diffusion across the cortical sheet, at rate D, and the influence of M molecular signalling fields, $\rho(\mathbf{x})$. The influence of a given field (indexed by j) on a given thalamic projection (indexed by i), is determined by $\gamma_{i,j}$, which may be positive or negative in order that axons may branch in the direction of either higher or

lower concentrations. The second term on the right represents an assumption that axon branching decreases when synaptogenesis increases, and vice versa, i.e., that axon branches are required for synaptic connections to form. Here $\chi_i = 0$ is a placeholder.

Software for modelling the cortical sheet as a hexagonal lattice with no-flux boundary conditions and an arbitrary boundary shape, for computing the flux using a finite difference method, for numerical integration by the fourth-order Runge-Kutta method, and for reproducing these results is available at https://github.com/ABRG-Models/BarrelEmerge.

Results

First we verified that all results established by (12) for a 1D axis could be reproduced using our extension to a 2D cortical sheet. Using an elliptical boundary with $M = 3$ offset guidance gradients aligned to the longer axis, $N = 5$ thalamocortical projections gave rise to five distinct cortical fields at locations that preserved the topographic ordering defined by the original γ values. However, we found that specifying N ordered areas required $M \approx (N + 1)/2$ signalling fields. As the number of distinct guidance fields is unlikely to approach the number of barrels, further extensions were required.

The term in parentheses in Eq. 1 represents competition between thalamocortical projections for a limited availability of cortical connections. To introduce competition also in terms of axon branching we redefined

$$\chi_i(\mathbf{x}, t) = -\frac{\epsilon a_i}{N - 1} \sum_{j \neq i}^{N} a_j, \qquad [3]$$

which reduces branching for each projection where the branches of other projections are dense. Divisive normalization keeps the axon branch density bounded on each iteration to the initial density at $t = 0$;

$$a_i(\mathbf{x}, t) = \int_\Omega a_i(\mathbf{x}, 0) d\sigma \, \frac{a_i'(\mathbf{x}, t)}{\int_\Omega a_i'(\mathbf{x}, t) d\sigma}, \qquad [4]$$

where prime symbols indicate the use of intermediate values computed from Eqs. 1–3 at time $t - \delta t$.

The only differences between thalamic projections are their strengths of interaction with the guidance fields, γ, hence any reliable differences between emergent cortical fields must be due to differences in these values only. We speculate that a projection interacts with a given ephrin field in the cortical subplate with a strength determined by the concentration of a similar molecule at its thalamic origin, i.e., the putative barreloid center. As such, two orthogonal linear thalamic gradients were defined, from which 41 pairs of γ values were sampled, at the coordinates of 41 barreloid centers recreated from (14) (Fig. 1B).

A cortical boundary enclosing 41 corresponding barrels was traced from a cytochrome oxidase stain from (13), and Eqs. 1–4 were solved for $N = 41$ projections on the resulting domain, using $M = 2$ linear signalling gradients aligned with the anterior-posterior and medial-lateral axes. From random initial conditions for $a(\mathbf{x}, 0)$ and $c(\mathbf{x}, 0)$, a clear Voronoi-like tesselation emerged (Fig. 1C; see Movie S1). A reduction in the Honda delta metric (see (10)) confirmed that a 'good' Voronoi pattern emerged within ≈ 3000 iterations (see

Fig. 1D). A measure of the difference in shape between real and simulated barrels revealed a strong correspondence (see Fig. 1D).

To further investigate the interplay of genetic and intrinsic mechanisms we simulated a seminal barrel duplication experiment (8), first by solving on a cortical domain comprising two separate, identically shaped cortical boundaries. In this case one distinct cortical field per thalamic projection emerged, within one of the two boundaries only, i.e., no duplication. However, merging the two fields to create an extended mirror-symmetric boundary shape, with an anterior-posterior guidance field reflected at the join, gave rise to two mirror-symmetrical barrel fields comprising 2N barrels, i.e., two cortical fields for each thalamic projection (Fig. 1E). Together these results suggest that by the misexpression of Fgf8, Shimogori & Grove (8) effectively created one large barrel field rather than two distinct cortical fields.

Movie S1 caption. Movie corresponding to Fig. 1C in the main paper. Simulation parameters were $N = 41$, $\alpha = 3$, $\beta = 20$, $k = 3$, $D = 0.2$, $\gamma \in \pm 2$, $\epsilon = 150$ and $\delta t = 0.0001$. Colours indicate the thalamic projection for which the connection density is maximal, black lines delineate boundaries, and overlaid contours show $c > 0.5$. The final frame in the movie is step 25000 of the simulation.

Discussion

The present results suggest that the key requirements for the emergence of realistic barrel patterning are i) at each cortical location thalamocortical projections compete for a limited number of available synaptic connections (Eqs. 1–2), ii) at each location the branching rate of a given projection is reduced by the density of other projections (Eq. 3), and iii) the branch density of each projection is conserved over time (Eq. 4).

The emergence of barrels in simulation required competition between thalamic projections in terms of synaptic connectivity, and also in terms of cortical space (as represented by χ). This requirement may account for the absence of barrels in rodents with larger brains, such as capybara, for which competition for space is presumably weaker (15). Hence, irrespective of whether barrels are necessary for adaptive whisker function, the emergence of somatotopically ordered modular structures may be an inevitable consequence of local competition for cortical territory driven by input from an array of discrete sensory organs (1).

It is important to emphasize that the formulation of the model is entirely local, insofar as simulation requires no information to be communicated from a given cortical grid cell to any but those immediately adjacent (via diffusion). Hence the simulations demonstrate how a self-organising system, constrained by genetically specified guidance cues and by the shape of the cortical field boundary, can faithfully reproduce an arrangement of cell aggregates in one neural structure as a topographic map in another.

Moreover, the present results confirm that somatotopic map formation does not require the pre-specification of centre-points by as yet undetermined additional developmental mechanisms.

ACKNOWLEDGMENTS. The authors thank Jason Berwick at the University of Sheffield for advice and for access to the rat

Fig. 1. **A** Left shows a cytochrome oxidase stain obtained from rat S1 by ([13]), with black lines to delineate barrels and to measure departure (Honda-δ; see ([10])) from a perfect Voronoi tesselation. Right shows the initial distribution of axon branching density (a) for one thalamocortical projection, and two molecular guidance fields (ρ). **B** The strengths of interaction γ with fields ρ_1 and ρ_2 are indicated for each of 41 projections by the lengths of green and blue arrows respectively, assuming that similar fields aligned to the posterior-anterior and medial-lateral axes in the ventroposterior medial nucleus of the thalamus are sampled at the locations of putative barreloid centers (reconstructed from ([14])). **C** Simulation results for parameters $N = 41$, $\alpha = 3$, $\beta = 20$, $k = 3$, $D = 0.2$, $\gamma \in \pm 2$, $\epsilon = 150$ and $\delta t = 0.0001$. Colours indicate the thalamic projection for which the connection density is maximal, black lines delineate boundaries, and overlaid contours show $c > 0.5$ (see Movie S1). **D** Red dots show the Honda-δ metric obtained from simulation approaching that obtained from barrels in **A** (dotted line); black squares measure the correspondence between the real and simulated barrel shapes, η (the product of the sum of squared differences between real and simulated centers and the sum of differences in area; units mm^4). **E** Guidance fields and emergent barrel pattern in a Fgf8 misexpression experiment (c.f. ([9])), simulated by reflecting ρ_1 at the join ($\epsilon = 80$). All scale bars 1mm.

barrel stains used to construct Fig. 1A. This work was supported by a Collaborative Activity Award, *Cortical Plasticity Within and Across Lifetimes*, from the James S. McDonnell Foundation (grant 220020516).

1. Purves D, Riddle D, LaMantia AS (1992) Iterated patterns of brain circuitry (or how the cortex gets its spots). *Trends in Neurosciences* 15(10):362–368.
2. Woolsey TA, Van der Loos H (1970) The structural organization of layer IV in the somatosensory region (S I) of mouse cerebral cortex: The description of a cortical field composed of discrete cytoarchitectonic units. *Brain Research* 17(2):205–242.
3. Erzurumlu RS, Gaspar P (2012) Development and critical period plasticity of the barrel cortex. *European Journal of Neuroscience* 35(10):1540–1553.
4. Armstrong-James M, Fox K, Das-Gupta A (1992) Flow of excitation within rat barrel cortex on striking a single vibrissa. *Journal of Neurophysiology* 68(4):1345–1358.
5. Antón-Bolaños N, et al. (2019) Prenatal activity from thalamic neurons governs the emergence of functional cortical maps in mice. *Science* 364(6444):987–990.
6. Vanderhaeghen P, et al. (2000) A mapping label required for normal scale of body representation in the cortex. *Nature Neuroscience* 3(4):358–365.
7. Miller K, Kolk SM, Donoghue MJ (2006) EphA7-ephrin-A5 signaling in mouse somatosensory cortex: Developmental restriction of molecular domains and postnatal maintenance of functional compartments. *The Journal of Comparative Neurology* 496(5):627–642.
8. Shimogori T (2005) Fibroblast Growth Factor 8 Regulates Neocortical Guidance of Area-Specific Thalamic Innervation. *Journal of Neuroscience* 25(28):6550–6560.
9. Assimacopoulos S, Kao T, Issa NP, Grove EA (2012) Fibroblast Growth Factor 8 Organizes the Neocortical Area Map and Regulates Sensory Map Topography. *Journal of Neuroscience* 32(21):7191–7201.
10. Senft SL, Woolsey TA (1991) Mouse Barrel Cortex Viewed as Dirichlet Domains. *Cerebral Cortex* 1(4):348–363.
11. Agmon A, Yang LT, O'Dowd DK, Jones EG (1993) Organized growth of thalamocortical axons from the deep tier of terminations into layer IV of developing mouse barrel cortex. *Journal of Neuroscience* 13(12):5365–5382.
12. Karbowski J, Ermentrout G (2004) Model of the Early Development of Thalamo-Cortical Connections and Area Patterning via Signaling Molecules. *Journal of Computational Neuroscience* 17(3):347–363.
13. Zheng Y, Johnston D, Berwick J, Mayhew J (2001) Signal Source Separation in the Analysis of Neural Activity in Brain. *NeuroImage* 13(3):447–458.
14. Haidarliu S, Ahissar E (2001) Size gradients of barreloids in the rat thalamus. *Journal of Comparative Neurology* 429(3):372–387.
15. Woolsey TA, Welker C, Schwartz RH (1975) Comparative anatomical studies of the sml face cortex with special reference to the occurrence of barrels in layer iv. *Journal of Comparative Neurology* 164(1):79–94.