# Molecular Basis for the Evolved Instability of a Human G-Protein Coupled Receptor

Laura M. Chamness,[1] Nathan B. Zelt,[1] Charles P. Kuntz,[1] Brian J. Bender,[3] Wesley D. Penn,[1] Joshua J. Ziarek,[2] Jens Meiler,[3,4] and Jonathan P. Schlebach[1]*

[1]*Department of Chemistry, Indiana University, Bloomington, Indiana, USA*
[2]*Department of Molecular and Cellular Biochemistry, Indiana University, Bloomington, Indiana, USA*
[3]*Department of Chemistry, Vanderbilt University, Nashville, TN, USA*
[4]*Institut for Drug Development, Leipzig University, Leipzig, SAC, Germany*

KEYWORDS: GnRHR, GPCR, Membrane Protein Folding, Proteostasis, Epistasis

*Corresponding author*. E-mail address: jschleba@indiana.edu

# ABSTRACT

Eukaryotic membrane proteins are prone to misfolding and degradation within the endoplasmic reticulum (ER). This is particularly true for the mammalian forms of the gonadotropin-releasing hormone receptor (GnRHR). Though they function at the plasma membrane, mammalian GnRHRs tend to accumulate within the secretory pathway. Their apparent instability is believed to have evolved in response to a selection for GnRHRs with attenuated activity. Nevertheless, the structural basis of this adaptation remains unclear. An experimental characterization of natural GnRHRs reveals this adaptation coincides with a C-terminal truncation and an increase in the polarity of its transmembrane (TM) domains. We show that this enhanced polarity compromises the translocon-mediated cotranslational folding of two nascent TM domains. Moreover, replacing a conserved polar residue in TM6 with an ancestral hydrophobic residue partially restores GnRHR expression with minimal impact on function. Our findings suggest the marginal energetics of cotranslational folding can be exploited to tune membrane protein fitness.

## Introduction

Proteins must continually sample new mutations that modify their conformational equilibria in order to maximize their evolutionary fitness (Harms & Thornton, 2013). Most mutations destabilize native protein structures, and evolutionary pathways are limited to those in which the protein retains sufficient conformational stability and activity with each successive mutation (Tokuriki & Tawfik, 2009). Like water-soluble proteins, the sequences of integral membrane proteins (MPs) are also constrained by folding energetics (Marinko et al., 2019). However, their transmembrane (TM) domains must also retain sufficient hydrophobicity to partition into the membrane, which further constrains their evolutionary sequence space (Marinko et al., 2019). In recent investigations of the sequence constraints within the class A G-protein coupled receptor (GPCR) rhodopsin, we found its expression to be highly sensitive to mutations within a marginally hydrophobic TM domain (Roushar et al., 2019). Moreover, the natural sequence of this TM domain appears to be more polar than is necessary to support function, and its expression can be enhanced by functionally-neutral hydrophobic substitutions (Roushar et al., 2019). As a result of this instability, much of the nascent protein is retained in the endoplasmic reticulum (ER) (Roushar et al., 2019). As is true for water-soluble proteins, these observations suggest that naturally evolved MPs tend to be metastable, and have not evolved to maximize the efficiency of protein biogenesis. Nevertheless, it is unclear whether this metastability provides an evolutionary benefit, or if it is simply an emergent property that stems from the evolutionary process itself (Taverna & Goldstein, 2002).

A series of previous investigations of the gonadotropin releasing hormone receptor (GnRHR), a class A GPCR, revealed that the mammalian forms of these receptors exhibit a heightened tendency to misfold and accumulate within the ER (Janovick, Ulloa-Aguirre, & Conn, 2003). GnRHR plays a critical role in steroidogenesis, and its functional expression at the plasma membrane is critical for reproductive fitness (Janovick et al., 2013, 2006). Variations within the *GnRHR* gene have been associated with shifts in litter size and length of the luteal phase (Bemji, Isa, Ibeagha-Awemu, & Wheto, 2018; P. Michael Conn, Knollman, Brothers, & Janovick, 2006; Ulloa-Aguirre, Janovick, Miranda, & Conn, 2006; Yang et al., 2011). Additionally, numerous loss of function mutations in human *GnRHR* have been found to cause hypogonadotropic hypogonadism (HH), which is characterized by infertility and loss of gonadal function (Janovick et al., 2013). GnRHRs found in fish, which produce many offspring, appear to exhibit robust plasma membrane expression (PME) relative to those found in mammals, which have far fewer offspring (P. Michael Conn, Knollman, et al., 2006; Janovick et al., 2006). Based on various observations, it has been suggested that the reproductive selection pressures increased the relative fitness of mammals expressing less stable GnRHR variants with diminished PME (P. Michael Conn, Janovick, Brothers, & Knollman, 2006; Ulloa-Aguirre et al., 2006). It has also been speculated that the pool of immature GnRHRs in the ER may provide a regulatory benefit, as modifications to the proteostasis network can alter the flux of mature protein through the secretory pathway (P. Michael Conn, Knollman, et al., 2006; P. Michael Conn & Ulloa-Aguirre, 2011; Janovick et al., 2006). Thus, it appears as though nature may have exploited the instability of GnRHRs in order to tune their evolutionary fitness. Nevertheless, the nature of the conformational modifications involved in these evolutionary adaptations remain poorly understood.

To evaluate the evolutionary sequence modifications that coincide with the proteostatic divergence between the mammalian and non-mammalian GnRHRs, we first compiled and analyzed the sequences of known type I GnRHRs. Consistent with previous observations, we first show that a C-terminal truncation within the mammalian receptors appears to significantly contribute to the attenuation of the mammalian receptor PME. However, multiple sequence alignments also reveal that the transmembrane (TM) helices of mammalian GnRHRs are considerably more polar than those of non-mammalian GnRHRs. We show that this enhanced polarity compromises the translocon-mediated membrane integration of two of its seven TM domains, which demonstrates that some of these sequence modifications compromise the fidelity of cotranslational folding. Structural models of these receptors suggest two of the polar substitutions that compromise translocon-mediated membrane integration of the nascent chain occur at surface residues that are projected into the membrane core. Moreover, we show that re-introducing the ancestral hydrophobic side chain at one of these positions partially restores the PME of human GnRHR with minimal impact on receptor activation. Together, these findings provide new evidence that evolution has exploited the marginal cotranslational folding energetics of GnRHR in order to tune its fitness. These observations suggest the instability of natural proteins provides an additional avenue for evolutionary adaptation.

**Figure 1. Cellular trafficking of GnRHR variants in HEK293T cells.** Human (red), mouse (black), and catfish (blue) GnRHRs were transiently expressed in HEK293T cells, and the relative abundance of plasma membrane GnRHR and intracellular GnRHR was analyzed by flow cytometry. Contour plots show the distribution of cellular fluorescence intensities associated with immunostaining of plasma membrane (y-coordinate) and intracellular (x-coordinate) GnRHRs.

# Results

## *Cellular Expression of Natural GnRHRs*

Various lines of evidence suggest the selection for reduced GnRH signaling in higher mammals produced GnRHRs with diminished conformational stability and attenuated plasma membrane expression (PME) (Janovick et al., 2003; Lin et al., 1998). Nevertheless, most previous studies relied on activity as a proxy for PME (P. M. Conn, McArdle, Andrews, & Huckle, 1987; Janovick et al., 2006; Leaños-Miranda, Ulloa-Aguirre, Ji, Janovick, & Conn, 2003). To more directly probe differences in GnRHR expression, we employed immunostaining in conjunction with flow cytometry to quantitatively characterize the expression of three previously characterized GnRHRs (human, mouse, and catfish). Briefly, each of these receptors was transiently expressed in HEK293T cells prior to labeling plasma membrane and intracellular GnRHRs with distinct fluorescent antibodies, as previously described (Schlebach et al., 2015). Cellular fluorescence profiles were then analyzed by flow cytometry. A comparison of the distribution of single-cell fluorescence profiles reveals that larger proportions of the expressed mouse GnRHR (*Mus musculus*, mGnRHR) and catfish GnRHRs (*Clarius gariepinus*, cGnRHR) accumulate at the plasma membrane relative to human

GnRHR (*Homo sapiens*, hGnRHR, Figure 1). The mean fluorescence intensity associated with the surface immunostaining of hGnRHR at the plasma membrane is 21.5 ± 6.0 fold lower than that of mGnRHR and 92.0 ± 17.9 fold lower than that of cGnRHR. Overall, the total cellular expression of hGnRHR was 2.71 ± 0.04 fold lower than that of mGnRHR and 2.04 ± 0.35 fold lower than cGnRHR. These results show for the first time that cGnRHR exhibits robust expression and trafficking relative to the mammalian receptors under equivalent conditions. Nevertheless, the nature of the structural modifications responsible for this apparent proteostatic divergence remains unclear.

## *Impact of the C-terminal Tail in GnRHR Expression*

Evolutionary adaptations in mammalian GnRHRs coincided with a variety of sequence modifications. Most strikingly, mammalian GnRHRs feature a C-terminal deletion of a disordered loop as well as a conserved amphipathic helix (helix 8, H8) that contains two palmitoylation sites (Lin et al., 1998; Millar, 2005; Palczewski et al., 2000). Fusing the C-terminal domain of cGnRHR to the C-terminus hGnRHR was previously shown to enhance the activity of the human receptor (Janovick et al., 2003). Nevertheless, the extent to which the structural elements within the C-terminal region impact the PME remains unclear. We therefore assessed the effects of various C-terminal modifications on the PME of cGnRHR. To determine whether C-terminal palmitoylation impacts PME, we first

**Figure 2. Impact of C-terminal modifications on the plasma membrane expression of cGnRHR.** Catfish GnRHRs bearing C-terminal modifications were transiently expressed in HEK293T cells prior to analysis of surface immunostaining of plasma membrane GnRHR by flow cytometry. A bar graph depicts the mean fluorescence intensity associated with the surface immunostaining of a series of catfish GnRHR variants normalized relative to that of human GnRHR. Values reflect the average of three biological replicates, and error bars reflect the standard deviation.

**Figure 3. Evolutionary divergence of the topological energetics of GnRHR.** The predicted transfer free energy associated with the translocon-mediated membrane integration of each TM domain within 59 known GnRHRs was calculated using the ΔG predictor, (ref) and the average value for the seven TM domains within each receptor was projected onto a phylogenetic tree. The names of each species within the phylogenetic tree are colored according to the average ΔG value for the seven TM domains within the corresponding receptor. The phylogenetic tree was constructed using ML after MUSCLE alignment using MEGA7 software, and the branch lengths represent number of substitutions per site. A length unit of 0.1 substitutions per site corresponds to 10% divergence. The *Clarius gariepinus* (catfish) receptor is annotated as a type II GnRHR, and was therefore excluded from this analysis (see Methods).

160 characterized a double mutant of cGnRHR that lacks its two C-terminal palmitoylation sites (C339A, C341A).
161 Removal of these palmitoylation sites has minimal effect on the PME of cGnRHR (Figure 2), which suggests the
162 loss of these modifications is not responsible for the attenuated PME of the mammalian receptors. To determine
163 whether the disordered portion of the C-terminal tail impacts PME, we next characterized a cGnRHR variant with
164 a deletion downstream of H8 (Δ352-379). Truncation of these residues reduces the PME of cGnRHR 2.0 ± 0.2

fold, (Figure 2), which suggests these residues are important for efficient expression. Finally, to determine whether H8 impacts PME, we characterized a cGnRHR variant lacking the entire C-terminal tail (Δ329-379), which mimics the deletion found in mammalian receptors (Millar, 2005). This truncation reduces the PME of cGnRHR 19.4 ± 0.9 fold (Figure 2), which demonstrates that the truncation of the C-terminal domain is primarily responsible for the diminished PME of mammalian GnRHRs. Nevertheless, the PME of this truncated cGnRHR variant is still 4.2 ± 1.1 fold higher than that of hGnRHR. Therefore, it is likely that additional sequence modifications have helped to tune the PME of mammalian GnRHRs.

*Impact of Sequence Variations on the Topological Energetics of GnRHR*

Previous investigations have concluded that evolutionary modifications of PME arise from variations in the conformational stability of GnRHR (Janovick et al., 2003; Ulloa-Aguirre et al., 2006). Variations in conformational stability may impact the propensity of the receptor to misfold during translocon-mediated cotranslational folding (stage I folding) and/ or during post-translational folding (stage II folding) (Popot & Engelman, 1990). The efficiency of stage I folding primarily depends on the hydrophobicity of TM domains, which dictates the energetics of their translocon-mediated cotranslational membrane integration (Hessa et al., 2005). To assess whether evolutionary adaptations may impact the fidelity of stage I folding, we analyzed the sequences of cGnRHR and hGnRHR using a knowledge-based algorithm that predicts the free energy difference associated with the transfer of nascent TM domains from the translocon to the ER membrane (ΔG predictor) (Hessa et al., 2007). A scan of the cGnRHR sequence reveals that six of its seven TM domains have pronounced energetic minima, four of which have negative transfer free energies (Figure S1). This observation suggests that most TM domains within cGnRHR are sufficiently hydrophobic to undergo efficient translocon-mediated membrane integration. By comparison, only two of the seven TM domains within hGnRHR have negative transfer free energies (Figure S1), which suggests this protein may be more prone to the formation of topological defects during stage I folding.

To determine whether these differences are reflective of a wider evolutionary trend, we used the ΔG predictor to scan the sequences of a total 59 known GnRHRs (Table S1). Projection of the average predicted transfer free energies across the seven TM domains of each receptor onto a phylogenetic tree reveals stark contrasts in the topological energetics of mammalian and non-mammalian GnRHRs (Figure 3). The average predicted transfer free energies are significantly higher for mammalian GnRHRs relative to those of the non-mammalian receptors (Figure 4A, Mann-Whitney $p = 5 \times 10^{-14}$). A comparison of the distribution of predicted transfer free energies for individual domains reveals that evolutionary adaptations resulted in particularly stark increases in the polarity of TM2 and TM6 (Figure 4B). It is unclear how sequence modifications within these domains may have impacted the energetics of post-translational folding reactions or functional signaling. Nevertheless, heightened predicted transfer free energies of mammalian TM2 and TM6 suggest the modifications within these regions could potentially compromise the efficiency of the cotranslational folding of mammalian GnRHRs.

**Figure 4. Comparison of the topological energetics of mammalian and non-mammalian GnRHRs.** The distribution of predicted transfer free energies associated with the translocon-mediated membrane integration of the TM domains within 59 known GnRHRs are shown. A) A histogram depicts the distribution of the average transfer free energies across the seven TM domains of the non-mammalian (blue) and mammalian (red) GnRHRs. B) A series of violin plots depict the distribution of predicted transfer free energies for each individual TM domain found within the non-mammalian (blue) and mammalian (red) receptors. The position of the median value is indicated by a horizontal line within each distribution. The shapes of the histograms and violins were generated using a kernel smoothing function.

**Figure 5. Translocon-mediated membrane integration of TMs 2 & 6.** Differences in the sequences of TM2 and TM6 are analyzed in relation to differences in their efficiency of translocon-mediated membrane integration. A) Logo plots depict the most common amino acid at each position within TM2 of non-mammalian (top) and mammalian (bottom) GnRHRs. The positions of polar substitutions are indicated with a red box. B) Logo plots depict the most common amino acid at each position within TM6 of non-mammalian (top) and mammalian (bottom) GnRHRs. The positions of polar substitutions are indicated with a red box. C) A cartoon depicts the manner in which the translocon-mediated membrane integration of the guested TM domain within chimeric Lep proteins impacts their glycosylation. A failure of the guest TM domain to undergo translocon-mediated membrane integration results in the glycosylation of two residues (top), whereas the membrane integration of the guest domain results in a single glycosylation (bottom). D) Chimeric Lep proteins containing the mammalian or non-mammalian consensus sequences for TM2 and TM6 were translated in canine rough microsomes and analyzed by SDS-PAGE. Negative control reactions lacking RNA (first lane) or containing RNA but lacking microsomes (second lane) are shown for reference. The positions of the untargeted (no glycans), singly glycosylated (G1), and doubly glycosylated (G2) forms of the protein are indicated for reference.

To further explore these TM domains, we constructed logo plots that depict the most common amino acids found
at each positions within TM2 and TM6 of mammalian and non-mammalian GnRHRs. The sequences of the non-
mammalian TM domains are more diverse than those of the mammalian receptor (Figure 5 A & B), which reflects
the increased evolutionary distance between the non-mammalian sequences (Figure 3). Nevertheless, a
comparison of the most common amino acids at each position reveals the heightened transfer free energies of
mammalian GnRHRs primarily arise from three polar substitutions in TM2 and two polar substitutions in TM6
(Figure 5 A & B). To assess the impact of these substitutions on the fidelity of stage I folding, we compared the
translocon-mediated membrane integration of the consensus versions of the mammalian and non-mammalian
TM domains. Briefly, a series of chimeric leader peptidase (Lep) proteins containing each TM domain of interest
was produced by *in vitro* translation in canine rough microsomes. The membrane integration efficiency of each
TM domain can then be inferred from the glycosylation state of Lep; membrane integration of the TM domain
results in a single glycosylation whereas passage into the lumen generates two glycosyl modifications (Figure
5C). The Lep protein containing the non-mammalian TM2 is produced as a mix of both glycoforms (Figure 5D).
In contrast, the doubly glycosylated form predominates for the Lep protein containing the mammalian TM2
(Figure 5D), which demonstrates that the increased polarity of the mammalian TM2 compromises its recognition
by the translocon. Similarly, the membrane integration of the non-mammalian form of TM6 appears to be
significantly more efficient than that of the corresponding mammalian form (Figure 5D). The apparent transfer
free energies associated with the translocon-mediated membrane integration of these helices deviate slightly
from the predicted values (Table S2). Nevertheless, these predictions correctly predicted the manner in which
these mutations should impact the efficiency of membrane integration. Thus, consistent with computational
predictions (Figure 4B), these results show that the increased polarity of TM2 and TM6 of mammalian GnRHRs
decreases the efficiency of their translocon-mediated membrane integration.

*Structural Context of Polar Residues and Their Impacts on PME*

**Figure 6. Structural context and proteostatic effects of polar residues within TMs 2 & 7.** Structural models of catfish (blue) and human (red) GnRHR illustrate the structural context of polar side chains within TMs 2 & 6, and the effects of hydrophobic substitutions at these positions on the plasma membrane expression (PME) of human GnRHR is shown. A) Structural homology models of human (red) and catfish (blue) GnRHRs are overlaid for reference. B) A cutaway of the shows that the polar residues of interest within TM2 (K81, T84, and E90) are buried within the core of the human GnRHR protein. C) A side view shows that the polar residues of interest within TM6 of human GnRHR appear to be projected into the membrane core. D) Polar side chains within TMs 2 & 6 were replaced with the most common hydrophobic residues found within non-mammalian GnRHRs, and the effects of these substitutions one the plasma membrane expression (PME) of human GnRHR was measured in HEK293T cells by flow cytometry. A bar graph depicts the mean fluorescence intensity associated with the surface immunostaining of each variant normalized relative to that of WT human GnRHR. Values reflect the average of three biological replicates, and error bars reflect the standard deviation.

Logo plots show that several polar residues were introduced within TM2 and TM6 during the evolutionary
adaptation of mammalian GnRHRs (Figure 5 A & B). Though these mutations disrupt cotranslational folding
(Figure 5D), it is possible that they also help to stabilize the structure of the folded receptor and/ or enhance its
function. To gain insights into the structural context of these residues, we constructed comparative models of
both the cGnRHR and hGnRHR receptors. With the exception of the C-terminal tail of cGnRHR, both forms of
the receptor have a similar architecture (Figure 6A, Cα RMSD 2.95 Å). The model of hGnRHR reveals that each
of the three polar substitutions within TM2 occur at position that are buried within the protein core (Figure 6B).
Thus, these side chains are likely to play integral roles in the native structure and/ or function of mammalian
GnRHRs. In contrast, the two polar substitutions within TM6 appear to have occurred at surface residues that
are exposed to the hydrophobic core of the lipid bilayer (Figure 6C). Though a complex role of these side chains
in the native conformational dynamics cannot be ruled out, this observation raises the possibility that these
mutations primarily tune the fidelity of cotranslational folding and the corresponding PME without impacting
function.

If the inefficient cotranslational membrane integration of TM2 and/or TM6 contributes to the proteostatic
adaptations in mammalian GnRHRs, then substitutions that enhance the hydrophobicity of these domains could
potentially restore their PME. We therefore measured the PME of a series of hGnRHR variants containing
ancestral substitutions that partially restore the hydrophobicity of TM2 and TM6. In each case, replacing polar
residues in TM2 with the hydrophobic consensus non-mammalian residue resulted in a further reduction in the
PME of hGnRHR (Figure 6D). This result is perhaps unsurprising. While these substitutions are likely to enhance
cotranslational folding, the structural model of hGnRHR suggests they are also likely to introduce packing defects
in the native structure (Figure 6B). Thus, the structural basis for the net proteostatic impact of these substitutions
is unclear. In contrast, consensus hydrophobic substitutions at surface residues within TM6 appear to be well-
tolerated. Though T274L has no impact on PME, replacing T277 with an isoleucine enhances the PME of GnRHR
3.6 ± 0.5 fold (Figure 6D). Thus, restoring a hydrophobic side chain to a single lipid-exposed residue within TM6
(Figure 6C) is sufficient to partially recover the PME of hGnRHR. In conjunction with *in vitro* translation
measurements (Figure 5D), this observation implies that the enhanced cotranslational misfolding of mammalian
TM6 contributes to the attenuated PME of these proteins.

Our results collectively reveal that the enhanced polarity of TM6 compromises the cotranslational folding and expression of mammalian GnRHRs, and that the reversion of a single surface residue to its ancestral hydrophobic side chain (T277I) is sufficient to partially recover PME. Nevertheless, it is possible that this side chain was introduced to support GnRHR function. To determine whether this residue is important for GnRHR signaling, we compared the activity of WT and T277I hGnRHRs. Briefly, cells transiently expressing WT or T277I GnRHR were titrated with gonadotropin-releasing hormone (GnRH), and the receptor activation was indirectly measured by the magnitude of the resulting cytosolic calcium flux. Cells expressing these receptors exhibit robust response to GnRH, and the magnitude of the calcium flux was quite similar for WT and T277I hGnRHR (Figure 7). Moreover, the fitted $EC_{50}$ values for WT (0.61 ± 0.38 μM) and T277I (0.23 ± 0.18 μM) were found to be statistically indistinguishable (Figure 7). These findings demonstrate that T277 is not essential for hGnRHR function. Given that this non-essential polar side chain negatively impacts cotranslational folding and PME, our collective observations suggest that the evolved polarity of this segment serves to tune the PME of mammalian GnRHRs.

# Discussion

Previous investigations on the evolution of GnRHR have suggested that its activity has been down-regulated in mammals through various coding mutations that enhanced its propensity to misfold (Janovick et al., 2003; Leaños-Miranda et al., 2003; Lin et al., 1998). In this work, we followed up on these investigations in order to assess the molecular basis for this evolved instability. Using quantitative cellular measurements, we first confirmed that the catfish receptor exhibits more robust expression and cellular trafficking relative to mammalian GnRHRs. To determine which mutations contributed to the lapse in mammalian GnRHR proteostasis, we then characterized the effects of various sequence modifications on the PME of cGnRHR. Our results provide additional evidence that the C-terminal truncation in mammalian GnRHRs is likely a key factor leading to their attenuated PME (Figure 2). However, an analysis of known GnRHR sequences also shows that the TM domains of mammalian GnRHRs are more polar than their non-mammalian counterparts (Figures 3 & 4). We find that this increased polarity compromises the translocon-mediated cotranslational folding of at least two TM domains (Figure 5). Moreover, we show that restoring a hydrophobic amino acid at a surface residue within TM6 increases the PME of hGnRHR 3.6-fold with no impact on receptor activation (Figures 6 & 7). This modification is likely relevant to the evolutionary trajectory of GnRHR considering the mouse receptor, which exhibits a higher PME than the human receptor, features a hydrophobic residue at this position (V276).

**Figure 7. Impact of 277I on GnRHR function.** The activation of the WT and T277I forms of human GnRHR in response to human gonadotropin releasing hormone (GnRH) was measured in HEK293T cells. Cells transiently expressing each receptor were stimulated with GnRH, and signaling was measured by the change in the fluorescence intensity of a cytosolic calcium reporter. The average fluorescence intensities of cells expressing WT (●) or T277I (○) from three technical replicates are normalized relative to baseline and plotted against the corresponding concentration of hormone. Error bars reflect the standard deviation from three technical replicates. Curves reflect the fit of the WT (black) and T277I (gray) data to a single-site binding model.

There are several caveats to these investigations. First, it should be noted that epistatic interactions between some of these mutations may alter their effects on PME in the context of mammalian receptors. Furthermore, we cannot measure the PME of these receptors in the context of their native environment within the pituitary gland of each animal. It is likely that the magnitude of these proteostatic effects are distinct in the context of the native proteostasis networks that typically support GnRHR biogenesis. Nevertheless, to our knowledge mammalian GnRHRs are the only class A GPCRs that completely lack helix 8 and/ or a C-terminal tail (Millar, 2005). We suspect this drastic modification is likely to have consequences for the maturation and sorting of mammalian GnRHRs within the secretory pathway regardless of the cellular context. It should also be noted that the mechanism of the translocon is highly conserved, and the increased polarity of TM2 and TM6 are therefore likely to reduce the efficiency of cotranslational GnRHR folding in any cellular context (Denks et al., 2014). Indeed, the hydrophobicity of TM

domains is also known to be a critical factor that governs the expression of MPs in *E. coli* (Marshall et al., 2016; Michiel et al., 2017). Based on these considerations, it seems likely that both the C-terminal truncation and the enhanced the polarity of the TM domains of GnRHR are likely to have contributed to evolutionary modifications to the PME and net activity of mammalian GnRHRs.

Our findings provide additional evidence to suggest the activity of mammalian GnRHRs has been tuned through modulation of GnRHR folding rather than through transcriptional modifications. This is perhaps surprising given the metabolic cost of protein synthesis. Nevertheless, we believe this outcome is reasonable in light of certain evolutionary considerations. It should first be noted that the length of the open reading frame of GnRHR is roughly six times that of its promoter (Schang et al., 2012). Considering most coding variants are destabilizing, there are likely to be far more coding variants that decrease the PME of the receptor relative to the number that would simply decrease its transcription. If selection pressures simply favored attenuated GnRHR signaling, then it is perhaps most probable this would arise from mutations that destabilize the native GnRHR structure. Consistent with the observed proteostatic patterns (Figure 1), such mutations would result in both a decreased PME and an increased accumulation of the receptor within the secretory pathway. Though energetically wasteful, it is unclear whether the biosynthesis of misfolded GnRHRs necessarily imposes a significant fitness burden in this case, as this receptor is only expressed at moderate levels within the pituitary gland according to the human protein atlas (proteinatlas.org). Thus, it seems plausible that various destabilizing mutations fixed in mammals, provided the gain in reproductive fitness outweighs the metabolic costs associated with the synthesis and degradation of misfolded receptors.

Marginal conformational stability is an emergent property of naturally evolved proteins (Bloom, Labthavikul, Otey, Arnold, & Levitt, 2006; Taverna & Goldstein, 2002; Tokuriki & Tawfik, 2009). This instability has been attributed to the net-destabilizing effects of random mutations in conjunction with a general lack of selection pressure for hyper-stable proteins (Tokuriki & Tawfik, 2009). Nevertheless, the natural history of GnRHR suggests that the instability of native protein structures provides additional evolutionary opportunities to tune protein fitness. In the context of integral MPs, the evolutionary utility of sub-optimal cotranslational folding energetics is illustrated by the fact that the PME of rhodopsin is highly sensitive a wide variety of mutations within its TM domains (Penn et al., n.d.). The natural exploitation of the thin energetic margins involved in the cotranslational MP folding perhaps also provides an explanation for the fact that the hydrophobicity of rhodopsin's TM domains has not been optimized to promote its efficient biosynthesis (Roushar et al., 2019). However, this malleability of cotranslational MP folding energetics does not come without costs, as TM domains are generally less tolerant of genetic variation and mutations within TM domains give rise to numerous genetic diseases (Marinko et al., 2019; Schlebach & Sanders, 2015; Telenti et al., 2016). Together, these observations provide new insights into the evolutionary balance between the constraints of protein folding and function.

## Materials & Methods

*Plasmid Preparation and Mutagenesis*

A series of pcDNA5 FRT expression vector containing various GnRHR cDNAs containing an N-terminal influenza hemaglutinin (HA) epitope were used for the transient expression of GnRHR variants. GnRHR cDNAs in this vector are followed by an internal ribosome entry site (IRES) and eGFP sequence, which generates bicistronic GFP expression in positively transfected cells. Vectors containing GnRHRs from various species were generated using In-Fusion HD Cloning (Takara Bio, Shiga, Japan). Mutations were introduced by site-directed mutagenesis, and truncations were generated by In-Fusion HD cloning. To adapt these constructs for functional experiments, the HA epitope was deleted to minimize interferences with ligand binding, and the IRES eGFP sequence was deleted to prevent interference of eGFP with fluorescence measurements.

A previously described pGEM expression vector containing modified leader peptidase (Lep) cDNA was used for *in vitro* translation (Hessa et al., 2007). TM domains of interest were cloned into the H-segment position within the Lep gene using directional cloning at the *Spe*I and *Kpn*I restriction sites. All plasmids were prepared with the Endotoxin-Free Zymopure Midiprep or Miniprep Kit (Zymo Research, Irvine, CA).

*In vitro Translation of Chimeric Lep Proteins*

Messenger RNA (mRNA) was generated for each chimeric Lep gene using the RiboMAX SP6 kit (Promega, Madison, WI). Lep variants were then translated using rabbit reticulocyte lysate (Promega, Madison, WI) supplemented with canine rough microsomes (tRNA probes, College Station, TX), and EasyTag $^{35}$S-labeled methionine (PerkinElmer, Waltham, MA). Translation was carried out at 30 °C for 60 minutes. Reactions were diluted 1:4 in 1X SDS PAGE loading buffer and separated on a 12% SDS PAGE gel. Gels were then dried, exposed overnight on a phosphor imaging plate (GE Healthcare, New York, NY), and imaged on a Typhoon Imager (GE Healthcare, New York, NY). The ratio of singly (*G1*) to doubly (*G2*) glycosylated Lep protein was quantified by densitometry using ImageJ software. The *G1:G2* ratio represents an apparent equilibrium constant (*K$_{app}$*) for the transfer of the H-segment from the translocon to the membrane, as previously described Apparent transfer free energy values for the H-segments were calculated using the following equation:

$$\Delta G_{app} = -RT \, \ln(K_{app}) = -R \, T \, \ln(\frac{G1}{G2})$$

where Δ*G$_{app}$* represents the apparent free energy for the transfer of the H-segment into the membrane, *R* represents the universal gas constant, *T* represents the temperature, *K$_{app}$* represents the apparent equilibrium constant for the transfer of the H-segment from the translocon into the membrane, *G1* represents the intensity of the singly glycosylated band and *G2* represents the intensity of the doubly glycosylated band, as was previously described (Hessa et al., 2005). Reported transfer free energy values represent the average values from three experimental replicates.

*Cellular GnRHR Expression Measurements*

To quantitatively measure the cellular trafficking of GnRHR variants, plasma membrane and intracellular GnRHRs were differentially immunostained and analyzed by flow cytometry, as described previously (Schlebach et al., 2015). GnRHR variants were transiently expressed in HEK293T cells using Lipofectamine 3000 (Invitrogen, Carlsbad, CA). Two days after transfection, the cells were harvested with TrypLE Express protease (Gibco, Grand Island, NY). Plasma membrane GnRHRs were then immunostained for 30 minutes in the dark with a DyLight 550-conjugated anti-HA antibody (Invitrogen, Carlsbad, CA). Cells were fixed and permeabilized using a Fix and Perm kit (Invitrogen, Carlsbad, CA), and washed twice with 2% fetal bovine serum in phosphate-buffered saline (wash buffer). Intracellular GnRHRs were then immunostained for 30 minutes in the dark using an Alexa Fluor 647-conjugated anti-HA antibody (Invitrogen, Carlsbad, CA). Cells were washed twice in order to remove excess antibody prior to analysis of cellular fluorescence profiles. Fluorescence profiles were analyzed on a BD LSRII flow cytometer (BD Biosciences, Franklin Lakes, NJ). Forward and side scatter measurements were used to set a gate for intact single cells. eGFP intensities (488 nm laser, 530/30 nm emission filter) were then used to set a gate for positively-transfected cells. DyLight 550 (561 nm laser, 582/15 nm emission filter) and Alexa Fluor 647 (640 nm laser, 670/30 nm emission filter) intensities were then calculated for several thousand positively-transfected single cells within each biological replicate. Data were analyzed using FlowJo software (Treestar, Ashland, OR). Characterizations of GnRHR variant expression levels were carried out with at least three biological replicates each.

*Functional Measurements of GnRHRs*

GnRHR activity was measured in HEK293T cells by monitoring cytosolic calcium fluxes that occurred in response to gonadotropin-releasing hormone (GnRH, Sigma Aldrich, St. Louis, MO). GnRHR variants were transiently expressed in HEK293T cells using Lipofectamine 3000 (Invitrogen, Carlsbad, CA). Two days after transfection, the cells were harvested with TrypLE Express (Gibco, Grand Island, NY), then re-plated in 96-well plates (Corning, Big Flats, NY) coated with poly-D-lysine (Gibco, Grand Island, NY) at a density of 60,000 cells per well. Cells were then dosed the following day and assayed using the FLIPR Calcium 6-QF Assay Kit (Molecular Devices, San Jose, CA) according to the manufacturer's protocol. The fluorescence intensities of cells incubated in the calcium-sensitive FLIPR dye was measured for thirty seconds prior to dosing with GnRH using a Synergy Neo2 microplate reader (BioTek, Winooski, VT) using an excitation wavelength of 485/20 nm and an emission filter at 525/10 nm. Directly after dosing the cells, the change in fluorescence was measured for six minutes. The percent calcium flux under each condition was calculated for each well using the following equation:

$$\text{Percent Calcium Flux} = \frac{M - B}{B} \times 100$$

where $M$ is the maximum fluorescence value for the calcium flux and $B$ is the baseline signal as was determined from by averaging the fluorescence intensity before ligand addition. $EC_{50}$ values were determined by fitting titrations to the following function:

$$Y = A + \frac{B - A}{1 + m \cdot 10^{C-x}}$$

where $Y$ is the percent calcium flux, $A$ is the minimal curve asymptote, $B$ is the maximal curve asymptote, $m$ is the slope of the transition region, $C$ is the logarithm of the $EC_{50}$, and $x$ is the logarithm of the GnRH concentration (Motulsky & Christopoulos, 2003). Reported $EC_{50}$ values represent the average from three biological replicates.

*Selection and Analysis of GnRHR Sequences*

59 GnRHR sequences from different species were collected from the NCBI (https://www.ncbi.nlm.nih.gov) and Uniprot (https://www.uniprot.org) databases. Humans have only one type of GnRHR (GnRHR-I), while other species may have multiple types (Pawson, Morgan, Maudsley, & Millar, 2003; Tello, Wu, Rivier, & Sherwood, 2008). Sequences selected for phylogenetic analysis were therefore limited to those annotated as GnRHR-I in order to analyze trends across species.

A phylogenetic tree was generated from these sequences using MEGA7 software (megasoftware.net). A sequence alignment was generated using the MUSCLE alignment tool with default settings. This alignment was then used to construct a Maximum Likelihood tree (Kumar, Stecher, & Tamura, 2016). The positions of nascent TM domains within each sequence were then identified from energetic minima generated with a window scan function within the ΔG predictor, which sums depth-dependent free energies associated with the transfer of amino acids from the translocon to the ER membrane (http://dgpred.cbr.su.se/) (Hessa et al., 2007). The ΔG predictor was then used to calculate the free energy difference associated with the translocon-mediated membrane integration of each putative TM domain (Hessa et al., 2007). The phylogenetic tree and ΔG prediction data were then uploaded to the Interactive Tree of Life (https://itol.embl.de), where the ΔG predictions were displayed as color gradients on the phylogenetic tree (Letunic & Bork, 2016).

To generate logo plots, the GnRHR-I sequences were first aligned in ClustalOmega (Sievers et al., 2011). The positions of the TM domains within the hGnRHR sequence were determined by the ΔG predictor, and the transmembrane domains in other species were then identified by the corresponding positions in the alignment. Sequence logo plots were then generated for each transmembrane domain in the mammalian and non-mammalian sequences using the WebLogo application (https://weblogo.berkeley.edu/logo.cgi) (Crooks, Hon, Chandonia, & Brenner, 2004).

*Structural Modeling*

Comparative models of the human and catfish forms GnRHR were generated using multi-template comparative modeling in Rosetta. The GnRHR sequence was first aligned with sequences for 34 GPCR crystal structures obtained from GPCRdb (http://www.gpcrdb.org) (Isberg et al., 2016). Manual adjustments were then made to account for well-known conserved residues in loop regions and TM domains (Bender, Marlow, & Meiler, 2019). OCTOPUS was used to define the TM domains, and the two disulfide bonds were defined manually (Millar et al., 2004; Viklund & Elofsson, 2008). To generate a model of GnRHR in the inactive state, the sequences were threaded onto the antagonist-bound structures of several other Class A Group β GPCRs including the human OX2 orexin receptor (HCRTR2, PDB 4S0V, 2.5 Å), human OX1 orexin receptor (HCRTR1, PDB 4ZJC, 2.8 Å), human endothelin receptor type B (EDNRB, PDB 5X93, 2.2 Å), and human neuropeptide Y receptor Y1 (NPY1R, PDB 5ZBQ, 2.7 Å). Threading was completed using the *partial thread* application in RosettaCM (Song et al., 2013). 1,000 models were then generated using the *hybridize* application in RosettaCM and the TM domains were relaxed using a set of optimized RosettaMembrane weights that were modified from the Talaris scoring

function (Duran & Meiler, 2018). The models with the lowest Rosetta energy score were used for structural analysis.

## Acknowledgments

We thank Christiane Hassel and the Indiana University Bloomington Flow Cytometry Core Facility for their experimental support. This research was supported in part by grant from the National Institute of General Medical Sciences (NIGMS) under award numbers R01GM129261, R01GM080403, R01HL122010, and R01DA046138. This work was also supported by a National Science Foundation (NSF) Graduate Research Fellowship to L.M. C. (grant number 1342962). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the NSF or NIH.

## References

Bemji, M. N., Isa, A. M., Ibeagha-Awemu, E. M., & Wheto, M. (2018). Polymorphisms of caprine GnRHR gene and their association with litter size in West African Dwarf goats. *Molecular Biology Reports*, *45*, 63–69. https://doi.org/10.1007/s11033-017-4141-0

Bender, B. J., Marlow, B., & Meiler, J. (2019). RosettaGPCR: Multiple Template Homology Modeling of GPCRs with Rosetta. *BioRxiv*. https://doi.org/10.1101/2019.12.13.875237

Bloom, J. D., Labthavikul, S. T., Otey, C. R., Arnold, F. H., & Levitt, M. (2006). *Protein stability promotes evolvability*. Retrieved from www.pnas.orgcgidoi10.1073pnas.0510098103

Conn, P. M., McArdle, C. A., Andrews, W. V., & Huckle, W. R. (1987). The Molecular Basis of Gonadotropin-Releasing Hormone (GnRH) Action in the Pituitary Gonadotrope. *Biology of Reproduction*, *36*(1), 17–35. https://doi.org/10.1095/biolreprod36.1.17

Conn, P. Michael, Janovick, J. A., Brothers, S. P., & Knollman, P. E. (2006). "Effective inefficiency": Cellular control of protein trafficking as a mechanism of post-translational regulation. *Journal of Endocrinology*, *190*(1), 13–16. https://doi.org/10.1677/joe.1.06771

Conn, P. Michael, Knollman, P. E., Brothers, S. P., & Janovick, J. A. (2006). Protein folding as posttranslational regulation: Evolution of a mechanism for controlled plasma membrane expression of a G protein-coupled receptor. *Molecular Endocrinology*, *20*(12), 3035–3041. https://doi.org/10.1210/me.2006-0066

Conn, P. Michael, & Ulloa-Aguirre, A. (2011). Pharmacological Chaperones for Misfolded Gonadotropin-Releasing Hormone Receptors. *Advances in Pharmacology*, *62*(503), 109–141. https://doi.org/10.1016/B978-0-12-385952-5.00008-7

Crooks, G. E., Hon, G., Chandonia, J.-M., & Brenner, S. E. (2004). WebLogo: A Sequence Logo Generator. *Genome Research*, *14*, 1188–1190. https://doi.org/10.1101/gr.849004

Denks, K., Vogt, A., Sachelaru, I., Petriman, N.-A., Kudva, R., & Koch, H.-G. (2014). The Sec translocon mediated protein transport in prokaryotes and eukaryotes. *Molecular Membrane Biology*, *31*(3), 58–84. https://doi.org/10.3109/09687688.2014.907455

Duran, A. M., & Meiler, J. (2018). Computational design of membrane proteins using RosettaMembrane. *Protein Science*, *27*(1), 341–355. https://doi.org/10.1002/pro.3335

Harms, M. J., & Thornton, J. W. (2013). Evolutionary biochemistry: revealing the historical and physical causes of protein properties. *Nature Reviews Genetics*, *14*(8), 559–571. https://doi.org/10.1038/nrg3540

Hessa, T., Kim, H., Bihlmaier, K., Lundin, C., Boekel, J., Andersson, H., … Von Heijne, G. (2005). Recognition of transmembrane helices by the endoplasmic reticulum translocon. *Nature*, *433*. Retrieved from www.nature.com/nature

Hessa, T., Meindl-Beinker, N. M., Bernsel, A., Kim, H., Sato, Y., Lerch-Bader, M., … Gunnar Von Heijne, &. (2007). Molecular code for transmembrane-helix recognition by the Sec61 translocon. *Nature*, *450*. https://doi.org/10.1038/nature06387

Isberg, V., Mordalski, S., Munk, C., Rataj, K., Harpsøe, K., Hauser, A. S., … Gloriam, D. E. (2016). GPCRdb: an information system for G protein-coupled receptors. *Nucleic Acids Research*, *44*, D356–D364. https://doi.org/10.1093/nar/gkv1178

Janovick, J. A., David Stewart, M., Jacob, D., Martin, L. D., Deng, J. M., Stewart, A., … Conn, P. M. (2013). Restoration of testis function in hypogonadotropic hypogonadal mice harboring a misfolded GnRHR mutant by pharmacoperone drug therapy. *PNAS*, *110*(52), 21030–21035. https://doi.org/10.1073/pnas.1315194110

Janovick, J. A., Knollman, P. E., Brothers, S. P., Ayala-Yáñ Ez, R., Aziz, A. S., & Conn, P. M. (2006).

Regulation of G Protein-coupled Receptor Trafficking by Inefficient Plasma Membrane Expression MOLECULAR BASIS OF AN EVOLVED STRATEGY. *The Journal of Biological Chemistry*, *281*(13), 8417–8425. https://doi.org/10.1074/jbc.M510601200

Janovick, J. A., Ulloa-Aguirre, A., & Conn, P. M. (2003). Evolved Regulation of Gonadotropin-Releasing Hormone Receptor Cell Surface Expression. In *Endocrine* (Vol. 22).

Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, *33*(7), 1870–1874. https://doi.org/10.1093/molbev/msw054

Leaños-Miranda, A., Ulloa-Aguirre, A., Ji, T. H., Janovick, J. A., & Conn, P. M. (2003). Dominant-negative action of disease-causing gonadotropin-releasing hormone receptor (GnRHR) mutants: A trait that potentially coevolved with decreased plasma membrane expression of GnRHR in humans. *Journal of Clinical Endocrinology and Metabolism*, *88*(7), 3360–3367. https://doi.org/10.1210/jc.2003-030084

Letunic, I., & Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Research*, *44*. https://doi.org/10.1093/nar/gkw290

Lin, X., Janovick, J. A., Brothers, S., Blömenrohr, M., Bogerd, J., & Conn, P. M. (1998). Addition of catfish gonadotropin releasing hormone (GnRH) receptor intracellular carboxyl-terminal tail to rat GnRH receptor alters receptor expression and regulation. *Molecular Endocrinology*, *12*(2), 161–171. https://doi.org/10.1210/mend.12.2.0056

Marinko, J. T., Huang, H., Penn, W. D., Capra, J. A., Schlebach, J. P., & Sanders, C. R. (2019). Folding and Misfolding of Human Membrane Proteins in Health and Disease: From Single Molecules to Cellular Proteostasis. *Chemical Reviews*, *119*(9), 5537–5606. https://doi.org/10.1021/acs.chemrev.8b00532

Marshall, S. S., Niesen, M. J. M., Müller, A., Tiemann, K., Saladi, S. M., Galimidi, R. P., … Miller, T. F. (2016). A Link between Integral Membrane Protein Expression and Simulated Integration Efficiency. *Cell Reports*, *16*(8), 2169–2177. https://doi.org/10.1016/j.celrep.2016.07.042

Michiel, X., Niesen, J. M., Marshall, S. S., Thomas, X., Miller Iii, F., & Clemons, W. M. (2017). Improving membrane protein expression by optimizing integration efficiency. *J. Biol. Chem*, *292*(47), 19537–19545. https://doi.org/10.1074/jbc.M117.813469

Millar, R. P. (2005). GnRHs and GnRH receptors. *Animal Reproduction Science*, *88*(1-2 SPEC. ISS.), 5–28. https://doi.org/10.1016/j.anireprosci.2005.05.032

Millar, R. P., Lu, Z.-L., Pawson, A. J., Flanagan, C. A., Morgan, K., & Maudsley, S. R. (2004). Gonadotropin-Releasing Hormone Receptors. *Endocrine Reviews*, *25*(2), 235–275. https://doi.org/10.1210/er.2003-0002

Motulsky, H., & Christopoulos, A. (2003). *Fitting Models to Biological Data using Linear and Nonlinear Regression: A practical guide to curve fitting Contents at a Glance*. Retrieved from www.graphpad.com.

Palczewski, K., Kumasaka, T., Hori, T., Behnke, C. A., Motoshima, H., Fox, B. A., … Miyano, M. (2000). Crystal Structure of Rhodopsin: A G Protein-Coupled Receptor. *Science*, *289*(5480), 739–745. Retrieved from http://science.sciencemag.org/

Pawson, A. J., Morgan, K., Maudsley, S. R., & Millar, R. P. (2003). Type II gonadotrophin-releasing hormone (GnRH-II) in reproductive biology. *Reproduction*, *126*, 271–278.

Penn, W. D., McKee, A. G., Kuntz, C. P., Woods, H., Nash, V., Gruenhagen, T. C., … Schlebach, J. P. (n.d.). Probing Biophysical Sequence Constraints within the Transmembrane Domains of Rhodopsin by Deep Mutational Scanning. *Science Advances*, *In Press*.

Popot, J. L., & Engelman, D. M. (1990). Membrane Protein Folding and Oligomerization. *Biochemistry*, *29*(17), 4031–4037. https://doi.org/10.1021/bi00469a001

Roushar, F. J., Gruenhagen, T. C., Penn, W. D., Li, B., Meiler, J., Jastrzebska, B., & Schlebach, J. P. (2019). Contribution of Cotranslational Folding Defects to Membrane Protein Homeostasis. *Journal of the American Chemical Society*, *141*(1), 204–215. https://doi.org/10.1021/jacs.8b08243

Schang, A. L., Quérat, B., Simon, V., Garrel, G., Bleux, C., Counis, R., … Laverrière, J. N. (2012). Mechanisms underlying the tissue-specific and regulated activity of the Gnrhr promoter in mammals. *Frontiers in Endocrinology*, *3*, 1–19. https://doi.org/10.3389/fendo.2012.00162

Schlebach, J. P., Narayan, M., Alford, C., Mittendorf, K. F., Carter, B. D., Li, J., & Sanders, C. R. (2015). Conformational Stability and Pathogenic Misfolding of the Integral Membrane Protein PMP22. *Journal of the American Chemical Society*, *137*, 8758–8768. https://doi.org/10.1021/jacs.5b03743

Schlebach, J. P., & Sanders, C. R. (2015). Influence of Pathogenic Mutations on the Energetics of Translocon-Mediated Bilayer Integration of Transmembrane Helices. *Journal of Membrane Biology*, *248*(3), 371–381.

https://doi.org/10.1007/s00232-014-9726-0

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., … Higgins, D. G. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, *7*. https://doi.org/10.1038/msb.2011.75

Song, Y., DiMaio, F., Yu-Ruei Wang, R., Kim, D., Miles, C., Brunette, T., … Baker, D. (2013). High resolution comparative modeling with RosettaCM. *Structure*, *21*(10). https://doi.org/10.1016/j.str.2013.08.005

Taverna, D. M., & Goldstein, R. A. (2002). Why are proteins marginally stable? *Proteins: Structure, Function and Genetics*, *46*(1), 105–109. https://doi.org/10.1002/prot.10016

Telenti, A., Pierce, L. C. T., Biggs, W. H., Di Iulio, J., Wong, E. H. M., Fabani, M. M., … Venter, J. C. (2016). Deep sequencing of 10,000 human genomes. *PNAS*, *113*(42), 11901–11906. https://doi.org/10.1073/pnas.1613365113

Tello, J. A., Wu, S., Rivier, J. E., & Sherwood, N. M. (2008). Four functional GnRH receptors in zebrafish: analysis of structure, signaling, synteny and phylogeny. *Integrative and Comparative Biology*, *48*(5), 570–587. https://doi.org/10.1093/icb/icn070

Tokuriki, N., & Tawfik, D. S. (2009). Stability effects of mutations and protein evolvability. *Current Opinion in Structural Biology*, *19*(5), 596–604. https://doi.org/10.1016/j.sbi.2009.08.003

Ulloa-Aguirre, A., Janovick, J. A., Miranda, A. L., & Conn, P. M. (2006). G-protein-coupled receptor trafficking: understanding the chemical basis of health and disease. *ACS Chemical Biology*, *1*(10), 631–638. https://doi.org/10.1021/cb600360h

Viklund, H., & Elofsson, A. (2008). OCTOPUS: improving topology prediction by two-track ANN-based preference scores and an extended topological grammar. *Bioinformatics*, *24*(15), 1662–1668. https://doi.org/10.1093/bioinformatics/btn221

Yang, W., Tang, K., Zhang, C., Xu, D., Wen, Q., & Liguo, Y. (2011). Polymorphism of the GnRHR gene and its association with litter size in Boer goats. *South African Journal of Animal Science*, *41*(4). https://doi.org/10.4314/sajas.v41i4.10