Genetic erosion reduces biomass temporal stability in wild fish populations.

Authors:

Jérôme G. Prunier [1]*, Mathieu Chevalier [2]**, Allan Raffard [1]***, Géraldine Loot [3], Nicolas Poulet [4], Simon Blanchet[1,3]*

Affiliations :

[1] Centre National de la Recherche Scientifique (CNRS), Université Paul Sabatier (UPS); Station d'Ecologie Théorique et Expérimentale, UAR 2029, F-09200 Moulis, France

[2] Department of Ecology and Evolution, University of Lausanne, Biophore, CH-1015 Lausanne, Switzerland

[3] CNRS, UPS, École Nationale de Formation Agronomique (ENFA) ; UMR 5174 EDB (Laboratoire Évolution & Diversité Biologique), 118 route de Narbonne, F-31062 Toulouse cedex 4, France

[4] Pôle écohydraulique AFB-IMT, allée du Pr Camille Soula, 31400 Toulouse, France

** *present address :* IFREMER, Centre de Bretagne, DYNECO, Laboratoire d'Ecologie Benthique Côtière (LEBCO), 29280 Plouzané, France.

*** *present address :* Univ. Savoie Mont Blanc, INRAE, CARRTEL, 74200 Thonon-les-Bains, France

* **Corresponding authors:**

Jérôme G. Prunier and Simon Blanchet

Station d'Ecologie Théorique et Expérimentale, UMR 5321, F-09200 Moulis, France

2 route du CNRS, 09200 Moulis

Phone : (+33)561040361

Email : jerome.prunier@gmail.com, simon.blanchet@sete.cnrs.fr

ORCIDs : Jérôme G. Prunier 0000-0003-4110-2567, Simon Blanchet 0000-0002-3843-589X, Mathieu Chevalier 0000-0002-1170-5343

Keywords

Eco-evolutionary dynamics; Genetic erosion; Intraspecific diversity, Biodiversity-Ecosystem function.

ABSTRACT

Genetic diversity sustains species adaptation. However, it may also support key ecosystems functions and services that can be altered by the worldwide loss of genetic diversity. Here, using long-term wild fish data, we demonstrate that populations with high genetic diversity do not reach higher biomasses than populations with low genetic diversity. Nonetheless, populations with high genetic diversity have much more stable biomasses over recent decades than populations having suffered from genetic erosion, which has implications for the provision of ecosystem services and the risk of population extinction. Our results strengthen the importance of adopting prominent environmental policies to conserve this important biodiversity facet.

INTRODUCTION

Biodiversity sustains critical ecosystem services, such as water filtering, pollination or biomass production[1], that are directly compromised by the ongoing global biodiversity crisis[2]. By promoting trait complementarity or redundancy among species, interspecific diversity allows ecological communities to optimally capture essential resources, to transform those resources into biomass and to recycle them[3,4]. In species-rich communities, these ecological processes are maintained even in the face of environmental fluctuations, thus promoting ecosystem productivity and stability over time[5,6]: this is the insurance effect of species richness[7].

Although biodiversity erosion is often associated to species loss, another form of erosion is silently underway: the loss of intraspecific genetic diversity[8]. Intraspecific genetic diversity can play a role similar to species diversity in driving ecological processes at the basis of ecosystem services, such as biomass production[9,10]. Beyond its positive influence on individual fitness and thus on *per capita* biomass production, intraspecific genetic diversity may favor functional complementarity or redundancy among individuals, thereby fostering a more efficient exploitation of available resources[1,7,9]. Genetically-diversified populations are therefore predicted to harbor both larger individuals (higher *per capita* biomass) and higher and more stable levels of total biomass than genetically-impoverished populations[9,10]. This direct relationship between intraspecific genetic diversity and biomass is expected to be particularly strong in ecosystems where species diversity is naturally low, which is actually the norm in many temperate ecosystems[11]. In such cases, the

functioning of ecosystems probably depends more on the complementarity among genotypes than on the

complementarity among species[12,13], emphasizing the importance of maintaining genetic diversity to

preserve ecosystem functions and services[9].

Most studies investigating the relationship between intraspecific genetic diversity and key-ecological

parameters such as biomass are based on experimental or semi-experimental settings, where population

densities and levels of intraspecific genetic diversity are manipulated, while environmental conditions are

controlled and maintained constant over time[9,14]. However, observational studies conducted in natural

settings are still scarce and mostly concern plants[15]. Although these studies offer a number of advantages,

experiments do not make it possible to cover large spatial and temporal scales or to investigate the influence

of historical contingencies. Local levels of intraspecific genetic diversity indeed result from the interplay

between long-term evolutionary trajectories (e.g., localization and size of glacial refugia[16]) and more recent

–if not ongoing– ecological processes affecting individual life history traits or population demography (e.g.,

stressful environmental conditions[17], bottleneck events[18], or strong directional selection[19]). This natural

complexity cannot be fully grasped by experimental studies. Observational field surveys are on the contrary

more realistic and may provide important insights into the contribution of intraspecific genetic diversity, and

the loss of it, to biomass and biomass stability in natural settings[15]. They yet raise several difficulties. First,

assessing the influence of genetic diversity on biomass and biomass stability over several generations or

seasonal cycles implies long-term monitoring programs of both population density and biomass, but such

data are usually difficult to collect and are still scarce. Furthermore, the relationships between genetic

diversity and biomass in across-population studies may be masked by the interplay with other factors also

involved in biomass production, such as population density and environmental conditions, making it difficult

to disentangle their respective contributions[15]. This last issue may however be partly alleviated through the

use of causal modeling procedures (path analyses), making it possible to thoroughly confront theoretical

expectations and experimental findings with the "real world"[3,20].

Here, we capitalized on long-term field surveys of three parapatric non-commercial freshwater fish

species (*Phoxinus dragarum*, *Gobio occitaniae* and *Squalius cephalus*) from two large river basins in south-

western France (Fig. 1) to assess the relationships between intraspecific genetic diversity, total fish biomass

and biomass temporal stability (measured as the inverse of biomass variability[21]), while controlling for the

effect of environment (upstream-downstream gradient and eutrophication levels), of *per capita* biomass (or

its temporal stability) and of past demographic events, using path analyses[22]. Total fish biomass stood for

the total weight of all individuals from the three focal species collected at a given site (in $g.m^{-2}$), standardized

and then averaged across species and over years. Local levels of intraspecific genetic diversity were

computed for each species using both microsatellite and SNPs data and similarly averaged across species.

We addressed the following questions: Do the positive relationships found experimentally between

intraspecific genetic diversity and total biomass and total biomass stability hold true in natural settings? If

any, is the contribution of intraspecific genetic diversity to these ecosystem functions comparable in

magnitude to that of other environmental determinants? Finally, is it possible to detect the impact of

contemporary genetic erosion -i.e., the loss of intraspecific genetic diversity in response to a recent

reduction in population size- on biomass and biomass stability of fish populations? This latter point is of high

concern: with conservative estimates of 6-15% loss of intraspecific genetic diversity in wild organisms in the

Anthropocene[23,24], the impact of human-induced genetic erosion on natural ecosystems' capacity to provide

critical provisioning and regulating services to humanity may actually be much more important than

anticipated.

RESULTS

Drivers of intraspecific genetic diversity. We found that intraspecific genetic diversity increased

downstreamward (Fig. 2 and 3B; Supplementary Table 1), a classical pattern in rivers that could stem from

asymmetrical gene flow, the presence of glacial refuges and/or higher effective population sizes in

downstream areas[25]. Intraspecific genetic diversity was also indirectly impacted by water eutrophication,

through a higher probability of having suffered from a bottleneck as eutrophication increases (Fig. 2 and

3A). As expected, the bottleneck probability altered spatial patterns of intraspecific genetic diversity: the loss

of intraspecific genetic diversity associated with recent bottlenecks was particularly strong in downstream

populations (Fig. 3B). This context-dependency of contemporary genetic erosion may reflect the observation

that downstream areas are usually subject to multiple stressors (pollution, urbanization, channelization, …)

that may reinforce the link between recent demographic changes and intraspecific genetic diversity.

Drivers of total biomass. We found no significant relationship between intraspecific genetic diversity and

either total biomass or *per capita* biomass (Fig. 2A; Supplementary Table 1). Yet, total biomass was directly

linked to *per capita* biomass (Fig. 3D), indicating that total biomass stemmed from the presence of (a few)

large individuals rather than that of many small individuals. *Per capita* biomass decreased downstreamward

and increased with eutrophication (Fig. 3C), suggesting that, although eutrophication may indirectly have a

long-term negative influence on populations and associated intraspecific genetic diversity, it may locally

boost the overall system productivity by favoring individuals' body condition[26].

Drivers of biomass stability. Contrastingly, we found that fish populations with higher levels of intraspecific

genetic diversity displayed higher biomass stability over time than genetically-impoverished populations,

whatever the temporal stability in *per capita* biomass (i.e., positive relationship between biomass stability

and intraspecific genetic diversity: Fig. 2B and 3F; Supplementary Table 1). This important finding also held

true when each species was analyzed separately (Supplementary Table 2). By favoring functional

complementarity or redundancy among phenotypes[9,10,27], higher genetic diversity likely allows populations

to maintain an efficient exploitation of available resources in the face of natural environmental fluctuations,

ensuring a stable production of biomass[1,7]. Biomass stability also tended to decrease with eutrophication

(Fig. 2B and 3F; Supplementary Table 1), probably because biomass stability was negatively related to total

biomass (Supplementary Note 1), and thus indirectly to *per capita* biomass, the latter increasing with

eutrophication (Fig. 2A and 3). Fish biomasses were thus higher in the most eutrophic sites, but they were

less stable over time: were this finding to be confirmed by further studies, this dual and opposite effect of

eutrophication would have important implications for both the conservation of fish populations, since

population stability is generally associated with lower extinction risk[28]. It is also noteworthy that temporal

fluctuations in total fish biomass were unrelated to fluctuations in *per capita* biomass (that is, to temporal

fluctuations in the average mass of individuals), reinforcing the hypothesis that biomass stability was rather

indirectly fostered by mechanisms such as functional complementarity or redundancy among

phenotypes[9,10,27], acting as a biological insurance against natural environmental fluctuations[7].

Contribution of intraspecific genetic diversity to biomass stability. The contribution of intraspecific

genetic diversity to the overall variance in biomass stability (R^2 = 21.1%) was much higher than that of

considered environmental determinants (R^2 = 4%; Fig. 4). 84 % of the total explained variance in fish

biomass stability was hence attributed to intraspecific genetic diversity, which suggests that intraspecific

genetic diversity is a substantial driver of biomass stability in this area, although other unmeasured variables

also likely sustain variation in biomass stability since a non-negligible part of this variation (~75%; Fig. 4)

remained unexplained by our model. Moreover, first-order interactions between intraspecific genetic

diversity and environmental variables were not retained in final models (Fig. 2; but see the specific case of

minnows in Supplementary Table 2), suggesting that the influence of intraspecific genetic diversity on total

biomass stability may be predictable across environmental gradients, a result which is yet to be generalized

to different taxa and ecosystems.

DISCUSSION

Capitalizing on long-term demographic surveys, we report a positive relationship between

intraspecific genetic diversity and temporal biomass stability in three freshwater fish species. This

relationship indicates a buffering effect of intraspecific genetic diversity; genetically-impoverished

populations being less efficient in maintaining stable biomass levels over time than genetically-diversified

populations[7]. By favoring higher functional complementarity among phenotypes, higher genetic diversity

likely allows populations to maintain an efficient exploitation of available resources in the face of natural

environmental fluctuations, ensuring a stable production of biomass[1,4,7]. Interestingly, this buffering effect of

intraspecific genetic diversity did not come with a performance-enhancing effect on biomass production[7]

(Fig. 2A): genetically-diversified populations did not show higher biomass levels compared to genetically-

impoverished populations, mean total biomass being mostly driven by *per capita* biomass, and indirectly by

the environment (Fig. 4). Nevertheless, our study provides one of the first non-experimental evidence that

"real world" genetic diversity can directly promote temporal stability in biomass of wild organisms, in line

with both theoretical expectations and experimental evidence[9,14].

Our study being based on empirical data, it is not surprising that a large amount of variance in mean

biomass and in biomass stability remained unexplained by our models (60 and 75% respectively; Fig. 2 and

4). Mean biomass and biomass stability were also probably driven by factors that we could not consider in

this study, such as interspecific interactions at the community-level [29] or, at the ecosystem level, autotroph

primary production [30] or terrestrial subsidies[31]. Nevertheless, intraspecific genetic diversity accounted for

more than 21% of the variance in biomass stability, a contribution much higher than that of considered

environmental predictors. Our findings not only indicate that the relationship between intraspecific genetic

diversity and biomass stability holds true in natural ecosystems, but also that this relationship can be

substantial compared to the effects of other undisputable determinants of biomass, as recently shown for

interspecific diversity[5]. While species richness can buffer natural fish biomass production against

environmental variations[20], we argue that the intraspecific facet of biodiversity may actually also contribute

to biomass stability in the wild [9,14].

A corollary to this finding is that the loss of intraspecific genetic diversity might undermine the

temporal stability in biomass production. We notably detected a significant negative relationship between

intraspecific genetic diversity and the probability that populations experienced a recent demographic

contraction (i.e., recent bottleneck, Fig. 2), which is consistent with the imprint of recent human activities on

contemporary levels of genetic diversity[23,24]. This finding must be considered with caution, since recent

bottleneck probabilities were estimated using a modest number of microsatellite markers, which can

generate bias in demographic inferences[32–35]. Specifically, the limited number of loci sampled in the

genome[32] and the departure from the assumed mutation model[33,34] or from mutation-drift equilibrium[35]

generally increase the probability of inferring false signals of bottlenecks, i.e., of detecting a population

decline in a truly stable population. This type of bias could have inflated the relationship we found between

bottleneck probability and patterns of intraspecific genetic diversity. However, Paz-Vinas et al.[36]

demonstrated using simulations that, in river systems, demographic inferences based on microsatellite

markers are rather robust to this bias, and actually more likely to detect false signals of expansion, i.e.,

detect a population expansion in a truly stable population, than to detect false signal of bottlenecks.

Furthermore, to somehow limit the potential biases associated with the use of microsatellite markers, past

bottleneck inferences were based on three independent methods that yielded congruent estimates

(Supplementary Note 2). Although the links between past-demographic events and contemporary patterns

of intraspecific genetic diversity would merit further confirmation based on genomic data[32], we believe that

it is reasonable to consider the low levels of intraspecific genetic diversity observed in some populations to

stem -at least in part- from recent, possibly human-induced, demographic contractions, for instance

triggered by water eutrophication (Fig. 2 and 3A). Although alternative and non-exclusive historical

processes (e.g., post glacial colonization events) are also likely to explain observed spatial patterns of

intraspecific genetic diversity[25,37], this suggests that contemporary evolutionary processes can modulate

ecological dynamics in natural settings. Since the loss of intraspecific genetic diversity always precede the

loss of species[8], genetic erosion may adversely affect key-ecological functions long before the first species

of a community becomes extirpated. We therefore argue that the loss of intraspecific diversity observed

worldwide[23,24] may actually be responsible for a considerable alteration of many ecological processes in

nature, but that these adverse effects might have been underestimated. With a loss of 6-15% in intraspecific

genetic diversity[23,24], we estimated a similar 8-10% reduction in biomass stability across the river basin (-

8.9%; 95% confidence interval: [-10.1; -7.7]; Supplementary Note 3). This reduction in biomass stability was

calculated using conservative estimates of intraspecific genetic diversity loss[23,24]; we therefore anticipate

that this reduction could be much greater in species with weak conservation statuses, and that this reduction

in stability may have important cascading effects on trophic networks, ecosystem functioning and, in some

countries, food provisioning[1,38].

Future studies are needed to confirm the significance of these results in other taxa and other

ecosystems and to disentangle the relative contribution of intra- and interspecific diversity in explaining

biomass production in the wild[10]. Nevertheless, our work suggests that the impact of genetic erosion on

natural ecosystems' capacity to provide critical provisioning and regulating services to humanity is probably

more important than anticipated. This makes human-induced genetic erosion a critical conservation issue

and stresses the need for human societies to adopt prominent environmental policies favoring all facets of

biodiversity[39].

METHODS

Sampling stations and biological models. We selected 47 river stations evenly scattered across two large River basins in South-Western France (the Garonne River basin and the Dordogne River basin) to reflect the environmental variability existing along the upstream–downstream gradients. Fish communities in these basins are generally poorly diverse (3 to 15 species[40]), and we focused on the three most common species: the minnow *Phoxinus dragarum*, the gudgeon *Gobio occitaniae* and the chub *Squalius cephalus*. These generalist cyprinids vary in their mean body length (minnows: 80–90 mm; gudgeons: 120–150 mm; chubs: 300–500 mm)[41]. They mainly feed on invertebrates (although chubs can also consume small-bodied fish) but occupy different habitats: chubs are primarily found in downstream sections at relatively low densities (~0.01 ind.m^{-2}), minnows are primarily found in upstream sections at relatively high densities (~0.10 ind.m^{-2}), whereas gudgeons are found all along the river basin in various habitats and at relatively high densities (~0.08 ind.m^{-2})[41]. All stations are monitored yearly by the French Office for Biodiversity (OFB) with a constant sampling effort since 1990[42]. Demographic and biomass data from the three focal species were extracted from the OFB databases[43]. We only retained stations monitored from 1993 to 2020 with at least eight sampling sessions, resulting in 42 stations (Fig. 1; mean number of sampling sessions = 15.6; mean survey duration = 21.2 years; mean number of focal species per station = 2.0). The minimum pairwise distance among sampling stations was 18.5km (between TARMil and CENSai). This distance is higher than the maximum travelled distance recorded in chubs (16 km)[44], here considered as the most mobile species, and all stations could thus be considered independent.

Biomass data. For each species, station and year of survey, we collected local fish density (number of individuals per m²), total fish biomass (in g.m^{-2}) and computed *per capita* biomass (or mean individual biomass, in g) as total fish biomass divided by local fish density. For each species, both total fish biomass and *per capita* biomass were standardized to make data comparable across species. For each station, we computed (i) Mean Total Biomass (respectively, *per capita* Mean Biomass) as the mean of total fish biomass (respectively, of *per capita* biomass) across species and over years, and to capture temporal fluctuations in biomass measures, (ii) Mean Total Biomass Stability (respectively, *per capita* Mean Biomass Stability) as the inverse of the squared coefficient of variation of Mean Total Biomass (respectively, of *per capita* Mean Biomass Stability) over years[45]. *Per capita* Biomass (stability) was here considered to determine whether

Total Biomass (stability) was directly driven by the average individual biomass (stability) alone, or by other

mechanisms such as functional complementarity among phenotypes. Each station was also assigned a

sampling weight (ranging from 0.29 to 0.98) computed as the average of the relative local survey duration

(compared to the maximal duration across stations) and of the relative local number of sampling sessions

(compared to maximal number of sampling sessions across stations).

Genetic data, intraspecific genetic diversity and bottleneck probability. The 42 retained stations were

sampled in 2011 and 2014 with up to 30 adults from each species caught by electric-fishing. On the field, a

small piece of pelvic fin was collected from each individual and was preserved in 70% ethanol, before

releasing fish in situ. The genetic diversity of each population was characterized using both microsatellite

(13 to 17 loci after filtering) and SNP markers (1244 to 1892 loci after filtering) as a composite metric

extracted from both markers (see Supplementary Note 4 for details on DNA extraction, SNP calling,

microsatellite genotyping and computation of metrics of genetic diversity). We used principal component

analysis (PCA) to get a synthetic predictor of the cross-species level of intraspecific genetic diversity at each

site. Only the first component was retained, accounting for 75.9 % of variance in genetic data, with

genetically impoverished sites on the one hand (negative coordinates) and genetically diversified sites on

the other hand (Supplementary Note 4). To determine the probability that populations suffered from a recent

bottleneck, we used microsatellite data to compute three indirect yet complementary metrics of demographic

contraction (using M-ratios[46], VarEff[47] and Migraine[48]), and used PCA to get a synthetic predictor of the

cross-species local degree of genetic erosion (bottleneck probability) experienced by fish in recent

generations. Only the first component was retained, accounting for 61.3 % of variance in input data, with

sites with no sign of bottleneck on the one hand (negative coordinates) and sites having probably

experienced a recent bottleneck on the other hand (Supplementary Note 2).

Environment data. We similarly used PCA to get synthetic predictors of environment characteristics

(distance to the river mouth, distance to the tributary source, water temperature and Water Quality Index)

at the site level (Supplementary Note 5). The two first components were retained, accounting for 80.5 % of

the total variance in environmental variables. The first component (55 % of variance) stood for the upstream-

downstream gradient, with fresh upstream sites on the one hand (negative coordinates) and warmer

downstream sites on the other hand. The second component (25.5 % of variance) stood for a eutrophic

gradient, with oligotrophic river sites on the one hand (negative coordinates, low Water Quality Index) and

nutrient-rich (mesotrophic) sites on the other hand (high Water quality Index).

Path analyses. To investigate how intraspecific genetic diversity might influence Biomass and/or Biomass

Stability while accounting for the effects of both environment and *per capita* Biomass[20,49], we used path

analyses[50,51]. We designed two full causal models describing the expected direct and indirect links among

variables and their first-order interactions (Supplementary Note 6). Our main focus was on the direct links

between intraspecific genetic diversity (intraspecific genetic diversity, or cross-product interactions of

intraspecific genetic diversity with environmental variables[52]) and Total Biomass on the one hand (model A)

and Biomass Stability on the other hand (model B). To consider the influence of *per capita* biomass on total

biomass variables, we hypothesized that intraspecific genetic diversity would also indirectly promote Total

Biomass (respectively, Biomass Stability), through a pathway involving *per capita* Biomass (respectively,

per capita Biomass Stability). We further hypothesized that the environmental characteristics of stations

(upstream-downstream gradient UDG, eutrophic gradient EG and the corresponding cross-product

interaction UDGxEG) would affect *per capita* and total biomass variables both directly (for instance through

higher intraspecific competition in harsh conditions) and indirectly, through pathways involving intraspecific

genetic diversity (promoted for instance by higher proximity to glacial refuges[25]) as well as bottleneck

(triggered for instance by pollutants). All variables were standardized to z-scores before using weighted path

analyses[22] with 'MLM' maximum likelihood estimation, 'Satorra-Bentler' scaled test statistic and station

sampling weights. We simplified each model by removing non-significant paths one at a time, provided that

cross-products were always associated with their additive terms[53] and that removal led to an increase in the

relative fit of the model (i.e., a decrease in BIC score[54]). The validity of final models was assessed according

to their absolute fit (standardized root mean square residual (SRMR) < 0.09 and Robust Comparative Fit

Index (CFI) > 0.96)[55]. Additionally, we verified that there was no significant discrepancy between the sample

and the fitted covariance matrices[55] (p-value associated with the model χ^2 statistic > 0.05). All path analyses

were run using the R-package *lavaan*[56]. Predicted values of endogenous variables (Fig. 3) were obtained

from linear regression models with each beta coefficient (and associated standard deviation) fixed to the

value of the corresponding direct link within the final model. 95% confidence envelops were obtained by

bootstrap (1000 iterations).

Variance partitioning. For each total biomass variable, we computed (a) the amount of variance (R^2)

explained by each model. To assess the relative contribution of *per capita* biomass, intraspecific genetic

diversity and environment to the variance in total biomass variables, we computing R^2 from further simplified

models with (b) all variables related to *per capita* biomass being discarded (amount of variance explained

by both environment and intraspecific genetic diversity), then (c) with all variables related to intraspecific

genetic diversity (intraspecific genetic diversity and associated cross-products) being discarded (amount of

variance explained by environment only). The relative contributions of *per capita* biomass and intraspecific

genetic diversity to the variance in total biomass variables were respectively obtained by subtracting R^2 of

(b) from R^2 of (a) and by subtracting R^2 of (c) from R^2 of (b).

ACKNOWLEDGMENTS

315 This work was financially supported by the Office Français pour la Biodiversité (OFB) and the Agence

316 Nationale pour la Recherche. We warmly thank all the colleagues and students who helped with field

317 sampling and data analyses. The authors declare no competing interests.

AUTHORS CONTRIBUTIONS

320 J.G.P., N.P. and S.B. formulated the main idea. J.G.P. collected environmental data and performed

321 genetic/genomic analyses. M.C. and N.P. collected and prepared biomass data. J.G.P. performed data-

322 analyses. J.G.P. and S.B. wrote the first draft and all authors provided feedback on subsequent versions of

323 the manuscript.

COMPETING INTERESTS

326 The authors declare no competing interests.

DATA ACCESSIBILITY

329 Raw data and data used to perform path analyses[57]: Figshare doi: 10.6084/m9.figshare.13095380.v8

CODE ACCESSIBILITY

Code used to perform path analyses and produce figures[57]: Figshare doi: 10.6084/m9.figshare.13095380.v8

REFERENCES

1. Cardinale, B. J. *et al.* Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012).

2. Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51–

57 (2011).

3. De Meester, L. *et al.* Analysing eco-evolutionary dynamics—The challenging complexity of

the real world. *Funct. Ecol.* **33**, 43–59 (2019).

4. Loreau, M. Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* **91**,

3–17 (2000).

5. Duffy, J. E., Godwin, C. M. & Cardinale, B. J. Biodiversity effects in the wild are common and

as strong as key drivers of productivity. *Nature* **549**, 261–264 (2017).

6. Hooper, D. U. *et al.* Effects of Biodiversity on Ecosystem Functioning: A Consensus of

Current Knowledge. *Ecol. Monogr.* **75**, 3–35 (2005).

7. Yachi, S. & Loreau, M. Biodiversity and ecosystem productivity in a fluctuating environment:

The insurance hypothesis. *Proc. Natl. Acad. Sci.* **96**, 1463–1468 (1999).

8. Spielman, D., Brook, B. W. & Frankham, R. Most species are not driven to extinction before

genetic factors impact them. *Proc. Natl. Acad. Sci.* **101**, 15261–15264 (2004).

9. Hughes, A. R., Inouye, B. D., Johnson, M. T. J., Underwood, N. & Vellend, M. Ecological

consequences of genetic diversity. *Ecol. Lett.* **11**, 609–623 (2008).

10. Raffard, A., Santoul, F., Cucherousset, J. & Blanchet, S. The community and ecosystem

consequences of intraspecific diversity: a meta-analysis: The ecological effects of

intraspecific diversity. *Biol. Rev.* **94**, 648–661 (2019).

11. Olson, D. M. *et al.* Terrestrial Ecoregions of the World: A New Map of Life on Earth.

BioScience **51**, 933 (2001).

12. Reusch, T. B. H., Ehlers, A., Hammerli, A. & Worm, B. Ecosystem recovery after climatic

extremes enhanced by genotypic diversity. *Proc. Natl. Acad. Sci.* **102**, 2826–2831 (2005).

13. Vellend, M. & Geber, M. A. Connections between species diversity and genetic diversity.

Ecol. Lett. **8**, 767–781 (2005).

14. Forsman, A. & Wennersten, L. Inter-individual variation promotes ecological success of

populations and species: evidence from experimental and comparative studies. *Ecography*

39, 630–648 (2016).

15. Hendry, A. P. A critique for eco-evolutionary dynamics. *Funct. Ecol.* **33**, 84–94 (2019).

16. Petit, R. J. Glacial Refugia: Hotspots But Not Melting Pots of Genetic Diversity. *Science* **300**,

1563–1565 (2003).

17. Bestion, E. *et al.* Altered trophic interactions in warming climates: consequences for

predator diet breadth and fitness. *Proc. R. Soc. B Biol. Sci.* **286**, 20192227 (2019).

18. Jangjoo, M., Matter, S. F., Roland, J. & Keyghobadi, N. Connectivity rescues genetic diversity

after a demographic bottleneck in a butterfly population network. *Proc. Natl. Acad. Sci.* **113**,

10914–10919 (2016).

19. Coltman, D. W. *et al.* Undesirable evolutionary consequences of trophy hunting. *Nature* **426**,

655–658 (2003).

20. Duffy, J. E., Lefcheck, J. S., Stuart-Smith, R. D., Navarrete, S. A. & Edgar, G. J. Biodiversity

enhances reef fish biomass and resistance to climate change. *Proc. Natl. Acad. Sci.* **113**,

6230–6235 (2016).

21. Donohue, I. *et al.* On the dimensionality of ecological stability. *Ecol. Lett.* **16**, 421–429

(2013).

22. Grace, J. B. *Structural equation modelling and natural systems.* (Cambridge University Press,

2006).

23. Leigh, D. M., Hendry, A. P., Vázquez-Domínguez, E. & Friesen, V. L. Estimated six per cent

loss of genetic variation in wild populations since the industrial revolution. *Evol. Appl.* **12**,

1505–1512 (2019).

24. Exposito-Alonso, M. *et al.* Genetic diversity loss in the Anthropocene. *Science* 1431–1435

(2022).

25. Paz-Vinas, I., Loot, G., Stevens, V. M. & Blanchet, S. Evolutionary processes driving spatial

patterns of intraspecific genetic diversity in river ecosystems. *Mol. Ecol.* **24**, 4586–4604

(2015).

26. Hudon, C. *et al.* Oligotrophication from wetland epuration alters the riverine trophic

network and carrying capacity for fish. *Aquat. Sci.* **74**, 495–511 (2012).

27. Hughes, A. R. & Stachowicz, J. J. Genetic diversity enhances the resistance of a seagrass

ecosystem to disturbance. *Proc. Natl. Acad. Sci.* **101**, 8998–9002 (2004).

28. Finn, D. S., Bogan, M. T. & Lytle, D. A. Demographic Stability Metrics for Conservation

Prioritization of Isolated Populations. *Conserv. Biol.* **23**, 1185–1194 (2009).

29. Raffard, A., Santoul, F., Cucherousset, J. & Blanchet, S. The community and ecosystem

consequences of intraspecific diversity: a meta-analysis: The ecological effects of

intraspecific diversity. *Biol. Rev.* **doi: 10.1111/brv.12472**, (2019).

30. Delong, M. D. & Thorp, J. H. Significance of instream autotrophs in trophic dynamics of the

Upper Mississippi River. *Oecologia* **147**, 76–85 (2006).

31. Nakano, S. & Murakami, M. Reciprocal subsidies: Dynamic interdependence between

terrestrial and aquatic food webs. *Proc. Natl. Acad. Sci.* **98**, 166–170 (2001).

32. Hoban, S. M., Gaggiotti, O. E. & Bertorelle, G. The number of markers and samples needed

for detecting bottlenecks under realistic scenarios, with and without recovery: a simulation-

based study. *Mol. Ecol.* **22**, 3444–3450 (2013).

33. Peery, M. Z. *et al.* Reliability of genetic bottleneck tests for detecting recent population

declines. *Mol. Ecol.* **21**, 3403–3418 (2012).

34. Putman, A. I. & Carbone, I. Challenges in analysis and interpretation of microsatellite data

for population genetic studies. *Ecol. Evol.* 4399–4428 (2014) doi:10.1002/ece3.1305.

35. Chikhi, L., Sousa, V. C., Luisi, P., Goossens, B. & Beaumont, M. A. The confounding effects of

population structure, genetic diversity and the sampling scheme on the detection and

quantification of population size changes. *Genetics* **186**, 983–995 (2010).

36. Paz-Vinas, I., Quéméré, E., Chikhi, L., Loot, G. & Blanchet, S. The demographic history of

populations experiencing asymmetric gene flow: combining simulated and empirical data.

Mol. Ecol. **22**, 3279–3291 (2013).

37. De Kort, H. *et al.* Life history, climate and biogeography interactively affect worldwide

genetic diversity of plant and animal populations. *Nat. Commun.* **12**, 516 (2021).

38. Duffy, J. E., Lefcheck, J. S., Stuart-Smith, R. D., Navarrete, S. A. & Edgar, G. J. Biodiversity

enhances reef fish biomass and resistance to climate change. *Proc. Natl. Acad. Sci.* **113**,

6230–6235 (2016).

39. Laikre, L. *et al.* Post-2020 goals overlook genetic diversity. *Science* **367**, 1083–1085 (2020).

40. Blanchet, S., Helmus, M. R., Brosse, S. & Grenouillet, G. Regional *vs* local drivers of

phylogenetic and species diversity in stream fish communities. *Freshw. Biol.* **59**, 450–462

(2014).

41. Keith, P., Persat, H., Feunteun, E., Adam, B. & Geniez, M. *Les Poissons d'eau douce de*

France. (2011).

42. Poulet, N., Beaulaton, L. & Dembski, S. Time trends in fish populations in metropolitan

France: insights from national monitoring data. *J. Fish Biol.* **79**, 1436–1452 (2011).

43. Irz, P. *et al.* A long-term monitoring database on fish and crayfish species in French rivers.

Knowl. Manag. Aquat. Ecosyst. 25 (2022) doi:10.1051/kmae/2022021.

44. Fredrich, F., Ohmann, S., Curio, B. & Kirschbaum, F. Spawning migrations of the chub in the

River Spree, Germany. *J. Fish Biol.* **63**, 710–723 (2003).

45. Haegeman, B. *et al. Resilience, invariability, and ecological stability across levels of*

organization. http://biorxiv.org/lookup/doi/10.1101/085852 (2016) doi:10.1101/085852.

46. Garza, J. C. & Williamson, E. G. Detection of reduction in population size using data from

microsatellite loci. *Mol. Ecol.* **10**, 305–318 (2001).

47. Nikolic, N. & Chevalet, C. Detecting past changes of effective population size. *Evol. Appl.* **7**,

663–681 (2014).

48. Leblois, R. *et al.* Maximum-Likelihood Inference of Population Size Contractions from

Microsatellite Data. *Mol. Biol. Evol.* **31**, 2805–2823 (2014).

49. Oehri, J., Schmid, B., Schaepman-Strub, G. & Niklaus, P. A. Biodiversity promotes primary

productivity and growing season lengthening at the landscape scale. *Proc. Natl. Acad. Sci.*

114, 10160–10165 (2017).

50. Lefcheck, J. S. piecewiseSEM: Piecewise structural equation modeling in R for ecology,

evolution, and systematics. *Methods Ecol. Evol.* **7**, 573–579 (2015).

51. Fourtune, L. *et al.* Inferring Causalities in Landscape Genetics: An Extension of Wright's

Causal Modeling to Distance Matrices. *Am. Nat.* **191**, 491–508 (2018).

52. Aiken, L. S., West, S. G. & Reno, R. R. *Multiple regression: Testing and interpreting*

interactions. (SAGE Publications, 1991).

53. Batista-Foguet, J. M., Coenders, G., Saris, W. E. & Bisbe, J. Simultaneous Estimation of

Indirect and Interaction Effects using Structural Equation Models. *Metodološki Zv.* **1**, 163–

184 (2004).

54. Burnham, K. P. & Anderson, D. R. *Model selection and multimodel inference: a practical*

information-theoretic approach. (Springer, 2002).

55. Hu, L. & Bentler, P. M. Cutoff criteria for fit indexes in covariance structure analysis:

Conventional criteria versus new alternatives. *Struct. Equ. Model. Multidiscip. J.* **6**, 1–55

(1999).

56. Rosseel, Y. **lavaan** : An *R* Package for Structural Equation Modeling. *J. Stat. Softw.* **48**, (2012).

57. Prunier, J. G. *et al.* Data for 'Genetic erosion reduces biomass temporal stability in wild fish

populations.' (2022) doi:10.6084/m9.figshare.13095380.v8.

Figure 1. Study area and localization of sampled river stations and species. Geographic situation of

the Garonne-Dordogne River basin in South-Western France and localization of the 42 unique river stations,

with pie charts indicating species (co-)occurrence within each station. The black arrow indicates the location

of the river mouth. Background is a shaded relief map.

Figure 2. Final causal graphs. a Causal graph depicting the retained links among environmental (purple),

bottleneck (blue), genetic (light green), and mean biomass variables (per capita and total; yellow). **b** Causal

graph depicting the retained links among environmental (purple), bottleneck (blue), genetic (light green),

and biomass stability variables (per capita and total; yellow). In each panel, retained first-order interactions

are represented by rounded rectangles. Blue and red arrows represent negative and positive significant

paths, respectively, with the width of arrows proportional to the absolute value of the corresponding path

coefficient. Also provided is the amount of variance (R^2) explained in each endogenous variable. Non-

significant paths are not represented, for the sake of clarity (see Supplementary Table 1 for detailed results).

Figure 3. **Predicted values and 95% confidence intervals of all endogenous variables given the**

retained links shown in Fig.1. When two predictors were considered (panels B, C and F), predictions were

performed using the full range of the first predictor (x-axis) and the minimal (purple), mean (blue) and

maximal (green) values of the second predictor.

Figure 4. Variance partitioning. Contributions of environment variables (purple), *per capita* biomass

variables (yellow) and intraspecific genetic diversity (light green) to the variance (R2) in Total Biomass (R^2

= 0.398) and Total Biomass stability (R^2 = 0.251). The contributions to the explained variance (in %) are

indicated into brackets.