Loss of genetic diversity in wild fish populations reduces biomass stability over time.

Authors:

Jérôme G. Prunier [1] *, Mathieu Chevalier [2], Allan Raffard [1], Géraldine Loot [3], Nicolas Poulet [4], Simon Blanchet[1,3]*

Affiliations :

[1] Centre National de la Recherche Scientifique (CNRS), Université Paul Sabatier (UPS); Station d'Ecologie Théorique et Expérimentale, UMR 5321, F-09200 Moulis, France

[2] Department of Ecology and Evolution, University of Lausanne, Biophore, CH-1015 Lausanne, Switzerland

[3] CNRS, UPS, École Nationale de Formation Agronomique (ENFA) ; UMR 5174 EDB (Laboratoire Évolution & Diversité Biologique), 118 route de Narbonne, F-31062 Toulouse cedex 4, France

[4] Pôle écohydraulique AFB-IMT, allée du Pr Camille Soula, 31400 Toulouse, France

* Corresponding authors:

Jérôme G. Prunier and Simon Blanchet

Station d'Ecologie Théorique et Expérimentale, UMR 5321, F-09200 Moulis, France

2 route du CNRS, 09200 Moulis

Phone: (+33)561040361

Email: jerome.prunier@gmail.com, simon.blanchet@sete.cnrs.fr

ORCIDs: Jérôme G. Prunier 0000-0003-4110-2567, Simon Blanchet 0000-0002-3843-589X

Keywords

Eco-evolutionary dynamics; Population performance; Intraspecific diversity, Biodiversity-Ecosystem function.

Abstract

Biodiversity sustains key ecosystem functions, and ongoing species losses have tremendous impacts on

good and services provided by biodiversity. Before species are extirpated, they yet lose intraspecific

genetic diversity. The loss of intraspecific diversity has implications for the evolutionary potential of

species, but may also alter ecological and ecosystem functions such as biomass productivity. However,

there have been little attempts to empirically test whether intraspecific genetic diversity actually promotes

productivity and stability in wild populations, and, if so, to quantify its relative importance compared to

other determinants. Capitalizing on 20-year demographic surveys in wild fish meta-populations, we

demonstrated that intraspecific genetic diversity did not increase mean biomass production in local

populations, but strongly and consistently stabilized biomass production over time. Overall, genetic

diversity accounted for about 20% of explained variance in biomass stability across species, a significant

contribution that was about half that of environment and demography (40% each). Populations having

suffered from past demographic bottlenecks harbored lower levels of genetic diversity and showed less

stability in biomass production over the last 20 years. Our study demonstrates that the loss of genetic

diversity can destabilize biomass productivity in natural vertebrate populations in just a few generations,

strengthening the importance for human societies to adopt prominent environmental policies to favor all

facets of biodiversity.

Introduction

There is now unequivocal evidence that biodiversity sustains critical ecosystem services that benefit directly to humanity, such as water filtering, pollination and biomass production (1–3). By promoting higher complementarity among species (4, 5), interspecific diversity allows ecological communities to optimally capture essential resources, transform those resources into biomass and recycle them. In species-rich communities, these ecological processes are maintained even in face of environmental variations, thus promoting ecosystem productivity and stability over time (6–10). The drastic erosion in species diversity observed worldwide is hence prejudicial, as it is expected to deeply impair natural ecosystems' capacity to durably sustain human health and wellbeing (2, 3, 11). However, the loss of species is only the tip of the iceberg. In most taxa, another form of biodiversity erosion is indeed silently underway: the loss of intraspecific genetic diversity (12, 13).

Beyond its positive influence on individual fitness and evolutionary rescue (14), intraspecific genetic diversity is expected to play a role similar to species diversity in driving key ecological functions such as biomass production (1). Assuming that genetic variation promotes intraspecific phenotypic variation and thus complementarity among individuals (15), genetically-diversified populations are predicted to have higher and more stable biomass yields than genetically impoverished populations, notably under stressful environmental conditions (16, 17). This direct relationship between genetic diversity and biomass production is expected to be particularly strong in ecosystems where species diversity is naturally low, which is actually the norm in many temperate ecosystems (18). Indeed, the functioning of such ecosystems probably depends more on complementarity among genotypes than on complementarity among species (19–21), emphasizing the importance of maintaining genetic diversity to preserve ecosystem functions and services (16).

However, most studies investigating the relationship between genetic diversity and ecological functions such as biomass production are based on experimental or semi-experimental designs, where population densities and levels of genetic diversity are manipulated, and environmental conditions are controlled and maintained constant over time (22). Contrary to field surveys, and although they offer a number of advantages (16, 22), experiments do not allow spanning large spatial and temporal scales or to investigate the influence of historical contingencies. Local levels of genetic diversity indeed result from the

interplay between long-term evolutionary trajectories (e.g., localization and size of glacial refugia (23)) and more recent (if not ongoing) ecological processes affecting individual life history traits (e.g., stressful environmental conditions (24)) or population demography (e.g., bottleneck events (25)). This natural complexity cannot be fully grasped from experimental studies (26). Observational field surveys are on the contrary more realistic and may provide important insights into the contribution of genetic diversity, and the loss of it, to biomass production and stability in natural settings, in accordance or in disagreement with theoretical expectations and experimental findings (16, 27, 28). They yet raise several difficulties. First, assessing the influence of genetic diversity on the stability of biomass production over several generations or over several seasonal cycles implies long-term monitoring programs of both population density and biomass production, but such data are still scarce or difficult to collect in many situations. Furthermore, genetic diversity may be correlated with variables that are also involved in biomass production, such as population density and environmental heterogeneity, making it difficult to disentangle their respective contributions (16). This last issue may however be partly alleviated through the use of causal modeling procedures (29), making it possible to thoroughly confront theoretical expectations and experimental findings with the "real world" (30, 31).

Capitalizing on long-term demographic surveys (up to 22 years) conducted in 47 rivers from a large watershed (the Garonne-Dordogne water basin, South-Western France; Figure 1), we assessed the relationship between genetic diversity and biomass yields (mean biomass production and temporal stability in biomass production) at the population level in three parapatric freshwater fish species. To that aim, we designed a causal model depicting all the expected causal links between biomass variables, genetic diversity and several environmental and demographic predictors. Rivers are characterized by strong environmental gradients that we synthetized in the form of an upstream-downstream gradient and a eutrophic gradient. The former is a critical feature known to shape both spatial patterns of genetic diversity and biomass yields in dendritic networks (32, 33) whereas the latter may act as a possible stressor and affect both population dynamics and productivity (34). As local population dynamics may also influence both genetic diversity and biomass yields, we considered several demographic covariates: mean population density, long-term stability in population density as well as bottleneck probability, the latter indicating past changes in population size (35). By investigating this causal model, we answered the

following questions: do the positive relationships between genetic variation and biomass production and

stability hold in natural settings? If any, is the contribution of genetic variation to biomass

production/stability similar across species, and comparable in magnitude to that of other environmental

and demographic determinants? Finally, and most importantly, is it possible to detect the impact of the

ongoing genetic erosion, e.g., the loss of genetic diversity in response to a recent reduction in population

size, on biomass production and stability? This latter point remains controversial but of high concern: with

an estimated 6% loss of intraspecific genetic diversity in wild organisms since the industrial revolution

(36), the impact of human-induced genetic erosion on natural ecosystems' capacity to provide critical

provisioning and regulating services to humanity may actually be much more important than anticipated,

making it a critical conservation issue.

Materials and Methods

Study area and biological models

The study was carried out in the Garonne-Dordogne river basin (South-Western France; Figure 1). We

selected 47 river sections evenly scattered across the whole basin to reflect the environmental variability

existing along the upstream–downstream gradient. All these sites are yearly monitored by the French

Agency for Biodiversity (AFB) since 1989 (50). Fish communities in this river basin are generally poorly

diverse (3 to 15 species in general) and we focused on three of the most common species (51): the

Eurasian minnow *Phoxinus phoxinus*, the Languedoc gudgeon *Gobio occitaniae* and the chub *Squalius*

cephalus. These cypriniform fish species vary in their mean body length (minnows: 80–90 mm; gudgeons:

120–150 mm; chubs: 300–500 mm; (52)). They are all generalist species mainly feeding on invertebrates

(although chubs can also predate on small-bodied fish) but occupy different habitats (51): chubs are

primarily found in downstream sections at relatively low densities (~0.01 ind.m-2), minnows are primarily

found in upstream sections at relatively high densities (~0.10 ind.m-2), whereas gudgeons are found all

along the river basin in various habitats and at relatively high densities (~0.08 ind.m-2; (42, 52)).

Demographic and biomass data

Demographic and biomass data were extracted from the AFB databases (53). We only retained sites

monitored for 10 to 22 years (mean = 15.02) with a constant sampling effort. For each species, site and

year of survey, we collected the local fish density (number of individuals per m²) and the local fish

biomass, expressed in g.m-2. For each species and each site, we computed the local population density

and the local biomass production as the mean of density and biomass values over years. Metrics of

density stability and biomass stability were both computed as the inverse of the squared coefficient of

variation of density (respectively, biomass) over years, following (54). All variables related to density

(population density and density stability) and biomass (biomass production and biomass stability) were

log-transformed before analyses to meet normality assumptions.

Genetic data

The 47 river stations were sampled once in 2014 with up to 30 adults from each species caught by

electrofishing, resulting in a set of 35, 37 and 21 sampled populations in minnows, gudgeons and chubs,

respectively. On the field, a small piece of pelvic fin was collected from each individual and was preserved

in 70% ethanol, before releasing fish in situ. Genomic DNA was extracted using a salt-extraction protocol

(55) and used to obtain, for each species, population-based allelic frequencies following a paired-end,

pool-seq procedure (56), as detailed in Appendix S3a. Read assembly, SNP calling and filtering (see

Appendix S3b for details) generated a total of 1244 SNPs in minnows, 1892 SNPs in gudgeons and 1847

SNPs in chubs. Genomic data were further completed with microsatellite genotypes from individuals

sampled in 2011 (Appendix S2), with 17, 13 and 16 loci in minnows, gudgeons and chubs, respectively.

For each species and site, we computed a total of six metrics of genetic diversity. First, we used SNPs

allelic frequencies to compute two metrics in R (57): the expected level of heterozygosity across SNPs loci

(sHe) and the observed level of SNP polymorphism (sPo), computed as the number of non-fixed loci (0 <

allelic frequency < 1) divided by the total number of loci with non-missing data in a given population. We

then used microsatellite data to compute four additional metrics using the softwares GENETIX 4.3 (58)

and ADZE 1.0 (59): the expected (μHe) and observed (μHo) levels of heterozygosity across microsatellite

loci, the mean number of alleles per locus (μNam) and the standardized allelic richness (μAR) based on a

standardized sample size of 20.

To capture overall trends in species-specific patterns of genetic diversity and smooth possible

particularities of each metric (60), the six metrics of genetic diversity were compiled using a principal

component analysis (PCA). Only the first component was retained in each species, respectively

accounting for 75.7, 68.8 and 86 % of variance in genetic data in minnows, gudgeons and chubs,

respectively. In general, all variables equally contributed to the retained components, which were thus

considered as species-specific synthetic predictors for genetic diversity, with genetically impoverished

populations on the one hand (negative coordinates) and genetically diversified populations on the other

hand (positive coordinates; Appendix S7a).

As genetic data could not be collected over years, we also computed the M-ratio (35) to determine

whether sampled populations experienced recent demographic bottlenecks. The M-ratio is the ratio

between the number of observed alleles at a microsatellite locus and the allelic range of that locus, the

latter being supposed to decrease slower than the number of alleles during a demographic collapse. This

index ranges from 0 to 1: empirical M-ratio values lower than a critical value of 0.68 indicate significant

bottleneck (35). For each populations and each species, the M-ratio was computed for each microsatellite

locus and then averaged over loci following (61; Appendix S1). We used the one minus the M-ratio as a

proxy for demographic bottleneck probability.

Environmental data

Each site was described using 17 environmental variables susceptible to affect demography, biomass

yields and genetic diversity of fish populations. From the French Theoretical Hydrological Network (62),

we extracted seven variables related to river topography: distance from the mouth (in m), distance from

the source (in m), river width (in m), slope (in degrees), altitude (in m) and mean annual water flow (in

$m^3.s^{-1}$). Additionally, we computed the betweenness centrality index as a measure of local connectivity

along the dendritic network (ComplexNetGIS toolbox in ARCGIS; (63)). Nine other variables were related

to water quality and were obtained from the database of the Water Information System of the Adour

Garonne basin (http://adour-garonne.eaufrance.fr): Temperature (in °C), dissolved oxygen concentration

(in mg.L-1) and oxygen saturation (in %), as well as the concentrations (in mg.L-1) in nitrogen compounds

(total nitrogen N, ammonium NH4+, nitrates NO3- and nitrites NO2-) and in phosphorus compounds (total

phosphorus P and phosphate PO43-). Only data from years 2012 to 2015 and from June to August were

selected and averaged over months and years. Older data, although incomplete, suggested that water

quality remained relatively stable over the considered period (Appendix S4). Except for a few sites that

might be subject to local water pollution resulting from phosphorus inputs, all sites showed good to very

good water quality according to the French implementation of the European Water Framework Directive

2000/60/EC (Appendix S5). Additionally, we computed a quantitative index of anthropogenic river

alteration (ANTH) using a database from the AFB (64), which notably describes observed hydrological,

morphological and thermic alterations resulting from human activities such as damming, channelization or

urbanization (see Appendix S6 for details and computation). This index theoretically ranges from 0 (no

identified anthropogenic alteration) to 1 (maximum alteration) but did not exceed 0.525 in our dataset, with

a mean of 0.275. Environmental variables were log-transformed to meet normality assumption when

required.

The 17 environmental variables were synthetized using a PCA (Figure 5; Appendix S7b). The two first

components were retained, accounting for 58.5 % of the total variance in environmental variables. The

first component (37.2% of variance) stood for the upstream-downstream gradient, with altitude and slope

on the one hand (upstream stations with negative coordinates) and distance from the source,

betweenness, temperature, river width and flow regime on the other hand (downstream stations with

positive coordinates). The second component (21.3% of variance) stood for an eutrophic gradient, with

dissolved oxygen, oxygen saturation and distance from the river mouth on the one hand (nutrient-

impoverished and highly oxygenated river stations with negative coordinates) and total phosphorus, total

nitrogen, nitrates and nitrites on the other hand (nutrient-rich and poorly oxygenated stations with positive

coordinates). Ammonium and phosphate concentrations, as well as anthropogenic river alteration, did not

contribute substantially to any of the retained components.

Statistical analyses

For each species, genetic diversity and all variables related to environment (upstream-downstream

gradient and eutrophic gradient), demography (population density, density stability and bottleneck

probability) and biomass yields (biomass production and biomass stability) were standardized to z-scores.

Our objective was to investigate how genetic diversity might influence biomass production and stability in

each species, while accounting for the direct and indirect influence of demography and environment (31,

65). To that aim, we designed a full causal model describing the expected direct and indirect links among

all variables (Appendix S8). We hypothesized that genetic diversity would directly promote biomass

production and biomass stability, either in an additive way, or in interaction with environmental variables.

Interactions were simply computed as the cross-product of genetic diversity with each environmental

variable (66). All other paths were used to control for the possible direct and indirect influence of other

variables. We notably hypothesized that population density and density stability would respectively affect

biomass production and biomass stability, both directly (biomass varying in proportion to density) and

indirectly, through pathways involving genetic diversity (low densities or low stability favoring genetic drift

(67)). Density stability was also expected to indirectly affect genetic diversity through demographic

bottlenecks: in the event of recent demographic collapse (in the 20 years preceding genetic sampling),

density stability should correlate negatively with the bottleneck probability, which is in turn expected to

decrease genetic diversity. Finally, we hypothesized that the environmental characteristics of river stations

(upstream-downstream gradient, eutrophic gradient and the corresponding first-order interaction

computed as their cross-product) would affect variations in biomass yields both directly (for instance

through higher intraspecific competition in harsh conditions) and indirectly, through pathways involving

genetic diversity (promoted for instance by higher proximity to glacial refuges or higher local connectivity;

(32)) and/or population dynamics (eutrophic conditions for instance leading to an increase in fish mortality

or oligotrophic conditions leading to an increase in competition for resources).

We then used a causal modeling procedure to infer actual causal relationships among variables.

Specifically, we used path analyses (29) with maximum likelihood estimation and 'Huber-White' robust

standard errors to assess, in each species, the validity of the full model and to simplify it when needed.

Model simplification was performed by removing non-significant paths one at a time, provided that cross-

products were always associated with their additive terms (68) and that removal led to an increase in the

relative fit of the model (i.e., a decrease in AIC score (69)). Simplification was stopped as soon as the

model reached a satisfactory absolute fit. Following Hu and Bentler (70), a model absolute fit was

considered as satisfactory when the standardized root mean square residual (SRMR) and the

Comparative Fit Index (CFI) were lower than ≤ 0.09 and higher than ≥ 0.96, respectively. Additionally, we

checked that the p-value associated with the model χ^2 statistic, a classical fit index measuring the

magnitude of discrepancy between the sample and the fitted covariance matrices (70), was non-significant

($p > 0.05$). To avoid model overfitting, we proceeded in two stages. In each species, we first considered a

subset of the full causal model excluding biomass variables and all related links, and performed model

simplification to get estimates for all important direct and indirect relationships among environmental

gradients, demographic variations and genetic diversity. Retained path coefficients, set to their mean

estimate, were subsequently considered as fixed parameters within the full causal model and a second set

of model simplification was performed to get estimates for all important direct and indirect relationships

pointing towards biomass variables. All path analyses were run using the R function lavaan (package

lavaan; (71)).

For each response variable, we (a) computed the amount of variance (R^2) explained by the retained

model in each species. To assess the relative contribution of genetic diversity, demography and

environment to the variance in biomass variables, we computing R^2 from further simplified models with (b)

all variables related to density being discarded (amount of variance explained by both environment and

genetic diversity), then (c) with all variables related to genetic diversity (genetic diversity and associated

cross-products) being discarded (amount of variance explained by environment only). The relative

contributions of demography and genetic diversity to the variance in biomass variables were respectively

obtained by subtracting R^2 of (b) from R^2 of (a) and by subtracting R^2 of (c) from R^2 of (b). We finally

averaged these contributions across species and build 100% stacked barplots to illustrate the overall

contribution of genetic diversity, demography and environment to the total explained variance in each

biomass variable.

Results

Contrary to our predictions, we did not find evidence that genetic variation was related to mean biomass production at the local scale (Figure 2 and Figure 3A). However, we demonstrated that, in the three considered species, local populations with higher levels of genetic diversity displayed less variability in biomass production over time than genetically impoverished populations, sustaining the idea that intraspecific genetic diversity stabilizes biomass production over time. In the chub *Squalius cephalus* (Figure 2A), biomass stability increased with genetic diversity, particularly in the most eutrophic (i.e. stressful) habitats (Figure 4A). In the Languedoc gudgeon *Gobio occitaniae* (Figure 2B), biomass stability similarly increased with increasing levels of genetic diversity and in the most upstream habitats, but decreased with eutrophication (Figure 4B). In the minnow *Phoxinus phoxinus* (Figure 2C), the retained interaction between the upstream-downstream gradient and genetic diversity actually stemmed from an uneven distribution of populations within the parameter space, with no genetically impoverished populations in the downstream part of the basin (Figure 4C). As a result, biomass stability actually increased both upstreamward and with increasing levels of genetic diversity, as observed in the Languedoc gudgeon. These striking findings were supported by robust simplified causal models that correctly represented observed data in each species (satisfactory absolute fit indices after model simplification; Tables 1 and 2; see Appendix S9 for detailed results).

The relative contribution of genetic diversity to the overall explained variance in biomass stability across species was of 21.7%, a non-negligible value, although about twice lower than that of environmentally- and demographically-related variables (37.7% and 40.6% respectively; Figure 3B). These contributions were yet highly species-dependent. Genetic diversity accounted for 23% of the total variance in biomass stability in chubs, a contribution twice higher than that of demography (11.3%) and much higher than that of environment (2.6%). In gudgeons, the contribution of genetic diversity was comparable to that of demography (11.2%) but lower than that of the environment (50.1%). It was the lowest in minnows (5%), environment and demography accounting for 15.2% and 49.5% of variance in biomass stability, respectively.

These idiosyncrasies in the contribution of intraspecific diversity to biomass stability may be related to the evolutionary history of each species within the basin, as indicated by the systematic negative

relationships between bottleneck probability and genetic diversity in all three species (Figure 2). In

minnows, the species with the lowest contribution, monitored populations showed high levels of genetic

variation (e.g., 5.9 ± 0.8 alleles per microsatellite locus on average) and only 5.7% of them experienced a

significant bottleneck (Fig. 3C-D). On the contrary, low levels of genetic variation (3.6 ± 0.6 alleles per

locus) and a high proportion of populations having experienced a significant bottleneck (up to 85.7%)

indicate that chubs, the species with the highest genetic contribution to biomass stability, suffered from

strong genetic erosion in the past. Finally, only in gudgeon populations did we find a significant negative

link between population stability (as measured over about 20 years) and bottleneck probability (Figure

2B). Together with a moderate proportion of populations having experienced a significant bottleneck

(29.7%) and high levels of genetic diversity (as high as in minnows; 5.6 ± 0.7 alleles per locus), these

results reveal signs of incipient genetic erosion in gudgeons.

In addition to revealing the consistent contribution of genetic diversity, or the loss of it, to biomass

stability, our modeling framework allowed identifying other recurrent and theoretically-grounded

relationships between variables. As expected, biomass yields were always positively linked to their

demographic counterparts (i.e., biomass production and biomass stability were positively related to

population density and population stability, respectively; Figure 2), explaining from 45.7% to 85.5 % of the

variance in biomass production (Figure 3A) and from 11.4% to 49.5% of variance in biomass stability

(Figure 3B). Similarly, we found a systematic downstream increase in genetic diversity, although this

pattern only held true in oligotrophic stations in the case of minnows (Figure 4D). In gudgeons, genetic

diversity was also positively influenced by the eutrophic gradient. We finally identified several significant

paths linking environment variables to population dynamics and biomass yields. For instance, population

density was found to increase downstreamward in chubs but upstreamward in gudgeons. Density stability

and biomass production were found to decrease downstreamward and biomass stability to decrease with

eutrophication in both minnows and gudgeons (Figure 2). All these expected findings confirm that causal

models were biologically sound and robust for all species.

Discussion

By promoting phenotypic diversity (15), intraspecific genetic diversity is expected to favor niche complementarity and facilitative interactions among individuals. It may therefore reduce intra- but also interspecific competition and allow a more efficient exploitation of available resources (10, 16, 28). This positive influence of genetic diversity on ecological properties of populations, but also of communities and ecosystems (37), is expected to be particularly important in harsh or unpredictable environmental conditions (22). Many experimental studies demonstrated such benefits of genetic diversity under controlled or semi-controlled conditions (see (28) for a review) , but observational studies conducted in natural settings are still scarce and only concern plants (26, 38). Capitalizing on long-term surveys, we here report systematic positive relationships between intraspecific genetic diversity and temporal stability in the biomass production of wild populations of freshwater fish species. These findings provide the first evidence that the relationship between intraspecific genetic diversity and biomass yields holds true in natural ecosystems.

Overall, intraspecific genetic diversity accounted for more than 20% of the explained variance in biomass stability across species. Although this contribution was about twice lower than that of other predictors, it indicates that intraspecific diversity is actually a substantial driver of stability in fish biomass production, somehow as important as other determinants such as environment and demography. It thus plays a role similar to that of interspecific diversity: species richness has been shown to buffer natural fish biomass production from environmental variations (31) but our findings suggest that both facets of biodiversity may actually contribute to biomass stability in the wild (16, 22). Interestingly, we did not identify any relationship between genetic diversity and mean biomass production over years, although our models explained about 60% of variance in chubs, and up to 95% in gudgeons and minnows: biomass production was here mainly explained by demography and, particularly in gudgeons, by environmental variation. With only biomass stability being identified as driven by intraspecific genetic diversity, our study highlights the usefulness of long-term field surveys when investigating the determinants of ecological dynamics in natural settings (39). The contribution of genetic diversity to the variance in biomass stability was yet highly species-dependent, which seems to be partly explained by the evolutionary trajectory of each species, and notably by the severity of past genetic erosion. The genetic contribution was indeed the

highest for the species that suffered from the strongest and widespread past genetic erosion (the chub),

intermediate for the species showing signs of incipient erosion (the Languedoc gudgeon) and the lowest

for the species that did not suffer from past erosion (the minnow). Although these results would deserve

further investigation (for instance with studies focusing on other species/taxa or other ecosystems), they

strongly suggest that the contemporary loss of genetic diversity observed worldwide (36) may impact

ecological functions in the same way as the loss of species (8), with possible critical consequences in

terms of ecosystem functioning and provision of services (37).

A major difficulty when investigating biodiversity-functions relationships in the wild is to control for the

direct and indirect effects of factors (other than biodiversity) that can affect ecosystem functions. This is

notably the case in spatially-structured landscapes such as rivers in which both spatial patterns of genetic

diversity and biomass yields are expected to depend on co-varying environmental factors along an

upstream-downstream gradient (32, 40, 41). This gradient is for instance known to shape evolutionary

processes such as downstream-biased gene flow and upstream-biased genetic drift, generating patterns

of downstream increase in genetic diversity (32). We accordingly detected a classical downstream

increase in genetic diversity in all three species, indicating that the river network topology also indirectly

contributed to a downstream increase in biomass stability. The upstream-downstream gradient might also

affect ecological processes such as foraging success, recruitment and competition, in turn causing spatial

variations in population dynamics and biomass yields (42, 43). In minnows and gudgeons for instance, two

species whose biological requirements are respectively fulfilled in the upstream and the intermediate parts

of the studied basin (42), the upstream-downstream gradient was responsible for a downstream decrease

in both biomass production and biomass stability, either directly or indirectly, through demography.

Fish species may also react differently to eutrophication levels. Associated with hypoxia and possible

exposure to sub-lethal concentrations of pollutants, eutrophication has indeed been shown to affect

individual behavior and physiology in different freshwater species, in turn impacting key-ecological

processes such as recruitment (44, 45). We for instance demonstrated that eutrophication indirectly

influenced biomass stability through a decrease in density stability in the two most sensitive fish species

(minnows and gudgeons). In chubs, a species that can tolerate degraded water quality (46), the positive

relationship between genetic diversity and biomass stability was particularly pronounced in the most

eutrophic habitats (Figure 3B). In accordance with empirical findings showing that among-individual

functional complementarity promoted by genetic and phenotypic variation is of greater importance under

more stressful than under benign conditions (22), this result suggests that genetic diversity may locally

contribute to higher individual tolerance and benefit to populations by buffering the possible negative

aftermaths of eutrophication or other anthropogenic stressors.

In line with both theoretical expectations and collective experimental evidence (22), we found strong

evidence that "real world" genetic diversity directly promotes temporal stability in biomass production,

particularly under the most stressful conditions. Similarly to what was demonstrated in natural settings at

the interspecific level (8), we showed that genetic diversity was somehow as important as other predictors

such as environment and demography to sustain biomass production over time. Given both the

abundance and the importance of studied species within trophic chains, we suggest that genetic diversity

might also indirectly promote community and ecosystem stability (28, 37), although deciphering these

relationships lied beyond the scope of our study. Future studies are now needed to confirm the

significance of these results in other taxa and to disentangle the relative contribution of intra- and

interspecific diversity in explaining biomass yields in the wild (21), notably in temperate ecosystems where

species diversity is naturally low. We yet also demonstrated that genetic erosion may jeopardize these

direct benefits in just a few generations. Considering the dramatic erosion of biodiversity worldwide and

the associated risks for the functioning and productivity of ecosystems (36, 47), our study adds to the

growing body of knowledge and stress the need for human societies to adopt prominent environmental

policies favoring all facets of biodiversity (48, 49).

Acknowledgments

We warmly thank all the colleagues and students who helped with field sampling. We are also grateful to

J. Canal for providing insightful details and explanations about the ecological state of studied rivers in

regard of the French implementation of the European Water Framework Directive 2000/60/EC. This work

has been financially supported by grants to SB from the Agence Française pour la Biodiversité

(DYNAGEN).

References

1. B. Worm, *et al.*, Impacts of Biodiversity Loss on Ocean Ecosystem Services. *Science* **314**, 787–790 (2006).

2. B. J. Cardinale, *et al.*, Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012).

3. G. M. Mace, K. Norris, A. H. Fitter, Biodiversity and ecosystem services: a multilayered relationship. **27**, 8 (2012).

4. A. Hector, The Effect of Diversity on Productivity: Detecting the Role of Species Complementarity. *Oikos* **82**, 597 (1998).

5. M. Loreau, C. de Mazancourt, Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. *Ecology Letters* **16**, 106–115 (2013).

6. F. van der Plas, Biodiversity and ecosystem functioning in naturally assembled communities. *Biological Reviews*, brv.12499 (2019).

7. D. Craven, *et al.*, Multiple facets of biodiversity drive the diversity–stability relationship. *Nature Ecology & Evolution* **2**, 1579–1587 (2018).

8. J. E. Duffy, C. M. Godwin, B. J. Cardinale, Biodiversity effects in the wild are common and as strong as key drivers of productivity. *Nature* **549**, 261–264 (2017).

9. K. Gross, *et al.*, Species Richness and the Temporal Stability of Biomass Production: A New Analysis of Recent Biodiversity Experiments. *The American Naturalist* **183**, 1–12 (2014).

10. D. U. Hooper, *et al.*, Effects of Biodiversity on Ecosystem Functioning: A Consensus of Current Knowledge. *Ecological Monographs* **75**, 3–35 (2005).

11. A. D. Barnosky, *et al.*, Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51–57 (2011).

12. D. Spielman, B. W. Brook, R. Frankham, Most species are not driven to extinction before genetic factors impact them. *Proceedings of the National Academy of Sciences* **101**, 15261–15264 (2004).

13. I. Saccheri, *et al.*, Inbreeding and extinction in a butterfly metapopulation. *Nature* **392**, 491–494 (1998).

14. D. H. Reed, R. Frankham, Correlation between fitness and genetic diversity. *Conservation Biology* **17**, 230–237 (2003).

15. S. M. Evans, E. A. Sinclair, A. G. B. Poore, K. F. Bain, A. Vergés, Genotypic richness predicts phenotypic variation in an endangered clonal plant. *PeerJ* **4**, e1633 (2016).

16. A. R. Hughes, B. D. Inouye, M. T. J. Johnson, N. Underwood, M. Vellend, Ecological consequences of genetic diversity. *Ecology Letters* **11**, 609–623 (2008).

17. D. I. Bolnick, *et al.*, Why intraspecific trait variation matters in community ecology. *Trends in*
Ecology & Evolution **26**, 183–192 (2011).

18. D. M. Olson, *et al.*, Terrestrial Ecoregions of the World: A New Map of Life on Earth. *BioScience* **51**,
933 (2001).

19. T. B. H. Reusch, A. Ehlers, A. Hammerli, B. Worm, Ecosystem recovery after climatic extremes
enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences* **102**, 2826–2831
(2005).

20. A. Siefert, *et al.*, A global meta-analysis of the relative extent of intraspecific trait variation in plant
communities. *Ecol Lett* **18**, 1406–1419 (2015).

21. M. Vellend, M. A. Geber, Connections between species diversity and genetic diversity. *Ecology*
Letters **8**, 767–781 (2005).

22. A. Forsman, L. Wennersten, Inter-individual variation promotes ecological success of populations
and species: evidence from experimental and comparative studies. *Ecography* **39**, 630–648 (2016).

23. R. J. Petit, Glacial Refugia: Hotspots But Not Melting Pots of Genetic Diversity. *Science* **300**, 1563–
1565 (2003).

24. E. Bestion, *et al.*, Altered trophic interactions in warming climates: consequences for predator diet
breadth and fitness. *Proc. R. Soc. B* **286**, 20192227 (2019).

25. M. Jangjoo, S. F. Matter, J. Roland, N. Keyghobadi, Connectivity rescues genetic diversity after a
demographic bottleneck in a butterfly population network. *Proceedings of the National Academy of*
Sciences **113**, 10914–10919 (2016).

26. A. P. Hendry, A critique for eco-evolutionary dynamics. *Functional Ecology* **33**, 84–94 (2019).

27. G. M. Crutsinger, *et al.*, Plant Genotypic Diversity Predicts Community Structure and Governs an
Ecosystem Process. *Science* **313**, 966–968 (2006).

28. A. Raffard, F. Santoul, J. Cucherousset, S. Blanchet, The community and ecosystem consequences of
intraspecific diversity: a meta-analysis: The ecological effects of intraspecific diversity. *Biological*
Reviews **doi: 10.1111/brv.12472** (2019).

29. J. B. Grace, *Structural equation modelling and natural systems.* (Cambridge University Press, 2006).

30. L. De Meester, *et al.*, Analysing eco-evolutionary dynamics—The challenging complexity of the real
world. *Funct Ecol* **33**, 43–59 (2019).

31. J. E. Duffy, J. S. Lefcheck, R. D. Stuart-Smith, S. A. Navarrete, G. J. Edgar, Biodiversity enhances reef
fish biomass and resistance to climate change. *Proceedings of the National Academy of Sciences*
113, 6230–6235 (2016).

32.   I. Paz-Vinas, G. Loot, V. M. Stevens, S. Blanchet, Evolutionary processes driving spatial patterns of

intraspecific genetic diversity in river ecosystems. *Molecular Ecology* **24**, 4586–4604 (2015).

33.   P. M. Kiffney, C. M. Greene, J. E. Hall, J. R. Davies, Tributary streams create spatial discontinuities in

habitat, biological productivity, and diversity in mainstem rivers. *Can. J. Fish. Aquat. Sci.* **63**, 2518–

2530 (2006).

34.   V. H. Smith, G. D. Tilman, J. C. Nekola, Eutrophication: impacts of excess nutrient inputs on

freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* **100**, 179–196 (1999).

35.   J. C. Garza, E. G. Williamson, Detection of reduction in population size using data from

microsatellite loci. *Molecular Ecology* **10**, 305–318 (2001).

36.   D. M. Leigh, A. P. Hendry, E. Vázquez-Domínguez, V. L. Friesen, Estimated six per cent loss of

genetic variation in wild populations since the industrial revolution. *Evol Appl*, eva.12810 (2019).

37.   E. P. Palkovacs, M. M. Moritsch, G. M. Contolini, F. Pelletier, Ecology of harvest-driven trait changes

and implications for ecosystem management. *Front Ecol Environ* **16**, 20–28 (2018).

38.   C. Gerstenmaier, S. Krueger-Hadfield, E. Sotka, Genotypic diversity in a non-native ecosystem

engineer has variable impacts on productivity. *Marine Ecology Progress Series* **556**, 79–89 (2016).

39.   T. W. Schoener, The newest synthesis: understanding the interplay of evolutionary and ecological

dynamics. *science* **331**, 426–429 (2011).

40.   F. Altermatt, Diversity in riverine metacommunities: a network perspective. *Aquatic Ecology* **47**,

365–377 (2013).

41.   J. G. Prunier, V. Dubut, G. Loot, L. Tudesque, S. Blanchet, The relative contribution of river network

structure and anthropogenic stressors to spatial patterns of genetic diversity in two freshwater

fishes: A multiple-stressors approach. *Freshwater Biology* **63**, 6–21 (2018).

42.   L. Buisson, L. Blanc, G. Grenouillet, Modelling stream fish species distribution in a river network:

the relative effects of temperature versus physical factors. *Ecol Freshwater Fish* **17**, 244–257

(2008).

43.   S. Nakano, M. Murakami, Reciprocal subsidies: Dynamic interdependence between terrestrial and

aquatic food webs. *Proceedings of the National Academy of Sciences* **98**, 166–170 (2001).

44.   S. M. Adams, *et al.*, The use of bioindicators for assessing the effects of pollutant stress on fish.

Marine Environmental Research **28**, 459–464 (1989).

45.   M. S. Pollock, L. M. J. Clarke, M. G. Dubé, The effects of hypoxia on fishes: from ecological

relevance to physiological effects. *Environ. Rev.* **15**, 1–14 (2007).

46. C. Liu, Y. Chen, J. R. Britton, Influences of environmental and chemical parameters on the spatial
growth patterns of four riverine cyprinid fishes. *Knowl. Manag. Aquat. Ecosyst.*, 12 (2015).

47. A. Miraldo, *et al.*, An Anthropocene map of genetic diversity. *Science* **353**, 1532–1535 (2016).

48. W. J. Ripple, *et al.*, World Scientists' Warning to Humanity: A Second Notice. *BioScience* **67**, 1026–
1028 (2017).

49. O. Rey, *et al.*, Linking epigenetics and biological conservation: Toward a *conservation epigenetics*
perspective. *Funct Ecol*, 1365-2435.13429 (2019).

50. N. Poulet, L. Beaulaton, S. Dembski, Time trends in fish populations in metropolitan France: insights
from national monitoring data. *Journal of Fish Biology* **79**, 1436–1452 (2011).

51. L. Fourtune, I. Paz-Vinas, G. Loot, J. G. Prunier, S. Blanchet, Lessons from the fish: a multi-species
analysis reveals common processes underlying similar species-genetic diversity correlations.
Freshwater Biology **61**, 1830–1845 (2016).

52. P. Keith, H. Persat, E. Feunteun, B. Adam, M. Geniez, *Les Poissons d'eau douce de France*, Muséum
National d'Histoire Naturelle and Publications Biotope (2011).

53. M. Chevalier, L. Comte, P. Laffaille, G. Grenouillet, Interactions between species attributes explain
population dynamics in stream fishes under changing climate. *Ecosphere* **9**, e02061 (2018).

54. B. Haegeman, *et al.*, "Resilience, invariability, and ecological stability across levels of organization"
(Ecology, 2016) https:/doi.org/10.1101/085852 (April 25, 2019).

55. S. M. Aljanabi, I. Martinez, Universal and rapid salt-extraction of high quality genomic DNA for PCR-
based techniques. *Nucleic acids research* **25**, 4692–4693 (1997).

56. A. Futschik, C. Schlotterer, The Next Generation of Molecular Markers From Massively Parallel
Sequencing of Pooled DNA Samples. *Genetics* **186**, 207–218 (2010).

57. R Development Core Team, *R: A Language and Environment for Statistical Computing, R*
Foundation for Statistical Computing (2014).

58. K. Belkhir, P. Borsa, L. Chikhi, N. Raufaste, F. Bonhomme, *GENETIX 4.03, logiciel sous WindowsTM*
pour la génétique des populations. Laboratoire Génome, Populations, Interactions CNRS UMR 5000,
Université de Montpellier II, Montpellier (France). (2004).

59. Z. A. Szpiech, M. Jakobsson, N. A. Rosenberg, ADZE: A rarefaction approach for counting alleles
private to combinations of populations. *Bioinformatics* **24**, 2498–2504 (2008).

60. A. R. Hughes, J. J. Stachowicz, Ecological impacts of genotypic diversity in the clonal seagrass
Zostera marina. *Ecology* **90**, 1412–1419 (2009).

61. E. Mathieu-Bégné, G. Loot, M. Chevalier, I. Paz-Vinas, Blanchet Simon, Resources for "Demographic
and genetic collapses in spatially-structured populations: Insights from a long-term survey in wild
fish metapopulations." 107781 Bytes (2018).

62. H. Pella, J. Lejot, N. Lamouroux, T. Snelder, Le réseau hydrographique théorique (RHT) français et
ses attributs environnementaux. *Géomorphologie : relief, processus, environnement* **3** (2012).

63. S. Caschili, "ComplexNetGIS: a tool for the analysis of complex spatial networks" in *Informatica e*
Pianificazione Urbana e Territoriale, G. Las Casas, P. Potrandolfi, B. Murgante, Eds. (Melfi, 2010),
pp. 233–242.

64. D. Pont, O. Delaigue, J. Belliard, A. Marzin, M. Logez, "Programme IPR+ : révision de l'indice poisson
rivière pour l'application de la DCE - version V.2.0 de l'indicateur." (IRSTEA-ONEMA, 2013).

65. J. Oehri, B. Schmid, G. Schaepman-Strub, P. A. Niklaus, Biodiversity promotes primary productivity
and growing season lengthening at the landscape scale. *Proceedings of the National Academy of*
Sciences **114**, 10160–10165 (2017).

66. L. S. Aiken, S. G. West, R. R. Reno, *Multiple regression: Testing and interpreting interactions* (SAGE
Publications, 1991).

67. F. W. Allendorf, Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biology* **5**, 181–190
(1986).

68. J. M. Batista-Foguet, G. Coenders, W. E. Saris, J. Bisbe, Simultaneous Estimation of Indirect and
Interaction Effects using Structural Equation Models. *Metodološki zvezki* **1**, 163–184 (2004).

69. K. P. Burnham, D. R. Anderson, *Model selection and multimodel inference: a practical information-*
theoretic approach, 2nd ed (Springer, 2002).

70. L. Hu, P. M. Bentler, Cutoff criteria for fit indexes in covariance structure analysis: Conventional
criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal* **6**, 1–55
(1999).

71. Y. Rosseel, **lavaan** : An *R* Package for Structural Equation Modeling. *Journal of Statistical Software*
48 (2012).

566 **Figure 1.** Geographic situation of the Garonne-Dordogne river basin in South-Western France and

567 localization of the 47 unique river stations, with pie charts indicating species (co-)occurrence within each

568 station. The black arrow indicates the location of the river mouth. Background is a shaded relief map.

A

B

C

Figure 2. Simplified causal graphs depicting retained links among environmental (blue rectangles),

demographic (purple rectangles), genetic (yellow rectangles) and biomass variables (green rectangles) in

chubs (A), gudgeons (B) and minnows (C). First-order interactions are represented by rounded rectangles

(washed-out when not involved in any causal link). Red and green arrows represent negative and positive

significant paths, respectively, with the width of arrows proportional to the absolute value of the

corresponding path coefficient. Washed-out green arrows stand for direct positive links between density

and biomass variables. Dashed grey arrows represent non-significant paths. Covariations and paths

removed during the simplification procedure are not displayed, for the sake of clarity.

Figure 3. Upper panels: Contributions of demographic variables (mean density and density stability),

environmental gradients (upstream-downstream gradient and eutrophic gradient) and genetic diversity

(genetic diversity and associated interactions) to biomass production (A) and temporal stability in biomass

production (B) in each species. The average contributions of each predictor to the explained variance in

biomass yields across species is shown in the form of 100% stacked barplots (%R²). Colors are the same

as in Figure 2. Lower panels: Distribution of levels of genetic diversity (C; here, microsatellite standardized

allelic richness) and bottleneck probability (D) as measured at the population-level in each species. The

horizontal black line in D indicates the threshold above which populations are supposed to have

experienced recent and severe bottleneck (M-ratio < 0.68).

Figure 4. Continuous representations of additive relationships or first-order interactions associated with biomass stability (A to C) and genetic diversity (D) in final simplified models in chubs (A), gudgeons (B) and minnows (C and D). Predicted values of biomass stability and genetic diversity were computed across all possible values of the two corresponding additive terms and plotted as a colored surface with contours. Horizontal and vertical black dashed lines indicate additive trends (i.e., the predicted relationship between the response and a given predictor when the other predictor is set to zero). Black dots, standing for observed data, and lateral shaded surfaces, standing for kernel density estimates of data points along each additive variable, allow visualizing the occupied parameter space. The white dashed line delimits an empty space in panel C. White arrows indicate noticeable trends. Biomass stability increased with genetic

604   diversity in all species (A-C) but mostly in eutrophic habitats in chubs (A) and in upstream populations in

605   minnows (C). Biomass stability also decreased with eutrophication in gudgeons (B). Minnows showed a

606   classical downstream increase in genetic diversity, but under the less eutrophic conditions only (D).

Figure 5. Characteristics of the two first principal components based on environmental data. a) Eigenvalues corresponding to the amount of the variation explained by each principal component. b) Visualization of the two first principal components (PC), with PC1 standing for the upstream-downstream gradient and PC2 standing for the eutrophic gradient. Percentage of variance explained by each component is also indicated. BET: Betweenness; DFM: Distance from the river mouth; DFS: Distance from the source; ANTH: anthropogenic river alteration. Solid black arrows: variables with a significant contribution to PC1; dashed black arrows: variables with a significant contribution to PC2; grey arrows: variables that did not contribute to any of the two first components.

Table 1. Main results from the first set of causal models, excluding biomass variables. For each species,

the table provides absolute fit indices (CFI, SRMR and p-value; see text for details), amounts of explained

variance (R^2) in response variables (indicated with an arrow) and estimates of path coefficients (indicated

by arrows). Significant path coefficients threshold are in bold (see Appendix S9 for a table with 95%

confidence intervals). Slashes indicate links that were discarded in the course of model simplification. Only

links retained in at least one species are shown. UDG: upstream-downstream gradient; EG: Eutrophic

gradient.

		S. cephalus	*G. occitaniae*	*P. Phoxinus*
	χ² p-value	0.908	0.638	0.986
Model fit indices	CFI	1	1	1
	SRMR	0.037	0.075	0.017
Upstream-downstream gradient	→ Genetic diversity	**0.469**	**0.372**	**0.311**
Eutrophic gradient	→ Genetic diversity	/	**0.444**	0.249
UDG x EG	→ Genetic diversity	/	/	**-0.35**
Bottleneck	→ Genetic diversity	**-0.451**	**-0.555**	-0.296
	R^2	0.548	0.548	0.353
Density stability	→ Bottleneck	/	**-0.309**	/
	R^2	/	0.095	/
Upstream-downstream gradient	→ Mean density	**0.544**	**-0.372**	/
	R^2	0.296	0.272	/
Upstream-downstream gradient	→ Density stability	/	**-0.415**	-0.241
Eutrophic gradient	→ Density stability	/	**-0.5**	**-0.319**
	R^2	/	0.371	0.129

630 **Table 2.** Main results from the second set of causal models including biomass variables, with estimates

631 from the first set of causal models being fixed. GD: Genetic diversity. See table 1 for complete legend.

		S. cephalus	*G. occitaniae*	*P. Phoxinus*
	χ² p-value	0.83	0.957	0.119
Model fit indices	CFI	1	1	0.961
	SRMR	0.067	0.079	0.084
Mean density → Biomass production		**0.733**	**0.737**	**0.932**
Upstream-downstream gradient → Biomass production		/	**-0.406**	-0.296
Eutrophic gradient → Biomass production		/	**-0.221**	-0.22
	R²	0.598	0.909	0.959
Density stability → Biomass stability		**0.393**	**0.509**	**0.754**
Upstream-downstream gradient → Biomass stability		/	**-0.486**	-0.114
Eutrophic gradient → Biomass stability		-0.096	**-0.253**	**-0.158**
Genetic diversity → Biomass stability		0.134	**0.198**	-0.079
GD x UDG → Biomass stability		/	/	**-0.188**
GD x EG → Biomass stability		**0.292**	/	/
	R²	0.363	0.738	0.705