Sympatric ecological divergence with coevolution of niche preference

Pavel Payne[1,2], Jitka Polechová[3]

[1] Charles University in Prague, Department of Zoology

[2] Institute of Vertebrate Biology, Czech Academy of Sciences

[3] University of Vienna, Faculty of Mathematics

Keywords

Sympatric speciation, maintenance of polymorphism, polygenic trait, Levene model, epistasis, trade-off

Abstract

Reinforcement, the increase of assortative mating driven by selection against unfit hybrids, is conditional on pre-existing divergence. Yet, for ecological divergence to precede the evolution of assortment, strict symmetries between fitnesses in niches must hold, and/or there must be low gene flow between the nascent species. It has thus been argued that conditions favouring sympatric speciation are rarely met in nature. Indeed, we show that under disruptive selection, violating symmetries in niche sizes and increasing strength of the trade-off in selection between the niches quickly leads to loss of genetic variation, instead of evolution of specialists. The region of the parameter space where polymorphism is maintained further narrows with increasing number of loci encoding the diverging trait and the rate of recombination between them. Yet, evolvable assortment and pre-existing assortment both substantially broaden the parameter space within which polymorphism is maintained. Notably, pre-existing niche preference speeds up further increase of assortment, thus facilitating reinforcement in the later phases of speciation. We conclude that in order for sympatric ecological divergence to occur, niche preference must co-evolve throughout the divergence process. Even if populations come into secondary contact, having

diverged in isolation, niche preference substantially broadens the conditions for coexistence

in sympatry and completion of the speciation process.

Introduction

Darwin observed that many closely related species occupied the same habitat. However, he

considered the sympatric origin of species by ecological divergence due to an advantage of

specialists over generalists rather unlikely [1]. Since then, the contribution of sympatric

speciation to observed biodiversity has remained controversial [2–5]. Speciation in a well-

mixed, panmictic population is difficult for two main reasons. First, gene flow and

recombination oppose divergence in polygenic traits as well as preventing reinforcement

(the build-up of associations between the loci contributing to pre- and post-zygotic isolation)

[6,7]. Second, the diverging types must be able to coexist. It is often thought that ecological

divergence must precede the evolution of mating assortment [8], as assortment is

reinforced by selection against maladapted hybrids. However, this order of events is not

necessary: Assortment can precede divergence, or co-evolve with ecological loci. It is the

combined effect of assortment and ecological divergence which contributes to the isolation

of the nascent species [9–11]. Yet, in contrast to ecological selection, the loss of fitness due

to missed heterospecific matings is typically compensated by an increase in conspecific

matings. This stabilises polymorphism at the loci determining assortment and can protect

polymorphism at other loci diverging under natural selection.

Habitat choice is an important source of assortment when mating occurs locally. Host-

specific mating is prevalent in nature, as in phytophagous insects (reviewed in [12]). Habitat

fidelity based on a preference for the hatching site is common for birds [13], another group

rich in sympatric species. In general, habitat choice can be based on learned characteristics

of the hatching site, on specific preference loci, or on an association with another

(ecological) phenotype [14]. The first two drove the classic sympatric speciation process in

the experiment by Rice and Salt [15], where fruit flies learned to choose based on phototaxis

and chemotaxis. In this experiment, 60 mated females from each of two extreme habitats

founded the next generation. Severe disruptive selection on multiple traits, coupled with

independent regulation within equally sized niches, led to nearly complete reproductive

isolation over 30 generations. Effectively, behavioural allopatry evolved (c.f. [8]). Further

divergence in other traits, both selective and neutral, could have followed. While the

plausibility of sympatric speciation is undisputed, how common can it really be? Selection is

rarely that severe in nature, and even if niches are independently regulated, they will very

rarely be perfectly symmetric.

A recognition and preference for the correct habitat is a common example of assortment by

association with another phenotype. While preference for food source is omnipresent,

strong [16] trade-offs in specialization of the sympatric-species-rich phytophagous insects

have been hard to find, triggering a substantial controversy [17,18]. Yet, strong trade-offs

must be reasonably common – otherwise, all habitats would be colonised by one generalist

species. Indeed, such trade-offs have often been recovered when life-time fitness and/or

performance of F1 hybrids on both hosts is assessed [19–23]. Moreover, a preference-

performance correlation seems to be common in herbivorous insects [24,25].

Here we use mathematical modelling to investigate how genetically encoded preferences for

the most favourable habitat facilitate sympatric speciation by ecological divergence. We

assume that there is a strong (convex) trade-off in a polygenic trait, leading to disruptive

selection such that generalists have a lower mean fitness than specialists. Assortment arises

via preference for the correct niche. Because regulation is independent within niches (soft

selection, [26]), the frequency of a type well adapted to its niche rises faster when this type

is rare than when it is common. This protects the polymorphism in the ecological loci

(coexistence of the specialists) via negative frequency-dependence [26].

We consider a polygenic trait under ecological divergence, and focus on assortment via

preference for the niche the individual is best adapted to. The assortment allele can

therefore be seen as a type of a 'modifier-allele' [8], realized via niche choice. The

population-wide matching of the habitat thus increases with the frequency of this modifier

allele. The modifier represents an 'one-allele model' [7], where recombination cannot break

down the allele-specific association with the choice-locus. Thereby, the constraints on

symmetries, which render sympatric speciation implausible, are relaxed. The modifier-allele

niche choice contrasts with two-allele models, where habitat or mate choice is determined

by loci independent of the ecological adaptation. Then, sympatric speciation under strong

(convex) trade-off is implausible as variation is quickly lost (c.f. [7,11,27]). Such an

assortment only enables evolution of coexisting specialists when trade-offs are weak

(concave) [16]. Note that modifier-allele niche choice is not a fixed 'matching' habitat choice,

where an ecological allele also directly increases the probability of getting in to the right

niche (magic niche choice), which always leads to coexistence [16].

How do two nascent species evolve towards coexistence? Coexistence during divergence in

sympatry is often overlooked in speciation models, side-lined by focus on growth of

associations between and within polygenic traits, assuming symmetries are protected [11].

The importance of pre-existing ecological divergence and/or of pre-existing assortment for

further diversification in sympatry has not been much studied. Hence, it is not clear whether

divergence and coexistence in sympatry is considerably more difficult than partial

divergence in allopatry or broad parapatry, followed by sympatry as assortment increases.

When neutral divergence in allopatry or broad parapatry does not lead to a sufficient niche

separation then (similarly to instant assortment by chromosomal variations; [28])

coexistence will not be possible. Yet, coexistence in sympatry is essential: without it, every

speciation in sympatry or parapatry would lead to a subdivision of the existing range – and

through time, species' ranges would become ever smaller.

Here we aim to elucidate the importance of niche recognition in facilitating divergence

under disruptive ecological selection on a polygenic trait. In other words, we study how pre-

existing prezygotic isolation (niche recognition) increases and coevolves jointly with

postzygotic isolation driven by a strong (convex) trade-off in adaptation to different

ecological niches. Does sympatric speciation remain feasible when negative epistasis

between loci increases (and hence the trade-off strengthens), and when with increasing

number of loci, selection becomes weaker relative to recombination?

Methods and results

Model

Since we focus on the effects of disruptive selection on sympatric divergence, we use a

haploid biallelic version of Levene's [29] model with mating within niches, which is more

favourable to speciation [30]. In the lifecycle of the model, individuals with non-overlapping

generations are – in the basic model without assortment – randomly distributed into two

niches, where selection acts. Alleles of the first type (capital letters) are beneficial in one

niche and alleles of the second type (small letters) in the other niche. The individuals who

survive mate within the niches and then each niche contributes to a common pool

proportionally to its size. From the common pool, individuals are again randomly distributed

into the two niches.

For further analysis, it is convenient to define fitness of the genotypes to be between zero

and one. The presence of an allele which is deleterious in one of the niches reduces the

fitness of its carrier by selection coefficient s, which is equal for all loci and both niches. The

epistatic coefficient ϵ determines how much the intermediate generalist genotypes (e.g., aB

and Ab) are disadvantaged relative to the specialist genotypes (e.g., AB and ab for $\epsilon < 0$)

and is defined as the deviation from additivity of deleterious alleles in each niche if more

than one of these alleles is present. Here we are assuming a special kind of symmetric

epistasis, which acts among alleles deleterious in one niche and also those in the other

niche. Fitnesses of all genotypes considered in this study are defined in detail further in the

text and Tables $1 - 4$.

Analytical solutions for polymorphic equilibria of our model are generally impossible to find.

Instead, we have analysed the instability of the two monomorphic equilibria for allele

frequencies in ecological and niche preference loci at 1 and 0. We assume that an

equilibrium polymorphic at the ecological loci exists between them. This approach was used

by Levene [29], Prout [31] and Hoekstra et al. [32]. Gliddon and Strobeck [33] analysed the

haploid analogue of Levene's [29] model with multiple niches, and proved that conditions

for instability of the monomorphic equilibria are also necessary and sufficient conditions for

stability of the unique polymorphic equilibrium. Although this conclusion is not valid for

general fitnesses, such as in the presence of epistasis and/or when assortment is present,

analysis of local (in)stability of the monomorphic equilibria still allows us to estimate the

stability of the polymorphic equilibrium for the ecological loci in the full model for most of

the parameter range. The stability is then tested numerically. We discuss a small parameter

region where the local equilibrium also depends on the initial conditions. Depending on the

initial allele frequencies, the system evolves either to a polymorphic equilibrium in the

ecological loci, or to another equilibrium where ecological loci are fixed and the locus for

assortment is polymorphic, or where all polymorphism is lost.

We complement the 2-3 locus analytical and numerical results by an individual-based

haploid simulation, with up to 16 ecological loci under disruptive selection (strong trade-off),

and 10 loci which gradually change the bias in the niche choice from 0 to 1. Within niches,

each individual is selected at random (with replacement) to pair up, with a probability

proportional to its fitness. The recombination rate r is uniform across loci, as is the mutation,

set to $\mu = 10^{-3}$. Each of the N_i pairs has exactly one offspring, the size of niche i is thus kept

constant at N_i. The juveniles are then redistributed between habitats, according to their

preference, which reflects linearly their number of matching alleles. For example, an A-B-c-

M-M-m-[..]-m type will add to its preference $\mathcal{A}(z) = {}^2\!/_3 (\alpha_1 + \alpha_2)$, and thus choose niche

1 with a probability of $\frac{c(1+\mathcal{A}(z))}{1+(2c-1)\mathcal{A}(z)}$, where c is the relative size of niche 1. The α_i sum to at

most 1 over all 'M' loci (whilst $\alpha_i = 0$ for all 'm' loci). Generations are discrete and non-

overlapping.

Ecological divergence at a polygenic trait

Ecological divergence is often driven by disruptive selection acting on polygenic traits.

Therefore, we first set out to analyse how increasing the number of loci influences the ability

of the model to maintain polymorphism under disruptive selection. We study two to three

loci analytically and for up to 16 loci using individual-based simulations. In particular, we

focus on violation of symmetries in niche proportions, which are defined as $c = c_I = 1 - c_{II}$

for niche I and II, respectively ($c \in (0,1)$). In order to be able to compare the two- ($n_{eco} = 2$)

and three- ($n_{eco} = 3$) locus models, we normalise the strength of selection (s) and epistasis

(ϵ) in the models with more than two ecological loci such that the mean per-trait selection

and epistasis remain the same ($s^*_{n_{eco}} = \frac{2\,s_{2loci}}{n_{eco}}$ and $\epsilon^*_{n_{eco}} = \frac{1}{\binom{n_{eco}}{2}}\epsilon_{2loci}$). This normalisation

assures that fitnesses of the specialists are always equal to $1 - 2s - \epsilon$, whilst preserving the

scaling of selection independently of epistasis. This leads to a slight weakening of the trade-

off as the number of loci increases (see also Fig. S1).

For simplicity, we assume that all pairwise epistatic effects have the same value, and neglect

higher-order interactions between alleles. Fitnesses of the individual genotypes for the two-

and three-locus model are defined in Table 1A and Table 1B, respectively.

Graphical representation of the trade-offs between fitnesses of individual genotypes in niche

I and niche II is shown discussed in Fig. S1 and the corresponding supplementary text.

A

genotype	niche I	niche II
AB	1	$1 - 2s - \epsilon$
aB, Ab	$1 - s$	$1 - s$
ab	$1 - 2s - \epsilon$	1

B

genotype	niche I	niche II
ABC	1	$1 - 2s - \epsilon$
ABc, AbC, aBC	$1 - \frac{2}{3}s$	$1 - \frac{4}{3}s - \frac{\epsilon}{3}$
Abc, aBc, abC	$1 - \frac{4}{3}s - \frac{\epsilon}{3}$	$1 - \frac{2}{3}s$
abc	$1 - 2s - \epsilon$	1

***Table 1**. **Fitnesses of individual genotypes in the model without niche preference.** A. Two-locus model. B.*

Three-locus model. Selection coefficient s determines how the deleterious alleles in niche I (a, b, c) and the

deleterious alleles niche II (A, B, C) reduce the fitness relative to the well adapted genotype. The pairwise

epistatic coefficient ϵ is defined as the deviation from additivity of the two deleterious alleles present in a

genotype together. We only consider $\epsilon < 0$ as it generates a convex trade-off and hence disruptive selection. In

the three-locus model the selection and epistasis are normalised as described above. Note that the epistatic

coefficient in the intermediate genotypes in the three-locus model is present only when the genotype contains

two alleles that are deleterious in the particular niche. In the other niche, the epistatic coefficient for these

genotypes is absent as there is only one selected allele.

First, we show that the region of parameter space where polymorphism is maintained is

highly sensitive to violation of symmetry in niche proportions, c, even if loci are completely

linked. Also, increasing convexity of the trade-off (i.e., more negative ϵ, more disruptive

selection) further reduces the parameter space with maintained polymorphism (Fig. 1A).

Both when the loci are completely linked (r=0) and when they are freely recombining (r=0.5),

there is no difference between two and three loci when mean per-trait selection and

epistasis remain the same (see above). The regions of parameter space with maintained

polymorphism coincide and shift towards stronger selection as epistasis increases (Fig. 1C).

When recombination between loci is low (r=0.01, i.e., 1cM), the regions of parameter space

with maintained polymorphism shift closer towards those of the free recombination regime

in the three-locus model than in the two-locus model, as in the three-locus case the per-

locus strength of selection is lower (Fig. 1B). Interestingly, there is a threshold for

recombination rate, above which the parameter space where polymorphism is maintained is

independent of the number of loci, which we elaborate on more in the Supplementary

Information. Below we provide the analytical expressions for the boundaries of the stable

regions, and the recombination threshold. The conditions are for protected polymorphism in

the ecological loci, obtained by assessing local instability of the monomorphic equilibria.

Note that as we have disruptive selection with symmetric selection coefficients, all

ecological loci, when polymorphic, evolve to the same allele frequencies.

In both models with normalized selection and $\epsilon = 0$ a polymorphic equilibrium is stable if

$$\frac{1-2s}{2-2s} < c < \frac{1}{2-2s}.$$

In the case with no recombination (Fig. 1A) and $\epsilon < 0$, polymorphism is maintained if

$$\frac{2s+\epsilon-1}{2s+\epsilon-2} < c < \frac{1}{2-\epsilon-2s},$$

and with free recombination (Fig. 1C) if

$$\frac{s(1-2s-\epsilon)}{(1-s)(2s+\epsilon)} < c < \frac{s+\epsilon}{(1-s)(2s+\epsilon)}.$$

These conditions hold for the two-locus model for recombination rates $r > \frac{-\epsilon}{1-s-\epsilon}$. If

recombination rate is low ($0 < r < \frac{-\epsilon}{1-s-\epsilon}$; Fig. 1B), the conditions above do not hold

anymore and maintenance of polymorphism is determined by

$$\frac{(1-2s-\epsilon)(2s+\epsilon(1-r)+r(1-2s))}{(1-r)(4(1-s)s+2\epsilon(1-2s)-\epsilon^2)} < c < \frac{2s+\epsilon-r}{(1-r)(4(1-s)s+2\epsilon(1-2s)-\epsilon^2)}.$$

Since the equations describing the conditions for the three-locus model with low

recombination and all other models presented from this point on exceed the width of a

page, we confine them to the Supplementary material, where we also give more details on

the stability analysis.

Figure 1. **Range of niche proportions where polymorphism is maintained: where ecological divergence**

towards two specialists is stable. The upper half of the graphs show conditions for the two-locus model, the

lower half for the three-locus model. On the x-axis is the normalized strength of selection (symmetric across

loci) and on the y-axis are niche proportions. A polymorphic equilibrium is achieved for the parameter

combinations of s^* and c between the black curves and the white axis at $c = 0.5$ (shaded). Note that we are

showing only one half of the parameter space for each model as the conditions are symmetrical. Therefore,

every condition at a value of c has its symmetric counterpart at $1 - c$. The outer solid curves represent linear

trade-offs ($\epsilon = 0$), the dashed curves are for ($\epsilon = -0.25s$), and the dash-dotted curves for ($\epsilon = -0.5s$). A) No

recombination; B) Low recombination ($r = 0.01$), and C) Free recombination ($r = 0.5$). It is noteworthy that in

the regime with low recombination (B), the region of parameter space where polymorphism is maintained

decreases with increasing number of loci if selection and epistasis is normalized as described in the main text.

Ecological divergence with niche choice

Next, we analyse how the presence of a niche preference allele influences the sensitivity of

polymorphism to violation of symmetry in niche proportions. We assume that habitat choice

depends both on the presence of a choosy (modifier) allele M and the phenotype under

natural divergent selection. For the individuals with wildtype modifier allele m, there is no

bias in selecting niches. When the modifier allele is M, choice depends on the selected loci

in the following manner: intermediate generalist genotypes (*Ab*, *aB*) disperse to both niches

with equal probability, and specialist genotypes (*AB*, *ab*) choose with probability $1 + \alpha$ the

niche which they are adapted to, and with probability $1 - \alpha$ the niche where they are

maladapted. Since in our model mating occurs within niches, this translates into a direct

modification of fitnesses of the individual genotypes, as defined in Table 2.

genotype	niche I	niche II
ABM	$1 + \alpha$	$(1 - 2s - \epsilon)(1 - \alpha)$
ABm	1	$1 - 2s - \epsilon$
Abm, AbM, aBm, aBM	$1 - s$	$1 - s$
abm	$1 - 2s - \epsilon$	1
abM	$(1 - 2s - \epsilon)(1 - \alpha)$	$1 + \alpha$

Table 2. *Fitnesses of individual genotypes modified by niche choice parameter α. Selection and epistasis modify*

fitnesses of individual genotypes as in table 1A. Genotype ABM goes to niche I with a probability increased by

the factor α and to niche II decreased by α. Similarly, genotype abM preferentially goes to niche II instead of

niche I.

In the presence of niche preference, the parameter space where polymorphism is

maintained broadens (Fig. 2). As allele *M* is beneficial for both specialist genotypes (*AB* and

ab), it quickly goes to fixation when polymorphisms for the other two loci are maintained –

hence, two specialists coexist. This holds in the case of complete linkage (Fig. 2A) and low

recombination (Fig. 2B). In the scenario with free recombination, maintenance of

polymorphism at the ecological loci depends on the initial frequencies of alleles *A, B* and *M*

in the population. With weaker selection against the maladapted genotypes, s^*, the model is

more sensitive to initial conditions, as indicated by the fading grey colour in the lower left

part of Fig. 2C. Within this region, depending on the initial allele frequencies, either the

preference allele M goes to fixation and protects polymorphism in the other loci, or the

ecological loci go to fixation and allele M remains polymorphic and converges to its

equilibrium frequency $\frac{|1-2c|}{\alpha}$. Figure 2D shows which combinations of initial frequencies of

the ecological ($p_1 = p_2$) and preference (p_3) alleles lead to which equilibria for four points

(a, b, c, d) in the parameter space in Fig. 2C.

Figure 2. Ranges of niche proportions where ecological polymorphism is maintained (i.e., two specialists evolve

and coexist), under coevolution of niche preference with ecological divergence.

A-C: Plots showing how niche recognition allele M with niche preference $\alpha = 0.2$ broadens the parameter space where polymorphism at two ecological loci is maintained, relative to the model without preference (upper half of the plot; axes and all other parameters are identical to Fig 1).

D: Streamline plots showing how maintenance of polymorphism depends on initial allele frequencies at ecological and niche preference loci for four points in Fig 2C (**a**, s=0.01; **b**, s=0.1; **c**, s=0.2; **d**, s=0.3; r=0.5, c=0.46). Black streamlines represent combinations of allele frequencies where ecological polymorphism is maintained, grey streamlines where either only ecological or all polymorphisms are lost. Since the plots were generated assuming linkage disequilibrium between ecological loci goes to 0 (i.e. loss of ecological polymorphism), the true equilibrium ecological allele frequencies, which are slightly offset with the black streamline attractors, are represented by black dots at the top of plots (**D.a-D.d**).

Increase of assortment

Once such a niche preference modifier allele gets fixed in a population, it not only inflates

the parameter space where polymorphism is maintained but it also always favours fixation

of another modifier allele, which then reinforces the divergence process. In order to analyse

such an increase of assortment, we again redefined fitnesses of the genotypes as shown in

Table 3. The fitnesses of the specialist genotypes (*AB* and *ab*) are now defined such that the

niche preference allele *M* from the previous model is fixed in the population. Therefore,

both specialist genotypes have their probability of going to the right niche increased by α_1.

The third locus can now be polymorphic for another modifier allele, which further increases

the probability of going to the right niche by α_2. In Fig. 3A-C we show how the parameter

space where polymorphism is maintained further broadens in the presence of another

modifier allele, increasing the assortment.

Furthermore, we numerically tested how different levels of pre-existing niche preference

affect time to fixation of a new niche preference allele. In Fig. 3D-G we show that with

increasing initial niche preference (α_1) the time to fixation of a new niche preference allele

($\alpha_2 = 0.2$) decreases. In particular, this effect is most profound when selection (s^*) is weak

and recombination (r) is low (Fig 3D, E).

331
332

Figure 3. Range of niche proportions where polymorphism is maintained when assortment further increases.

A-C: Plots showing how the parameter space where polymorphism at two ecological loci is maintained with

already fixed niche preference $\alpha_1 = 0.2$, and an additional allele M_2 with niche preference $\alpha_2 = 0.2$. In Figure

3C, which depicts the parameter space for free recombination ($r = 0.5$), the region where maintenance of

ecological polymorphism depends on initial allele frequencies is indicated by a shading grey area (as in Fig. 2C).

D-G: Time to fixation (y-axis), of the additional niche preference allele M_2 with initial frequency $p_3 = 0.01$ for

various combinations of initial niche preference, α_1 (x-axis), strengths of selection, s^*, and recombination

rates, r. All plots are for the same disruptiveness of selection, $\epsilon = -0.5s$. The time to fixation is reduced most

when selection is weak ($s^* = 0.01, 0.05$), and recombination rate is low ($r = 0.01, 0.05$) (**D, E**). In contrast, with

moderate or strong selection ($s^* = 0.1, 0.2$) and low recombination ($r = 0.01$) the time to fixation of the niche

preference allele M_2 either decreases very little or even slightly increases with increasing initial niche

preference, α_1. Note that the values on the y-axis (time to fixation) differ in D-G.

genotype	niche I	niche II
ABM_2	$1 + \alpha_1 + \alpha_2$	$(1 - 2s - \epsilon)(1 - \alpha_1 - \alpha_2)$
ABm_2	$1 + \alpha_1$	$(1 - 2s - \epsilon)(1 - \alpha_1)$
$AbM_2, aBM_2, Abm_2, aBm_2$	$1 - s$	$1 - s$
abm_2	$(1 - 2s - \epsilon)(1 - \alpha_1)$	$1 + \alpha_1$
abM_2	$(1 - 2s - \epsilon)(1 - \alpha_1 - \alpha_2)$	$1 + \alpha_1 + \alpha_2$

Table 3. Fitnesses of individual genotypes modified by a fixed niche choice parameter α_1 and a polymorphic

allele M_2, which increases niche preference for the better niche by α_2. The genotype ABM_2 goes to niche I with

a probability increased by the factor $\alpha_1 + \alpha_2$ and to niche II decreased by $\alpha_1 + \alpha_2$. Similarly, genotype ABM_2

preferentially goes to niche II instead of niche I. Preferences of other genotypes are identical as in Table 2.

Secondary sympatry

It should be noted that all conditions presented above are independent of the initial linkage

disequilibrium between the ecological loci. Therefore, for maintenance of genetic

polymorphism in the ecological loci it is irrelevant whether the population starts in a Hardy-

Weinberg linkage equilibrium or if it consists of only two specialist genotypes as it would

during a secondary contact after divergence in allopatry.

Cost of niche choice

It has been suggested by de Meeûs et al. [34] that in a model with soft selection and habitat

preference, the conditions under which polymorphism is maintained appear to be slightly

broadened but only if there is no cost to habitat selection. We implemented a cost of niche

choice, γ, into our model, which is defined as a fraction of the selection coefficient (fitnesses

are described in Table 4). The cost is associated with presence of the allele that allows for

niche discrimination and choice, therefore, it is present even if the individual genotype has

no preference for any of the niches but carries the niche preference allele M. In other words,

it can be seen as a cost of an ability to discriminate (and choose).

As analytical solutions are rather difficult to obtain, we numerically assessed how various

values of γ affect maintenance of polymorphism in the ecological loci. In Fig. S2 we show

that for low recombination (around $r = 0.01$), certain cost is robustly tolerated in a large

region of the parameter space. E.g., in an example with $s = 0.1, c = 0.46$ and $\epsilon = -0.5s$,

cost $\gamma = 0.05s$ is tolerated for $\alpha_2 \geq 0.1$, cost $\gamma = 0.1s$ for $\alpha_2 \geq 0.2$, and cost $\gamma = 0.2s$ for

$\alpha_2 \geq 0.3$.

genotype	niche I	niche II
ABM_2	$1 + \alpha_1 + \alpha_2 - \gamma$	$(1 - 2s - \epsilon)(1 - \alpha_1 - \alpha_2 - \gamma)$
ABm_2	$1 + \alpha_1$	$(1 - 2s - \epsilon)(1 - \alpha_1)$
AbM_2, aBM_2	$1 - s - \gamma$	$1 - s - \gamma$
Abm_2, aBm_2	$1 - s$	$1 - s$
abm_2	$(1 - 2s - \epsilon)(1 - \alpha_1)$	$1 + \alpha_1$
abM_2	$(1 - 2s - \epsilon)(1 - \alpha_1 - \alpha_2 - \gamma)$	$1 + \alpha_1 + \alpha_2 - \gamma$

Table 4. Fitnesses of individual genotypes modified by a fixed niche choice parameter α_1 a polymorphic allele

M_2, which increases niche preference for the better niche by α_2, and a cost of this allele, γ.

Individual-based simulations: polygenic traits with more than 3 loci

We use individual-based simulations to illustrate that initial degree of assortment is essential

to allow for the evolution of coupling between ecological loci. When assortment has to

evolve from low frequencies, even with strong recombination (r = 0.01), selection has to be

strong for the coupling to evolve (c.f. Fig. 3 for 2 ecological loci loci, approx. s > 0.1). Weak

pre-existing niche preference (one locus out of ten with α = 0.1 is near fixation) allows for

evolution of two coexisting specialist if linkage is strong and the number of loci is low (Fig 4

A,B – top, n_{eco} = 3 and 6). The number of loci, among which coupling evolves, increases

readily with higher initial niche-recognition (Fig. 4 vs 5). For example, with two loci out of

ten with α = 0.1 near fixation, variation is already maintained for a moderate number of loci

(Fig 5), though coupling between the ecological loci (in colour) stays weak when

recombination is high (r=0.1) (Fig 5B, bottom rows). As initial assortment increases further,

conditions for coexistence broaden considerably (see Fig S4) – two nascent specialist can

then diverge and coexist even for a polygenic trait under a strong trade-off.

For more ecological loci - assuming variation is not lost faster - we see a slow build-up of

coupling between ecological loci (Fig 5C, second row). This can be seen as an abrupt phase

change (tipping point, c.f. [35]). For our scenario, however, the crucial question is whether

enough assortment can build up before variation is lost. Simply because the rate of change

of allele frequencies grows with its heterozygosity (i.e., when their frequencies are close to

½), we see a slow increase in niche recognition initially, when its allele frequencies are low

(black). It is to be noted that the niche-preference alleles are growing from low variation, not

from zero – this would need different time scales (even with high mutation of $\mu = 10^{-3}$), and

the waiting time does further narrow the parameter range for speciation towards

coexistence. Here we focus on the later stages, assuming some variation is present.

To help the curious reader to understand the patterns in relations to other changes in the

parameters, in the supplement we also include results in the absence of epistasis (when

variation is maintained more easily), stronger selection (*ditto*), and a lower population size

(converse). Curiously, even with total population size of N=10 000 and selection per locus $s_i >$

0.01 (16 loci), genetic drift is still important. Unfortunately, detailed exploration of these

patterns is beyond the scope of this paper. We provide the source code (for Mathematica,

Wolfram) and are happy to deliver further specific reference-simulations upon request.

Figure 4. Comparison of individual based simulations with variable numbers of ecological loci (columns) and

recombination between them (top two vs. bottom two rows). With moderate initial niche recognition,

associations (LD) build up among multiple loci under disruptive ecological selection, and both niche recognition

and ecological divergence increase jointly. In contrast, when initial niche recognition is weak, variation is

quickly lost (c.f. Figure 5). First row shows the change in allele frequencies for ecological loci (in colour) and

niche preference loci (black), second row shows the dynamics of the pairwise linkage disequilibria LD between

ecological loci (colour), niche preference loci (black) and between ecological and niche preference loci (grey),

assuming strong linkage (r = 0.01). Notice that for low number of loci (strong selection), LD starts building up

immediately; whereas for higher number of loci LD only starts to build up after niche preference increases. For

even higher number of loci (and/or recombination), variation is lost before assortment builds up. The quick loss

of variation is facilitated by epistasis (c.f. Figure S5), genetic drift (c.f. Figure S6,S7) and recombination. Third

and fourth rows depicts allele frequencies and LDs when loci are only weakly linked, r = 0.1, where joint

diversification and evolution of niche preference only occurs for lower number of loci (stronger selection). Due

to epistasis, loss of variation is actually faster for 12 than 16 loci. Note that both selection and epistasis are

normalized with respect to the baseline two-locus case, which leads to a slight flattening of the trade-off curve

as the number of loci increases (see Model and Figure S1). Parameters: s = 0.1, ε = - 0.05, slightly asymmetric

niche sizes with c = 0.46, population size across niches N = 10 000. 10 niche preference loci with alpha = 0.1

(average initial niche preference for the specialist is $\mathcal{A}$ = 0.22 for 2 niche-preference locus near fixation.

Selection per locus is 0.1, 0.067, 0.033, 0.017, 0.0125 and pairwise epistasis is 0.05, 0.017, 0.0033, 0.00075,

0.00042 for 2,3,6,12 and 16 loci, respectively (2 loci are the reference value).

Figure 5. When initial niche preference is weak, divergence is only possible if ecological trait is determined by

few loci, with strong selection per locus relative to recombination. With more than 6 loci (selection per locus

weaker than 0.03), variation is quickly lost even when linkage is strong (r=0.01, top rows). When linkage is

weak (bottom rows, r=0.1), variation is lost already for 3 loci (s = 0.67) but not 2 loci (s=0.1; not shown).

Parameters are the same as in Figure 4 except here, only one of the niche-preference loci is near fixation

initially, giving average initial niche preference for the specialist $\mathcal{A}$ = 0.135.

Figure 4.

Figure 5.

Discussion

Maintenance of polymorphism and the plausibility of sympatric speciation are some of the most persistent questions in evolutionary biology. An arguable assumption frequently made in many sympatric speciation models is their restriction to one or rarely several loci, whereas ecological adaptation of a population often involves a gradual change in a polygenic trait [36]. An obvious reason for this common restriction is the analytical difficulty of such models, especially if additional evolutionary forces such as epistasis and a nonlinear trade-off between viabilities are included. Notably, the multilocus version of Levene's model – with a trade-off in adaptation to two different niches – has been analysed [37,38] but no epistasis was allowed (linear trade-off). An exception is Barton's [39] analyses of adaptation in a quantitative trait under a strong (convex) trade-off. He assumed the symmetries in allele frequencies were maintained under disruptive and negative frequency dependent selection [40]. Yet, when niches are not perfectly symmetric, independent regulation within niches does not always generate sufficient negative frequency-dependence to stabilise ecological polymorphism. Evolution of a quantitative trait in a Levene's model with soft selection has been also addressed using adaptive dynamics, most notably by Kisdi and Geritz [41]. They show that successive invasions and substitutions of new alleles of small effect lead to a stable polymorphism, even when niches are considerably asymmetric. Yet, branching cannot be equated with speciation. The question we are asking here is: will these branches diverge and then remain stable in the presence of recombination?

In order for specialists to be favoured over generalists in the absence of niche choice, trade-off in fitness between niches must be strong, such that a small change of the fitness in one habitat leads to a large change of fitness on the other (the fitness curve is convex). For a polygenic trait, this implies less-than-additive epistasis and/or heterozygote disadvantage. Yet, most early models of sympatric speciation focused on maintenance of polymorphism at a single diploid locus under heterozygote advantage (concave trade-off), and on coupling of such polymorphism with a locus for assortment [8,32,42]. Felsenstein [7] analysed coupling of two loci under an ecological trade-off with an independent self-recognising locus for assortment and concluded that assortment can only increase for linear or concave (weak)

trade-offs. With negative epistasis (strong, convex trade-off), polymorphism was not

maintained and assortment could not evolve ([7] , p. 130). It has been recognised that

asymmetry in niche proportions significantly influences the ability of these models to

maintain genetic polymorphism – and that symmetric selection and concave trade-offs in

fitness make such maintenance easier [7,8,16,26]. Curiously, Barton [39] showed that

assortment can readily increase in a population under a convex trade-off, provided the

starting genetic variance is large enough. However, that model assumed symmetric niches,

and that symmetry in allelic frequencies is maintained by frequency-dependent selection

(hypergeometric model, [40]).

We show that specialists can readily evolve even when the trade-off in fitness between

niches is strong, provided that polymorphism is stabilised by assortment arising from a

preference for the "correct" niche (to which the individual is better adapted), and there is

some initial niche-recognition. As the number of loci determining the ecological trait

increases, higher initial assortment is necessary to facilitate divergence. When the number

of loci determining the ecological trait is small, and selection is large relative to

recombination, associations (linkage disequilibria - LD) between ecological loci increase

readily (c.f. [43]). With a higher number of loci (weaker per locus selection), initial niche

recognition is essential to prevent loss of the variation driven by the negative epistasis

(strong trade-off). This limits the scope for lingering before linkage disequilibrium starts

building up (tipping points, as popularized by [35]). If selection is large relative to

recombination, associations among ecological loci start building up straight away. As

selection weakens relative to negative epistasis (i.e. disruptiveness of selection), variability

gets quickly lost. Hence in the presence of epistasis, we only observe the so-called tipping

points in the build-up of LD for moderate number of loci, not too strong recombination and

moderate initial assortment (see Figure 4C).

In our analysis, we have neglected unprotected polymorphism. It has been shown that in

Levene's model with soft selection, as long as fitness is not a linear function of allele

frequency, there are 'extra' unprotected equilibria, reachable only from surrounding allele

frequencies [44]. Independently, Novak [45] showed that up to three polymorphic equilibria

may coexist between 2 demes for 1 diploid bi-allelic locus (two of them unprotected). While

we would have missed such unprotected equilibria by only testing for instability of

monomorphism, our extensive numerical analysis and simulations do not suggest that these

are biologically relevant in our model system (indeed, the population state space around

such unprotected equilibria is expected to be very narrow [45]).

Prezygotic isolation due to pre-existing niche recognition both speeds up further evolution

of niche preference and allows ecological polymorphism to increase. As it stabilises

polymorphism in divergent populations under secondary contact, it is replacing the previous

assortment due to geographic isolation. Similar reduction of gene flow can arise via

philopatry (learned habitat preference) – but as the association with fitness is indirect, the

effect is likely to be weaker [46,47].

We argue that niche recognition and preference is widespread across all life forms. Even

plants' roots have growth oriented towards nutrient-rich parts of the rhizosphere [48],

motile unicellular organisms, including even bacteria, can move towards their preferred food

source by chemotaxis [49,50] and among (in)vertebrates it is generally known that they can

recognise and choose their preferred food or habitat. Therefore, this pre-existing ability to

recognise the preferred niche may in fact "hijack" arising ecological divergence, protect it,

and even further reinforce the newly arising reproductive isolation by strengthening the

assortment. Furthermore, the degree of niche preference does not necessarily tightly

correlate with the fitness advantage, which the niche provides to the organism. For example,

bacteria and yeast tend to prefer one food source over all others (a phenomenon known as

a diauxic shift), which in fact means a 100% preference even if the fitness advantage in

growth rate is only minor [51,52]. In a phytophagous butterfly *Melitaea cinxia,* host plant

preference has been observed but no measurable fitness advantage was detected on the

more preferred host compared to the less preferred one [53]. In many other cases a

preference-performance correlation has been detected [24,25]. While the genetic basis of

preference remains mostly unclear, our model is consistent with many observations in

ecology where animals tend to choose a diet, which is optimal for them [54].

Acknowledgement

We would like to thank Nick Barton, Joachim Hermisson, Stuart Baird, Tiago Paixão and

Tadeas Priklopil for their insightful comments on parts and/or earlier stages of this study.

Also, we are grateful for many valuable comments and suggestions from the reviewers

during the peer review process. PP was supported by a bilateral Czech-Austrian program

Aktion, and both JP and PP were supported by Austrian National Agency (FWF P-32896B).

References

1.   Darwin C. 1859 *On the origin of species by means of natural selection, or the*
preservation of favoured races in the struggle for life. London: John Murray.

2.   Mayr E. 1963 *Animal species and evolution*. Cambridge, MA: The Belknap Press of
Harvard University Press.

3.   Jiggins CD. 2006 Sympatric speciation: why the controversy? *Curr Biol* **16**, R333-334.
(doi:10.1016/j.cub.2006.03.077)

4.   Bolnick DI, Fitzpatrick BM. 2007 Sympatric speciation: models and empirical evidence.
Annu Rev Ecol Evol Syst **38**, 459–487. (doi:10.1146/annurev.ecolsys.38.091206.095804)

5.   Via S. 2001 Sympatric speciation in animals: the ugly duckling grows up. *Trends in*
Ecology & Evolution **16**, 381–390.

6.   Dobzhansky Th. 1940 Speciation as a stage in evolutionary divergence. *The American*
Naturalist **74**, 312–321.

7.   Felsenstein J. 1981 Skepticism towards santa rosalia, or why are there so few kinds of
animals? *Evolution* **35**, 124–138. (doi:10.2307/2407946)

8.   Maynard-Smith JM. 1966 Sympatric speciation. *The American Naturalist* **100**, 637–650.

9.   Udovic D. 1980 Frequency-dependent selection, disruptive selection, and the evolution
of reproductive isolation. *The American Naturalist* **116**, 621–641.

10. Gavrilets S. 2003 Perspective: models of speciation: what have we learned in 40 years?
Evolution **57**, 2197–2215.

11. Barton NH, de Cara MAR. 2009 The evolution of strong reproductive isolation. *Evolution*
63, 1171–1190. (doi:10.1111/j.1558-5646.2009.00622.x)

12. Berlocher SH, Feder JL. 2002 Sympatric speciation in phytophagous insects: moving
beyond controversy? *Annu Rev Entomol* **47**, 773–815.
(doi:10.1146/annurev.ento.47.091201.145312)

13. Greenwood PJ. 1980 Mating systems, philopatry and dispersal in birds and mammals.
Animal Behaviour **28**, 1140–1162. (doi:10.1016/S0003-3472(80)80103-5)

14. Lavagnino N, Serra F, Arbiza L, Dopazo H, Hasson E. 2012 Evolutionary genomics of genes
involved in olfactory behavior in the drosophila melanogaster species group.
Evolutionary Bioinformatics (doi:10.4137/EBO.S8484)

15. Rice WR, Salt GW. 1990 The evolution of reproductive isolation as a correlated character
under sympatric conditions: experimental evidence. *Evolution* **44**, 1140–1152.
(doi:10.1111/j.1558-5646.1990.tb05221.x)

16. Ravigné V, Dieckmann U, Olivieri I. 2009 Live where you thrive: joint evolution of habitat
choice and local adaptation facilitates specialization and promotes diversity. *The*
American Naturalist **174**, E141–E169. (doi:10.1086/605369)

17. Butlin R. 1987 A new approach to sympatric speciation. *Trends in Ecology & Evolution* **2**,
310–311. (doi:10.1016/0169-5347(87)90085-1)

18. Jaenike J, Holt RD. 1991 Genetic variation for habitat preference: evidence and
explanations. *The American Naturalist* **137**, S67–S90. (doi:10.1086/285140)

19. Via S. 1991 The genetic structure of host plant adaptation in a spatial patchwork:
demographic variability among reciprocally transplanted pea aphid clones. *Evolution* **45**,
827–852. (doi:10.1111/j.1558-5646.1991.tb04353.x)

20. Mackenzie A. 1996 A trade-off for host plant utilization in the black bean aphid, Aphis
fabae. *Evolution* **50**, 155–162. (doi:10.1111/j.1558-5646.1996.tb04482.x)

21. Sezer M, Butlin RK. 1998 The genetic basis of host plant adaptation in the brown
planthopper (Nilaparvata lugens). *Heredity* **80**, 499–508. (doi:10.1046/j.1365-
2540.1998.00316.x)

22. Rossi AM, Stiling P, Cattell MV, Bowdish TI. 1999 Evidence for host-associated races in a
gall-forming midge: trade-offs in potential fecundity. *Ecological Entomology* **24**, 95–102.
(doi:10.1046/j.1365-2311.1999.00164.x)

23. Craig TP, Horner JD, Itami JK. 2007 Genetics, experience, and host-plant preference in
Eurosta solidaginis: implications for host shifts and speciation. *Evolution* **55**, 773–782.
(doi:10.1111/j.0014-3820.2001.tb00813.x)

24. Forister ML, Dyer LA, Singer MS, Stireman JO, Lill JT. 2012 Revisiting the evolution of
ecological specialization, with emphasis on insect–plant interactions. *Ecology* **93**, 981–
991. (doi:10.1890/11-0650.1)

25. Gripenberg S, Mayhew PJ, Parnell M, Roslin T. 2010 A meta-analysis of preference–
performance relationships in phytophagous insects. *Ecology Letters* **13**, 383–393.
(doi:10.1111/j.1461-0248.2009.01433.x)

26. Christiansen FB. 1975 Hard and soft selection in a subdivided population. *The American*
Naturalist **109**, 11–16.

27. Ravigné V, Olivieri I, Dieckmann U. 2004 Implications of habitat choice for protected
polymorphisms. *Evolutionary Ecology Research* , 125–145.

28. Rieseberg LH. 2001 Chromosomal rearrangements and speciation. *Trends in Ecology &*
Evolution **16**, 351–358. (doi:10.1016/s0169-5347(01)02187-5)

29. Levene H. 1953 Genetic equilibrium when more than one ecological niche is available.
The American Naturalist **87**, 331.

30. Deakin MAB. 1966 Sufficent conditions for genetic polymorphism. *The American*
Naturalist **100**, 690–692. (doi:10.1086/282462)

31. Prout T. 1968 Sufficient conditions for multiple niche polymorphism. *The American*
Naturalist **102**, 493–496.

32. Hoekstra RF, Bijlsma R, Dolman AJ. 1985 Polymorphism from environmental
heterogeneity: models are only robust if the heterozygote is close in fitness to the
favoured homozygote in each environment. *Genetics Research* **45**, 299–314.
(doi:10.1017/S001667230002228X)

33. Gliddon C, Strobeck C. 1975 Necessary and sufficient conditions for multiple-niche
polymorphism in haploids. *The American Naturalist* **109**, 233–235.

34. de Meeûs T, Michalakis Y, Renaud F, Olivieri I. 1993 Polymorphism in heterogeneous
environments, evolution of habitat selection and sympatric speciation: Soft and hard
selection models. *Evol Ecol* **7**, 175–198. (doi:10.1007/BF01239387)

35. Nosil P, Feder JL, Flaxman SM, Gompert Z. 2017 Tipping points in the dynamics of
speciation. *Nat Ecol Evol* **1**, 0001. (doi:10.1038/s41559-016-0001)

36. Wilson DS, Turelli M. 1986 Stable underdominance and the evolutionary invasion of
empty niches. *The American Naturalist* **127**, 835–850. (doi:10.1086/284528)

37. Nagylaki T. 2009 Evolution under the multilocus Levene model without epistasis. *Theor*
Popul Biol **76**, 197–213. (doi:10.1016/j.tpb.2009.07.003)

38. Bürger R. 2010 Evolution and polymorphism in the multilocus Levene model with no or
weak epistasis. *Theor Popul Biol* **78**, 123–138. (doi:10.1016/j.tpb.2010.06.002)

39. Barton NH. 2010 What role does natural selection play in speciation? *Philos Trans R Soc*
B **365**, 1825–1840. (doi:10.1098/rstb.2010.0001)

40. Shpak M, Kondrashov AS. 1999 Applicability of the hypergeometric phenotypic model to
haploid and diploid populations. *Evolution* **53**, 600–604. (doi:10.1111/j.1558-
5646.1999.tb03794.x)

41. Kisdi É, Geritz SAH. 1999 Adaptive dynamics in allele space: evolution of genetic
polymorphism by small mutations in a heterogeneous environment. *Evolution* **53**, 993–
1008. (doi:10.1111/j.1558-5646.1999.tb04515.x)

42. Maynard-Smith J, Hoekstra R. 1980 Polymorphism in a varied environment: how robust
are the models? *Genetical Research* **35**, 45–57. (doi:10.1017/S0016672300013926)

43. Barton NH. 1983 Multilocus clines. *Evolution* **37**, 454. (doi:10.2307/2408260)

44. Priklopil T. 2012 On invasion boundaries and the unprotected coexistence of two
strategies. *J Math Biol* **64**, 1137–1156. (doi:10.1007/s00285-011-0448-y)

45. Novak S. 2011 The number of equilibria in the diallelic Levene model with multiple
demes. *Theor Popul Biol* **79**, 97–101. (doi:10.1016/j.tpb.2010.12.002)

46. Beltman JB, Haccou P, Cate CT. 2004 Learning and Colonization of New Niches: A First
Step toward Speciation. *Evolution* **58**, 35–46.

47. Beltman JB, Metz JAJ. 2005 Speciation: more likely through a genetic or through a
learned habitat preference? *Proc R Soc B* **272**, 1455–1463. (doi:10.1098/rspb.2005.3104)

48. Wells DM, French AP, Naeem A, Ishaq O, Traini R, Hijazi H, Bennett MJ, Pridmore TP.
2012 Recovering the dynamics of root growth and development using novel image
acquisition and analysis methods. *Philos Trans R Soc B* **367**, 1517–1524.
(doi:10.1098/rstb.2011.0291)

49. Arkowitz RA. 1999 Responding to attraction: chemotaxis and chemotropism in
Dictyostelium and yeast. *Trends Cell Biol.* **9**, 20–27. (doi:10.1016/s0962-8924(98)01412-
3)

50. Xie L, Altindal T, Chattopadhyay S, Wu X-L. 2011 Bacterial flagellum as a propeller and as
a rudder for efficient chemotaxis. *PNAS* **108**, 2246–2251.
(doi:10.1073/pnas.1011953108)

51. Berthels NJ, Cordero Otero RR, Bauer FF, Thevelein JM, Pretorius IS. 2004 Discrepancy in
glucose and fructose utilisation during fermentation by Saccharomyces cerevisiae wine
yeast strains. *FEMS Yeast Res* **4**, 683–689. (doi:10.1016/j.femsyr.2004.02.005)

52. Bren A, Park JO, Towbin BD, Dekel E, Rabinowitz JD, Alon U. 2016 Glucose becomes one
of the worst carbon sources for E.coli on poor nitrogen sources due to suboptimal levels
of cAMP. *Sci Rep* **6**, 1–10. (doi:10.1038/srep24834)

53. Nouhuys SV, Singer MC, Nieminen M. 2003 Spatial and temporal patterns of caterpillar
performance and the suitability of two host plant species. *Ecological Entomology* **28**,
193–202. (doi:10.1046/j.1365-2311.2003.00501.x)

54. Sinervo B. 1997 Optimal foraging theory: constraints and cognitive processes. *Behavioral
ecology* , 105–130.