Dynamics of T cell activation mediated by eIF3 interactions with T cell receptor mRNAs

Dasmanthie De Silva[1,2], Lucas Ferguson[1], Benjamin E. Smith[3], Grant H. Chin[1], Ryan A.

Apathy[4], Theodore L. Roth[4], Marek Kudla[1], Alexander Marson[4,5,6,7,8,9], Nicholas T.

Ingolia[1,10], Jamie H. D. Cate[1,2,9,11,12]*

1. Department of Molecular and Cell Biology, University of California-Berkeley, Berkeley, California 94720, United States.
2. The J. David Gladstone Institutes, San Francisco, CA 94158 USA
3. School of Optometry, University of California, Berkeley, Berkeley, CA 94720, USA.
4. Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, CA, USA.
5. Diabetes Center, University of California, San Francisco, San Francisco, CA, USA.
6. Chan Zuckerberg Biohub, San Francisco, CA, USA
7. Department of Medicine, University of California, San Francisco, San Francisco, CA, USA.
8. Parker Institute for Cancer Immunotherapy, San Francisco, CA, USA.
9. Innovative Genomics Institute, University of California, Berkeley, Berkeley, CA, USA.
10. California Institute for Quantitative Biosciences, University of California, Berkeley, Berkeley, United States.
11. Department of Chemistry, University of California-Berkeley, Berkeley, California 94720, United States.
12. Molecular Biophysics and Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, United States.

* Corresponding author: jcate@lbl.gov

Abstract

Activation of T cells requires a global surge in cellular protein synthesis, which is

accompanied by a large increase in translation initiation[1–4]. A central component of the

translation initiation machinery–the multi-subunit eukaryotic initiation factor 3 (eIF3)–is

rapidly turned on when quiescent T cells are stimulated[3]. However, the precise role eIF3

plays in activated T cells is not known. Using a global transcriptome cross-linking

approach, we show that human eIF3 interacts with a distinct set of mRNAs in activated

Jurkat cells. A subset of these mRNAs, including those encoding the T cell receptor (TCR)

subunits TCRA and TCRB, crosslink to eIF3 across the entire length of the mRNA.

Fluorescence in situ hybridization studies reveal both *TCRA* and *TCRB* mRNAs form

granules, potentially acting as translation "hot-spots." We further find that elements in the

3'-untranslated regions (3'-UTRs) of the *TCRA* and *TCRB* mRNAs that directly interact

with eIF3 control the early dynamics of TCR translation and the extent of T cell activation.

These results highlight a new role for eIF3 in regulation of translation dynamics and

provide insights that can guide engineering of T cells for cell immunotherapy applications.

Main

Several lines of evidence indicate eIF3 serves specialized roles in cellular translation, by recognizing specific RNA structures in the 5'-untranslated regions (5'-UTRs) of target mRNAs[5], binding the 7-methyl-guanosine (m^7G) cap[6] or through interactions with *N*-6-methyl-adenosine (m^6A) post-transcriptional modifications in mRNAs[7]. Binding to these *cis*-regulatory elements in mRNA can lead to translation activation or repression, depending on the RNA sequence and structural context[5,7,8]. These functions for eIF3 can aid cell proliferation[5], or allow cells to rapidly adapt to stress such as heat shock[7]. Additionally, eIF3 plays an important role in the development of specific tissues[9–11].

In the immune system, T cell activation involves a burst in translation[1–4] and correlates with assembly of subunit EIF3J with the main eIF3 complex[3]. However, whether eIF3 serves a general or more specific role in T cell activation is not known. To delineate how eIF3 contributes to T cell activation, we first identified mRNAs that directly interact with eIF3 in Jurkat cells activated for 5 hours with ionomycin and phorbol 12-myristate 13-acetate (I+PMA), or in non-activated Jurkat cells as a control, using photoactivatable ribonucleoside-enhanced crosslinking and immunoprecipitation (PAR-CLIP)[5,12,13] (**Fig. 1a**). We used Jurkat cells as a model for T cells, as PAR-CLIP experiments require a large number of cells labeled with 4-thiouridine at a non-toxic concentration[14]. Jurkat cells also have a defined T cell receptor and transcriptome, avoiding the donor-to-donor variability of primary T cells. In the Jurkat PAR-CLIP experiments, RNA crosslinked to eight of the thirteen eIF3 subunits, as identified by mass spectrometry: subunits EIF3A, EIF3B, EIF3D and EIF3G as seen in HEK293T cells[5], as well as subunits EIF3C, EIF3E,

EIF3F, and EIF3L (**Fig. 1b, Extended Data Fig. 1, Supplementary Table 1**). In activated

Jurkat cells, eIF3 crosslinked to a substantially larger number of mRNAs compared to the

non-activated cells (**Extended Data Figs. 2 and 3, Supplementary Tables 2 and 3**).

Notably, eIF3 interacted with a completely new suite of mRNAs in activated Jurkat cells,

compared to those identified in HEK293T cells (**Fig 1c, Extended Data Fig. 4,**

Supplementary Tables 4 and 5). These mRNAs are enriched for those encoding

proteins central to immune cell function (**Supplementary Table 6**). Moreover, eIF3

crosslinking does not correlate with mRNA abundance (**Extended Data Fig. 4**),

suggesting that eIF3 is involved in specific regulation of T cell activation.

In activated Jurkat cells, eIF3 crosslinked to mRNAs by a multitude of patterns (**Fig 2a,**

b, Extended Data Fig. 5), consistent with varied roles for eIF3 in T cell activation and

function. Many of the mRNAs have a single PAR-CLIP site in the 5'-UTR as observed in

HEK293T cells[5] (**Fig. 2a**, **Extended Data Figs. 2, Supplementary Table 6**). However,

eIF3 crosslinked to some mRNAs across their entire length, from the beginning of the 5'-

UTR through the 3'-UTR (**Fig. 2a, b, Extended Data Fig. 5**). This "pan-mRNA" pattern of

eIF3 crosslinking–which includes polyadenylated mRNAs as well as histone mRNAs–has

not been observed before. Notably, a number of these mRNAs encode proteins important

for T cell activation, including both the alpha and beta subunits of the TCR, subunits TCRA

and TCRB (**Fig. 2b**, **Supplementary Table 6**).

Crosslinking in PAR-CLIP experiments requires direct interaction between the RNA and

protein of interest[14]. Thus, the pan-mRNA pattern of crosslinking between eIF3 and

certain mRNAs suggests formation of ribonucleoprotein complexes (RNPs) highly

enriched in eIF3. Remarkably, the pan-mRNA crosslinking pattern in the *TCRA* and *TCRB*

mRNAs occurs in activated but not non-activated Jurkat cells. As previously known RNPs

such as processing bodies (P bodies) and stress granules contain translationally

repressed mRNAs, we performed fluorescence in situ hybridization (FISH) on both *TCRA*

and *TCRB* mRNAs in activated Jurkat cells. Since Jurkat cells have a defined TCR, we

were able to design FISH probes across the entire *TCRA* and *TCRB* transcripts. We also

probed for P body and stress granule proteins DCP1 and G3BP1, respectively[15–18]. Both

the *TCRA* and *TCRB* mRNAs formed distinct puncta, but did not co-localize with either P

bodies or stress granules, or with each other (**Fig. 2c, Extended Data Fig. 6**). Taken

together, these results suggest that the *TCRA* and *TCRB* mRNA puncta represent eIF3-

mRNA concentrated regions that may act as translation "hot spots."

Both the *TCRA* and *TCRB* mRNAs include prominent sites of eIF3 crosslinking in their

respective 3'-UTR elements. To test the functional importance of these sites of eIF3

interaction, we used CRISPR-Cas9 genome editing with guide RNAs (gRNAs) targeted

to the two boundaries of the eIF3 PAR-CLIP sites in each of the 3'-UTRs of *TCRA* and

TCRB mRNA (**Extended Data Fig. 7, Supplementary Table 7**). We nucleofected these

pairs of gRNA/Cas9 RNPs into primary human T cells from two donors, to delete the

entire PAR-CLIP site in each 3'-UTR. PCR analysis showed successful deletion of the

eIF3 PAR-CLIP site in the *TCRA* 3'-UTR or in the *TCRB* 3'-UTR (*TCRA ΔPAR* or *TCRB*

ΔPAR, respectively) in approximately 50-60% of the cells (**Extended Data Fig. 7**). For

comparison with the *TCRA ΔPAR* or *TCRB ΔPAR* cell populations, we edited cells from

both donors using each gRNA targeting the *TCRA* 3'-UTR and *TCRB* 3'-UTR region

separately (single gRNA experiments), a gRNA targeting the coding sequence (CDS) of

TCRA which knocks out TCR expression with high efficiency (Δ*TCR*)[19], a scrambled

gRNA (SC) which does not cut anywhere in the human genome, and cells mixed with the

gRNA/Cas9 RNPs but not nucleofected.

We first measured total TCR protein levels in both the *TCRA ΔPAR* and *TCRB ΔPAR* cell

populations by western blot, after different times following T cell stimulation using anti-

CD3 and anti-CD28 antibodies (**Fig. 3a**). We used an anti-TCRA antibody to measure

TCR levels in both *TCRA ΔPAR* and *TCRB ΔPAR* cells, since the formation of an intact

TCR is required to stabilize both the TCRA and TCRB subunits[20,21]. In both *TCRA ΔPAR*

and *TCRB ΔPAR* cell populations, TCRA protein levels were nearly absent or clearly

reduced at early time points after activation (~1 hr). By contrast, SC control cells–which

should behave as wild-type T cells–exhibited a substantial burst in TCRA protein levels

immediately after activation (**Fig. 3a**). Only at later time points did TCRA levels in the

TCRA ΔPAR and *TCRB ΔPAR* cell populations begin to increase. Next we checked

whether the reduced TCR protein levels in *TCRA ΔPAR* and *TCRB ΔPAR* cell populations

at early time points after activation with anti-CD3 and anti-CD28 antibodies affected the

rate of TCR clustering, as occurs in immune synapse formation[22]. We used the same cell

populations as those for western blot analysis (**Fig. 3a**), to correlate the TCRA protein

levels observed in the western blots with TCR cluster formation. Using TCR protein

staining, we observed a decrease in the proportion of cells that formed TCR clusters in

the *TCRA ΔPAR* and *TCRB ΔPAR* cell populations compared to SC control cells, with

more severe defects seen at the earlier time points (**Fig. 3b**). We also stained cells with

a fluorescently tagged anti-TCR antibody and used flow cytometric analysis to measure

the levels of TCR on the cell surface with and without activation with I+PMA (**Fig. 3c, d,**

Extended Data Fig. 8). A substantially lower percentage of the *TCRA ΔPAR* and *TCRB*

ΔPAR cell populations expressed TCR on the cell surface compared to SC control cells

or cells edited with a single gRNA (**Fig. 3c, d, Extended Data Fig. 8**). Taken together,

these results suggest that regulation of TCR protein synthesis mediated by the *TCRA* and

TCRB mRNA 3'-UTR elements that interact with eIF3 is required for the robust TCR

clustering that is central to immune synapse formation[22].

To test whether the defect in TCR clustering in the *TCRA ΔPAR* and *TCRB ΔPAR* cell

populations reflects a general deficiency in T cell activation, we measured the T cell

activation markers CD69 and CD25 by flow cytometry and secreted cytokines IL2 and

IFNγ using ELISA (**Fig. 4, Extended Data Figs. 9–12**). Both the CD4+ and CD8+

ΔTCR cell populations failed to express significant amounts of CD69 and CD25 (**Fig.**

4a,b, Extended Data Fig. 10, Extended Data Fig. 11), and the *ΔTCR* cells only weakly

upregulated IL2 and IFNγ, particularly at the 24 hour time point (**Fig. 4c, Extended**

Data Fig. 12). Notably, fewer cells in the *TCRA ΔPAR* and *TCRB ΔPAR* CD4+ and

CD8+ cell populations expressed CD69 after 5 and 8 hours after activation, compared

to SC control CD4+ and CD8+ cells (**Fig. 4a,b, Extended Data Fig. 10, Extended Data**

Fig. 11). At later time points, fewer cells in the *TCRA ΔPAR* and *TCRB ΔPAR* cell

populations expressed both CD69 and CD25 activation markers compared to SC

control cells. Furthermore, the *TCRA ΔPAR* and *TCRB ΔPAR* T cell populations

secreted lower amounts of cytokines IL2 and IFNγ, compared to the SC control cells

(**Fig. 4c, Extended Data Fig. 12**). Taken together, the *TCRA ΔPAR* and *TCRB ΔPAR*

primary T cell populations exhibited activation defects at early and later time points, and

align with the decreased TCR levels observed by western blot, detected by TCR

clustering and by flow cytometry analyses (**Fig. 3**), consistent with eIF3-mediated

regulation of T cell activation through interactions with the *TCRA* and *TCRB* mRNA 3'-

UTR PAR-CLIP sites, in both CD4+ and CD8+ T cells.

To obtain a deeper mechanistic understanding of the role of eIF3 in the regulation of

TCRA and TCRB translation, we recreated the deletions of the eIF3 PAR-CLIP sites in

both the *TCRA* and *TCRB* mRNA 3'-UTRs (**Extended Data Fig. 7**) in Jurkat cells, which

can be clonally propagated for detailed analysis. As observed in edited primary T cells,

TCRA ΔPAR and *TCRB ΔPAR* Jurkat cell populations expressed lower levels of the TCR

proteins compared to wild-type (WT) cells, and failed to show a burst in TCR expression

at early time points after anti-CD3 and anti-CD28 activation (**Extended Data Fig. 13a**).

Notably, *TCRA* and *TCRB* mRNA levels were unaffected or even increased in the *TCRA*

ΔPAR and *TCRB ΔPAR* Jurkat cells (**Extended Data Fig. 13b, c**). Importantly, the

amount of *TCRA* and *TCRB* mRNAs interacting with eIF3 was higher in WT cells than in

the *TCRA ΔPAR* and *TCRB ΔPAR* cells after activation (**Extended Data Fig. 14**),

indicating that deleting the eIF3 PAR-CLIP sites disrupts eIF3 interactions with these

mRNAs. Additionally, the levels of eIF3-mRNA interaction closely mirrored the total *TCRA*

protein levels at both 5 hours and 8 hours for WT, *TCRA ΔPAR* and *TCRB ΔPAR* cells,

suggesting that eIF3 interactions with the 3'-UTR sequences in the *TCRA* and *TCRB*

mRNAs is tightly coupled with *TCRA* and *TCRB* protein levels. The *TCRA ΔPAR* and

TCRB ΔPAR Jurkat cells also showed a delay in TCR clustering (**Extended Data Fig.**

15), had lower cell populations expressing CD69 and CD25, and secreted lower amounts

of IL2 compared to WT cells (**Extended Data Fig. 16**), mirroring the results obtained in

primary T cells (**Fig. 4, Extended Data Fig. 10-12**). These results verify that Jurkat cells

recapitulate the response of primary T cells, and further support the model that eIF3

interactions with the PAR-CLIP sites in the *TCRA* and *TCRB* mRNA 3'-UTRs mediate

translational regulation at early times after TCR activation.

In order to understand the functional role of eIF3 interactions with the 3'-UTRs of the

TCRA and *TCRB* mRNAs, we constructed nanoluciferase reporters with the 5'-UTR of

the human beta globin gene (*HBB*) and a PEST destabilization domain. The PEST

domain reduces nanoluciferase protein half-life[23,24] and provides better time resolution of

nanoluciferase expression after T cell activation. The reporter coding sequence (CDS)

was then fused to the wild-type *TCRA* or *TCRB* mRNA 3'-UTR sequence (WT), or 3'-UTR

with the eIF3 PAR-CLIP site deleted (*ΔPAR*), or 3'-UTR with the reversed sequence of

the eIF3 PAR-CLIP site (R*PAR, i.e. 5'-3' sequence reversed to 3'-5' direction) (**Fig. 5a**),

and stably transduced into Jurkat cells using a lentiviral vector. Cells expressing

nanoluciferase from reporters with a deletion or reversal of the eIF3 PAR-CLIP site

sequence led to reduction of the luminescence signal compared to cells expressing

nanoluciferase with the WT 3'-UTR sequences (**Fig. 5b**). Furthermore, similar to the

TCRA ΔPAR and *TCRB ΔPAR* Jurkat cell lines, there was no significant defect in the

TCRA ΔPAR and *TCRA* R*PAR nanoluciferase mRNA levels compared to intact *TCRA*

3'-UTR mRNA, or for *TCRB ΔPAR* and R*PAR nanoluciferase mRNA levels compared to

intact *TCRB* 3'-UTR mRNA through 5 hrs, when the biggest defect in luciferase translation

is observed (**Fig. 5c**)**.** Immunoprecipitation of eIF3 followed by qRT-PCR quantification of

nanoluciferase mRNA showed that less nanoluciferase mRNA was bound to eIF3 when

the 3'-UTR PAR-CLIP site was either deleted or reversed, compared to nanoluciferase

mRNAs carrying the WT *TCRA* or *TCRB* 3'-UTR (**Fig. 5d**). These results support the

model that eIF3 binding to the PAR-CLIP sites in the *TCRA* and *TCRB* mRNA 3'-UTRs is

necessary for eIF3-mediated temporal regulation of TCRA and TCRB translation in

activated T cells.

Recent experiments indicate that T cells must cross a "threshold" of T cell receptor

signaling to commit to activation and proliferation[25,26], setting up a "digital" response to

antigen recognition[25–28]. The response threshold involves integration of intensity and

duration of TCR signaling[25,26,28], and spans a wide range of TCR antigen affinity[25–28].

Notably, T cell commitment to clonal expansion and differentiation can occur with as little

as 1 to 2 hours of TCR stimulation for effector CD4+ and naive CD8+ T cells[29,30]. This

time period spans the burst in TCR protein synthesis mediated by eIF3 interactions with

the *TCRA* and *TCRB* 3'-UTR elements (**Fig. 3a**). Naive CD4+ T cells require a longer

duration of TCR signaling of ~20 hours[29,31]. Although we were not able to distinguish

levels of TCR translation in isolated CD8+ and CD4+ cells (**Fig. 3**), subsequent events in

T cell activation including CD69 and CD25 expression, and IL2 and IFNγ secretion, were

equally affected in CD8+ and CD4+ cells in which the 3'-UTR PAR-CLIP sites were

deleted (**Fig. 3, Fig. 4**). Taken together, our results indicate eIF3 controls *TCRA* and

TCRB mRNA translation during the first critical hours after TCR stimulation that lead to

subsequent T cell commitment to proliferation and differentiation. These results also

suggest that eIF3-responsive mRNA 3'-UTR elements could be used to improve chimeric

antigen receptor (CAR) expression and CAR-T cell responsiveness[32,33].

The eIF3 PAR-CLIP experiment we present here provides a snapshot of eIF3-RNA

interactions that occur at the time TCR translation is most sensitive to eIF3 regulation (5

hours in Jurkat cells, **Fig. 2b**, **Extended Data Fig. 13**). At this point in time, eIF3

crosslinks to multiple mRNAs encoding proteins involved in immune cell function

(**Supplementary Table 6**). Interestingly, the patterns of eIF3 crosslinking, which for a

number of mRNAs include interactions with the protein coding sequence and 3'-UTR (**Fig.**

2, **Extended Data Fig. 5b**), suggest an active role for eIF3 in promoting translation of

these mRNAs. In support of this model, the two examples of pan-mRNAs we examined

here (*TCRA* and *TCRB*) reside in puncta distinct from P bodies and stress granules (**Fig.**

2c). However, this eIF3-mediated translation activation is transient, lasting only 1-2 hours

in primary T cells (**Fig. 3b**, **Extended Data Fig. 13a**). Recent evidence suggests that eIF3

can remain associated with translating ribosomes[34,35,36] a phenomenon that may be

enhanced for the pan-mRNAs identified here. Notably, additional layers of translation

regulation also contribute to T cell function[37], particularly with respect to mTOR

signaling[3,38] and carbon metabolism[4,39]. It should now be possible to use the present

PAR-CLIP experiments to elucidate additional roles for eIF3-mediated translation

regulation and to more fully map the system-wide role of translation regulation in T cell

activation.

Figures

Figure 1. PAR-CLIP analysis of eIF3 interactions with RNAs in Jurkat cells. a,

Schematic of Jurkat cell treatment with 4-thiouridine, activation, and cell harvesting for

PAR-CLIP analysis. **b**, Phosphorimage of SDS polyacrylamide gel resolving 5′ ^{32}P-

labeled RNAs crosslinked to eIF3 subunits in activated and non-activated Jurkat cells.

c, Venn diagram of genes identified by eIF3 PAR-CLIP in the two biological replicates of

activated Jurkat cells, compared with the eIF3 PAR-CLIP mRNAs identified in HEK293T

cells. The same number of mRNAs from the Jurkat cell PAR-CLIP libraries, ranked by

total reads mapped to a given gene, are compared to the number of mRNAs identified

in the HEK293T PAR-CLIP experiments.

Figure 2. Varied mRNA crosslinking patterns to eIF3 in activated Jurkat cells. a,

Cumulative plot showing mRNA crosslinking in sample EIF3A/C/B to predominantly the

5'-UTR (n = 396, 414 mRNAs in replicates 1 and 2, respectively), and across the entire

length of some mRNAs (n = 634, 621 mRNAs). We defined mRNAs having a ratio of

normalized 5'-UTR reads divided by 3'-UTR reads of 20 or more as "5'-UTR enriched"

with respect to eIF3 crosslinking. All others were categorized as "pan-mRNAs," with

eIF3 crosslinking across the entire length of the mRNA. The cut-off value 20 is not a

sharp boundary between the two categories of mRNA (**Extended Data Fig. 5a**). The 5'-

UTR, CDS, and 3'-UTR regions are shown normalized by length. **b**, Crosslinking of eIF3

subunits EIF3A/C/B, EIF3B/C and EIF3D/L across the entire *TCRA* and *TCRB* mRNAs.

5'-UTR, coding sequence (CDS), and 3'-UTR elements for the mapped TCR genes in

Jurkat cells are shown. Counts, PAR-CLIP reads from deep sequencing library

analysis[40]. **c**, FISH analysis of *TCRA* and *TCRB* mRNAs (yellow and magenta,

respectively) and P bodies (on the left) marked by the location of DCP1 (cyan) and

stress granules (on the right) marked by the location of G3BP1 (cyan), in activated

Jurkat cells. Graphs to the right of the images indicate Pearson's correlation coefficients

(PCCs) of *TCRA* and *TCRB* mRNA with each other or with P bodies or stress granules.

Labels on the x axis are, B: tetraSpeck microsphere beads, C: activated Jurkat cells. (*n*

= 5, P <0.008, for PCC values of cells relative to bead colocalization, across all the

channels tested, using the Wilcoxon rank-sum test). Images are representative of one

experiment of the five independent experiments in the graphs.

Figure 3. TCR levels in activated primary human T cells with eIF3 3'-UTR PAR-

CLIP sites deleted. a, Western blots measuring TCRA protein levels as a function of

time after anti-CD3/anti-CD28 activation. Cell lines used are labeled on the left: *TCRA*

ΔPAR (*TCRA* gRNA 1+2), *TCRB ΔPAR* (*TCRB* gRNA 1+2), and SC (scrambled gRNA).

HSP90 was used as a loading control. Schematics of *TCRA* and *TCRB* mRNAs with

and without eIF3 PAR-CLIP sites are shown above. SC control cells have the wild-type

(WT) 3'-UTRs for *TCRA* and *TCRB* mRNAs. **b**, The number of T cells with one or more

TCR clusters measured by anti-TCRA/anti-TCRB protein staining and epifluorescence

microscopy as a function of time after anti-CD3/anti-CD28 activation. A total of 100 cells

from each donor were imaged for *TCRA ΔPAR* (*n* = 2 donors, stained with anti-TCRA

antibody), *TCRB ΔPAR* (*n* = 2 donors, stained with anti-TCRB antibody), and SC cell

lines (*n* = 2 donors, each stained separately with anti-TCRA and anti-TCRB antibodies).

Values are mean ± standard deviation. P < 0.005 comparing *TCRA ΔPAR* and *TCRB*

ΔPAR to SC after 1 hour of activation. P ≤ 0.0002 comparing *TCRA ΔPAR* and *TCRB*

ΔPAR to SC after 3 hours of activation, using two-way ANOVA. **c,** Representative

density plots showing the percentage of I+PMA activated T cells expressing TCR on the

cell surface 3 hours after activation, measured by flow cytometric analysis for one of two

donors. The cell lines shown are *TCRA ΔPAR, TCRB ΔPAR, ΔTCR* (negative control),

and SC. **d**, Percent cells expressing TCR on the cell surface after I+PMA activation for 3

hours (*n = 2* donors, with mean and standard deviation shown), reported in bar graph

form. P < 0.0001 for *TCRA ΔPAR* or *TCRB ΔPAR* relative to SC, using one-way

ANOVA.

Figure 4. Phenotypic analysis of primary human T cells. a, Flow cytometric analysis

measuring T cell activation markers CD69 (early activation marker) and CD25 (mid-

activation marker) in *TCRA ΔPAR, TCRB ΔPAR, ΔTCR* and SC CD8+ T cells,

quantifying mean percent cells expressing CD69+ only (left) and CD69+ CD25+ (right)

from (*n = 2* donors). **b**, Density plots showing cell populations expressing CD69+ only,

CD69+ CD25+ or CD25+ only from one of the donors. Cells sorted as shown in

Extended Data Fig. 9. **c**, Quantification of secreted IL2 and IFNγ from cell lines *TCRA*

ΔPAR, TCRB ΔPAR, ΔTCR and SC cell populations from (*n = 2* donors) at different

time points after stimulation with anti-CD3/anti-CD28 antibodies, as determined by

ELISA. For both figures **a,** and **c,** Values are mean ± standard deviation, ****P < 0.0001

relative to SC, using two-way ANOVA. Differences in expression of activation markers

and secreted cytokines in comparisons of *TCRA ΔPAR* with *TCRB ΔPAR* were not

significant.

Figure 5. Role of the eIF3 PAR-CLIP sites in the 3'-UTRs of *TCRA* and *TCRB*,

measured by nanoluciferase reporters. a, Schematic of the nanoluciferase reporters

stably expressed in Jurkat cells. WT, intact 3'-UTR from either *TCRA* or *TCRB* mRNA;

ΔPAR, 3'-UTR of *TCRA* or *TCRB* with the eIF3 PAR-CLIP site deleted (nucleotides 102-

338 in the 3'-UTR of *TCRA* mRNA or nucleotides 16-161 in the 3'-UTR of *TCRB*

mRNA); R*PAR, reversed PAR-CLIP sequence in the 3'-UTR of *TCRA* or *TCRB* mRNA.

b, Luciferase activity in anti-CD3/anti-CD28 activated Jurkat cells, relative to non-

activated controls ($n = 3$, with mean and standard deviations shown). $P < 0.0001$ for

TCRA ΔPAR and R*PAR relative to WT at 5 hr and 8 hr, and $P \leq 0.0002$ for 10 hr, using

two-way ANOVA. $P < 0.0001$ for *TCRB ΔPAR* and R*PAR relative to WT for 1 hr - 10

hr. **c**, Changes in nanoluciferase mRNA levels in cells stably expressing the

nanoluciferase reporters activated with anti-CD3/anti-CD28 antibodies, as determined

by qRT-PCR. $P < 0.0001$ for *TCRA ΔPAR* relative to WT at 8 hr, and for *TCRA ΔPAR*

and R*PAR relative to WT at 10 hr, using two-way ANOVA. $P \leq 0.0002$ for *TCRB ΔPAR*

and R*PAR relative to WT at 8 hr and 10 hr. **d**, Immunoprecipitation of eIF3 using an

anti-EIF3B antibody[5], followed by qRT-PCR to quantify the amount of nanoluciferase

mRNA bound to eIF3 ($n = 3$, with mean and standard deviations shown). The schematic

of the experiment is indicated on the left. For *TCRA* and *TCRB* mRNAs, $P < 0.0001$ for

ΔPAR and R*PAR relative to WT, using two-way ANOVA. P values for IL2 mRNAs for

ΔPAR and R*PAR relative to WT were not significant.

Materials and methods

Jurkat cell culture

Human Jurkat, Clone E6-1 (ATCC TIB-152) was purchased from American Type

Culture Collection (ATCC) and was maintained in RPMI 1640 Medium (ATCC

modification) with 10% FBS (VWR Life Science Seradigm) and 0.01% Penicillin-

Streptomycin (10,000 U/mL). The cells were maintained between 1 million to 8 million

cells ml^{-1}. When cells were stimulated they were always maintained at 8 million

cells ml^{-1}.

Isolation of human primary T cells

Primary human T cells were isolated from healthy human donors from leukoreduction

chambers after Trima Apheresis (Vitalant, formerly Blood Centers of the Pacific).

Peripheral blood mononuclear cells (PBMCs) were isolated from whole blood samples

by Ficoll centrifugation using SepMate tubes (STEMCELL) per manufacturer's

instructions. T cells were isolated from PBMCs from all cell sources by magnetic

negative selection using an EasySep Human T Cell Isolation Kit (STEMCELL), per

manufacturer's instructions. Unless otherwise noted, isolated T cells were stimulated as

described below and used immediately.

Primary human T cell culture

Bulk T cells were cultured in XVivo15 medium (STEMCELL) with 5% fetal bovine serum

(FBS), 50 µM 2-mercaptoethanol, and 10 µM *N*-acetyl-cystine. Immediately after

isolation, T cells were stimulated for 2 days with anti-human CD3/CD28 magnetic

Dynabeads (ThermoFisher) at a bead to cell concentration of 1:1, along with cytokine

IL-2 at 500 U ml^{-1} (UCSF Pharmacy). After electroporation, T cells were cultured in

media with IL-2 at 500 U ml^{-1}. Throughout the culture period T cells were maintained at

an approximate density of 1 million cells per mL of media. Every 2–3 days after

electroporation, additional media was added, along with additional fresh IL-2 to bring the

final concentration to 500 U/mL, and the cells were transferred to larger culture flasks as

necessary to maintain a density of 1 million cells ml^{-1}.

For the T cell stimulation assay, edited primary T cells were transferred to fresh media

lacking IL2 after 9 days of culturing. The T cells were then stimulated with anti-CD3 and

anti-CD28 antibodies using flat bottom plates coated with anti-CD3 antibody (at a 10 µg

ml^{-1} concentration), and anti-CD28 antibody added to the cell culture media at a

concentration of 5 µg ml^{-1}.

Jurkat cell stimulation

The Jurkat cells used for the for the PAR-CLIP experiment were stimulated with 1X Cell

Stimulation Cocktail, containing phorbol 12-myristate 13-acetate (PMA) and ionomycin

(Thermofisher, Catalog number: 00-4970-93). Unless otherwise stated all other Jurkat

cell simulations were performed with anti-CD3 and anti-CD28 antibodies (Tonbo). Flat

bottom plates were coated with anti-CD3 antibody (at a 10 µg ml^{-1} concentration), and

anti-CD28 antibody was added to the cell culture media at a concentration of 5 µg ml^{-1}.

4-thiouridine optimization

Jurkat cells were seeded, so that they reached 8×10^5 cells ml^{-1} seeding density on the

day of the experiment. Varying concentrations of s4U were added to the cells (50 µM,

75 µM, 100 µM, or none as a negative control). The cells were then incubated for

different time points: 8 hours, 10 hours, 12 hours, or 16 hours. After each incubation

time cell viability was determined using the CellTiter-Glo assay (Promega), according to

the manufacturer's instructions. Concentrations at which the relative luminescence in

the presence of s4U (luminescence of the s4U treated cells/luminescence of the

untreated cells) exceeded 95% were considered non-toxic. Based on these

measurements, we used 50 µM of 4-thiouridine for PAR-CLIP experiments.

PAR-CLIP

Two biological replicates were used to perform PAR-CLIP analysis as described in[5],

with modifications for Jurkat cells. 50 µM of 4-thiouridine was determined as non-toxic to

Jurkat cells over the time course of the PAR-CLIP experiments[14]. A total of 55 million

Jurkat cells seeded at 8 million cells ml^{-1} was treated with 50 µM of 4-thiouridine

(Sigma) for 7 hours, then stimulated with 1X Cell Stimulation Cocktail for 5 hours in

media containing 50 µM of 4-thiouridine (**Fig. 1a**). The same number of cells were

treated with 50 µM of 4-thiouridine (Sigma) for 12 hours without stimulation as a non-

activated control. The cells were then crosslinked on ice with 365 nm UV irradiation at

an energy dose of 0.2 J cm^{-2}. The cells were pelleted by centrifugation at 100 x g for 15

min at 4 °C, and the pellet was resuspended in three volumes of NP40 lysis buffer (50

mM HEPES-KOH pH 7.5, 150 mM KCl, 2 mM EDTA, 0.5% Nonidet P-40 alternative, 0.5

mM dithiothreitol (DTT), 1 Complete Mini EDTA-free Protease Inhibitor Cocktail tablet

(Roche)). The cell suspension was then incubated on ice for 10 min, passed through an

18G needle five times, and centrifuged at 13,000 x g for 15 min at 4 °C and RNAs were

lightly digested by treatment with MNase (Thermo Scientific) at a final concentration of

0.05 U µl⁻¹ for 20 min at 16 °C. For each PAR-CLIP assay 1000 µl of Dynabeads

(Invitrogen) and 800 µl of anti-EIF3B antibody (Bethyl A301-761A) were used. The

remaining steps of the PAR-CLIP analysis was performed exactly as described in [5,41]

with the exception of using MNase at 5 U µl⁻¹ for the on-bead digestion step.

PAR-CLIP computational analysis

PAR-CLIP cDNA libraries were sequenced on an Illumina HiSeq 2500 platform. To

eliminate potential PCR biases during PAR-CLIP library preparation, a random bar code

was introduced into the 3' adapter and all the reads that matched the random barcode

were collapsed into single reads. Clusters of overlapping sequence reads mapped

against the human genome version hg38 were generated using the PARalyzer software

[40] incorporated into the PARpipe pipeline (https://ohlerlab.mdc-

berlin.de/software/PARpipe_119/, [13]) with the settings below. Binding sites were

categorized using the Gencode GRCh38.p12 GTF annotations

(gencode.v21.annotation.gtf), https://www.gencodegenes.org/human/.

The PARpipe settings used were:

Conversion = T>C

Minimum read count per group = 5

Minimum read count per cluster = 7

Minimum read count for kde = 3

Minimum cluster size = 11

Minimum conversion locations for cluster = 2

Minimum conversion count for cluster = 2

Minimum read count for cluster inclusion = 1

Minimum read length = 20

Maximum number of non conversion mismatches = 1

Comparison of eIF3 PAR-CLIP results in Jurkat and HEK293T cells

To compare RNAs in activated Jurkat cells crosslinked to eIF3 with those crosslinked to

eIF3 in HEK293T cells[5], the gene cluster lists (*.gene_cl.csv) from APA_REP1,

APA_REP2, APB_REP1, APB_REP2, APD_REP1, and APD_REP2 were used (see

Supplementary Table 3). The genes were first sorted by total read counts

("ReadCountSum") from high to low for each library to obtain the top candidate genes.

Then, the same number of top candidate genes from these sorted lists as the number of

genes identified in HEK293T cells, for eIF3 subunit EIF3A, EIF3B, and EIF3D

crosslinked to RNA[5], were chosen for comparison.

PAR-CLIP Pathway analysis

To determine biological pathways enriched in the set of mRNAs that crosslinked to eIF3

in activated Jurkat cells, genes with at least 100 total aligned reads were used, as

determined in the PARpipe analysis described above[13], from the EIF3A/C/B samples.

Since PAR-CLIP reads are short, it is not possible to determine with certainty which

mRNA transcript isoform cross-linked with eIF3. Therefore the most abundant mRNA

transcript isoform for each gene was chosen, as determined by transcriptome profiling

using kallisto (protein_coding category)[42], as described in the Transcriptome Profiling

section. Even with this choice, eIF3 crosslinks to mRNAs do not correlate with mRNA

abundance (**Extended Data Figure 4b**). Human genome GRCh38.p13 annotation was

used to extract mRNA lengths by 5'-UTR, coding region and 3'-UTR (Ensembl

Biomart)[43]. These genes were then sorted by the density of PAR-CLIP reads in the

mRNA 5'-UTR region, prior to mapping pathways of transcripts that cross-linked to eIF3.

Due to the complexity of *TCRA* and *TCRB* transcript annotation, these transcripts were

excluded from the normalization calculation, but included in the pathway analysis. The

top 500 genes from the resulting EIF3A/C/B PAR-CLIP gene lists were used, sorted as

described above, to analyze gene enrichment profiles in the STRING database[44]. The

top tissue-specific categories involve the immune system and T cell function

(**Supplementary Table 6**). Note that the STRING database does not include TCR

subunits in its analysis.

Metagene analysis

The PAR-CLIP genes sorted as described above in the "PAR-CLIP pathway analysis"

were used and mapped against the most abundant mRNA transcript isoforms to

generate cumulative plots of the reads. Reads for the *TCRA* and *TCRB* mRNAs were

manually extracted from the Bowtie version 1.0.0 Hg38 alignment of the eIF3 PAR-CLIP

reads. These reads were combined with the mapped reads in the *.read.csv files

generated by Parpipe. The combined reads were then sorted to extract only reads

annotated as 5'-UTR, start codon, coding, stop codon, and 3'-UTR. The most abundant

transcript isoform, as identified in the transcriptome profiling using kallisto (described

below) was used. Reads mapped to the 5'-UTR and start codon were normalized by the

length of the 5'-UTR. Reads mapped to the coding region and stop codon were

normalized by the length of the coding region. Finally, reads mapped to the 3'-UTR

were normalized to the length of the 3'-UTR. Relative positions of the mapped reads

along a metagene were computed based on the locations each mapped read fell within

its respective feature. Relative positions were from -1 to 0 for the 5'-UTR, 0 to 1 for the

coding region, and 1 to 2 for the 3'-UTR. 5'-UTR values were computed by multiplying

the relative position by -1, whereas the 3-UTR values were were computed by adding 1

to the relative position. Coding region relative positions were unchanged. The empirical

cumulative distribution frequency function from R package ggplot2[45] was used to build

the metagene plots from the vector of relative positions for the reads which mapped to a

given set of reads.

We defined mRNAs having a ratio of normalized 5'-UTR reads divided by 3'-UTR reads

of 20 or more as 5'-UTR enriched with respect to eIF3 crosslinking. All others were

categorized as mRNAs with eIF3 crosslinking across the entire length of the mRNA.

The cut-off value 20 is not a sharp boundary between the two categories of mRNA

(**Extended Data Fig. 5a**).

Transcriptome Profiling

RNA samples were extracted from HEK293T cells, non-activated Jurkat cells, and

Jurkat cells activated for 5 hr with I+PMA, using the Direct-zol RNA Miniprep kit

(Zymoresearch). The libraries were prepared using TruSeq Stranded RNA LT Ribo-Zero

Gold kit (Illumina) following the manufacturer's instructions, with two biological

replicates per condition. Cutadapt (version 2.6)[46] with a minimum read length of 20, 5'

end with a cutoff of 15 and the 3' end with a cutoff of 10 in paired-end mode was used

to remove adapters. RNA-seq reads were pseudoaligned using kallisto v.0.46.0 run in

quant mode with default parameters to estimate transcript abundance (transcripts per

million, TPM)[42]. The transcript index for kallisto was made with default parameters and

GENCODE Release 32 (GRCh38.p13) FASTA file[47].

To compare HEK293T or non-activated Jurkat mRNA abundance with that in activated

Jurkat cells, the output from kallisto was used for analysis using DeSeq2 (ref [48]). All

protein-coding transcripts (protein_coding) were then selected for the comparisons,

using tximport[49] as part of the Bioconductor environment in R. All genes with ≥ 50 reads

summed across all four samples in a comparison (two biological replicates per

condition) were then used. Prior to making MA plots, the Approximate Posterior

Estimation for generalized linear model (apeglm) shrinkage estimator for log fold

changes was used[50] DeSeq2 output in **Supplementary Table 5** is presented prior to

shrinkage analysis.

Mass spectrometry

To identify eIF3 subunits that crosslinked with RNAs in the PAR-CLIP experiments, a

portion of eIF3 immunoprecipitated using Dynabeads as described above were treated

with nonradioactive ATP during the T4 PNK labeling step. The nonradioactive samples

were then run on the same gel next to the radiolabeled PAR-CLIP samples, Coomassie

stained (Pierce) and the bands that matched with the phosphorimager printout were

excised from the gel and submitted for identification using one-dimensional LC-MS/MS

(**Supplementary Table 1**).

sgRNA/Cas9 RNP production

The sgRNA/Cas9 RNPs used to edit Jurkat cells were produced by complexing sgRNA

(Synthego) to Cas9 as described[51] while RNPs to edit Primary Human T cells were

produced by complexing a two-component gRNA (crRNA and tracrRNA, Dharmacon) to

Cas9 as described in[19]. The targeting sequences for the sgRNAs and crRNA are given

in **Supplementary Table 7**. Recombinant Cas9-NLS was obtained from MacroLab in

the California Institute for Quantitative Biosciences.

Jurkat and primary T cell genome editing

Jurkat cells used for electroporation were collected at passage 5 or lower and were

maintained at a seeding density of 8 million cells ml^{-1} or lower. Primary T cells were

isolated as described above. Prior to electroporation the Primary T cells were stimulated

with magnetic anti-CD3/anti-CD28 Dynabeads (ThermoFisher) for 48 hours. After 48

hours these beads were removed from the cells by placing cells on an EasySep cell

separation magnet for 2 min before electroporation. One million Jurkat (not activated)

or primary T cells cells (activated with anti-CD3/anti-CD28 Dynabeads for 48 hours)

were rinsed with PBS and then resuspended in 20 µl of Lonza electroporation buffer P3.

The cells were then mixed with 2.5 µl Cas9 RNPs (50 pmol total) along with 2 µl of a

127-nucleotide non-specific single-stranded DNA oligonucleotide at 2 µg µl^{-1} (4 µg

ssDNA oligonucleotide total). The cells were then electroporated per well using a Lonza

4D 96-well electroporation system with pulse code DN100 for Jurkat cells and EH115

for primary human T cells. Immediately after electroporation, 80 µl of pre-warmed

media (without cytokines) was added to each well, and the cells were allowed to rest for

15 min at 37 °C in a cell culture incubator while remaining in electroporation cuvettes.

After 15 min, cells were moved to final culture flasks.

Western Blot

Western blot analysis was performed using the following antibodies: anti-TCRA (SC-

515719), anti-HSP90 (SC-69703) (**Supplementary Table 7**).

Total mRNA isolation and quantitative RT-PCR analysis

Total RNA was isolated from whole cells for qRT-PCR using Quick RNA miniprep plus

kit from Zymo Research following the manufacturer's instructions. Quantitative RT-PCR

analysis was performed using the Power SYBR Green RNA-to-Ct 1-Step kit (Applied

Biosystems) according to the manufacturer's instructions, and the QuantStudio™ 3

Real-Time PCR System (ThermoFisher). Each target mRNA was quantified in three

biological replicates, with each biological replicate having three technical replicates.

Plasmids

The *TCRA* 3'-UTR and *TCRB* 3'-UTR sequences were amplified from Jurkat genomic

DNA. The nanoluciferease sequence fused to a PEST domain was amplified from

pNL1.2[*NlucP*] Vector Sequence (Promega) and was cloned into a modified CD813A

vector (System Biosciences) containing the beta-globin (*HBB*) 5'-UTR using the In-

Fusion® HD Cloning Kit (Takara). The subsequent mutations in the *TCRA* and *TCRB*

3'-UTRs were generated using these initial constructs. Both *TCRA/TCRB ΔPAR or*

R*PAR sequences were purchased as gblocks from IDT and were cloned into this

plasmid backbone.

Generation of stable Jurkat cell lines expressing nanoluciferase reporter

For lentiviral production HEK293T cells were plated at a density of 80% in T-75 flasks

the night before transfection. The cells were then transfected with plasmids: expressing

the nanoluciferase, PsPAX2 and pCMV-VSV-G using the Lipofectamine 3000 reagent

(ThermoFisher) following the manufacturer's instructions. Forty-eight hours after

transfection, the viral supernatant was collected, filtered and concentrated using PEG-it

Virus Precipitation Solution (System Biosciences) following the manufacturer's

instructions. The virus pellets were then resuspended in RPMI-1640 media and stored

in -80°C.

The Jurkat cell transductions were done with multiple viral titers using TransDux™ MAX

Lentivirus Transduction Reagent (System Biosciences) following the manufacturer's

instructions. To test the viral transduction efficiency of the cells, forty-eight hours after

viral transduction the percent of cells expressing GFP was measured by FACS analysis

and cells expressing less than 15% GFP was treated with 1µg ml^{-1} puromycin to the

media. The cells were maintained in media with 1µg ml^{-1} puromycin, but transferred into

media without puromycin the day before the cells were used for assays.

Luciferase reporter assay

The stable cell lines expressing the Nanoluciferease reporter were stimulated with anti-

CD3 and anti-CD28 antibodies and the nanoluciferase activity was assayed after 1 hr, 3

hr, 5 hr, 8 hr, 10 hr and 24 hrs after stimulation using Nano-Glo® Luciferase Assay

System (Promega). For each time point 200,000 cells were tested in triplicate for each

cell line.

RNA immunoprecipitation and qPCR

The EIF3B-RNA immunoprecipitations were carried out following the exact same

conditions used for the PAR-CLIP analysis with the following changes. For each

immunoprecipitation, cell lysates were prepared in NP40 lysis buffer (50 mM HEPES-

KOH pH 7.5, 150 mM KCl, 2 mM EDTA, 0.5% Nonidet P-40 alternative, 0.5 mM DTT, 1

Complete EDTA-free Proteinase Inhibitor Cocktail tablet per 50 ml of buffer) with 4

million cells. The lysates were then incubated with 50 µl Protein G Dynabeads

conjugated with 20 µl of anti-EIF3B antibody (Bethyl A301-761A) for two hours at 4 °C.

After incubation, the flow through was removed and the beads were washed three times

with 1 ml NP40 lysis buffer for each wash. The beads were then resuspended in 400 µl

TRIzol reagent (ThermoFisher) and vortexed for 1 minute. The RNA was extracted

following the manufacturer's instructions and qPCR was performed as described above

using primers listed in **Supplementary Table 7**.

Flow cytometry and cell sorting

Flow cytometric analysis was performed on an Attune NxT Acoustic Focusing

Cytometer (ThermoFisher). Surface staining for flow cytometry and cell sorting was

performed by pelleting cells and resuspending in 50 µl of FACS buffer (2% FBS in PBS)

with antibodies at the indicated concentrations (**Supplementary Table 7**) for 20 min at

4 °C in the dark. Cells were washed twice in FACS buffer before resuspension.

ELISA

The cell suspensions were collected after each time point of activation with anti-

CD3/anti-CD28 antibodies for WT, *TCRA ΔPAR* or *TCRB ΔPAR* cells. For each

timepoint the same number of cells were used from each strain to be able to compare

across strains and time points. The amount of secreted IL2 in the cell suspensions after

activation anti-CD3/anti-CD28 antibodies for WT, *TCRA ΔPAR* or *TCRB ΔPAR* cells

were measured by ELISA MAX™ Deluxe Set Human IL-2 (BioLegend) according to the

manufacturer's instructions.

RNA-FISH and immunofluorescence

Jurkat cells were washed with PBS, fixed with 3.7% (vol./vol.) formaldehyde for 10 min

at room temperature and washed three times with PBS. PBS was discarded and 1 ml

70% ethanol was added. The cells were incubated at 4 °C for 16 hours. The 70%

ethanol was aspirated and the cells were washed once with 0.5 ml Stellaris RNA wash

buffer A (Biosearch technologies). The cells were then incubated with 100 µl Stellaris

hybridization buffer (Biosearch Technologies) containing Stellaris RNA FISH probes

(Biosearch Technologies) at a final concentration of 125 nM (**Supplementary Table 7)**

and with the relevant antibody (**Supplementary Table 7)** for 16 hours at 28 °C. The

cells were then washed twice with 0.5 ml Stellaris RNA wash buffer A containing

secondary antibody conjugated with a fluorophore for 30 minutes at 37 °C in the dark.

The second Stellaris RNA wash buffer A contained DAPI in addition to the secondary

antibody. Finally the cells were washed once with 0.5 mL Stellaris RNA wash buffer B

and mounted with mounting solution (Invitrogen). All high resolution images were taken

using confocal ZEISS LSM 880 in Airyscan super-resolution mode, equipped withA

Plan-Apochromat 63x/1.4 Oil objective (Zeiss). To measure colocalization of *TCRA* and

TCRB mRNAs with each other or with P bodies (using DCP1 antibody) or stress

granules (G3BP1 antibody) the cells were mounted along with 0.1 µm TetraSpeck™

microspheres (ThermoFisher) adhered to the slide according to manufacturer's

instructions, to be able to account for the chromatic shift during image acquisition. For

the counting of cells containing TCR clusters, a Revolve Epi-Fluorescence microscope

(Echo), equipped with an A Plan-Apochromat 40x objective (Olympus) was used.

Colocalization analysis

To measure colocalization of *TCRA* and *TCRB* mRNAs with each other or with P bodies

(using DCP1 antibody) or stress granules (G3BP1 antibody), immunofluorescently

labelled cells (see above) were mounted along with 0.1 µm TetraSpeck™ microspheres

(ThermoFisher) adhered to the slide according to manufacturer's instructions, to be able

to account for the lateral and axial chromatic aberrations during image acquisition.

Z-stacks were acquired with 35 nm x 35 nm x 190 nm voxels on a ZEISS LSM 880 in

Airyscan super-resolution mode, equipped with a Plan-Apochromat 63x/1.4 Oil objective

(Zeiss). The images were then deconvolved to a lateral resolution of ≈150 nm and an

axial resolution of ≈500 nm (as confirmed by observing the discrete Fourier transform of

the z-stacks). After imaging a single cell, beads that were on the slide axial to the cell

were imaged to measure the corresponding lateral and axial chromatic aberrations.

To quantify the relative colocalization, we developed an automated processing and

analysis pipeline in ImageJ 1.52p available on github:

https://github.com/Llamero/TCR_colocalization_analysis-macro. Specifically, the

chromatic aberrations in the z-stacks were compensated for by registering the channels

of the bead z-stack to one another, and then applying the same registration vectors to

the corresponding channels in the cell z-stack[52]. Each channel of a z-stack was then

thresholded to remove background in the image, and then the colocalization between

each pair of images was measured using the Pearson's correlation coefficient.

Samples in which any pair of channels in the bead z-stack had a correlation of less than

0.45 were removed from final analysis, as this suggested that the images had

insufficient dynamic range in at least one of the channels for an accurate deconvolution.

Data and Code Availability

Sequencing data supporting the findings in this study have been deposited in the Gene

Expression Omnibus (GEO) Database (https://www.ncbi.nlm.nih.gov/geo/), with

accession codes _____. Code used to analyze the microscopy images is available on

github at https://github.com/Llamero/TCR_colocalization_analysis-macro. Raw data for

Figure 3a, **Extended Data Figure 1b**, and **Extended Data Figure 13a** are provided in

Supplementary Figures 1-3.

Acknowledgements

We thank M. Hafner for advice on PAR-CLIP methodology and data analysis, H. Nolla

and A. Valeros at the UC Berkeley flow cytometry facility for helping out with the FACS

analysis and single cell sorting, F. Ives and H. L. Aaron at the UC Berkeley Imaging

center for help with microscopy, J. Lui for advice on planning experiments and for

suggestions on the manuscript, and M. Mignardi for advice on FISH experiments. This

work was supported by NIH grant R01-GM065050 (J.H.D.C.) and a grant from the Tang

Prize for Biopharmaceutical Science (J.H.D.C.). N.T.I. is a Damon Runyon-Rachleff

Innovator (DRR#37-15) and NIH New Innovator (DP2 CA195768). A.M. holds a Career

Award for Medical Scientists from the Burroughs Wellcome Fund, is an investigator at

the Chan Zuckerberg Biohub and has received funding from the Innovative Genomics

Institute (IGI) and the Parker Institute for Cancer Immunotherapy (PICI). B.E.S. is

supported by NIH grant P30EY003176. Imaging experiments were conducted at the

Cancer Research Laboratory Molecular Imaging Center, supported by Helen Wills

Neuroscience Institute. Sequencing was carried out at the UCSF Center for Advanced

Technology and at the Vincent J. Coates Genomics Sequencing Laboratory at UC

Berkeley, supported by NIH grant S10 OD018174.

Author Contributions

Project and experiments were conceived and designed by D.D.S. and J.H.D.C. D.D.S.

carried out all experiments, with assistance from B.E.S. and G.H.C. for image

acquisition and analysis, from T.R., R.A.A., and A.M. for primary T cell genome editing

and cell culture protocols, and from L.F., M.K., N.T.I. and J.H.D.C. for analysis of the

deep sequencing-based experiments. The manuscript was written by D.D.S. and

J.H.D.C., with editing by all authors.

Competing Interests

The authors declare competing financial interests: T.L.R. and A.M. are co-founders of

Arsenal Therapeutics. A.M. is a co-founder of Spotlight Therapeutics. A.M. serves on the

scientific advisory board of PACT Pharma, is an advisor to Trizell and was a former

advisor to Juno Therapeutics. The Marson Laboratory has received sponsored research

support from Juno Therapeutics, Epinomics, Sanofi and a gift from Gilead. A provisional

patent application has been filed on some of the work presented herein.

Materials and Correspondence

Correspondence and requests for materials should be addressed to J.H.D.C.,

jcate@lbl.gov.

Bibliography

1. Ahern, T., Sampson, J. & Kay, J. E. Initiation of protein synthesis during

lymphocyte stimulation. *Nature* **248**, 519–521 (1974).

2. Kleijn, M. & Proud, C. G. The regulation of protein synthesis and translation factors

by CD3 and CD28 in human primary T lymphocytes. *BMC Biochem.* **3**, 11 (2002).

3. Miyamoto, S., Patel, P. & Hershey, J. W. B. Changes in ribosomal binding activity

of eIF3 correlate with increased translation rates during activation of T

lymphocytes. *J. Biol. Chem.* **280**, 28251–28264 (2005).

4. Ricciardi, S. *et al.* The Translational Machinery of Human CD4+ T Cells Is Poised

for Activation and Controls the Switch from Quiescence to Metabolic Remodeling.

Cell Metab. **28**, 895–906.e5 (2018).

5. Lee, A. S. Y., Kranzusch, P. J. & Cate, J. H. D. eIF3 targets cell-proliferation

messenger RNAs for translational activation or repression. *Nature* **522**, 111–114

(2015).

6. Lee, A. S., Kranzusch, P. J., Doudna, J. A. & Cate, J. H. D. eIF3d is an mRNA cap-

binding protein that is required for specialized translation initiation. *Nature* **536**, 96–

99 (2016).

7. Meyer, K. D. *et al.* 5' UTR m(6)A Promotes Cap-Independent Translation. *Cell* **163**,

999–1010 (2015).

8. de la Parra, C. *et al.* A widespread alternate form of cap-dependent mRNA

translation initiation. *Nat. Commun.* **9**, 3068 (2018).

9. Curran, S. P. & Ruvkun, G. Lifespan regulation by evolutionarily conserved genes

essential for viability. *PLoS Genet.* **3**, e56 (2007).

10. Choudhuri, A., Maitra, U. & Evans, T. Translation initiation factor eIF3h targets

specific transcripts to polysomes during embryogenesis. *Proc. Natl. Acad. Sci. U.*

S. A. **110**, 9818–9823 (2013).

11. Cattie, D. J. *et al.* Mutations in Nonessential eIF3k and eIF3l Genes Confer

Lifespan Extension and Enhanced Resistance to ER Stress in Caenorhabditis

elegans. *PLoS Genet.* **12**, e1006326 (2016).

12. Hafner, M. *et al.* Transcriptome-wide identification of RNA-binding protein and

microRNA target sites by PAR-CLIP. *Cell* **141**, 129–141 (2010).

13. Mukherjee, N. *et al.* Deciphering human ribonucleoprotein regulatory networks.

Nucleic Acids Res. **47**, 570–581 (2019).

14. Ascano, M., Hafner, M., Cekan, P., Gerstberger, S. & Tuschl, T. Identification of

RNA-protein interaction networks using PAR-CLIP. *Wiley Interdiscip. Rev. RNA* **3**,

159–177 (2012).

15. Hubstenberger, A. *et al.* P-Body Purification Reveals the Condensation of

Repressed mRNA Regulons. *Mol. Cell* **68**, 144–157.e5 (2017).

16. Aizer, A. *et al.* Quantifying mRNA targeting to P-bodies in living human cells

reveals their dual role in mRNA decay and storage. *J. Cell Sci.* **127**, 4443–4456

(2014).

17. Aulas, A. *et al.* G3BP1 promotes stress-induced RNA granule interactions to

preserve polyadenylated mRNA. *J. Cell Biol.* **209**, 73–84 (2015).

18. Protter, D. S. W. & Parker, R. Principles and Properties of Stress Granules. *Trends*

Cell Biol. **26**, 668–679 (2016).

19. Roth, T. L. *et al.* Reprogramming human T cell function and specificity with non-

viral genome targeting. *Nature* **559**, 405–409 (2018).

20. Ohashi, P. S. *et al.* Reconstitution of an active surface T3/T-cell antigen receptor by

DNA transfer. *Nature* **316**, 606–609 (1985).

21. Koning, F., Lew, A. M., Maloy, W. L., Valas, R. & Coligan, J. E. The biosynthesis

and assembly of T cell receptor alpha- and beta-chains with the CD3 complex. *J.*

Immunol. **140**, 3126–3134 (1988).

22. Cochran, J. R., Aivazian, D., Cameron, T. O. & Stern, L. J. Receptor clustering and

transmembrane signaling in T cells. *Trends Biochem. Sci.* **26**, 304–310 (2001).

23. Voon, D. C. *et al.* Use of mRNA- and protein-destabilizing elements to develop a

highly responsive reporter system. *Nucleic Acids Res.* **33**, e27 (2005).

24. Hall, M. P. *et al.* Engineered luciferase reporter from a deep sea shrimp utilizing a

novel imidazopyrazinone substrate. *ACS Chem. Biol.* **7**, 1848–1857 (2012).

25. Au-Yeung, B. B. *et al.* A sharp T-cell antigen receptor signaling threshold for T-cell

proliferation. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E3679–88 (2014).

26. Au-Yeung, B. B. *et al.* IL-2 Modulates the TCR Signaling Threshold for CD8 but Not

CD4 T Cell Proliferation on a Single-Cell Level. *J. Immunol.* **198**, 2445–2456

(2017).

27. Allison, K. A. *et al.* Affinity and dose of TCR engagement yield proportional

enhancer and gene activity in CD4+ T cells. *Elife* **5**, (2016).

28. Richard, A. C. *et al.* T cell cytolytic capacity is independent of initial stimulation

strength. *Nat. Immunol.* **19**, 849–858 (2018).

29. Iezzi, G., Karjalainen, K. & Lanzavecchia, A. The duration of antigenic stimulation

determines the fate of naive and effector T cells. *Immunity* **8**, 89–95 (1998).

30. van Stipdonk, M. J., Lemmens, E. E. & Schoenberger, S. P. Naïve CTLs require a

single brief period of antigenic stimulation for clonal expansion and differentiation.

Nat. Immunol. **2**, 423–429 (2001).

31. Schrum, A. G., Palmer, E. & Turka, L. A. Distinct temporal programming of naive

CD4+ T cells for cell division versus TCR-dependent death susceptibility by

antigen-presenting macrophages. *Eur. J. Immunol.* **35**, 449–459 (2005).

32. Eyquem, J. *et al.* Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances

tumour rejection. *Nature* **543**, 113–117 (2017).

33. Watanabe, K., Kuramitsu, S., Posey, A. D., Jr & June, C. H. Expanding the

Therapeutic Window for CAR T Cell Therapy in Solid Tumors: The Knowns and

Unknowns of CAR T Cell Biology. *Front. Immunol.* **9**, 2486 (2018).

34. Mohammad, M. P., Munzarová Pondelícková, V., Zeman, J., Gunišová, S. &

Valášek, L. S. In vivo evidence that eIF3 stays bound to ribosomes elongating and

terminating on short upstream ORFs to promote reinitiation. *Nucleic Acids Res.* **45**,

2658–2674 (2017).

35. Lin, Y. *et al.* eIF3 associates with 80S ribosomes to promote translation elongation,

mitochondrial homeostasis, and muscle health. *bioRxiv* 651240 (2019)

doi:10.1101/651240.

36. Bohlen, J., Fenzl, K., Kramer, G., Bukau, B. & Teleman, A. A. Selective 40S

footprinting reveals that scanning ribosomes remain cap-tethered in human cells.

bioRxiv 806364 (2019) doi:10.1101/806364.

37. Tan, H. *et al.* Integrative Proteomics and Phosphoproteomics Profiling Reveals

Dynamic Signaling Networks and Bioenergetics Pathways Underlying T Cell

Activation. *Immunity* **46**, 488–503 (2017).

38. Myers, D. R., Wheeler, B. & Roose, J. P. mTOR and other effector kinase signals

that impact T cell function and activity. *Immunol. Rev.* **291**, 134–153 (2019).

39. Manfrini, N. *et al.* High levels of eukaryotic Initiation Factor 6 (eIF6) are required for

immune system homeostasis and for steering the glycolytic flux of TCR-stimulated

CD4+ T cells in both mice and humans. *Dev. Comp. Immunol.* **77**, 69–76 (2017).

40. Corcoran, D. L. *et al.* PARalyzer: definition of RNA binding sites from PAR-CLIP

short-read sequence data. *Genome Biol.* **12**, R79 (2011).

41. Danan, C., Manickavel, S. & Hafner, M. PAR-CLIP: A Method for Transcriptome-

Wide Identification of RNA Binding Protein Interaction Sites. *Methods Mol. Biol.*

1358, 153–173 (2016).

42. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-

seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

43. Cunningham, F. *et al.* Ensembl 2019. *Nucleic Acids Res.* **47**, D745–D751 (2019).

44. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with

increased coverage, supporting functional discovery in genome-wide experimental

datasets. *Nucleic Acids Res.* **47**, D607–D613 (2019).

45. Wickham, H., Chang, W. & Others. ggplot2: Create elegant data visualisations

using the grammar of graphics. *R package version* **2**, (2016).

46. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10–12 (2011).

47. Frankish, A. *et al.* GENCODE reference annotation for the human and mouse

genomes. *Nucleic Acids Res.* **47**, D766–D773 (2019).

48. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

49. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. *F1000Res.* **4**, 1521

(2015).

50. Zhu, A., Ibrahim, J. G. & Love, M. I. Heavy-tailed prior distributions for sequence

count data: removing the noise and preserving large differences. *Bioinformatics* **35**,

2084–2092 (2019).

51. Schumann, K. *et al.* Generation of knock-in primary human T cells using Cas9

ribonucleoproteins. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10437–10442 (2015).

52. Parslow, A., Cardona, A. & Bryson-Richardson, R. J. Sample drift correction

following 4D confocal time-lapse imaging. *J. Vis. Exp.* (2014) doi:10.3791/51086.

53. Noguchi, S. *et al.* FANTOM5 CAGE profiles of human and mouse samples. *Sci*

Data **4**, 170112 (2017).

Extended Data Figures

Extended Data Figure 1. Composition of eIF3 in the Jurkat PAR-CLIP experiments.

a, Composition of eIF3 in I+PMA activated and non-activated Jurkat cells after anti-

EIF3B immunoprecipitation (IP), identified by mass spectrometry. Shown is an SDS

polyacrylamide gel stained with Coomassie Brilliant Blue. **b**, Composition of eIF3 in

activated Jurkat cells determined by western blot after anti-EIF3B IP. IN: input; FT: flow-

through from anti-EIF3B IP beads; ELU: elution of eIF3 from anti-EIF3B IP beads. Anti-

EIF4G1 and anti-RPS19 western blots confirm stringency of bead wash steps[5].

Extended Data Figure 2. Categories of RNA crosslinked to eIF3 by PAR-CLIP in

Jurkat cells. PAR-CLIP crosslinking sites to eIF3, defined by clusters, divided into RNA

categories, in I+PMA activated Jurkat cells (left) and non-activated Jurkat cells (right). In

all panels, RNA categories include: 5'-UTR, 5'-UTR of mRNA; 3'-UTR, 3'-UTR of

mRNA; CDS, protein coding region of mRNA; Start codon, beginning of mRNA CDS;

Stop codon, stop codon region of mRNA; Intron, regions of pre-mRNA; Other, other

classes of RNA; No annotation, reads that map to the human genome but that have no

annotation assigned. Clusters in different mRNA categories may occur in the same

mRNA species.

Extended Data Figure 3. Properties of PAR-CLIP sequence reads. a, The length

distribution of PAR-CLIP clusters mapped to RNAs crosslinked to eIF3 in the EIF3A/C/B

samples, from activated (left) and non-activated (right) Jurkat cells. **b**, The frequency

distribution of eIF3 PAR-CLIP clusters mapped to RNAs in the EIF3A/C/B samples,

from activated (left) and non-activated (right) Jurkat cells. In all panels, the distributions

are the average of both biological replicates. RNA PAR-CLIP properties observed in the

EIF3A/C/B sample are representative of those seen in PAR-CLIP sequence reads of

RNAs crosslinked to other eIF3 subunits.

Extended Data Figure 4. Comparison of transcription profiles and PAR-CLIP

reads. MA plots of **a**, non-activated Jurkat cells compared to Jurkat cells activated with

I+PMA for 5 hours (left) and HEK293T cells compared to Jurkat cells activated with

I+PMA for 5 hours (right). In **a**, M = $\log_2$(HEK293T mRNA TPM/Jurkat mRNA TPM),

and A = mean(TPM) for activated Jurkat mRNAs. TPM, transcripts per million. The FDR

was set to 0.01, and the blue dots indicate significant fold changes. **b,** Scatterplot of

TPM of mRNAs expressed in activated Jurkat cells (most abundant isoform, see

Methods) versus mRNAs identified by PAR-CLIP as crosslinked to eIF3, with ≥ 100 total

read counts.

Extended Data Figure 5. eIF3-mRNA crosslinking patterns in activated Jurkat

cells. a, mRNAs sorted by increasing value of the ratio of normalized 5'-UTR reads to

normalized 3'-UTR reads. Values of the ratio of 5'-UTR/3'-UTR normalized reads > 200

have been truncated to a value of 200 in the plot (asterisk). The arrow indicates the

threshold used to create cumulative plots shown in **Figure 2a**. **b**, Examples of eIF3

crosslinking to histone mRNAs *HIST1H2AI* and *HIST1H3H* are shown. Transcription

start sites were determined from the FANTOM5 Database[53].

Extended Data Figure 6. *TCRA* and *TCRB* mRNA localization patterns. a, Single

plane image of TetraSpeck microspheres (100 nm) adjacent to Jurkat cell probed for P

bodies (far left). Scatter plots of pixel intensities in different emission channels are

shown: channel 1, 670 nm; channel 2, 570 nm; channel 3, 488 nm. Plots left to right are:

channel 1 vs. channel 2, channel 1 vs. channel 3, and channel 2 vs. channel 3. **b**,

Single plane image of Jurkat cell probed for *TCRA* mRNA (channel 1), *TCRB* mRNA

(channel 2), and DCP1 to mark P bodies (channel 3). Scatter plots to the right are

shown as in **a**. **c**, Single plane image and scatter plots for TetraSpeck microspheres as

in **a**, but adjacent to Jurkat cell probed for stress granules. **d**, Single plane image of

Jurkat cell probed for *TCRA* mRNA (channel 1), *TCRB* mRNA (channel 2), and G3BP1

to mark stress granules (channel 3). Scatter plots to the right are shown as in **a**.

Extended Data Figure 7. CRISPR/Cas9 RNP mediated genome editing at the *TCRA*

and *TCRB* genomic loci in primary human T cells. a, Location of gRNA sites

targeting the *TCRA* 3'-UTR, to generate *TCRA ΔPAR* cells. In *TCRA*, gRNA1 and

gRNA2 target hg38 genomic locations chr14:22,551,700 and chr14:22,552,073,

respectively. **b**, PCR primers flanking the *TCRA* editing sites. **c**, Analysis of *TCRA* 3'-

UTR PAR-CLIP site deletion efficiency. Total genomic DNA extracted from WT, *TCRA*

ΔPAR, TCRB ΔPAR (here serving as a negative control), scrambled gRNA

nucleofected (S) and non-nucleofected (N) cells was analyzed by PCR to measure the

editing efficiency. *TCRA ΔPAR* cells produced a 1283 bp PCR product compared to

1676 bp in S, N or *TCRB ΔPAR* cells. **d**, Location of gRNA sites of targeting the *TCRB*

3'-UTR, to generate *TCRB ΔPAR* cells. In *TCRB*, gRNA1 and gRNA2 target hg38

genomic locations chr7:142,802,561 and chr7:142,802,695. **e**, PCR primers flanking

the *TCRB* editing sites. **f**, Analysis of *TCRB* 3'-UTR PAR-CLIP site deletion efficiency.

Total genomic DNA extracted from WT, *TCRA ΔPAR* (here serving as a negative

control)*, TCRB ΔPAR*, scrambled gRNA nucleofected (S) and non-nucleofected (N)

cells were analyzed by PCR to measure the editing efficiency. *TCRB ΔPAR* cells

produced a 1022 bp PCR product compared to 1190 bp in S, N or *TCRA ΔPAR* cells.

Extended Data Figure 8. TCR levels on the cell surface of primary human T cells.

a, Percentage of cells expressing TCR on the cell surface measured by flow cytometric

analysis (*n = 2* donors). TCR expressing cells in the cell populations tested: *1gB, TCRB*

gRNA 1; *2gB, TCRB* gRNA 2; *1+2 gB, TCRB* gRNA 1+2 (i.e. *TCRB ΔPAR*); *1gA, TCRA*

gRNA 1; *2gA, TCRA* gRNA 2; *1+2 gA, TCRA* gRNA 1+2 (i.e. *TCRA ΔPAR*); *ΔTCR*,

TCR gRNA targeting the CDS of *TCRA*; *N*, Non-nucleofected cells; SC, Scrambled

gRNA. (*n* = 2 donors, with standard deviation shown.) Left: I+PMA activated T cells,

after 3 hours of activation; Right: non-activated T cells. P values for *TCRA ΔPAR, TCRB*

ΔPAR, ΔTCR were < 0.0001 relative to SC in both activated and non-activated cells,

using one-way ANOVA. The P values for single guide edited non-activated cells relative

to SC were not significant, while in activated cells P = 0.02 – 0.05 relative to SC. **b,**

Representative density plots showing the percentage of TCR on the cell surface in non

activated T cells for one of the two donors. The cell lines shown are: *TCRA ΔPAR,*

TCRB ΔPAR, ΔTCR (negative control), and SC.

Extended Data Figure 9. Gating strategy for flow cytometric analysis of CD69 and

CD25 levels. Primary human T cells were gated to isolate lymphocytes, followed by

isolation of single cells, then live cells. Next, cells were gated to separate CD4+ and

CD8+ cells expressing T cell activation markers CD69 and CD25 after activation with

anti-CD3/anti-CD28 antibodies. Shown is the workflow of the FACS gating, with an

example of T cells activated with anti-CD3/anti-CD28 antibodies for 8 hours.

Extended Data Figure 10. Phenotypic analysis of CD8+ primary human T cells

with eIF3 3'-UTR PAR-CLIP sites deleted. Flow cytometric analysis of CD8+ T cells,

showing quantification of T cell activation markers CD69 (early activation marker) and

CD25 (mid-activation marker) after activation with anti-CD3/anti-CD28 antibodies at

different time points, for all cell populations tested: *1gB*, *TCRB* gRNA 1; *2gB*, *TCRB*

gRNA 2; *1+2 gB*, *TCRB* gRNA 1+2 (i.e. *TCRB ΔPAR*); *1gA*, *TCRA* gRNA 1; *2gA*, *TCRA*

gRNA 2; *1+2 gA*, *TCRA* gRNA 1+2 (i.e. *TCRA ΔPAR*); *ΔTCR* gRNA targeting the CDS

of *TCRA*; N, Non-nucleofected cells; SC, Scrambled gRNA. (*n = 2* donors, with mean

and standard deviation shown). P values for *TCRA ΔPAR, TCRB ΔPAR, ΔTCR* were <

0.0001 relative to SC, while single gRNA edited cell lines had P < 0.0001 to P < 0.01,

relative to SC as calculated by two-way ANOVA.

Extended Data Figure 11. Phenotypic analysis of CD4+ primary human T cells

with eIF3 3'-UTR PAR-CLIP sites deleted. a, Flow cytometric analysis of CD4+ T

cells, showing quantification of T cell activation markers CD69 (early activation marker)

and CD25 (mid-activation marker) after activation with anti-CD3/anti-CD28 antibodies at

different time points, for all cell populations tested: *1gB*, *TCRB* gRNA 1; *2gB*, *TCRB*

gRNA 2; *1+2 gB*, *TCRB* gRNA 1+2 (i.e. *TCRB ΔPAR*); *1gA*, *TCRA* gRNA 1; *2gA*, *TCRA*

gRNA 2; *1+2 gA*, *TCRA* gRNA 1+2 (i.e. *TCRA ΔPAR*); *ΔTCR* gRNA targeting the CDS

of *TCRA*; N, Non-nucleofected cells; SC, Scrambled gRNA. (*n = 2* donors, with mean

and standard deviation shown). P values for *TCRA ΔPAR*, *TCRB ΔPAR, ΔTCR* were <

0.0001 relative to SC, while single gRNA edited cell lines had P < 0.0001 to P < 0.01,

relative to SC as calculated by two-way ANOVA. **b**, Representative density plots

showing the percentage of cells expressing CD69, CD69 and CD25 or CD25 only, for

one of two donors. The cell lines shown are *TCRA ΔPAR*, *TCRB ΔPAR*, *ΔTCR*

(negative control), and SC.

Extended Data Figure 12. Cytokine secretion profile of primary human T cells with

with eIF3 3'-UTR PAR-CLIP sites deleted . Quantification of **a**, secreted IL2 and **b,**

secreted IFNγ, for all cell populations tested, as determined by ELISA. Cell populations

tested were: *1gB*, *TCRB* gRNA 1; *2gB*, *TCRB* gRNA 2; *1+2 gB*, *TCRB* gRNA 1+2 (i.e.

TCRB ΔPAR); *1gA*, *TCRA* gRNA 1; *2gA,* *TCRA* gRNA 2; *1+2 gA*, *TCRA* gRNA 1+2 (i.e.

TCRA ΔPAR); *ΔTCR* gRNA targeting the CDS of *TCRA*; N, Non-nucleofected cells; SC,

Scrambled gRNA. (*n = 2* donors, with mean and standard deviation shown.) P values

for *TCRA ΔPAR*, *TCRB ΔPAR, ΔTCR* were < 0.0001 relative to SC, while single gRNA

edited cell lines had P < 0.0001 to P < 0.01, relative to SC as calculated by two-way

ANOVA.

1010

Extended Data Figure 13. TCR protein and RNA levels in Jurkat cells with eIF3 3'-

UTR PAR-CLIP sites deleted. a, Western blots measuring TCRA protein levels as a

function of time after anti-CD3/anti-CD28 activation. Cell lines are indicated by transcript

structure above the western blots. HSP90 was used as a loading control. **b**, *TCRA* and

TCRB mRNA levels in WT Jurkat cells, or Jurkat cells with the *TCRA* eIF3 3'-UTR PAR-

CLIP site deleted, as determined by qRT-PCR at different time points after anti-

CD3/anti-CD28 activation. **c**, *TCRA* and *TCRB* mRNA levels in WT Jurkat cells, or

Jurkat cells with the *TCRB* eIF3 3'-UTR PAR-CLIP site deleted, as determined by qRT-

PCR at different time points after anti-CD3/anti-CD28 activation. (*n = 2*, with mean ±

standard deviations shown), **** P < 0.0001, *** P < 0.001, ** P < 0.01 for *TCRA ΔPAR*

or *TCRB ΔPAR* relative to WT, using two-way ANOVA.

1022

Extended Data Figure 14. TCR mRNA binding to eIF3 in Jurkat cells.

Immunoprecipitation of eIF3 using an anti-EIF3B antibody[5] , followed by qRT-PCR to

quantify the amount of **a**, *TCRA* or **b**, *TCRB* mRNA bound to eIF3. Jurkat cells with or

without the eIF3 PAR-CLIP site in the 3'-UTR were analyzed after activation with anti-

CD3/anti-CD28 antibodies for 5 hours (left) and 8 hours (right). Percent mRNA bound to

anti-EIF3B beads relative to total mRNA isolated from the cells (*TCRA* or *TCRB* on the

left, *IL2* on the right). (*n = 3*, with mean and standard deviations shown). For *TCRA* and

TCRB mRNAs, P < 0.0001 relative to WT. P values for IL2 mRNAs for *ΔPAR* relative

to WT were not significant using two-way ANOVA. The schematic of the experiment is

shown in **Figure 5d**.

1033

1034 **Extended Data Figure 15. TCR clustering in Jurkat cells.** WT, *TCRA ΔPAR, TCRB*

1035 *ΔPAR* cells exhibiting TCR clustering, as determined using anti-TCRA/anti-TCRB

1036 protein staining and Airyscan confocal microscopy. Cells were activated with anti-

1037 CD3/anti-CD28 antibodies for 5 hours (left) and 8 hours (right). (n = 5 for each time

1038 point after activation for all cell lines, with representative images shown)

1039

Extended Data Figure 16. Phenotypic analysis of Jurkat cells. a, Flow cytometric

analysis of cells expressing T cell activation markers CD69 (early activation marker) and

CD25 (mid-activation marker) in WT, *TCRA ΔPAR* or *TCRB ΔPAR* Jurkat cells at

different time points after stimulation with anti-CD3/anti-CD28 antibodies. (n = 3, with

mean and standard deviations shown). Left graph, P < 0.0001 for CD69 expression in

TCRA ΔPAR and *TCRB ΔPAR* cells relative to WT cells using one-way ANOVA. Right

graph, P < 0.001 for CD69 and CD25 expression of *TCRA ΔPAR* and *TCRB ΔPAR*

relative to WT using two-way ANOVA. **b**, Density plots showing cell populations

expressing CD69+ only, CD69+ CD25+ or CD25+ only from WT, *TCRA ΔPAR* or *TCRB*

ΔPAR Jurkat cells at different time points after stimulation with anti-CD3/anti-CD28

antibodies, (n = 3, with representative plots shown). (**c**) Quantification of secreted IL2

from WT, *TCRA ΔPAR* or *TCRB ΔPAR* cells at different time points after stimulation

with anti-CD3/anti-CD28 antibodies, as determined by ELISA. (n = 3, with mean and

standard deviation shown.) P < 0.0001 for *TCRA ΔPAR* and *TCRB ΔPAR* cells relative

to WT cells, as determined by two-way ANOVA.

1055

1056

1057   **Supplementary Information**

1058   Supplementary Figures 1-3: Raw images of gels used for Fig. 1b, Fig. 3a, Extended

1059   Data Fig. 1, and Extended Data Fig. 13a.

1060

1061   Supplementary Table 1: Subunits in eIF3 crosslinked to RNA in activated Jurkat cells.

1062   Lists the eIF3 subunits, percent sequence coverage and number of identified peptides.

1063

1064   Supplementary Table 2: PARpipe statistics for eIF3 PAR-CLIP samples. Samples are

1065   indexed in the first tab, including both biological replicates for activated and non-

1066   activated Jurkat cells. Statistics are given for each library at the read, cluster and group

1067   level.

1068

Supplementary Table 3: PAR-CLIP mapping to individual genes for each eIF3 PAR-

CLIP sample. Samples are indexed in the first tab, including both biological replicates

for activated and non-activated Jurkat cells. First lists the gene name and number of

clusters identified. The statistics also include: Sum, sum of that statistic over all sites for

that gene; Med, median of that statistic for all sites for that gene; ReadCount, total reads

mapping to the gene; T2Cfraction, number of reads with T-to-C conversions / number of

reads; ConversionSpecificity, log(number of reads with T-to-C conversions / number of

reads with other conversions); UniqueReads, reads collapsed to single copies. Also

included: 5'utr/Intron/Exon/3'utr/Start_codon/Stop_codon, number of sites mapping to

that annotation category; Junction, number of sites mapping to a junction between

categories (coding-intron, coding-3'utr, etc.); GeneType, as described in the gene_type

category for this gene in the .gtf file used.

1081

1082   Supplementary Table 4. Transcriptome analysis of HEK293T cells and non-activated or

1083   activated Jurkat cells. Each tab lists transcript name and version, gene name, type of

1084   transcript, length of transcript, and mean transcripts per million, calculated from two

1085   biological replicates.

1086

1087   Supplementary Table 5. Differential expression analysis of HEK293T or non-activated

1088   vs. activated Jurkat cells. Each tab lists the gene name, mean of normalized counts,

1089 $\log_2$-fold change, $\log_2$-fold change standard error, Wald statistic, Wald test P-value, and

1090   Benjamini-Hochberg adjusted P-value.

1091

Supplementary Table 6. Pathway enrichment analysis. Lists for both biological

replicates of the EIF3A/C/B PAR-CLIP libraries are included (genes with >= 100 reads),

along with associated transcript names, lengths in nts of the 5'-UTR, coding region, and

3'-UTR, and reads normalized to the lengths of the transcript regions. Tabs also include

the top tissue-specific pathway enrichment categories determined using the STRING

Database. These list: the Gene Ontology (GO) number, GO description, observed gene

count, background gene count, false discovery rate, and matching proteins in the

network by Ensembl protein ID, and by gene name.

1100

1101   Supplementary Table 7. Reagent information for experiments. Lists include antibodies

1102   used, PCR primers, qPCR primers, gRNA targeting sequences, and FISH probes.

Figure 1

Figure 2

a

b

c

Figure 3

Figure 4

a

b

c

a

b

c

d

Extended Data Fig. 1

a

b

Extended Data Fig. 2

Extended Data Fig. 3

a

b

a

Jurkat cells
Non-activated vs activated

HEK293T vs activated Jurkat

b

Replicate 1

Replicate 2

Extended Data Fig. 5

a

b

Extended Data Fig. 6

Extended Data Fig. 8

a Activated — % TCR+

TCRA gRNA, TCRB gRNA, ΔTCR gRNA, Non-nucleofected and Scrambled gRNA

x-axis: 1 gA, 2 gA, 1+2 gA, 1 gB, 2 gB, 1+2 gB, Δ, N, SC

b Non-Activated — % TCR+

x-axis: 1 gA, 2 gA, 1+2 gA, 1 gB, 2 gB, 1+2 gB, Δ, N, SC

Legend:
- TCRA gRNA
- TCRB gRNA
- ΔTCR gRNA
- Non-nucleofected and Scrambled gRNA

c

FCS-A vs TCR

| TCRA ΔPAR | TCRB ΔPAR | ΔTCR | SC |
| 50.0 | 59.4 | 5.8 | 94.2 |

Extended Data Fig. 9

a

Lymphocytes

SSC - A / FSC - A

Lymphocytes
32.5%

Single Cells

FSC - H / FSC - A

Single cells
91.7%

Live cells

FSC - H / VL1-A Live Dead-Pacific Blue - A

Live cells
87.8%

CD4+ and CD8+ cells

YL4-A CD4 - PE-Cy7 - A / BL3-A CD8 - PerCP-Cy5.5 - A

CD4+
31.3%

C8+
53.9%

CD4+ cells CD69+ CD25+

RL1-A CD25- APC - A / YL1-A CD69 PE - A

15.2%   15.0%

57.0%   12.8%

CD8+ cells CD69+ CD25+

RL1-A CD25- APC - A / YL1-A CD69 PE - A

6.89%   24.3%

42.6%   26.2%

Extended Data Fig. 10

a

b

Extended Data Fig. 12

a

b

IL2 pg ml^{-1}

IFNγ pg ml^{-1}

a

Extended Data Fig. 14

Extended Data Fig. 15

Extended Data Fig. 16

a

% of Jurkat cells CD69+ (5 hours)

% of Jurkat cells CD69+ CD25+ (8 hours, 24 hours, 48 hours)

Legend:
- *TCRB ΔPAR*
- *TCRA ΔPAR*
- WT

b

	TCRA ΔPAR	*TCRB ΔPAR*	WT
5 hours	21.7	24.4	48.3
8 hours	15.2	26.0	42.6
24 hours	44.5	46.1	66.0
48 hours	41.6	39.5	52.9

CD25 (y-axis) / CD69 (x-axis)

c

IL2 pg ml^{-1} at 5 hour, 8 hours, 24 hours, 48 hours, 72 hours