OBERON3 and SUPPRESSOR OF MAX2 1-LIKE proteins form a regulatory module specifying phloem identity

Eva-Sophie Wallner[1,5], Nina Tonn[1], Dongbo Shi[1,2,3], Laura Luzzietti[1], Friederike Wanke[4], Pascal Hunziker[1], Yingqiang Xu[1], Ilona Jung[1], Vadir Lopéz-Salmerón[1,6], Michael Gebert[1], Christian Wenzl[1], Jan U. Lohmann[1], Klaus Harter[4], Thomas Greb[1]*

[1]Centre for Organismal Studies (COS), Heidelberg University, 69120 Heidelberg, Germany

[2]Japan RIKEN Center for Sustainable Resource Science (CSRS), Yokohama, Japan

[3]Japan Science and Technology Agency (JST), Saitama, Kawaguchi, Japan

[4]Center for Plant Molecular Biology (ZMBP), University of Tübingen, 72076 Tübingen, Germany

[5]Current address: Gilbert Biological Sciences, Stanford University, Stanford, CA 94305-5020, USA

[6]Current address: BD Bioscience, 69126 Heidelberg, Germany

*Corresponding author. Tel.: +49 6221 54 5524; Fax: +49 6221 54 6424; Email: thomas.greb@cos.uni-heidelberg.de

Abstract

Spatial specificity of cell fate decisions is central for organismal development. The phloem

tissue mediates long-distance transport of energy metabolites along plant bodies and is

characterized by an exceptional degree of cellular specialization. How a phloem-specific

developmental program is implemented is, however, unknown. Here we reveal that the

ubiquitously expressed PHD-finger protein OBE3 forms a central module with the phloem-

specific SMXL5 protein for establishing phloem identity in *Arabidopsis thaliana*. By protein

interaction studies and phloem-specific ATAC-seq analyses, we show that OBE3 and SMXL5

proteins form a complex in nuclei of phloem stem cells where they establish a phloem-specific

chromatin profile. This profile allows expression of *OPS*, *BRX*, *BAM3*, and *CVP2* genes acting

as mediators of phloem differentiation. Our findings demonstrate that OBE3/SMXL5 protein

complexes establish nuclear features essential for determining phloem cell fate and highlight

how a combination of ubiquitous and local regulators generate specificity of developmental

decisions in plants.

Keywords: sieve element, phloem, Arabidopsis thaliana, root, stem cell, ATAC-seq

Introduction

Growth and body shape of multicellular organisms largely depend on a functional long-distance transport of energy metabolites to fuel stem cell activity. In vascular plants, sugars are photosynthetically produced in source organs, such as leaves, and delivered via the phloem to sink organs where they are allocated to storage tissues or stem cell niches, such as the root apical meristem (RAM) (Oparka and Turgeon 1999; De Schepper et al. 2013). The dividing stem cells of the RAM are located next to a mostly dormant organizer, known as quiescent center (QC) (van den Berg et al. 1997). These stem cells divide and differentiate in a strictly controlled manner to give rise to two phloem poles which ensure a steady energy supply to the RAM during root growth (Rodriguez-Villalon et al. 2014; Wallner et al. 2017). One phloem pole comprises a protophloem and a metaphloem strand, each forming a sieve element (SE) and a companion cell (CC) lineage (Lucas et al. 2013). During differentiation, SEs degrade most of their organelles to build connected sieve tubes for intracellular allocation of sugars, hormones, proteins and RNAs (Furuta et al. 2014). This is why functional SEs are metabolically sustained by CCs via intercellular channels named plasmodesmata (Ross-Elliott et al. 2017). Underlining the importance of the phloem, defects in protophloem development impair root growth, possibly, as a consequence of RAM starvation (Depuydt et al. 2013; Rodriguez-Villalon et al. 2014).

Due to the remarkable transition of phloem stem cells to cells holding an extreme degree of specialization, gaining insights into phloem formation and identifying its molecular regulators is highly instructive for our general understanding of cell fate regulation and differentiation (Marhava et al. 2018; Gujas et al. 2020; Marhava et al. 2020). Moreover, due to the importance of the phloem for plant growth and physiology, revealing mechanisms of phloem formation holds great promises for crop production and may increase our understanding of plant evolution and of the adaptation to environmental conditions (López-Salmerón et al. 2019). Importantly, although several genes, including *ALTERED PHLOEM*

DEVLEOPMENT (*APL*), *OCTOPUS* (*OPS*), *BREVIS RADIX* (*BRX*), *BARELY ANY*

MERISTEM3 (*BAM3*), and *COTYLEDON VASCULAR PATTERN2* (*CVP2*) have been

identified to regulate different aspects of phloem formation (Bonke et al. 2003; Depuydt et al.

2013; Rodriguez-Villalon et al. 2014; Anne et al. 2015; Rodriguez-Villalon et al. 2015; Hazak

et al. 2017; Marhava et al. 2018), those genes seem to act downstream of phloem specification

leaving the question open of how a phloem-specific developmental program is initiated.

Recently, we revealed a central role of the SUPPRESSOR OF MAX2 1-LIKE (SMXL)

protein family members SMXL3, SMXL4 and SMXL5 in phloem formation (Wallner et al. 2017;

Wallner et al. 2020). SMXL proteins are well-conserved nuclear localized developmental

regulators and, in *Arabidopsis thaliana* (Arabidopsis), form a protein family of eight members

sub-divided into different sub-clades based on phylogeny and function (Zhou et al. 2013;

Soundappan et al. 2015; Liang et al. 2016; Wallner et al. 2016; Walker and Bennett 2017).

Among those, SMXL6, SMXL7, and SMXL8 are proteolytic targets of the strigolactone

signaling pathway which bind directly to promoter regions of downstream target genes and,

thereby, repress their transcription (Soundappan et al. 2015; Wang et al. 2015; Liang et al.

2016; Ma et al. 2017; Wang et al. 2020). In comparison, SMXL3/4/5 proteins act independently

from strigolactone signaling as central regulators of phloem formation (Wallner et al. 2017; Wu

et al. 2017; Cho et al. 2018). Their redundant and dose-dependent functions become obvious

in double and triple mutants which are completely deprived of protophloem formation within

the RAM resulting in root growth termination a few days after germination (Wallner et al. 2017).

Despite their fundamental role in phloem formation the mechanism of SMXL3/4/5 protein

action remained obscure.

In contrast to SMXL proteins whose activity is spatially highly restricted, OBERONs

(OBEs) are a family of four ubiquitously expressed, nuclear-localized proteins essential for

tissue specification and meristem maintenance starting from the earliest stages of embryo

development (Saiga et al. 2008; Thomas et al. 2009; Saiga et al. 2012; Lin et al. 2016). This

role is reflected by mutants deficient for either of the two *OBE* sub-families which are embryo

lethal (Saiga et al. 2008; Thomas et al. 2009; Saiga et al. 2012). In the shoot apical meristem

(SAM), *OBE3* (also known as *TITANIA1* (*TTA1*))*,* interacts genetically with the homeobox

transcription factor gene *WUSCHEL* (*WUS*) in stem cell regulation (Lin et al. 2016).

Additionally, *OBE1* and *OBE2* are associated with vascular patterning in the embryo (Thomas

et al. 2009). Interestingly, OBEs carry a highly conserved plant homeodomain (PHD)-finger

domain known to bind di- and trimethylated histone H3 which allows recruitment of chromatin

remodeling complexes and transcription factors (Sanchez and Zhou 2011). Indeed, OBE

proteins show chromatin binding and remodeling activities important for root initiation during

embryogenesis (Saiga et al. 2008; Saiga et al. 2012). Taken together, OBEs have versatile

roles associated with cell fate regulation in plants (Saiga et al. 2008; Thomas et al. 2009; Saiga

et al. 2012; Lin et al. 2016) but, as for SMXL proteins, their specific roles in distinct tissues and

their mode of action is unknown.

Here, we report that OBE3 and SMXL5 proteins physically interact forming a functional

unit during protophloem formation in the RAM. We provide evidence that SMXL5 and OBE3

proteins are instrumental for the establishment of a phloem-specific chromatin configuration

and for the expression of phloem-associated regulators. By characterizing the *SMXL3/4/5-*

OBE3 interaction and of phloem-specific chromatin conformation we provide insights into

molecular mechanisms of cell specification and the establishment of a highly specialized and

central plant tissue.

Results

SMXL4 and *SMXL5* promote expression of early phloem markers

To map the function of the *SMXL4* and *SMXL5* genes within the process of phloem formation, we introgressed a series of developmental markers visualizing early steps of phloem formation (Rodriguez-Villalon 2016) into the *smxl4;smxl5* double mutants showing severe defects in protophloem formation (Wallner et al. 2017). Analysis of root tips two days after germination when the overall anatomy of the *smxl4;smxl5* RAM is comparable to wild type RAMs (Wallner et al. 2017), showed that *OPS:OPS-GFP, BRX:BRX-CITRINE*, *BAM3:BAM3-CITRINE*, or *CVP2:NLS-VENUS* marker activities (Rodriguez-Villalon et al. 2014) were reduced or not detectable in *smxl4;smxl5* plants (Figure 1, A-H). This reduction was found along the entire strand of the developing protophloem and included SE-procambium stem cells located immediately proximal to the quiescent center (QC). In these founder cells of the phloem lineage, we observed accumulation of OPS-GFP and BRX-CITRINE fusion proteins in wild type which was hardly detectable in *smxl4;smxl5* double mutants (Figure 1, I-P). In contrast to markers associated with early stages of phloem development, activity of the *APL* promoter marking differentiating SEs and CCs in wild type (Bonke et al. 2003) was not detectable in root tips of *smxl4;smxl5* mutants (Supplementary Figure 1). These observations argued for an early stem cell-associated role of *SMXL4* and *SMXL5* in establishing a general phloem-specific developmental program including *OPS*, *BRX*, *BAM3*, and *CVP2* gene activities. Supporting this conclusion, *smxl4;smxl5;bam3* triple mutants showed root growth defects similar to *smxl4;smxl5* double mutants and stimulating the BAM3 pathway by CLE45 treatments had no effect on *smxl4;smxl5* roots (Supplementary Figure 1). Together with the reduced *BAM3* reporter activity (Figure 1, E,F), this indicated that the phloem-specific BAM3 pathway which counteracts phloem development (Depuydt et al. 2013) is less active in *smxl4;smxl5* mutants and not causing the developmental defects observed in *smxl4;smxl5*.

Activity of *SMXL* genes upstream of *OPS* and *BRX* is required for phloem formation

To challenge the idea of an early role of *SMXL5*, we tested the capacity of *SMXL5* to

restore protophloem formation when expressed under the control of promoters active during

different phases of protophloem development (Rodriguez-Villalon 2016). Root length served

as a fast and efficient read-out for phloem defects (Depuydt et al. 2013; Wallner et al. 2017).

Supporting the need of *SMXL5* activity during early phases of phloem development, reduced

root length and impaired SE formation usually found in *smxl4;smxl5* mutants was not observed

when they expressed *SMXL5* under the control of the early *OPS*, *BAM3*, or *CVP2* promoters

(Figure 2A, Supplementary Figure 1). In contrast, driving *SMXL5* expression by the late *APL*

promoter did not restore of root length or SE formation (Figure 2A, Supplementary Figure 1).

To see whether the reduced activity of regulators like *BRX* is causative for reduced root

length of *smxl4;smxl5* double mutants, we expressed *BRX-VENUS* in *smxl4;smxl5* mutant

backgrounds under the control of the protophloem-specific *SMXL4* promoter (Supplementary

Figure 1F) (Wallner et al. 2017). Indeed, root length of *SMXL4:BRX-VENUS/smxl4;smxl5* lines

was comparable to the length of wild type roots (Figure 2A) indicating that *BRX* acts

downstream of *SMXL4* and that reduced *BRX* activity is one reason for disturbed phloem

development in *smxl4;smxl5* mutants. Further supporting a role of the *SMXL4* and *SMXL5*

genes upstream of *OPS*, visualization of SMXL4 and SMXL5 proteins in *OPS*-deficient

backgrounds by respective reporters (Wallner et al. 2017) did neither reveal a reduced level

nor altered localization of SMXL4 or SMXL5 proteins in protophloem cells (Figure 2, B-E). This

suggested that in contrast to a positive effect of *SMXL4* and *SMXL5* on the activity of *OPS* and

BRX genes (Figure 1), *OPS* was not important for stimulating *SMXL4* or *SMXL5* activity and

that *OPS* and *SMXL* genes function at distinct steps during phloem formation.

SMXL genes act on different steps of phloem formation than _OPS_ and _BRX_

Our interpretation that _SMXL_ genes on the one side and _OPS_ and _BRX_ on the other

side act on different steps of phloem development, was confirmed by investigating their genetic

interaction. The OPS protein is required for SE formation in the protophloem by counteracting

the BAM3/CLAVATA3/ESR-related 45 (CLE45) pathway (Breda et al. 2019). Due to enhanced

activity of the BAM3/CLE45 pathway, _ops_ mutants develop 'gap cells' within protophloem

strands in which SE establishment fails (Truernit et al. 2012; Rodriguez-Villalon et al. 2014).

Interestingly, root length of _ops;smxl5_ double mutants was similarly reduced as in _smxl4;smxl5_

double mutants although _smxl5_ single mutants were not affected and _ops_ single mutants were

very variable in this regard (Figure 3A). This finding suggested an additive effect of _SMXL_ and

OPS-dependent pathways on phloem formation. Indeed, when phloem development was

carefully analyzed in the respective mutant backgrounds, we observed a variation of phloem

defects in _ops_ single mutants ranging from the appearance of gap cells to the complete

absence of SEs in a small fraction of plants (Figure 3B, Supplementary Figure 2). In

comparison, 60 % of the _ops;smxl5_ mutants displayed complete SE deficiency demonstrating

that both genes contribute to a robust phloem development. A similar trend was observed for

brx;smxl5 double mutants. Like _OPS_, the _BRX_ gene ensures continuous SE formation, in this

case, by downregulation of _BAM3_ transcription and steepening the auxin gradient in

developing phloem cells (Scacchi et al. 2009; Depuydt et al. 2013; Marhava et al. 2018;

Marhava et al. 2020). Similar to _ops;smxl5_ double mutants, _brx;smxl5_ plants developed shorter

roots than _brx_ and _smxl5_ single mutants and largely failed to differentiate SEs (Figure 3A,

Supplementary Figure 3). Importantly, in those _ops;smxl5_ plants developing SEs, gap cell

formation was comparable to _ops_ and _brx_ single mutants (Figure 3B, Supplementary Figures

1). These observations suggested that, although all three genes are important for stable SE

formation, _SMXL5_ and _OPS_/_BRX_ genes play roles at different steps during phloem formation

with _SMXL5_ acting upstream.

193   **SMXL5 proteins interact with OBE3 proteins in nuclei of plant cells**

194   To find indications for how SMXL proteins fulfil their early role, we isolated interacting

195   proteins by a Yeast-Two-Hybrid-based screen of a cDNA expression library generated from

196   *Arabidopsis* seedlings (Legrain et al. 2001; Cifuentes-Esquivel et al. 2013) using the full length

197   SMXL5 protein as a bait. After testing 84 million individual protein-protein interactions, we

198   identified OBE3 as one out of five 'very high confident' candidate interactors with 24 isolated

199   independent cDNA clones (Supplementary Figure 4). After confirming the yeast-based

200   interaction of OBE3 with SMXL5 in independent experiments (Figure 4A), we tested whether

201   both proteins also interact *in planta*. To this end, we transiently expressed SMXL5 fused to a

202   triple human influenza hemagglutinin (HA) affinity tag and OBE3 fused to a six fold c-Myc

203   epitope tag in *Nicotiana benthamiana* (*N. benthamiana*) leaves under the control of the

204   Cauliflower Mosaic Virus (CaMV) *35S* promoter (Benfey and Chua 1990). In raw protein

205   extracts before ('input') and after ('unbound') immunoprecipitation (IP) using HA-affinity beads

206   and in the precipitate itself ('IP: α HA'), the SMXL5-3xHA protein was detected with the

207   expected size of approximately 120 kDa in Western analyses (Figure 4B). Importantly, the

208   6xMyc-OBE3 fusion protein co-immunoprecipitated with the SMXL5-3xHA protein and did not

209   show unspecific binding to the HA-affinity beads, indicating that SMXL5-3xHA and 6xMyc-

210   OBE3 proteins interacted in plant cells.

211

212   To compare the sub-cellular localizations of SMXL5 and OBE3 proteins, we transiently

213   expressed the SMXL5 protein fused to monomeric Cherry (SMXL5-mCherry) together with the

214   OBE3 protein fused to monomeric GFP (OBE3-mGFP) again in *N. benthamiana* leaves.

215   Initially, nuclear localization of the OBE3 protein was confirmed by co-expressing OBE3-GFP

216   with mCherry fused to a nuclear localization signal (mCherry-NLS). Interestingly, while the

217   mCherry-NLS signal was homogenously distributed within the nucleus, the OBE3-mGFP

218   protein appeared in nuclear subdomains (Figure 4, C-E). Co-expression of SMXL5-mCherry

219   and OBE3-GFP revealed a co-localization of both proteins within these domains (Figure 4, F-

H) which were distinct from the whole nucleus highlighted by an *mGFP-mCherry-NLS* fusion

protein expressed under the control of the *ubiquitin 10* (*UBI10*) promoter (Figure 4, I-K). Next,

we evaluated our yeast two-hybrid and co-immunoprecipitation data by performing Förster

resonance energy transfer (FRET)-fluorescence lifetime imaging microscopy (FLIM) analysis

as an *in planta* assay for protein-protein association. In transiently transformed *N. benthamiana*

epidermal leaf cells, FRET-FLIM analysis detected a significant decrease in the lifetime of the

donor OBE3-mGFP fusions in the nucleus when co-expressed with SMXL5-mCherry (Figure

4L, M). In contrast, we did not observe significant mGFP lifetime changes when OBE-mGFP

was co-expressed with NLS-mCherry (Figure 4L, M). Taken these observations together, we

concluded that OBE3 interacts directly with SMXL5 in plant cell nuclei.

The *OBE3* gene acts together with *SMXL3*, *SMXL4* and *SMXL5*

Since physical interaction and subcellular co-localization suggested a common action

of SMXL5 and OBE3 proteins, we investigated whether the corresponding genes are

functionally connected by again using root length as a first read-out for potential phloem

defects. As before, *smxl4;smxl5* double mutants were short rooted, while root lengths of *smxl4*

and *smxl5* single mutants were similar to wild type (Wallner et al. 2017) (Figure 5, A and B).

Similarly, roots from *obe1, obe2*, *obe3* and *obe4* single mutants resembled wild type roots. In

contrast, *smxl4;obe3, smxl5;obe3* and *smxl3;obe3* double mutants had short roots just as

smxl4;smxl5 (Figure 5, A-D) suggesting a concerted action of *OBE3* and *SMXL3*, *SMXL4* or

SMXL5 genes during primary root growth. Of note, we only detected a genetic interaction

between *SMXL3/4/5* and *OBE3* and not between *SMXL4/5* and other *OBE* family members

(Supplementary Figure 5).

OBE3 locally promotes early stages of phloem development

Because reduced root length of *obe3;smxl* double mutants suggested a role of *OBE3*

in phloem development, we next tested whether *OBE3* is expressed in developing phloem cells

by comparing the activity pattern of a translational *SMXL4:SMXL4-YFP* reporter (Wallner et al.

2017) with patterns of a translational *OBE3:OBE3-GFP* reporter (Saiga et al. 2012) (Figure 6,

A-D). As reported previously (Wallner et al. 2017), the SMXL4-YFP protein accumulated

specifically in nuclei of the protophloem lineage identified by enhanced Direct Red 23 staining

(Figure 6A). In comparison, the *OBE3:OBE3-GFP* reporter revealed OBE3-GFP protein

accumulation in nuclei of all cell types of the root tip including developing protophloem cells

(Figure 6B) expressing SMXL3, SMXL4 and SMXL5 proteins (Figure 6A) (Wallner et al. 2017).

We thus concluded that SMXL3/4/5 and OBE proteins had the potential to interact during early

phases of phloem formation. We could not detect differences in activity patterns between

OBE3:OBE3-GFP and *OBE4:OBE4-GFP* (Saiga et al. 2012) reporters (Figure 6, B-C) which

argues against the possibility that differences in expression are the reason why *OBE3*, but not

OBE4, genetically interacted with *SMXL3/4/5* genes.

To evaluate whether growth defects observed in *smxl;obe3* roots are correlated with

the same type of protophloem defects observed in *smxl4;smxl5* mutants, we analyzed phloem

development in the respective genetic backgrounds. During protophloem development, SE

procambium-precursors divide periclinally to give rise to procambium and SE precursor cells.

After 2-3 anticlinal divisions, SE precursor cells divide again periclinally to initiate meta- and

protophloem cell files that subsequently undergo gradual differentiation (Figure 6D) (Wallner

et al. 2017). Our analysis revealed that in *obe3* mutant both the periclinal cell divisions and the

onset of SE differentiation appeared as in wild type roots (Figure 6, E, F, K). In contrast, in

smxl4;obe3 and *smxl5;obe3* mutants, the onset of the second periclinal division initiating meta-

and protophloem cell lineages was similarly delayed as in *smxl4;smxl5* mutants and the

enhanced mPS-PI staining visualizing differentiated SEs was likewise absent (Figure 6 G-I, K

and Supplementary Figure 6). This observation demonstrated that, like the more locally

expressed *SMXL3/4/5* genes, *OBE3* substantially contributes to phloem formation. Moreover,

because respective single mutants did not show these defects (Figure 6 and Supplementary

Figure 6), we concluded that *OBE3* or *SMXL5*-deficient plants represent sensitized

backgrounds for the functional loss of the other regulator. In accordance with the wild type-like

root growth of those plants, protophloem formation in *smxl5;obe4* double was indistinguishable

from wild type (Figure 6, E and J).

Because *OBE3* is broadly expressed, phloem defects observed in *smxl5;obe3* mutants

could arise due to a function of *OBE3* in other tissues than developing phloem cells, which

would contradict a direct interaction of SMXL5 and OBE3 proteins. To address this concern

and to determine whether *OBE3* acts cell-autonomously on phloem development, we

expressed *OBE3* exclusively in developing phloem cells by introducing a transgene driving an

OBE3-turquoise fusion protein under the control of the *SMXL5* promoter (*SMXL5:OBE3-*

turquoise) into a *smxl5;obe3* double mutant background. Microscopic analysis of root tips from

smxl5;obe3/SMXL5:OBE3-turquoise lines confirmed the presence of the OBE3-turquoise

protein in nuclei of developing protophloem cells (Figure 7, A and B) as described for the

SMXL5 protein expressed under the control of the same promoter (Wallner et al. 2017). When

comparing *smxl5;obe3/SMXL5:OBE3-turquoise* lines with *smxl5;obe3* mutants, we observed

that expression of *OBE3-turquoise* within the *SMXL5* domain was indeed sufficient to restore

root length in *smxl5;obe3* double mutants (Figure 7, C and D). Additionally, enhanced Direct

Red staining of the mature protophloem indicating SE differentiation was recovered in

smxl5;obe3 carrying the *SMXL5:OBE3-turquoise* transgene (Figure 7, A and B). To see

whether the predominant reduction of *OBE3* activity in developing phloem cells is furthermore

sufficient for generating phloem defects, we designed two artificial microRNAs targeting the

OBE3 mRNA (*obe3-miRNAs*) (Schwab et al. 2006) and expressed them independently under

the control of the *SMXL5* promoter in *smxl5* mutant plants. As expected, the majority of those

plants was short rooted, indicating that *OBE3* knock-down within the *SMXL5* domain was

sufficient to evoke a *smxl5;obe3*-like phenotype in *smxl5* mutants (Figure 7, C and D). We thus

concluded that *OBE3* fulfils a cell autonomous and *SMXL5*-dependent role in protophloem

formation.

SMXL and OBE3 genes determine chromatin structure in phloem cells

To probe the common role of SMXL and OBE3 proteins and their putative effect on

chromatin signatures (Saiga et al. 2012; Ma et al. 2017; Wang et al. 2020), we performed

phloem-specific Assays for Transposase-Accessible Chromatin using sequencing (ATAC-seq)

revealing chromatin structure in a genome-wide fashion (Buenrostro et al. 2015). To this end,

we fluorescently labeled nuclei in phloem-associated cells in wild type, *smxl5*, *smxl4;smxl5*

and *smxl5;obe3* plants by expressing a histone H4-GFP protein fusion under the control of the

SMXL5 promoter (*SMXL5:H4-GFP*, (Shi et al. 2019), Supplementary Figure 7A). Isolation of

nuclei from those lines and their fluorescence-activated sorting into GFP-positive and GFP-

negative populations (FANS (Shi et al. 2021), Supplementary Figure 7B) allowed ATAC-seq

analyses separately on phloem-related and non-phloem-related cells. Thereby, we detected

36,925 to 40,168 open chromatin regions (OCRs) in all sample types mostly located in 5' and

3' regions of respective gene bodies (Figure 8A, B, Supplementary Figures 8, 9 and 10).

Importantly, the overall conformation was similar in all samples (Supplementary Figure 8)

suggesting, by large, a comparable chromatin structure in phloem and non-phloem cells and

an independency of most chromatin domains from *SMXL4*, *SMXL5* or *OBE3* activity. Among

38,562 OCRs detected in phloem cells from wild type, 1488 OCRs associated with 1072 genes

showed a significant increase of read alignment in comparison to non-phloem cells (Figure 8A;

Supplementary Dataset 1 and 2; fold change > 2, poisson enrichment *p*-value < 0,05, using 40

M reads). These genes included OCRs in promoter regions of the phloem-specific *OPS*, *BAM3*,

CVP2, and *APL* genes (Supplementary Figures 9 and 10, Supplementary Dataset 1 and 2)

suggesting that we succeeded in phloem-specific ATAC-seq analysis. In accordance with the

wild type-like phenotype of *smxl5* mutants, 1055 of 36,925 OCRs associated with 809 genes

showed a significant increase of read alignment in the *smxl5* phloem in comparison to non-

phloem cells in *smxl5* again including *OPS*, *BAM3*, *CVP2*, and *APL* (Figure 8A, Supplementary

Figures 9 and 10, Supplementary Datasets 1 and 2). In contrast, only 48 and 316 phloem-

associated OCRs were detected in *smxl4;smxl5* and *smxl5;obe3* double mutants, respectively

(Figure 8A, Supplementary Datasets 1 and 2), suggesting a loss of phloem-specific chromatin

signatures in both backgrounds.

This conclusion was also supported by direct comparison of the OCR profile obtained

from GFP-positive samples. In total, only 60 OCRs showed significantly more or fewer aligned

reads comparing *smxl5* mutants with wild type (Figure 8C, Supplementary Dataset 3, fold

change >2, poisson enrichment *p*-value < 0.05, using 40 M reads) arguing for a high similarity

of the chromatin profile in phloem-related cells from both backgrounds. In contrast,

smxl4;smxl5 and *smxl5;obe3* double mutants differed considerably from both wild type and

smxl5 mutants with regard to their phloem-related chromatin profile. *smxl4;smxl5* mutants

showed 3283 OCRs in total with significant differences in read alignment in comparison to wild

type and, for *smxl5;obe3* mutants, 1902 differential OCRs were found. A similar situation was

found for both double mutants in comparison to *smxl5* mutants (Figure 8C, Supplementary

Dataset 3). Importantly, the number of differential OCRs was lower when comparing

smxl4;smxl5 and *smxl5;obe3* samples directly. In this case, only 575 differential OCRs were

detected (Figure 8C) suggesting that both double mutants differed similarly from wild type and

smxl5 mutants with regard to their chromatin profile in phloem-related cells. Beyond the overall

OCR profile, the chromatin around phloem-related genes like *OPS*, *BAM3*, *CVP2*, and *APL*

displayed a more condensed conformation in GFP-positive nuclei from *smxl4;smxl5* and

smxl5;obe3 and the difference between GFP-positive and -negative nuclei was less

pronounced in these cases (Supplementary Figure 9 and 10).

When comparing the OCR profile obtained from GFP-negative samples, the difference

between the genetic backgrounds was substantially lower with a maximum of 141 differential

OCRs comparing *smxl5* and *smxl5;obe3* backgrounds (Figure 8D, Supplementary Dataset 4).

Together, these comparisons suggested a considerable and mostly phloem-specific difference

between chromatin conformation in phloem-related cells when comparing wild type and *smxl5*

mutants on the one side and *smxl4;smxl5* and *smxl5;obe3* double mutants on the other side.

Moreover, we concluded that phenotypic changes are similar in *smxl4;smxl5* and *smxl5;obe3*

mutants, underlining the functional relatedness of *SMXL* and *OBE3* genes.

To see how chromatin conformation differed overall between the different genotypes,

we defined a group of 'phloem' genes (combined SUC2, APL and S32 domain genes according

to (Brady et al. 2007), Supplementary Dataset 5) and 'non-phloem' genes (combined M1000,

cortex, COBL9, GL2, AGL42, PET111, and LRC domain genes according to (Brady et al.

2007), Supplementary Dataset 5). When analyzing chromatin conformation associated with

the different gene groups, a similar pattern in wild type and *smxl5* mutants was observed with

a more open chromatin around transcriptional start sites (TSS) of non-phloem genes in GFP-

negative samples compared to GFP-positive samples (Figure 8B). Interestingly, in GFP-

positive samples from *smxl4;smxl5* and *smxl5;obe3* double mutants, overall chromatin

conformation of phloem genes was similar to wild type. In contrast, chromatin of non-phloem

genes was more open around TSSs in GFP-positive samples and the conformation resembled

the pattern in GFP-positive samples (Figure 8B). This observation again indicated that phloem-

specific chromatin signatures were equally reduced in phloem-related cells of *smxl4;smxl5* and

smxl5;obe3 double mutants reflecting a comparable function of both SMXL4/5 and OBE3

proteins.

Discussion

Cell type specification is fundamental for establishing multicellular organisms and, in recent years, the phloem has become an instructive model for studying this aspect in plants (Anne and Hardtke 2017; Blob et al. 2018; Gujas et al. 2020; Marhava et al. 2020). With our study we provide new insights into the regulation of (proto)phloem formation by revealing a role of the putative chromatin remodeling protein OBE3 (Saiga et al. 2012) and a direct interaction between the OBE3 protein and the central phloem regulator SMXL5. Based on our findings, we propose that SMXL proteins fulfil their role in phloem formation by creating a distinct chromatin signature important for establishing phloem identity.

SMXL3/4/5 and OBE3 proteins act together on early stages of phloem development

SMXL3/4/5 and OBE3 proteins are already expressed in phloem stem cells (Wallner et al. 2017), which was so far not described for other phloem regulators (Blob et al. 2018). Here, we detected OPS and BRX protein accumulation already in those stem cells raising the question of functional interdependence. The positive effect of *SMXL4* and *SMXL5* gene functions on OPS and BRX protein accumulation in those and more mature phloem cells suggests that *SMXL* genes are required for the establishment of a phloem-specific developmental program including *OPS* and *BRX* gene activities. The different subcellular localization of OPS, BRX and BAM3 proteins on the one side and SMXL proteins on the other side argues against a more interconnected mechanism of both groups of regulators. The conclusion that both groups act on different aspects of phloem formation is furthermore supported by our genetic analyses which revealed a combination of distinct phloem defects in respective double mutants. Here, we propose that SMXL proteins fulfil their role in the nucleus of phloem stem cells and beyond by direct interaction with OBE3. In fact, the *SMXL3/4/5-OBE3* interaction seems to be a prerequisite for both the initiation of phloem cell fate and a timely onset of differentiation. On the genetic level, *smxl4;obe3*, *smxl5;obe3* and the *smxl4;smxl5* double mutants share the same phloem defects meaning that they are deprived of protophloem

formation within the RAM. Other reported phloem mutants have either problems with

completing phloem differentiation in general, which is the case in mutants of the *ALTERED*

PHLOEM DEVELOPMENT (*APL*) gene (Bonke et al. 2003; Truernit et al. 2008; Kondo et al.

2016), or they develop 'gap cells' in which SE-differentiation is disturbed as in *ops* or *brx* single

or in *cvp2;cvp2-like1* (*cvl1*) double mutants (Depuydt et al. 2013; Anne et al. 2015; Rodriguez-

Villalon et al. 2015; Marhava et al. 2018; Breda et al. 2019). In contrast, *smxl5;obe3*,

smxl4;obe3 and multiple mutants of the *SMXL3/4/5* genes show absence of all morphological

hallmarks of phloem formation within the RAM. This suggests that *SMXL3/4/5* and *OBE3* act

together on the establishment of a phloem-specific developmental program. Interestingly,

complete suppression of protophloem formation was so far only reported for roots treated with

certain CLAVATA3/ESP-RELATED (CLE) peptides, such as CLE45, which signal through the

leucine-rich repeat receptor like kinase BAM3 and the pseudokinase CORYNE (CRN)

(Depuydt et al. 2013; Hazak et al. 2017). However, the formation of 'gap cells' in *ops* or *brx*

mutants is suppressed in *BAM3*-deficient backgrounds (Rodriguez-Villalon et al. 2014). This

shows that none of those factors is required to obtain protophloem cell identity and proper

differentiation in the first place (Depuydt et al. 2013; Rodriguez-Villalon et al. 2014). Moreover,

the epistatic relationship between *SMXL4/5* and *BAM3* genes as observed in this study shows

that the SMXL and BAM3 pathways are functionally independent.

A putative role of a SMXL3/4/5/OBE3 protein complex in chromatin remodeling

PHD-finger motifs as carried by the OBE3 protein are known to be epigenetic readers

binding to histone H3 tails carrying distinct post-translational modifications such as

trimethylation of lysine 4 (H3K4me3) or lysine 9 (H3K9me) marking actively transcribed or

silent chromatin regions, respectively (Sanchez and Zhou 2011). Although PHD finger proteins

themselves are not necessarily activating or repressing, they can indirectly modify transcription

by recruiting chromatin modifying complexes (De Lucia et al. 2008). Indeed, OBE proteins

have been proposed to remodel chromatin structure during embryogenesis and thereby

transcriptionally activate RAM initiation factors (Saiga et al. 2012). SMXL proteins share an

ethylene-responsive element binding factor-associated amphiphilic repression (EAR) motif

which interacts with transcriptional regulators of the TOPLESS (TPL) family (Pauwels et al.

2010) and they have recently shown to bind DNA directly (Wang et al. 2020). Indeed, the

strigolactone signaling mediators SMXL6, SMXL7, SMXL8 and DWARF53 (D53), an SMXL

protein from rice, directly interact with TPL-like proteins (Soundappan et al. 2015; Wang et al.

2015; Ma et al. 2017). TPLs are transcriptional co-repressor which can recruit histone

deacetylases (HDAC) and, thereby, induce chromatin condensation and transcriptional

suppression (Krogan and Long 2009; Ma et al. 2017). In addition to EAR motifs, SMXLs share

a conserved double caseinolytic protease (Clp) domain with ATPase activity that resembles

heat shock protein 101 (HSP101) (Jiang et al. 2013; Zhou et al. 2013). As recently proposed,

the p-loop ATPase domain of D53 fosters the formation of TPL hexamers and the threading of

DNA through a central pore of this hexamer inducing nucleosome repositioning and/or higher-

order chromatin reorganization (Ma et al. 2017).

Judging from the protein domains, their sub-nuclear localization and the similar loss of

phloem-specific chromatin signatures in *smxl4;smxl5* and *smxl5;obe3* mutants, we

hypothesize that SMXL3/4/5 and OBE3 form a protein complex that is involved in chromatin

remodeling and/or transcriptional regulation of downstream targets. In summary, our study

reveals the functional interaction between *SMXL3/4/5* and *OBE3*, an important step in

understanding plant tissue formation.

Material and methods

Plant material

Genotypes of plant species *Arabidopsis thaliana* (L.) Heynh. of the ecotype Columbia

(Col) used for genetic analysis are listed in the Key Resource Table. Sterile seeds were

stratified in microcentrifuge tubes containing dH$_2$O at 4 °C in the dark for 3 days and then sown

in rows on ½ Murashige and Skoog (MS) medium-plates supplemented with 1 % sucrose and

grown vertically. Seedlings were grown in long day (LD, 16 h light and 8 h dark) conditions at

21 °C for 2-10 days. Flowering plants were transformed by the floral dip method described

earlier (Clough and Bent 1998). Seeds were liquid sterilized by 70 % ethanol supplemented

with 0.2 % Tween-20 for 15 min, washed twice with 100 % ethanol and air dried under sterile

conditions. *bam3-2* (SALK_044433), *obe1-1* (SALK_075710), *obe3-2* (SALK_042597) and

obe4-1 (SALK_082338) mutants were obtained from the NASC stock center. *smxl3-1*

(SALK_024706), *smxl4-1* (SALK_037136), *smxl5-1* (SALK_018522), and *obe2-2* mutants

were described before (Thomas et al. 2009; Wallner et al. 2017). *smxl4-1;smxl5-1;bam3-3*

triple mutants were generated by CRISPR/Cas9 targeting *BAM3* in a *smxl4-1;smxl5-1* mutant

background (creating the *bam3-3* allele), leading to a G insertion after position 707 of the CDS

and to the generation of a stop codon after 255 aa of the BAM3 protein (i.e. in the LRR receptor

domain). All other lines are referenced in the text.

Yeast-Two-Hybrid

The yeast-based screen for proteins interacting with SMXL5 was performed by

Hybrigenics (Evry, France) as described before (Legrain et al. 2001). The yeast strain *AH109*

was used for the yeast two-hybrid assay according to Matchmaker™ Two-Hybrid System 3

(Clontech, Palo Alto) and grown on YPD (full medium) or SD (selective medium)-agar plates

for 3-5 days at 28°C, then stored at 4°C and stroked onto new plates every 10 days. Dilution

series (OD600 1-0.001) of transformed yeast strains were grown for 3 days on selective

medium (SD) -Leu/-Trp/-His/-Ade selecting for protein interaction or SD -Leu/-Trp selecting for

the presence of the plasmids. When grown in liquid YPD medium for transformation, yeast was

grown over night at 28 °C with shaking at 250 rpm. For expressing the GAL4BD-SMXL5 fusion

protein in yeast the open reading frame of the *SMXL5* gene was cloned into XmaI/BamHI sites

of the *pGBKT7* plasmid (Clontech, Palo Alto) resulting in *pEW6*. For expressing the GAL4AD-

OBE3 fusion protein, the open reading frame of the *OBE3* gene was cloned into XmaI/BamHI

sites of the *pGADT7* plasmid (Clontech, Palo Alto) resulting into *pEW11*.

Agrobacterium tumefaciens

The *Agrobacterium tumefaciens* genotypes C58C1: RifR with pSoup plasmid (TetR) or

ASE: KanR, CamR with pSoup+ plasmid (TetR) were used for transformation of *Arabidopsis*

thaliana or infiltration of *N. benthamiana* leaves and grown at 28 °C over night in liquid YEB

medium on a shaker (180 rpm to an OD_{600} > 1) or plated on YEB-plates and grown in an

incubator (Fraley et al. 1983; Ashby et al. 1988; Hellens et al. 2000). Antibiotics were used for

plasmid selection.

Root length measurements

For measuring root lengths, seedlings were scanned by a commercial scanner and

analyzed using ImageJ 1.49d (Schindelin et al. 2012). For CLE45 treatments, plants were

germinated on normal MS media (mock) or media containing 50 nM CLE45 and root length

measurements were performed after five days.

Genotyping

Genotyping was performed by PCR using primers listed in Supplementary Table 1.

Further information about standard DNA extraction and genotyping can be found in (Wallner

2018).

Transient protein expression in *N. benthamiana*

N. benthamiana plants were used for transient protein expression and grown in the

greenhouse at approximately 25 °C and watered daily. Transformed *Agrobacteria* were stored

as glycerol stocks and grown in a 10 ml YEB liquid culture prior to use. The densely grown

culture was centrifuged at 4000 rpm for 5 min at RT. The supernatant was removed and the

pellet was washed with 5 ml induction buffer and re-suspended in 10 ml induction buffer.

Culture densities were adjusted to an OD_{600} of 1. Prior to infiltration, these bacterial solutions

were mixed with *Agrobacteria* expressing *35S:P19* in a ratio 1:2 and incubated in the dark for

2-3 h (Voinnet et al. 2003; Scholthof 2006). *N. benthamiana* leaves were infiltrated with the

mixtures using a 1 ml syringe (Becton Dickinson S.A., Heidelberg, Germany). Leaves were

harvested three days after infiltration.

Protein extraction, immunoprecipitation and Western blot

Infiltrated *N. benthamiana* leaves were frozen in liquid nitrogen and ground by a mortar.

Proteins were extracted by mixing the leaf powder 1:1 with extraction buffer (50 mM Na_3PO_4,

150 mM NaCl, 10% glycerol, 5 mM EDTA, 10 mM β-mercaptoethanol, 0.1% triton X-100, 2

mM $NaVO_4$, 2 mM NaF, 20 μM MG-132, 1 mM PMSF, 1x cOmpleteTM Protease Inhibitor

Cocktail (Roche; Basel, Switzerland)). Each sample was vortexed for 10 sec and centrifuged

at 13000 rpm for 10 min at 4 °C. The protein extract was retrieved by sieving it through a nylon

mesh. Protein quantities were measured by Bradford assays according to the manual provided

with the Bio-Rad Protein Assay Dye Reagent Concentrate (Bio-Rad Laboratories; Hercules,

533 USA). Proteins were immunoprecipitated by 50 µl Anti-HA MicroBeads (Miltenyi Biotec,

534 Bergisch Gladbach, Germany) after incubation for 2.5 h at 4 °C while slowly rotating. Beads

535 were captured by µ Columns (Miltenyi Biotec, Bergisch Gladbach, Germany) on magnetic

536 stands by following the user manual and washed three times by 200 µl Wash buffer I

537 (extraction buffer without β-mercaptoethanol) and two times by 200 µl Wash buffer II (50 mM

538 Na_3PO_4, 150 mM NaCl, 10% glycerol, 5 mM EDTA). Proteins were eluted by 2x Laemmli buffer

539 (95 °C) and separated by size on a SDS-PAGE with subsequent western blotting. Detailed

540 procedures can be found in (Wallner 2018). SMXL5-3xHA and 6xMyc-OBE3 bands were

541 detected by antibodies Anti-HA-Peroxidase High Affinity (3F10) (Roche; Basel, Switzerland)

542 or c-Myc Antibody (9E10) sc-40 HRP (Santa Cruz Biotechnology, Santa Cruz, USA),

543 respectively and visualized by chemiluminescence agents SuperSignal™ West Femto

544 Maximum Sensitivity Substrate (Thermo-Scientific; Waltham, USA) by an Advanced

545 Fluorescence and ECL Imager (Intas Science Imaging Instruments, Göttingen, Germany).

FRET-FLIM analyses

548 FRET-FLIM analyses were performed principally as described previously (Ladwig et al.

549 2015; Arongaus et al. 2018). Briefly, measurements were performed using a Leica TCS SP8

550 microscope (Leica Microsystems, Germany) equipped with a rapidFLIM unit (PicoQuant).

551 Images were acquired using a 63x/1.20 water immersion objective. For the excitation and

552 emission of fluorescent proteins, the following settings were used: mGFP at excitation 488 nm

553 and emission 500-550 nm; and mCherry at excitation 561 nm and emission 600-650 nm. The

554 lifetime τ [ns] of either the donor only expressing cells or the cells expressing the indicated

555 combinations was measured with a pulsed laser at an excitation light source of 470 nm and a

556 repetition rate of 40 MHz (PicoQuant Sepia Multichannel Picosecond Diode Laser, PicoQuant

557 Timeharp 260 TCSPC Module and Picosecond Event Timer). The acquisition was performed

558 until 500 photons in the brightest pixel were reached. To obtain the GFP fluorescence lifetime,

559 data processing was performed with SymPhoTime software and bi-exponential curve fitting

and a correction for the instrument response function. Statistical analysis was carried out using

the JMP 14 software (JMP, USA).

Direct Red 23 staining

To preserve fluorescent signals in roots, seedlings were fixed in a vacuum chamber for

1 h by 4 % (w/v) PFA dissolved in PBS. The tissue was washed twice by PBS and cleared with

ClearSee solution for a minimum of two days according to (Kurihara et al. 2015). Cleared

seedlings were stained by 0.01 % (w/v) Direct Red 23 in ClearSee solution for 1 h. Excess

staining was removed by clearing once again in pure ClearSee solution for 1 h.

mPS-PI staining

The mPS-PI staining of roots was carried out as described before (Truernit et al. 2008).

Confocal microscopy

For confocal microscopy, TCS SP5 or SP8 microscopes (Leica Microsystems;

Mannheim, Germany) were used. GFP signals were excited at excited by an argon laser at

488 nm, collecting the emission between 500-575 nm. YFP was excited by an argon laser at

514 nm and the emission detected in a range of 520-540 nm. DirectRed stained tissue was

excited at 561 nm (DPSS laser) and emission was detected at wavelengths >660 nm. mPS-PI

stained tissue was excited at 561 nm (DPSS laser) and emission was detected at 590-690 nm.

Molecular cloning and miRNA generation

OPS:SMXL5-VENUS (*pNT52*), *OPS:ER-VENUS* (*pNT53*), *BAM3:SMXL5-VENUS*

(*pNT49*), *BAM3:ER-VENUS* (*pNT50*), *CVP2:SMXL5-VENUS* (*pNT16*), *CVP2:ER-VENUS*

(*pNT69*), *APL:SMXL5-VENUS* (*pNT10*), *APL:ER-VENUS* (*pNT68*), *SMXL4:BRX-VENUS*

(*pNT72*), *SMXL5:OBE3-turquoise* (*pEW72*)*, 35S:5xc-Myc-OBE3* (*pEW78*)*, 35S:SMXL5-*

mCherry (*pVL122*)*, 35S:OBE3-mGFP* (*pVL127*)*, 35S:mCherry-NLS* (*pMG103*) and

UBI10:mGFP-mCherry-NLS (*pCW194*) constructs were generated by using appropriate

modules according to the GreenGate manual described in (Lampropoulos et al. 2013).

Destination modules, entry modules, and correlating primers for amplifying DNA fragments for

generating entry modules are depicted in Supplementary Table 1. In case reporter proteins

were targeted to the endoplasmatic reticulum (ER), they were fused to the appropriate motifs

(Haseloff et al. 1997). *miRNAs* targeting *OBE3* transcripts were designed and cloned

according to the manual provided by the WMD3 - Web MicroRNA Designer Version 3 (Max

Planck Institute for Developmental Biology, Tübingen. http://www.weigelworld.org) with

primers listed in Supplementary Table 1. To generate *35S:SMXL5-3xHA* (*pEW33*), the *SMXL5*

CDS was amplified by primers listed In Supplementary Table 1 and cloned into BamHI/XbaI

sites of *pGreen0229:35S* (Hellens et al. 2000) resulting in *pKG33*. Next, ssDNA sequences

coding for 3xHA (Supplementary Table 1) were annealed by gradual cool-down from 80 °C to

50 °C and inserted into vector *pKG33* using BamHI/XmaI sites, resulting in plasmid *pEW31*.

Further information about detailed cloning procedures can be found in (Wallner 2018).

Fluorescence Activated Nucleus Sorting (FANS) and ATAC-seq

Nucleus extraction for FANS/ATAC-seq was based on (Thibivilliers et al. 2020) and

carried out on ice. Approximately 1 gr of three week-old Arabidopsis seedlings grown in ½ MS

LD conditions were collected in a 60 mm petri dish and thoroughly chopped using a razor blade

for five minutes in nucleus isolation buffer (1x NIB, CelLytic™-PN Isolation/Extraction buffer,

Sigma-Aldrich, cat.no. CELLYTPN1) supplemented with 10 µg/ml Hoechst 33342 (Sigma-

Aldrich, B2261) as the final concentration. After incubation for 15 min at 4°C in darkness, the

nucleus suspension was applied to 50 µm nylon strainers mounted into a 30 µm nylon strainers

at the top (Sysmex, CellTrics), and filtered further by a FACS tube with 35 µm strainer cap

(Corning, #352235). 15,000 nuclei for each sample were then sorted according to their GFP

signal levels into a 300 μL collection buffer (15 mM TRIS-HCl ph 7.5, 20 mM NaCl, 80 mM KCl,

0.1 % Triton) by a BD FACSAriaTM IIIu cell sorter using a 100 μm sort nozzle and a 30 kHz

drop drive frequency. The gate for GFP+ nuclei was set using wild type as a reference

(Supplementary Figure 7B). Next, samples were centrifuged at 3000 g for 10 minutes at 4°C.

The supernatant was partially removed and the nuclei resuspended in 300 μL Tris-Mg buffer

(Tris HCl pH8 10 mM, MgCl$_2$ 5 mM). A second washing step was performed using

centrifugation at 1000 g for 10 minutes at 4°C and the supernatant was removed completely.

For tagmentation, the TDE1 (Nextera Tn5 Transposase) Tagment DNA Enzyme (Illumina kit

#20034198) was immediately applied to the samples by resuspending the nuclei in the

transposition reaction mix and the suspension was incubated for 30 minutes at 37°C while

gently shaking (Buenrostro et al. 2015). The samples were then purified (NEB, Monarch

Nucleic Acid Purification Kit #20034198), and 12 μl of the transposed DNA were used for

amplification by PCR (8 μl H2O, 2.5 μl 25 μM Customr Nextera PCR primer 1 (Buenrostro et

al. 2013), 2.5 μl 25 μM Custom Nextera PCR Primer 2 and 25 μl NEB Next High Fidelity 2x

PCR Master Mix (NEB, #M0541) with 1 cycle of (72°C for 5 min, 98°C for 30 sec) and 16 cycles

of (98°C for 10 sec, 63°C for 30 sec, 72°C for 1 min). The library was cleaned up using the

Coulter Agencourt AMPure XP kit (Beckman-Coulter, #10136224) applying a 1:1 vol:vol (PCR

product:beads) and eluted in 15 μl 15 mM Tris. The library was sequenced using NextSeq 550

(Illumina) using High-Output with 40 cycles Paired End mode.

Analysis of sequencing data

Reads were mapped to Arabidopsis TAIR 10 genome by bowtie2 (v2.2.6) (Langmead

and Salzberg 2012). Output sam files were converted to bam files and then trimed with "-q 2"

and "1 2 3 4 5" parameters in samtools (v0.1.19) (Li et al. 2009) to remove mitochondrial and

chloroplast sequence. Output bam files were treated with picard MarkDuplicates (v2.25.5) to

remove duplicated reads. Then, output files were passed through samtools view again, to

obtain 40 M reads (targeted) for further analysis. For the profiles, bigWig files were generated

by bamCoverage in deepTools (v3.5.1) (Ramírez et al. 2014) from bam files with 10 bp bin

size, and visualized by igv (Thorvaldsdóttir et al. 2013). For peak calling, bam files were

converted to TagDirectory by homer (v4.10.3) (Heinz et al. 2010), and peaks were called by

findPeaks in homer with "-region -minDist 150" option. Differential peaks were obtained by

getDifferentialPeaks function with "-F 2.0 -P 0.05" option. TSS profiles were made by ngs.plot

(v2.47) (Shen et al. 2014). Raw data were deposited to the NCBI GEO archive (Barrett et al.

2013)   under   the   accession   GSE184344

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE184344).

Quantification and statistical analyses

Statistical analyses were performed using IBM SPSS Statistics for Windows, Version

22.0. Armonk, NY: IBM Corp or using GraphPad Prism version 6.01 (GraphPad Software, La

Jolla, USA). Means were calculated from measurements with sample sizes as indicated in the

respective figure legends. In general, all displayed data represents at least three independent,

technical repetitions, unlike otherwise indicated. Error bars represent ± standard deviation. All

analyzed datasets were prior tested for homogeneity of variances by the Levene statistic. One-

way ANOVA was performed, using a confidence interval (CI) of 95% and a post-hoc Tukey

HSD for comparisons of five or more data sets of homogenous variances or a post-hoc

Tamhane-T2 in case variances were not homogenous. Graphs were generated in GraphPad

Prism version 6.01 or 9.2 (GraphPad Software, La Jolla, USA) or in Excel (Microsoft,

Redmond, USA).

Acknowledgements

This work was supported by the SFB 1101 to K.H., J.U.L, T.G. and SFB 873 (Deutsche

Forschungsgemeinschaft, DFG) to J.U.L. and T.G., a Heisenberg Professorship (DFG, GR

2104/5-2), and an ERC Consolidator grant (PLANTSTEMS, #647148) to T.G. , a postdoctoral

fellowship of the Japan Society for the Promotion of Science [JSPS Overseas Research

Fellowships 201960008 to D.S.], and a PhD student fellowship of the Cusanuswerk to N.T..

We are grateful to Christian Hardtke (University of Lausanne, Switzerland), Dolf Weijers,

Shunsuke Saiga (both at Wageningen University, The Netherlands), Andy Maule (John Innes

Centre, UK), Sebastain Wolf and Karin Schumacher (both Heidelberg University, Germany)

for providing seed material and constructs. We are also grateful to Monika Langlotz (Cell

Networks Flow Cytometry & FACS Core Facility, Heidelberg University, Germany) and David

Ibberson (Cell Networks Deep Sequencing Core Facility, Heidelberg University, Germany) for

providing excellent technical support.

Author Contributions

Conceived and designed the experiments: EW, NT, FW, VL, DS, LL, KH, TG. Performed

experiments: EW, NT, FW, LL, IJ, VL, PH, YX, MG, CW. Analysed the data: EW, NT, FW, KH,

JUL, DS, TG. Wrote the paper: EW, TG.

Conflict of Interest

The authors declare no conflict of interest.

References

Anne P, Azzopardi M, Gissot L, Beaubiat S, Hematy K, Palauqui JC. 2015. OCTOPUS Negatively Regulates BIN2 to Control Phloem Differentiation in Arabidopsis thaliana. *Current Biology* **25**: 2584-2590.

Anne P, Hardtke CS. 2017. Phloem function and development-biophysics meets genetics. *Curr Opin Plant Biol* **43**: 22-28.

Arongaus AB, Chen S, Pireyre M, Glockner N, Galvao VC, Albert A, Winkler JB, Fankhauser C, Harter K, Ulm R. 2018. Arabidopsis RUP2 represses UVR8-mediated flowering in noninductive photoperiods. *Genes Dev* **32**: 1332-1343.

Ashby AM, Watson MD, Loake GJ, Shaw CH. 1988. Ti plasmid-specified chemotaxis of Agrobacterium tumefaciens C58C1 toward vir-inducing phenolic compounds and soluble factors from monocotyledonous and dicotyledonous plants. *Journal of bacteriology* **170**: 4181-4187.

Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M et al. 2013. NCBI GEO: archive for functional genomics data sets--update. *Nucleic acids research* **41**: D991-995.

Benfey PN, Chua NH. 1990. The Cauliflower Mosaic Virus 35S Promoter: Combinatorial Regulation of Transcription in Plants. *Science* **250**: 959-966.

Blob B, Heo JO, Helariutta Y. 2018. Phloem differentiation: an integrative model for cell specification. *J Plant Res* **131**: 31-36.

Bonke M, Thitamadee S, Mähönen AP, Hauser MT, Helariutta Y. 2003. APL regulates vascular tissue identity in Arabidopsis. *Nature* **426**: 181-186.

Brady SM, Orlando DA, Lee JY, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN. 2007. A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* **318**: 801-806.

Breda AS, Hazak O, Schultz P, Anne P, Graeff M, Simon R, Hardtke CS. 2019. A Cellular Insulator against CLE45 Peptide Signaling. *Current biology : CB* **29**: 2501-2508.

Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature methods* **10**: 1213-1218.

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. 2015. ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**: 21.29.21-21.29.29.

Cho H, Cho HS, Nam H, Jo H, Yoon J, Park C, Dang TVT, Kim E, Jeong J, Park S et al. 2018. Translational control of phloem development by RNA G-quadruplex-JULGI determines plant sink strength. *Nature plants* **4**: 376-390.

Cifuentes-Esquivel N, Bou-Torrent J, Galstyan A, Gallemi M, Sessa G, Salla Martret M, Roig-Villanova I, Ruberti I, Martinez-Garcia JF. 2013. The bHLH proteins BEE and BIM positively modulate the shade avoidance syndrome in Arabidopsis seedlings. *The Plant journal : for cell and molecular biology* **75**: 989-1002.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *The Plant journal : for cell and molecular biology* **16**: 735-743.

De Lucia F, Crevillen P, Jones AM, Greb T, Dean C. 2008. A PHD-polycomb repressive complex 2 triggers the epigenetic silencing of FLC during vernalization. *Proceedings of the National Academy of Sciences of the United States of America* **105**: 16831-16836.

De Schepper V, De Swaef T, Bauweraerts I, Steppe K. 2013. Phloem transport: a review of mechanisms and controls. *Journal of experimental botany* **64**: 4839-4850.

Depuydt S, Rodriguez-Villalon A, Santuari L, Wyser-Rmili C, Ragni L, Hardtke CS. 2013. Suppression of Arabidopsis protophloem differentiation and root meristem growth by CLE45 requires the receptor-like kinase BAM3. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 7074-7079.

Fraley RT, Rogers SG, Horsch RB, Sanders PR, Flick JS, Adams SP, Bittner ML, Brand LA,
Fink CL, Fry JS et al. 1983. Expression of bacterial genes in plant cells. *Proceedings*
of the National Academy of Sciences of the United States of America **80**: 4803-4807.
Furuta KM, Hellmann E, Helariutta Y. 2014. Molecular control of cell specification and cell
differentiation during procambial development. *Annual review of plant biology* **65**:
607-638.
Gujas B, Kastanaki E, Sturchler A, Cruz TMD, Ruiz-Sola MA, Dreos R, Eicke S, Truernit E,
Rodriguez-Villalon A. 2020. A Reservoir of Pluripotent Phloem Cells Safeguards the
Linear Developmental Trajectory of Protophloem Sieve Elements. *Current biology :*
CB **30**: 755-766.e754.
Haseloff J, Siemering KR, Prasher DC, Hodge S. 1997. Removal of a cryptic intron and
subcellular localization of green fluorescent protein are required to mark transgenic
Arabidopsis plants brightly. *Proceedings of the National Academy of Sciences of the*
United States of America **94**: 2122-2127.
Hazak O, Brandt B, Cattaneo P, Santiago J, Rodriguez-Villalon A, Hothorn M, Hardtke CS.
2017. Perception of root-active CLE peptides requires CORYNE function in the
phloem vasculature. *EMBO reports* **18**: 1367-1381.
Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,
Glass CK. 2010. Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities.
Molecular cell **38**: 576-589.
Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM. 2000. pGreen: a versatile
and flexible binary Ti vector for Agrobacterium-mediated plant transformation. *Plant*
molecular biology **42**: 819-832.
Jiang L, Liu X, Xiong G, Liu H, Chen F, Wang L, Meng X, Liu G, Yu H, Yuan Y et al. 2013.
DWARF 53 acts as a repressor of strigolactone signalling in rice. *Nature* **504**: 401-
405.
Kondo Y, Nurani AM, Saito C, Ichihashi Y, Saito M, Yamazaki K, Mitsuda N, Ohme-Takagi
M, Fukuda H. 2016. Vascular Cell Induction Culture System Using Arabidopsis
Leaves (VISUAL) Reveals the Sequential Differentiation of Sieve Element-like Cells.
Plant Cell **28**: 1250-1262.
Krogan NT, Long JA. 2009. Why so repressed? Turning off transcription during plant growth
and development. *Curr Opin Plant Biol* **12**: 628-636.
Kurihara D, Mizuta Y, Sato Y, Higashiyama T. 2015. ClearSee: a rapid optical clearing
reagent for whole-plant fluorescence imaging. *Development (Cambridge, England)*
142: 4168-4179.
Ladwig F, Dahlke RI, Stuhrwohldt N, Hartmann J, Harter K, Sauter M. 2015. Phytosulfokine
Regulates Growth in Arabidopsis through a Response Module at the Plasma
Membrane That Includes CYCLIC NUCLEOTIDE-GATED CHANNEL17, H+-ATPase,
and BAK1. *Plant Cell* **27**: 1718-1729.
Lampropoulos A, Sutikovic Z, Wenzl C, Maegele I, Lohmann JU, Forner J. 2013. GreenGate-
--a novel, versatile, and efficient cloning system for plant transgenesis. *PloS one* **8**:
e83043.
Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature*
methods **9**: 357-359.
Legrain P, Wojcik J, Gauthier JM. 2001. Protein--protein interaction maps: a lead towards
cellular functions. *Trends Genet* **17**: 346-352.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R.
2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford,*
England) **25**: 2078-2079.
Liang Y, Ward S, Li P, Bennett T, Leyser O. 2016. SMAX1-LIKE7 Signals from the Nucleus
to Regulate Shoot Development in Arabidopsis via Partially EAR Motif-Independent
Mechanisms. *Plant Cell* **28**: 1581-1601.
Lin TF, Saiga S, Abe M, Laux T. 2016. OBE3 and WUS Interaction in Shoot Meristem Stem
Cell Regulation. *PloS one* **11**: e0155657.

López-Salmerón V, Cho H, Tonn N, Greb T. 2019. The Phloem as a Mediator of Plant
Growth Plasticity. *Current biology : CB* **29**: R173-r181.
Lucas WJ, Groover A, Lichtenberger R, Furuta K, Yadav SR, Helariutta Y, He XQ, Fukuda H,
Kang J, Brady SM et al. 2013. The plant vascular system: evolution, development
and functions. *Journal of integrative plant biology* **55**: 294-388.
Ma H, Duan J, Ke J, He Y, Gu X, Xu TH, Yu H, Wang Y, Brunzelle JS, Jiang Y et al. 2017. A
D53 repression motif induces oligomerization of TOPLESS corepressors and
promotes assembly of a corepressor-nucleosome complex. *Science advances* **3**:
e1601217.
Marhava P, Aliaga Fandino AC, Koh SWH, Jelínková A, Kolb M, Janacek DP, Breda AS,
Cattaneo P, Hammes UZ, Petrášek J et al. 2020. Plasma Membrane Domain
Patterning and Self-Reinforcing Polarity in Arabidopsis. *Developmental cell* **52**: 223-
235.e225.
Marhava P, Bassukas AEL, Zourelidou M, Kolb M, Moret B, Fastner A, Schulze WX,
Cattaneo P, Hammes UZ, Schwechheimer C et al. 2018. A molecular rheostat adjusts
auxin flux to promote root protophloem differentiation. *Nature* **558**: 297-300.
Oparka KJ, Turgeon R. 1999. Sieve elements and companion cells-traffic control centers of
the phloem. *Plant Cell* **11**: 739-750.
Pauwels L, Barbero GF, Geerinck J, Tilleman S, Grunewald W, Perez AC, Chico JM,
Bossche RV, Sewell J, Gil E et al. 2010. NINJA connects the co-repressor TOPLESS
to jasmonate signalling. *Nature* **464**: 788-791.
Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. 2014. deepTools: a flexible platform for
exploring deep-sequencing data. *Nucleic acids research* **42**: W187-191.
Rodriguez-Villalon A. 2016. Wiring a plant: genetic networks for phloem formation in
Arabidopsis thaliana roots. *New Phytol* **210**: 45-50.
Rodriguez-Villalon A, Gujas B, Kang YH, Breda AS, Cattaneo P, Depuydt S, Hardtke CS.
2014. Molecular genetic framework for protophloem formation. *Proceedings of the*
National Academy of Sciences of the United States of America **111**: 11551-11556.
Rodriguez-Villalon A, Gujas B, van Wijk R, Munnik T, Hardtke CS. 2015. Primary root
protophloem differentiation requires balanced phosphatidylinositol-4,5-biphosphate
levels and systemically affects root branching. *Development (Cambridge, England)*
142: 1437-1446.
Ross-Elliott TJ, Jensen KH, Haaning KS, Wager BM, Knoblauch J, Howell AH, Mullendore
DL, Monteith AG, Paultre D, Yan D et al. 2017. Phloem unloading in Arabidopsis
roots is convective and regulated by the phloem-pole pericycle. *Elife* **6**: e24125.
Saiga S, Furumizu C, Yokoyama R, Kurata T, Sato S, Kato T, Tabata S, Suzuki M, Komeda
Y. 2008. The Arabidopsis OBERON1 and OBERON2 genes encode plant
homeodomain finger proteins and are required for apical meristem maintenance.
Development (Cambridge, England) **135**: 1751-1759.
Saiga S, Moller B, Watanabe-Taneda A, Abe M, Weijers D, Komeda Y. 2012. Control of
embryonic meristem initiation in Arabidopsis by PHD-finger protein complexes.
Development (Cambridge, England) **139**: 1391-1398.
Sanchez R, Zhou MM. 2011. The PHD finger: a versatile epigenome reader. *Trends*
Biochem Sci **36**: 364-372.
Scacchi E, Osmont KS, Beuchat J, Salinas P, Navarrete-Gomez M, Trigueros M, Ferrandiz
C, Hardtke CS. 2009. Dynamic, auxin-responsive plasma membrane-to-nucleus
movement of Arabidopsis BRX. *Development (Cambridge, England)* **136**: 2059-2067.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B et al. 2012. Fiji: an open-source platform for
biological-image analysis. *Nature methods* **9**: 676-682.
Scholthof HB. 2006. The Tombusvirus-encoded P19: from irrelevance to elegance. *Nat Rev*
Microbiol **4**: 405-411.
Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D. 2006. Highly Specific Gene
Silencing by Artificial MicroRNAs in Arabidopsis. *Plant Cell* **18**: 1121-1133.

Shen L, Shao N, Liu X, Nestler E. 2014. ngs.plot: Quick mining and visualization of next-
generation sequencing data by integrating genomic databases. *BMC genomics* **15**:
284.
Shi D, Jouannet V, Agustí J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T. 2021.
Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem reveals
local cellular signatures. *Plant Cell* **33**: 200-223.
Shi D, Lebovka I, Lopez-Salmeron V, Sanchez P, Greb T. 2019. Bifacial cambium stem cells
generate xylem and phloem during radial plant growth. *Development (Cambridge,*
England) **146**.
Soundappan I, Bennett T, Morffy N, Liang Y, Stanga JP, Abbas A, Leyser O, Nelson DC.
2015. SMAX1-LIKE/D53 family members enable distinct MAX2-dependent responses
to strigolactones and karrikins in Arabidopsis. *Plant Cell* **27**: 3143-3159.
Thibivilliers S, Anderson D, Libault M. 2020. Isolation of Plant Root Nuclei for Single Cell
RNA Sequencing. *Current protocols in plant biology* **5**: e20120.
Thomas CL, Schmidt D, Bayer EM, Dreos R, Maule AJ. 2009. Arabidopsis plant
homeodomain finger proteins operate downstream of auxin accumulation in
specifying the vasculature and primary root meristem. *The Plant journal : for cell and*
molecular biology **59**: 426-436.
Thorvaldsdóttir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Briefings in bioinformatics*
14: 178-192.
Truernit E, Bauby H, Belcram K, Barthelemy J, Palauqui JC. 2012. OCTOPUS, a polarly
localised membrane-associated protein, regulates phloem differentiation entry in
Arabidopsis thaliana. *Development (Cambridge, England)* **139**: 1306-1315.
Truernit E, Bauby H, Dubreucq B, Grandjean O, Runions J, Barthelemy J, Palauqui JC.
2008. High-resolution whole-mount imaging of three-dimensional tissue organization
and gene expression enables the study of Phloem development and structure in
Arabidopsis. *Plant Cell* **20**: 1494-1503.
van den Berg C, Willemsen V, Hendriks G, Weisbeek P, Scheres B. 1997. Short-range
control of cell differentiation in the Arabidopsis root meristem. *Nature* **390**: 287-289.
Voinnet O, Rivas S, Mestre P, Baulcombe D. 2003. An enhanced transient expression
system in plants based on suppression of gene silencing by the p19 protein of tomato
bushy stunt virus. *The Plant journal : for cell and molecular biology* **33**: 949-956.
Walker C, Bennett T. 2017. Reassessing the evolution of strigolactone synthesis and
signalling. *bioRxiv*: 228320.
Wallner ES. 2018. Early events in phloem formation: Exploring the molecular network of
SMXL3/4/5. in *Faculty for Biosiences*, p. 128. University of Heidelberg, Heidelberg.
Wallner ES, Lopez-Salmeron V, Belevich I, Poschet G, Jung I, Grunwald K, Sevilem I,
Jokitalo E, Hell R, Helariutta Y et al. 2017. Strigolactone- and Karrikin-Independent
SMXL Proteins Are Central Regulators of Phloem Formation. *Current biology : CB* **27**:
1241-1247.
Wallner ES, Lopez-Salmeron V, Greb T. 2016. Strigolactone versus gibberellin signaling:
reemerging concepts? *Planta* **243**: 1339-1350.
Wallner ES, Tonn N, Shi D, Jouannet V, Greb T. 2020. SUPPRESSOR OF MAX2 1-LIKE 5
promotes secondary phloem formation during radial stem growth. *The Plant journal :*
for cell and molecular biology **in press**.
Wang L, Wang B, Jiang L, Liu X, Li X, Lu Z, Meng X, Wang Y, Smith SM, Li J. 2015.
Strigolactone signaling in Arabidopsis regulates shoot development by targeting D53-
Like SMXL repressor proteins for ubiquitination and degradation. *Plant Cell* **27**: 3128-
3142.
Wang L, Wang B, Yu H, Guo H, Lin T, Kou L, Wang A, Shao N, Ma H, Xiong G et al. 2020.
Transcriptional regulation of strigolactone signalling in Arabidopsis. *Nature* **583**: 277-
281.
Wu YY, Hou BH, Lee WC, Lu SH, Yang CJ, Vaucheret H, Chen HM. 2017. DCL2- and
RDR6-dependent transitive silencing of SMXL4 and SMXL5 in Arabidopsis dcl4

mutants causes defective phloem transport and carbohydrate over-accumulation. *The*
Plant journal : for cell and molecular biology **90**: 1064-1078.
Zhou F, Lin Q, Zhu L, Ren Y, Zhou K, Shabek N, Wu F, Mao H, Dong W, Gan L et al. 2013.
D14-SCF(D3)-dependent degradation of D53 regulates strigolactone signalling.
Nature **504**: 406-410.

Main figure titles and legends

Figure 1: Phloem-related genes are less active in *smxl4;smxl5* mutants

(A – H) Comparison of *OPS:OPS-GFP* (A, B), *BRX:BRX-CITRINE* (C, D), *BAM3:BAM3-*

CITRINE, and *CVP2:NLS-VENUS* (G, H) reporter activities in wild type (A, C, E, G) and

smxl4;smxl5 double mutants (B, D, F, H). Asterisks depict the first differentiating SE. Arrows

point to earliest detectable reporter activities. Scale bars represent 50 μm. Note that the

fluorescent signal in the root cortex visible in F is autofluorescence due to tissue disruption.

(I – P) *OPS:OPS-GFP* (I-L) and *BRX:BRX-CITRINE* (M-P) close to the QC in wild type (I, J,

M, N) and *smxl4;smxl5* mutants (K, L, O, P). In J, L, N, and P, fluorescent signals are depicted

without counterstaining. Asterisks indicate QC cells. Arrows point to earliest detectable

reporter activities. Scale bar in (I) represents 50 μm. Same magnification in I-P.

Figure 2 : Analysis of interaction between *SMXL4* and *SMXL5* with other phloem

regulators

(**A**) Root length of five day-old plants. n = 37–75. Statistical groups determined by one-way

ANOVA and post-hoc Tukey's test (95 % CI). Shown is one representative experiment of three

repetitions.

(B – E) Comparison of *SMXL4:SMXL4-YFP* and *SMXL5:SMXL5-YFP* reporter activities in

smxl4;smxl5 and *smxl;ops* mutants. Arrows indicate the signals closest to the QC. Scale bars

represent 50 μm.

Figure 3: Genetic interaction of *SMXL5* with *OPS* and *BRX*

(**A**) Root length of five day-old wild type (WT), *smxl5*, *brx*, *ops*, *smxl4;smxl5*, *brx;smxl5*, and

ops;smxl5 plants. n = 38–76 for each group. Statistical groups determined by one-way ANOVA

and post-hoc Tukey's HSD test (95 % CI). Shown is one representative experiment of three

repetitions.

(B) Phenotypic characterization of phloem development of two day-old wild type, *smxl5, brx,*

ops, smxl4;smxl5, brx;smxl5, and *ops;smxl5* plants. n = 97–129 for each group.

Figure 4: SMXL5 and OBE3 proteins interact

(A) The SMXL5 protein was expressed in yeast fused to the GAL4 DNA binding domain (BD)

and OBE3 fused to the GAL4 activation domain (AD). All strains contained AD- and BD-

expressing plasmids either alone or fused with SMXL5, OBE3 or control proteins, respectively.

Growth on SD-leu-trp indicate the presence of both plasmids, growth on SD-leu-trp-his-ade

medium indicate the presence of plasmids and protein interaction.

(B) Interaction of SMXL5-3xHA and 6xMyc-OBE3 proteins by co-immunoprecipitation (co-IP)

and subsequent Western blot analysis after transient overexpression in *N. benthamiana*. 'Input'

represents unprocessed protein extracts, 'unbound' show proteins that remained in the extract

after IP and 'IP: α HA' depicts samples after immunoprecipitation by α-HA-beads. Western

blots were probed by α HA or α Myc antibodies, respectively. Signals revealed an expected

SMXL5-3xHA protein size of approximately 120 kDa. The size of the detected 6xMyc-OBE3

protein (~100 kDa) exceeded the expected size of 92 kDa.

(C – K) Fluorescent signals and bright field images of epidermal *N. benthamiana* nuclei

transiently co-expressing OBE3-mGFP/mCherry (C-E), OBE3-mGFP/SMXL5-mCherry (F-H)

and mGFP-NLS/mCherry-NLS (I-K). The dashed yellow line indicates the outlines of nuclei in

merged images (E, H, K). Scale bars represent 5 μm.

(L) FRET-FLIM analysis of transiently transformed *N. benthamiana* epidermal leaf cells

expressing the OBE3-mGFP donor without an mCherry acceptor or in the presence of NLS-

mCherry or SMXL5-mCherry. Error bars indicate standard deviation. Data were derived from

three biological replicates with n = 27–80. To test for homogeneity of variance, a Brown-

Forsythe test was applied. As the variances are not homogenous a Wilcoxon / Kruskal-Wallis

test was carried out followed by a Dunn post hoc analysis.

(M) Heat maps of representative nuclei used for FLIM measurements. The donor lifetimes of

OBE3-mGFP are color-coded according to the scale at the bottom. Size bars represent 4 µm.

Figure 5: *OBE3* genetically interacts with *SMXL3/4/5* in root growth regulation

(A) 10 day-old wild type, *smxl4, smxl5, smxl4;smxl5, obe3, obe4, smxl4;obe3, smxl4;obe4,*

smxl5;obe3, smxl5;obe4 seedlings are shown from left to right. Scale bar represents 1 cm.

(B) Quantification of root length depicted in A. Mean values of four independent experiments

(n = 15-55 per experiment and genotype) were analyzed by a one-way ANOVA with post-hoc

Tukey HSD (95 % CI). Statistical groups are marked by letters.

(C) Quantification of root length depicted in D. Mean values of three independent experiments

(n = 62–74 per experiment and genotype) were analyzed by a one-way ANOVA with post-hoc

Tamhane-T2 (95 % CI). Statistical groups are marked by letters.

(D) 10 day-old wild type, *smxl3, smxl5, smxl3;smxl5, obe3, smxl3;obe3, smxl5;obe3* seedlings

are shown from left to right. Scale bar represents 1 cm.

Figure 6: *OBE3* interacts with *SMXL5* in protophloem formation

(A − C) Phloem-specific activity of *SMXL4:SMXL4-YFP* (A) coincides with activity of

OBE3:OBE3-GFP (B) and *OBE4:OBE4-GFP* (C) reporters in the developing phloem.

Fluorescent signals (green) and cell wall staining by Direct Red 23 (magenta) were detected

by confocal microscopy. Pink arrows point to the first differentiated SE indicated by enhanced

cell wall staining. Scale bars represent 50 µm.

(D) Schematic representation of one developing phloem pole at the root tip. Two periclinal cell

divisions generate SE precursor and procambium (orange arrow) and proto- and metaphloem

cell lineages (blue arrow), respectively. Differentiated SEs are marked by a pink arrow. The

QC is marked by a yellow asterisk.

(E − J) Phloem formation in 2 day-old wild type (E), *obe3* (F), *smxl4;smxl5* (G), *smxl4;obe3*

(H), *smxl5;obe3* (I) and *smxl5;obe4* (J) root tips. Cell walls were stained by mPS-PI (white).

Yellow asterisks mark the QC. Enhanced mPS-PI staining indicates differentiation of SEs (pink

arrows) in wild type (E), *obe3* (F) and *obe4;smxl5* (G). Orange and blue arrows mark the first

and second periclinal division, respectively, in the developing phloem cell lineage. Scale bars

represent 20 µm.

(K) The distance from the QC to the first and second periclinal division shown in D-I was

quantified (n = 9–11). Statistical groups are indicated by letters and were determined by a one-

way ANOVA with post-hoc Tukey HSD (95 % CI). Distances of 1^{st} cell divisions and 2^{nd} cell

divisions were compared independently.

Figure 7: Phloem-specific *OBE3* expression is sufficient for promoting root growth

(A – B) Seven day-old *smxl5;obe3* root tips carrying an *SMXL5:OBE3-turquoise* reporter

(green signal, A) were compared to a non-transformed wild type (WT) root tip (B). Cell walls

were stained by DirectRed (magenta). White arrows mark differentiated SEs. Scale bars

indicate 20 µm.

(C) Root length quantification of plants depicted in D. Results from one representative

experiment out of three independent experiments (n = 61–71 per experiment and genotype)

are shown. Mean values were analyzed by one-way ANOVA with post-hoc Tamhane-T2 (95 %

CI). Statistical groups are indicated by letters.

(D) 10 day-old wild type, *smxl5, obe3, smxl5;obe3, SMXL5:OBE3-turquoise/smxl5;obe3*,

SMXL5:obe3-miRNA3/smxl5 and *SMXL5:obe3-miRNA4/smxl5* seedlings are shown from left

to right. Scale bar represents 1 cm.

1008

1009

Figure 8: *SMXL* and *OBE3* genes determine phloem-related chromatin profile

(A) Summary of ATAC-seq results for the different samples. Specific OCRs were identified by

comparing GFP-positive and GFP-negative samples of the respective genetic backgrounds.

After initial alignment, OCR detection and other subsequent analyses were conducted using

40 M reads aligning to the nuclear genome excluding duplicates for each sample.

(B) Chromatin conformation profiles for all genes found in the Arabidopsis genome or only for

phloem and non-phloem genes according to Brady et al., 2007. Read alignment was adjusted

to transcriptional start sites (TSS) and transcriptional end sites (TES). X-axes show distance

to TSSs and TES in base-pairs. Inserts in all graphs are enlargements of the peaks identified

close to TSSs.

(C, D) Comparison of OCR profiles found in GFP-positive (C) and GFP-negative (D) samples.

Numbers indicate the number of OCRs with significantly more ('up') or less ('down') aligned

reads.

Supplementary item titles and legends

1024

Supplementary Figure 1: Expression of SMXL5-VENUS protein fusions in *smxl4;smxl5*

mutants in comparison to promoter reporters.

(**A – E**) Expression of SMXL5-VENUS protein fusions under the control of different

heterologous promoters. Differentiated SEs are indicated by asterisks.

(**F**) Detection of the BRX-VENUS protein fusion expressed under the control of the *SMXL4*

promoter. Differentiated SEs are indicated by the asterisk.

(**G – I**) Detection of *OPS:ER-VENUS*, *BAM3:ER-VENUS*, and *CVP2:ER-VENUS* reporter

activities in *smxl4;smxl5* mutants. Arrow in G points to a weakly detectable reporter-derived

signal.

(**J – M**) Detection of *APL:ER-VENUS* reporter activity in wild type (J, K) and *smxl4;smxl5*

mutants (L, M). K and M show the same root tip as in J and L, respectively, without the counter

stain signal. Scale bars in all pictures represent 50 μm. Arrows in J and K point to a weakly

detectable reporter-derived signal.

(**N**) Root length measurements in different genetic backgrounds in the presence and absence

of exogenous CLE45 five days after germination. Results from one representative experiment

out of two independent experiments (n = 23–58 per experiment and genotype, except n = 6 for

bam3-2) are shown. Mean values were analyzed by one-way ANOVA with post-hoc Tukey

HSD (95 % CI). Statistical groups are indicated by letters.

1043

Supplementary Figure 2: Characterization of phloem development in *smxl* and *ops*

mutants.

(**A – K**) Phloem development is monitored in wild type (A), *smxl5* (B), *smxl4;smxl5* (C), *ops* (D-

G) and *ops;smxl5* (H-K) mutant backgrounds by confocal analysis of mPS-PI-stained root tips.

Asterisks indicate the most apical appearance of differentiated SEs. In case 'gaps' are

observed, more than one asterisk is depicted.

1050

Supplementary Figure 3: Characterization of phloem development in *smxl* and *brx*

mutants.

(**A – H**) Phloem development is monitored in wild type (A), *smxl5* (B), *smxl4;smxl5* (C), *brx* (D-

E) and *brx;smxl5* (F-H) mutant backgrounds by confocal analysis of mPS-PI-stained root tips.

Asterisks indicate the most apical appearance of differentiated SEs. In case 'gaps' are

observed, more than one asterisk is depicted.

1057

Supplementary Figure 4: *OBE3* cDNA clones isolated when screening for SMXL5

interactors using the Yeast-Two-Hybrid system.

(**A**) Alignment of isolated clones (in green) to the full *OBE3* cDNA sequence (top) using CLC

Main Workbench Version 7.6.1 (CLC Bio Qiagen, Aarhus, Denmark). Protein domains

predicted for OBE3 are indicated above. The sequence area present in all isolated clones

(selected interacting domain, SID) is marked by a purple arrow and flanked by red lines. The

OBE3 protein contains two predicted nuclear localization signals (NLS) identified by the cNLS

Mapper (Kosugi et al. 2009, NLS_bipartite and NLS_monopartite, light green arrows), a PHD-

finger domain (blue arrow) with a histone 3 (H3) binding motif (light blue arrow) and a coiled

coil domain (orange arrow) predicted by InterPro (EMBL-EBI, Cambridgeshire, UK). The OBE3

open reading frame is flanked by a 3' and a 5' untranslated region (3'UTR and 5'UTR, green

arrows), respectively.

(**B**) Amino acid sequence encoded by the SID sequence, present in all isolated clones.

1071

Supplementary Figure 5: *SMXL4* or *SMXL5* genes do not genetically interact with *OBE1*

or *OBE2* genes

(**A**) 10 day-old wild type, *smxl4, smxl5, smxl4;smxl5, obe1, obe2, smxl4;obe1, smxl4;obe2,*

smxl5;obe1, smxl5;obe2 seedlings are shown from left to right. Scale bar represents 1 cm.

(**B**) Quantification of root lengths depicted in A. Mean values of three independent experiments

(n = 34-75 per experiment and genotype) were analyzed by a one-way ANOVA with post-hoc

Tukey HSD (95 % CI). Statistical groups are indicated by letters.

1079

Supplementary Figure 6: SEs form in *smxl4* and *smxl5* single mutants

(**A – D**) 2 day-old mPS-PI-stained root tips of wild type (A), *smxl4* (B), *smxl5* (C) and

smxl4;smxl5 (D) plants. Differentiating SEs were observed in wild type, *smxl4* and *smxl5* (pink

arrows). Periclinal cell divisions are marked by orange and blue arrows. The QC is indicated

by a yellow asterisk. Scale bars represent 20 µm.

(**E**) The distance of the first and second periclinal division from the QC for plants shown in A-

D was quantified (n = 10-12). Statistical groups marked by letters were determined by one-way

ANOVA with post-hoc Tamhane-T2 (95 % CI). Distances of 1^{st} cell divisions and 2^{nd} cell

divisions were compared independently.

1089

Supplementary Figure 7: Activity of the *SMXL5:H4-GFP* transgene in wild type, *smxl5*,

***smxl4;smxl5* and *smxl5;obe3* backgrounds and sorting of GFP-positive and GFP-**

negative nucleus fractions

(**A**) Activity of the *SMXL5:H4-GFP* transgene in root tips of wild type, *smxl5, smxl4;smxl5* and

smxl5;obe3 seedlings two days after germination. Scale bars represent 20 µm.

(**B**) Sorting plots displaying gates for identifying GFP-positive (P4) and GFP-negative (P3)

nuclei comparing wild type without transgene (wt ctrl) and wild type, *smxl5, smxl4;smxl5* and

smxl5;obe3 plants carrying the *SMXL5:H4-GFP* transgene. 10,000 sorting events were

processed for display in all cases. P3 fractions included 2.2 to 2.9 % of all events and P4

fractions 0.1 to 0.2 % (0 % for wild type without the transgene). X-axes indicate RFP

fluorescence (561 nm), Y-axes indicate GFP fluorescence (488 nm).

1101

Supplementary Figure 8: ATAC-seq analysis comparing GFP-positive and GFP-negative

samples from wild type, *smxl5*, *smxl4;smxl5* and *smxl5;obe3* seedlings

Profile of read alignment (dark and light green) and OCR detection (red) across all

chromosomes for all samples processed. Dark green: GFP-positive samples. Light green:

GFP-negative samples. Centromeric regions are indicated by reduced read alignment and

OCR detection.

1108

1109   **Supplementary Figure 9: Chromatin conformation profile for *OPS* and *BAM3* gene**

1110   **regions.**

1111   Profile of read alignment (dark and light green) and OCR detection (red) for all samples are

1112   shown. Dark green: GFP-positive samples. Light green: GFP-negative samples. Gene

1113   structures are indicated at the bottom (dark blue).

1114

1115   **Supplementary Figure 10: Chromatin conformation profile for *CVP2* and *APL* gene**

1116   **regions.**

1117   Profile of read alignment (dark and light green) and OCR detection (red) for all samples are

1118   shown. Dark green: GFP-positive samples. Light green: GFP-negative samples. Gene

1119   structures are indicated at the bottom (dark blue).

1120

1121   **Supplementary Dataset 1: OCRs and associated genes specific for GFP-positive or**

1122   **GFP-negative fractions from wild type, *smxl5*, *smxl4;smxl5* or *smxl5;obe3* backgrounds,**

1123   **respectively.**

1124

1125 **Supplementary Datset 2: Genes with OCRs specific for GFP-positive or GFP-negative**

1126 **fractions for wild type, *smxl5*, *smxl4;smxl5* and *smxl5;obe3* backgrounds.**

1127

1128 **Supplementary Dataset 3: OCRs and associated genes specific for wild type, *smxl5*,**

1129 ***smxl4;smxl5* and *smxl5;obe3* plants comparing GFP-positive fractions from the**

1130 **respective backgrounds.**

1131

1132 **Supplementary Dataset 4: OCRs and associated genes specific for wild type, *smxl5*,**

1133 ***smxl4;smxl5* and *smxl5;obe3* plants comparing GFP-negative fractions from the**

1134 **respective backgrounds.**

1135

1136 **Supplementary Dataset 5: Phloem and non-phloem genes according to Brady et al.,**

1137 **2007.**

1138

1139 **Supplementary Table 1: Oligonucleotides used in this study.**

Figure 1: Phloem-related genes are less active in *smxl4;smxl5* mutants

(A – H) Comparison of *OPS:OPS-GFP* (A, B), *BRX:BRX-CITRINE* (C, D), *BAM3:BAM3-CITRINE*, and *CVP2:NLS-VENUS* (G, H) reporter activities in wild type (A, C, E, G) and *smxl4;smxl5* double mutants (B, D, F, H). Asterisks depict the first differentiating SE. Arrows point to earliest detectable reporter activities. Scale bars represent 50 μm.

(I – P) *OPS:OPS-GFP* (I-L) and *BRX-:BRX-CITRINE* (M-P) close to the QC in wild type (I, J, M, N) and *smxl4;smxl5* mutants (K, L, O, P). In J, L, N, and P, fluorescent signals are depicted without counterstaining. Asterisks indicate QC cells. Arrows point to earliest detectable reporter activities. Scale bar in (I) represents 50 μm. Same magnification in I-P.

A

B *SMXL4:SMXL4-YFP smxl4;5*

D *SMXL5:SMXL5-YFP smxl4;5*

C *SMXL4:SMXL4-YFP ops;smxl4*

E *SMXL5:SMXL5-YFP ops;smxl5*

Figure 2 : Analysis of interaction between *SMXL4* and *SMXL5* with other phloem regulators
(A) Root length of five day-old plants. n = 37–75. Statistical groups determined by one-way ANOVA and post-hoc Tukey's test (95 % CI). Shown is one representative experiment of three repetitions.
(B – E) Comparison of *SMXL4:SMXL4-YFP* and *SMXL5:SMXL5-YFP* reporter activities in *smxl4;smxl5* and *smxl;ops* mutants. Arrows indicate the signals closest to the QC. Scale bars represent 50 μm.

Figure 3: Genetic interaction of *SMXL5* with *OPS* and *BRX*

(A) Root length of five day-old wild type (WT), *smxl5*, *brx*, *ops*, *smxl4;smxl5*, *brx;smxl5*, and *ops;smxl5* plants. n = 38–76 for ech group. Statistical groups determined by one-way ANOVA and post-hoc Tukey's HSD test (95 % CI). Shown is one representative experiment of three repetitions.

(B) Phenotypic characterization of phloem development of two day-old wild type, *smxl5*, *brx*, *ops*, *smxl4;smxl5*, *brx;smxl5*, and *ops;smxl5* plants. n = 97–129 for each group.

Figure 4: SMXL5 and OBE3 proteins interact

(A) The SMXL5 protein was expressed in yeast fused to the GAL4 DNA binding domain (BD) and OBE3 fused to the GAL4 activation domain (AD). All strains contained AD- and BD-expressing plasmids either alone or fused with SMXL5, OBE3 or control proteins, respectively. Growth on SD-leu-trp indicate the presence of both plasmids, growth on SD-leu-trp-his-ade medium indicate the presence of plasmids and protein interaction.

(B) Interaction of SMXL5-3xHA and 6xMyc-OBE3 proteins by co-immunoprecipitation (co-IP) and subsequent Western blot analysis after transient overexpression in *N. benthamiana*. 'Input' represents unprocessed protein extracts, 'unbound' show proteins that remained in the extract after IP and 'IP: α HA' depicts samples after immunoprecipitation by α-HA-beads. Western blots were probed by α HA or α Myc antibodies, respectively. Signals revealed an expected SMXL5-3xHA protein size of approximately 120 kDa. The size of the detected 6xMyc-OBE3 protein (~100 kDa) exceeded the expected size of 92 kDa.

(C – K) Fluorescent signals and bright field images of epidermal *N. benthamiana* nuclei transiently co-expressing OBE3-mGFP/mCherry (C-E), OBE3-mGFP/SMXL5-mCherry (F-H) and mGFP-NLS/mCherry-NLS (I-K). The dashed yellow line indicates the outlines of nuclei in merged images (E, H, K). Scale bars represent 5 µm.

(L) FRET-FLIM analysis of transiently transformed *N. benthamiana* epidermal leaf cells expressing the OBE3-mGFP donor without an mCherry acceptor or in the presence of NLS-mCherry or SMXL5-mCherry. Error bars indicate standard deviation. Data were derived from three biological replicates with n = 27–80. To test for homogeneity of variance, a Brown-Forsythe test was applied. As the variances are not homogenous a Wilcoxon / Kruskal-Wallis test was carried out followed by a Dunn post hoc analysis.

(M) Heat maps of representative nuclei used for FLIM measurements. The donor lifetimes of OBE3-mGFP are color-coded according to the scale at the bottom. Size bars represent 4 µm.

Figure 5: *OBE3* genetically interacts with *SMXL3/4/5* in root growth regulation

(A) 10 day-old wild type, *smxl4*, *smxl5*, *smxl4;smxl5*, *obe3*, *obe4*, *smxl4;obe3*, *smxl4;obe4*, *smxl5;obe3*, *smxl5;obe4* seedlings are shown from left to right. Scale bar represents 1 cm.

(B) Quantification of root length depicted in A. Mean values of four independent experiments (n = 15-55 per experiment and genotype) were analyzed by a one-way ANOVA with post-hoc Tukey HSD (95 % CI). Statistical groups are marked by letters.

(C) Quantification of root length depicted in D. Mean values of three independent experiments (n = 62–74 per experiment and genotype) were analyzed by a one-way ANOVA with post-hoc Tamhane-T2 (95 % CI). Statistical groups are marked by letters.

(D) 10 day-old wild type, *smxl3*, *smxl5*, *smxl3;smxl5*, *obe3*, *smxl3;obe3*, *smxl5;obe3* seedlings are shown from left to right. Scale bar represents 1 cm.

Figure 6: *OBE3* interacts with *SMXL5* during protophloem formation

(A – C) Phloem-specific activity of *SMXL4:SMXL4-YFP* (A) coincides with activty of *OBE3:OBE3-GFP* (B) and *OBE4:OBE4-GFP* (C) reporters in the developing phloem. Fluorescent signals (green) and cell wall staining by Direct Red 23 (magenta) were detected by confocal microscopy. Pink arrows point to the first differentiated SE indicated by enhanced cell wall staining. Scale bars represent 50 µm.

(D) Schematic representation of one developing phloem pole at the root tip. Two tangential cell divisions generate SE precursor and procambium (orange arrow) and proto- and metaphloem cell lineages (blue arrow), respectively. Differentiated SEs are marked by a pink arrow. The QC is marked by a yellow asterisk.

(E – J) Phloem formation in 2 day-old wild type (E), *obe3* (F), *smxl4;smxl5* (G), *smxl4;obe3* (H), *smxl5;obe3* (I) and *smxl5;obe4* (J) root tips. Cell walls were stained by mPS-PI (white). Yellow asterisks mark the QC. Enhanced mPS-PI staining indicates differentiation of SEs (pink arrows) in wild type (E), *obe3* (F) and *obe4;smxl5* (G). Orange and blue arrows mark the first and second tangential division, respectively, in the developing phloem cell lineage. Scale bars represent 20 µm.

(K) The distance from the QC to the first and second tangential division shown in D-I was quantified (n = 9–11). Statistical groups are indicated by letters and were determined by a one-way ANOVA with post-hoc Tukey HSD (95 % CI). Distances of 1st cell divisions and 2nd cell divisions were compared independently.

Figure 7: Phloem-specific *OBE3* expression is sufficient for promoting root growth
(A – B) Seven day-old *smxl5;obe3* root tips carrying an *SMXL5:OBE3-turquoise* reporter (green signal, B) were compared to a non-transformed wild type (WT) root tip (B). Cell walls were stained by DirectRed (magenta). White arrows mark differentiated SEs. Scale bars indicate 20 µm.
(C) Root length quantification of plants depicted in D. Results from one representative experiment out of three independent experiments (n = 61–71 per experiment and genotype) are shown. Mean values were analyzed by one-way ANOVA with post-hoc Tamhane-T2 (95 % CI). Statistical groups are indicated by letters.
(D) 10 day-old wild type, *smxl5*, *obe3*, *smxl5;obe3*, *SMXL5:OBE3-turquoise/smxl5;obe3*, *SMXL5:obe3-miRNA3/smxl5* and *SMXL5:obe3-miRNA4/smxl5* seedlings are shown from left to right. Scale bar represents 1 cm.

Figure 8: *SMXL* and *OBE3* genes determine phloem-related chromatin profile
(A) Summary of ATAC-seq results for the different samples. Specific OCRs were identified by comparing GFP-positive and GFP-negative samples of the respective genetic backgrounds. After initial alignment, OCR detection and other subsequent analyses were conducted using 40 M reads aligning to the nuclear genome excluding duplicates for each sample.
(B) Chromatin conformation profiles for all genes found in the Arabidopsis genome or only for phloem and non-phloem genes according to Brady et al., 2007. Read alignment was adjusted to transcriptional start sites (TSS) and transcriptional end sites (TES). X-axes show distance to TSSs and TES in base-pairs. Inserts in all graphs are enlargements of the peaks identified close to TSSs.
(C, D) Comparison of OCR profiles found in GFP-positive (C) and GFP-negative (D) samples. Numbers indicate the number of OCRs with significantly more ('up') or less ('down') aligned reads.