**1**    **Ecological and evolutionary drivers of hemoplasma infection and bacterial genotype**

**2**    **sharing in a Neotropical bat community**

Daniel J. Becker[1,2*], Kelly A. Speer[3-5], Alexis M. Brown[6], M. Brock Fenton[7], Alex D.

Washburne[8], Sonia Altizer[2,9], Daniel G. Streicker[9-11], Raina K. Plowright[8], Vladimir E.

Chizhikov[12§], Nancy B. Simmons[3,13], Dmitriy V. Volokhov[12]

[1]Department of Biology, Indiana University, Bloomington, IN, USA

[2]Center for the Ecology of Infectious Disease, University of Georgia, Athens, GA, USA

[3]Richard Gilder Graduate School, American Museum of Natural History, New York, NY, USA

[4]Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian

Institution, Washington, D.C., USA

[5]Center for Conservation Genomics, Smithsonian Conservation Biology Institute, National

Zoological Park, Washington, D.C., USA

[6]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA

[7]Department of Biology, Western University, London, Ontario, Canada

[8]Department of Microbiology and Immunology, Montana State University, Bozeman, MT USA

[9]Odum School of Ecology, University of Georgia, Athens, GA USA

[10]MRC–University of Glasgow Centre for Virus Research, Glasgow, United Kingdom

[11]Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

United Kingdom

[12]Center for Biologics Evaluation and Research, Food and Drug Administration, Silver Spring,

MD, USA

[13]Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural

History, New York, NY

[*]Corresponding author, danbeck@iu.edu

[§]Currently retired

**Running head**: Ecology and evolution of bat hemoplasmas

**Keywords:** 16S rRNA; cophylogeny; host shifts; parasite sharing; bacterial zoonosis; host

specificity

**Abstract**

Most emerging pathogens can infect multiple species, underscoring the importance of understanding the ecological and evolutionary factors that allow some hosts to harbor greater infection prevalence and share pathogens with other species. However, our understanding of pathogen jumps is primarily based around viruses, despite bacteria accounting for the greatest proportion of zoonoses. Because bacterial pathogens in bats (Order: Chiroptera) can have conservation and human health consequences, studies that examine the ecological and evolutionary drivers of bacterial prevalence and barriers to pathogen sharing are crucially needed. We here studied hemotropic *Mycoplasma* spp. (i.e., hemoplasmas) across a species-rich bat community in Belize over two years. Across 469 bats spanning 33 species, half of individuals and two-thirds of species were hemoplasma positive. Infection prevalence was higher for males and for species with larger body mass and colony sizes. Hemoplasmas displayed high genetic diversity (21 novel genotypes) and strong host specificity. Evolutionary patterns supported co-divergence of bats and bacterial genotypes alongside phylogenetically constrained host shifts. Bat species centrality to the network of shared hemoplasma genotypes was phylogenetically clustered and unrelated to prevalence, further suggesting rare—but detectable—bacterial sharing between species. Our study highlights the importance of using fine phylogenetic scales when assessing host specificity and suggests phylogenetic similarity may play a key role in host shifts for not only viruses but also bacteria. Such work more broadly contributes to increasing efforts to understand cross-species transmission and epidemiological consequences of bacterial pathogens.

**Introduction**

Most pathogens that cause disease in humans, domestic animals, and wildlife are capable of infecting multiple host species (Woolhouse, Taylor, & Haydon, 2001). However, predicting which hosts maintain pathogens and identifying their role in cross-species transmission can be challenging, as many hosts can be infected but not play key roles in the reservoir community (A. Fenton, Streicker, Petchey, & Pedersen, 2015). Pathogen jumps between species depends on infection prevalence in the donor host, transmission opportunities between donor and recipient species, and suitability of the recipient host for pathogen replication (Plowright et al., 2017). Each of these steps can be shaped by ecological and evolutionary factors (VanderWaal & Ezenwa, 2016). For example, small-bodied species can have greater competence, the ability to transmit new infections, than larger species (Downs, Schoenle, Han, Harrison, & Martin, 2019), and host switching is often constrained by phylogeny, owing to similarity in immunological barriers to pathogen replication between closely related species (Streicker et al., 2010). Identifying the ecological and evolutionary factors that allow some species to harbor greater prevalence and have facilitated pathogen sharing can improve our general understanding of disease emergence (Fountain-Jones et al., 2018). Examining evolutionary associations between hosts and pathogens can further uncover factors favoring host shifts versus codivergence and assess the frequency of cross-species transmission (Geoghegan, Duchêne, & Holmes, 2017).

Given the public health and agricultural burdens of many zoonotic pathogens such as SARS coronavirus, avian influenza virus, and rabies virus, many investigations of pathogen prevalence and emergence focus on viruses (Geoghegan et al., 2017; Olival et al., 2017). However, more zoonoses are caused by bacteria than other pathogen taxa (Han, Kramer, & Drake, 2016), and bacterial pathogens can negatively impact newly infected host species (e.g.,

*Mycoplasma galliscepticum*, a poultry pathogen, caused rapid population declines in wild house

finches; Hochachka & Dhondt, 2000). More attention to bacteria and their propensity for host

specificity versus generalism is accordingly important for understanding whether factors that

govern cross-species transmission of viruses can be extended to other pathogens (Bonneaud,

Weinert, & Kuijper, 2019). Bacterial pathogens have been especially understudied for bats

(Mühldorfer, 2013), in contrast to intensive studies of zoonotic viruses across the Chiroptera

(Luis et al., 2015). However, many bacterial pathogens are likely important to both bat

conservation and human health due to pathogenic effects on bats themselves as well as their

zoonotic potential (Becker et al., 2018; Evans, Bown, Timofte, Simpson, & Birtles, 2009).

To determine the ecological and evolutionary drivers of bacterial prevalence and barriers

to pathogen sharing, we focused on hemotropic *Mycoplasma* spp. (i.e., hemoplasmas) in a

species-rich bat community in Belize (Fenton et al., 2001; Herrera, Duncan, Clare, Fenton, &

Simmons, 2018). The Neotropics have remarkable bat diversity owing to adaptive radiation in

the Phyllostomidae (Gunnell & Simmons, 2012), producing a range of feeding strategies (e.g.,

frugivory, carnivory, sanguivory), body sizes, and roost preferences (Monteiro & Nogueira,

2011). Hemoplasmas are intracellular erythrocytic bacteria transmitted through direct contact

(Cohen et al., 2018; Museux et al., 2009) and also possibly arthropod vectors (Willi, Boretti,

Meli, et al., 2007). Hemoplasmas can cause acute and chronic anemia, especially for

immunocompromised hosts; however, many animals develop inapparent infections and are

asymptomatic (Messick, 2004). As *Mycoplasma* spp. lack many of the metabolic pathways

associated with energy production and synthesis of cell components found in other bacteria, they

are fully dependent on host cells (Citti & Blanchard, 2013). Hemoplasmas have therefore been

described as mostly host specialists (Pitcher & Nicholas, 2005), yet interspecies and potentially

zoonotic transmission can occur (Maggi et al., 2013; Willi, Boretti, Tasker, et al., 2007).

Hemoplasmas are common and genetically diverse in bats (Di Cataldo, Kamani, Cevidanes,

Msheliza, & Millán, 2020; Ikeda et al., 2017; Mascarelli et al., 2014; Millán et al., 2019, 2019;

Volokhov et al., 2017), which offers an ideal model system for identifying the ecological and

evolutionary factors structuring bacterial infection risks within and between host species.

Many cross-species comparisons of pathogen infection risks and sharing have used less-

diverse host communities (Johnson et al., 2012; VanderWaal, Atwill, Isbell, & McCowan, 2014)

or global datasets of host–pathogen associations that can be limited by heterogeneous sampling

effort and variation in pathogen detection methods (Dallas et al., 2019; Huang, Bininda-Emonds,

Stephens, Gittleman, & Altizer, 2014). Our focus on a widespread pathogens group in a highly

diverse host community allowed us to capitalize on strong host trait variation while controlling

for sampling effort and diagnostics methods (Becker, Crowley, Washburne, & Plowright, 2019;

Han, Kramer, et al., 2016). Past work has also used host–pathogen networks to characterize

contemporary or historic transmission at often coarse taxonomic scales (e.g., pathogen species

complexes or genera; Blyton, Banks, Peakall, Lindenmayer, & Gordon, 2014; VanderWaal et al.,

2014). However, as bat species can be infected by multiple hemoplasma genotypes, and because

genotypes with ≥99% sequence identity of their 16S rRNA genes can represent different

bacterial species (Volokhov, Hwang, Chizhikov, Danaceau, & Gottdenker, 2017; Volokhov,

Simonyan, Davidson, & Chizhikov, 2012), focusing on genotypes provides finer-scale resolution

to determine the ecological and evolutionary features of species that facilitate pathogen sharing

and to identify likely maintenance hosts of bacterial infections (Fountain-Jones et al., 2018).

We asked three questions about the relative contribution of ecological traits and

evolutionary history to structuring infection patterns and pathogen sharing. First, what are the

individual and ecological predictors of hemoplasma infection in a Neotropical bat community?

Second, does the distribution of hemoplasma genotypes across the bat community map onto the

bat phylogeny, as predicted by host–pathogen codivergence? Lastly, if genotype sharing among

host species occurs, which host clades or traits best predict species ability to share pathogens?

We predicted that ecological covariates such as ectoparasitism and large host colonies could

increase bacteria risk through vector-borne or density-dependent transmission (McCallum,

Barlow, & Hone, 2001; Willi, Boretti, Meli, et al., 2007). We also expected hemoplasma

genotypes would be specific to particular host species but that more closely related bats would

share hemoplasma genotypes, indicating phylogenetically restricted host shifts (Pitcher &

Nicholas, 2005). However, ecological traits that increase the risk of pathogen exposure between

species, such as occupying a greater diversity of roosting habitats, could also facilitate pathogen

genotype sharing among less closely related hosts (McKee et al., 2019).

**Methods**

*Bat capture and sampling*

From April 24 to May 6 2017 and from April 23 to May 5 2018, we sampled 469 bats from 33

species captured in two adjacent areas in the Orange Walk District of Belize: Lamanai

Archaeological Reserve (LAR) and Ka'Kabish (KK; Fig. 1). The LAR is bordered by the New

River Lagoon, forest, and agriculture, while KK is a remnant patch of forest surrounded by

agriculture located 10 km away. We consider these sites to be independent, as the small home

ranges of many Neotropical bat species and lack of continuous forest between LAR and KK has

likely restricted most individual bat inter-site movement (Jones, Hämsch, Page, Kalko, &

O'mara, 2017; Loayza & Loiselle, 2008); however, some species may have greater connectivity

(e.g., inter-roost movement of *Desmodus rotundus* occurs rarely). At least 44 of the 70 bat

species in Belize have been recorded in this region (Herrera et al., 2018; Reid, 1997). Bats were

captured with mist nets primarily along flight paths and occasionally at the exits of roosts from

19:00 until 22:00. Harp traps were also set from 18:00 to 05:00. In KK, *Desmodus rotundus*,

*Trachops cirrhosus*, and *Chrotopterus auritus* were captured at their shared roost. In the LAR,

*Desmodus rotundus*, *Saccopteryx bilineata*, and *Glossophaga soricina* were sampled from a

shared roost, although individuals were also sampled along flight paths across the greater site.

More broadly, patterns of roost sharing of bat species in northern Belize remain elusive.

Bats were placed in cloth bags until processing and were identified to species (and sex)

based on morphology (Reid, 1997). Reproductive activity was indicated by the presence of

scrotal testes in males and by the evidence of pregnancy or lactation in females; across bat

species, 69% of males and 42% of females were in reproductive condition. We also visually

screened bats for the presence of ectoparasites (i.e., bat flies, ticks, bat bugs, mites; Ter Hofstede,

Fenton, & Whitaker, 2004). We collected 3–30 μL of blood (volumes were dependent on bat

mass) by lancing the propatagial vein with a sterile needle. Blood was collected with heparinized

capillary tubes and stored on Whatman FTA cards to preserve bacterial DNA. Field procedures

followed guidelines for safe and humane handling of bats published by of the American Society

of Mammalogists (Sikes, Care, & Mammalogists, 2016) and were approved by the Institutional

Animal Care and Use Committees of the University of Georgia (A2014 04-016-Y3-A5) and

American Museum of Natural History (AMNHIACUC-20170403 and AMNHIACUC-

20180123). Fieldwork was authorized by the Belize Forest Department under permits

WL/2/1/17(16), WL/2/1/17(19), and WL/2/1/18(16). Sample size was similar between years

(2017=202, 2018=267) but varied by site (LAR=365, KK=101). More species were sampled for

blood at LAR (*n*=33) than KK (*n*=17; Fig. 1), reflecting site differences in species richness

(Herrera et al., 2018). We sampled 1–139 individuals per bat species (the maximum being

*Desmodus rotundus*), with a mean of 14 individuals per bat species (Table S1).

*DNA extraction, PCR amplification, and amplicon sequencing*

Genomic DNA was extracted from blood on FTA cards using QIAamp DNA Investigator Kits

(Qiagen). We tested DNA for hemoplasmas using PCR with primers and procedures described in

prior analyses (Volokhov et al., 2017). We included blank FTA punches as an extraction control,

ultrapure water as a negative control, and *Candidatus* Mycoplasma haemozalophi DNA as a

positive control (Volokhov et al., 2011). Amplicons from PCR-positive samples were purified by

electrophoresis and extracted using the QIAquick Gel Extraction Kit (Qiagen).

To determine hemoplasma infection status, all 16S rRNA amplicons were directly

sequenced by Macrogen (https://www.macrogenusa.com/). Amplicons were sequenced with the

same primers used for PCR amplification and then with internal (walking) primers when needed

(Volokhov et al., 2017). Negative DNA samples were tested for amplification quality using the

universal PCR primers targeting the mammal mitochondrial 16S rRNA gene (Volokhov, Kong,

George, Anderson, & Chizhikov, 2008) or the mitochondrial cytochrome *c* oxidase subunit 1

gene (Clare, Lim, Engstrom, Eger, & Hebert, 2007); all hemoplasma-negative DNA samples

gave positive signal in the mitochondrial 16S rRNA gene- and/or the COI-specific PCR assays.

All amplified sequences were subjected to chimeric sequence analysis using DECIPHER

(Wright, Yilmaz, & Noguera, 2012) and UCHIME (Edgar, Haas, Clemente, Quince, & Knight,

2011). All hemoplasma sequences have been deposited in GenBank under accession numbers

MH245119–MH245194 and MK353807–MK353892; four positive samples were identified as

*Bartonella* spp. during sequencing and were considered hemoplasma negative in our analyses.

*Bat phylogenetic data*

We used the *rotl* and *ape* packages in R to extract a bat phylogeny from the Open Tree of Life

and to calculate branch lengths with Grafen's method (Michonneau, Brown, & Winter, 2016;

Paradis, Claude, & Strimmer, 2004). To assess hemoplasma genotype sharing as a function of

host phylogenetic similarity, we derived pairwise phylogenetic distances between the 33 sampled

bat species (Fig. S1). Because more evolutionarily distant and distinct species could display less

frequent bacterial genotype sharing owing to ecological and immunological barriers to pathogen

exposure and replication (Huang, Drake, Gittleman, & Altizer, 2015), we used the *picante*

package and our bat phylogeny to derive evolutionary distinctiveness (Kembel et al., 2010).

*Host species trait data*

We obtained species-level data on host traits relevant to pathogen transmission from previously

published sources (Table S2). We obtained fecundity (litter size, litters per year), body mass, and

diet from the Amniote Life History and EltonTraits databases (Myhrvold et al., 2015; Wilman et

al., 2014). For foraging ecology, which could affect bacterial exposure (e.g., trophic interactions;

Kellner et al., 2018), we defined three dietary guilds: frugivory (including nectarivory; $n=11$),

insectivory ($n=18$), and carnivory (including sanguivory and piscivory, $n=4$; González-Salazar,

Martínez-Meyer, & López-Santiago, 2014). We also considered the proportion of plant-based

items in diet. We simplified foraging strata into aerial ($n=14$), arboreal ($n=16$, including

scansorial), and ground- or aquatic-level foraging ($n=3$). We also expanded prior compilations of

wing aspect ratios and roost preferences to serve as proxies for ecological overlap among species

(Fenton et al., 2001; Herrera et al., 2018; Reid, 1997). Roost type was simplified to open (e.g.,

only foliage; *n*=6) or closed (e.g., hollows, caves; *n*=27), and roost flexibility was simplified to

using one (*n*=16) or multiple roost types (*n*=17). We classified maximum colony sizes as small-

to-medium (e.g., under 100 individuals; *n*=20) or large (e.g., hundreds to thousands; *n*=13; Reid,

1997; Santana, Dial, Eiting, & Alfaro, 2011), as most values were reported in ranges. We did not

record pairwise sympatry (e.g., Luis et al., 2015; McKee et al., 2019) given that all species occur

in Belize (Fig. S2). Yet because more widely distributed species could have more opportunities

for pathogen sharing due to range overlap, we used the *geosphere* package and data from the

International Union for Conservation of Nature to derive geographic range size (Baillie, Hilton-

Taylor, & Stuart, 2004; Hijmans, Williams, Vennes, & Hijmans, 2019). Missing species-level

traits were taken from other databases, primary literature, or closely related species (Table S2).

*Individual-level analysis of bat infection status*

We first used the *prevalence* package to estimate hemoplasma infection prevalence and its 95%

confidence interval (Wilson method). We then used phylogenetic generalized linear mixed

models (GLMMs) to test if infection status varied by sex, reproductive status, year, site, and

ectoparasite presence while accounting for bat phylogenetic relatedness. We fit candidate

GLMMs that considered all fixed effects as well as interactions between sex and reproduction

and between site and year. We also considered a model that excluded site to account for possible

non-independence of LAR and KK alongside an intercept-only model. As vampire bats were

banded for a mark–recapture study (Volokhov et al., 2017) and some were sampled between and

within years (*n*=14), we randomly selected one of each recapture. After removing recaptures and

missing values ($n$=323), we fit the phylogenetic GLMMs using the *brms* package, default priors,

and infection status as a Bernoulli-distributed response. We included random effects for bat

species and phylogeny, the latter of which used the phylogenetic covariance matrix (Bürkner,

2017). We ran four chains for 20,000 iterations with a burn-in period of 10,000, thinned every 10

steps, for a total 4,000 samples. We compared GLMMs using the leave-one-out cross-validation

(LOOIC) and assessed fit with a Bayesian $R^2$, including the total modeled variance and that

attributed to only the fixed effects (Gelman, Goodrich, Gabry, & Vehtari, 2019; Vehtari,

Gelman, & Gabry, 2017). We then estimated fixed effects (means and 95% highest density

intervals [HDI]) from the posterior distributions of each predictor from the top GLMM. Lastly,

we assessed the sensitivity of individual-level analyses to overrepresentation of *Desmodus*

*rotundus* by randomly subsampling this species by the maximum $n$ for other bat species.

*Species-level analysis of hemoplasma prevalence*

We next calculated infection prevalence per species, using the *metafor* package to estimate logit-

transformed proportions and sampling variances (Viechtbauer, 2010). We used the *nlme* package

to estimate phylogenetic signal as Pagel's λ with a weighted phylogenetic generalized least

squares (PGLS) model to account for within-species variance (Garamszegi, 2014). We next used

a graph-partitioning algorithm, phylogenetic factorization, to flexibly identify clades with

significantly different prevalence estimates at various taxonomic depths. We used the *taxize*

package to obtain a bat taxonomy from the National Center for Biotechnology Information

(Chamberlain & Szöcs, 2013) and used the *phylofactor* package to partition prevalence as a

Bernoulli-distributed response in a GLM (Washburne et al., 2019). We determined the number of

significant bat clades using Holm's sequentially rejective test with a 5% family-wise error rate.

To identify species trait correlates of prevalence, we fit 11 PGLS models (weighted by

sampling variance) with body mass, annual fecundity (litters per year * pups per litter), dietary

guild, quantitative diet, foraging strata, aspect ratio, roost type, roost flexibility, colony size,

geographic range size, and evolutionary distinctiveness as predictors. We also fit PGLS models

with only sample size or an intercept. We compared models with Akaike information criterion

corrected for small sample sizes (AICc) and estimated $R^2$ (Burnham & Anderson, 2002).

*Hemoplasma phylogenetic analyses and genotype assignment*

We compared our 16S rRNA sequences to those in GenBank (Volokhov et al., 2017; Volokhov

et al., 2011). Briefly, we aligned sequences using Clustal X, and inter- and intra-species

similarity values were generated using BioEdit. Genetic distances were calculated with the

Kimura two-parameter and Tamura–Nei models, and the phylogeny was constructed using

MEGA X with the minimum evolution algorithm (Kumar, Stecher, Li, Knyaz, & Tamura, 2018).

We assigned hemoplasma genotypes to positive bats based on analysis of the 16S rRNA

partial gene (860–1000 bp) sequences in GenBank and their clustering on the phylogeny.

Genotypes were designated as novel if (*i*) sequences differed from the closest hemoplasma

sequences in GenBank by ≥1.5% and/or (*ii*) if sequence similarity was <1.5% but genotype-

specific reproducible mutations (at least two per sequence) were observed between hemoplasma

sequences from at least two independent bat samples and the nearest GenBank hemoplasma

sequences. These genotype-specific mutations were further used to differentiate closely related

hemoplasma genotypes from our sample. We caution that genotype is not synonymous with

species, as analysis of the 16S rRNA gene alone is insufficient for accurate species identification

of *Mycoplasma* spp. (Volokhov et al., 2012). Future studies using genomics or housekeeping

genes may identify independent but closely related hemoplasma species in our genotypes.

To assess if hemoplasma genotype assignments were associated with site and year, we

used $\chi^2$ tests with *p* values generated through a Monte Carlo procedure. Prior to our phylogenetic

and network analyses of genotype distributions across bat species (see below), we used another

$\chi^2$ test to assess the association between hemoplasma genotype identify and bat host identity.

*Evolutionary relationships between bats and hemoplasmas*

To determine the degree to which bat hemoplasma genotypes display host specificity and to

describe their evolutionary relationships with host species, we used our bat and hemoplasma

phylogenies to construct a binary association matrix. To test the dependence of the hemoplasma

phylogeny upon the bat phylogeny and thus assess evidence of evolutionary codivergence, we

applied the Procrustes Approach to Cophylogeny (PACo) using distance matrices and the *paco*

package (Hutchinson, Cagua, Balbuena, Stouffer, & Poisot, 2017). We used a jackknife

procedure to estimate the degree to which each bat–genotype link supported a hypothesis of

phylogenetic congruence; links were supported if their upper 95% confidence interval was below

the mean of all squared jackknife residuals (Balbuena, Míguez-Lozano, & Blasco-Costa, 2013).

*Hemoplasma genotype sharing among bat species*

We used hemoplasma genotype assignments to create a network, with each node representing a

bat species and edges representing shared genotypes among bat species pairs. We built an

adjacency matrix using the *igraph* package and used the Louvain method to assess the structure

of bat–hemoplasma communities within this network (Csardi & Nepusz, 2006). To test whether

the distribution of hemoplasma genotypes across our Neotropical bat species is shaped by host

phylogeny, we used two GLMs to predict counts of shared genotypes (Poisson errors) and the

presence of sharing (binomial errors) by phylogenetic distance between bat species. We assessed

statistical significance with a quadratic assignment procedure via the *sna* package (Butts, 2008).

We calculated two metrics of network centrality to quantify different aspects of how

important a node (bat species) is to hemoplasma genotype sharing: degree and eigenvector

centrality (Bell, Atkinson, & Carlson, 1999). Whereas degree indicates the number of other

species with which a host shares bacterial genotypes (i.e., links per node), eigenvector centrality

indicates the tendency for a host to share genotypes with species that also share more genotypes

(i.e., connectivity). Eigenvector centrality is thus an extension of degree that can identify hubs of

parasite sharing (Gómez, Nunn, & Verdú, 2013). These two metrics were moderately correlated

($\rho$=0.59), with many non-zero degree species displaying zero eigenvector centrality. To examine

spatial and temporal patterns in host centrality, we built separate adjacency networks per each

site and year. We fit separate GLMs to ask how hemoplasma sharing centrality was predicted by

site, year, and the two-way interaction. Degree was modeled as a Poisson-distributed response,

while eigenvector centrality was logit-transformed and used Gaussian errors. We next applied

phylogenetic factorization to both metrics and weighted the algorithms by the square-root sample

size per species (Garamszegi, 2014). We then fit the same PGLS models used in our prevalence

analysis to identify the most competitive trait predictors of bat species centrality to hemoplasma

sharing. To lastly assess whether network centrality is associated with hemoplasma prevalence,

we fit two weighted PGLS models with each centrality metric as a univariate predictor.

**Results**

*Hemoplasma infection status*

We detected sequence-confirmed hemoplasma infection in 239 of 469 individuals (51%; 95%

CI: 46–55%), with positive individuals in 23 of the 33 sampled bat species (Table S3). The most

parsimonious phylogenetic GLMM explained 22% of the modeled variance in infection status

and included only sex, reproductive status, ectoparasites, and year ($\Delta$LOOIC=0.16, $w_i$=0.27;

Table S4), suggesting site to be uninformative. Males had higher odds of infection than females

(OR=2.35, 95% HDI: 1.36–4.03), but risk was unrelated to ectoparasitism, reproduction, or year

(Fig. 2A). This result was weakly sensitive to overrepresentation by *Desmodus rotundus*, as

effect size and the prevalence sex difference was weaker when we randomly subsampled this

species (Fig. 2). Across all models, fixed effects only explained up to 7% of the modeled

variance, suggesting more variation explained by the species and phylogeny random effects.

*Inter-species variation in hemoplasma prevalence*

Across bat species, hemoplasma prevalence ranged from 0% to 100% ($\tilde{x}$=0.37). We estimated

Pagel's $\lambda$ in logit-transformed prevalence to be 0.39, indicating moderate phylogenetic signal.

Similarly, phylogenetic factorization identified one bat clade with significantly lower prevalence

compared to the paraphyletic remainder: the Emballonuridae (12% infected; Fig. 3A). Our trait-

based analysis showed that relatively larger species ($\beta$=1.48, $p$=0.01, $R^2$=0.24) and those with

larger colonies ($\beta$=0.67, $p$=0.06, $R^2$=0.20) had higher prevalence (Fig. 3B; Table 1). Relatively

heavier ($\geq$ 20 g) and larger colony species included *Desmodus rotundus*, *Molossus rufus*, and

*Pteronotus mesoamericanus*, for which prevalence was greater than 58%. Although these three

species were also heavily sampled, other well-sampled species such as *Sturnira parvidens* and

*Carollia sowelli* had lower prevalence, and sample size did not predict prevalence (Table 1).

*Hemoplasma genotype diversity*

Our phylogenetic analysis identified 29 *Mycoplasma* genotypes in the Belize bat community

(Table 2), including three previously identified from vampire bats (VBG1–3; Volokhov et al.,

2017). All genotypes demonstrated minor levels of sequence variability (99.5–100%; Table 2).

Based on comparisons with sequences from GenBank, 21 of these genotypes represent novel

hemoplasmas, five are closely related to non-hemoplasma mycoplasmas, and many are

phylogenetically related to previously identified *Mycoplasma* spp. and hemoplasmas identified

from other bat species (e.g., Di Cataldo et al., 2020; Millán et al., 2019, 2015), bat ticks (e.g.,

Hornok et al., 2019), primates including humans (e.g., Alcorn et al., 2020; Bonato, Figueiredo,

Gonçalves, Machado, & André, 2015; Hattori et al., 2020), and rodents (e.g., Gonçalves et al.,

2015; Goto, Yasuda, Hayashimoto, & Ebukuro, 2010; Vieira et al., 2009). A more detailed

description of these 29 bacterial genotypes is provided in the Supporting Information (Fig. S3).

After controlling for multiple comparisons, our 29 bacterial genotypes were associated

with site ($\chi^2$=47.11, $p$<0.01) and year ($\chi^2$=40.40, $p$<0.01). Genotype composition was more

diverse at LAR (Fig. S4), and KK hemoplasmas were dominated by vampire bat genotypes

(VBG1–3). Genotype composition was more idiosyncratic by study year. However, these 29

bacterial genotypes were most strongly associated with bat species ($\chi^2$=3532, $p$<0.01; Fig. S4).

*Bat–hemoplasma evolutionary relationships*

Although some hemoplasma genotypes were shared between bat species (i.e., VBG1, CS2, PPM,

EF1, AH1–2, MYE, PLU, SP; *n*=9), most showed strong host specificity (*n*=20; Table 2). Our

coevolutionary analysis (PACo) supported congruence between the bat and hemoplasma

phylogenies ($m^2_{XY}$=36.21, $p$=0.02, $n$=1000; Fig. 4), suggesting that hemoplasma evolution has

mostly tracked bat speciation. However, PACo also demonstrated that only 56% of the 41 unique

bat–hemoplasma links displayed significant evidence of coevolution (Fig. S5), and these patterns

were almost exclusively found within the Phyllostomidae (with the exception of *Saccopteryx*

*bilineata* and its *Mycoplasma muris*–like bacterial genotype). The other 18 bat–hemoplasma

links therefore displayed evidence of phylogenetic incongruence and thus likely host shifts.

*Hemoplasma genotype sharing networks*

Within our bat–hemoplasma network, genotype sharing was restricted to five host communities,

whereas six genotypes were each restricted to a single bat species (Fig. 5A). GLMs showed that

both the frequency and presence of genotype sharing declined with phylogenetic distance

between bat species (Poisson: $p$<0.001, $R^2$=0.08; binomial: $p$<0.001, $R^2$=0.51; Fig. 5B).

Bat species shared hemoplasma genotypes with zero to five other species (i.e., degree),

and most hosts were not central to the network of genotype sharing (i.e., eigenvector centrality of

zero). Six bat species had non-zero eigenvector centrality values that ranged from 37% to 100%,

indicating that these hosts generally shared more hemoplasma genotypes with other highly

connected hosts. Stratifying our hemoplasma genotype network across sites and years showed

that centrality measures varied by space but not time (Fig. S7, Table S5). We observed no

hemoplasma genotype sharing at KK, likely reflecting lower host diversity (Herrera et al., 2018).

Phylogenetic factorization identified similar bat clades with significantly different

centrality compared to the paraphyletic remainder (Fig. 6A–B). For degree, the algorithm only

identified *Artibeus lituratus* as being more central (x☐=5) than other bats (x☐=1.14). However,

phylogenetic factorization identified three taxa in the subfamily Stenodermatinae that had

significantly elevated eigenvector centrality: the genera *Artibeus* and *Dermanura* ($\bar{x}$=0.67

compared to $\bar{x}$=0.02 in all other bats), the species *Artibeus lituratus* ($\bar{x}$=1 compared to

$\bar{x}$=0.12), and the species *Sturnira parvidens* ($\bar{x}$=0.37 compared to $\bar{x}$=0.15). Mirroring these

results, phylogenetic signal was absent for degree ($\lambda$=0) but high for eigenvector centrality

($\lambda$=0.93).

Trait-based analyses showed that degree centrality was best predicted by diet (Table S6);

bat species feeding more heavily on fruit and nectar shared more bacterial genotypes with other

species ($\beta$=0.004, $p$<0.001, $R^2$=0.20; Fig. 6C). Similarly, eigenvector centrality was best

predicted by bat colony size and diet (Table S7); highly central species had small colonies

($\beta_{large}$=–1.93, $p$=0.05, $R^2$=0.13) and fed more on plants ($\beta$=0.03, $p$<0.01, $R^2$=0.10; Fig. 6D).

As a final analysis, we assessed whether network centrality (i.e., a bat species' role in

hemoplasma genotype sharing) predicted contemporary infection prevalence (Fig. S8). However,

we found no associations between species-level infection prevalence and centrality as measured

by degree ($\beta$=–0.13, $R^2$=0.03, $p$=0.42) or eigenvector centrality ($\beta$=–0.20, $R^2$<0.01, $p$=0.79).

**Discussion**

By examining the prevalence and distribution of a common bacterial pathogen (hemoplasmas) in

a diverse bat community, we expanded analysis of the ecological and evolutionary predictors of

bat infection and pathogen sharing beyond viruses. Across the bat community, hemoplasma

infection risk was weakly higher for males but was better predicted by phylogeny, with large-

bodied and large-colony bat species showing greater prevalence. Hemoplasmas showed high

diversity and mostly strict host associations, with strong congruence between the bat and

hemoplasma phylogenies. Although codivergence was supported by our analyses, we also

observed hemoplasma genotype sharing and evidence of historical host shifts between closely

related bats. Species most central to this hemoplasma sharing network displayed taxonomic

clustering and were disproportionately frugivores and nectarivores. Yet these highly central bat

species did not also have the highest hemoplasma prevalence, reinforcing mostly infrequent

bacterial sharing between species. Our work reveals phylogenetic patterns in hemoplasma

infection in a diverse bat community while contributing to broader efforts to understand the host

specificity of bacterial pathogens and their cross-species transmission patterns in wildlife.

Whereas many bacterial pathogens, including hemoplasmas, are common in bats (Bai et

al., 2011; Becker et al., 2018; Ikeda et al., 2017; Mascarelli et al., 2014; Millán et al., 2015;

Volokhov et al., 2017), the factors that confer high infection probability are poorly understood.

In the Belize bat community, the odds of hemoplasma infection tended to be higher in males,

suggesting male-biased transmission as detected in feline and canine systems (Soto et al., 2017;

Walker Vergara et al., 2016). Such patterns could stem from males mounting weaker immune

responses than females (Kelly, Stoehr, Nunn, Smyth, & Prokop, 2018) or to male defense of

multi-female roosts in many Neotropical bat species (Voigt, von Helversen, Michener, & Kunz,

2001). Direct transmission of hemoplasmas has been demonstrated in feline and rodent systems

(Cohen et al., 2018; Museux et al., 2009) but only inferred in bats from metagenomic studies

detecting these bacteria in saliva (Volokhov et al., 2017). We found weak support for the

hypothesis that ectoparasites play a role in infection risks (Hornok et al., 2019; Willi, Boretti,

Meli, et al., 2007). The weak male bias in infection could cast further doubt on vector-borne

transmission, as females in some bat species have elevated ectoparasitism (Frank, Mendenhall,

Judson, Daily, & Hadly, 2016); however, a secondary GLMM testing this hypothesis found

generally weak support for a female bias in ectoparasitism in our system (Fig. S9). Future work

assessing ectoparasite burdens could better elucidate the roles of vectors in hemoplasma risk.

Across Neotropical bats sampled in Belize, we found phylogeny to be a better predictor

of hemoplasma risk than individual traits, site, or year. Phylogenetic factorization identified one

clade, the Emballonuridae, with significantly lower prevalence than all other bats in the

community. This moderate phylogenetic signal mirrors comparable effects of phylogeny for bat

viruses (Guy, Thiagavel, Mideo, & Ratcliffe, 2019), similarly suggesting potential for innate

differences in species susceptibility or pathogen exposure. Trait-based analyses revealed that this

taxonomic pattern was driven by heavier and large-colony species having greater hemoplasma

prevalence. Small-bodied species could have low prevalence due to small blood volumes and

low bacterial titers (Volokhov et al., 2017). Alternatively, the positive, saturating relationship

between body mass and bacterial prevalence could be driven by allometric patterns in

competence (Downs et al., 2019), in contrast to weak or opposite relationships between mass and

viral richness across bats (Guy et al., 2019; Han, Schmidt, et al., 2016). As larger-bodied bat

species can also be more abundant in Neotropical habitat fragments (Herrera et al., 2018), these

results suggest land conversion could increase the frequency of bat species most capable of

maintaining hemoplasma infection. Similarly, positive relationships between colony size and

prevalence could support density-dependent transmission of bacteria (McCallum et al., 2001),

whereas mixed support has been found for bat viruses (Streicker et al., 2012; Webber, Fletcher,

& Willis, 2017). Future work could test how community-wide infection patterns vary across

broader habitat gradients and use multiple bacteria to assess the generality of these trends.

Approximately two-thirds of the Neotropical bat species sampled in Belize were infected

by hemoplasmas, for which we observed high genetic diversity consistent with other studies of

this pathogen in bats (Mascarelli et al., 2014; Millán et al., 2015; Volokhov et al., 2017).

However, these bacterial genotypes were mostly novel and only weakly related to hemoplasmas

described elsewhere in Latin America (Ikeda et al., 2017; Millán et al., 2019), with the exception

of those previously identified from vampire bats (Volokhov et al., 2017). When considering the

phylogenetic scale of genotypes, most hemoplasmas were host specific. Over half our

hemoplasma communities consisted of a single bat–genotype association, matching the degree of

host specificity observed more generally for *Mycoplasma* spp. (Citti & Blanchard, 2013; Pitcher

& Nicholas, 2005). When we did detect genotype sharing between species, this occurred mostly

between closely related hosts (e.g., PPM was detected in *Pteronotus mesoamericanus* and *P.*

*fulvus*), indicating bat phylogenetic distance decreased the probability of bacterial transfer.

Analyses to characterize species centrality to the hemoplasma genotype sharing network

showed that one species (*Artibeus lituratus*) and the subfamily Stenodermatinae played key

roles. This clade, and especially the genera *Artibeus* and *Dermanura* (formerly all classified in

*Artibeus*), was the only taxon with non-zero connectivity, and this pattern was reflected in fruit-

and nectar-based diets and small colonies being the primary predictors of centrality. The strictly

frugivorous Stenodermatinae represents a recent divergence in the Phyllostomidae (Botero-

Castro et al., 2013), and high centrality of these species may indicate weaker phylogenetic

barriers for bacterial transmission between hosts in this clade. Two other analyses reinforced

infrequent and conserved hemoplasma sharing between species. First, phylogenetic patterns in

prevalence were distinct from those in genotype sharing centrality (e.g., large-colony species had

higher prevalence but lower connectivity), and prevalence accordingly did not predict centrality.

Second, we found general congruence between bat and hemoplasma phylogenies. Although this

shows codivergence is a strong evolutionary force, congruence can also stem from preferential

jumps to closely related hosts (De Vienne et al., 2013). Though we cannot rule out that some

host shifts may be artefacts of the limited resolution of both the phylogenies, our evolutionary

analyses and genotype sharing results imply that hemoplasma host shifts are possible yet rare.

By sampling a diverse assemblage of bacterial genotypes in an ecologically and

evolutionary rich host community, our work has broader implications for our understanding of

disease emergence. Many bacterial pathogens are thought to be generalists and relatively

unlikely to specialize in a novel host (Pedersen, Altizer, Poss, Cunningham, & Nunn, 2005;

Woolhouse & Gowtage-Sequeria, 2005), in contrast to many viruses in which host shifts are

more common owing to high mutation rates and short infectious periods (Geoghegan et al., 2017;

Longdon, Brockhurst, Russell, Welch, & Jiggins, 2014). Recent theoretical work suggests host

shift speciation may be less common for bacteria because of higher phenotypic plasticity (e.g.,

the ability to reside in diverse habitats) and a slower tempo of evolution (Bonneaud et al., 2019).

Obligate reliance of *Mycoplasma* spp. on host cells and more chronic infections likely explains

their propensity to specialize (Citti & Blanchard, 2013; Cohen et al., 2018). More broadly,

however, using genetics to infer pathogen sharing, rather than coarser phylogenetic scales (e.g.,

species complexes or genera), is increasingly showing that many bacterial strains may be more

host specific (Withenshaw, Devevey, Pedersen, & Fenton, 2016). The high specialism of bat

hemoplasma genotypes thus underscores the importance of using finer phylogenetic scales in the

study of infectious disease (Fountain-Jones et al., 2018; Graham, Storch, & Machac, 2018).

Comparative analyses of viruses have suggested that phylogenetically conserved

pathogen jumps between species may be a broader generality in the study of disease emergence

(Albery, Eskew, Ross, & Olival, 2019; Luis et al., 2015; Streicker et al., 2010). With few

exceptions, our results on hemoplasma genotype sharing between Neotropical bat species are

generally consistent with this pattern for a bacterial pathogen. Two cases in which hemoplasmas

were shared between more distantly related species included the VBG1 genotype in *Desmodus*

*rotundus* and *Pteronotus fulvus* (Phyllostomidae and Mormoopidae) and the EF1 genotype in

*Glossophaga soricina* and *Saccopteryx bilineata* (Phyllostomidae and Emballonuridae). For the

latter, both bat species co-roost in the LAR, which suggests an ecological context for pathogen

exposure over current timescales. However, other genetic markers (e.g., *rpoB, rpoC, gyrB*)

would be necessary to infer contemporary cross-species transmission (Kämpfer & Glaeser, 2012;

Volokhov et al., 2012), as analysis of the 16s rRNA gene alone is insufficient for hemoplasma

species identification (Volokhov et al., 2012). If hemoplasmas are more likely to specialize

rather than expand their range into new and unrelated species, genotype sharing between

unrelated bats could represent more transient spillovers (Bonneaud et al., 2019). As specialized

pathogens could be more transmissible than generalists (Garamszegi, 2006), species with high

infection prevalence of specialist genotypes could be prioritized for bacterial surveillance.

In conclusion, our analysis of a diverse community of bats and their pathogen genotypes

identifies several key ecological and evolutionary factors structuring bacterial infection within

and between species and provides a starting point for contrasts with such patterns for viruses.

Similar to bat viruses, we found moderate phylogenetic signal in hemoplasma prevalence.

However, these phylogenetic patterns in prevalence were decoupled from those describing bat

species centrality in sharing hemoplasmas, such that genotype sharing was generally restricted

by bat phylogeny. These findings imply codivergence of bats and their bacterial pathogens

alongside rare and phylogenetically constrained host shifts. Future work more broadly

characterizing the ecological and evolutionary determinants of bacterial infections in diverse host

communities will improve our understanding of cross-species transmission beyond viruses and

contribute to efforts to understand the epidemiological consequences of bacterial pathogens.

**Competing interests**

We have no competing interests.

**Data accessibility**

Individual-level data are available in Dryad

(https://datadryad.org/stash/share/7qOGkejyIldAcD9FEZHNNVCc_4k0lmmiABy0EiRwZoo).

Bat species trait data are available in Table S2. Hemoplasma sequences are available in GenBank

(MH245119–MH245194 and MK353807–MK353892).

**Funding**

DJB was funded by the ARCS Foundation, American Society of Mammalogists, Odum School

of Ecology, Explorer's Club, and NSF DEB-1601052. RKP was supported by NSF DEB-

1716698, the Defense Advanced Research Projects Agency (DARPA D16AP00113 and the

DARPA PREEMPT program Cooperative Agreement D18AC00031), U.S. National Institutes of

General Medical Sciences IDeA Program (P20GM103474 and P30GM110732), and the USDA

National Institute of Food and Agriculture (Hatch project 1015891). DJB and KAS were both

supported by the American Museum of Natural History Theodore Roosevelt Memorial Fund.

NBS was supported by the American Museum of Natural History Taxonomic Mammalogy Fund.

DGS was supported by a Sir Henry Dale Fellowship, jointly funded by the Wellcome Trust and

Royal Society (102507/Z/13/Z). The views, opinions, and/or findings expressed are those of the

authors and should not be interpreted as representing the official views or policies of the

Department of Defense or the U.S. Government.

**Acknowledgements**

For assistance with field logistics, bat sampling, and permits, we thank Mark Howells, Neil

Duncan, James Herrera, Melissa Ingala, and staff of the Lamanai Field Research Center. We also

thank the many colleagues who helped net bats during 2017–2018 bat research in Belize. We

thank Seth Walk, Susan Broadway, and Jonathan Martinson for laboratory access and assistance.

Lastly, we thank two anonymous reviewers for helpful suggestions to improve this manuscript.

**Statement of authorship**

DJB, KAS, and AMB collected samples; NBS and MBF coordinated fieldwork; RKP, SA, and

DGS supported laboratory analyses; DVV and VEC conducted molecular and phylogenetic

analyses; and ADW and DGS assisted with statistical analyses. DJB analyzed data, produced

figures, and wrote the manuscript. All authors provided critical review of the manuscript.

**References**

1.  Albery, G. F., Eskew, E. A., Ross, N., & Olival, K. J. (2019). Predicting the global

mammalian viral sharing network using phylogeography. *BioRxiv*, 732255.

2.  Alcorn, K., Gerrard, J., Cochrane, T., Graham, R., Jennison, A., Irwin, P. J., & Barbosa, A.

D. (2020). First report of Candidatus Mycoplasma haemohominis infection in Australia

causing persistent fever in an animal carer. *Clinical Infectious Diseases: An Official*

*Publication of the Infectious Diseases Society of America*. doi: 10.1093/cid/ciaa089

3.  Bai, Y., Kosoy, M., Recuenco, S., Alvarez, D., Moran, D., Turmelle, A., … Rupprecht, C.

(2011). Bartonella spp. in Bats, Guatemala. *Emerging Infectious Diseases*, *17*(7), 1269–1272.

doi: 10.3201/eid1707.101867

4.  Baillie, J., Hilton-Taylor, C., & Stuart, S. N. (2004). *IUCN red list of threatened species: a*

*global species assessment*. IUCN.

5.  Balbuena, J. A., Míguez-Lozano, R., & Blasco-Costa, I. (2013). PACo: A Novel Procrustes

Application to Cophylogenetic Analysis. *PLOS ONE*, *8*(4), e61048. doi:

10.1371/journal.pone.0061048

6.  Becker, D. J., Bergner, L. M., Bentz, A. B., Orton, R. J., Altizer, S., & Streicker, D. G.

(2018). Genetic diversity, infection prevalence, and possible transmission routes of

Bartonella spp. in vampire bats. *PLOS Neglected Tropical Diseases*, *12*(9), e0006786. doi:

10.1371/journal.pntd.0006786

7.  Becker, D. J., Crowley, D. E., Washburne, A. D., & Plowright, R. K. (2019). Temporal and

spatial limitations in global surveillance for bat filoviruses and henipaviruses. *Biology*

*Letters*, *15*(12), 20190423. doi: 10.1098/rsbl.2019.0423

8.  Bell, D. C., Atkinson, J. S., & Carlson, J. W. (1999). Centrality measures for disease

transmission networks. *Social Networks*, *21*(1), 1–21. doi: 10.1016/S0378-8733(98)00010-0

9.  Blyton, M. D. J., Banks, S. C., Peakall, R., Lindenmayer, D. B., & Gordon, D. M. (2014).

Not all types of host contacts are equal when it comes to E. coli transmission. *Ecology*

*Letters*, *17*(8), 970–978. doi: 10.1111/ele.12300

10. Bonato, L., Figueiredo, M. A. P., Gonçalves, L. R., Machado, R. Z., & André, M. R. (2015).

Occurrence and molecular characterization of Bartonella spp. and hemoplasmas in

neotropical primates from Brazilian Amazon. *Comparative Immunology, Microbiology and*

*Infectious Diseases*, *42*, 15–20. doi: 10.1016/j.cimid.2015.09.001

11. Bonneaud, C., Weinert, L., & Kuijper, B. (2019). Understanding the emergence of bacterial

pathogens in novel hosts. *Phil. Trans. R. Soc. B*. doi: 10.1098/rstb.2018.0328

12. Botero-Castro, F., Tilak, M., Justy, F., Catzeflis, F., Delsuc, F., & Douzery, E. J. P. (2013).

Next-generation sequencing and phylogenetic signal of complete mitochondrial genomes for

resolving the evolutionary history of leaf-nosed bats (Phyllostomidae). *Molecular*

*Phylogenetics and Evolution*, *69*(3), 728–739. doi: 10.1016/j.ympev.2013.07.003

13. Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan.

*Journal of Statistical Software*, *80*(1), 1–28.

14. Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: a*

*practical information-theoretic approach*. Springer Science & Business Media.

15. Butts, C. T. (2008). Social network analysis with sna. *Journal of Statistical Software*, *24*(6),

1–51.

16. Chamberlain, S. A., & Szöcs, E. (2013). taxize: taxonomic search and retrieval in R.

*F1000Research*, *2*.

17. Citti, C., & Blanchard, A. (2013). Mycoplasmas and their host: emerging and re-emerging

minimal pathogens. *Trends in Microbiology*, *21*(4), 196–203. doi: 10.1016/j.tim.2013.01.003

18. Clare, E. L., Lim, B. K., Engstrom, M. D., Eger, J. L., & Hebert, P. D. (2007). DNA

barcoding of Neotropical bats: species identification and discovery within Guyana.

*Molecular Ecology Notes*, *7*(2), 184–190.

19. Cohen, C., Shemesh, M., Garrido, M., Messika, I, Einav, M., Khokhlova, I., … Hawlena, H.

(2018). Haemoplasmas in wild rodents: Routes of transmission and infection dynamics.

*Molecular Ecology*, *27*(18), 3714–3726. doi: 10.1111/mec.14826

20. Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research.

*InterJournal, Complex Systems*, *1695*(5), 1–9.

21. Dallas, T. A., Han, B. A., Nunn, C. L., Park, A. W., Stephens, P. R., & Drake, J. M. (2019).

Host traits associated with species roles in parasite sharing networks. *Oikos*, *128*(1), 23–32.

22. De Vienne, D. M., Refrégier, G., López-Villavicencio, M., Tellier, A., Hood, M. E., &

Giraud, T. (2013). Cospeciation vs host-shift speciation: methods for testing, evidence from

natural associations and relation to coevolution. *New Phytologist*, *198*(2), 347–385.

23. Di Cataldo, S., Kamani, J., Cevidanes, A., Msheliza, E. G., & Millán, J. (2020). Hemotropic

mycoplasmas in bats captured near human settlements in Nigeria. *Comparative Immunology,*

*Microbiology and Infectious Diseases*, *70*, 101448. doi: 10.1016/j.cimid.2020.101448

24. Downs, C. J., Schoenle, L. A., Han, B. A., Harrison, J. F., & Martin, L. B. (2019). Scaling of

Host Competence. *Trends in Parasitology*, *0*(0). doi: 10.1016/j.pt.2018.12.002

25. Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME

improves sensitivity and speed of chimera detection. *Bioinformatics*, *27*(16), 2194–2200. doi:

10.1093/bioinformatics/btr381

26. Evans, N. J., Bown, K., Timofte, D., Simpson, V. R., & Birtles, R. J. (2009). Fatal

Borreliosis in Bat Caused by Relapsing Fever Spirochete, United Kingdom. *Emerging*

*Infectious Diseases*, *15*(8), 1331–1333. doi: 10.3201/eid1508.090475

27. Fenton, A., Streicker, D. G., Petchey, O. L., & Pedersen, A. B. (2015). Are All Hosts Created

Equal? Partitioning Host Species Contributions to Parasite Persistence in Multihost

Communities. *The American Naturalist*, *186*(5), 610–622. doi: 10.1086/683173

28. Fenton, M. B., Bernard, E., Bouchard, S., Hollis, L., Johnston, D. S., Lausen, C. L., …

Zigouris, J. (2001). The bat fauna of Lamanai, Belize: roosts and trophic roles. *Journal of*

*Tropical Ecology*, *17*(4), 511–524.

29. Fountain-Jones, N. M., Pearse, W. D., Escobar, L. E., Alba-Casals, A., Carver, S., Davies, T.

J., … Worsley-Tonks, K. (2018). Towards an eco-phylogenetic framework for infectious

disease ecology. *Biological Reviews*, *93*(2), 950–970.

30. Frank, H. K., Mendenhall, C. D., Judson, S. D., Daily, G. C., & Hadly, E. A. (2016).

Anthropogenic impacts on Costa Rican bat parasitism are sex specific. *Ecology and*

*Evolution*, *6*(14), 4898–4909. doi: 10.1002/ece3.2245

31. Garamszegi, L. Z. (2006). The evolution of virulence and host specialization in malaria

parasites of primates. *Ecology Letters*, *9*(8), 933–940. doi: 10.1111/j.1461-

0248.2006.00936.x

32. Garamszegi, L. Z. (2014). *Modern Phylogenetic Comparative Methods and Their*

*Application in Evolutionary Biology: Concepts and Practice*. Springer.

33. Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian

Regression Models. *The American Statistician*, *73*(3), 307–309. doi:

10.1080/00031305.2018.1549100

34. Geoghegan, J. L., Duchêne, S., & Holmes, E. C. (2017). Comparative analysis estimates the

relative frequencies of co-divergence and cross-species transmission within viral families.

*PLOS Pathogens*, *13*(2), e1006215. doi: 10.1371/journal.ppat.1006215

35. Gómez, J. M., Nunn, C. L., & Verdú, M. (2013). Centrality in primate–parasite networks

reveals the potential for the transmission of emerging infectious diseases to humans.

*Proceedings of the National Academy of Sciences*, *110*(19), 7738–7741. doi:

10.1073/pnas.1220716110

36. Gonçalves, L. R., Roque, A. L. R., Matos, C. A., Fernandes, S. de J., Olmos, I. D. F.,

Machado, R. Z., & André, M. R. (2015). Diversity and molecular characterization of novel

hemoplasmas infecting wild rodents from different Brazilian biomes. *Comparative*

*Immunology, Microbiology and Infectious Diseases*, *43*, 50–56. doi:

10.1016/j.cimid.2015.10.006

37. González-Salazar, C., Martínez-Meyer, E., & López-Santiago, G. (2014). A hierarchical

classification of trophic guilds for North American birds and mammals. *Revista Mexicana de*

*Biodiversidad*, *85*(3), 931–941. doi: 10.7550/rmb.38023

38. Goto, K., Yasuda, M., Hayashimoto, N., & Ebukuro, S. (2010). Morphological and sequence

analysis of Mycoplasma sp. isolated from the oral cavity of a house musk shrew (Suncus

murinus). *The Journal of Veterinary Medical Science*, *72*(1), 109–111.

39. Graham, C. H., Storch, D., & Machac, A. (2018). Phylogenetic scale in ecology and

evolution. *Global Ecology and Biogeography*, *27*(2), 175–187. doi: 10.1111/geb.12686

40. Gunnell, G. F., & Simmons, N. B. (2012). *Evolutionary history of bats: fossils, molecules*

*and morphology*. Cambridge University Press.

41. Guy, C., Thiagavel, J., Mideo, N., & Ratcliffe, J. M. (2019). Phylogeny matters: revisiting 'a

comparison of bats and rodents as reservoirs of zoonotic viruses.' *Royal Society Open*

*Science*, *6*(2), 181182.

42. Han, B. A., Kramer, A. M., & Drake, J. M. (2016). Global Patterns of Zoonotic Disease in

Mammals. *Trends in Parasitology*, *32*(7), 565–577.

43. Han, B. A., Schmidt, J. P., Alexander, L. W., Bowden, S. E., Hayman, D. T., & Drake, J. M.

(2016). Undiscovered bat hosts of filoviruses. *PLOS Negl Trop Dis*, *10*(7), e0004815.

44. Hattori, N., Kuroda, M., Katano, H., Takuma, T., Ito, T., Arai, N., … Niki, Y. (2020).

Candidatus Mycoplasma haemohominis in Human, Japan. *Emerging Infectious Diseases*,

*26*(1), 11–19. doi: 10.3201/eid2601.190983

45. Herrera, J. P., Duncan, N., Clare, E., Fenton, M. B., & Simmons, N. (2018). Disassembly of

Fragmented Bat Communities in Orange Walk District, Belize. *Acta Chiropterologica*, *20*(1),

147–159. doi: 10.3161/15081109ACC2018.20.1.011

46. Hijmans, R. J., Williams, E., Vennes, C., & Hijmans, M. R. J. (2019). Package 'geosphere.'

*Spherical Trigonometryv*, 1–5.

47. Hochachka, W. M., & Dhondt, A. A. (2000). Density-dependent decline of host abundance

resulting from a new infectious disease. *Proceedings of the National Academy of Sciences*,

*97*(10), 5303–5306.

48. Hornok, S., Szőke, K., Meli, M. L., Sándor, A. D., Görföl, T., Estók, P., … Hofmann-

Lehmann, R. (2019). Molecular detection of vector-borne bacteria in bat ticks (Acari:

Ixodidae, Argasidae) from eight countries of the Old and New Worlds. *Parasites & Vectors*,

*12*(1), 50. doi: 10.1186/s13071-019-3303-4

49. Huang, S., Bininda-Emonds, O. R., Stephens, P. R., Gittleman, J. L., & Altizer, S. (2014).

Phylogenetically related and ecologically similar carnivores harbour similar parasite

assemblages. *Journal of Animal Ecology*, *83*(3), 671–680.

50. Huang, S., Drake, J. M., Gittleman, J. L., & Altizer, S. (2015). Parasite diversity declines

with host evolutionary distinctiveness: A global analysis of carnivores. *Evolution*, *69*(3),

621–630. doi: 10.1111/evo.12611

51. Hutchinson, M. C., Cagua, E. F., Balbuena, J. A., Stouffer, D. B., & Poisot, T. (2017). paco:

implementing Procrustean Approach to Cophylogeny in R. *Methods in Ecology and*

*Evolution*, *8*(8), 932–940.

52. Ikeda, P., Seki, M. C., Carrasco, A. O. T., Rudiak, L. V., Miranda, J. M. D., Gonçalves, S. M.

M., … others. (2017). Evidence and molecular characterization of Bartonella spp. and

hemoplasmas in neotropical bats in Brazil. *Epidemiology & Infection*, 1–15.

53. Johnson, P. T. J., Rohr, J. R., Hoverman, J. T., Kellermanns, E., Bowerman, J., & Lunde, K.

B. (2012). Living fast and dying of infection: host life history drives interspecific variation in

infection and disease risk. *Ecology Letters*, *15*(3), 235–242. doi: 10.1111/j.1461-

0248.2011.01730.x

54. Jombart, T., Kendall, M., Almagro☐Garcia, J., & Colijn, C. (2017). treespace: Statistical

exploration of landscapes of phylogenetic trees. *Molecular Ecology Resources*, *17*(6), 1385–

1392. doi: 10.1111/1755-0998.12676

55. Jones, P. L., Hämsch, F., Page, R. A., Kalko, E. K., & O'mara, M. T. (2017). Foraging and

roosting behaviour of the fringe-lipped bat, Trachops cirrhosus, on Barro Colorado Island,

Panamá. *Acta Chiropterologica*, *19*(2), 337–346.

56. Kämpfer, P., & Glaeser, S. P. (2012). Prokaryotic taxonomy in the sequencing era–the

polyphasic approach revisited. *Environmental Microbiology*, *14*(2), 291–317.

57. Kellner, A., Carver, S., Scorza, V., McKee, C. D., Lappin, M., Crooks, K. R., … Antolin, M.

F. (2018). Transmission pathways and spillover of an erythrocytic bacterial pathogen from

domestic cats to wild felids. *Ecology and Evolution*, *8*(19), 9779–9792. doi:

10.1002/ece3.4451

58. Kelly, C. D., Stoehr, A. M., Nunn, C., Smyth, K. N., & Prokop, Z. M. (2018). Sexual

dimorphism in immunity across animals: a meta-analysis. *Ecology Letters*, *21*(12), 1885–

1894. doi: 10.1111/ele.13164

59. Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D.,

… Webb, C. O. (2010). Picante: R tools for integrating phylogenies and ecology.

*Bioinformatics*, *26*(11), 1463–1464.

60. Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and*

*Evolution*, *35*(6), 1547–1549. doi: 10.1093/molbev/msy096

61. Loayza, A. P., & Loiselle, B. A. (2008). Preliminary information on the home range and

movement patterns of Sturnira lilium (Phyllostomidae) in a naturally fragmented landscape

in Bolivia. *Biotropica*, *40*(5), 630–635.

62. Longdon, B., Brockhurst, M. A., Russell, C. A., Welch, J. J., & Jiggins, F. M. (2014). The

Evolution and Genetics of Virus Host Shifts. *PLOS Pathogens*, *10*(11), e1004395. doi:

10.1371/journal.ppat.1004395

63. Luis, A. D., O'Shea, T. J., Hayman, D. T. S., Wood, J. L. N., Cunningham, A. A., Gilbert, A.

T., … Webb, C. T. (2015). Network analysis of host–virus communities in bats and rodents

reveals determinants of cross-species transmission. *Ecology Letters*, *18*(11), 1153–1162. doi:

10.1111/ele.12491

64. Maggi, R. G., Compton, S. M., Trull, C. L., Mascarelli, P. E., Mozayeni, B. R., &

Breitschwerdt, E. B. (2013). Infection with hemotropic Mycoplasma species in patients with

or without extensive arthropod or animal contact. *Journal of Clinical Microbiology*, *51*(10),

3237–3241.

65. Mascarelli, P. E., Keel, M. K., Yabsley, M., Last, L. A., Breitschwerdt, E. B., & Maggi, R.

G. (2014). Hemotropic mycoplasmas in little brown bats (Myotis lucifugus). *Parasit Vectors*,

*7*, 117.

66. McCallum, H., Barlow, N., & Hone, J. (2001). How should pathogen transmission be

modelled? *Trends in Ecology & Evolution*, *16*(6), 295–300. doi: 10.1016/S0169-

5347(01)02144-9

67. McKee, C. D., Krawczyk, A. I., Sándor, A., Görföl, T., Földvári, M., & Földvári, G. (2019).

Host phylogeny, geographic overlap, and roost sharing shape parasite communities in

European bats. *Frontiers in Ecology and Evolution*, *7*.

68. Messick, J. B. (2004). Hemotrophic mycoplasmas (hemoplasmas): a review and new insights

into pathogenic potential. *Veterinary Clinical Pathology*, *33*(1), 2–13. doi: 10.1111/j.1939-

165X.2004.tb00342.x

69. Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl□: an R package to interact with

the Open Tree of Life data. *Methods in Ecology and Evolution*. doi: 10.1111/2041-

210X.12593

70. Millán, J., Cevidanes, A., Sacristán, I., Alvarado, M., Sepúlveda, G., Ramos-Mella, C. A., &

Lisón, F. (2019). Detection and Characterization of Hemotropic Mycoplasmas in Bats in

Chile. *Journal of Wildlife Diseases*. doi: 10.7589/2018-12-290

71. Millán, J., López-Roig, M., Delicado, V., Serra-Cobo, J., & Esperón, F. (2015). Widespread

infection with hemotropic mycoplasmas in bats in Spain, including a hemoplasma closely

related to "Candidatus Mycoplasma hemohominis." *Comparative Immunology, Microbiology*

*and Infectious Diseases*, *39*, 9–12. doi: 10.1016/j.cimid.2015.01.002

72. Monteiro, L. R., & Nogueira, M. R. (2011). Evolutionary patterns and processes in the

radiation of phyllostomid bats. *BMC Evolutionary Biology*, *11*, 137. doi: 10.1186/1471-2148-

11-137

73. Mühldorfer, K. (2013). Bats and Bacterial Pathogens: A Review. *Zoonoses and Public*

*Health*, *60*(1), 93–103. doi: 10.1111/j.1863-2378.2012.01536.x

74. Museux, K., Boretti, F. S., Willi, B., Riond, B., Hoelzle, K., Hoelzle, L. E., … Hofmann-

Lehmann, R. (2009). In vivo transmission studies of "Candidatus Mycoplasma turicensis" in

the domestic cat. *Veterinary Research*, *40*(5), 1, 45–14. doi: 10.1051/vetres/2009028

75. Myhrvold, N. P., Baldridge, E., Chan, B., Sivam, D., Freeman, D. L., & Ernest, S. K. M.

(2015). An amniote life-history database to perform comparative analyses with birds,

mammals, and reptiles. *Ecology*, *96*(11), 3109–3109. doi: 10.1890/15-0846R.1

76. Olival, K. J., Hosseini, P. R., Zambrana-Torrelio, C., Ross, N., Bogich, T. L., & Daszak, P.

(2017). Host and viral traits predict zoonotic spillover from mammals. *Nature*, *546*(7660),

646–650. doi: 10.1038/nature22975

77. Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution

in R language. *Bioinformatics*, *20*(2), 289–290.

78. Pedersen, A. B., Altizer, S., Poss, M., Cunningham, A. A., & Nunn, C. L. (2005). Patterns of

host specificity and transmission among parasites of wild primates. *International Journal for*

*Parasitology*, *35*(6), 647–657.

79. Pitcher, D. G., & Nicholas, R. A. J. (2005). Mycoplasma host specificity: Fact or fiction? *The*

*Veterinary Journal*, *170*(3), 300–306. doi: 10.1016/j.tvjl.2004.08.011

80. Plowright, R. K., Parrish, C. R., McCallum, H., Hudson, P. J., Ko, A. I., Graham, A. L., &

Lloyd-Smith, J. O. (2017). Pathways to zoonotic spillover. *Nature Reviews Microbiology*,

*15*(8), 502–510.

81. Reid, F. (1997). *A Field Guide to the Mammals of Central America and Southeast Mexico*.

OUP USA.

82. Santana, S. E., Dial, T. O., Eiting, T. P., & Alfaro, M. E. (2011). Roosting Ecology and the

Evolution of Pelage Markings in Bats. *PLOS ONE*, *6*(10), e25845. doi:

10.1371/journal.pone.0025845

83. Sikes, R. S., Care, A., & Mammalogists, U. C. of the A. S. of. (2016). 2016 Guidelines of the

American Society of Mammalogists for the use of wild mammals in research and education.

*Journal of Mammalogy*, *97*(3), 663–688.

84. Soto, F., Walker, R., Sepulveda, M., Bittencourt, P., Acosta-Jamett, G., & Müller, A. (2017).

Occurrence of canine hemotropic mycoplasmas in domestic dogs from urban and rural areas

of the Valdivia Province, southern Chile. *Comparative Immunology, Microbiology and*

*Infectious Diseases*, *50*, 70–77. doi: 10.1016/j.cimid.2016.11.013

85. Streicker, D. G., Recuenco, S., Valderrama, W., Benavides, J. G., Vargas, I., Pacheco, V., …

Altizer, S. (2012). Ecological and anthropogenic drivers of rabies exposure in vampire bats:

implications for transmission and control. *Proc. R. Soc. B*, *279*(1742), 3384–3392. doi: 10.1098/rspb.2012.0538

86. Streicker, D. G., Turmelle, A. S., Vonhof, M. J., Kuzmin, I. V., McCracken, G. F., & Rupprecht, C. E. (2010). Host phylogeny constrains cross-species emergence and establishment of rabies virus in bats. *Science*, *329*(5992), 676–679.

87. Ter Hofstede, H. M., Fenton, M. B., & Whitaker, J. O. (2004). Host and host-site specificity of bat flies (Diptera: Streblidae and Nycteribiidae) on Neotropical bats (Chiroptera). *Canadian Journal of Zoology*, *82*(4), 616–626.

88. VanderWaal, K. L., Atwill, E. R., Isbell, L. A., & McCowan, B. (2014). Quantifying microbe transmission networks for wild and domestic ungulates in Kenya. *Biological Conservation*, *169*(Supplement C), 136–146. doi: 10.1016/j.biocon.2013.11.008

89. VanderWaal, K. L., & Ezenwa, V. O. (2016). Heterogeneity in pathogen transmission: mechanisms and methodology. *Functional Ecology*, *30*(10), 1606–1622.

90. Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

91. Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, *36*(3), 1–48.

92. Vieira, R. F. C., Molento, M. B., dos Santos, L. C., Moraes, W., Cubas, Z. S., Santos, A. P., … Messick, J. B. (2009). Detection of a novel hemoplasma based on 16S rRNA gene DNA in captive and free-ranging capybaras (Hydrochaeris hydrochaeris). *Veterinary Microbiology*, *139*(3–4), 410–413. doi: 10.1016/j.vetmic.2009.06.018

93. Voigt, C. C., von Helversen, O., Michener, R., & Kunz, T. H. (2001). The economics of

harem maintenance in the sac-winged bat, Saccopteryx bilineata (Emballonuridae).

*Behavioral Ecology and Sociobiology*, *50*(1), 31–36.

94. Volokhov, D. V., Becker, D. J., Bergner, L. M., Camus, M. S., Orton, R. J., Chizhikov, V. E.,

… Streicker, D. G. (2017). Novel hemotropic mycoplasmas are widespread and genetically

diverse in vampire bats. *Epidemiology & Infection*, *145*(15), 3154–3167.

95. Volokhov, Dmitriy V., Hwang, J., Chizhikov, V. E., Danaceau, H., & Gottdenker, N. L.

(2017). Prevalence, Genotype Richness, and Coinfection Patterns of Hemotropic

Mycoplasmas in Raccoons (Procyon lotor) on Environmentally Protected and Urbanized

Barrier Islands. *Applied and Environmental Microbiology*, *83*(9). doi: 10.1128/AEM.00211-

17

96. Volokhov, Dmitriy V., Kong, H., George, J., Anderson, C., & Chizhikov, V. E. (2008).

Biological enrichment of Mycoplasma agents by cocultivation with permissive cell cultures.

*Applied and Environmental Microbiology*, *74*(17), 5383–5391. doi: 10.1128/AEM.00720-08

97. Volokhov, Dmitriy V., Norris, T., Rios, C., Davidson, M. K., Messick, J. B., Gulland, F. M.,

& Chizhikov, V. E. (2011). Novel hemotrophic mycoplasma identified in naturally infected

California sea lions (Zalophus californianus). *Veterinary Microbiology*, *149*(1–2), 262–268.

doi: 10.1016/j.vetmic.2010.10.026

98. Volokhov, Dmitriy V., Simonyan, V., Davidson, M. K., & Chizhikov, V. E. (2012). RNA

polymerase beta subunit (rpoB) gene and the 16S–23S rRNA intergenic transcribed spacer

region (ITS) as complementary molecular markers in addition to the 16S rRNA gene for

phylogenetic analysis and identification of the species of the family Mycoplasmataceae.

*Molecular Phylogenetics and Evolution*, *62*(1), 515–528.

99. Walker Vergara, R., Morera Galleguillos, F., Gómez Jaramillo, M., Pereira Almosny, N. R.,

Arauna Martínez, P., Grob Behne, P., … Müller, A. (2016). Prevalence, risk factor analysis,

and hematological findings of hemoplasma infection in domestic cats from Valdivia,

Southern Chile. *Comparative Immunology, Microbiology and Infectious Diseases*, *46*, 20–26.

doi: 10.1016/j.cimid.2016.03.004

100. Washburne, A. D., Silverman, J. D., Morton, J. T., Becker, D. J., Crowley, D., Mukherjee,

S., … Plowright, R. K. (2019). Phylofactorization: a graph partitioning algorithm to identify

phylogenetic scales of ecological data. *Ecological Monographs*, *0*(0), e01353. doi:

10.1002/ecm.1353

101. Webber, Q. M. R., Fletcher, Q. E., & Willis, C. K. R. (2017). Viral Richness is Positively

Related to Group Size, but Not Mating System, in Bats. *EcoHealth*, 1–10. doi:

10.1007/s10393-017-1276-3

102. Willi, B., Boretti, F. S., Meli, M. L., Bernasconi, M. V., Casati, S., Hegglin, D., …

Hofmann-Lehmann, R. (2007). Real-time PCR investigation of potential vectors, reservoirs,

and shedding patterns of feline hemotropic mycoplasmas. *Applied and Environmental*

*Microbiology*, *73*(12), 3798–3802. doi: 10.1128/AEM.02977-06

103. Willi, B., Boretti, F. S., Tasker, S., Meli, M. L., Wengi, N., Reusch, C. E., … Hofmann-

Lehmann, R. (2007). From Haemobartonella to hemoplasma: Molecular methods provide

new insights. *Veterinary Microbiology*, *125*(3–4), 197–209. doi:

10.1016/j.vetmic.2007.06.027

104. Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., & Jetz, W.

(2014). EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals.

*Ecology*, *95*(7), 2027–2027. doi: 10.1890/13-1917.1

105. Withenshaw, S. M., Devevey, G., Pedersen, A. B., & Fenton, A. (2016). Multihost

Bartonella parasites display covert host specificity even when transmitted by generalist

vectors. *Journal of Animal Ecology*, *85*(6), 1442–1452.

106. Woolhouse, M. E. J., & Gowtage-Sequeria, S. (2005). Host Range and Emerging and

Reemerging Pathogens. *Emerging Infectious Diseases*, *11*(12), 1842–1847. doi:

10.3201/eid1112.050997

107. Woolhouse, M. E., Taylor, L. H., & Haydon, D. T. (2001). Population biology of multihost

pathogens. *Science*, *292*(5519), 1109–1112.

108. Wright, E. S., Yilmaz, L. S., & Noguera, D. R. (2012). DECIPHER, a Search-Based

Approach to Chimera Identification for 16S rRNA Sequences. *Applied and Environmental*

*Microbiology*, *78*(3), 717–725. doi: 10.1128/AEM.06516-11

    **Tables**

897    Table 1. Competing weighted phylogenetic generalized least squares models predicting

898    hemoplasma infection prevalence (logit-transformed) across the Belize bat community. Models

899    are ranked by ΔAICc with the number of coefficients ($k$), Akaike weights ($w_i$), and a likelihood

900    ratio test pseudo-$R^2$.

| Model structure | $k$ | ΔAICc | $w_i$ | $R^2$ |
|---|---|---|---|---|
| ~ log body mass | 2 | 0.00 | 0.40 | 0.24 |
| ~ maximum colony size | 2 | 1.71 | 0.17 | 0.20 |
| ~ roost type | 2 | 2.82 | 0.10 | 0.18 |
| ~ foraging strata | 3 | 2.88 | 0.09 | 0.24 |
| ~ roost flexibility | 2 | 3.56 | 0.07 | 0.16 |
| ~ percent plants in diet | 2 | 4.30 | 0.05 | 0.14 |
| ~ dietary guild | 3 | 5.46 | 0.03 | 0.17 |
| ~ log evolutionary distinctiveness | 2 | 5.50 | 0.02 | 0.11 |
| ~ log aspect ratio | 2 | 5.61 | 0.02 | 0.10 |
| ~ log annual fecundity | 2 | 6.13 | 0.02 | 0.09 |
| ~ square-root geographic range size | 2 | 6.27 | 0.02 | 0.08 |
| ~ 1 (intercept only) | 1 | 6.77 | 0.01 | 0.00 |
| ~ sample size | 2 | 8.38 | 0.01 | 0.02 |

902 Table 2. Hemoplasma genotypes identified from the Belize bat community. Genotypes are given

903 with their bat host species, representative GenBank numbers, and intra-genotype variability.

| Genotype | Host species | Representative GenBank number | Mean intra-genotype sequence variability |
|---|---|---|---|
| VBG1 | *Desmodus rotundus, Pteronotus fulvus*[***] | KY932701 | 99.8 |
| VBG2 | *Desmodus rotundus* | KY932678 | 99.9 |
| VBG3 | *Desmodus rotundus* | KY932722 | 99.6 |
| CS1[**] | *Carollia sowelli* | MK353833 | 100 |
| CS2[**] | *Carollia sowelli, C. perspicillata* | MH245134 | 99.7 |
| MR1[**] | *Molossus nigricans* | MH245174 | 99.7 |
| MR2[**] | *Molossus nigricans* | MH245151 | NA[*] |
| PPM[**] | *Pteronotus mesoamericanus, P. fulvus*[***] | MH245159 | 99.9 |
| EF1[**] | *Eptesicus furinalis, Saccopteryx bilineata*[***], *Glossophaga soricina*[***] | MH245147 | 99.6 |
| EF2[**] | *Eptesicus furinalis* | MH245131 | 99.9 |
| NM[**] | *Natalus mexicanus*[***] | MK353818 | NA[*] |
| LE[**] | *Lophostoma evotis* | MK353892 | 99.9 |
| TC1[**] | *Trachops cirrhosus* | MH245145 | 99.8 |
| TC2[**] | *Trachops cirrhosus* | MK353860 | 99.8 |
| APH1[**] | *Dermanura phaeotis, D. watsoni, A. lituratus*[***] | MH245132 | 100 |
| APH2[**] | *Artibeus jamaicensis, A. lituratus, A. intermedius* | MH245187 | 99.9 |
| APH3[**] | *Artibeus intermedius* | MH245186 | 99.8 |
| GLS[**] | *Glossophaga soricina* | MK353874 | 99.5 |
| MYE[**] | *Myotis elegans, Myotis pilosatibialis*[***] | MK353840 | 100 |
| MYK[**] | *Myotis pilosatibialis*[***] | MH245153 | NA[*] |
| UB[**] | *Uroderma convexum* | MK353869 | 99.8 |
| PLU[**] | *Platyrrhinus helleri*[***], *Uroderma convexum*[***] | MK353883 | 99.6 |
| SP[**] | *Sturnira parvidens; A. lituratus*[***] | MH245168 | 99.4 |
| RHN[**] | *Rhynchonycteris naso*[***] | MK353871 | NA[*] |
| *M. moatsii*-like 1[****] | *Pteronotus mesoamericanus*[***] | MK353864 | NA[*] |
| *M. moatsii*-like 2[****] | *Myotis pilosatibialis*[***] | MK353862 | NA[*] |
| *M. moatsii*-like 3[****] | *Rhynchonycteris naso*[***] | MH245146 | NA[*] |
| *M. lagogenitalium*-like[****] | *Glossophaga soricina*[***] | MH245140 | NA[*] |
| *M. muris*-like[****] | *Saccopteryx bilineata*[***] | MH245138 | NA[*] |

*Intra-genotype sequence variability could not be assessed, as only one sequence was identified

**Novel hemoplasma genotypes

***Genotypes were detected in only one individual of these bat species

****Non-hemoplasma *Mycoplasma* genotypes

**Figures and legends**

Figure 1. Study sites in northern Belize. The shaded inset shows the location of Orange Walk

District. Borders show the boundaries of the LAR (Lamanai Archaeological Reserve) and KK

(Ka'Kabish). White and tan shading indicates agricultural and urban development, while dark

green shading represents intact forest. Satellite imagery was derived from Google Maps. Stacked

bar plots display the relative abundance of each sampled bat family per study site.

Figure 2. Predictors of individual bat hemoplasma infection status. (A) Odds ratios and 95%

HDIs from the most parsimonious phylogenetic GLMM (Table S4). Estimates that do not

overlap with 1 (dashed line) are displayed in black. Reference levels for the odds ratios include

bats sampled at LAR, females, reproductive bats, absence of ectoparasites, and bats sampled in

2017. (B) Infection prevalence and 95% confidence intervals (Wilson method) stratified by sex.

Results are shown for the full dataset and after randomly subsampling *Desmodus rotundus*.

Figure 3. Predictors of species-level hemoplasma prevalence across the Belize bat community.

(A) Clades with significantly different prevalence are highlighted. (B) Results from the top

PGLS models predicting prevalence as a function of mass and colony size. Model fit and 95%

confidence intervals are shown overlaid with data scaled by sample size; species from the clade

identified through phylogenetic factorization are colored as in panel A. Species identified by

phylogenetic factorization (*Saccopteryx bilineata* and *Rhynchonycteris naso*) and with larger

body mass, colony size, and hemoplasma prevalence (*Desmodus rotundus*, *Molossus nigricans*,

and *Pteronotus mesoamericanus*) are shown to the right (photographs by Brock Fenton).

Figure 4. Evolutionary relationships between Belize bats and hemoplasma genotypes. The

cophylogeny plot shows the bat phylogeny on the left and the hemoplasma genotype phylogeny

on the right. We used the *treespace* package to collapse our complete hemoplasma phylogeny

(Fig. S3) to only the 29 bacterial genotypes (Jombart, Kendall, Almagro☐Garcia, & Colijn,

2017). Lines display bat–hemoplasma associations and are shaded by the inverse of the squared

residuals from PACo (i.e., dark lines show small residuals more indicative of coevolution).

Figure 5. Patterns of hemoplasma genotype sharing across the Belize bat community. (A) Nodes

in the genotype network represent bat species (abbreviated by Latin binomials), and edges

represent a shared genotype. Nodes are colored by communities identified with the Louvain

method and are scaled by the number of individuals per species. (B) The matrix shows pairwise

hemoplasma genotype sharing, colored by the number of genotypes shared between bat species.

Figure 6. Phylogenetic patterns in hemoplasma genotype networks for Belize bat species (A)

degree and (B) eigenvector centrality. Clades showing significantly different centrality metrics

are highlighted, and points are scaled by observed values. Results from the top PGLS models

predicting both centrality metrics as a function of bat species traits (C–D). Model fit and 95%

confidence intervals are shown overlaid with data scaled by sample size; species from the clades

identified through phylogenetic factorization are colored as in panels A–B.

relative abundance of sampled bat families per site

(A) odds of hemoplasma infection

(B) bat sex

(A) prevalence

○ 0.00
○ 0.25
○ 0.50
○ 0.75
○ 1.00

A. lituratus
A. intermedius
A. jamaicensis
D. phaeotis
D. watsoni
U. convexum
P. helleri
S. parvidens
C. perspicillata
C. sowelli
G. keenani
L. evotis
M. cozumelae
T. cirrhosus
C. auritus
G. soricina
D. rotundus
M. schmidtorum
P. fulvus
P. mesoamericanus
M. megalophylla
N. leporinus
S. bilineata
R. naso
B. dubiaquercus
R. aeneus
E. furinalis
L. ega
M. elegans
M. pilosatibialis
M. nigricans
E. nanus
N. mexicanus

(B)

hemoplasma prevalence

body mass

max colony size

small/medium    large

(A) degree

(B) eigenvector centrality

(C) percent plants in diet

(D) percent plants in diet