Alteration of the premature tRNA landscape by gammaherpesvirus infection

Jessica Tucker[1], Aaron M. Schaller[2], Ian Willis[3], and Britt A. Glaunsinger[1,2,4*]

[1]Department of Plant and Microbial Biology, University of California Berkeley, CA, USA
[2]Department of Molecular and Cell Biology, University of California Berkeley, CA, USA
[3]Departments of Biochemistry and Systems and Computational Biology, Albert Einstein
College of Medicine, Bronx NY, USA
[4]Howard Hughes Medical Institute, University of California Berkeley, CA, USA
*Address correspondence to: Britt Glaunsinger, glaunsinger@berkeley.edu
9
10

Abstract

Transfer RNAs (tRNAs) are transcribed by RNA polymerase III (RNAPIII) and play a

central role in decoding our genome. Many DNA tumor viruses enhance the activity of RNAPIII,

yet whether infection alters tRNA expression is largely unknown. Here, we present the first

genome wide analysis of how viral infection alters the tRNAome. Using a tRNA-specific

sequencing method (DM-tRNA-seq), we find that the murine gammaherpesvirus MHV68

induces global changes in pre-tRNA expression, indicating that differential tRNA gene induction

is a characteristic of DNA virus infection. Elevated pre-tRNA expression corresponds to

increased RNAPIII occupancy, but post-transcriptional mechanisms also contribute. Pre-tRNA

accumulation, but not RNAPIII recruitment, requires the virally encoded mRNA endonuclease

muSOX, suggesting interplay between the different facets of virus-induced gene regulation.

Collectively, these findings reveal pervasive changes to tRNA expression during DNA virus

infection and highlight the potential of using viruses to explore tRNA biology.

Impact

The first genome-wide analysis of virus-induced tRNA expression changes reveals that

tRNA transcription and processing are perturbed by the gammaherpesvirus MHV68.

Introduction

The elegant design of the tRNA decoder, an adaptor molecule which converts genetic information into protein, was proposed over 60 years ago (1). Although tRNAs were the first non-coding RNA described, our understanding of tRNA biology, especially their functions outside of protein translation and what dictates their expression, remains limited. This is partly due to the difficulties in applying RNA sequencing and analysis platforms to tRNAs, but recent methodological advances have resulted in a surge of new studies that help to resolve these complexities (2-4). Non-canonical functions of tRNA transcripts have been described, namely the discovery that both premature and mature tRNAs undergo fragmentation into shorter RNAs with numerous regulatory functions in response to stress, cancer, and viral infection (reviewed in (5, 6)). Given the evidence that the ~400 predicted tRNA genes in mammalian genomes are differentially expressed across cell lines (7-9) and that viral infection can lead to post-transcriptional modulation of tRNAs (10, 11), defining the tRNAome under varying conditions will be central to our understanding of tRNA function, canonical and otherwise.

RNA polymerase III (RNAPIII) transcribes tRNA genes and exhibits enhanced activity during infection with DNA viruses (12-17), many of which encode their own RNAPIII genes. Though RNAPIII activation stimulates expression of viral RNAPIII genes, concomitant accumulation of host RNAPIII transcripts can trigger antiviral immune responses. For example, the induction or misprocessing of RNAPIII-generated host 5S RNA pseudogene and vault RNA transcripts during Herpes Simplex Virus-1 (HSV-1) and Kaposi's sarcoma-associated herpesvirus (KSHV) infection, respectively, activates RNA sensing pattern recognition receptors (18, 19). The RNAPIII-driven B2 SINE family of retrotransposons are also induced and sensed during infection with the murine gammaherpesvirus MHV68, although MHV68 has uniquely evolved to hijack components of the antiviral NF-κB signaling pathway to instead promote viral gene expression and replication (20-24). While it is clear that induced RNAPIII transcripts can have functional consequences during infection, little is known about the mechanisms underlying

the selectivity of RNAPIII activation, including identification of the specific RNAPIII genes

induced by viral infection on a genome-wide scale.

DNA viruses upregulate RNAPIII transcription by increasing the abundance of limiting

RNAPIII transcription factor complexes specific for certain RNAPIII promoter types (reviewed in

(25)). RNAPIII genes have one of three different promoters that recruit a unique combination of

the transcription factor complexes, TFIIIA (Gtf3a), TFIIIB (TBP, Brf1 or Brf2, and Bdp1), TFIIIC

(Gtf3c1-6) or SNAPc (Snapc1-5) plus RNAPIII for gene expression (26). tRNAs use the

abundant RNAPIII promoter type II, which consists of internal A and B boxes that bind TFIIIC

and ultimately recruit TFIIIB and RNAPIII to initiate transcription. 7SL, short interspersed nuclear

elements (SINEs), vault RNAs, and viral RNAPIII genes also use type II promoters. 5S rRNA

genes have an internal type I promoter, which binds TFIIIA (Gtf3a) and requires TFIIIC, TFIIIB,

and RNAPIII for transcription. In contrast, type III promoters found in U6 and 7SK genes have

external promoters that are TFIIIC-independent and bind the SNAPc complex and a unique

TFIIIB complex containing Brf2 protein, rather than Brf1. Defining virally-induced RNAPIII

transcript classes and/or transcription factor complexes has been essential for understanding

how viruses enhance RNAPIII activity.

The breadth of tRNA function in cells and the fact that many DNA viruses increase

RNAPIII activity suggest that regulation of tRNA expression may be a significant point of control

during infection. The repertoire of differentially expressed tRNA genes during DNA virus

infection is not known, although individual tRNA transcripts are upregulated during EBV and

SV40 infection (12, 27). Here, we present the first genome-wide analysis of how infection alters

tRNA abundance using the model gammaherpesvirus MHV68. Using a tRNA-specific

sequencing method (DM-tRNA-seq) and analysis pipeline, we found that lytic MHV68 infection

leads to a $\geq$3-fold increase in pre-tRNA levels for approximately 14% of tRNA genes. Notably,

infection-induced pre-tRNAs do not appear to undergo canonical maturation, as mature tRNA

levels remain largely constant. The accumulation of pre-tRNAs occurs through a combination of

transcriptional induction, as evidenced by increased RNAPIII occupancy at activated loci, and

post-transcriptional mechanisms. In particular, pre-tRNA accumulation is linked to the activity of

the viral endonuclease muSOX/ORF37, which cleaves host mRNAs and downregulates protein

synthesis during infection in a process called host shutoff. Given that muSOX is not required for

RNAPIII recruitment to tRNA genes, we instead hypothesize that degradation of host mRNAs

depletes factors involved in tRNA processing and turnover, contributing to the accumulation of

pre-tRNAs. The alterations in tRNA expression during MHV68 infection described here highlight

the potential of using viruses to explore mechanistic details of tRNA maturation and turnover.

Results

Pre-tRNAs increase in abundance in MHV68-infected cells

Given that MHV68 infection upregulates RNAPIII-driven B2 SINEs (20), which are

evolutionarily related to tRNAs, we hypothesized that gammaherpesviral infection may also alter

cellular tRNA expression. We first measured pre-tRNA levels by RT-qPCR, using primer sets

that were specific for premature tRNAs (pre-tRNAs) produced from intron-containing tRNA-Tyr

and tRNA-Leu genes (Figure 1A). During MHV68 infection of NIH3T3 mouse fibroblasts, pre-

tRNA levels increased by greater than 15-fold (Figure 1A). This induction appeared specific to

tRNAs, as other RNAPIII transcripts (*7SK*, *U6*, *5S*, and *7SL*) did not show differential expression

during infection. This is in contrast to Epstein-Barr virus, a gammaherpesvirus that is associated

with increased expression of many RNAPIII gene classes due to upregulation of TFIIIB/TFIIIC

transcription factor complex subunits at both the mRNA and protein level (12). Accordingly, we

did not detect changes in TFIIIB/C subunit mRNA levels upon MHV68 infection (Supplemental

Figure 1), and in fact these transcripts were less abundant during infection. This is in line with

the general downregulation of host mRNA transcripts that occurs during MHV68-induced host

shutoff (28). A time course of MHV68 infection showed that pre-tRNA-Tyr and pre-tRNA-Leu

levels were increased by 12 hours post infection (hpi), with their levels accumulating through

late stages of infection at 24 hpi (Figure 1B). Notably, the timing of pre-tRNA induction

corresponded well with the induction pattern for B2 SINEs. Finally, we also confirmed that pre-

tRNA-Tyr and pre-tRNA-Leu are upregulated by northern blot, a method that does not rely on

cDNA production (Figure 1C). Mature tRNA-Tyr transcript levels remained constant during

infection, suggesting that infection-induced pre-tRNAs may not be processed into mature

tRNAs.

Given the kinetics of induction, we hypothesized that events occurring mid to late

infection were responsible for pre-tRNA accumulation, as opposed to a cellular response to viral

binding and entry. In agreement with this hypothesis, infection of MC57G fibroblasts with UV-

inactivated MHV68 did not lead to pre-tRNA-Tyr induction (Figure 1D). Additionally, wild type

MHV68 infection of the MC57G fibroblast cell line led to an induction similar to NIH3T3s. UV-

inactivation prevented post-entry viral gene expression, as demonstrated by the absence of

expression of the immediate early viral gene ORF50. To narrow down the gene class

responsible for the kinetics of tRNA induction, we performed infections in the presence of

phosphonoacetic acid (PAA), which blocks viral DNA replication and prevents the expression of

viral late genes, such as gB. Although late gene expression was inhibited, pre-tRNA-Tyr

accumulated to similar levels in cells infected in the presence of PAA (Figure 1E). Thus, one or

more effects of early viral gene expression underlie pre-tRNA accumulation during MHV68

infection. Together, these data demonstrate that early viral gene expression drives the

increased abundance of host pre-tRNAs.

Dm-tRNA-seq reveals upregulation of premature tRNAs during infection

Extensive base modification of tRNAs can prevent reverse transcription of full-length

cDNA products. Although there are some modifications present in pre-tRNAs, they likely have

not undergone extensive modification and therefore are more efficiently amplified relative to

mature tRNAs during reverse transcription (4, 29). Amplification of full-length cDNA products

 from tRNAs can be improved by using the highly processive thermostable group II-intron

Figure 1

Figure 1. MHV68 infection results in increased pre-tRNA expression.
(A) Pre-tRNA-Tyr-GTA-1-3 and pre-tRNA-Tyr-Leu-CAA-1-1 are transcribed with flanking and intron sequences (depicted in red), which are removed during maturation. Pre-tRNAs were detected with forward or reverse primers with 3' ends complementary to the intron by AMV RT-qPCR. Relative abundance of pre-tRNAs and other RNAPIII transcripts was measured by RT-qPCR in mock and MHV68-infected NIH3T3s. Expression was normalized to 18S rRNA and compared to values from mock-infected cells. (B) RNA was extracted from NIH3T3 cells following a time course of MHV68 infection and used for both AMV RT-qPCR to detect pre-tRNAs and primer extension to detect B2 SINE and 7SK transcripts. (C) Northern blot was performed on RNA from mock or MHV68-infected MC57Gs. Transcripts produced from tRNA-Leu-CAA-1-1 were detected with intron and exon-specific probes. tRNA-Tyr transcripts were detected with a probe that anneals with a conserved 5' region in the tRNA-Tyr family. Probe binding is depicted on the right. (E) RNA from mock and MHV68-infected MC57Gs +/- UV-inactivation was used for AMV RT-qPCR. ORF50 is an early viral gene. (F) Mock and MHV68-infected MC57Gs were treated with 200 ng/l phosphonoacetic acid (PAA) to block viral DNA replication, and extracted RNA was used for RT-qPCR analysis. gB is a late viral gene whose expression is dependent on successful viral DNA replication. All infections were done at an MOI=5 for 24 hours unless otherwise noted. RT-qPCR experiments were done in triplicate at a minimum. Error bars show SD and statistics were calculated using an unpaired t- test. *P<0.05, **P<0.01. ***P<0.001.

containing reverse transcriptase (TGIRT) (30) and by partial removal of tRNA strong-stop

modifications by treating with AlkB demethylases (2, 3). To define the extent of tRNA

upregulation genome-wide, we therefore generated sequencing libraries using TGIRT and AlkB-

treated small RNA (a method called DM-tRNA-seq (3)) in mock versus MHV68 infected MC57G

cells. Our analysis pipeline was designed to distinguish reads from pre-tRNAs and mature

tRNAs, as our preliminary data with tRNA-Leu and tRNA-Tyr suggested that MHV68 may

elevate pre-tRNA but not mature tRNA levels (Figure 1C). Reads were first aligned to mature

tRNA sequences, which contained predicted tRNA genes from tRNAscan-SE (31, 32) modified

by intron removal and the addition of mature –CCA 3' ends, yielding counts for mature tRNAs

(Figure 2A). The remaining reads were aligned to a masked genome appended with pre-tRNA

sequences including intact introns and 50 nt of upstream and downstream flanking sequence.

Thus, classification as a pre-tRNA required the absence of a 3' –CCA and the presence of 5'

leader, intron, or 3' trailer sequence.

tRNAscan-SE yields a list of genes containing both true tRNAs and tRNA-like sequences

such as pseudo-tRNAs and B2 SINE elements. Among these, we focused on the high

confidence set of tRNA genes as defined by the genomic tRNA database (n=405) (31, 32). Due

to unique genomic sequences found in 5' leader and 3' trailers, pre-tRNA reads were less likely

to multimap to the genome and were used as a proxy for expression of the respective tRNA

genes. Of the 405 predicted high confidence tRNA genes, we were able to map pre-tRNAs

originating from 313 tRNA genes using Salmon (Supplemental File 1) (33). Additionally, we

used a different mapping strategy analyzing only the uniquely mapping reads in order to more

confidently identify the locus of origin for expressed pre-tRNAs, although this strategy may

underestimate the total number of expressed pre-tRNAs. Restricting the data to uniquely

mapping reads yielded a total of 244 tRNA genes with detectable pre-tRNAs from infected cells.

Figure 2

Figure 2. DM-tRNA-seq reveals differential expression of the pretRNAome upon infection.
(A) Reads from DM-tRNA-seq (MC57G +/- 24 h MHV68 infection) were aligned to mature tRNA sequences, generating the mature tRNAome. Other small RNAs and pre-tRNAs were mapped by aligning the remaining reads over the masked mm10 reference genome with appended pre-tRNA sequences. All tRNA sequences were predicted by tRNAscanSE. (B) Differential gene expression analysis from DM-tRNA-seq was plotted as fold change (FC) versus the false discovery rate (FDR). Pre-tRNAs (high confidence in yellow, other in blue), mature tRNAs (high confidence only in red), and other small RNA reads (gray) are annotated and the dotted lines represent the boundaries of FC=2 and FDR=0.05. High confidence refers to the tRNA gene set predicted by GtRNAdb. (C) Raw read counts for mature tRNAs, pre-tRNAs, or high confidence pre-tRNAs from DM-tRNA-seq of MHV68-infected MC57Gs were compared to those of uninfected cells. Error bars show SD and statistics were calculated using an unpaired t- test on raw read counts from four replicates. (D) A pie chart shows the percentage of high confidence pre-tRNAs detected by DM-tRNA-seq that were significantly upregulated (dark orange, FC>3 and FDR<0.05) versus upregulated but not significant (light orange, FC>2 and FDR>0.05) versus unregulated (gray, FC<2). (E) A heat map of log2FC changes of pre-tRNA levels by DM-tRNA-seq by isotype. (F) RNA from mock or MHV68-infected MC57Gs was used for northern blot using probes specific for either the 3' trailer or gene body of tRNA-Leu-TAA-3-1. (G) RNA from mock or MHV68-infected MC57Gs was used for northern blot using probes specific for the gene body of the tRNA-Phe-GAA, tRNA-Cys-GCA, and tRNA-Lys-TTT families. Probes detected tRNAs from multiple loci, and those for tRNA-Phe-GAA and tRNA-Cys-GCA also detected pre-tRNAs induced during infection. All infections were done in MC57G fibroblasts at an MOI=5 for 24 hours.

Differential gene expression analysis revealed that pre-tRNAs are more abundant during

MHV68 infection (Figure 2B, blue and orange circles). In fact, raw read count ratios alone (sum

of total reads per replicate compared to mock) displayed an increase in pre-tRNA reads present

in infected cells compared to uninfected cells (Figure 2C). It is likely that many of the highly

expressed pre-tRNA genes (Figure 2B, blue circles) that are not in the high confidence tRNA

gene list (Figure 2B, orange circles) are SINE retrotransposons, as B2 SINEs are robustly

induced during infection (20, 34). Focusing on high confidence tRNA genes, we found many of

these loci produce more pre-tRNAs during infection (Figure 2B, yellow circles; Figure 2C). Yet,

while total levels of pre-tRNAs were higher in infected cells, we observed selectivity in which

tRNA genes were upregulated. Of the high confidence pre-tRNAs detectably expressed in

uninfected MC57G cells, approximately one-fifth were significantly upregulated during MHV68

infection, depending on the mapping strategy used (Figure 2D shows results with Salmon: 43

out of 313 (13.7%), unique mappers: 63 out of 244 (25.8%), FDR<0.05). However, a general

trend of increased expression of pre-tRNAs was observed as judged by a fold change greater

than 2 (Figure 2D shows results with Salmon: 141 out of 314 (44.9%); unique mappers: 158 out

of 244 (64.8%)). In general, at least one member of each tRNA isotype (tRNAs encoding the

same amino acid) were present in the differential hits, but not all isodecoders (tRNAs with the

same anticodon) were represented (Figure 2E, Supplemental Figure 2). A triple A box motif that

is present in each RNAPIII-driven TMER locus encoded by MHV68 and in 10% of host tRNA

genes (35) was not enriched in the differential hits (Supplemental Figure 3).

Unlike pre-tRNAs, the levels of most mature tRNAs did not change in a statistically

significant manner, suggesting that the upregulated pre-tRNAs do not undergo canonical

maturation (Supplemental File 1). In fact, only two high confidence tRNA genes (tRNA-Lys-TTT-

2-2 and tRNA-Leu-TAA-3-1) showed significant differential expression at the level of the mature

tRNA transcript (Figure 2B, green circles FDR<0.05). To validate these findings, we designed

northern blot probes to the 3' trailer or gene body specific to transcripts originating from tRNA-

Leu-TAA-3-1 (Figure 2F). While pre-tRNA transcripts were elevated from the tRNA-Leu-TAA-3-1

gene in infected cells, levels of mature tRNAs did not change upon infection. We were not able

to design a probe specific for tRNA-Lys-TTT-2-2 transcripts, but using a probe that recognizes

members of the tRNA-Lys-TTT family did not show any differences in mature tRNA-Lys-TTT

expression during MHV68 infection (Figure 2G). Similarly, other tRNA families that we tested by

northern blotting, including tRNA-Phe-GAA and tRNA-Cys-GCA families, showed upregulation

at the level of the full-length pre-tRNA only (Figure 2G).

RNA Polymerase III is recruited to upregulated tRNA genes during infection

The distinct subset of tRNA genes that have higher expression at the pre-tRNA level during

MHV68 infection could be reflective of increased transcription, stability, or a combination of

both. To determine if the increase in pre-tRNA transcripts during infection results from

transcriptional upregulation, we performed chromatin immunoprecipitation followed by qPCR

(ChIP-qPCR) to measure occupancy of the RNAPIII subunit Polr3A at several loci in mock and

MHV68-infected MC57G fibroblasts. RNAPIII abundance specifically increased during MHV68

infection at each of the tested differentially expressed loci (tRNA-Tyr-GTA-1-3, tRNA-Leu-CAA-

1-1, and tRNA-Leu-TAA-3-1) but remained unchanged at a tRNA gene not differentially

expressed (tRNA-Ile-TAT-2-3) and at 7SK (Figure 3A). RNAPIII abundance at the RNAPII-

dependent GAPDH promoter was included as a negative control to show specificity of the

RNAPIII ChIP signal. These data suggest that there is an increase in RNAPIII recruitment to

select tRNA genes during MHV68 infection.

Maf1 is a well-studied negative regulator of transcription by RNAPIII whose activity is

controlled by phosphorylation and cellular localization in response to nutrients and stress

conditions (36, 37). When active, Maf1 directly targets the Brf1 subunit of TFIIIB and RNAPIII to

inhibit transcription. In addition, in proliferating and differentiated mammalian cells, Maf1

Figure 3. Polr3A is recruited to induced tRNA loci in infected cells.
(A) RNAPIII occupancy was measured in mock and MHV68-infected MC57Gs using ChIP-qPCR with antibodies against the Polr3A subunit at three tRNA genes upregulated at the pre-tRNA level by DM-tRNA-seq (tRNA-Tyr-GTA-1-3, tRNA-Leu-CAA-1-1, tRNA-Leu-TAA-3-1), a tRNA gene not associated with increased pre-tRNA levels (tRNA-Ile-TAT-2-3), 7SK, and the promoter of RNAPII-driven Gapdh. (B) RT-qPCR was performed with RNA extracted from wild type and Maf1-/- MHV68-infected bone marrow derived macrophages (BMDMs). (C) RNAPIII occupancy was measured using Polr3A ChIP-qPCR on wild type and Maf1-/- BMDMs at tRNA-GTA-1-3, a tRNA gene also upregulated during MHV68 infection, and 7SK. Results are reported in relative % input. (D) RNA from mock or MHV68-infected MC57Gs was extracted at the indicated hours post infection (hpi) and subjected to RT-qPCR. All infections were done in MC57G fibroblasts at an MOI=5 for 24 hours except where indicated. Error bars show SD and statistics were calculated using an unpaired t- test (RT-qPCR) or paired t-test on raw % input values (ChIP-qPCR). *P<0.05, **P<0.01. ***P<0.001.

chronically represses RNAPIII transcription to balance RNA synthesis with cellular demand (38,

39).

Thus, if the mechanism underlying MHV68 induction of pre-tRNAs involved antagonizing

Maf1-dependent repression, we reasoned that cells lacking Maf1 should no longer display a

difference in pre-tRNA levels between uninfected and infected cells. To test this hypothesis, we

derived primary bone marrow macrophages (BMDM) from WT and *Maf1-/-* mice and infected

them with MHV68 at an MOI=5. MHV68 infection of BMDMs resulted in ~10-fold increase in pre-

tRNA-Tyr upon MHV68 infection, demonstrating that increased pre-tRNA expression during

infection occurs in primary cells and is not unique to fibroblast cell lines (Figure 3B). MHV68-

infected WT and *Maf1-/-* BMDMs showed similar levels of pre-tRNA expression, indicating that

viral induction of pre-tRNA-Tyr levels was not dependent on *Maf1* regulation of RNAPIII. Similar

to other published reports (40), uninfected *Maf1*-/- cells showed a ~2-fold increase in pre-tRNA-

Tyr compared to WT BMDMs as well as increased RNAPIII promoter occupancy in ChIP

assays, confirming that RNAPIII activity is elevated in the absence of this negative regulator

(Figure 3B, 3C).

Even though the relative increases in pre-tRNA-Tyr levels were dramatically larger in

MHV68-infected cells compared to cells lacking Maf1 (Figure 3B,D), we were struck that the

increase in RNAPIII occupancy in these two scenarios was comparable (Figure 3A,C). To

explore the contribution of RNAPIII recruitment to pre-tRNA levels, we compared pre-tRNA

abundance by RT-qPCR (Figure 3D) and RNAPIII promoter occupancy by Polr3A ChIP-qPCR

(Figure 3A) across a time course of MHV68 infection in MC57G fibroblasts. Pre-tRNA-Tyr levels

were induced about 5-fold at 16 and 20 hpi but increased to a nearly 15-fold induction at 24 hpi,

whereas *7SK* levels remained constant (Figure 3D). In contrast to the RNA abundance data,

RNAPIII occupancy at the pre-tRNA-Tyr promoter was modestly increased by 16hpi, but did not

further increase at later time points (Figure 3A). Thus, after 16 hpi, it is likely that post-

transcriptional mechanisms underlie the continued accumulation of pre-tRNA-Tyr.

Virus-induced mRNA degradation contributes to post-transcriptional accumulation of pre-tRNAs

We reasoned that selective post-transcriptional accumulation of pre-tRNAs might occur if pre-tRNA maturation or turnover were impaired, for example if the virus down-regulated factors involved in these processes. A well-characterized phenotype of MHV68 infection is the widespread degradation of host mRNAs, which is caused by the viral mRNA-specific endonuclease muSOX (28). Notably, the levels of the RNAPIII-transcribed viral tRNAs (vtRNAs) are decreased in cells infected with the MHV68 muSOX point mutant R443I (MHV68.R443I), which has reduced host shutoff activity (41). muSOX is expressed with early kinetics, but its mRNA depletion phenotype is most prominent at late stages of infection, coincident with maximal pre-tRNA accumulation. We tested whether virus-induced mRNA decay contributes to post-transcriptional accumulation of cellular pre-tRNAs by measuring pre-tRNA abundance in MC57G cells infected with MHV68.R443I or the mutant revertant (MR) virus in which the R443I was restored to wild type (Figure 4A). Similar to the vtRNAs, the levels of cellular pre-tRNA-Tyr and pre-tRNA-Leu were lower levels in cells infected with the R443I mutant compared to MR MHV68 (Figure 4A). In contrast, the viral ORF50 mRNA was slightly increased during the R443I mutant infection, consistent with previous reports (41). To determine if these changes reflected transcriptional output, we measured RNAPIII binding at two differentially regulated host tRNA genes and the viral TMER2 locus during MR vs R443I infection (Figure 4B). RNAPIII abundance was similar during infection with MR and R443I viruses, in stark contrast to the difference in pre-tRNA and vtRNA transcript abundance. This result suggests that the decreased abundance of these transcripts in total RNA is not due to decreased levels of transcription. Collectively, these data suggest that the mRNA decay phenotype associated with WT MHV68 infection is not involved in transcriptional activation of tRNA loci, but instead stabilizes the unprocessed pre-tRNAs.

Figure 4

Figure 4. muSOX activity is required for pre-tRNA accumulation, but not for RNAPIII recruitment.
(A) MC57Gs were infected with muSOX R443I or mutant revertant (MR) MHV68 and RT-qPCR was performed with extracted RNA. (B) RNAPIII occupancy was measured using ChIP-qPCR with antibodies against the Polr3A subunit of RNAPIII at induced tRNA loci, viral TMER2, and 7SK using the cells described in (A). All infections were done at an MOI=5 for 24 hours. Error bars show SD and statistics were calculated using an unpaired t- test (RT-qPCR) or paired t-test on raw % input values (ChIP-qPCR). *P<0.05, **P<0.01. ***P<0.001.

Discussion

This study represents the first genome-wide analysis of how infection modulates the

tRNAome, revealing that the model gammaherpesvirus MHV68 causes extensive changes to

pre-tRNA but not mature tRNA abundance. We did not detect increased expression across all

tRNA genes, but instead found a dramatic accumulation of pre-tRNAs from a subset of loci,

which may reflect differences in promoter accessibility, regulation and/or pre-tRNA processing.

Collectively, our data support a model whereby pre-tRNA accumulation during MHV68 infection

is driven by both RNAPIII recruitment and RNA stabilization (Figure 5). We hypothesize that

early events in infection stimulate RNAPIII-driven transcription of specific tRNA loci, while

muSOX-induced depletion or relocalization of pre-tRNA processing machinery stalls further

canonical processing of these transcripts, leading to their build up late in infection (Figure 5).

These findings are bolstered by early reports showing enhanced RNAPIII activity during DNA

virus infection, including for select tRNAs (12-17, 27), suggesting that manipulation of the pre-

tRNA landscape is likely to be a conserved feature of numerous viruses.

Figure 5

Figure 5. MHV68 infection induces pre-tRNA accumulation by transcriptional and post-transcriptional mechanisms. RNAPIII is recruited to specific tRNA genes by a Maf1-independent mechanism, increasing the output of pre-tRNAs. Pre-tRNAs accumulate in infected cells due to depletion of maturation or turnover machinery during muSOX-mediated mRNA decay (host shutoff).

Pre-tRNA steady state levels are dependent on both the production and stability of the

pre-tRNA transcripts. We performed Polr3A ChIP-qPCR and found that, for several tRNA

genes, increased pre-tRNA abundance was associated with higher levels of Polr3A occupancy

during infection. In contrast, a tRNA gene with constant pre-tRNA expression did not show

changes in Polr3A binding. This confirms that transcriptional increases- at least at some tRNA

genes- contribute to pre-tRNA accumulation during infection. Whether this is true for all tRNA

genes associated with increased pre-tRNA levels remains to be determined. Additionally,

increased RNAPIII occupancy during infection suggests a mechanism involving RNAPIII

recruitment to specific tRNA genes, and raises the possibility that these tRNA genes are not

expressed to their full potential in uninfected cell populations. In support of this, there is

considerable overlap between tRNA genes upregulated during MHV68 infection with those

upregulated in cells lacking Maf1, a negative regulator of RNAPIII that prevents polymerase

recruitment (59 of 79 pre-tRNAs induced by MHV68 infection of MC57G fibroblasts were also

increased in *Maf1-/-* versus WT liver in (40)). However, Maf1 does not appear to play a pivotal

role in pre-tRNA accumulation during MHV68 infection.

The mechanisms employed during MHV68 infection to increase RNAPIII recruitment

have not been identified. During EBV infection, TFIIIC subunit and Bdp1 protein levels increase,

along with transcripts for RNAPIII genes with type I and II promoters (*7SL*, *5S*, tRNAs), but not

those with type III promoters (*7SK*, *U6*, *MRP*) (12). In contrast, MHV68 infection leads to

transcriptional induction of B1/B2 SINEs (20, 34) and certain tRNA genes, all of which have type

II promoters. TFIIIB/C subunit mRNA levels do not increase during MHV68 infection, and

quantitative mass spectrometry of MHV68-infected cells indicates that total TFIIIB/C protein

levels are not elevated, although there may be enrichment of these factors in the nucleus (42).

Thus, RNAPIII regulation during MHV68 infection appears distinct from that observed with EBV

and may therefore involve novel mechanisms. Like EBV, SV40 infection leads to increased

TFIIIC protein levels, but additionally results in higher availability of TFIIIB by large T-antigen

sequestration of the retinoblastoma tumor suppressor protein (RB), a negative regulator of

RNAPIII transcription (43). The changes in RNAPIII output in mouse fibroblasts are similar

between SV40 and MHV68 infections (B1/B2 SINEs and certain tRNAs are induced (27, 44)),

plus mass spectrometry data suggests that RB subcellular concentration changes during

MHV68 infection (42). Thus, future studies will explore how RB is regulated during MHV68

infection and whether this protein is an important player in gammaherpesvirus-induced RNAPIII

activation. These could include the posttranslational regulation of transcription factor activities,

altered factor availability or epigenetic changes influencing gene accessibility.

With the exception of selenocysteine tRNA (7), all tRNAs genes use the same internal

A/B box promoter architecture for transcription. However, based on differential binding of

RNAPIII, many studies have revealed that tRNA genes are not equally transcribed in

mammalian cells, as measured by the differential binding of RNAPIII (7-9). Genome-wide

studies examining chromatin accessibility, modification and transcription factor binding have

suggested that a driver of RNAPIII occupancy at tRNA genes is open and active chromatin,

typically in the vicinity of RNAPII transcription sites, yet the basis for differential RNAPIII binding

remains to be determined. (7, 8, 45, 46). Further studies are also needed to assess how host

chromatin accessibility and modifications change during MHV68 infection. Interestingly, MHV68

infection leads to depletion of RNAPII-specific subunits, resulting in less RNAPII transcription of

the host genome during infection (42). There is some evidence of transcriptional interference

between RNAPIII and RNAPII bound at nearby genes, as depletion of the Rbp1 subunit of

RNAPII leads to upregulation of a subset of tRNA genes (47). It is possible that depletion of

RNAPII allows for increased recruitment of RNAPIII machinery. We hypothesize that a

combination of genome accessibility and transcription factor availability are associated with

tRNA transcriptional increases during infection.

Although tRNA gene transcription is elevated due to RNAPIII recruitment, the magnitude

of pre-tRNA abundance is unlikely to be explained by transcriptional activation alone. Instead,

our data suggest that pre-tRNA stability is also increased. While we have not been able to

measure pre-tRNA half-life directly (due to difficulty in robustly inhibiting RNAPIII in the required

short time frame of pre-tRNA processing), it is clear that pre-tRNAs accumulate dramatically

between 20-24 hpi, a time frame in which RNAPIII occupancy is unchanged. Infection-induced

pre-tRNAs do not seem to be processed into mature tRNAs, as we found no evidence of

differential expression at the mature tRNA transcript level. Similarly, *Maf1-/-* tissues exhibit

upregulation of pre-tRNAs without significant changes to mature tRNA levels, suggestive of a

homeostatic mechanism to maintain the size of mature tRNA pools when tRNA gene

transcription is increased (40). In both of these cases, it is possible that the increase of pre-

tRNAs upon transcriptional induction saturates processing machinery, an outcome well

established in yeast (48, 49), resulting in their clearance (degradation) from cells. It was striking

that some pre-tRNA transcripts revealed in our study were much more highly expressed than

others. These pre-tRNAs might have specific modifications, unique structural characteristics, or

processing fates that render them inherently more stable in the cell or during infection. Knowing

the identity of these potentially hyper-stable pre-tRNAs will help guide studies to investigate

unique features and broaden our understanding of tRNA maturation and turnover.

We hypothesized that tRNA processing and turnover machinery might be depleted in

infected cells due to gammaherpesvirus-induced host shutoff implemented at late stages of viral

infection. Widespread mRNA decay is initiated by a gammaherpesvirus-encoded endonuclease,

called SOX in KSHV and muSOX in MHV68. SOX and its homologs cleave RNAPII-transcribed

mRNAs, leading to subsequent downregulation of host protein synthesis (28, 50). Depletion of

host proteins might further increase the ratio of pre-tRNA to available processing and turnover

machinery, leading to increased pre-tRNA half-lives. Though total protein levels of tRNA

processing machinery are not dramatically altered during a 24-hour lytic infection cycle, there

are some decreases in nuclear abundance of tRNA-related factors during infection, including

the nuclear exosome (42). We hypothesize that modest abundance changes of many tRNA-

related processing and turnover complexes could have an additive effect on the stability of pre-

tRNAs in infected cells. Moreover, pre-tRNAs were less abundant in cells infected with the host

shut-off defective R443I virus, a phenotype not explained by changes in nascent pre-tRNA

levels or Polr3A abundance at tRNA genes. Viral TMER transcripts are also less abundant

under these conditions (41), suggesting that the factors involved with pre-tRNA stability during

infection may also be involved in viral TMER transcript stability. TMERs consist of a tRNA-like

sequence with 1-2 miRNA stem loops that are processed by the host tRNA 3' processing

enzyme tRNAse Z (51, 52). Whether tRNAse Z or other host factors are involved in stabilizing

host pre-tRNA and viral TMERs is an important question for the future.

Boosting RNAPIII activity is thought to promote the expression of RNAPIII-transcribed

genes encoded by some DNA viruses (e.g., Epstein-Barr virus EBERs, Adenoviruses VA RNAs,

MHV68 TMERs) -- however, not all DNA viruses encode RNAPIII genes, suggesting that

regulation of RNAPIII could be advantageous to the virus or host for other reasons. For

example, because nascent RNAPIII transcripts contain 5' triphosphate ends, they can and do

serve as substrates for RNA sensors in the cell, such as RIG-I (18, 19). These studies show that

while typically the 5' ends of RNAPIII transcripts are processed or protected from sensing by

protein binding partners, HSV-1 and KSHV infection tips the balance and results in exposure of

5' triphosphate ends of RNAPIII transcripts for RIG-I recognition. Additionally, increased tRNA

expression could contribute to tRNA fragmentation reported during viral infection and cancer;

there are increasing examples of these fragments functioning in gene regulation, cell

proliferation, and stress responses (5, 6). As DM-tRNA-seq is not optimized for analysis of tRNA

fragments, it is not yet clear if tRNA fragments are produced during MHV68 infection. However,

some tRNA fragments are processed from pre-tRNA transcripts, including pre-tRNA-Tyr (53,

54), and thus it would not be surprising if pre-tRNA-derived fragments also increased upon

infection. In summary, exploring viral modulation of host tRNAomes could lead to exciting

insights both for tRNA biology and for the role of RNAPIII activation during DNA virus infection.

Materials and Methods

Cell and Virus Culture

NIH3T3 and MC57G cell lines were obtained from ATCC, maintained in Dulbecco's modified

Eagle medium (DMEM; Gibco) supplemented with 10% fetal bovine serum (FBS; VWR), and

screened regularly for mycoplasma by PCR. Wild-type and *Maf1* knockout primary bone

marrow-derived macrophages (BMDMs) were differentiated from adult 3-6 month old C57BL/6

mice (40). Provided femurs and tibia were ground in a sterile tissue-culture hood using a mortar

and pestle in DMEM before being passed through a 70 μm filter to remove debris. Cells were

differentiated in macrophage media + PenStrep (High glucose DMEM, 10% FBS, 10% MCSF,

1% PenStrep, 1% GlutaMax) for 7 days before being frozen for later use. After thaw, cells were

not subjected to any media containing antibiotics. MHV68 was amplified in NIH3T12 cells and

TCID50 was measured on NIH3T3s by limiting dilution. GFP-expressing wild type (55) and

GFP-expressing ΔHS (SOX R443I (56)) MHV68 viruses were incubated with cells for 1 hour at

an MOI of 5 to allow viral entry, and then the virus-containing media was removed and replaced

with fresh media. PAA was used at a concentration of 200 μg/ml and was added at the start of

the infection. UV-inactivation of MHV68 was performed by autocrosslinking twice in a

Stratalinker 2400 (1200uJX100). For infection of BMDMs, virus was added to cells in serum-free

DMEM for 4 hours in non-TC treated plates. Virus containing media was then aspirated and

replaced with macrophage media without antibiotics.

RNA isolation and analysis

Total RNA was isolated from cells using TRIzol (Invitrogen), treated with Turbo DNase (Ambion)

and reverse transcribed with AMV RT (Promega) primed with random 9-mers. For TGIRT-

qPCR, cDNA was synthesized with the TGIRT™-III Enzyme (InGex) following manufacturer's

instructions. Quantitative PCR analysis was performed with iTaq Universal SYBR Green

Supermix (Bio-Rad) using the primers listed in Table 1. qPCR is performed on at least three

biological replicates and Ct values were measured from three technical replicates per biological

sample. Fold change was calculated by ΔΔCT method. For northern blots, 20 μg of RNA was

loaded into 8% PAGE/7 M Urea gels, transferred to Hybond-XL (GE) membranes using a

Trans-blot Turbo Transfer system (Bio-Rad). Blots were pre-hybridized in ULTRAhyb buffer

(Thermo Fisher) at 42 °C for 1 hour before adding radiolabeled probe. Probes were generated

by end-labeling oligos listed in Table 1 using T4 PNK and $[\gamma\text{-}^{32}P]$-ATP. Blots were probed

overnight at 42 °C, washed 3 times for 5 minutes in 0.5x SSC and 0.1% SDS. Primer extension

was performed on 10 µg of RNA using a 5' fluorescein labeled oligo specific for B2 SINEs or

7SK. RNA was ethanol precipitated and resuspended in 9 µl of annealing buffer (10 mM Tris-

HCl, pH7.5, 0.3 M KCl, 1 mM EDTA) plus 1 µl of 5' fluorescein labeled oligo. Sample was briefly

heated to 95 °C, then incubated at 55 °C for 1 hour. 40 µl of extension buffer (10 mM Tris-HCl,

pH 8.8, 5 mM MgCl2, 5 mM DTT, 1 mM dNTP) and 1 µl of AMV RT (Promega) was added and

the sample was incubated at 42 °C for 1 hour. Nucleic acids were precipitated, resuspended in

RNA loading dye, and separated in 8% PAGE/ 7 M Urea gels.

DM-tRNA-seq library preparation

Protocol was modified from (3). His-tagged wild-type and D135S AlkB plasmids were obtained

from Addgene (#79050 and #79051) and proteins were purified by a Ni-NTA column followed by

cation exchange. Total RNA was spiked with *in vitro* transcribed *E. coli* tRNA-Lys, *E. coli* tRNA-

Tyr, and *S. cerevisiae* tRNA-Phe transcripts at 0.01 pmol IVT tRNAs per µg total RNA. RNA was

deacylated in 0.1M Tris-HCl, pH 9 at 37 °C for 45 min, ethanol purified, and then

dephosphorylated with PNK. Deacylated and dephosphorylated RNAs were then purified with a

mirVANA small RNA purification kit (Ambion) to isolate RNAs ≤ 200 nucleotides. Small RNAs

were demethylated in 300 mM NaCl, 50 mM MES pH 5, 2 mM $MgCl_2$, 50 µM ferrous ammonium

sulfate, 300 µM 2-ketoglutarate, 2 mM L-ascorbic acid, 50 µg/ml BSA, 1U/µl SUPERasin, 2X

molar ratio of wt AlkB, and 4X molar ratio of D135S AlkB for 2 hours at room temperature.

Reaction was quenched with 5 mM EDTA and purified with Trizol. 100 ng demethylated small

RNAs were used for library prep with a TGIRT Improved Modular Template-Switching RNA-seq

Kit (InGex) following the manufacturer's instructions. The included RNA/DNA heteroduplex for

the template switching reaction by TGIRT contains a 3' N-overhang on the DNA primer to

promote reverse transcription of all small RNAs (as opposed to the T-overhang for targeting

mature tRNAs described in (3)). PCR amplification was performed with Phusion polymerase

(Thermo Fisher) with Illumina multiplex and barcoded primers synthesized by IDT for 12 cycles.

Library was sequenced on a HiSeq4000. Data has been deposited at NCBI GEO, series

GSE142393.

DM-tRNA-seq bioinformatic analysis

tRNAscan-SE(57) was used to create the predicted tRNA gene library. Pre-tRNA sequences

were defined by adding 50bp of genomic sequences on 5' and 3' ends of the predicted tRNA

genes. Mature tRNAs were appended with a CCA tail and were clustered as described in (58).

Illumina 150bp paired-end sequence data were subjected to quality control using BBDuk

(https://jgi.doe.gov/data-and-tools/bbtools/). Sequencing adapters were removed from raw

reads. Bases that have quality score lower than 30 were trimmed. Any reads that are at least 50

bp in length after quality control were aligned to mature tRNA reference using end-to-end

mapping mode in Bowtie2 (59) (version 2.3.4.1). Reads that did not map to mature tRNAs were

then mapped to the masked mm10 genome appended with pre-tRNA sequences. Two

approaches were used for this mapping step. One is to use Bowtie2, with reads that map to

multiple locations excluded from downstream analysis, followed by quantification of expression

using SALMON (33). The other approach used SALMON to map the reads and quantify the

expression levels, which includes the reads that map to multiple locations in estimating the

expression levels. Raw counts were normalized using spike-in RNA species using R package

RUVSeq (60). A generalized linear model was built to test for differential expression for each

RNA species, using R package edgeR (61, 62).

Chromatin immunoprecipitation

Cells from a 10 cm dish were crosslinked in 1% formaldehyde in PBS for 5 min at room temperature, quenched in 0.125 M glycine, and washed twice with PBS. Crosslinked cells were lysed with 1 ml ChIP lysis buffer (50 mM HEPES pH 7.9, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP40, 0.25% Triton X-100) by rotation for 10 min. Nuclei were collected by centrifugation at 1700 x g for 5 min at 4°C, washed once with ChIP shearing buffer (50 mM Tris Cl, pH 7.5, 10 mM EDTA, 0.1% SDS), then resuspended in 1 ml of ChIP shearing buffer. Chromatin was sheared for 8 minutes using a Covaris S220 Focused-ultrasonicator at 140 power, 5% duty cycle, 200 bursts / cycle. Chromatin was spun at 15,000 x g for 5 min at 4°C and the pellet was discarded. 40 µg of chromatin was incubated with 10 µg rabbit polyclonal anti-POLR3A (Abcam ab96328) or rabbit IgG (Southern Biotech) overnight. 30 µl of mixed protein A and G dynabeads (Thermo Fisher) were added and tubes were rotated for 2 h at 4 °C. Beads were washed with low salt immune complex (20 mM Tris pH 8.0, 1% Triton X-100, 2 mM EDTA, 150 mM NaCl, 0.1% SDS), high salt immune complex (20 mM Tris pH 8.0, 1% Triton X-100, 2 mM EDTA, 500 mM NaCl, 0.1% SDS), lithium chloride immune complex (10 mM Tris pH 8.0, 0.25 M LiCl, 1% NP-40, 1% Deoxycholic acid, 1 mM EDTA), and Tris-EDTA for 5 min each at 4°C with rotation. DNA was eluted from the beads using 100 µl of elution buffer (150 mM NaCl, 50 µg/ml proteinase K) and incubated at 50°C for 2 hr, then 65°C overnight. DNA was purified using a Zymo Oligo Clean and Concentrator kit and used for qPCR analysis using the primers listed in Table 1.

Replicates and statistical analysis

This work contains cell culture-based assays, where biological replicates are defined as experiments performed on at least three distinct samples (cells maintained in different flasks). Technical replicates are defined as experiments performed on the same biological sample at

494 least three times. The datapoint for each biological replicate performed is depicted in the

495 Figures. Statistical analysis was performed using Prism 7 (Version 7.0c) software (GraphPad

496 Software), and the exact test performed is described in the Figure legends. The criteria for

497 including data involved the quality of the data for positive and negative controls.

500 **Table 1. Primers used in this study.**

Primer extension oligos (mouse)	
B2_PXT	TACACTGTAGCTGTCTTCAGACA
7SK_PXT	GAGCTTGTTTGGAGGTTCT
RT-qPCR primers (mouse)	
Mm_pre-tRNA-Tyr-GTA-1-3_F	CCTTCGATAGCTCAGTTGGTAGAGC
Mm_pre-tRNA-Tyr-GTA-1-3_R	CTTCGAGCCGGATTCGAACCAGCGAC
Mm_pre-tRNA-Leu-CAA-1-1_F	CGCCAGACTCAAGCTATCGC
Mm_pre-tRNA-Leu-CAA-1-1_R	TGTCAGAAGTGGGATTCGAACC
7SK_F	CCCCTGCTAGAACCTCCAAAC
7SK_R	CACATGCAGCGCCTCATTT
U6_F	CGCTTCGGCAGCACATATAC
U6_R	AAAATATGGAACGCTTCACGA
5S_F	TCTCGTCTGATCTCGGAAGC
5S_R	AGCCTACAGCACCCGGTATT
7SL_F	ATCGGGTGTCCGCACTAAGTT
7SL_R	CAGCACGGGAGTTTTGACCT
GTF3C1_F	TTCACACACCGCTTTCAAG
GTF3C1_R	CTCCAGTAGGTAATCGGATCACA

GTF3C2_F	ACAGCAAGAGGCGCTAAAAGG
GTF3C2_R	GGACTGATCTAAGAGACCAGGAA
GTF3C3_F	ATGGGGTGAGTAAGTCAGTTCCA
GTF3C3_R	TCCAACACAAATACATCGCCTG
GTF3C4_F	CCCAATGCGGGGTTTCAAATA
GTF3C4_R	CTGCACAGTCAGGCGATTGT
GTF3C5_F	TTGGTGTGCGTGGAGTACC
GTF3C5_R	CCCAGGGTCTGTAGCATCT
GTF3C6_F	GCGTCTCCGAACGCTATGG
GTF3C6_R	CCTCTCCGTGTCAATTCCCA
BDP1_F	TGCTTCTTCTGAGATAGGGCA
BDP1_R	GAGGGCTTCTTTAGCAGCACT
RT-qPCR primers (MHV68)	
MHV68_orf50_F	GGCCGCAGACATTTAATGAC
MHV68_orf50_R	GCCTCAACTTCTCTGGATATGCC
MHV68_gB_F	GGCCCAAATTCAATTTGCCT
MHV68_gB_R	CCCTGGACAACTCCTCAAGC
MHV68_vtRNA2_F	GGTAGAGCAGCGGTTCCT
MHV68_vtRNA2_R	ACTCCCCCTCTCAACCA
MHV68_vtRNA4_F	AGCGGCAGGCTCATC
MHV68_vtRNA4_R	ATCTCAACTCTGCGTCGG
ChIP-qPCR primers (mouse)	
Mm_tRNA-Tyr-GTA-1-3_F	CCTTCGATAGCTCAGTTGGTAGAGC
Mm_tRNA-Tyr-GTA-1-3_R	CTTCGAGCCGGATTCGAACCAGCGAC
Mm_tRNA-Leu-CAA-1-1_F	CGCCAGACTCAAGCTATCGC

Mm_tRNA-Leu-CAA-1-1_R	TGTCAGAAGTGGGATTCGAACC
Mm_tRNA-Leu-TAA-3-1_F	ACTGGGATGGCTGAGTGGTTAAGGC
Mm_tRNA-Leu-TAA-3-1_R	GCAGACAGTCGAGCAGTACA
Mm_tRNA-Ile-TAT-2-3_F	GCTCCAGTGGCGCAATCGGTTAGC
Mm_tRNA-Ile-TAT-2-3_R	GCCTAGCTCCAAAAGGATCG
7SK_F	CCCCTGCTAGAACCTCCAAAC
7SK_R	CACATGCAGCGCCTCATTT
GAPDH_promoter_F	GGTTATCACCAGGCCAGCTA
GAPDH_promoter_R	GGAAGGGAGAAAAGGCATTC
ChIP-qPCR primers (MHV68)	
MHV68_vtRNA2_F (TMER2)	GGTAGAGCAGCGGTTCCT
MHV68_vtRNA2_R (TMER2)	ACTCCCCCTCTCAACCA

Supplemental File 1. Differential gene expression analysis output from DM-tRNA-seq

comparing MHV68-infected vs mock-infected MC57G fibroblasts.

Acknowledgements

We thank Jie Li and the UC Davis Bioinformatics Core for valuable assistance with the

bioinformatics analysis and Ella Hartenian for providing samples. We thank members of the

Glaunsinger lab and Coscoy lab for discussion and comments on the manuscript. This research

was supported by an American Cancer Society Postdoctoral Award 131370-PF-17-245-01-MPC

to JMT, and National Institutes of Health grants CA136367 and AI147183 to BG and GM120358

to IW. BG is an investigator of the Howard Hughes Medical Institute.

Competing Interests

None

Supplemental Figure 1

Supplemental Figure 1. The levels of mRNAs encoding TFIIIB and TFIIIC subunits are decreased upon MHV68 infection. RT-qPCR was performed on MC57Gs infected at an MOI=5 for 24 h and were done in triplicate. Error bars show SD and statistics were calculated using an unpaired t- test on raw ΔCt values. *P<0.05.

Supplemental Figure 2

Supplemental Figure 2. A variety of tRNA isotypes are upregulated at the pre-tRNA level in response to MHV68 infection. Pre-tRNAs detected by DM-tRNA-seq from each tRNA isodecoder are shown in black, while the number with increased pre-tRNA expression in response to MHV68 are shown in are pink. Most isotypes (tRNAs encoding the same amino acid), but not all isodecoders (tRNAs with same anticodon), are represented.

Supplemental Figure 3

All high confidence tRNA genes

10.54% triple A motif
89.46% no triple A motif

Total=408

tRNA genes induced by MHV68

12.66% triple A motif
87.34% no triple A motif

Total=79

Supplemental Figure 3. A triple A-box motif is not enriched in MHV68-induced tRNA genes.
The triple A-box motif (TRGYNNARNTGGTRGARNAGNNG) identified in MHV68 TMERs is present
in ~11% of mouse high confidence tRNA genes as previously reported (35, 63). Both uniquely
mapped and Salmon-identified MHV68-induced tRNA genes (n=79 genes) were analyzed for the
triple A-box motif.

52R   References

1.   **Hoagland MB, Stephenson ML, Scott JF, Hecht LI, Zamecnik PC.** 1958. A soluble ribonucleic acid intermediate in protein synthesis. J Biol Chem **231:**241-257.

2.   **Cozen AE, Quartley E, Holmes AD, Hrabeta-Robinson E, Phizicky EM, Lowe TM.** 2015. ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. Nat Methods **12:**879-884.

3.   **Zheng G, Qin Y, Clark WC, Dai Q, Yi C, He C, Lambowitz AM, Pan T.** 2015. Efficient and quantitative high-throughput tRNA sequencing. Nat Methods **12:**835-837.

4.   **Gogakos T, Brown M, Garzia A, Meyer C, Hafner M, Tuschl T.** 2017. Characterizing Expression and Processing of Precursor and Mature Human tRNAs by Hydro-tRNAseq and PAR-CLIP. Cell Rep **20:**1463-1475.

5.   **Oberbauer V, Schaefer MR.** 2018. tRNA-Derived Small RNAs: Biogenesis, Modification, Function and Potential Impact on Human Disease Development. Genes (Basel) **9**.

6.   **Li S, Xu Z, Sheng J.** 2018. tRNA-Derived Small RNA: A Novel Regulatory Small Non-Coding RNA. Genes (Basel) **9**.

7.   **Oler AJ, Alla RK, Roberts DN, Wong A, Hollenhorst PC, Chandler KJ, Cassiday PA, Nelson CA, Hagedorn CH, Graves BJ, Cairns BR.** 2010. Human RNA polymerase III transcriptomes and relationships to Pol II promoter chromatin and enhancer-binding factors. Nat Struct Mol Biol **17:**620-628.

8.   **Raha D, Wang Z, Moqtaderi Z, Wu L, Zhong G, Gerstein M, Struhl K, Snyder M.** 2010. Close association of RNA polymerase II and many transcription factors with Pol III genes. Proc Natl Acad Sci U S A **107:**3639-3644.

9.   **Canella D, Praz V, Reina JH, Cousin P, Hernandez N.** 2010. Defining the RNA polymerase III transcriptome: Genome-wide localization of the RNA polymerase III transcription machinery in human cells. Genome Res **20:**710-721.

10.   **Li M, Kao E, Gao X, Sandig H, Limmer K, Pavon-Eternod M, Jones TE, Landry S, Pan T, Weitzman MD, David M.** 2012. Codon-usage-based inhibition of HIV protein synthesis by human schlafen 11. Nature **491:**125-128.

11.   **Wang Q, Lee I, Ren J, Ajay SS, Lee YS, Bao X.** 2013. Identification and functional characterization of tRNA-derived RNA fragments (tRFs) in respiratory syncytial virus infection. Mol Ther **21:**368-379.

12.   **Felton-Edkins ZA, Kondrashov A, Karali D, Fairley JA, Dawson CW, Arrand JR, Young LS, White RJ.** 2006. Epstein-Barr virus induces cellular transcription factors to allow active expression of EBER genes by RNA polymerase III. J Biol Chem **281:**33871-33880.

13.   **Felton-Edkins ZA, White RJ.** 2002. Multiple mechanisms contribute to the activation of RNA polymerase III transcription in cells transformed by papovaviruses. J Biol Chem **277:**48182-48191.

14.   **Gaynor RB, Feldman LT, Berk AJ.** 1985. Transcription of class III genes activated by viral immediate early proteins. Science **230:**447-450.

15.   **Hoeffler WK, Roeder RG.** 1985. Enhancement of RNA polymerase III transcription by the E1A gene product of adenovirus. Cell **41:**955-963.

16.   **Jang KL, Latchman DS.** 1989. HSV infection induces increased transcription of Alu repeated sequences by RNA polymerase III. FEBS Lett **258:**255-258.

17.   **Panning B, Smiley JR.** 1994. Activation of RNA polymerase III transcription of human Alu
elements by herpes simplex virus. Virology **202:**408-417.

18.   **Chiang JJ, Sparrer KMJ, van Gent M, Lassig C, Huang T, Osterrieder N, Hopfner KP, Gack**
MU. 2018. Viral unmasking of cellular 5S rRNA pseudogene transcripts induces RIG-I-
mediated immunity. Nat Immunol **19:**53-62.

19.   **Zhao Y, Ye X, Dunker W, Song Y, Karijolich J.** 2018. RIG-I like receptor sensing of host
RNAs facilitates the cell-intrinsic immune response to KSHV infection. Nat Commun
9:4841.

20.   **Karijolich J, Abernathy E, Glaunsinger BA.** 2015. Infection-Induced Retrotransposon-
Derived Noncoding RNAs Enhance Herpesviral Gene Expression via the NF-kappaB
Pathway. PLoS Pathog **11:**e1005260.

21.   **Dong X, Feng H, Sun Q, Li H, Wu TT, Sun R, Tibbetts SA, Chen ZJ, Feng P.** 2010. Murine
gamma-herpesvirus 68 hijacks MAVS and IKKbeta to initiate lytic replication. PLoS
Pathog **6:**e1001001.

22.   **Dong X, Feng P.** 2011. Murine gamma herpesvirus 68 hijacks MAVS and IKKbeta to
abrogate NFkappaB activation and antiviral cytokine production. PLoS Pathog
7:e1002336.

23.   **Krug LT, Collins CM, Gargano LM, Speck SH.** 2009. NF-kappaB p50 plays distinct roles in
the establishment and control of murine gammaherpesvirus 68 latency. J Virol **83:**4732-
4748.

24.   **Krug LT, Moser JM, Dickerson SM, Speck SH.** 2007. Inhibition of NF-kappaB activation in
vivo impairs establishment of gammaherpesvirus latency. PLoS Pathog **3:**e11.

25.   **Tucker JM, Glaunsinger BA.** 2017. Host noncoding retrotransposons induced by DNA
viruses: a SINE of infection? J Virol doi:10.1128/JVI.00982-17.

26.   **Schramm L, Hernandez N.** 2002. Recruitment of RNA polymerase III to its target
promoters. Genes Dev **16:**2593-2620.

27.   **White RJ, Stott D, Rigby PW.** 1990. Regulation of RNA polymerase III transcription in
response to Simian virus 40 transformation. EMBO J **9:**3713-3721.

28.   **Covarrubias S, Richner JM, Clyde K, Lee YJ, Glaunsinger BA.** 2009. Host shutoff is a
conserved phenotype of gammaherpesvirus infection and is orchestrated exclusively
from the cytoplasm. J Virol **83:**9554-9566.

29.   **Torres AG, Pineyro D, Rodriguez-Escriba M, Camacho N, Reina O, Saint-Leger A,**
Filonava L, Batlle E, Ribas de Pouplana L. 2015. Inosine modifications in human tRNAs
are incorporated at the precursor tRNA level. Nucleic Acids Res **43:**5145-5157.

30.   **Mohr S, Ghanem E, Smith W, Sheeter D, Qin Y, King O, Polioudakis D, Iyer VR, Hunicke-**
Smith S, Swamy S, Kuersten S, Lambowitz AM. 2013. Thermostable group II intron
reverse transcriptase fusion proteins and their use in cDNA synthesis and next-
generation RNA sequencing. RNA **19:**958-970.

31.   **Chan PP, Lowe TM.** 2009. GtRNAdb: a database of transfer RNA genes detected in
genomic sequence. Nucleic Acids Res **37:**D93-97.

32.   **Chan PP, Lowe TM.** 2016. GtRNAdb 2.0: an expanded database of transfer RNA genes
identified in complete and draft genomes. Nucleic Acids Res **44:**D184-189.

33.   **Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C.** 2017. Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods **14:**417-419.

34. **Karijolich J, Zhao Y, Alla R, Glaunsinger B.** 2017. Genome-wide mapping of infection-
induced SINE RNAs reveals a role in selective mRNA export. Nucleic Acids Res **45:**6194-
6208.
35. **Diebel KW, Claypool DJ, van Dyk LF.** 2014. A conserved RNA polymerase III promoter
required for gammaherpesvirus TMER transcription and microRNA processing. Gene
544:8-18.
36. **Graczyk D, Ciesla M, Boguta M.** 2018. Regulation of tRNA synthesis by the general
transcription factors of RNA polymerase III - TFIIIB and TFIIIC, and by the MAF1 protein.
Biochim Biophys Acta Gene Regul Mech **1861:**320-329.
37. **Willis IM, Moir RD.** 2018. Signaling to and from the RNA Polymerase III Transcription
and Processing Machinery. Annu Rev Biochem **87:**75-100.
38. **Orioli A, Praz V, Lhote P, Hernandez N.** 2016. Human MAF1 targets and represses active
RNA polymerase III genes by preventing recruitment rather than inducing long-term
transcriptional arrest. Genome Res **26:**624-635.
39. **Bonhoure N, Praz V, Moir RD, Willemin G, Mange F, Moret C, Willis IM, Hernandez N.**
2019. Chronic repression by MAF1 supports futile RNA cycling as a mechanism for
obesity resistance. bioRxiv doi:https://doi.org/10.1101/775353.
40. **Bonhoure N, Byrnes A, Moir RD, Hodroj W, Preitner F, Praz V, Marcelin G, Chua SC, Jr.,**
Martinez-Lopez N, Singh R, Moullan N, Auwerx J, Willemin G, Shah H, Hartil K,
Vaitheesvaran B, Kurland I, Hernandez N, Willis IM. 2015. Loss of the RNA polymerase
III repressor MAF1 confers obesity resistance. Genes Dev **29:**934-947.
41. **Abernathy E, Clyde K, Yeasmin R, Krug LT, Burlingame A, Coscoy L, Glaunsinger B.**
2014. Gammaherpesviral gene expression and virion composition are broadly controlled
by accelerated mRNA degradation. PLoS Pathog **10:**e1003882.
42. **Hartenian E, Glaunsinger B.** 2019. Herpesvirus infection reduces Pol II occupancy of
host promoters but spares viral promoters. bioRxiv doi:https://doi.org/10.1101/585984
.
43. **Larminie CG, Sutcliffe JE, Tosh K, Winter AG, Felton-Edkins ZA, White RJ.** 1999.
Activation of RNA polymerase III transcription in cells transformed by simian virus 40.
Mol Cell Biol **19:**4927-4934.
44. **Carey MF, Singh K, Botchan M, Cozzarelli NR.** 1986. Induction of specific transcription
by RNA polymerase III in transformed cells. Mol Cell Biol **6:**3068-3076.
45. **Barski A, Chepelev I, Liko D, Cuddapah S, Fleming AB, Birch J, Cui K, White RJ, Zhao K.**
2010. Pol II and its associated epigenetic marks are present at Pol III-transcribed
noncoding RNA genes. Nat Struct Mol Biol **17:**629-634.
46. **Yeganeh M, Praz V, Carmeli C, Villeneuve D, Rib L, Guex N, Herr W, Delorenzi M,**
Hernandez N, Cycli Xc. 2019. Differential regulation of RNA polymerase III genes during
liver regeneration. Nucleic Acids Res **47:**1786-1796.
47. **Gerber A, Ito K, Chu C, Roeder RG.** 2019. Induced RPB1 depletion reveals a direct gene-
specific control of RNA Polymerase III function by RNA Polymerase II. bioRxiv
doi:https://doi.org/10.1101/764605.
48. **Arimbasseri AG, Blewett NH, Iben JR, Lamichhane TN, Cherkasova V, Hafner M, Maraia**
RJ. 2015. RNA Polymerase III Output Is Functionally Linked to tRNA Dimethyl-G26
Modification. PLoS Genet **11:**e1005671.

49. **Karkusiewicz I, Turowski TW, Graczyk D, Towpik J, Dhungel N, Hopper AK, Boguta M.**
2011. Maf1 protein, repressor of RNA polymerase III, indirectly affects tRNA processing.
J Biol Chem **286:**39478-39488.
50. **Abernathy E, Glaunsinger B.** 2015. Emerging roles for RNA degradation in viral
replication and antiviral defense. Virology **479-480:**600-608.
51. **Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grasser FA, van Dyk LF,**
Ho CK, Shuman S, Chien M, Russo JJ, Ju J, Randall G, Lindenbach BD, Rice CM, Simon V,
Ho DD, Zavolan M, Tuschl T. 2005. Identification of microRNAs of the herpesvirus
family. Nat Methods **2:**269-276.
52. **Bogerd HP, Karnowski HW, Cai X, Shin J, Pohlers M, Cullen BR.** 2010. A mammalian
herpesvirus uses noncanonical expression and processing mechanisms to generate viral
MicroRNAs. Mol Cell **37:**135-142.
53. **Hanada T, Weitzer S, Mair B, Bernreuther C, Wainger BJ, Ichida J, Hanada R, Orthofer**
M, Cronin SJ, Komnenovic V, Minis A, Sato F, Mimata H, Yoshimura A, Tamir I, Rainer J,
Kofler R, Yaron A, Eggan KC, Woolf CJ, Glatzel M, Herbst R, Martinez J, Penninger JM.
2013. CLP1 links tRNA metabolism to progressive motor-neuron loss. Nature **495:**474-
480.
54. **Karaca E, Weitzer S, Pehlivan D, Shiraishi H, Gogakos T, Hanada T, Jhangiani SN,**
Wiszniewski W, Withers M, Campbell IM, Erdin S, Isikay S, Franco LM, Gonzaga-
Jauregui C, Gambin T, Gelowani V, Hunter JV, Yesil G, Koparir E, Yilmaz S, Brown M,
Briskin D, Hafner M, Morozov P, Farazi TA, Bernreuther C, Glatzel M, Trattnig S, Friske
J, Kronnerwetter C, Bainbridge MN, Gezdirici A, Seven M, Muzny DM, Boerwinkle E,
Ozen M, Baylor Hopkins Center for Mendelian G, Clausen T, Tuschl T, Yuksel A, Hess A,
Gibbs RA, Martinez J, Penninger JM, Lupski JR. 2014. Human CLP1 mutations alter tRNA
biogenesis, affecting both peripheral and central nervous system function. Cell **157:**636-
650.
55. **Adler H, Messerle M, Wagner M, Koszinowski UH.** 2000. Cloning and mutagenesis of
the murine gammaherpesvirus 68 genome as an infectious bacterial artificial
chromosome. J Virol **74:**6964-6974.
56. **Richner JM, Clyde K, Pezda AC, Cheng BY, Wang T, Kumar GR, Covarrubias S, Coscoy L,**
Glaunsinger B. 2011. Global mRNA degradation during lytic gammaherpesvirus infection
contributes to establishment of viral latency. PLoS Pathog **7:**e1002150.
57. **Lowe TM, Eddy SR.** 1997. tRNAscan-SE: a program for improved detection of transfer
RNA genes in genomic sequence. Nucleic Acids Res **25:**955-964.
58. **Hoffmann A, Fallmann J, Vilardo E, Morl M, Stadler PF, Amman F.** 2018. Accurate
mapping of tRNA reads. Bioinformatics **34:**2339.
59. **Langmead B, Salzberg SL.** 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods **9:**357-359.
60. **Risso D, Ngai J, Speed TP, Dudoit S.** 2014. Normalization of RNA-seq data using factor
analysis of control genes or samples. Nat Biotechnol **32:**896-902.
61. **Robinson MD, McCarthy DJ, Smyth GK.** 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics **26:**139-
140.

62.   **McCarthy DJ, Chen Y, Smyth GK.** 2012. Differential expression analysis of multifactor
696   RNA-Seq experiments with respect to biological variation. Nucleic Acids Res **40:**4288-
697   4297.

698