1

SCSA: a cell type annotation tool for single-cell RNA-seq data

Yinghao Cao[1], Xiaoyue Wang[1]*, Gongxin Peng[1]*

[1]Center for Bioinformatics, Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences, School of Basic Medicine, Peking Union Medical College, Beijing, China

*Correspondence

Corresponding Author

Xiaoyue Wang

wxy@ibms.pumc.edu.cn

Gongxin Peng

penggongxin@ibms.pumc.edu.cn

Keywords: single-cell RNA sequencing, cell type annotation, CellMarker database, score annotation model, differentially expressed genes

Abstract

Currently most methods take manual strategies to annotate cell types after clustering the single-cell RNA sequencing (scRNA-seq) data. Such methods are labor-intensive and heavily rely on user expertise, which may lead to inconsistent results. We present SCSA, an automatic tool to annotate cell types from scRNA-seq data, based on a score annotation model combining differentially expressed genes (DEGs) and confidence levels of cell markers from both known and user-defined information. Evaluation on

real scRNA-seq datasets from different sources with other methods shows that SCSA

is able to assign the cells into the correct types at a fully automated mode with a

desirable precision.

Introduction

Recent development of scRNA-seq methods has enabled unbiased, high-resolution

transcriptomic analysis of individual cells in a heterogeneous cell population (Tang et

al., 2009; Picelli et al., 2013a; Kolodziejczyk et al., 2015; Haque et al., 2017). scRNA-

seq methods have been used to characterize thousands to millions of cells in developing

embryos (Chu et al., 2016), immune cells (Shalek et al., 2013), complex tissues such as

brain (Zhong et al., 2018) and tumor (Chung et al., 2017), which have greatly promoted

our understanding of human development and diseases.

At the core of myriad scRNA-seq applications is the ability to identify different cell

types and cellular states from a complex cell mixture based on gene expression profiles.

Recently, several computational methods have been developed for annotating cell types

in scRNA-seq using cell-based mapping strategy. For instance, SingleR (Aran et al.,

2019) infers the cell type for each of the single cells using a novel hierarchical clustering

method based on similarity to the reference transcriptomic datasets of purified cell types.

Similarly, scMatch (Hou et al., 2019) annotates single cells by identifying their closest

match in gene expression profiles of large reference dataset, such as the Functional

Annotation Of The Mammalian Genome 5 (FANTOM5) resource   (Brown et al., 2009;

Lizio et al., 2017). However, such approaches required transcriptome of purified cells

or pre-annotated scRNA-seq data, ideally under the same experimental design using the

same platform, which is often not available. Using prior knowledge of cell-type specific

marker genes increased the accuracy and efficiency of cell type assignment, allowing

for identification of both known and *de novo* cell types in scRNA-seq data of complex

tissues, as shown by CellAssign (Zhang et al., 2019a) and Garnett (Pliner et al., 2019a).

A more common practice for cell type annotation is the cluster-then-annotate

approach. Unsupervised clustering methods based on dimension reduction algorithms

such as principal component analysis (PCA) and t-distributed stochastic neighbor

embedding (t-SNE) have been developed to partition the cells based on the similarity

of their gene expression patterns (Bacher and Kendziorski, 2016; Kiselev et al., 2019).

And users could manually assign a cell type to each cluster based on differentially

expressed markers by consulting the literature for cell-type specific gene markers.

However, there is still some problem associated with the manual annotation step. For

example, the canonical marker genes used in the assignment process may have an

impact on the annotation accuracy, which may lead to biased results with uncontrolled

vocabularies for cell type labels in different datasets. Expert-curated knowledge

databases such as CellMarker (Zhang et al., 2019b) and CancerSEA (Yuan et al., 2019b),

have been developed to provide a comprehensive and unified resource of cell markers

for various cell types in human and mouse tissues. Yet it is still lack a method to

leverage the information in these databases for annotation. Furthermore, marker genes

could express in more than one cell type, make the annotation more complex for human

to process.

To overcome these difficulties and to streamline the cell type assignment process for

scRNA-seq data, we developed SCSA, an algorithm that can automatically assign cell

types for each cell cluster in scRNA-seq data. SCSA uses marker genes of known cell

types highly expressed in a cell cluster to label the cluster. It can be directly applied to

clustering results generated from other scRNA-seq analysis softwares such as

CellRanger (Zheng et al., 2017) and Seurat (Butler et al., 2018). To eliminate the bias

of marker selection, SCSA integrates all marker genes to cell-type matrix from

CellMarker (Zhang et al., 2019b) and CancerSEA (Yuan et al., 2019a) database, using

a score annotation model that accounts for quantitative information and discrepancies

information of marker genes. For cell clusters lacking known cell markers, SCSA will

also perform a Gene Ontology (GO) enrichment analysis and report the results to give

some clues to the user. Through extensive evaluation on several real scRNA-seq

datasets of both human and mouse origin generated from different platforms including

Smart-seq (Ramskold et al., 2012)，Smart-seq2 (Picelli et al., 2013b) and 10x Genomics

(Zheng et al., 2017). We demonstrated that SCSA has an excellent and unbiased

performance on cell type annotation with a desirable precision.

Materials and methods

Marker genes identification

The input of SCSA is a DEGs clusters matrix, in a format that is supported by the

clustering output of CellRanger or Seurat. Based on the matrix, SCSA identifies the

marker genes of each cell cluster through a filtration with log2-based fold-change (LFC)

value and P-value (LFC >= 1, P <= 0.05). For each cell cluster, a marker gene

identification vector is generated for j genes with LFC values, which is defined as $E_j =$

$\{e_1, e_2, \cdots e_j\} = (e)_{j\times1}$, here, e represents the absolute value of LFC multiplied by

mean of all.

Cell marker database

In order to improve the accuracy of cell cluster annotation for scRNA-seq data, SCSA

uses cell markers from two public databases: CellMarker and CancerSEA. SCSA

integrated 11,464 manually curated cell markers of 459 cell types in 158 human

tissues/sub-tissues and 7,855 cell markers of 385 cell types in 80 mouse tissues/sub-

tissues from CellMarker database. Also SCSA integrates 1,244 markers from

CancerSEA database, which consists of 14 functional states from 25 human cancer

types. Furthermore, SCSA can accept user-defined marker gene database as additional

information for cell cluster annotation. The user-defined marker gene database must

have two columns, with the name of cell types in the first column and marker gene for

each cell type in the second column. In that case, SCSA will combine both known

databases and the custom database to predict the annotations for cell clusters.

Annotation model construction

For those genes which existed in both the DEGs and known cell marker databases, SCSA constructs a cell-gene sparse matrix (defined as $M_1 = (a_{ij})_{c_1 \times g_1}$) with $c_1(c_1 \leq$ i) cells and $g_1(g_1 \leq j)$ genes as "marker evidence". Here, for each cell i and each gene j in the matrix M_1, a refers to the sum number of references in the CellMarker database. To eliminate the huge differences of marker evidence between the well-known gene and less-known genes, we transform the value to log2-based and plus a constant (0.05). Also, to represent the whole gene set for a certain cell, we create a cell type style vector which takes multiplication of standard deviation of the marker evidence and marker numbers (defined as $L_1 = \{l_1, l_2, \cdots l_{c1}\}$, where $l = std(a_{ij}) * num(a_{ij} > 0)$). So, for the known marker database, the raw score vector of a cell type is $S_{c_1} = M_1 \times E_{g1} * L_1 = \{s_1, s_2, \cdots s_{c_1}\}$.

For marker databases from multiple sources including known and user-defined ones, with k as the total number of databases, multiple cell-gene sparse matrices could be constructed according to the DEGs, which are defined as $M_k = (a_{ij})_{p \times q}$, $p \times q \in \{(c_1 \times g_1), (c_2 \times g_2), \cdots, (c_k \times g_k)\}$. With the definition of multiple gene expression vectors defined as $E_k = (e)_{q \times 1}, q \in \{g_1, g_2, \cdots, g_k\}, (q \leq j)$, k raw score vectors will be generated, which is defined as $S_k = M_k E_k L_k = (a_{ij})_{p \times q} \times (e)_{q \times 1} * (l)_{q \times 1} = (s)_{p \times 1}, p \in \{c_1, c_2, \cdots c_k\}$. Suppose S_k to be a function to the score vectors, which is defined as F_l. To eliminate the difference of those vectors and compare them with each other standardly, SCSA performs z-score normalization for them. In detail,

$$F_l = M_l E_l L_l = (f_l)_{p \times 1}, l \in [1, k]$$

$$Z_l = (z_{mn})_{p \times 1} = (f'_l)_{p \times 1}, l \in [1, k]$$

$$f'_l = \frac{f_l - \bar{f}}{d}, \bar{f} = \frac{1}{p} \sum_{l=1}^{p} f_l, d = \sqrt{\frac{1}{p-1} \sum_{l=1}^{p} (f_l - \bar{f})^2}$$

Notably, the score vectors derived from different kinds of databases may have

different lengths. To give a uniform score to a certain cell type, SCSA transforms them

to the same length:

$$Z'_l = (z_{m'n'})_{p'\times 1}, z_{m'n'} = \begin{cases} z_{mn}, & m = m', n = n' \\ 0, & m \neq m' \parallel n \neq n' \end{cases}, p' = num\{c_1 \cup c_2 \cup \cdots \cup c_k\}$$

Finally, an annotation model will be constructed by merging the database weight

coefficient matrix W and the last uniform score vector.

$$S' = (Z'_1, Z'_2, \cdots, Z'_k)W + b, W = (w_z)_{k\times 1}$$

GO enrichment analysis

Not all gene markers of cell types have been curated in known cell marker databases.

To solve this problem, SCSA employs a GO enrichment analysis to give some clues to

the user to identify new cell types. In detail, for a certain GO term, SCSA uses Fisher's

exact test to calculate the P-values, using DEGs of the selected cluster as foreground

values and DEGs in other clusters as background values, respectively. After that, P-

value was adjusted by the Benjamini-Hochberg (BH) method (Benjamini and Hochberg,

1995).

FANTOM5+SingleR dataset

FANTOM5+SingleR dataset was collected from GitHub repository of scMatch

(*https://github.com/forrest-lab/scMatch*). The dataset contained gene expression data

of 916 human samples (660 primary cell samples and 256 cell lines) and 821 mouse

samples (471 primary cell line samples, 302 tissue samples, and 48 transformed cell

lines) from FANTOM5 and 972 human samples and 1188 mouse samples in SingleR's

reference dataset. Annotation of these cells into certain categories were also

downloaded from the same repository. For human cells in FANTOM5+SingleR dataset,

terms of cell types associated with more than 10 samples were collected as the

FANTOM5_hs1 dataset. And cell types associated with more than 40 samples were

used FANTOM5_hs2 datasets. Similarly, FANTOM5_mm1 contained cell types with

more than 10 samples and FANTOM5_mm2 included cell types with more than 40

samples in the FANTOM5+SingleR mouse dataset.

Real scRNA-seq datasets

Four real datasets (GSE72056 (Tirosh et al., 2016b), GSE81861 (Li et al., 2018), E-

MTAB-6149 (Lambrechts et al., 2018), and the PBMCs dataset were used for

evaluation of SCSA. GSE72056, GSE81861 and E-MTAB-6149 datasets are single cell

data obtained from human tumors (melanoma, colorectal and lung) using Smart-seq2,

Smart-seq and 10X genomics technology, respectively. Normal cells in the GSE72056

dataset was downloaded from Gene Expression Omnibus (GEO) repository

(*https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE72056*)   and used to

optimize the parameters for SCSA. Normal cells in the GSE81861 dataset was

downloaded   from   GEO

(*https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE81861*).   E-MTAB-6149

was   downloaded   from   ArrayExpress   repository

(*https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-6149*).

The peripheral blood mononuclear cells (PBMCs) datasets, including four scRNA-

seq data of peripheral blood mononuclear cells (3k, 4k, 6k and 8k of PBMCs) were

downloaded   from   the   10X   Genomics   official   website

(*https://www.10xgenomics.com/resources/datasets*). The 3k and 6k data were blood

samples from one donor generated using the v1. chemistry and preprocessed with

CellRanger1.1.0. They were labeled "3k PBMCs from a healthy donor" and "6k

PBMCs from a healthy donor", respectively. The 4k and 8k datasets, under the label

"4k PBMCs from a healthy donor" and "8k PBMCs from a healthy donor", were

samples collected from one donor, generated using the v2. Chemistry and preprocessed

using CellRanger2.1.0.

Evaluation of SCSA performance on real datasets

For the known cell type datasets, we defined cell type cluster by their real cell types.

Then we generated calculated the DEGs for each clusters using two methods. One is

traditional Student's t-test perform by in-house script and the other using Seurat. We

obtained the final two lists of DEGs for the input of SCSA through setting the threshold

with P-value of 0.001 and LFC value of 1.

For the 4 PBMCs datasets, data preprocessing, normalization and unsupervised

clustering were already performed by CellRanger workflow from its website. Since

10X Genomics official website illustrated a workflow example using 3k PBMCs from

a healthy donor containing 5 cell clusters and gave a final annotation results contained

monocytes, T cells, NK cells, megakaryocytes, and B cells, respectively, we choose

results with five clusters to do the further evaluation. SCSA identified the DEGs of each

cluster through the LFC (LFC>=1.5) value and P-value (P<=0.05) and predicted the

cell types according to the clusters.

To evaluate the stability of SCSA in annotating the cell type of a cluster, a heat map

was generated using hierarchical clustering method for all cell types of top five scores

in a cell cluster. To further demonstrate the accuracy of SCSA, we calculated the

percentage of the five clusters cell types to measure the abundance of the same cell type

based on the prediction of SCSA in the four PBMCs datasets, respectively.

Comparison with other cell type annotation tools

Three annotation tools, scMatch (Hou et al., 2019), CellAssign (Zhang et al., 2019a)

and Garnett (Pliner et al., 2019a) were used for comparison. scMatch was run with

default parameters using FANTOM5 as the reference dataset. For CellAssign (version

0.99.2), a build-in set of markers contain 8 common cell types in human tumor

microenvironment were used to test for human datasets. For Garnett (version 0.1.14),

two pre-trained classifiers, trained from human PBMCs tissue (Garnett_human_pbmc)

and lung tissue (Garnett_human_lung), were downloaded from (*https://cole-trapnell-*

lab.github.io/garnett/classifiers) and used for classification. Because CellAssign and

Garnett only annotate cell types for each of single cells instead of cell cluster, to make

the evaluation at the same level, all cells of a certain cluster were predicted and the cell

type with the maximum occurrence was chosen as the final predicted result. Three tags

were used to compare the results. "Positive" meant correct prediction to the cell type.

"Negative" meant incorrect prediction cell type. "Missed" meant no clear cell type

prediction. Accuracy of cluster was calculated as sum of clusters with "Positive" tag

divided by sum of all clusters.

Software availability

SCSA is implemented in Python3 as an open source software under the GNU General

Public License, and the source code is freely available together with full documentation

at *https://github.com/bioinfo-ibms-pumc/SCSA*.

Results

Design principles of SCSA

The SCSA algorithm is a three-step procedure that includes marker genes identification,

annotation model construction, and GO enrichment analysis (Figure 1). First, the input

of SCSA is a gene expression matrix with cell cluster information (such as the output

results of CellRanger or Seurat). SCSA identifies a group of marker genes for each

cluster from input expression matrix by differential gene expression analysis. Next, for

each cluster, genes identified as marker genes that have one or more linked cell types

in a database will be used to generate a cell-gene matrix for that cluster. For each cell

type in the matrix, SCSA then used a decision model to assign a score by combining

the enrichment of marker gene expression and the strength of evidence for the marker

genes in the database. SCSA could also take marker gene information from multiple

databases and assign different weights to them. Finally, SCSA provides an alternative

gene ontology enrichment analysis step to give some clues to the user for the function

of a cell cluster in addition to its annotation.

Evaluation performance of SCSA on known datasets

We first evaluated the performance of SCSA on three scRNA-seq datasets with known

cell types (Table S1). The GSE72056 dataset include 2,840 single cells isolated from

normal tissues of 19 melanoma patients (Tirosh et al., 2016a). These cells have been

annotated manually by experts into six cell types, namely B cells, T cells, Cancer-

associated fibroblast (CAF), Macrophage, Endothelial cells and Natural killer cells.

SCSA accurately annotated five of the six cell types, except for the "Cancer-associated

fibroblast (CAF)" cells (Table S2, S3). Instead, SCSA gave the CAF cluster a different

label named "Mesenchymal stem cell". It has been reported that fibroblasts shared more

common features with mesenchymal stem cells (Haniffa et al., 2009) by expressing

similar cell immunophenotypic markers, as well as the genes that are known to be

expressed in stem cells (Brohem et al., 2013). Therefore, we considered the

"Mesenchymal stem cell (MSC)" label SCSA reported concordant with the "CAF"

label assigned by human expert. Similarly, for seven known clusters in GSE81861

dataset (Li et al., 2017), which contains 265 single cells derived from normal tissues

adjacent to 11 colorectal tumors, SCSA correctly predicted six of the seven clusters and

identified the "Fibroblast" group as "MSC" (Table S2, S3). In the lung cancer dataset

of 45,232 normal cells containing 7 known cell types (Lambrechts et al., 2018), SCSA

accurately assigned B cells, Endothelial cells, Epithelial cells, Myeloid and T cells, and

identified "Fibroblast" group as "MSC" and "Alveolar" as "Epithelial Cell" (Table S2,

S3).

In order to evaluate the robustness of SCSA on large scRNA-seq datasets, we used

four PBMCs (3k, 4k, 6k and 8k) datasets from 10X genomics website. We collected all

possible cell types of a cell cluster according to the top five scores under the score

annotation model of SCSA. The correlation of all cell types and scores were calculated

and compared. As shown in Figure 3A, based on the five annotated cell types

(monocyte cells, T cells, NK cells, megakaryocytes cells, and B cells) of CellRanger,

SCSA achieved a great consistency in the four PBMCs datasets. Notably, the

"macrophage cell" type was predicted as the second top score for the cluster, which was

annotated as "monocytes" by SCSA due to the reason that they share many marker

genes.

To further demonstrate the robustness of SCSA over the five annotated cell types

(monocytes cells, T cells, NK cells, megakaryocytes cells, and B cells), we compared

their abundance in each cluster using the four PBMCs datasets. As shown in Figure 3B,

the percentages of cell numbers for five cell types annotated by SCSA remained stable

across these datasets. T cells occupied half of the PBMCs, and monocytes cell

represented another 25%, B cells and NK cells had similar levels, while

megakaryocytes cell has the lowest number among all the five cell types. Meanwhile,

as shown in Figure 3C, SCSA can predicted the five cell types consistent with the

reference information of 4 PBMCs.

In addition, to test the GO module in SCSA, which is designed to give some clues

on cluster cell function in addition to its label, we collected gene expression profiles of

primary cells and cell lines from FANTOM5+SingleR dataset, which contains both

human and mouse cell types (Table S8). For cell types with 10 or more samples, SCSA

achieved a 60% (15 of 25) and a 57% (12 of 21) accuracy for human and mouse data

respectively. For cell types containing more than 40 samples, the accuracy of SCSA

were improved to 73% (8 of 11) for human data and achieved 56% (5 of 9) for mouse

data, respectively (Table S2). For the "Aortic smooth muscle cell" cell type, which was

not identified correctly due to lack of evidence in the CellMarker reference database, a

GO analysis step in SCSA revealed a functional enrichment in the term "extracellular

matrix structural constituent", suggesting its role in regulating cell shape and cross-talk

with extracellular matrix (Owens et al., 2004).

Comparison to other cell type annotation tools

We compared the performance of SCSA with three other tools (scMatch (Hou et al.,

2019), CellAssign (Zhang et al., 2019a) and Garnett (Pliner et al., 2019a)) using 3

known cell type datasets. Since scMatch, CellAssign and Garnett annotate each of the

single cells instead of cell clusters, all cells were first annotated and a cell type with the

maximum occurrence in the cluster was defined as the final predicted result of the

cluster (Table S2-S8). As showed in Figure 2, for the three human tumor datasets

(GSE72756, GSE81861 and E-MTAB-6149), SCSA achieved the highest accuracy.

scMatch owned the similar accuracy with Garnett when using the predefined lung

classifier (Garnett_human_lung) on the three human tumor datasets. CellAssign had a

better prediction results than Garnett on GSE72756 and GSE81861 datasets, with 83%

and 86% accuracy, respectively. However, for E-MTAB-6149 dataset with more than

45,000 cells, CellAssign only yielded 43% accuracy which was lower than Garnett

using pre-trained classifier from human lung (Garnett_human_lung) (71%). A possible

explanation for these might be that CellAssign was not suitable to annotate cell types

for large datasets. Another possible explanation for these was that E-MTAB-6149

dataset is the training-set of pre-trained classifier of Garnett.

Cell-based annotation approaches could assign multiple cell type labels to one

cluster due to cell heterogeneity in the clusters. Evaluation results of CellAssign in E-

MTAB-6149 illustrated that only 8 (0.1%) of B cells were correctly assigned for true B

cell cluster, whereas 2,861 (51.1%) cells were mistakenly assigned as myofibroblast

cells and 1,755 (31.3%) cells were not able to assign a clear cell type. In general，most

predicted cell types in the cluster (>50%) from these tools were not consistent with true

cell type. For example, the T cells of E-MTAB-6149 dataset was not annotated by

CellAssign, the fibroblasts of GSE81861 dataset was missed by scMatch, and the

natural killer cells of GSE72056 dataset were not identified by CellAssign and scMatch

(Table S2, S4-S6). Moreover, Garnett failed to assign epithelial cells and T cells of E-

MTAB-6149 dataset although using classifier trained from the same dataset.

Discussion

Currently, for scRNA-seq data, cell type annotation of cell clusters after unsupervised

clustering is mainly conducted manually. The limitation of the manual procedure makes

it impossible to generate high-quality, reproducible, and standardized annotation results

for the growing number of scRNA-seq datasets.

To address this challenge, we presented a novel tool, SCSA, for automatic annotating

the cell types from single-cell RNA sequencing data, which can be applied directly on

the output generated from CellRanger or Seurat. By introducing the newly developed

annotation model merging DEGs and cell markers reference information to replace the

manual steps, and based on the extensive evaluation of performance on 7 real known

datasets, as well as comparisons with alternative methods (CellAssign (Zhang et al.,

2019a), Garnett (Pliner et al., 2019a) and scMatch (Hou et al., 2019)), SCSA can

perform the annotation task at a high accuracy and efficient level and have a

preferentially choice on annotating cell types in tumor and rare cell type datasets.

In cell type annotation, it is usually hard to find high-quality marker genes to

describe a cell cluster. A strategy is to use genes specifically expressed in a cell cluster

to mark the cell type. However, using a few marker genes is often not sufficient to

distinguish a cell cluster from the others. In addition, using the whole expressed gene

sets may decrease the power to find the true patterns within each cell cluster. Therefore,

we used DEGs in the marker gene identification step in SCSA. This step avoids the

influence of ubiquitously expressed genes and collects the appropriate genes for

calculating the optimal score in the annotation model. There still exist some limitations,

which may influence the accuracy of cell type annotation using SCSA. First, the

quantity of marker genes in these cell marker databases greatly impacted the results of

cell type annotation. Since cell marker collection is far away from completion, it is

possible that some cell types are unclassifiable due to the lack of appropriate markers.

Specifically, this phenomenon is quite common for unknown tissues and novel sub-

clusters of cells at different states. User-defined marker combinations need to be

developed to solve this problem. SCSA can accept them as additional information to

improve the annotation results. Second, for complex tissues such as cancer tissues, the

accuracy of cell annotation is heavily relied on the clustering algorithms. Different

unsupervised clustering method could have different results, especially when the cluster

size is unevenly distributed in the population (Kiselev et al., 2019). In that situation,

algorithms using supervised clustering may be more appropriate for cell type

classification (Pliner et al., 2019b).

Compared with the results of SCSA over different datasets, SCSA exhibited a

reasonable accuracy and robustness in cell type annotation. Further efforts could be

made to improve the annotation ability of SCSA by taking into account more

information (e.g., the more accurate information of cell marker genes, the

comprehensive clustering algorithm). We believe that SCSA is an important addition to

the toolbox used for single-cell studies and will greatly improve our efficiency and

capacity to explore the functional potential of novel cell types.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

GP proposed the method and conceived of the study. YC optimized algorithm and

designed the program; YC and GP analyzed the data. YC, GP, and XW wrote the

manuscript. All authors discussed the results and commented on the manuscript. All

authors read and approved the final manuscript.

Funding

This work was supported by grants from the CAMS Innovation Fund for Medical

Sciences (2017-I2M-1-009) and National Key R&D Program of China

(2018YFA0109800).

Acknowledgment

We thank Dr. Fangqing Zhao for the insightful suggestions for finishing the manuscript.
This manuscript has been released as a pre-print at
https://doi.org/10.1101/2019.12.22.886481 (Cao et al., 2019).

Reference

Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., et al. (2019). Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* 20(2)**,** 163-172. doi: 10.1038/s41590-018-0276-y.

Bacher, R., and Kendziorski, C. (2016). Design and computational analysis of single-cell RNA-sequencing experiments. *Genome Biology* 17. doi: ARTN 63
10.1186/s13059-016-0927-y.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Statistical Methodology* 57(1)**,** 289-300.

Brohem, C.A., de Carvalho, C.M., Radoski, C.L., Santi, F.C., Baptista, M.C., Swinka, B.B., et al. (2013). Comparison between fibroblasts and mesenchymal stem cells derived from dermal and adipose tissue. *International Journal of Cosmetic Science* 35(5)**,** 448-457. doi: 10.1111/ics.12064.

Brown, S.D., Wurst, W., Kuhn, R., and Hancock, J.M. (2009). The functional annotation of mammalian genomes: the challenge of phenotyping. *Annu Rev Genet* 43**,** 305-333. doi: 10.1146/annurev-genet-102108-134143.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* 36(5)**,** 411-420. doi: 10.1038/nbt.4096.

Cao, Y., Wang, X., and Peng, G. (2019). SCSA: a cell type annotation tool for single-cell RNA-seq data. *bioRxiv***,** 2019.2012.2022.886481. doi: 10.1101/2019.12.22.886481.

Chu, L.F., Leng, N., Zhang, J., Hou, Z.G., Mamott, D., Vereide, D.T., et al. (2016).

Single-cell RNA-seq reveals novel regulators of human embryonic stem cell

differentiation to definitive endoderm. *Genome Biology* 17. doi: ARTN 173

10.1186/s13059-016-1033-x.

Chung, W., Eum, H.H., Lee, H.O., Lee, K.M., Lee, H.B., Kim, K.T., et al. (2017).

Single-cell RNA-seq enables comprehensive tumour and immune cell profiling

in primary breast cancer. *Nature Communications* 8. doi: ARTN 15081

10.1038/ncomms15081.

Haniffa, M.A., Collin, M.P., Buckley, C.D., and Dazzi, F. (2009). Mesenchymal stem

cells: the fibroblasts' new clothes? *Haematologica-the Hematology Journal*

94(2)**,** 258-263. doi: 10.3324/haematol.13699.

Haque, A., Engel, J., Teichmann, S.A., and Lonnberg, T. (2017). A practical guide to

single-cell RNA-sequencing for biomedical research and clinical applications.

Genome Medicine 9. doi: ARTN 75

10.1186/s13073-017-0467-4.

Hou, R., Denisenko, E., and Forrest, A.R.R. (2019). scMatch: a single-cell gene

expression profile annotation tool using reference datasets. *Bioinformatics*

35(22)**,** 4688-4695. doi: 10.1093/bioinformatics/btz292.

Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Challenges in unsupervised

clustering of single-cell RNA-seq data (vol 20, pg 273, 2019). *Nature Reviews*

Genetics 20(5)**,** 310-310. doi: 10.1038/s41576-019-0095-5.

Kolodziejczyk, A.A., Kim, J.K., Svensson, V., Marioni, J.C., and Teichmann, S.A.

(2015). The technology and biology of single-cell RNA sequencing. *Mol Cell*

58(4)**,** 610-620. doi: 10.1016/j.molcel.2015.04.005.

Lambrechts, D., Wauters, E., Boeckx, B., Aibar, S., Nittner, D., Burton, O., et al. (2018).

Phenotype molding of stromal cells in the lung tumor microenvironment.

Nature Medicine 24(8)**,** 1277-+. doi: 10.1038/s41591-018-0096-5.

Li, H., Courtois, E.T., Sengupta, D., Tan, Y., Chen, K.H., Goh, J.J.L., et al. (2017).

Reference component analysis of single-cell transcriptomes elucidates cellular

heterogeneity in human colorectal tumors. *Nat Genet* 49(5)**,** 708-718. doi:

10.1038/ng.3818.

Li, H.P., Courtois, E.T., Sengupta, D., Tan, Y., Chen, K.H., Goh, J.J.L., et al. (2018).

Reference component analysis of single-cell transcriptomes elucidates cellular

heterogeneity in human colorectal tumors (vol 50, pg 1754, 2018). *Nature*

Genetics 50(12)**,** 1754-1754. doi: 10.1038/s41588-018-0299-1.

Lizio, M., Harshbarger, J., Abugessaisa, I., Noguchi, S., Kondo, A., Severin, J., et al.

(2017). Update of the FANTOM web resource: high resolution transcriptome of

diverse cell types in mammals. *Nucleic Acids Res* 45(D1)**,** D737-D743. doi:

10.1093/nar/gkw995.

Owens, G.K., Kumar, M.S., and Wamhoff, B.R. (2004). Molecular regulation of

vascular smooth muscle cell differentiation in development and disease. *Physiol*

Rev 84(3)**,** 767-801. doi: 10.1152/physrev.00041.2003.

Picelli, S., Bjorklund, A.K., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg,

R. (2013a). Smart-seq2 for sensitive full-length transcriptome profiling in single

cells. *Nature Methods* 10(11)**,** 1096-1098. doi: 10.1038/Nmeth.2639.

Picelli, S., Bjorklund, A.K., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg,

R. (2013b). Smart-seq2 for sensitive full-length transcriptome profiling in

single cells. *Nat Methods* 10(11)**,** 1096-1098. doi: 10.1038/nmeth.2639.

Pliner, H.A., Shendure, J., and Trapnell, C. (2019a). Supervised classification enables

rapid annotation of cell atlases. *Nat Methods* 16(10)**,** 983-986. doi:

10.1038/s41592-019-0535-3.

Pliner, H.A., Shendure, J., and Trapnell, C. (2019b). Supervised classification enables

rapid annotation of cell atlases. *Nature Methods* 16(10)**,** 983-+. doi:

10.1038/s41592-019-0535-3.

Ramskold, D., Luo, S., Wang, Y.C., Li, R., Deng, Q., Faridani, O.R., et al. (2012). Full-

length mRNA-Seq from single-cell levels of RNA and individual circulating

tumor cells. *Nat Biotechnol* 30(8)**,** 777-782. doi: 10.1038/nbt.2282.

Shalek, A.K., Satija, R., Adiconis, X., Gertner, R.S., Gaublomme, J.T., Raychowdhury,

R., et al. (2013). Single-cell transcriptomics reveals bimodality in expression

and splicing in immune cells. *Nature* 498(7453)**,** 236-240. doi:

10.1038/nature12172.

Tang, F.C., Barbacioru, C., Wang, Y.Z., Nordman, E., Lee, C., Xu, N.L., et al. (2009).

mRNA-Seq whole-transcriptome analysis of a single cell. *Nature Methods* 6(5)**,**

377-U386. doi: 10.1038/Nmeth.1315.

Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., 2nd, Treacy, D., Trombetta, J.J.,

et al. (2016a). Dissecting the multicellular ecosystem of metastatic melanoma

by single-cell RNA-seq. *Science* 352(6282)**,** 189-196. doi:

10.1126/science.aad0501.

Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., Treacy, D., Trombetta, J.J., et al.

(2016b). Dissecting the multicellular ecosystem of metastatic melanoma by

single-cell RNA-seq. *Science* 352(6282)**,** 189-196. doi:

10.1126/science.aad0501.

Yuan, H., Yan, M., Zhang, G., Liu, W., Deng, C., Liao, G., et al. (2019a). CancerSEA:

a cancer single-cell state atlas. *Nucleic Acids Res* 47(D1)**,** D900-D908. doi:

10.1093/nar/gky939.

Yuan, H.T., Yan, M., Zhang, G.X., Liu, W., Deng, C.Y., Liao, G.M., et al. (2019b).

CancerSEA: a cancer single-cell state atlas. *Nucleic Acids Research* 47(D1)**,**

D900-D908. doi: 10.1093/nar/gky939.

Zhang, A.W., O'Flanagan, C., Chavez, E.A., Lim, J.L.P., Ceglia, N., McPherson, A., et

al. (2019a). Probabilistic cell-type assignment of single-cell RNA-seq for tumor

microenvironment profiling. *Nat Methods* 16(10)**,** 1007-1015. doi:

10.1038/s41592-019-0529-1.

Zhang, X., Lan, Y., Xu, J., Quan, F., Zhao, E., Deng, C., et al. (2019b). CellMarker: a

manually curated resource of cell markers in human and mouse. *Nucleic Acids*

Res 47(D1)**,** D721-D728. doi: 10.1093/nar/gky900.

Zheng, G.X., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., et al. (2017).

Massively parallel digital transcriptional profiling of single cells. *Nat Commun*

8**,** 14049. doi: 10.1038/ncomms14049.

Zhong, S.J., Zhang, S., Fan, X.Y., Wu, Q., Yan, L.Y., Dong, J., et al. (2018). A single-

cell RNA-seq survey of the developmental landscape of the human prefrontal

cortex. *Nature* 555(7697)**,** 524-+. doi: 10.1038/nature25980.

M: Cell-Gene Matrix   *E*: Expression Vector *L*: Cell Style Vector
S: Raw Score Vector   *W*: Database Weight Vector Z: Combined Raw Score Vector

Figure 1. Flowchart of the SCSA. First, DEGs of each cluster will be extracted and

filtered from gene expression file. Next, SCSA employs marker gene databases to

annotate cell clusters. In this step, both known marker gene database and user-defined

marker database could be used simultaneously. For each cluster each database, a cell-

gene matrix (*M*) with two vectors (*E, L)* will be generated to form a raw score vector

(*S*). If multiple databases were selected, vectors would be normalized and combined

together to make a new vector (*Z*), then multiplied with a database weight matrix (*W*)

to make the last uniform vector. In the last step, ranked cell type vector will be generated

according to the uniform score. In addition, SCSA employs GO enrichment analysis to

give users some clue for unidentified clusters.

Figure 2. Performance of SCSA in comparison with other methods (scMatch，CellAssign and Garnett) based on 3 known cell type datasets. Dataset identity was labeled on top of the panel with cluster numbers in brackets. For legends, "Positive" meant percentage of correctly predicted clusters, while "Negative" meant incorrectly predicted clusters and "Missed" meant predictions with uncertain cell types.

Figure 3. Cell components of PBMCs predicted by SCSA. A. Clustering of uniform

scores of the top 5 predicted cell types in 4 PBMCs datasets by SCSA. Each column

stands for one cluster of 4 PBMCs datasets and each row stands for one cell type.

Uniform scores were normalized using the z-score method to make clusters comparable.

B. Percentages of 4 different cell types in 4 PBMCs datasets based on SCSA's

prediction. C. Five cell types plotted by t-SNE based on the prediction of SCSA for 4

PBMCs datasets.